Zymoseptoria tritici proteins structurally related to UmV-KP4 and UmV-

KP6 are toxic to fungi, and define novel structural families of fungal

effectors

Karine de Guillen[1], Léa Mammri[1], Jérôme Gracy[1], André Padilla[1], Philippe Barthe[1],

François Hoh[1], Mounia Lahfa[1], Justine Rouffet[2], Yohann Petit-Houdenot [2], Thomas

Kroj[3], Marc-Henri Lebrun[2]

[1] Centre de Biologie Structurale, Université de Montpellier, INSERM U1054, CNRS

UMR 5048, 34090 Montpellier, France.

[2] Université Paris-Saclay, INRAE, UR BIOGER, Palaiseau, France

[3] PHIM Plant Health Institute, University of Montpellier, INRAE, CIRAD, Institut Agro,

IRD, 34060 Montpellier, France.

Corresponding Author:

Karine de Guillen

karine.deguillen@cbs.cnrs.fr

tel 33 467417708

Abstract

Fungal effectors play crucial roles in plant infection. Despite low sequence identity, their classification into families is feasible using structural homology. In this study, we have elucidated the structures of the two effectors Mycgr3-91409 and Zt-NIP1 from the wheat fungal pathogen *Zymoseptoria tritici* using X-ray crystallography and NMR. These proteins shared structural homology with, respectively, KP6α and KP4 killer toxins, encoded by UmV dsRNA viruses infecting the corn fungal pathogen *Ustilago maydis*. Consequently, Zt-Mycgr3-91409 and Zt-NIP1 were renamed Zt-KP6-1 and Zt-KP4-1. Orthologs and paralogs of Zt-KP6-1 and Zt-KP4-1 were identified in *Zymoseptoria* species, but not in other fungi, with the exception of ECP2 effectors related to Zt-KP4-1. A novel pipeline relying on Foldseek and cysteine-pattern constrained HMM searches identified a large number of proteins structurally related to Zt-KP6-1 and Zt-KP4-1 in fungi, and to Zt-KP4-1 in plants. These proteins were classified into distinct structural super families (4 KP4 and 3 KP6) including effectors not known as structurally related to KP4 (ECP2) or KP6 (BAS4). The biological activities of Zt-KP6-1 and Zt-KP4-1 were assessed on plants and fungi. They did not induce symptoms on wheat leaves. However, they inhibited the growth of *Botrytis cinerea* and *Z. tritici*, demonstrating their toxicity to fungi. These results highlight the importance of structure determination of effectors for their classification and the prediction of their function.

INTRODUCTION

Fungal plant pathogens express secreted proteins without enzymatic functions during host plant infection called effectors (Lo Presti *et al.*, 2015; Li *et al.*, 2024). Most fungal effectors display no or very limited sequence similarities (Outram *et al.*, 2022; Lazar *et al.*, 2022). However, comparison of their three-dimensional structures allowed their classification into families of structurally related proteins (Outram *et al.*, 2022). This strategy was successful in identifying a large number of Magnaporthe AVRs and ToxB-like (MAX) effectors from the fungal rice pathogen *Pyricularia oryzae* sharing the same conserved fold (de Guillen *et al.*, 2015; Le Naour—Vernet *et al.*, 2023; Lahfa *et al.*, 2024). The RNase-like proteins associated with haustoria (RALPH) effectors of the cereal powdery mildew fungus *Blumeria graminis* were also grouped into a family of structurally related proteins using modeling and experimental structure determination (Spanu, 2017; Pennington *et al.*, 2019; Bauer *et al.*, 2021; Cao *et al.*, 2023). Additional families of structurally related fungal effectors were identified using X-ray diffraction (XRD) and modeling in *Leptosphaeria maculans* (Blondeau *et al.*, 2015; Lazar *et al.*, 2020; Talbi *et al.*, 2023; Loubet *et al.*, 2024), *Fusarium oxysporum* (Teulet *et al.*, 2023; Yu *et al.*, 2023), and *Venturia inaequalis* (Rocafort *et al.*, 2022). Recently, the systematic prediction of structures of fungal secreted proteins with different programs including AlphaFold2 (AF2) (Jumper *et al.*, 2021), revealed novel families of structurally related fungal effectors (Seong & Krasileva, 2023; Derbyshire & Raffaele, 2023).

To identify novel families of fungal effectors, we elucidated the three-dimensional structures of the secreted proteins Zt-Mycgr3-91409 and Zt-NIP1 from the wheat fungal pathogen *Zymoseptoria tritici* by X-ray diffraction (XRD) and nuclear magnetic resonance (NMR). Zt-Mycgr3-91409 is an effector upregulated during infection of wheat leaves (Rudd *et al.*, 2015) suggested to be a MAX effector (de Guillen *et al.*, 2015). Zt-NIP1 is expressed during infection of wheat (Rudd *et al.*, 2015; Kilaru *et al.*, 2022) and induces mild chlorotic/necrotic lesions on leaves (Ben M'Barek *et al.*, 2015). Unexpectedly, the structures of Zt-Mycgr3-91409 and Zt-NIP1 shared structural similarities with UmV-KP6α and UmV-KP4 killer toxins encoded by dsRNA viruses infecting the corn fungal pathogen *Ustilago maydis*. UmV-KP6α and UmV-KP4 kill susceptible *U. maydis* strains, while UmV infected strains are resistant to the killer toxin they produce (Finkler *et al.*, 1992; Allen *et al.*, 2013b). Consequently, we renamed Zt-

Mycgr3-91409 to Zt-KP6-1 and Zt-NIP1 to Zt-KP4-1. Proteins structurally related to Zt-

KP6-1 or Zt-KP4-1 were widely distributed across fungi and they were classified in

families of related structures. Bioassays revealed that Zt-KP6-1 and Zt-KP4-1 were not

toxic to wheat leaves, but inhibited fungal growth.

RESULTS

Zt-Mycgr3-91409-2 is a structural homolog of UmV-KP6 killer toxin

Zt-Mycgr3-91409 gene model was curated according to transcript evidence (Lapalu *et*
al., 2023) (Sup Figure 1A). Zt-Mycgr3-91409-2 (SMR48303, A0A2H1G421) was
expressed in *Escherichia coli* and purified by size exclusion chromatography (SEC)
(Sup File S1, Sup Figure 2). High-resolution diffracting crystals were obtained after
crystallization screens and optimization of Li_2SO_4 concentration and pH then soaked
in tantalum. Analyses of XRD data at 1.36 Å resolution showed that crystals belonged
to the $P2_12_12_1$ space group with two copies of the protein per asymmetric unit. The
structure of Zt-Mycgr3-91409-2 solved by single-wavelength anomalous diffraction
(SAD) revealed that the protein had a typical α/β sandwich structure (Figure 1, Sup
Table 1). This fold has two α-helices and four β-strands arranged in an antiparallel β-
sheet with one hairpin connection formed by the short strands β2-β3 and two βαβ
connections, which placed the α-helices in an antiparallel manner on one side of the
β-sheet.

Figure 1

Figure 1. *Crystal structures of (A) Zt-KP6-1 (6QPK chain A) and (B) UmV-KP6α (4GVB chain A).*

(A) Zt-Mycgr3-91409-2 was renamed to Zt-KP6-1 according to its structural similarity to UmV-KP6α (see text). Structures were colored in rainbow colors from blue (N terminus) to red (C terminus). For Zt-KP6-1, the loop connecting β1 (in blue) to H1 (in cyan) was not observed in the density map and is shown as a blue dashed line. Disulphide bonds are shown in grey and labelled Dx. For the D1 (C26-C86) bond in Zt-KP6-1, both conformations are shown.

The α-helix H2 was preceded by a 3^{10} helix formed by residues 77 to 89 (numbering according to the mature protein). Both α-helices (H1 and H2) were tightly packed against the β-sheet, with one disulphide bond formed between cysteine residues in β1 (Cys26) and H2 (Cys86) and another disulphide bond formed between cysteine residues in H1 (Cys43) and β3 (Cys65), making the fold very compact. Alternative conformations were observed for Cys26 involving the disulphide bond with Cys86. The interface area between the two monomers observed in the crystal, determined by PDBePISA, was estimated to be 435.8Å^2. The Complexation Significance Score (CSS) was 0, indicating that the dimer has no biological relevance and resulted from crystal packing. NMR studies of the protein labelled with ^{15}N revealed a high similarity to the crystal structure, with an r.m.s.d. of 0.7 ± 0.6 Å on structured parts. However, the protein was not dimeric in solution, as shown by the narrowness of the ^{1}H-^{15}N HSQC spectrum peaks (Sup Figure 3 and Sup Table 2), and the single elution profile at 95 min observed in SEC corresponding to a retention time of a 9 kDa globular protein (Sup Figure 2).

A DALI analysis (Holm, 2020) revealed structural similarity between Zt-Mycgr3-91409-2 and UmV-KP6α (PDB 4GVB, Figure 1). UmV-KP6α is encoded by a dsRNA virus infecting *U. maydis,* a fungal pathogen causing corn smut disease (Allen *et al.*, 2013b). UmV-KP6α is produced together with UmV-KP6β by cleavage of the UmV-KP6 pre-protein encoded by the virus. UmV-KP6α has eight cysteine residues, while UmV-KP6β has six cysteines, all involved in intramolecular disulphide bonds. Both UmV-KP6α and UmV-KP6β have a highly similar fold (Li *et al.*, 1999; Allen *et al.*, 2013b). Zt-KP6-1 structure was more similar to UmV-KP6α (Z-score of 6.7 and r.m.s.d of 2.5 Å for 63 aligned residues). However, Zt-KP6-1 did not have the N-terminal hydrophobic

α-helix extension present in UmV-KP6α. The KP6 protein fold shared between UmV-KP6α, UmV-KP6β and Zt-KP6-1, is defined by two α-helices (H1 and H2) positioned on one side of a 4-stranded β-sheet, with a β4-β1-H1-β3-H2-β2topology (Sup Figure 4). All three proteins have two right handed β-α-β split crossovers, which distinguish the KP6 protein fold from toxins of similar size (Allen *et al.*, 2013a). The KP6 protein fold also shared two disulphide bonds that link the antiparallel α-helices to the central β-sheet of the protein (Sup. Fig. 4). According to these structural similarities, Zt-Mycgr3-91409-2 was renamed Zt-KP6-1.

Zt-NIP1 is a structural homolog of *UmV*-KP4 killer toxin

Zt-NIP1 was identified as a wheat necrosis-inducing protein produced by IPO323 *Z. tritici* isolate during wheat leaf infection (Ben M'Barek *et al.*, 2015). However, the CDS at the *Zt-NIP1* locus in IPO323 genome (*Zt-Mycgr3-106176*) encoded a protein with a sequence unrelated to NIP1. A new CDS, almost identical to NIP1 (99% identity), was predicted in agreement with transcript evidence (Lapalu *et al.*, 2023) (Sup Figure 1B). Zt-NIP1 (SMR6055, A0A2H1H404) was expressed in *Escherichia coli* and purified by SEC (Sup File S1). SEC revealed an equilibrium between Zt-NIP1 monomers and dimers in solution (Sup Figure 5). High-resolution diffracting Zt-NIP1 crystals were in the *H3* space group with 8 copies of the protein per asymmetric unit (Sup Table 1). The Zt-NIP1 structure was solved by molecular replacement using an AF2 model (Figure 2). The protein has a typical α/β sandwich structure with a single split βαβ motif and two disulphide bonds. A main antiparallel 5 stranded β-sheet is formed by β1-β3-β4-β6-β7, with a second small antiparallel 2 stranded β-sheet comprising β2 and β5 (with 4 residues each). Two α-helices (H1 and H2) close the structure. The fold is held together by two disulphide bonds formed between cysteine residues in β2 (Cys39) and β7 (Cys142), and H1 (Cys60) and β4 (Cys89). Chains E-H, G-B, F-C, and D-A formed dimers, through the main 5 stranded β-sheet, in the crystal structure, with an average interface area of 852.2 $Å^2$ determined by PDBePISA. These dimers exhibited a CSS score of 1.0, which indicates the biological significance of such a dimer (Sup Figure 6).

Figure 2

Figure 2. *Crystal structures of (A) Zt-KP4-1 (8ACX chain A) and (B) UmV-KP4 (1KPT chain A).*

(A) Zt-NIP1 was renamed to Zt-KP4-1 according to its structural similarity to UmV-KP4 using DALI (see main text). Structures were colored in rainbow colors from blue (N terminus) to red (C terminus). For Zt-KP4-1 the loops connecting β6 to β7 and β6 to H1 were not observed in the density map and were shown as dashed lines. Disulphide bonds are shown in grey and labelled Dx.

A DALI analysis (Holm, 2020), revealed structural similarity between Zt-NIP1 and UmV-KP4 (PDB 1KPT, Figure 2). UmV-KP4 is encoded by a dsRNA virus infecting *U. maydis* (Allen *et al.*, 2013b). The two proteins superposed with a Z-score of 7.3 and a r.m.s.d value of 2.4 Å (82 residues aligned). UmV-KP4 (Koltin, 1988; Gu *et al.*, 1995) belongs to the α/β sandwich family of proteins and is composed of seven β strands and three α-helices. Similar to Zt-NIP1, UmV-KP4 has two helices (H2 and H3) in a left-handed βαβ crossover conformation, which is a rare structural feature (Sup Figure 7) (Allen *et al.*, 2013a). The main difference between the two proteins is the presence of a β-strand (β1) at the N-terminus for Zt-NIP1 instead of a short α-helix (H1) for UmV-

184 KP4. The two proteins also differ in the number and location of disulphide bonds. While
185 Zt-NIP1 has two disulphide bonds (as described above), UmV-KP4 has five. Based on
186 the structural similarity of these proteins, Zt-NIP1 was renamed to Zt-KP4-1.
187
188 **Zt-KP4 and Zt-KP6 are toxic to fungi**
189
Zt-KP4-1 and Zt-KP6-1 (each at a concentration of 2 µM) were infiltrated into leaves of
the wheat cultivar Taichung 29 which is susceptible to *Z. tritici*. Neither of the two
proteins induced visible signs of toxicity such as leaf chlorosis or necrosis. In the same
experiment, infiltration of the necrotrophic effector ToxA from *P. nodorum* (Tan *et al.*,
2012) into Taichung 29 wheat leaves induced typical necrotic lesions at concentrations
ranging from 0.2 to 2 µM. To test if Zt-KP4-1 and Zt-KP6-1 are toxic to fungi, growth
assays were performed in liquid potato dextrose growth medium complemented with
recombinant Zt-KP4-1 or Zt-KP6-1 at different concentrations (0.55 to 5.5 µM). The *Z.*
tritici IPO323 isolate, which expresses both Zt-KP4-1 and Zt-KP6-1, and the
necrotrophic plant pathogen *Botrytis cinerea* were used for these bioassays. Zt-KP4-1
strongly inhibited the growth of both fungi at 5.5 µM (91-95 % inhibition, Table 1).
However, at 1.65 µM, only *Z. tritici* growth was inhibited (48 % inhibition, Table 1). Zt-
KP6-1 was as toxic as Zt-KP4-1 on *B. cinerea* (94 % inhibition at 5.5 µM, Table 1).
However, it only weakly inhibited the growth of *Z. tritici* (24 % inhibition at 5.5 µM, Table
1). These results demonstrate that Zt-KP4-1 and Zt-KP6-1 are not toxic to wheat leaves
but are toxic to fungi at concentrations below those measured for fungal toxic proteins
(20 µM) (Kettles *et al.*, 2018; Lima *et al.*, 2022). While Zt-KP4-1 displayed the same
toxicity to *Z. tritici* and *B. cinerea,* Zt-KP6-1 displayed some quantitative selectivity*,* as
Z. tritici was 4-fold less sensitive than *B. cinerea*.
209

		% inhibition	
		Botrytis cinerea	*Zymoseptoria tritici*
Protein concentration (µM)			
TOX A	5.50	25 ± 1.4	0
	1.65	0	0
	0.55	0	0
Zt-KP4-1	5.50	91 ± 3.2	95 ± 1.4
	1.65	0	48 ± 2.2
	0.55	0	25 ± 1.4
Zt-KP6-1	5.50	94 ± 2.0	24 ± 1.2
	1.65	0	0
	0.55	0	0

Table 1. *Toxicity of Zt-KP4-1 and Zt-KP6-1 to fungi*

The percentages of B. cinerea *and* Z. tritici *growth inhibition were calculated for different concentrations of ToxA, Zt-KP4-1and Zt-KP6-1. Values are the mean ± Standard Error of the Mean (SEM) from three experiments.*

Identification of Zt-KP6-1 homologs in *Zymoseptoria*

Proteins with sequences highly similar to Zt-KP6-1 were identified in other strains of *Z. tritici* (Badet *et al.*, 2020) and *Z. pseudotritici* (Zp-KP6-1, 92% identity, Figure 3), but not in other *Zymoseptoria* species. Another protein family with a relatively low sequence identity to Zt-KP6-1 (28-30%) was identified in *Z. tritici* (Zt-KP6-2) and other *Zymoseptoria* species (KP6-2 from *Z. brevis, Z. arbibiliaie* and *Z. pseudotritici*, Figure 3). The structure of Zt-KP6-2 protein predicted by AF2 was similar to Zt-KP6-1 (Z-score of 12.1; r.m.s.d of 1.9 Å for 73 aligned residues, Sup Figure 8). This comparison highlighted an 11 amino acid C-terminal sequence with two cysteine residues conserved in all KP6-2 proteins, but absent from KP6-1 proteins. Proteins similar to Zt-KP6-1 and Zt-KP6-2 were not identified in other fungi. The phylogenetic analysis of KP6-1 and KP6-2 proteins from different *Zymoseptoria* species highlighted two clades gathering orthologous proteins from *Zymoseptoria* (Figure 3). This topology suggested that KP6-1 and KP6-2 are paralogs originating from the duplication of an ancestral gene before the divergence of *Zymoseptoria* species.

Figure 3. Phylogenetic tree of *Zymoseptoria* KP6-1 and KP6-2

MAFFT alignment, BMGE cleaning, PhyML Phylogenetic tree using NGPhylogeny.fr.

(Lemoine *et al.*, 2019) Bootstrap values > 70% are indicated under each branch after

the supported node. Midpoint tree rooting. Zt-KP6-1_Ztritici_3D7: Uniprot

A0A1X7RM13 (identical to Zt-KP6-1_IPO323); Zt-KP6-1_Ztritici_1E4: Uniprot

A0A2H1G421, PDB 6QKP; Zp-KP6-1_Zpseudotritici: JGI Zymps1_802032; Zt-KP6-

2_Ztritici_1E4: Uniprot A0A2H1H0K4; Zt-KP6-2_Ztritici_IPO323: Uniprot F9XLY9; Zt-

KP6-2_Ztritici_3D7: Uniprot A0A1X7S5L2; Zb-KP6-2_Zbrevis: Uniprot A0A0F4GMG5;

Za-KP6-2_Zardabiliae: JGI Za_774343 ; Zp-KP6-2_Zpseudotritici: JGI

Zymps1_805921.

Using a pipeline based on FoldSeek and Hmmer (Sup Figure 10), we searched for

proteins structurally related to Zt-KP6-1 and Zt-KP6-2 in *Zymoseptoria*. The structures

of the retrieved proteins were predicted using AF2 and visually inspected for structural

overlaps with Zt-KP6-1. Three proteins were structurally related to Zt-KP6-1: Zt-KP6-

4, Zt-KP6-5 and Zb-KP6-6. However, Zt-KP6-3 did not share sufficient structural

similarity to KP6-1 (DALI score of 0) to be selected and it was excluded from further

analyses.

Identification of Zt-KP4-1 homologs in *Zymoseptoria*

Proteins with sequences highly similar to Zt-KP4-1 were identified in other strains of *Z.*

tritici (Badet *et al.*, 2020) (Figure 4), but not in other *Zymoseptoria* species. Two other

protein families with a relatively low sequence identity to Zt-KP4-1 (33-35%) were

identified in *Z. tritici* (Zt-KP4-2, 35% identity; Zt-KP4-3, 33% identity) and *Z. brevis* (Zb-

KP4-2; Zb-KP4-3). The structures of *Zymoseptoria* KP4-2 and KP4-3 proteins

predicted with AF2 were similar to Zt-KP4-1 (Sup Figure 9). A phylogenetic analysis of *Zymoseptoria* KP4-1, KP4-2 and KP4-3 was performed using related ECP2 proteins as outgroups (Van den Ackerveken *et al.*, 1993; Chang *et al.*, 2016; Carreón-Anguiano *et al.*, 2023). ECP2 proteins were retrieved as the top hits in BlastP searches of the proteomes of the corresponding fungi using Zt-KP4-1 and Zt-KP4-2 as queries, although with relatively low scores (28-32% identity, e-value 10^{-20}). The *Zymoseptoria* KP4-1, KP4-2 and KP4-3 proteins clustered as three independent clades gathering orthologous proteins from *Zymoseptoria*. These three clades were rooted by ECP2 proteins from other fungi (Figure 4). This topology suggested that KP4-1, KP4-2 and KP4-3 originated from duplications of an ancestral gene, prior to the divergence of *Zymoseptoria* species.

Figure 4: Phylogenetic tree of *Zymoseptoria* KP4-1, KP4-2 and KP4-3

MAFFT alignment, BMGE cleaning, PhyML Phylogenetic tree using NGPhylogeny.fr (Lemoine *et al.*, 2019). Bootstrap values > 70% are indicated under each branch after the supported node. Tree root with Ecp2_Passalora fulva and Ecp2_Pseudocercospora fijiensis as outgroups. Zt-KP4-1_Ztritici_IPO323: re-annotated NIP1/Mycgr3-106176 see Sup. Fig1; Zt-KP4-1_Ztritici_1E4: Uniprot A0A2H1H404; Zt-KP4-1_Ztritici_3D7: Uniprot A0A1X7S6U3; Zt-KP4-1_Ztritici_1A5: Uniprot A0A1Y6M012; Zt-KP4-2_Ztritici_IPO323: Mycgr3-107904, Uniprot F9X223; Zb-KP4-2_Zbrevis: KJY02238, Uniprot A0A0F4GYX0; Zt-KP4-3_Ztritici_IPO323: Mycgr3-111636, Uniprot F9XQ02; Zt-KP4-3_Ztritici_3D7: Uniprot A0A1X7S9R6; Zb-KP4-3_Zbrevis: KJX96086, Uniprot A0A0F4GF74; Ecp2_Passalora fulva: Uniprot

Q00365; Ecp2_Pseudocercospora fijiensis: Uniprot M2Z6V3; Ecp2_Sphaerulina

musiva: Uniprot N1QK49; Ecp2_Cochliobolus sativus: Uniprot M2TE34

Using a pipeline based on FoldSeek and Hmmer (Sup Figure 10), we searched for

proteins structurally related to Zt-KP4-1, Zt-KP4-2 and Zt-KP4-3 in *Zymoseptoria*.

Zymoseptoria KP4-1, KP4-2 and KP4-3 have an HCE2 domain (PF14856). This

domain was defined using proteins related to the fungal effector ECP2 from *P. fulvum*

(Stergiopoulos *et al.*, 2012). Therefore, *Zymoseptoria* proteins with an HCE2 domain

were retrieved from Uniprot. The HCE2 domain is a multi-gene superfamily that has

been named Hce2 for homologues of the *C. fulvum* effector Ecp2, which is widely

distributed in fungi, with a putative function as a necrosis-inducing factor. The

structures of these proteins were predicted using AF2 and visually inspected for

structural overlaps with Zt-KP4-1. This analysis identified four additional proteins

structurally related to Zt-KP4-1: Zt-KP4-5, Zt-KP4-6, Zt-KP4-7 and Zt-KP4-9. Two

proteins, Zt-KP4-4 and Zt-KP4-8, were not sufficiently structurally to KP4-1 to be

selected, despite the presence of the same number of α−helices and β strands

organized in the same order as in Zt-KP4-1. In Zt-KP4-4, the structures of the β2 and

β5 strands were not well defined (twisted), and the β2-β3 and β5-β6 interactions were

reduced. In Zt-KP4-8, the interactions between the β1 and β4 strands were missing.

These two proteins were excluded from further analyses.

Zymoseptoria *KP4* and *KP6* are expressed during infection

Available IPO323 RNA-Seq data (Rudd *et al.*, 2015) were used to retrieve the

expression patterns of Zt-KP4 and Zt-KP6 (Sup Fig. 11). *Zt-KP4-5, Zt-KP4-6* and *Zt-*

KP4-7 were expressed at low levels (1/10th of beta-tubulin *TUB1*) in all conditions, while

Zt-KP4-9 was not expressed. *Zt-KP4-1* was the most highly expressed *Zt-KP4* gene

during infection of wheat leaves (25 x *TUB1*), with maximum expression at 9 dpi. It was

also expressed during *in vitro* growth, although to a lesser extent. *Zt-KP4-3* was

expressed during infection at 9 and 14 dpi, and during *in vitro* growth. *Zt-KP4-2* was

constitutively expressed at a level similar to *TUB1*. *Zt-KP6-1*, *Zt-KP6-2*, *Zt-KP6-4* and

Zt-KP6-6 displayed a similar expression pattern with maximum expression during

infection at 9 dpi and highly reduced expression during *in vitro* growth (100 times less

than during infection). This survey showed that most *Zymoseptoria KP4* and *KP6* were

over-expressed during the switch from the asymptomatic phase to the necrotrophic phase of infection (9-14 dpi). This switch is associated with the massive colonization of the apoplast of the infected wheat leaf by infectious hyphae of *Z. tritici*, and their invasion of sub-stomatal cavities where they differentiate into pycnidia (Battache *et al.*, 2022).

Proteins with Zt-KP6-1 structural folds are widespread in fungi

Structure modelling of Ecp28-1, Ecp28-2 and Ecp28-3 effectors from the tomato pathogen *Passalora fulvum* (Mesarich *et al.*, 2018) suggested that they have a KP6 fold. AF2 structure modelling of these proteins supported this hypothesis (DALI Z score of 7.4). We used our pipeline (Sup Figure 10) to detect fungal proteins structurally related to Zt-KP6-1 in 1014 fungal genomes. The KP6 candidates were selected if their AlphaFold models had a TM-score above 0.5 compared to either Zt-KP6-1 or UmV-KP6α using TM-Align (Zhang & Skolnick, 2005). A total of 1506 KP6 candidates were identified and grouped into 81 AFDB50 clusters (kclXX-Y). The representative proteins of these AFDB50 clusters were merged into 46 meta-clusters (kclXX) (Sup Figure 10). The structure of one representative protein from each meta-cluster was visually inspected using PyMOL for the occurrence of a typical KP6 topology. This analysis rejected nine meta-clusters. The remaining meta-clusters consisted of 1148 proteins with a KP6 fold (sup Table 3) including ECP28 (kcl14), Zt-KP6-1 (kcl06), Zt-KP6-2 (kcl06), Zt-KP6-4 (kcl02), Zt-KP6-5 (kcl07) and Zt-KP6 (kcl07). Several fungal effectors have recently been described as KP6 proteins: AvrLm10A and AvrLm6 from *L. maculans* (Talbi *et al.*, 2023; Loubet *et al.*, 2024), SIX5 from *Fusarium oxysporum* (Talbi *et al.*, 2023), BAS4 from *Pyricularia oryzae* (Khang *et al.*, 2010), and Cb-Nip1 from *Cercospora beticola* (Ebert *et al.*, 2021). The systematic analysis of the structures of the effectors of *V. inaequalis* and *L. maculans* also revealed 108 and 13 KP6 proteins, respectively (Rocafort *et al.*, 2022; Loubet *et al.*, 2024). Only few of these proteins were identified in our analysis, since most were not in Uniprot. We added these proteins to meta-cluster representatives. A visual inspection of the structural overlaps between these additional proteins and Zt-KP6-1 rejected only Cb-NIP1, since its AF2 model was not sufficiently structured to allow its superposition to Zt-KP6-1.

The structures of the 36 selected KP6 proteins were compared using a principal

component analysis (PCA) with DALI Z-scores (Figure 5). This analysis highlighted three groups of structurally related KP6 proteins. Group-1 included AvrLm6 and AvrLm10A from *L. maculans* and SIX5 from *F. oxysporum,* but not KP6 proteins from *Zymoseptoria.* Group-2 contained Zt-KP6-4, ECP28, BAS4, four cluster representatives (kcl01-C, kcl02-G, kcl04-A, kcl11), and UmV-KP6β*.* Group-3 encompassed Zt-KP6-1, UmV-KP6α, Zt-KP6-2, Zt-KP6-6, and cluster representative kcl02-A.

Figure 5 Classification of fungal KP6 families according to their structures

DALI all by all analysis followed by a PCA using a DALI score matrix (Holm, 2020).

meta-cluster kclXX, cluster AFDB50 kclXX-Y, (number of proteins in AFDB50 cluster), Uniprot representative (example A0A2H1G421): kcl01-C, (100), A0A4V3HRP1, M1W1H7; kcl01-L, (65), SIX5 = A0A010RIE2, AvrLm10A = E5A720_LEPMJ; kcl02-A, (5), A0A4R8TM13; kcl02-D, (138), Zt-KP6-4 = A0A2H1GZA7; kcl02-G, (321), A0A135SXI2; BAS4 = BAS4_MAGO7, Lm-KP6 = E5A0W1_LEPMJ; kcl04-A, (1), F9FAC2; kcl04-C, (25), AvrLm6 = A0A010QJV2; AvrLm6 = E5R515_LEPMJ; kcl06-A, (6), Zb-KP6-2 = A0A0F4GMG5, Zt-KP6-2 = A0A2H1H0K4; kcl06-B, (3), Zt-KP6-1 = A0A2H1G421/6QKP; kcl07-A, (5), Zt-KP6-5 = A0A2H1FMK1, Zb-KP6-6 = A0A0F4GGQ1; kcl11-A, (6), A0A317AKI4; kcl14-A, (93), ECP28-2 = A0A1P8YXM1, ECP28-3 = A0A1P8YXM5; kcl14-B, (14), ECP28-1 = A0A1P8YXQ5; LmXXXX, *Leptosphaeria* KP6 proteins, (8 proteins, (Loubet *et al.*, 2024)); Vi-KP6-famXX, *Venturia* KP6 proteins, (4 families,(Rocafort *et al.*, 2022)); White circle, UmV-KP6α, 4gvbA,

375   White circle, UmV-KP6β, 4gvbB

Proteins with Zt-KP4-1 structural folds are also widely distributed in fungi

The HCE2 domain (PF14856) detected in KP4-1, KP4-2 and KP4-3 relies on a pattern

of three conserved cysteine residues and a W/F/Y motif (Stergiopoulos *et al.*, 2012).

The KP4 domain (PF09044) defined for *F. graminearum* proteins Fg-KP4-1, Fg-KP4-

2 and Fg-KP4-3 and UmV-KP4 relies on a pattern involving four conserved cysteine

residues, three conserved glycine residues and two W/F/Y motifs (Brown, 2011; Lu &

Faris, 2019). We did not detect overlaps in proteins retrieved independently from

Uniprot with either the HCE2 or the KP4 domain. Fungal proteins with an HCE2 domain

retrieved from Uniprot included Zt-KP4-1, Zt-KP4-2 and Zt-KP4-3. There were no

Zymoseptoria proteins with a KP4 domain. Additional KP4 proteins were detected

using our pipeline (Sup Figure 10) using Zt-KP4-1 and UmV-KP4 as references. A total

of 3722 KP4 candidates were identified and grouped into 153 AFDB50 clusters. The

representative proteins of these AFDB50 clusters were merged into 59 meta-clusters.

The structure of one representative protein from each meta-cluster was visually

inspected using PyMOL for the occurrence of a typical KP4 topology. This analysis

rejected 31 meta-clusters. The remaining meta-clusters contained 2825 proteins with

a KP4 fold (sup Table 4) including ECP2 (kcl01), Zt-KP4-1 (kcl01), Zt-KP4-2 (kcl07),

Zt-KP4-3 (kcl07), Zt-KP4-5 (kcl18), Zt-KP4-6 (kcl09), Zt-KP4-7 (kcl09) and Zt-KP4-9

(kcl02). We added Fg-KP4-1, Fg-KP4-2 and Fg-KP4-3, which were used to define the

KP4 domain, to our analysis (Brown, 2011; Lu & Faris, 2019). These proteins were

members of meta-cluster kcl02, but they were not selected as representative of this

cluster. Visual inspection of these proteins confirmed their structural similarity to Zt-

KP4-1.

The structures of the 26 selected KP4 proteins were compared using a PCA with DALI

Z-scores (Figure 6). This analysis highlighted four groups of structurally related KP4

families. Group-1 included Fg-KP4-1, Fg-KP4-2 and Fg-KP4-3, UmV-KP4 and kcl02

representative. All these proteins had a KP4 domain. Group-2 contained Zt-KP4-1, Zt-

KP4-2 and Zt-KP4-3 and ECP2, as well askcl07, kcl10 and kcl14 representatives. All

proteins from Group-2 had an HCE2 domain. Group-3 included the representatives of

the clusters kcl08-A and kcl38-A. Proteins from Group-3 did not have known Pfam

domains, except kcl38-A proteins which contained the domain PF15474. Group-4 contained all the other KP4 proteins, including Zt-KP4-5, Zt-KP4-6, Zt-KP4-7 and Zt-KP4-9. Proteins from Group-4 did not have a Pfam domain, except kcl21-A proteins which contained the domain PF21691.

Figure 6 Classification of fungal KP4 families according to their structures

DALI all by all analysis followed by a PCA using a DALI score matrix (Holm, 2020). meta-cluster kclXX, cluster AFDB50 kclXX-Y, (number of proteins in AFDB50 cluster), Uniprot representative (example A0A2H1H404): kcl01-D, (83), Zt-KP4-1 = A0A2H1H404/8ACX; kcl01-Q, (203), Sm-ECP2 = N1QK49, Psf-ECP2 = M2Z6V3, Paf-ECP2 = Q00365; kcl02-A, (15), A0A4U0XEI9; kcl02-E, (256), Fg-KP4-2 = A0A3S7E9S6, Fg-KP4-3 = A0A2H3GQS6; kcl02-I, (125), Fg-KP4-1 = A0A2H3GK41; klc02-0, (324), Zt-KP4-9 = A0A2H1GXX1; kcl05-A, (14), A0A6A6DAE0; kcl07-A, (53), A0A177DVW3; kcl07-C, (35), Zt-KP4-2 = A0A2H1FWY7; Zt-KP4-3 = A0A1Y6M296; kcl08-A, (5), S3D5P; kcl09-A, (14), M3BT27; kcl09-B, (131), Zt-KP4-6 = A0A2H1GNH3, Zt-KP4-7 = A0A2H1GND9; kcl10-A, (16), A0A1L9SH47; kcl12-A, (9), E3L0Y8; kcl14-C, (12), A0A0A1V5R3; kcl18-A, (9), Zt-KP4-5 = A0A2H1GP20; kcl21-A, (13), A0A0J9W663; kcl27-A, (39), A0A4T0WCM2; kcl38-A, (8), A0A017SS08; White circle, UmV-KP4 = kptA

DISCUSSION

Comparison of the structures of fungal effectors allowed their classification into families of proteins with related structures (de Guillen *et al.*, 2015; Seong & Krasileva, 2023; Derbyshire & Raffaele, 2023). In this study, the elucidation of the structure of two *Zymoseptoria tritici* effectors (Zt-Mycgr3-91409-2, Zt-NIP1) by X-ray crystallography showed their unexpected structural similarity to the UmV-KP6α and UmV-KP4 killer toxins from the dsRNA virus infecting the fungal pathogen *U. maydis*. Therefore, they were renamed Zt-KP4-1 and Zt-KP6-1. We showed that both effectors were toxic to fungi, but not to wheat leaves. In addition, we found that proteins structurally related to Zt-KP4-1 and Zt-KP6-1 are widespread in fungi, and could be classified into super families of structurally related proteins (4 KP4 and 3 KP6).

***UmV*-KP6 and *UmV*-KP4 killer toxins**

U. maydis secretes three killer proteins (UmV-KP1, KP4 and KP6) encoded by double-stranded RNA viruses (UmV) (Tao *et al.*, 1990; Gage *et al.*, 2001; Allen *et al.*, 2013b). They kill susceptible *U. maydis* strains, while UmV-infected strains are resistant to the killer protein they produce (Finkler *et al.*, 1992; Allen *et al.*, 2013b). UmV-KP4 inhibits fungal growth presumably by perturbing cellular calcium homeostasis (Gage *et al.*, 2001). UmV-KP4 also inhibited L-type voltage-gated (LVG) Ca^{2+} channels in mammalian cells (Gage *et al.*, 2002). UmV-KP4 inhibited *A. thaliana* root growth, and perturbed Ca^{2+} gradients of root hair cells (Allen *et al.*, 2008). The *P. patrens* moss KP4 protein also modified Ca^{2+} gradients of moss cells (Guan *et al.*, 2023). However, UmV-KP4 does not induce major growth defects, necrosis, chlorosis or cell death when over-expressed in transgenic wheat, corn or tobacco (Clausen *et al.*, 2000; Schlaich *et al.*, 2006; Quijano *et al.*, 2016; Deb *et al.*, 2018). UmV-KP6α and UmV-KP6β associate to form a stable heterodimer (Li *et al.*, 1999). UmV-KP6α appears necessary for recognition of susceptible cells, through a mechanism involving the N-terminal α helix, while UmV-KP6β is required for toxicity (Allen *et al.*, 2013b). However, the molecular mechanism underpinning the toxicity of UmV-KP6α/β to fungal cells is unknown (Allen *et al.*, 2013b), although it has been hypothesised that this complex forms membrane pores (Li *et al.*, 1999).

Zt-KP6-1 and Zt-KP4-1 are antifungal proteins

Neither Zt-KP4-1 nor Zt-KP6-1 induced necrosis when infiltrated into wheat leaves. Similarly, Fg-KP4L-2 from *F. graminearum* did not induce necrosis of wheat leaves (Lu & Faris, 2019). Still, recombinant Fg-KP4L-2 slightly inhibited the growth of wheat roots (Lu & Faris, 2019). This is reminiscent of the low activity of UmV-KP4 on *A. thaliana* root growth (Allen *et al.*, 2008). We showed that both Zt-KP4-1 and Zt-KP6-1 inhibited the growth of *Z. tritici* and *B. cinerea*. Within *U. maydis*, UmV-KP4 and UmV-KP6 were only toxic to susceptible strains (Allen *et al.*, 2013b). However, UmV-KP4 inhibited the growth of other fungi such as *Tilletia caries*, *Ustilago tritici*, *Alternaria alternata* and *Phoma exigua* (Deb *et al.*, 2018), but not the yeast *S. cerevisae* (Park *et al.*, 1994). When overexpressed in transgenic wheat or tobacco, UmV-KP4 conferred resistance to, respectively, *T. caries* and *U. tritici* (Clausen *et al.*, 2000; Schlaich *et al.*, 2006; Quijano *et al.*, 2016), and *A. alternata* and *P. exigua* (Deb *et al.*, 2018). UmV-KP6 was also toxic to the yeast *Brettanomyces bruxellensis* (Santos *et al.*, 2011), but not *S. cerevisae* (Tao *et al.*, 1990; Santos *et al.*, 2011). The structure-activity relationship of Zt-KP6-1 likely differs from UmV-KP6α/β as it did not require the formation of a heterodimer to be toxic (Li *et al.*, 1999; Allen *et al.*, 2013b). In addition, Zt-KP6-1 does not have the N-terminal α-helical extension of UmV-KP6α critical for its toxicity (Li *et al.*, 1999; Allen *et al.*, 2013b). Since structural homologs of Zt-KP4-1 and Zt-KP6-1 are widespread in fungi, it is tempting to speculate that some of these KP4 and KP6 proteins are toxic to fungi. Recent work on plant pathogenic fungi highlighted effectors with antimicrobial activities (Snelders *et al.*, 2022). For instance, the cotton pathogenic fungus *Verticiulium dahliae* secretes several effectors during host xylem invasion that have an antibacterial activity (Snelders *et al.*, 2020). These antimicrobial effectors changed the composition of the xylem and root bacterial microbiota of host plants (Snelders *et al.*, 2021, 2023). In particular, VdAve1, an effector with antibacterial activity, specifically inhibits the growth of *Sphingomonas* bacteria, which act antagonistically toward *V. dahliae* in the host plant rhizosphere (de Jonge *et al.*, 2012; Snelders *et al.*, 2020). AMAPEC, a machine-learning based computational software using shared structural and physicochemical properties of known antimicrobial proteins, identified a large number of candidate antimicrobial proteins among effectors of three different plant-associated fungi (Mesny & Thomma, 2024). This finding suggests that fungi have extended repertoires of antimicrobial effectors. We have used

AMAPEC to predict the antimicrobial activities of the KP4 and KP6 proteins we
identified. Interestingly most of these KP6 and KP4 proteins, including Zt-KP4-1 and
Zt-KP6-1, were predicted to have antimicrobial activities by AMAPEC (83% of KP6,
77% of KP4, Sup Table 5).

Fungal KP4 and KP6 proteins are wide spread in fungi and define novel
structural superfamilies

Using a novel pipeline, we identified a large number of KP4 and KP6 structural
homologs in fungi. This novel pipeline is based on (i) FoldSeek structure similarity
searches using AF2 structural models of fungal secreted proteins, combined with (ii)
HMM pattern searches relying on Zt-KP4/UmV-KP4 or Zt-KP6/UmV-KP6α conserved
cysteine patterns and (iii) structure-based manual curation (Sup. Fig. 11; FHAT
(Foldseek Hmmer Alphafold TMalign) in Figure 7). The topology and structure of KP4
and KP6 proteins were manually compared to Zt-KP4/UmV-KP4 or Zt-KP6/UmV-
KP6α. This procedure discriminated proteins with a true KP4 or KP6 structural fold
from proteins with limited or no structural similarity. To compare our dataset (FHAT)
with those of two recent systematic structural studies on fungal effectors (Seong:
Seong & Krasileva, 2023; Derbyshire: Derbyshire & Raffaele, 2023), we clustered all
the KP4 and KP6 proteins identified in these three studies into families of highly related
proteins (at least 90% sequence identity and 60% coverage). This analysis identified
1737 KP4 families for FHAT, 227 for Seong and 333 for Derbyshire, as well as 559
KP6 families for FHAT, 97 for Seong and 80 for Derbyshire. This result was expected
since we have analyzed more fungal species than the two previous studies. We have
then compared these datasets for the seven species shared across these three studies
(Figure 7). For example, in *Z. tritici*, 60% of the FHAT KP4 and KP6 were jointly
identified by Seong and Derbyshire (Figure 7). Adding KP4 and KP6 detected only by
FHAT and Seong and those detected only by FHAT and Derbyshire increased the
number of KP4 and KP6 shared between FHAT and either Seong or Derbyshire to
80%. Eleven putative KP4 and two putative KP6 were specifically detected by
Derbyshire in *Z. tritici*. However, the manual analysis of their structures showed that
they do not have the KP4 or the KP6 fold. Therefore, these proteins are likely false
positive from this pipeline. No false positive were detected in *Z. tritici* by Seong. The
same trend was observed for the six other species (Figure 7). This result suggested

that the FHAT method is more efficient than methods used in these two previous

analyses (slightly more KP4 and KP6 detected, less false positive).

KP4 and KP6 proteins were found in *ascomycotina* and *basidiomycotina* with different

lifestyles, including plant pathogens, plant symbionts and saprophytes. A notable

exception were animal fungal pathogens which did not have KP4 proteins. Contrary to

KP6 proteins, KP4 proteins were identified in plants. Some plant KP4 clustered with

Fg-KP4 and UmV-KP4 (kcl02-E), as observed by Brown (2011), Lu and Faris (2019),

and Guan (2023). Another cluster (kcl01-R) included 175 plant KP4 proteins containing

the same Pfam domain (PF1657) found in some plant antifungal proteins such as

Ginkgobilobin-2.

Figure 7. Venn Diagram comparing KP4 and KP6 proteins identified by different

structural classification methods.

Comparison between the FHAT protocol (Sup. Fig. 10) to other structural classification

methods (Seong:(Seong & Krasileva, 2023)), (Derbyshire:(Derbyshire & Raffaele,

2023)). KP4 and KP6 protein sequences identified in the seven fungal species

analysed by the three studies were clustered into families with at least 90% sequence

identity and 60% sequence overlap. The numbers of effector families detected by one

or more methods are indicated in the Venn diagrams for KP4 and KP6 proteins of either

the seven species or only Z. tritici included in the seven species analysed.

Classification of the KP4 proteins according to their structural similarities using PCA

and DALI Z-scores (see Fig. 6) revealed four structural super families not identified

before. Group-1 only contained proteins carrying the KP4 domain including Fg-KP4-1,

Fg-KP4-2, Fg-KP4-3, UmV-KP4, and 396 proteins from clusters kcl02-A, kcl02-E and

klc02-I. This group overlaps with the KP4 super family defined using Fg-KP4 and UmV-

KP4 by Lu and Faris (2019), and Brown (2011) that includes proteins carrying the KP4

Pfam domain and some plant KP4 (Guan *et al.*, 2023). Group-2 overlaps with the ECP2

super family that includes proteins carrying the HCE2 Pfam domain (Stergiopoulos *et*

al., 2012). However, ECP2 proteins were not yet predicted to have a KP4 fold. This

structural group included *Zt*-KP4-1, Zt-KP4-2 and Zt-KP4-3, ECP2 proteins, and 402

proteins from the clusters kcl01-D, kcl01-Q, kcl07-A, kcl10-A and kcl14-A. Among KP4

proteins from structural Groups-1 and 2, only UmV-KP4 (Group-1) and Zt-KP4-1

(Group-2) were tested for their toxicity to fungi. It is tempting to speculate that members

of these two structural groups, including ECP2, are fungitoxic. This hypothesis has to

be investigated by testing the biological activity of a wide array of KP4 proteins.

Groups-3 and -4 included proteins that did not have HCE2 or KP4 domains, and mostly

(90%) no Pfam domain. Since none of these KP4 proteins have been studied, it will be

interesting to assess their biological activity.

Classification of the KP6 proteins according to their structural similarities using PCA

and DALI Z-scores (Fig. 5) revealed three structural super families not identified before.

Group-1 gathered proteins structurally related to AvrLm10A and AvrLm6 from *L.*

maculans, SIX5 from *F. oxysporum*, and 25 fungal proteins from the kcl04-C cluster,

but no proteins from *Zymoseptoria*. AvrLm6 and AvrLm10A were already shown to

have a structural fold similar to Zt-KP6-1 and UmV-KP6α (Talbi *et al.*, 2023; Loubet *et*

al., 2024). AvrLm10A and SIX5 physically interact with effectors generally encoded by

neighboring genes (Ma *et al.*, 2015; Petit-Houdenot *et al.*, 2019; Yu *et al.*, 2023; Talbi

et al., 2023), and they facilitate the symplastic transport of their partner protein in host

plant tissues (Ma *et al.*, 2015; Talbi *et al.*, 2023). The biological activity of Group-1 KP6

proteins towards fungi is unknown. Group-2 included proteins structurally related to
known effectors such as BAS4 from *P. oryzae* (Khang *et al.*, 2010) and *ECP28* from
P. fulvum (Mesarich *et al.*, 2018). BAS4 is expressed during the biotrophic stage of
rice infection by *P. oryzae*, and restricted to the biotrophic interface between the fungal
cell wall and the plant membrane of the infected rice cells (Khang *et al.*, 2010). This
structural group includes Zt-KP6-4, and 420 KP6 proteins from clusters kcl01-C, kcl02-
G, kcl04-A, kcl11, and UmV-KP6β. Group-3 contained proteins related to Zt-KP6-1 and
UmV-KP6α. It also includes Zt-KP6-2, Zt-KP6-6, and five proteins from cluster kcl02-
A. Since Group-2 and -3 include three proteins with antifungal activities (UmV-KP6β,
Zt-KP6-1, UmV-KP6α), one may predict that some proteins from these two groups will
be fungitoxic. This hypothesis needs to be confirmed by testing the biological activity
of a wide array of KP6 proteins on fungi.
596
597

MATERIELS AND METHODS

Protein expression and purification

The Zt-KP6 mature protein (residues 21-90) was expressed with the pET-SP vector,

produced and purified as previously described (de Guillen *et al.*, 2015). The Zt-KP4

protein was expressed with the pET derived plasmid pdbccdb-3C-His, produced and

purified as previously described (Lahfa *et al.*, 2024) (Supplementals File S1) .

Nuclear magnetic resonance spectroscopy

To generate isotopically labelled samples for NMR spectroscopy, we used $^{15}NH_4Cl$, as

the unique source of nitrogen. The samples containing 0,7 mM of purified protein in

10 % D_2O and 1 µM DSS as a reference were analyzed using an 800 MHz Avance

Bruker spectrometer and a TCI probe at 303 K. Spectra were recorded with Topspin

3.2 (Bruker) and referenced to DSS (4,4-dimethyl-4-silapentane-1-sulfonic acid) for the

^1H dimension and indirectly for the ^{15}N dimension. NMR data were processed using

Topspin 3.6 (Bruker) and analyzed with the in-house software Cindy 2.1 (Padilla,

www.cbs.cnrs.fr). Side chain assignments were carried out using 2D-NOESY, 2D-

TOCSY and COSY-DQF experiments with D_2O samples, combined with ^{15}N-NOESY-

HSQC and ^{15}N-TOCSY-HSQC 3D spectra. The CYANA 3 program (Güntert, 2004)

was used for automatic NOE assignments and structure calculations. The CANDID

procedure of CYANA was used to assign the 3D-peaks list from the ^{15}N-NOESY-HSQC

spectra. Structures were analyzed with PROCHECK (Laskowski *et al.*, 1993).

Refinement statistics for the structure are listed in Sup. Table 1. The structure of Zt-

KP6 was deposited in the Protein Data Bank and in BMRB with accession number

9GWD and ID 34963 respectively.

X-ray crystallography

The purified protein Zt-KP6 was concentrated to 10 mg/ml in 20 mM Tris-HCl, pH 8.0,

150 mM NaCl, 1 mM DTT. The purified protein Zt-KP4 was concentrated to 8 mg/ml in

20 mM NaCi, pH 5.6, 150 mM NaCl, 1 mM DTT. Crystallization trials were performed

at 20°C by the hanging-drop vapor-diffusion method in 96-well microplates using a

Mosquito HTS robot with 100 nl of protein mixed with 100 nl of reservoir. After 3 weeks,

Zt-KP6 crystals were obtained from the condition 0.01 M $NiCl_2$, 0.1 M Tris pH 8.5, 1M

$LiSO_4$ (from classic kit, Hampton research®). After refinement and soaking in tantalum,

diffraction crystals were obtained in 0.1M Tris pH 9, 1.5M LiSO$_4$, 100mM ZnSO$_4$. After

2 weeks, small crystals of Zt-KP4 were obtained from one condition of the Peg kit

(Qiagen®) containing 0.1M sodium acetate pH 4.6 and 30%(v/v) PEG 300. All crystals

were flash frozen in liquid nitrogen.

Data collection, processing and structure determination

Crystal diffraction datasets were collected at the European Synchrotron Radiation

Facility (ESRF, Grenoble). For Zt-KP6, data were collected at beam line ID30A-1

(Svensson *et al.*, 2015, 2018; Bowler *et al.*, 2015, 2016; Nurizzo *et al.*, 2016) using a

Pixel detector (Pilatus3-2M) and processed by XDS (Kabsch, 2010) and AIMLESS

(Evans, 2011), from the CCP4/CCP4i programs suite (Potterton *et al.*, 2003; Winn *et*

al., 2011). For Zt-KP4, data were collected at beam line ID30B (Mueller-Dieckmann *et*

al., 2015; McCarthy *et al.*, 2018) using a Pixel detector (PILATUS3 6M) and auto

processed by XDSAPP (Krug *et al.*, 2012; Sparta *et al.*, 2016). Zt-KP6 crystals belong

to the $P2_12_12_1$ space group and contain two molecules in the asymmetric unit. The

structure of Zt-KP6 was determined to 1.36 Å by the single wavelength anomalous

diffraction (SAD) method using HKL2MAP (Pape & Schneider, 2004; Sheldrick, 2010).

Zt-KP4 crystals belong to the H3 space group and contain eight molecules in the

asymmetric unit. The structure of Zt-KP4 was determined by molecular replacement

with Molrep (Vagin & Teplyakov, 1997) from CCP4I2 (Winn *et al.*, 2011) using the

model determined by AF2 (Jumper *et al.*, 2021). After modelbuilding using arp/warp

(Langer *et al.*, 2008), Coot (Emsley *et al.*, 2010) and refinement by REFINE from

PHENIX (Adams *et al.*, 2010; Afonine *et al.*, 2012) final structures had R(%) / R(%)free

of 14 / 18.3 for Zt-KP6 and 17.5 / 21.9 for Zt-KP4. Data collection and

refinement statistics for the structures are listed in Sup. Table 2. Figures were

generated using PyMol v1.6 (http://www.pymol.org/). The atomic coordinates and

structure factors of Zt-KP6 and Zt-KP4, were deposited in the Protein Data Bank with

accession numbers 6QPK and 8ACX respectively.

Biological assays of Zt-KP4-1 and Zt-KP6-1

Zt-KP4-1 and Zt-KP6-1 were infiltrated into leaves of the wheat cultivar Taichung 29 at

a concentration of 2 µM. As a control, the necrotrophic effector ToxA from *P. nodorum*

(Tan *et al.*, 2012) was infiltrated into Taichung 29 wheat leaves at concentrations from

0.2 to 2 µM. *Z. tritici* IPO323 isolate and the necrotrophic plant pathogen *Botrytis*

cinerea were used for growth assays in 96 well microtiter plates containing liquid potato dextrose growth medium (PDA). Each well of the plate was filled with 100µL of PDA containing either 10E6 *Z. tritici* spores/mL or 10E6 *B. cinerea* spores/mL and different concentrations of Zt-KP4-1 or Zt-KP6-1 (0.55, 1.65 and 5.5 µM). 96 well plates were incubated in the dark at either 18°C (*Z. tritici*) or 25°C (*B. cinerea*) with shaking (50 rpm). After 7 days of growth, the OD at 580 nm was recorded on a spectrophotometer. The % of inhibition was calculated according to the formula: % inhibition = ((OD treated/OD control)-1)*100*. Values displayed in Table 1 were the mean ± Standard Error Deviation (SED) from three independent experiments.

FHAT protocol for the detection of fungal KP4 and KP6 proteins

Fungal KP4 and KP6 proteins were retrieved from public databases according to the following protocol (Sup Figure 10) and result files for each step are available at URLs https://pat.cbs.cnrs.fr/kp4/fhat_kp4 and https://pat.cbs.cnrs.fr/kp6/fhat_kp6. The protocol for the detection of fungal KP4 and KP6 proteins is fully described in the legend of Sup Figure 10.

PCA analysis

To classify KP4 and KP6 proteins according to their structural similarities, an "all against all" analysis was performed for both families using the DALI server (Holm, 2020). This computing delivered matrices of all the Z-scores for each pair of proteins. If a pair of proteins had a Z-score of 0.1, they were excluded from the following step. A Principal Component Analysis (PCA) was performed using these Z-scores matrices using *Statistic Kingdom* (https://www.statskingdom.com/410multi_linear_regression.html). Protein names were directly labeled on the graphs.

Acknowledgements

The CBS is a member of the France-BioImaging (FBI) and the French Infrastructure for Integrated Structural Biology (FRISBI), 2 national infrastructures supported by the French National Research Agency (ANR-10-INBS-0004 and ANR-10-INBS-0005, respectively). We thank R. Lambert for assistance with protein production and x-ray crystallography. We are grateful to Matthew Bowler and Didier Nurizzo at the European Synchrotron Radiation Facility (ESRF), Grenoble, France for providing assistance in

700   using beamline ID30B and ID30A-1 and the ESRF for provision of synchrotron
701   radiation facilities via Block Allocation Group beamtime.
702
703   **Data availability**
704
705   The data that support the findings of this study are openly available at the following
706   URLs:
707   https://pat.cbs.cnrs.fr/kp4/fhat_kp4
708   https://pat.cbs.cnrs.fr/kp6/fhat_kp6
709
710   **Authors contributions**
711   KDG, AP, TK and MHL planned and designed the research. KDG, LM, JG, PB, FH,
712   ML, JR, YPH and MHL performed experiments and analysed data. KDG, AP, TK and
713   MHL wrote the manuscript.
714

the supported node. Tree root with Ecp2_Passalora fulva and
Ecp2_Pseudocercospora fijiensis as outgroups. Zt-KP4-1_Ztritici_IPO323: re-
annotated NIP1/Mycgr3-106176 see Sup. Fig1; Zt-KP4-1_Ztritici_1E4: Uniprot
A0A2H1H404; Zt-KP4-1_Ztritici_3D7: Uniprot A0A1X7S6U3; Zt-KP4-1_Ztritici_1A5:
Uniprot A0A1Y6M012; Zt-KP4-2_Ztritici_IPO323: Mycgr3-107904, Uniprot F9X223;
Zb-KP4-2_Zbrevis: KJY02238, Uniprot A0A0F4GYX0; Zt-KP4-3_Ztritici_IPO323:
Mycgr3-111636, Uniprot F9XQ02; Zt-KP4-3_Ztritici_3D7: Uniprot A0A1X7S9R6; Zb-
KP4-3_Zbrevis: KJX96086, Uniprot A0A0F4GF74; Ecp2_Passalora fulva: Uniprot
Q00365; Ecp2_Pseudocercospora fijiensis: Uniprot M2Z6V3; Ecp2_Sphaerulina
musiva: Uniprot N1QK49; Ecp2_Cochliobolus sativus: Uniprot M2TE34
759
Figure 5. Classification of fungal KP6 families according to their structures
DALI all by all analysis followed by a PCA using a DALI score matrix (Holm, 2020).
meta-cluster kclXX, cluster AFDB50 kclXX-Y, (number of proteins in AFDB50 cluster),
Uniprot representative (example A0A2H1G421): kcl01-C, (100), A0A4V3HRP1,
M1W1H7; kcl01-L, (65), SIX5 = A0A010RIE2, AvrLm10A = E5A720_LEPMJ; kcl02-A,
(5), A0A4R8TM13; kcl02-D, (138), Zt-KP6-4 = A0A2H1GZA7; kcl02-G, (321),
A0A135SXI2; BAS4 = BAS4_MAGO7, Lm-KP6 = E5A0W1_LEPMJ; kcl04-A, (1),
F9FAC2; kcl04-C, (25), AvrLm6 = A0A010QJV2; AvrLm6 = E5R515_LEPMJ; kcl06-A,
(6), Zb-KP6-2 = A0A0F4GMG5, Zt-KP6-2 = A0A2H1H0K4; kcl06-B, (3), Zt-KP6-1 =
A0A2H1G421/6QKP; kcl07-A, (5), Zt-KP6-5 = A0A2H1FMK1, Zb-KP6-6 =
A0A0F4GGQ1; kcl11-A, (6), A0A317AKI4; kcl14-A, (93), ECP28-2 = A0A1P8YXM1,
ECP28-3 = A0A1P8YXM5; kcl14-B, (14), ECP28-1 = A0A1P8YXQ5; LmXXXX,
Leptosphaeria KP6 proteins, (8 protein, Loubet et al. 2024); Vi-KP6-famXX, *Venturia*
KP6 proteins, (4 families, Rocafort et al. 2022); White circle, UmV-KP6α, 4gvbA, White
circle, UmV-KP6β, 4gvbB
775
Figure 6. Classification of fungal KP4 families according to their structures
DALI all by all analysis followed by a PCA using a DALI score matrix (Holm, 2020).
meta-cluster kclXX, cluster AFDB50 kclXX-Y, (number of proteins in AFDB50 cluster),
Uniprot representative (example A0A2H1H404): kcl01-D, (83), Zt-KP4-1 =
A0A2H1H404/8ACX; kcl01-Q, (203), Sm-ECP2 = N1QK49, Psf-ECP2 = M2Z6V3, Paf-
ECP2 = Q00365; kcl02-A, (15), A0A4U0XEI9; kcl02-E, (256), Fg-KP4-2 =
A0A3S7E9S6, Fg-KP4-3 = A0A2H3GQS6; kcl02-I, (125), Fg-KP4-1 = A0A2H3GK41;

klc02-0, (324), Zt-KP4-9 = A0A2H1GXX1; kcl05-A, (14), A0A6A6DAE0; kcl07-A, (53),

A0A177DVW3; kcl07-C, (35), Zt-KP4-2 = A0A2H1FWY7; Zt-KP4-3 = A0A1Y6M296;

kcl08-A, (5), S3D5P; kcl09-A, (14), M3BT27; kcl09-B, (131), Zt-KP4-6 =

A0A2H1GNH3, Zt-KP4-7 = A0A2H1GND9; kcl10-A, (16), A0A1L9SH47; kcl12-A, (9),

E3L0Y8; kcl14-C, (12), A0A0A1V5R3; kcl18-A, (9), Zt-KP4-5 = A0A2H1GP20; kcl21-

A, (13), A0A0J9W663; kcl27-A, (39), A0A4T0WCM2; kcl38-A, (8), A0A017SS08;

White circle, UmV-KP4 = kptA.

Figure 7. Venn Diagram comparing KP4 and KP6 fungal proteins identified by

different pipelines

KP4 and KP6 fungal proteins identified by recent systematic structural studies on

fungal effectors (Seong & Krasileva, 2023; Derbyshire & Raffaele, 2023) were

compared to our dataset (FHAT). Zt-KP6-1 to 5 and Zt-KP4-1 to 5 were not detected

by Krasileva (2023) and Raffaele (2023).

Table1. Toxicity of Zt-KP4-1 and Zt-KP6-1 to fungi

The percentage of growth inhibition of *B. cinerea* and *Z. tritici* were plotted for 3

different concentrations of ToxA from *Phaeosphaeria nodorum*, Zt-KP4-1 and Zt-KP6-

1. Values are the mean ± standard deviation from three experiments.

List of supplementary figures and tables:

Supplementary Figure 1. Annotation of IPO323 *Z. tritici* Zt-Mycgr3-91409 (Zt-KP6-
1) and Zt-Mycgr3-106176 (Zt-KP4-1)

Sup. Fig.1-A Annotation of IPO323 *Z. tritici* CDS Zt-Mycgr3-91409 (Zt-KP6-1)

Sup. Fig.1-B Annotation of IPO323 *Z. tritici* CDS Zt-Mycgr3-106176 (Zt-KP4-1)

Supplementary Figure 2 Purification of Zt-KP6-1 by gel filtration
Gel filtration profile and SDS-PAGE analysis of purified Zt-KP6-1. Sizes of the
molecular mass markers are indicated on the left of the gel.

Supplementary Figure 3 NMR analysis of Zt-KP6-1
HSQC spectra of Zt-KP6-1 at, 0.7 mM in 20 mM NaCitrate pH 5.4, 150 mM NaCl, 1
mM DTT, 303 K, 800 MHz. Cross-peak assignments are indicated using one-letter
amino acid and number (the asterisk shows a folded red peak). Missing residue N36.

Supplementary Figure 4 Topology diagrams of Zt-KP6-1 and UmV-KP6α
The structures were processed by PDBsum and the secondary structures were colored
from N-terminus (blue) to C-terminus (red) according to Figure 1. Disulfide Bonds (D)
are represented as gray lines.

Supplementary Figure 5 Purification of Zt-KP4-1 by gel filtration
Gel filtration profile and SDS-PAGE analysis of purified Zt-KP4-1. (a) dimer, (b)
monomer. Sizes of the molecular mass markers are indicated on the left of the gel.

Supplementary Figure 6 Dimer of Zt-KP4-1 between chains B (cyan) and G (blue)
Residues involved in hydrogen bonds at the dimer interface are labelled and
represented by sticks. Hydrophobics residues involved in the pocket are highlighted in
lines and coloured in yellow. According to the GenBank sequence SMR60557.1, the
residue numbering must start after the signal peptide at residue S20. T33 in the
structure corresponds to T48 in the GenBank sequence.

Supplementary Figure 7 Topology diagrams of Zt-KP4 and UmV-Kp4

The structures were processed by PDBsum and the secondary structure were colored

from N-terminus (blue) to C-terminus (red) according to the Figure 2. Disulfide Bonds

(D) are represented as gray lines.

Supplementary Figure 8 Structure and topology of Zt-KP6-1 and Zt-KP6-2

Zt-KP6-2 structure was predicted by AlphaFold-2. The structures were processed by

PDBsum and the secondary structure were coloured from N-terminus (blue) to C-

terminus (red) according to Figure 1.

Supplementary Figure 9 Structure and topology of Zt-KP4-1, Zt-KP4-2 and Zt-

KP4-3

Zt-KP4-1, Zt-KP4-2 and Zt-KP4-3 structures were predicted by AlphaFold-2. The

structures were processed by PDBsum and the secondary structure were coloured

from N-terminus (blue) to C-terminus (red) according to the Figure 2.

Supplementary Figure 10 Flow chart of the pipeline used to identify fungal

proteins structurally related to Zt-KP6-1 and Zt-KP4-1

Supplementary Figure 11 Expression of Zt-KP4-1, Zt-KP4-2, Zt-KP4-3, Zt-KP6-1,

Zt-KP6-2, Zt-KP6-4 and Zt-KP6-6 during infection of wheat leaves infection.

RNAseq data from Rudd et al. (2015). RPKM (fungal RNAseq Reads Per Kilobase

Million)

Supplementary Table 1. NMR and refinement statistics for Zt-KP6-1

(Zt-KP6-1, 0.7 mM, 20 mM NaCitrate pH 5.4, 150 mM NaCl, 1 mM DTT, 303 K, 800

MHz) PDB:9GWD BMRB ID 34963

Supplementary Table 2. X-ray data collection and refinement statistics for Zt-

KP6-1 and Zt-KP4-1

#Rmerge = Shkl Si |Ii(hkl) - (I(hkl))| / Shkl Si Ii(hkl), where *Ii(hkl)* is the *i*th observation

of reflection *hkl* and *<I(hkl)>* is the weighted average intensity for all observations of

reflection *hkl*.

Supplementary Table 3 List of the proteins structurally related to Zt-KP6-1

detected using our pipeline

Supplementary Table 4 List of the proteins structurally related to Zt-KP4-1

detected using our pipeline

Supplementary Table 5 Prediction of the antimicrobial activity of fungal KP4 and

KP6 proteins using AMAPEC (Mesny & Thomma, 2024)

Supplementary file S1 Nucleotide sequence of synthetic genes and plasmids

used in this study

References

Adams PD, Afonine PV, Bunkóczi G, Chen VB, Davis IW, Echols N, Headd JJ, Hung L-W, Kapral GJ, Grosse-Kunstleve RW, _et al._ 2010. *PHENIX* : a comprehensive Python-based system for macromolecular structure solution. *Acta Crystallographica Section D Biological Crystallography* **66**: 213–221.

Afonine PV, Grosse-Kunstleve RW, Echols N, Headd JJ, Moriarty NW, Mustyakimov M, Terwilliger TC, Urzhumtsev A, Zwart PH, Adams PD. 2012. Towards automated crystallographic structure refinement with *phenix.refine*. *Acta Crystallographica Section D Biological Crystallography* **68**: 352–367.

Allen A, Chatt E, Smith TJ. 2013a. The Atomic Structure of the Virally Encoded Antifungal Protein, KP6. *Journal of Molecular Biology* **425**: 609–621.

Allen A, Islamovic E, Kaur J, Gold S, Shah D, Smith TJ. 2013b. The virally encoded killer proteins from Ustilago maydis. *Fungal Biology Reviews* **26**: 166–173.

Allen A, Snyder AK, Preuss M, Nielsen EE, Shah DM, Smith TJ. 2008. Plant defensins and virally encoded fungal toxin KP4 inhibit plant root growth. *Planta* **227**: 331–339.

Badet T, Oggenfuss U, Abraham L, McDonald BA, Croll D. 2020. A 19-isolate reference-quality global pangenome for the fungal wheat pathogen Zymoseptoria tritici. *BMC biology* **18**: 12.

Battache M, Lebrun M-H, Sakai K, Soudière O, Cambon F, Langin T, Saintenac C. 2022. Blocked at the Stomatal Gate, a Key Step of Wheat Stb16q-Mediated Resistance to Zymoseptoria tritici. *Frontiers in Plant Science* **13**.

Bauer S, Yu D, Lawson AW, Saur IML, Frantzeskakis L, Kracher B, Logemann E, Chai J, Maekawa T, Schulze-Lefert P. 2021. The leucine-rich repeats in allelic barley MLA immune receptors define specificity towards sequence-unrelated powdery mildew avirulence effectors with a predicted common RNase-like fold (B Kobe, Ed.). *PLOS Pathogens* **17**: e1009223.

Ben M'Barek S, Cordewener JHG, Tabib Ghaffary SM, van der Lee TAJ, Liu Z, Mirzadi Gohari A, Mehrabi R, America AHP, Robert O, Friesen TL, _et al._ 2015. FPLC and liquid-chromatography mass spectrometry identify candidate necrosis-inducing proteins from culture filtrates of the fungal wheat pathogen Zymoseptoria tritici. *Fungal Genetics and Biology* **79**: 54–62.

Blondeau K, Blaise F, Graille M, Kale SD, Linglin J, Ollivier B, Labarde A, Lazar N, Daverdin G, Balesdent M, _et al._ 2015. Crystal structure of the effector A vr L m4–7 of *Leptosphaeria maculans* reveals insights into its translocation into plant cells and recognition by resistance proteins. *The Plant Journal* **83**: 610–624.

Bowler MW, Nurizzo D, Barrett R, Beteva A, Bodin M, Caserotto H, Delagenière S, Dobias F, Flot D, Giraud T, _et al._ 2015. MASSIF-1: a beamline dedicated to the fully automatic characterization and data collection from crystals of biological macromolecules. *Journal of Synchrotron Radiation* **22**: 1540–1547.

Bowler MW, Svensson O, Nurizzo D. 2016. Fully automatic macromolecular crystallography: the impact of MASSIF-1 on the optimum acquisition and quality of data. *Crystallography Reviews* **22**: 233–249.

Brown DW. 2011. The KP4 killer protein gene family. *Current Genetics* **57**: 51–62.

Cao Y, Kümmel F, Logemann E, Gebauer JM, Lawson AW, Yu D, Uthoff M, Keller B, Jirschitzka J, Baumann U, _et al._ 2023. Structural polymorphisms within a common powdery mildew effector scaffold as a driver of coevolution with cereal immune receptors. *Proceedings of the National Academy of Sciences of the United States of America* **120**: e2307604120.

Carreón-Anguiano KG, Gómez-Tah R, Pech-Balan E, Ek-Hernández GE, De los Santos-Briones C, Islas-Flores I, Canto-Canché B. 2023. Pseudocercospora fijiensis Conidial

Germination Is Dominated by Pathogenicity Factors and Effectors. *Journal of Fungi* **9**.

Chang T-C, Salvucci A, Crous PW, Stergiopoulos I. **2016**. Comparative Genomics of the Sigatoka Disease Complex on Banana Suggests a Link between Parallel Evolutionary Changes in Pseudocercospora fijiensis and Pseudocercospora eumusae and Increased Virulence on the Banana Host. *PLoS genetics* **12**: e1005904.

Clausen M, Kräuter R, Schachermayr G, Potrykus I, Sautter C. **2000**. Antifungal activity of a virally encoded gene in transgenic wheat. *Nature biotechnology* **18**: 446–449.

Deb D, Shrestha A, Maiti IB, Dey N. **2018**. Recombinant Promoter (MUASCsV8CP) Driven Totiviral Killer Protein 4 (KP4) Imparts Resistance Against Fungal Pathogens in Transgenic Tobacco. *Frontiers in Plant Science* **9**: 278.

Derbyshire MC, Raffaele S. **2023**. *Surface frustration re-patterning underlies the structural landscape and evolvability of fungal orphan candidate effectors*. Genomics.

Ebert MK, Rangel LI, Spanner RE, Taliadoros D, Wang X, Friesen TL, de Jonge R, Neubauer JD, Secor GA, Thomma BPHJ, *et al.* 2021. Identification and characterization of Cercospora beticola necrosis-inducing effector CbNip1. *Molecular Plant Pathology* **22**: 301–316.

Emsley P, Lohkamp B, Scott WG, Cowtan K. **2010**. Features and development of *Coot*. *Acta Crystallographica Section D Biological Crystallography* **66**: 486–501.

Evans PR. **2011**. An introduction to data reduction: space-group determination, scaling and intensity statistics. *Acta Crystallographica Section D Biological Crystallography* **67**: 282–292.

Finkler A, Peery T, Tao J, Bruenn J, Koltin igal. **1992**. Immunity and resistance to the KP6 toxin of Ustilago maydis. *Molecular and General Genetics MGG* **233**: 395–403.

Gage MJ, Bruenn J, Fischer M, Sanders D, Smith TJ. **2001**. KP4 fungal toxin inhibits growth in Ustilago maydis by blocking calcium uptake: KP4 fungal toxin mechanism. *Molecular Microbiology* **41**: 775–785.

Gage MJ, Rane SG, Hockerman GH, Smith TJ. **2002**. The Virally Encoded Fungal Toxin KP4 Specifically Blocks L-Type Voltage-Gated Calcium Channels. *Molecular Pharmacology* **61**: 936–944.

Gu F, Khimani A, Rane SG, Flurkey WH, Bozarth RF, Smith TJ. **1995**. Structure and function of a virally encoded fungal toxin from Ustilago maydis: a fungal and mammalian Ca2+ channel inhibitor. *Structure* **3**: 805–814.

Guan Y, Ma L, Wang Q, Zhao J, Wang S, Wu J, Liu Y, Sun H, Huang J. **2023**. Horizontally acquired fungal killer protein genes affect cell development in mosses. *The Plant Journal* **113**: 665–676.

de Guillen K, Ortiz-Vallejo D, Gracy J, Fournier E, Kroj T, Padilla A. **2015**. Structure Analysis Uncovers a Highly Diverse but Structurally Conserved Effector Family in Phytopathogenic Fungi. *PLOS Pathog* **11**: e1005228.

Güntert P. **2004**. Automated NMR structure calculation with CYANA. *Methods in molecular biology (Clifton, N.J.)* **278**: 353–378.

Holm L. **2020**. DALI and the persistence of protein shape. *Protein Science* **29**: 128–140.

de Jonge R, van Esse HP, Maruthachalam K, Bolton MD, Santhanam P, Saber MK, Zhang Z, Usami T, Lievens B, Subbarao KV, *et al.* 2012. Tomato immune receptor Ve1 recognizes effector of multiple fungal pathogens uncovered by genome and RNA sequencing. *Proceedings of the National Academy of Sciences of the United States of America* **109**: 5110–5115.

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek A, Potapenko A, *et al.* 2021. Highly accurate protein structure prediction with AlphaFold. *Nature* **596**: 583–589.

Kabsch W. **2010**. Integration, scaling, space-group assignment and post-refinement. *Acta*

Crystallographica Section D Biological Crystallography **66**: 133–144.

Kettles GJ, Bayon C, Sparks CA, Canning G, Kanyuka K, Rudd JJ. **2018**. Characterization of an antimicrobial and phytotoxic ribonuclease secreted by the fungal wheat pathogen Zymoseptoria tritici. *New Phytologist* **217**: 320–331.

Khang CH, Berruyer R, Giraldo MC, Kankanala P, Park S-Y, Czymmek K, Kang S, Valent B. **2010**. Translocation of Magnaporthe oryzae Effectors into Rice Cells and Their Subsequent Cell-to-Cell Movement. *The Plant Cell* **22**: 1388–1403.

Kilaru S, Fantozzi E, Cannon S, Schuster M, Chaloner TM, Guiu-Aragones C, Gurr SJ, Steinberg G. **2022**. Zymoseptoria tritici white-collar complex integrates light, temperature and plant cues to initiate dimorphism and pathogenesis. *Nature Communications* **13**: 5625.

Koltin Y. **1988**. The killer system of Ustilago maydis: secreted polypeptides encoded by viruses. *Viruses of fungi and simple eukaryotes*: 209–242.

Krug M, Weiss MS, Heinemann U, Mueller U. **2012**. *XDSAPP* : a graphical user interface for the convenient processing of diffraction data using *XDS*. *Journal of Applied Crystallography* **45**: 568–572.

Lahfa M, Barthe P, De Guillen K, Cesari S, Raji M, Kroj T, Le Naour—Vernet M, Hoh F, Gladieux P, Roumestand C, *et al.* 2024. The structural landscape and diversity of Pyricularia oryzae MAX effectors revisited (F Mesny, Ed.). *PLOS Pathogens* **20**: e1012176.

Langer G, Cohen SX, Lamzin VS, Perrakis A. **2008**. Automated macromolecular model building for X-ray crystallography using ARP/wARP version 7. *Nature Protocols* **3**: 1171–1179.

Lapalu N, Simon A, Demenou B, Paumier D, Guillot M-P, Gout L, Suffert F, Valade R. **2023**. Complete Genome Sequences of Septoria linicola: A Resource for Studying a Damaging Flax Pathogen. *Molecular plant-microbe interactions : MPMI* **36**: 59–63.

Laskowski RA, MacArthur MW, Moss DS, Thornton JM. **1993**. PROCHECK: a program to check the stereochemical quality of protein structures. *Journal of Applied Crystallography* **26**: 283–291.

Lazar N, Mesarich CH, Petit-Houdenot Y, Talbi N, Li De La Sierra-Gallay I, Zélie E, Blondeau K, Gracy J, Ollivier B, Blaise F, *et al.* 2022. A new family of structurally conserved fungal effectors displays epistatic interactions with plant resistance proteins (B Kobe, Ed.). *PLOS Pathogens* **18**: e1010664.

Lazar N, Mesarich CH, Petit-Houdenot Y, Talbi N, de la Sierra-Gallay IL, Zélie E, Blondeau K, Gracy J, Ollivier B, Blaise F, *et al.* 2020. *A new family of structurally conserved fungal effectors displays epistatic interactions with plant resistance proteins*. Plant Biology.

Le Naour—Vernet M, Charriat F, Gracy J, Cros-Arteil S, Ravel S, Veillet F, Meusnier I, Padilla A, Kroj T, Cesari S, *et al.* 2023. Adaptive evolution in virulence effectors of the rice blast fungus Pyricularia oryzae (MF Seidl, Ed.). *PLOS Pathogens* **19**: e1011294.

Lemoine F, Correia D, Lefort V, Doppelt-Azeroual O, Mareuil F, Cohen-Boulakia S, Gascuel O. **2019**. NGPhylogeny.fr: new generation phylogenetic services for non-specialists. *Nucleic acids research* **47**: W260–W265.

Li N, Erman M, Pangborn W, Duax WL, Park C-M, Bruenn J, Ghosh D. **1999**. Structure of Ustilago maydis Killer Toxin KP6 α-Subunit. *Journal of Biological Chemistry* **274**: 20425–20431.

Li G, Newman M, Yu H, Rashidzade M, Martínez-Soto D, Caicedo A, Allen KS, Ma L-J. **2024**. Fungal effectors: past, present, and future. *Current Opinion in Microbiology* **81**: 102526.

Lima AM, Azevedo MIG, Sousa LM, Oliveira NS, Andrade CR, Freitas CDT, Souza PFN. **2022**. Plant antimicrobial peptides: An overview about classification, toxicity and clinical applications. *International journal of biological macromolecules* **214**: 10–21.

Lo Presti L, Lanver D, Schweizer G, Tanaka S, Liang L, Tollot M, Zuccaro A,
Reissmann S, Kahmann R. **2015**. Fungal Effectors and Plant Susceptibility. *Annual Review*
of Plant Biology **66**: 513–545.

Loubet C, Li de la Sierra-Gallay I, van Tilbeurgh H, Fudal I. **2024**. Decoding plant
pathogen strategies: exploring Leptosphaeria maculans effector structural insights for
enhanced plant resistance. *Submitted*.

Lu S, Faris JD. **2019**. Fusarium graminearum KP4-like proteins possess root growth-
inhibiting activity against wheat and potentially contribute to fungal virulence in seedling rot.
Fungal Genetics and Biology **123**: 1–13.

Ma L, Houterman PM, Gawehns F, Cao L, Sillo F, Richter H, Clavijo-Ortiz MJ,
Schmidt SM, Boeren S, Vervoort J, et al. **2015**. The AVR2–SIX5 gene pair is required to
activate I-2-mediated immunity in tomato. *New Phytologist* **208**: 507–518.

McCarthy AA, Barrett R, Beteva A, Caserotto H, Dobias F, Felisaz F, Giraud T,
Guijarro M, Janocha R, Khadrouche A, et al. **2018**. ID30B – a versatile beamline for
macromolecular crystallography experiments at the ESRF. *Journal of Synchrotron Radiation*
25: 1249–1260.

Mesarich CH, Ökmen B, Rovenich H, Griffiths SA, Wang C, Karimi Jashni M,
Mihajlovski A, Collemare J, Hunziker L, Deng CH, et al. **2018**. Specific Hypersensitive
Response–Associated Recognition of New Apoplastic Effectors from *Cladosporium fulvum* in
Wild Tomato. *Molecular Plant-Microbe Interactions* **31**: 145–162.

Mesny F, Thomma BP. **2024**. *AMAPEC: accurate antimicrobial activity prediction for*
fungal effector proteins. Microbiology.

Mueller-Dieckmann C, Bowler MW, Carpentier P, Flot D, McCarthy AA, Nanao MH,
Nurizzo D, Pernot P, Popov A, Round A, et al. **2015**. The status of the macromolecular
crystallography beamlines at the European Synchrotron Radiation Facility. *The European*
Physical Journal Plus **130**: 70.

Nurizzo D, Bowler MW, Caserotto H, Dobias F, Giraud T, Surr J, Guichard N, Papp G,
Guijarro M, Mueller-Dieckmann C, et al. **2016**. RoboDiff: combining a sample changer
and goniometer for highly automated macromolecular crystallography experiments. *Acta*
Crystallographica Section D Structural Biology **72**: 966–975.

Outram MA, Figueroa M, Sperschneider J, Williams SJ, Dodds PN. **2022**. Seeing is
believing: Exploiting advances in structural biology to understand and engineer plant
immunity. *Current Opinion in Plant Biology*: 8.

Pape T, Schneider TR. **2004**. ıt HKL2MAP: a graphical user interface for macromolecular
phasing with ıt SHELX programs. *Journal of Applied Crystallography* **37**: 843–844.

Park C-M, Bruenn JA, Ganesa C, Flurkey WF, Bozarth RF, Koltin Y. **1994**. Structure
and heterologous expression of the Ustilago maydis viral toxin KP4. *Molecular Microbiology*
11: 155–164.

Pennington HG, Jones R, Kwon S, Bonciani G, Thieron H, Chandler T, Luong P,
Morgan SN, Przydacz M, Bozkurt T, et al. **2019**. The fungal ribonuclease-like effector
protein CSEP0064/BEC1054 represses plant immunity and interferes with degradation of host
ribosomal RNA. *PLOS Pathogens* **15**: e1007620.

Petit-Houdenot Y, Degrave A, Meyer M, Blaise F, Ollivier B, Marais C-L, Jauneau A,
Audran C, Rivas S, Veneault-Fourrey C, et al. **2019**. A two genes – for – one gene
interaction between Leptosphaeria maculans and Brassica napus. *New Phytologist* **223**: 397–
411.

Potterton E, Briggs P, Turkenburg M, Dodson E. **2003**. A graphical user interface to the
CCP4 program suite. *Acta crystallographica. Section D, Biological crystallography* **59**:
1131–1137.

Quijano CD, Wichmann F, Schlaich T, Fammartino A, Huckauf J, Schmidt K, Unger C,

Broer I, Sautter C. **2016**. KP4 to control Ustilago tritici in wheat: Enhanced greenhouse resistance to loose smut and changes in transcript abundance of pathogen related genes in infected KP4 plants. *Biotechnology Reports* **11**: 90–98.

Rocafort M, Bowen JK, Hassing B, Cox MP, McGreal B, de la Rosa S, Plummer KM, Bradshaw RE, Mesarich CH. **2022**. The Venturia inaequalis effector repertoire is dominated by expanded families with predicted structural similarity, but unrelated sequence, to avirulence proteins from other plant-pathogenic fungi. *BMC Biology* **20**: 246.

Rudd JJ, Kanyuka K, Hassani-Pak K, Derbyshire M, Andongabo A, Devonshire J, Lysenko A, Saqi M, Desai NM, Powers SJ, *et al.* 2015. Transcriptome and metabolite profiling of the infection cycle of Zymoseptoria tritici on wheat reveals a biphasic interaction with plant immunity involving differential pathogen chromosomal contributions and a variation on the hemibiotrophic lifestyle definition. *Plant physiology* **167**: 1158–1185.

Santos A, Navascués E, Bravo E, Marquina D. **2011**. Ustilago maydis killer toxin as a new tool for the biocontrol of the wine spoilage yeast Brettanomyces bruxellensis. *International journal of food microbiology* **145**: 147–154.

Schlaich T, Urbaniak BM, Malgras N, Ehler E, Birrer C, Meier L, Sautter C. **2006**. Increased field resistance to Tilletia caries provided by a specific antifungal virus gene in genetically engineered wheat. *Plant biotechnology journal* **4**: 63–75.

Seong K, Krasileva KV. **2023**. Prediction of effector protein structures from fungal phytopathogens enables evolutionary analyses. *Nature Microbiology* **8**: 174–187.

Sheldrick GM. **2010**. Experimental phasing with *SHELXC / D / E* : combining chain tracing with density modification. *Acta Crystallographica Section D Biological Crystallography* **66**: 479–485.

Snelders NC, Boshoven JC, Song Y, Schmitz N, Fiorin GL, Rovenich H, van den Berg GCM, Torres DE, Petti GC, Prockl Z, *et al.* 2023. A highly polymorphic effector protein promotes fungal virulence through suppression of plant-associated Actinobacteria. *The New phytologist* **237**: 944–958.

Snelders NC, Petti GC, van den Berg GCM, Seidl MF, Thomma BPHJ. **2021**. An ancient antimicrobial protein co-opted by a fungal plant pathogen for in planta mycobiome manipulation. *Proceedings of the National Academy of Sciences of the United States of America* **118**.

Snelders NC, Rovenich H, Petti GC, Rocafort M, van den Berg GCM, Vorholt JA, Mesters JR, Seidl MF, Nijland R, Thomma BPHJ. **2020**. Microbiome manipulation by a soil-borne fungal plant pathogen using effector proteins. *Nature plants* **6**: 1365–1374.

Snelders NC, Rovenich H, Thomma BPHJ. **2022**. Microbiota manipulation through the secretion of effector proteins is fundamental to the wealth of lifestyles in the fungal kingdom. *FEMS microbiology reviews* **46**.

Spanu PD. **2017**. Cereal immunity against powdery mildews targets RNase-Like Proteins associated with Haustoria (RALPH) effectors evolved from a common ancestral gene. *New Phytologist* **213**: 969–971.

Sparta KM, Krug M, Heinemann U, Mueller U, Weiss MS. **2016**. *XDSAPP2.0. Journal of Applied Crystallography* **49**: 1085–1092.

Stergiopoulos I, Kourmpetis YAI, Slot JC, Bakker FT, De Wit PJGM, Rokas A. **2012**. In Silico Characterization and Molecular Evolutionary Analysis of a Novel Superfamily of Fungal Effector Proteins. *Molecular Biology and Evolution* **29**: 3371–3384.

Svensson O, Gilski M, Nurizzo D, Bowler MW. **2018**. Multi-position data collection and dynamic beam sizing: recent improvements to the automatic data-collection algorithms on MASSIF-1. *Acta Crystallographica Section D Structural Biology* **74**: 433–440.

Svensson O, Malbet-Monaco S, Popov A, Nurizzo D, Bowler MW. **2015**. Fully automatic characterization and data collection from crystals of biological macromolecules. *Acta*

Crystallographica Section D Biological Crystallography **71**: 1757–1767.

Talbi N, Fokkens L, Audran C, Petit-Houdenot Y, Pouzet C, Blaise F, Gay EJ, Rouxel T, Balesdent M, Rep M, *et al.* 2023. The neighbouring genes *AvrLm10A* and *AvrLm10B* are part of a large multigene family of cooperating effector genes conserved in Dothideomycetes and Sordariomycetes. *Molecular Plant Pathology* **24**: 914–931.

Tan K-C, Ferguson-Hunt M, Rybak K, Waters ODC, Stanley WA, Bond CS, Stukenbrock EH, Friesen TL, Faris JD, McDonald BA, *et al.* 2012. Quantitative Variation in Effector Activity of ToxA Isoforms from Stagonospora nodorum and Pyrenophora tritici-repentis. *Molecular Plant-Microbe Interactions®* **25**: 515–522.

Tao J, Ginsberg I, Banerjee N, Held W, Koltin Y, BRUENNl JA. 1990. Ustilago maydis KP6 Killer Toxin: Structure, Expression in Saccharomyces cerevisiae, and Relationship to Other Cellular Toxins. *MOL. CELL. BIOL.* **10**: 9.

Teulet A, Quan C, Evangelisti E, Wanke A, Yang W, Schornack S. 2023. A pathogen effector FOLD diversified in symbiotic fungi. *New Phytologist*: nph.18996.

Vagin A, Teplyakov A. 1997. *MOLREP* : an Automated Program for Molecular Replacement. *Journal of Applied Crystallography* **30**: 1022–1025.

Van den Ackerveken GF, Van Kan JA, Joosten MH, Muisers JM, Verbakel HM, De Wit PJ. 1993. Characterization of two putative pathogenicity genes of the fungal tomato pathogen Cladosporium fulvum. *Molecular plant-microbe interactions : MPMI* **6**: 210–215.

Winn MD, Ballard CC, Cowtan KD, Dodson EJ, Emsley P, Evans PR, Keegan RM, Krissinel EB, Leslie AGW, McCoy A, *et al.* 2011. Overview of the *CCP* 4 suite and current developments. *Acta Crystallographica Section D Biological Crystallography* **67**: 235–242.

Yu DS, Outram MA, Smith A, McCombe CL, Khambalkar PB, Rima SA, Sun X, Ma L, Ericsson DJ, Jones DA, *et al.* 2023. *The structural repertoire of Fusarium oxysporum f. sp. lycopersici effectors revealed by experimental and computational studies.* elife.

Zhang Y, Skolnick J. 2005. TM-align: a protein structure alignment algorithm based on the TM-score. *Nucleic acids research* **33**: 2302–2309.