1                 Human Brain-Wide Activation of Sleep Rhythms

Haiteng Wang[1,2#], Qihong Zou[3,4#], Jinbo Zhang[1,2], Jia-Hong Gao[3,4,5*], Yunzhe Liu[1,2*]

1. State Key Laboratory of Cognitive Neuroscience and Learning, IDG/McGovern Institute for Brain

Research, Beijing Normal University, Beijing, China

2. Chinese Institute for Brain Research, Beijing, China

3. Center for MRI Research, Academy for Advanced Interdisciplinary Studies, Peking University,

Beijing, China

4. Beijing City Key Lab for Medical Physics and Engineering, Institute of Heavy Ion Physics, School

of Physics, Peking University, Beijing, China

5. McGovern Institute for Brain Research, Peking University, Beijing, China

# These authors contributed equally

* Correspondence: jgao@pku.edu.cn (J. G.),    yunzhe.liu@bnu.edu.cn (Y.L.)

**Abstract**

During sleep, our brain undergoes highly synchronized activity, orchestrated by distinct neural rhythms.

Little is known about the associated brain activation during these sleep rhythms, and even less about

their functional implications. In this study, we investigated the brain-wide activation underlying human

sleep rhythms by employing simultaneous electroencephalography (EEG) and functional magnetic

resonance imaging (fMRI) in 107 participants during nocturnal nap (first half of the night). We

identified robust coupling between slow oscillations (SOs) and fast spindles during deep non-rapid eye

movement (NREM) sleep (N2/3 stages), with spindle peaks consistently occurring just before the SO

UP-state. This SO-spindle coupling was linked to elevated activation in both the thalamus and

hippocampus, alongside increased functional connectivity from the hippocampus to the thalamus and

from the thalamus to the medial prefrontal cortex (mPFC). An open-ended cognitive state decoding

analysis suggested that these activations may relate to episodic memory processes, yet were distinct

from task-related networks. Together, these findings highlight the thalamus as a key coordinator of

hippocampal-cortical communication during sleep and provide new insights into the mechanisms by

which synchronized sleep rhythms may support memory consolidation.

**Keywords**: slow oscillation (SO), spindle, hippocampus, thalamus, simultaneous EEG-fMRI

**Introduction**

During sleep, the human brain engages in highly synchronized activity driven by distinct neural rhythms. In non-rapid eye movement (NREM) sleep, the emergence and coordination of slow oscillations (SOs) and sleep spindles are key features (Hahn et al., 2020; Helfrich et al., 2018; Ngo, Fell, & Staresina, 2020; Schreiner et al., 2022; Staresina et al., 2015; Staresina et al., 2023). However, the brain activation patterns associated with these rhythms, and especially their coordination, remain largely unexplored, and their functional implications are even less understood.

SOs (~1 Hz) are characterized by fluctuations in neuronal membrane potential between periods of silence ('hyperpolarization', or 'DOWN' state) and excitation ('depolarization', or 'UP' state) (Amzica & Steriade, 2002; Steriade, McCormick, & Sejnowski, 1993). These oscillations originate in neocortical regions, particularly the medial prefrontal cortex (mPFC), during NREM sleep (Dang-Vu et al., 2008; Massimini et al., 2004; Nir et al., 2011). The depolarization phase of SOs is believed to trigger sleep spindles in the thalamus, producing 11–16 Hz oscillations (Fernandez & Lüthi, 2020; Mak-McCully et al., 2017). Thalamic spindles tend to align with the excitable UP-states of SOs and can synchronize with hippocampal ripples (Buzsáki, 2015; Helfrich et al., 2019; Joo & Frank, 2018; Ngo, Fell, & Staresina, 2020). Rodent optogenetics studies have shown that only thalamic spindles phase-locked to the UP-state of cortical SOs enhance memory consolidation, while out-of-phase spindles do not (Latchoumane et al., 2017). This finding is consistent with intracranial evidence from human epilepsy patients, suggesting the thalamus acts as a key relay between the hippocampus and cortex, coordinating their interaction (Coulon, Budde, & Pape, 2012; Ferraris et al., 2021; Schreiner et al., 2022; Staresina et al., 2015).

The triple coupling of SOs, spindles, and ripples is believed to facilitate memory consolidation by synchronizing neuronal activity across brain regions and replaying memory traces from the hippocampus to the neocortex (Diekelmann & Born, 2010; Latchoumane et al., 2017; Liu et al., 2022; Singh, Norman, & Schapiro, 2022; Staresina et al., 2015; Staresina et al., 2023), possibly through relay from thalamic spindles. However, the exact mechanisms of this inter-regional communication during

coordinated sleep rhythms remain unclear. Understanding the brain-wide activation associated with these rhythms is crucial, as it reveals how information flows among brain regions, especially during the critical time windows when these rhythms align.

Investigating these sleep-related neural processes in humans is challenging because it requires tracking transient sleep rhythms while simultaneously assessing their widespread brain activation. Recent advances in simultaneous EEG-fMRI techniques provide a unique opportunity to explore these processes. EEG allows for precise event-based detection of neural signal, while fMRI provides insight into the broader spatial patterns of brain activation and functional connectivity (Horovitz et al., 2008; Huang et al., 2024; Laufs, 2008; Laufs, Walker, & Lund, 2007; Schabus et al., 2007; Spoormaker et al., 2010). Previous EEG-fMRI studies on sleep have focused on classifying sleep stages or examining the neural correlates of specific waves (Bergmann et al., 2012; Caporro et al., 2012; Czisch et al., 2009; Fogel et al., 2017; Hale et al., 2016; Ilhan-Bayrakcı et al., 2022; Moehlman et al., 2019; Picchioni et al., 2011). These studies have generally reported that slow oscillations are associated with widespread cortical and subcortical BOLD changes, whereas spindles elicit activation in the thalamus, as well as in several cortical and paralimbic regions. Although these findings provide valuable insights into the BOLD correlates of sleep rhythms, they often do not employ sophisticated temporal modeling (Huang et al., 2024), to capture the dynamic interactions between different oscillatory events, e.g., the coupling between SOs and spindles.

Understanding sleep rhythm-related brain activation is also essential for uncovering its potential cognitive functions. Since performing cognitive tasks during sleep is limited, most neuroimaging studies on human sleep rely on the targeted memory reactivation (TMR) paradigm, where participants are re-exposed to sensory cues during sleep to trigger associated memory reactivation (Cousins et al., 2016; Hu et al., 2020; Oudiette & Paller, 2013; Siefert et al., 2024). For instance, fMRI studies have shown significant hippocampal activation in response to odour re-exposure during sleep (Rasch et al., 2007), while EEG studies suggest that this reactivation is more likely during the UP-state of SOs (Cousins et al., 2016; Göldi et al., 2019; Xia et al., 2023). However, while TMR provides insight into

externally induced memory processing during sleep, it offers limited understanding of the endogenous neural mechanisms underlying natural sleep rhythms.

To address this gap, our study investigates brain-wide activation and functional connectivity patterns associated with SO-spindle coupling, and employs a cognitive state decoding approach (Margulies et al., 2016; Yarkoni et al., 2011)—albeit indirectly—to infer potential cognitive functions. In the current study, we used simultaneous EEG-fMRI recordings during nocturnal naps (detailed sleep staging results are provided in the Methods and **Table S1**) in 107 participants. Although directly detecting hippocampal ripples using scalp EEG or fMRI is challenging, we expected that hippocampal activation in fMRI would coincide with SO-spindle coupling detected by EEG, given that SOs, spindles, and ripples frequently co-occur during NREM sleep. We also anticipated a critical role of the thalamus, particularly thalamic spindles, in coordinating hippocampal-cortical communication.

We found significant coupling between SOs and spindles during NREM sleep (N2/3), with spindle peaks occurring slightly before the SO peak. This coupling was associated with increased activation in both the thalamus and hippocampus, with functional connectivity patterns suggesting thalamic coordination of hippocampal-cortical communication. These findings highlight the key role of the thalamus in coordinating hippocampal-cortical interactions during human sleep and provide new insights into the neural mechanisms underlying sleep-dependent brain communication. A deeper understanding of these mechanisms may contribute to future neuromodulation approaches aimed at enhancing sleep-dependent cognitive function and treating sleep-related disorders.

## Results

Simultaneous EEG-fMRI data during nocturnal sleep were collected from 138 healthy subjects (age, 22.56 ± 0.30; 81 females). After excluding subjects due to excessive head motion (mean framewise displacement > 0.5) or insufficient scanning sleep duration (<10 min), 107 participants (age, 22.40 ± 0.34; 63 females) were retained for subsequent analysis, with an average scanning time of sleep of 3.01 ± 0.14 hours (see Methods and **Table S1**). Since NREM sleep predominantly occurs in early sleep (Carskadon & Dement, 2005), we conducted our scans during the first half of the night to ensure sufficient coverage of NREM sleep, this is also the period when sleep rhythms are most consistently observed and studied in relation to memory consolidation (Maingret et al., 2016; Molle et al., 2011; Schreiner et al., 2021; Staresina et al., 2015; Staresina et al., 2023).

## Sleep stages and sleep rhythms

The primary objective of the EEG analysis was to examine the intrinsic characteristics of sleep rhythms and to identify the timing of these SO and spindle events for subsequent fMRI analyses.

We first removed MRI gradient noise from the EEG signal (detailed in Methods, see **Fig. S1**) and then applied an automated sleep staging algorithm (Vallat & Walker, 2021). The staging results were manually reviewed by two sleep experts to ensure the accurate classification of each sleep stages (**Fig. 1b**). Upon validation, we confirmed that N2/3 stages predominated in the dataset, with a significantly higher proportion of N2/3 sleep (75.77% ± 1.50%) compared to N1 (14.57% ± 1.06%, $t_{(106)}$ = 25.04, $p$ < 1e-4, paired-samples $t$ test) and REM sleep (9.66% ± 0.90%, $t_{(106)}$ = 29.60, $p$ < 1e-4, **Fig. 1d, Table S1**).

Each sleep stage is characterized by distinct spectral properties and rhythmic waveforms, serving as physiological markers (**Fig. 1c**). As a sanity check, we compared SO and spindle amplitudes and densities across stages. Both were significantly higher during N2/3 sleep (SO: 25.59 ± 1.49 μV, spindle: 7.39 ± 0.27 μV) compared to N1 (SO: 18.87 ± 1.08 μV, spindle: 6.72 ± 0.25 μV) and REM (SO: 16.67

± 1.14 µV, spindle: 6.23 ± 0.23 µV, all $p$ < 1e-4, **Fig. 1e, Fig. S2**). This aligns with the established understanding that N2/3 stages feature more pronounced slow-wave activity and robust spindles (Gaillard & Blois, 1981). Furthermore, we found that the amplitude of the SO DOWN-state (13.62 ± 0.67 µV) was significantly higher than the UP-state (12.98 ± 0.66 µV, $t_{(106)}$ = 4.82, $p$ < 1e-4), consistent with previous work (Cash et al., 2009; Dang-Vu et al., 2008). Based on this, we used the DOWN-state to lock the timing of SOs for subsequent EEG-informed fMRI analysis. Descriptive information on sleep rhythms during these stages for all 107 subjects can be found in **Table S2-S4**.

**Fig. 1 | Sleep stages and sleep rhythms in 107 subjects. a,** Sleep rhythms and task schematic. Subjects slept first half of the night with simultaneous EEG-fMRI recordings. Since detecting hippocampal ripples directly from scalp EEG is challenging, our focus was on capturing SOs, spindles, and their couplings. Regions of interest (ROIs) are colour-coded: green for the thalamus (spindle), purple for the mPFC (SOs), and orange for the hippocampus (ripples). **b,** Sleep staging and EEG spectrogram. N2/3 sleep stages (red line) were initially identified using an offline automatic sleep staging algorithm (Vallat & Walker, 2021) and then manually validated. (c) Schematic of EEG data across different sleep stages, using preprocessed data from the C3 electrode. (d) Proportion of each sleep stage in the dataset. (e) Amplitudes (µV) of SOs (left) and spindles (right) across sleep stages. The SO amplitudes used here are from 0.16-1.25 Hz band EEG data, and the spindle amplitudes used here are from 12-16 Hz band EEG data. Each dot represents an individual participant. Error bars indicate SEM. *** $p < 0.001$.

## SO-spindle coupling

SO-spindle coupling is considered important for sleep-dependent memory consolidation. In current study, we found that this coupling primarily occurs during N2/3 sleep stages ($2.46 \pm 0.06$ times/min), significantly more than in N1 ($2.23 \pm 0.09$ times/min, $t_{(106)} = 2.81$, $p = 0.0059$) and REM stages ($2.32 \pm 0.09$ times/min, $t_{(106)} = 1.99$, $p = 0.0492$, **Fig. 2b, Table S2-S4**), consistent with previous findings (Ngo et al., 2013; Staresina et al., 2015). After extracting all EEG epochs where SO-spindle coupling occurred, we analysed their spectral and phase characteristics. The spindles were most likely to occur slightly before the UP state peak of SOs (**Fig. 2a, e**), aligning with results from both animal studies (Maingret et al., 2016) and human research (Staresina et al., 2015). In our data, this pattern was consistent across subjects (**Fig. 2d**, Rayleigh test: $z = 9.51$, $p < 1e-4$), with the peak of the spindle aligned at an SO phase of $-41.61 \pm 0.86°$ (the SO UP state peak is $0°$).

Interestingly, spindle events coupled with SOs had higher amplitudes ($7.69 \pm 0.26$ µV) compared to those not coupled ($7.33 \pm 0.25$ µV, $t_{(106)} = 4.65$, $p < 1e-4$, **Fig. 2c**, left). Similarly, SO events coupled with spindles exhibited higher amplitudes ($26.28 \pm 1.34$ µV) than those not coupled ($24.56 \pm 1.33$ µV, $t_{(106)} = 6.86$, $p < 1e-4$, **Fig. 2c**, right). These findings suggest coordinated neural processing during SO-spindle coupling. Next, we explore the brain-wide activation associated with these sleep rhythms.

**Fig. 2 | Sleep rhythms and SO-spindle coupling. a,** The SO-spindle coupling in the temporal frequency domain. The upper two rows illustrates the spindle (12-16 Hz) phase-locked in the transition to UP-state of SO (0.16-1.25 Hz). The bottom row shows the averaged temporal frequency pattern across all instances of SO-spindle coupling and over all subjects. **b,** SO-spindle coupling density across different sleep stages. **c,** Differences between coupled and uncoupled sleep rhythms. The left panel shows the difference in amplitude between spindles coupled with SOs (Coupling) and spindles not coupled with SOs (Other). The right panel displays the difference in amplitude between SOs coupled with spindles (Coupling) and SOs not coupled with spindles (Other). **d,** Phase modulation of SO-spindle coupling. Spindle peaks cluster slightly before the UP-state peak of SO (i.e., 0°), where –π/2 reflects the transition from DOWN to UP-state. The histogram represents the distribution of coupling directions across all subjects, with the red line showing the mean. Coupling phases for each subject are plotted on the circle, with coupling strength color-coded. **e,** Distribution of spindle peaks on the SO phase during all SO-spindle coupling events across participants. The distribution is represented by a probability density function, and the density is evaluated at 100 equally spaced points covering the data range. Each dot represents data from an individual subject. Error bars indicate the SEM. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.

**Sleep rhythms associated brain activation**

To investigate brain-wide activity during sleep rhythms, we modeled the fMRI blood oxygen level-dependent (BOLD) signal based on EEG-identified timing of SOs and spindle events during N2/3 sleep stages. The design matrix included the main effects of SOs, spindles, and their coupling (**Fig. 3a**, detailed in Methods). This method is validated in our previous work to effectively capture replay-aligned brain activation during rest (Huang et al., 2024). We found that SOs were associated with positive activation in the thalamus (ROI analysis, $t_{(106)} = 2.38$, $p = 0.0096$, one-sample $t$ test) and significant deactivation in the neocortex (**Fig. 3b**, see also **Fig. S5**), particularly in the default mode network (DMN), which includes the mPFC and posterior cingulate cortex (PCC), consistent with previous research (Dang-Vu et al., 2008; Picchioni et al., 2011; Tagliazucchi et al., 2013). This deactivation of the neocortex, particularly the DMN, at the trough of SOs is intriguing, given that the DMN is thought to encode the internal model of the world or cognitive map (Baldassano, Hasson, & Norman, 2018; Constantinescu, O'Reilly, & Behrens, 2016; Park, Miller, & Boorman, 2021).

Spindles were correlated with positive activation in the thalamus (ROI analysis, $t_{(106)} = 15.39$, $p < 1$e-4), the anterior cingulate cortex (ACC), and the putamen, alongside deactivation in the DMN (**Fig. 3c**). Notably, SO-spindle coupling was linked to significant activation in both the thalamus (ROI analysis, $t_{(106)} = 3.38$, $p = 0.0005$) and the hippocampus (ROI analysis, $t_{(106)} = 2.50$, $p = 0.0070$, **Fig. 3d**). However, no decrease in DMN activity was found during SO-spindle coupling, and DMN activity during SO was significantly lower than during coupling (ROI analysis, $t_{(106)} = -4.17$, $p < 1$e-4). For more detailed activation patterns, see **Table S5-S7**. We also varied the threshold used to detect SO events to assess its effect on hippocampal activation during SO-spindle coupling and observed that hippocampal activation remained significant when the percentile thresholds for SO detection ranged between 71% and 80% (see **Fig. S6**).

To understand the functional implications of these sleep rhythm-related brain activation, we performed open-ended cognitive state decoding using the NeuroSynth database (Yarkoni et al., 2011), examining associations between topic terms and regions of interest derived from the activation patterns of the sleep

rhythms. Topic terms were ranked by their $z$-scores across the rhythms to assess overall functional relevance (see Methods for details). For positive activation patterns (**Fig. S7a**), SOs were associated with declarative ($z = 6.31$, $p < 1e-4$) and episodic memory ($z = 5.08$, $p < 1e-4$), while spindles aligned most with working memory ($z = 4.08$, $p < 1e-4$). SO-spindle coupling showed the strongest similarity to episodic ($z = 3.66$, $p = 0.0003$) and declarative memory ($z = 2.21$, $p = 0.0271$). For negative (deactivation) patterns (**Fig. S7b**), SOs were linked to declarative memory ($z = 7.61$, $p < 1e-4$), while spindles resembled task-related activations ($z = 7.05$, $p < 1e-4$). SO-spindle coupling had the highest similarity to task-related ($z = 9.40$, $p < 1e-4$) and working memory ($z = 9.15$, $p < 1e-4$). These findings suggest that internal memory-related regions are more likely to reactivate during SO-spindle coupling, whereas task-related activations, like working memory, are less prominent.

Fig. 3 | **Brain-wide activation associated with sleep rhythms. a,** Simultaneous EEG-fMRI analysis framework for detecting brain-wide activation during sleep rhythms. Detected SOs, spindles, and their coupling were convolved with the hemodynamic response function (HRF) and downsampled to match fMRI temporal resolution. These events formed the design matrix for the general linear model (GLM) analysis of fMRI activity during sleep, linking the EEG-derived timing of sleep rhythms to the corresponding brain responses in fMRI. **b,** Brain-wide activation associated with SOs. The upper row illustrates SOs, and the lower row shows the fMRI activation pattern during SO events, whole-brain family-wise error (FWE) corrected at the cluster level ($p < 0.05$) with a cluster-forming voxel threshold of $p < 0.001$. **c,** Brain-wide activation associated with spindles. Same as panel b, but for spindle events. **d,** Brain-wide activation associated with SO-spindle coupling (compared to non-coupling events).

**Functional connectivity during SO-spindle coupling**

Previous studies have indicated that the hippocampal-cortical dialogue during sleep memory consolidation relies on the thalamus (Coulon, Budde, & Pape, 2012; Ferraris et al., 2021; Latchoumane et al., 2017). To explore this in the human brain, we examined how functional connectivity (FC) changes when spindles couple (versus not) with SOs during sleep, using psychophysiological interaction (PPI) analysis (Friston et al., 1997). Building on the previous GLM (**Fig. 3a**), this model incorporated two additional regressors: the BOLD signal of the ROI seed and its interaction with SO-spindle events (**Fig. 4a**, detailed in Methods).

We first examined whole-brain FC (physiological term) in three ROIs: the hippocampus (bilateral, anatomically defined), thalamus (bilateral, anatomically defined), and mPFC (bilateral, functionally defined). The hippocampus-based FC was mainly connected with the DMN, particularly the mPFC (**Fig. 4b**). The thalamus-based FC was primarily linked to the basal ganglia and ACC (**Fig. 4c**), and the mPFC-based FC showed strong connections with the hippocampus and PCC (**Fig. 4d**).

Crucially, the interaction between FC and SO-spindle coupling revealed that only the functional connectivity of hippocampus -> thalamus (ROI analysis, $t_{(106)} = 1.86$, $p = 0.0328$) and thalamus -> mPFC (ROI analysis, $t_{(106)} = 1.98$, $p = 0.0251$) significantly increased during SO-spindle coupling, with no significant changes in all other pathways (**Fig. 4e**). We also conducted PPI analyses for the other two events (SOs and spindles), and neither yielded significant connectivity changes in the three ROIs, as all failed to survive whole-brain FWE correction at the cluster level ($p < 0.05$). Together, these findings suggest that the thalamus, likely via spindles, coordinates hippocampal-cortical communication selectively during SO-spindle coupling, but not isolated SOs or spindle events alone.

**Fig. 4 | Functional connectivity changes during SO-spindle coupling. a,** The PPI analysis framework for detecting brain-wide connectivity changes during SO-spindle coupling. This starts by setting a specific ROI (e.g., the hippocampus) as the seed to extract the BOLD signal (physiological condition) and using identified SO-spindle coupling events as the psychological condition to compute the interaction term. The design matrix includes the main effects of the physiological and psychological conditions, along with their interaction. This analysis examines whether whole-brain communication with the hippocampus changes as a function of SO-spindle coupling. **b,** Hippocampus-based functional connectivity with the whole brain (main effect of hippocampus BOLD signal in PPI analysis). The hippocampus ROI is bilateral, anatomically defined (bottom, orange colour). Brain-wide connectivity is shown with whole-brain FWE correction at the cluster level ($p < $ 1e-7) with a cluster-forming voxel threshold of $p_{unc.} < 0.001$ for visualization purpose. **c,** Same with panel b, but based on thalamus (bilateral anatomically defined ROI). **d,** Same with panel b, but based on the mPFC (bilateral functionally defined ROI, detailed in Methods). **e,** Functional connectivity changes during SO-spindle coupling for hippocampus-based (left bottom, orange colour), thalamus-based (middle, green colour), and mPFC-based (right bottom, purple colour) connectivity. The results of ROI analysis for each direction are shown on the arrows. * $p < 0.05$, ns., not significant. Abbreviations: FC - functional connectivity, PPI - psychophysiological interaction.

**Discussion**

Using simultaneous EEG and fMRI during nocturnal naps in 107 subjects, we identified robust coupling between spindles and the UP-state of SOs in deep NREM sleep (N2/3 stages). This coupling was linked to distinct neural activation patterns, including significant thalamic activation during spindle occurrences, reduced activity in the DMN during SOs trough, and selective hippocampal activation when spindles and SOs coupled. Furthermore, functional connectivity analyses revealed increased connectivity from the hippocampus to the thalamus and from the thalamus to the mPFC, only during SO-spindle coupling, but not isolated SOs or spindle events alone. These results shed light on how the thalamus may coordinate hippocampal-cortical interactions during NREM sleep.

Our data provide insights into the neurobiological underpinnings of these sleep rhythms. SOs, originating mainly in neocortical areas such as the mPFC, alternate between DOWN- and UP-states. The thalamus generates sleep spindles, which in turn couple with SOs. Our finding that spindle peaks consistently occurred slightly before the UP-state peak of SOs (in 83 out of 107 participants), concurs with prior studies, including Schreiner et al. (2021). Yet it differs from some results suggesting spindles might peak right at the SO UP-state (Hahn et al., 2020; Helfrich et al., 2018). Such discrepancies could arise from differences in detection algorithms, participant age (Helfrich et al., 2018), or subtle variations in cortical-thalamic timing. Nonetheless, these results underscore the importance of coordinated SO-spindle interplay in supporting sleep-dependent processes.

When modeling the timing of these sleep rhythms in the fMRI, we observed hippocampal activation selectively during SO-spindle events. This suggests the possibility of triple coupling (SOs–spindles–ripples), even though our scalp EEG was not sufficiently sensitive to detect hippocampal ripples—key markers of memory replay (Buzsáki, 2015). Recent iEEG evidence indicates that ripples often co-occur with both spindles (Ngo, Fell, & Staresina, 2020) and SOs (Staresina et al., 2015; Staresina et al., 2023). Therefore, the hippocampal involvement during SO-spindle events in our study may reflect memory replay from the hippocampus, propagated via thalamic spindles to distributed cortical regions.

The thalamus, known to generate spindles (Halassa et al., 2011), plays a key role in producing and

coordinating sleep rhythms (Coulon, Budde, & Pape, 2012; Crunelli et al., 2018), while the

hippocampus is found essential for memory consolidation (Buzsáki, 2015; Diba & Buzsáki, 2007;

Singh, Norman, & Schapiro, 2022). The increased hippocampal and thalamic activity, along with

strengthened connectivity between these regions and the mPFC during SO-spindle events, underscores

a hippocampal-thalamic-neocortical information flow. This aligns with recent findings suggesting the

thalamus orchestrates neocortical oscillations during sleep (Schreiner et al., 2022). The thalamus and

hippocampus thus appear central to memory consolidation during sleep, guiding information transfer

to the neocortex, e.g., mPFC.

An intriguing aspect of our findings is the reduced DMN activity during SOs when modeled at the SO

trough (DOWN-state). This reduced DMN activity may reflect large-scale neural inhibition

characteristic of the SO trough. The DMN is typically active during internally oriented cognition (e.g.,

self-referential processing or mind-wandering) and is suppressed during external stimuli processing

(Yeshurun, Nguyen, & Hasson, 2021). It is unlikely, however, that this suppression of DMN during SO

events is related to a shift from internal cognition to external responses given it is during deep sleep

time. Instead, it could be driven by the inherent rhythmic pattern of SOs, which makes it difficult to

separate UP- from DOWN-states (the two temporal regressors were highly correlated, and similar brain

activation during SOs events was obtained if modelled at the SO peak instead, **Fig. S5**). Since the

amplitude at the SO trough is consistently larger than that at the SO peak, the neural activation we

detected may primarily capture the large-scale inhibition from DOWN-state. Interestingly, no such

DMN reduction was found during SO-spindle coupling, implying that coupling may involve distinct

neural dynamics that partially re-engage DMN-related processes, possibly reflecting memory-related

reactivation. Future research using high-temporal-resolution techniques like iEEG could clarify these

possibilities.

To explore functional relevance, we employed an open-ended cognitive state decoding approach using

meta-analytic data (NeuroSynth: Yarkoni et al. (2011)). Although this method usefully generates

hypotheses about potential cognitive processes, particularly in the absence of a pre- and post-sleep

memory task, it is inherently indirect. Many cognitive terms showed significant associations (16 of 50), such as "episodic memory," "declarative memory," and "working memory." We focused on episodic/declarative memory given the known link with hippocampal reactivation (Diekelmann & Born, 2010; Staresina et al., 2015; Staresina et al., 2023). Nonetheless, these inferences regarding memory reactivation should be interpreted cautiously without direct behavioral measures. Future research incorporating explicit tasks before and after sleep would more rigorously validate these potential functional claims.

Despite providing new insights, our study has several limitations. First, our scalp EEG did not directly capture hippocampal ripples, preventing us from conclusively demonstrating triple coupling. Second, the combination of EEG-fMRI and the lack of a memory task limit our ability to parse fine-grained BOLD responses at the DOWN- vs. UP-states of SOs and link observed activations to behavioral outcomes. Third, the use of large anatomical ROIs may mask subregional contributions of specific thalamic nuclei or hippocampal subfields. Finally, without a memory task, we cannot establish a direct behavioral link between sleep-rhythm-locked activation and memory consolidation. Future studies combining techniques such as ultra-high-field fMRI or iEEG with cognitive tasks may refine our understanding of subregional network dynamics and functional significance during sleep.

In summary, we found that SO-spindle coupling during deep NREM sleep is associated with distinct activation in both the thalamus and hippocampus, accompanied by increased connectivity from the hippocampus to the thalamus and from the thalamus to the mPFC. These results emphasize the central role of the thalamus in orchestrating hippocampal-cortical interactions and highlight potential neural substrates underlying the complex interplay of SOs, spindles, and, possibly, ripples. Our open-ended decoding analyses suggest that these coordinated sleep events may be relevant for memory-related processes; however, future experiments incorporating explicit behavioral measures are necessary to confirm their functional significance. By revealing how these rhythms modulate widespread brain networks, our findings open avenues for developing targeted neuromodulatory strategies to enhance sleep-dependent cognitive functions and potentially address sleep-related disorders.

## Methods

### Participants & Protocol

A total of 138 healthy adults (age: $22.56 \pm 0.30$; 81 females) were recruited for the study. All participants had either normal vision or vision corrected to normal standards. None had any past psychiatric or neurological disorders. Prior to participation, they were screened for eligibility to undergo magnetic resonance imaging (MRI). The experiment received approval from Peking University ethics committee (reference number: #2015-26, #2018-11-05). Every participant provided written informed consent.

At the start of the experiment, subjects arrived early at the laboratory for registration. They signed informed consent forms, washed their hair, changed their clothes, and put on the EEG cap in preparation for the experiment. After these preliminary steps, participants entered the MRI scanning room by around 20:30, settled down for sleep, and data collection began. Given concern for potential data drift and decline in the signal-to-noise ratio due to extended and continuous fMRI scanning, the data acquisition was segmented into several sessions for data collection. A "sleep" session was concluded either when a participant was fully awake (unable to fall asleep again or needed to use the restroom) or after 8192 seconds (the maximum scanning time of the scan, equating to 4096 volumes with 2 seconds TR).

After removing subjects with excessive head movements (mean FD > 0.5) or those whose sleep durations were under 10 minutes, the final participant count for further analyses was 107 (age: $22.40 \pm 0.34$; 63 females) with a mean scanning duration of sleep of $3.01 \pm 0.14$ hours.

### EEG data acquisition

EEG was recorded simultaneously with fMRI data using an MR-compatible EEG amplifier system (BrainAmps MR-Plus, Brain Products, Germany), along with a specialized electrode cap. The recording was done using 64 channels in the international 10/20 system, with the reference channel positioned at FCz. In order to adhere to polysomnography (PSG) recording standards, six electrodes were removed from the EEG cap: one for electrocardiogram (ECG) recording, two for electrooculogram (EOG) recording, and three for electromyogram (EMG) recording. EEG data was recorded at a sample rate of

5000 Hz, the resistance of the reference and ground channels was kept below 10 kΩ, and the resistance

of the other channels was kept below 20 kΩ. To synchronize the EEG and fMRI recordings, the

BrainVision recording software (BrainProducts, Germany) was utilized to capture triggers from the

MRI scanner. The EEG sampling was synchronized to the MR scanner's 10 MHz gradient system clock,

ensuring a stable gradient artifact shape over time and enabling accurate artifact removal. This was

achieved via the standard clock synchronization interface of the EEG amplifier, minimizing timing jitter

and drift.

**MRI data acquisition**

All MRI data were acquired using a 20-channel head coil on a research-dedicated 3-Tesla Siemens

Magnetom Prisma MRI scanner. Earplugs and cushions were provided for noise protection and head

motion restriction. We chose the 20-channel head coil because it was compatible with our EEG system

and ensured sufficient signal-to-noise ratio (SNR) for both EEG and fMRI acquisition. The EEG

electrode cables were guided through the lateral and posterior openings of the head coil, secured with

foam padding to reduce motion and minimize MR-related artifacts. Moreover, given the extended nature

of nocturnal sleep recordings, the 20-channel coil helped maintain participant comfort while still

achieving high-quality simultaneous EEG-fMRI data.

For registration purposes, high-resolution T1-weighted anatomical images were acquired using a three-

dimensional magnetization prepared rapid acquisition gradient-echo sequence (sequence specification:

number of slices = 192, repetition time (TR) = 2,530 ms, echo time (TE) = 2.98 ms, inversion time =

1,100 ms, voxel size = $0.5 \times 0.5 \times 1$ mm$^3$, flip angle (FA) = 7°). The participants were asked to lie

quietly in the scanner during data acquisition.

Then, the "sleep" session began after the participants were instructed to try and fall asleep. For the

functional scans, whole-brain images were acquired using k-space and steady-state T2*-weighted

gradient echo-planar imaging (EPI) sequence that is sensitive to the BOLD contrast. This measures

local magnetic changes caused by changes in blood oxygenation that accompany neural activity

(sequence specification: 33 slices in interleaved ascending order, TR = 2000 ms, TE = 30 ms, voxel size

= 3.5 × 3.5 × 4.2 mm$^3$, FA = 90°, matrix = 64 × 64, gap = 0.7 mm). A relatively long TR and larger

voxel size were chosen to optimize SNR and reduce susceptibility-related distortions, which are critical

in EEG-fMRI sleep studies where head motion and physiological noise can be substantial. The longer

TR allowed whole-brain coverage with sufficient temporal resolution, while the larger voxel size helped

enhance BOLD sensitivity and minimize partial volume effects in deep brain structures (e.g., the

thalamus and hippocampus), which are key regions of interest in this study.

**EEG data preprocessing and sleep stage scoring**

EEG data collected inside MRI scanner was contaminated by imaging, ballistocardiographic (BCG)

and ocular artifacts. To deal with these artifacts, the preprocessing processes was divided into two parts:

automated processing and manual processing.

The automated processing was carried out using custom Python scripts and the MNE toolkit (Gramfort

et al., 2013). This encompassed the following procedures: **I.** The raw EEG data were downsampled

from 5000 Hz to 2048 Hz. This downsampled rate accelerates subsequent analyses while retaining the

necessary precision to distinguish MR noise. **II.** Imaging artifacts were removed using the Average

Artefact Subtraction method (Allen, Josephs, & Turner, 2000). **III.** We then applied 0.1 - 30 Hz band-

pass finite impulse response (FIR) filters to all channels. The EEG and EOG electrodes were re-

referenced by subtracting the value from the M1 or M2 electrode (whichever was from the opposite

hemisphere). The outer EMG electrodes were re-referenced to the middle one. **IV.** The signal space

projection algorithm was applied to extract all epochs based on TRs and remove residual MR noise, as

well as automatically extracting all heartbeat epochs using ECG electrode data for the removal of BCG

noise. **V.** The data were further downsampled to 100 Hz and subjected to Independent Component

Analysis (ICA).

Following this, the manual processing is carried out with labelling and removing the physiological

artifacts ICs belonging to eye, muscle and residual BCG through the use of custom MATLAB scripts

and the EEGLAB toolkit (Delorme & Makeig, 2004). Sleep stages were then scored by YASA toolkit

(Vallat & Walker, 2021) after preprocessing, necessitating data from electrodes C3, left EOG, and left

EMG. The classification of N2/3 stages were further verified by experts for validity.

**MRI data preprocessing**

Results included in this manuscript come from preprocessing performed using *fMRIPrep* 21.0.4

(Esteban et al., 2020; Esteban et al., 2019), which is based on *Nipype* 1.6.1 (Gorgolewski et al., 2011).

Many internal operations of *fMRIPrep* use *Nilearn* 0.8.1 (Abraham et al., 2014), mostly within the

functional processing workflow. For more details of the pipeline, see

https://fmriprep.readthedocs.io/en/latest/workflows.html.

*Conversion of data to the brain imaging data structure standard.* To facilitate further analysis and

sharing of data, all study data were arranged according to the Brain Imaging Data Structure (BIDS)

specification using dcm2bids tool, which is freely available from

https://unfmontreal.github.io/Dcm2Bids/.

*Anatomical data preprocessing.* The T1-weighted (T1w) image was corrected for intensity non-

uniformity (INU) with N4BiasFieldCorrection (Tustison et al., 2010), distributed with ANTs 2.3.3

(Avants et al., 2008), and used as T1w-reference throughout the workflow. The T1w-reference was then

skull-stripped with a *Nipype* implementation of the antsBrainExtraction.sh workflow (from ANTs),

using OASIS30ANTs as target template. Brain tissue segmentation of cerebrospinal fluid (CSF), white-

matter (WM) and gray-matter (GM) was performed on the brain-extracted T1w using fast (FSL

6.0.5.1:57b01774, RRID:SCR_002823) (Zhang, Brady, & Smith, 2001). Volume-based spatial

normalization to one standard space (MNI152NLin2009cAsym) was performed through nonlinear

registration with antsRegistration (ANTs 2.3.3), using brain-extracted versions of both T1w reference

and the T1w template. The following template was selected for spatial normalization: *ICBM 152*

*Nonlinear Asymmetrical template version 2009c* (Fonov, 2009) [RRID:SCR_008796; TemplateFlow

ID: MNI152NLin2009cAsym].

*Functional data preprocessing.* First, a reference volume and its skull-stripped version were generated

using a custom methodology of *fMRIPrep*. Head-motion parameters with respect to the BOLD reference

(transformation matrices, and six corresponding rotation and translation parameters) are estimated

before any spatiotemporal filtering using mcflirt (Jenkinson et al., 2002) [FSL 6.0.5.1:57b01774].

BOLD runs were slice-time corrected to 0.96s (0.5 of slice acquisition range 0s-1.92s) using 3dTshift

from AFNI (Cox & Hyde, 1997). The BOLD time-series (including slice-timing correction when

applied) were resampled onto their original, native space by applying the transforms to correct for head-

motion. These resampled BOLD time-series will be referred to as *preprocessed BOLD in original space*,

or just *preprocessed BOLD*. The BOLD reference was then co-registered to the T1w reference using

mri_coreg (FreeSurfer) followed by flirt (Jenkinson & Smith, 2001) [FSL 6.0.5.1:57b01774] with the

boundary-based registration (Greve & Fischl, 2009) cost-function. Co-registration was configured with

six degrees of freedom. Several confounding time-series were calculated based on the *preprocessed*

*BOLD*: framewise displacement (FD), DVARS and three region-wise global signals. FD was computed

using two formulations following Power (absolute sum of relative motions) (Power et al., 2014) and

Jenkinson (relative root mean square displacement between affines, Jenkinson et al. (2002)). FD and

DVARS are calculated for each functional run, both using their implementations in *Nipype* (following

the definitions by Power et al. (2014)). The three global signals are extracted within the CSF, the WM,

and the whole-brain masks. Additionally, a set of physiological regressors were extracted to allow for

component-based noise correction (*CompCor)* (Behzadi et al., 2007). Principal components are

estimated after high-pass filtering the *preprocessed BOLD* time-series (using a discrete cosine filter

with 128s cut-off) for the two *CompCor* variants: temporal (tCompCor) and anatomical (aCompCor).

tCompCor components are then calculated from the top 2% variable voxels within the brain mask. For

aCompCor, three probabilistic masks (CSF, WM and combined CSF+WM) are generated in anatomical

space. The implementation differs from that of Behzadi et al. (2007) in that instead of eroding the masks

by 2 pixels on BOLD space, the aCompCor masks are subtracted a mask of pixels that likely contain a

volume fraction of GM. This mask is obtained by thresholding the corresponding partial volume map

at 0.05, and it ensures components are not extracted from voxels containing a minimal fraction of GM.

Finally, these masks are resampled into BOLD space and binarized by thresholding at 0.99 (as in the

original implementation). Components are also calculated separately within the WM and CSF masks.

For each CompCor decomposition, the *k* components with the largest singular values are retained, such

that the retained components' time series are sufficient to explain 50 percent of variance across the nuisance mask (CSF, WM, combined, or temporal). The remaining components are dropped from consideration. Frames that exceeded a threshold of 0.5 mm FD or 1.5 standardised DVARS were annotated as motion outliers.

The BOLD time-series were resampled into standard space, generating a *preprocessed BOLD run in MNI152NLin2009cAsym space*. First, a reference volume and its skull-stripped version were generated using a custom methodology of *fMRIPrep*. All resamplings can be performed with *a single interpolation step* by composing all the pertinent transformations (i.e., head-motion transform matrices, susceptibility distortion correction when available, and co-registrations to anatomical and output spaces). Gridded (volumetric) resamplings were performed using antsApplyTransforms (ANTs), configured with Lanczos interpolation to minimize the smoothing effects of other kernels (Lanczos, 1964). Non-gridded (surface) resamplings were performed using mri_vol2surf (FreeSurfer).

**EEG sleep rhythms analysis**

In line with other studies examining the coordination between SOs and spindles, particularly their spatial distribution (Massimini et al., 2004; Molle et al., 2011), EEG data from the F3 electrode were used to quantify the occurrence of SOs and spindles. Using the established detection algorithms (detailed below), we identified SOs and sleep spindles for each subject.

*Detection of SOs*. Data were first bandpass-filtered between 0.16 and 1.25 Hz (Butterworth filter, order 3, bidirectional filtering for zero phase). After identifying all positive-to-negative zero crossings, potential SOs were defined based on the interval between consecutive zero crossings (ranging from 0.8 s to 3 s). For each potential SO, we calculated the amplitude range (peak minus trough). Only those exceeding the 75th percentile in amplitude were labeled as SOs, following previous work (Schreiner et al., 2021).

*Detection of sleep spindles*. Data were bandpass-filtered between 12 and 16 Hz (Butterworth filter, order 3, bidirectional filtering for zero phase). The root mean square (RMS) of the filtered signal was computed with a 200 ms sliding time window. The amplitude threshold for spindles was set at the 75th

percentile of RMS values. Any RMS values that surpassed this threshold for 0.5 s to 3 s were identified

as spindles (Staresina et al., 2015).

*Detection of SO-spindle couplings.* From the detected SOs and spindles, we identified the peak time of

each spindle. Within each SO interval, we checked for a spindle peak; if found, it was recorded as an

SO-spindle coupling event. For every SO-spindle coupling event, an epoch was created time-locked to

the SO trough (as the central reference), following Schreiner et al. (2021). We extracted data in a [-4 s

to 4 s] window around this point, forming the epoch for each coupling event.

We note that the above methods for detecting SOs, spindles, and their couplings were originally

developed for N2 and N3 sleep data, based on the specific characteristics of these stages. These methods

are widely recognized in sleep research (Hahn et al., 2020; Helfrich et al., 2019; Helfrich et al., 2018;

Ngo, Fell, & Staresina, 2020; Schreiner et al., 2022; Schreiner et al., 2021; Staresina et al., 2015;

Staresina et al., 2023). However, because this percentile-based detection approach will inherently

identify a certain number of events if applied to other stages (e.g., N1 and REM), the nature of these

events in those stages remains unclear compared to N2/N3. We nevertheless identified and reported the

detailed descriptive statistics of these sleep rhythms in all sleep stages, under the same operational

definitions, both for completeness and as a sanity check. Within the same subject, there should be more

SOs, spindles, and their couplings in N2/N3 than in N1 or REM (see also **Figure S2-S4**, **Table S1-S4**).

*Event-related potentials (ERP) analysis.* After completing the detection of each sleep rhythm event, we

performed ERP analyses for SOs, spindles, and coupling events in different sleep stages. Specifically,

for SO events, we took the trough of the DOWN-state of each SO as the zero-time point, then extracted

data in a [-2 s to 2 s] window from the broadband (0.1–30 Hz) EEG and used [-2 s to -0.5 s] for baseline

correction; the results were then averaged across 107 subjects (see **Fig. S2a**). For spindle events, we

used the peak of each spindle as the zero-time point and applied the same data extraction window and

baseline correction before averaging across 107 subjects (see **Fig. S2b**). Finally, for SO-spindle

coupling events, we followed the same procedure used for SO events (see **Fig. 2a**, **Figs. S3–S4**).

*Time-frequency analysis.* We performed this analysis on the SO-spindle coupling epochs using the FieldTrip toolbox (Oostenveld et al., 2011). Fourier analysis was executed on the data, utilizing a sliding time window with a forward shift of 50 ms. The window length was defined as five cycles of the present frequency (frequency range: 1-30 Hz, step size: 1 Hz). Before Fourier analysis, windowed data segments underwent multiplication by a Hanning taper. Subsequently, power values were normalized to z-scores within the time range of [-4 s to 4 s], selecting a longer duration to prevent edge artifacts from low-frequency activity in the target time window of [-1.5 s to 1.5 s].

*Preferred phase analysis.* All SO-spindle coupling epochs were filtered to the SO frequency band (0.16-1.25 Hz, 3rd-order Butterworth filter), and the instantaneous SO phase was determined through the Hilbert transform. Using the saved spindle peak times for each SO-spindle coupling, the preferred phase was deduced. The distribution of these phases across participants was then tested for uniformity using the Rayleigh test.

**fMRI GLM analysis**

All fMRI GLM analyses were conducted using only data from the N2/3 stages of NREM sleep. Segments of sleep data from these stages were concatenated for each participant. After pre-processing, the fMRI data was smoothed with an 8 mm FWHM kernel. All images underwent a temporal high-pass filter (width of 100 s), and the autocorrelation of the hemodynamic response was modelled via an AR(1) model. We then conducted General Linear Model (GLM) analyses to identify whole-brain activations associated with sleep rhythms and their coupling. We incorporated nuisance regressors estimated during the preprocessing with fMRIprep: six rigid-body motion-correction parameters identified during realignment (comprising three translation and rotation parameters), the mean White Matter, and the mean Cerebral Spinal Fluid signals. The cosine drift model was employed to detect and eliminate data fluctuations, thereby enhancing the accuracy of actual experimental effects. The GLMs were designed to evaluate the influence of experimental conditions on BOLD responses. All whole-brain analyses were thresholded using whole-brain FWE corrected at the cluster level ($p < 0.05$), with a cluster-forming voxel threshold of $p_{unc.} < 0.001$, unless otherwise specified.

The fMRI GLM analysis was designed to look for brain activations pertinent to SO, spindle, and SO-spindle coupling. Drawing from the EEG results concerning the timing of SO and spindle, we identified the trough of the SO as the onset for SO events and the peak of the spindle as the onset for spindle events. We chose the trough of the SO and the peak of the spindle as the onset points, primarily due to the larger amplitudes observed at these moments (thereby more robust to noise). This was also the common choice in previous EEG studies on SO and spindles (Ngo, Fell, & Staresina, 2020; Staresina et al., 2015; Staresina et al., 2023). To model SO-spindle coupling in fMRI, we followed Schreiner et al. (2021), and defining SO-spindle complex as the peak of spindle appeared within 1.5 seconds after SO trough. When such a peak was found, that particular SO was labelled as an SO-spindle event, with its SO trough labelled as the event onset. After convolving these events with the canonical Hemodynamic Response Function (HRF) of SPM and downsampling them to align with the fMRI time resolution, they were integrated as regressors in the design matrix for GLM analysis, allowing us to determine the effects of SO and spindle, as well as the SO-spindle coupling (**Fig. 3b-d**).

We have also conducted a control analysis aimed to distinguish brain activations associated with DOWN- vs. UP state of SOs. Drawing again from EEG data on SO and spindle, we randomly selected half of the SO events using the peak as the onset and the other half using the trough, to reduce collinearity between the two regressors. As with the main analysis, we identified the spindle peak as the onset for spindle events (**Fig. S5**).

**ROI analysis**

The objective of our ROI analysis was to test the activation of specific brain regions during distinct sleep rhythms. Within each ROI, the beta values at the subject-level were averaged across all voxels to facilitate subsequent statistical inferences (one-sample $t$ test). We defined the mDMN and mPFC ROIs functionally based on the meta-analysis results in Neurosynth (Yarkoni et al., 2011), by the keyword "medial default mode network" and "ventromedial prefrontal cortex" respectively. In addition to mDMN and mPFC, we defined other ROIs anatomically, using the high-resolution Harvard-Oxford Atlas probabilistic atlas (Desikan et al., 2006).

**Open-ended Cognitive State Functional Decoding**

For functional decoding, we followed the approach in Margulies et al. (2016). We used the NeuroSynth (Yarkoni et al., 2011) meta-analytic database (www.neurosynth.org) to assess topic terms associated with the sleep rhythms activation patterns. ROI masks were created from the whole-brain activation patterns associated with the three sleep rhythms in **Fig. 3b-d**. These activation patterns were binarized using a threshold of $p < 0.001$, forming a brain mask for each sleep rhythm.

For each ROI mask, the analysis output was a z-statistic associated with a feature term, and the terms were ranked based on the weighted sum of their z-scores. The feature terms were derived from the most general set of 50 topic terms available in the NeuroSynth v3 dataset (https://github.com/NeuroanatomyAndConnectivity/gradient_analysis/blob/master/gradient_data/neurosynth/v3-topics-50-keys.txt).

From the 50 topics, 16 exceeded the threshold of $z > 1.96$ ($p < 0.05$). Two terms were removed as they did not represent coherent cognitive functions, leaving 14 relevant topic terms (**Fig. S7**). These included: episodic memory, declarative memory, working memory, task representation, language, learning, faces, visuospatial processing, category recognition, cognitive control, reading, cued attention, inhibition, and action.

**PPI analysis**

PPI (psychophysiological interaction) analysis is designed to quantify context-dependent alterations in effective connectivity between brain regions. For instance, it can determine whether hippocampus-based functional connectivity with the thalamus varies based on the occurrence of an SO-spindle coupling event. We conducted nine whole-brain PPI analyses with *nilearn* to investigate interactions based in the hippocampus, thalamus or mPFC during SO-spindle coupling, isolated SO and isolated spindle, utilizing the same ROIs. Both the main effects of physiological and physiological condition, as well as their interaction were modelled in the design matrix for PPI analysis. The main effect of physiological term produced functional connectivity maps of the entire brain in relation to the specified

610    ROI (**Fig. 4b-d**). While the interaction term asked for the *change* of functional connectivity along with

611    the SO-spindle coupling (**Fig. 4e**).

## References

Abraham, A., Pedregosa, F., Eickenberg, M., Gervais, P., Mueller, A., Kossaifi, J., Gramfort, A., Thirion, B., & Varoquaux, G. (2014). Machine learning for neuroimaging with scikit-learn. *Frontiers in neuroinformatics*, *8*, 14.

Allen, P. J., Josephs, O., & Turner, R. (2000). A method for removing imaging artifact from continuous EEG recorded during functional MRI. *NeuroImage*, *12*(2), 230-239.

Amzica, F., & Steriade, M. (2002). The functional significance of K-complexes. *Sleep medicine reviews*, *6*(2), 139-149.

Avants, B. B., Epstein, C. L., Grossman, M., & Gee, J. C. (2008). Symmetric diffeomorphic image registration with cross-correlation: evaluating automated labeling of elderly and neurodegenerative brain. *Medical image analysis*, *12*(1), 26-41.

Baldassano, C., Hasson, U., & Norman, K. A. (2018). Representation of real-world event schemas during narrative perception. *Journal of Neuroscience*, *38*(45), 9689-9699.

Behzadi, Y., Restom, K., Liau, J., & Liu, T. T. (2007). A component based noise correction method (CompCor) for BOLD and perfusion based fMRI. *NeuroImage*, *37*(1), 90-101.

Bergmann, T. O., Mölle, M., Diedrichs, J., Born, J., & Siebner, H. R. (2012). Sleep spindle-related reactivation of category-specific cortical regions after learning face-scene associations. *NeuroImage*, *59*(3), 2733-2742.

Buzsáki, G. (2015). Hippocampal sharp wave-ripple: A cognitive biomarker for episodic memory and planning. *Hippocampus*, *25*(10), 1073-1188.

Caporro, M., Haneef, Z., Yeh, H. J., Lenartowicz, A., Buttinelli, C., Parvizi, J., & Stern, J. M. (2012). Functional MRI of sleep spindles and K-complexes. *Clinical neurophysiology*, *123*(2), 303-309.

Carskadon, M. A., & Dement, W. C. (2005). Normal human sleep: an overview. *Principles and practice of sleep medicine*, *4*(1), 13-23.

Cash, S. S., Halgren, E., Dehghani, N., Rossetti, A. O., Thesen, T., Wang, C., Devinsky, O., Kuzniecky, R., Doyle, W., & Madsen, J. R. (2009). The human K-complex represents an isolated cortical down-state. *Science*, *324*(5930), 1084-1087.

Constantinescu, A. O., O'Reilly, J. X., & Behrens, T. E. (2016). Organizing conceptual knowledge in humans with a gridlike code. *Science*, *352*(6292), 1464-1468.

Coulon, P., Budde, T., & Pape, H.-C. (2012). The sleep relay—the role of the thalamus in central and decentral sleep regulation. *Pflügers Archiv-European Journal of Physiology*, *463*, 53-71.

Cousins, J. N., El-Deredy, W., Parkes, L. M., Hennies, N., & Lewis, P. A. (2016). Cued reactivation of motor learning during sleep leads to overnight changes in functional brain activity and connectivity. *PLoS biology*, *14*(5), e1002451.

Cox, R. W., & Hyde, J. S. (1997). Software tools for analysis and visualization of fMRI data. *NMR in Biomedicine: An International Journal Devoted to the Development and Application of Magnetic Resonance In Vivo*, *10*(4-5), 171-178.

Crunelli, V., Lőrincz, M. L., Connelly, W. M., David, F., Hughes, S. W., Lambert, R. C., Leresche, N., & Errington, A. C. (2018). Dual function of thalamic low-vigilance state oscillations: rhythm-regulation and plasticity. *Nature Reviews Neuroscience*, *19*(2), 107-118.

Czisch, M., Wehrle, R., Stiegler, A., Peters, H., Andrade, K., Holsboer, F., & Sämann, P. G. (2009). Acoustic oddball during NREM sleep: a combined EEG/fMRI study. *PloS one*, *4*(8), e6749.

Dang-Vu, T. T., Schabus, M., Desseilles, M., Albouy, G., Boly, M., Darsaud, A., Gais, S., Rauchs, G., Sterpenich, V., Vandewalle, G., Carrier, J., Moonen, G., Balteau, E., Degueldre, C., Luxen, A., Phillips, C., & Maquet, P. (2008). Spontaneous neural activity during human slow wave sleep. *Proceedings of the National Academy of Sciences*, *105*(39), 15160-15165.

Delorme, A., & Makeig, S. (2004). EEGLAB: an open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of neuroscience methods*, *134*(1), 9-21.

Desikan, R. S., Segonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., Buckner, R. L., Dale, A. M., Maguire, R. P., Hyman, B. T., Albert, M. S., & Killiany, R. J. (2006). An automated

labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *NeuroImage*, *31*(3), 968-980.

Diba, K., & Buzsáki, G. (2007). Forward and reverse hippocampal place-cell sequences during ripples. *Nature Neuroscience*, *10*(10), 1241.

Diekelmann, S., & Born, J. (2010). The memory function of sleep. *Nature Reviews Neuroscience*, *11*(2), 114-126.

Esteban, O., Ciric, R., Finc, K., Blair, R. W., Markiewicz, C. J., Moodie, C. A., Kent, J. D., Goncalves, M., DuPre, E., Gomez, D. E. P., Ye, Z., Salo, T., Valabregue, R., Amlien, I. K., Liem, F., Jacoby, N., Stojic, H., Cieslak, M., Urchs, S., . . . Gorgolewski, K. J. (2020). Analysis of task-based functional MRI data preprocessed with fMRIPrep. *Nature protocols*, *15*(7), 2186-2202.

Esteban, O., Markiewicz, C. J., Blair, R. W., Moodie, C. A., Isik, A. I., Erramuzpe, A., Kent, J. D., Goncalves, M., DuPre, E., Snyder, M., Oya, H., Ghosh, S. S., Wright, J., Durnez, J., Poldrack, R. A., & Gorgolewski, K. J. (2019). fMRIPrep: a robust preprocessing pipeline for functional MRI. *Nature methods*, *16*(1), 111-116.

Fernandez, L. M., & Lüthi, A. (2020). Sleep spindles: mechanisms and functions. *Physiological Reviews*, *100*(2), 805-868.

Ferraris, M., Cassel, J.-C., de Vasconcelos, A. P., Stephan, A., & Quilichini, P. P. (2021). The nucleus reuniens, a thalamic relay for cortico-hippocampal interaction in recent and remote memory consolidation. *Neuroscience & Biobehavioral Reviews*, *125*, 339-354.

Fogel, S., Albouy, G., King, B. R., Lungu, O., Vien, C., Bore, A., Pinsard, B., Benali, H., Carrier, J., & Doyon, J. (2017). Reactivation or transformation? Motor memory consolidation associated with cerebral activation time-locked to sleep spindles. *PloS one*, *12*(4), e0174755.

Fonov, V., AC Evans, RC McKinstry, CR Almli, and DL Collins. (2009). Unbiased Nonlinear Average Age-Appropriate Brain Templates from Birth to Adulthood. *NeuroImage*, *47*, S102.

Friston, K. J., Buechel, C., Fink, G. R., Morris, J., Rolls, E., & Dolan, R. J. (1997). Psychophysiological and modulatory interactions in neuroimaging. *NeuroImage*, *6*(3), 218-229.

Gaillard, J. M., & Blois, R. (1981). Spindle density in sleep of normal subjects. *Sleep*, *4*(4), 385-391.

Göldi, M., van Poppel, E. A. M., Rasch, B., & Schreiner, T. (2019). Increased neuronal signatures of targeted memory reactivation during slow-wave up states. *Scientific reports*, *9*(1), 2715.

Gorgolewski, K., Burns, C. D., Madison, C., Clark, D., Halchenko, Y. O., Waskom, M. L., & Ghosh, S. S. (2011). Nipype: a flexible, lightweight and extensible neuroimaging data processing framework in python. *Frontiers in neuroinformatics*, *5*, 13.

Gramfort, A., Luessi, M., Larson, E., Engemann, D. A., Strohmeier, D., Brodbeck, C., Goj, R., Jas, M., Brooks, T., Parkkonen, L., & Hamalainen, M. (2013). MEG and EEG data analysis with MNE-Python. *Frontiers in neuroscience*, *7*, 267.

Greve, D. N., & Fischl, B. (2009). Accurate and robust brain image alignment using boundary-based registration. *NeuroImage*, *48*(1), 63-72.

Hahn, M. A., Heib, D., Schabus, M., Hoedlmoser, K., & Helfrich, R. F. (2020). Slow oscillation-spindle coupling predicts enhanced memory formation from childhood to adolescence. *Elife*, *9*, e53730.

Halassa, M. M., Siegle, J. H., Ritt, J. T., Ting, J. T., Feng, G., & Moore, C. I. (2011). Selective optical drive of thalamic reticular nucleus generates thalamic bursts and cortical spindles. *Nature Neuroscience*, *14*(9), 1118-1120.

Hale, J. R., White, T. P., Mayhew, S. D., Wilson, R. S., Rollings, D. T., Khalsa, S., Arvanitis, T. N., & Bagshaw, A. P. (2016). Altered thalamocortical and intra-thalamic functional connectivity during light sleep compared with wake. *NeuroImage*, *125*, 657-667.

Helfrich, R. F., Lendner, J. D., Mander, B. A., Guillen, H., Paff, M., Mnatsakanyan, L., Vadera, S., Walker, M. P., Lin, J. J., & Knight, R. T. (2019). Bidirectional prefrontal-hippocampal dynamics organize information transfer during sleep in humans. *Nature Communications*, *10*(1), 3572.

Helfrich, R. F., Mander, B. A., Jagust, W. J., Knight, R. T., & Walker, M. P. (2018). Old brains come uncoupled in sleep: slow wave-spindle synchrony, brain atrophy, and forgetting. *Neuron*, *97*(1), 221-230. e224.

Horovitz, S. G., Fukunaga, M., de Zwart, J. A., van Gelderen, P., Fulton, S. C., Balkin, T. J., & Duyn, J. H. (2008). Low frequency BOLD fluctuations during resting wakefulness and light sleep: A simultaneous EEG-fMRI study. *Human brain mapping*, *29*(6), 671-682.

Hu, X., Cheng, L. Y., Chiu, M. H., & Paller, K. A. (2020). Promoting memory consolidation during
sleep: A meta-analysis of targeted memory reactivation. *Psychological bulletin*, *146*(3), 218.
Huang, Q., Xiao, Z., Yu, Q., Luo, Y., Xu, J., Qu, Y., Dolan, R., Behrens, T., & Liu, Y. (2024). Replay-
triggered brain-wide activation in humans. *Nature Communications*, *15*(1), 7185.
Ilhan-Bayrakcı, M., Cabral-Calderin, Y., Bergmann, T. O., Tüscher, O., & Stroh, A. (2022). Individual
slow wave events give rise to macroscopic fMRI signatures and drive the strength of the BOLD
signal in human resting-state EEG-fMRI recordings. *Cerebral Cortex*, *32*(21), 4782-4796.
Jenkinson, M., Bannister, P., Brady, M., & Smith, S. (2002). Improved optimization for the robust and
accurate linear registration and motion correction of brain images. *NeuroImage*, *17*(2), 825-841.
Jenkinson, M., & Smith, S. (2001). A global optimisation method for robust affine registration of brain
images. *Medical image analysis*, *5*(2), 143-156.
Joo, H. R., & Frank, L. M. (2018). The hippocampal sharp wave–ripple in memory retrieval for
immediate use and consolidation. *Nature Reviews Neuroscience*, *19*(12), 744-757.
Lanczos, C. (1964). Evaluation of Noisy Data. *Journal of the Society for Industrial and Applied
Mathematics Series B Numerical Analysis*, *1*(1), 76–85.
Latchoumane, C.-F. V., Ngo, H.-V. V., Born, J., & Shin, H.-S. (2017). Thalamic spindles promote
memory formation during sleep through triple phase-locking of cortical, thalamic, and
hippocampal rhythms. *Neuron*, *95*(2), 424-435. e426.
Laufs, H. (2008). Endogenous brain oscillations and related networks detected by surface EEG-
combined fMRI. *Human brain mapping*, *29*(7), 762-769.
Laufs, H., Walker, M. C., & Lund, T. E. (2007). 'Brain activation and hypothalamic functional
connectivity during human non-rapid eye movement sleep: an EEG/fMRI study'—its
limitations and an alternative approach. *Brain*, *130*(7), e75.
Liu, Y., Nour, M. M., Schuck, N. W., Behrens, T. E., & Dolan, R. J. (2022). Decoding cognition from
spontaneous neural activity. *Nature Reviews Neuroscience*, *23*(4), 204-214.
Maingret, N., Girardeau, G., Todorova, R., Goutierre, M., & Zugaro, M. (2016). Hippocampo-cortical
coupling mediates memory consolidation during sleep. *Nature Neuroscience*, *19*(7), 959-964.
Mak-McCully, R. A., Rolland, M., Sargsyan, A., Gonzalez, C., Magnin, M., Chauvel, P., Rey, M.,
Bastuji, H., & Halgren, E. (2017). Coordination of cortical and thalamic activity during non-
REM sleep in humans. *Nature Communications*, *8*(1), 15499.
Margulies, D. S., Ghosh, S. S., Goulas, A., Falkiewicz, M., Huntenburg, J. M., Langs, G., Bezgin, G.,
Eickhoff, S. B., Castellanos, F. X., & Petrides, M. (2016). Situating the default-mode network
along a principal gradient of macroscale cortical organization. *Proceedings of the National
Academy of Sciences*, *113*(44), 12574-12579.
Massimini, M., Huber, R., Ferrarelli, F., Hill, S., & Tononi, G. (2004). The sleep slow oscillation as a
traveling wave. *Journal of Neuroscience*, *24*(31), 6862-6870.
Moehlman, T. M., de Zwart, J. A., Chappel-Farley, M. G., Liu, X., McClain, I. B., Chang, C.,
Mandelkow, H., Özbay, P. S., Johnson, N. L., & Bieber, R. E. (2019). All-night functional
magnetic resonance imaging sleep studies. *Journal of neuroscience methods*, *316*, 83-98.
Molle, M., Bergmann, T. O., Marshall, L., & Born, J. (2011). Fast and slow spindles during the sleep
slow oscillation: disparate coalescence and engagement in memory processing. *Sleep*, *34*(10),
1411-1421.
Ngo, H.-V., Fell, J., & Staresina, B. (2020). Sleep spindles mediate hippocampal-neocortical coupling
during long-duration ripples. *Elife*, *9*, e57011.
Ngo, H. V., Martinetz, T., Born, J., & Molle, M. (2013). Auditory closed-loop stimulation of the sleep
slow oscillation enhances memory. *Neuron*, *78*(3), 545-553.
Nir, Y., Staba, R. J., Andrillon, T., Vyazovskiy, V. V., Cirelli, C., Fried, I., & Tononi, G. (2011). Regional
slow waves and spindles in human sleep. *Neuron*, *70*(1), 153-169.
Oostenveld, R., Fries, P., Maris, E., & Schoffelen, J. M. (2011). FieldTrip: Open source software for
advanced analysis of MEG, EEG, and invasive electrophysiological data. *Computational
intelligence and neuroscience*, *2011*, 156869.
Oudiette, D., & Paller, K. A. (2013). Upgrading the sleeping brain with targeted memory reactivation.
*Trends in Cognitive Sciences*, *17*(3), 142-149.
Park, S. A., Miller, D. S., & Boorman, E. D. (2021). Inferences on a multidimensional social hierarchy
use a grid-like code. *Nature Neuroscience*, *24*(9), 1292-1301.

Picchioni, D., Horovitz, S. G., Fukunaga, M., Carr, W. S., Meltzer, J. A., Balkin, T. J., Duyn, J. H., & Braun, A. R. (2011). Infraslow EEG oscillations organize large-scale cortical–subcortical interactions during sleep: a combined EEG/fMRI study. *Brain research*, *1374*, 63-72.

Power, J. D., Mitra, A., Laumann, T. O., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2014). Methods to detect, characterize, and remove motion artifact in resting state fMRI. *NeuroImage*, *84*, 320-341.

Rasch, B. r., Büchel, C., Gais, S., & Born, J. (2007). Odor cues during slow-wave sleep prompt declarative memory consolidation. *Science*, *315*(5817), 1426-1429.

Schabus, M., Dang-Vu, T. T., Albouy, G., Balteau, E., Boly, M., Carrier, J., Darsaud, A., Degueldre, C., Desseilles, M., & Gais, S. (2007). Hemodynamic cerebral correlates of sleep spindles during human non-rapid eye movement sleep. *Proceedings of the National Academy of Sciences*, *104*(32), 13164-13169.

Schreiner, T., Kaufmann, E., Noachtar, S., Mehrkens, J.-H., & Staudigl, T. (2022). The human thalamus orchestrates neocortical oscillations during NREM sleep. *Nature communications*, *13*(1), 5231.

Schreiner, T., Petzka, M., Staudigl, T., & Staresina, B. P. (2021). Endogenous memory reactivation during sleep in humans is clocked by slow oscillation-spindle complexes. *Nature Communications*, *12*(1), 3112.

Siefert, E. M., Uppuluri, S., Mu, J., Tandoc, M. C., Antony, J. W., & Schapiro, A. C. (2024). Memory reactivation during sleep does not act holistically on object memory. *Journal of Neuroscience*, *44*(24).

Singh, D., Norman, K. A., & Schapiro, A. C. (2022). A model of autonomous interactions between hippocampus and neocortex driving sleep-dependent memory consolidation. *Proceedings of the National Academy of Sciences*, *119*(44), e2123432119.

Spoormaker, V. I., Schröter, M. S., Gleiser, P. M., Andrade, K. C., Dresler, M., Wehrle, R., Sämann, P. G., & Czisch, M. (2010). Development of a large-scale functional brain network during human non-rapid eye movement sleep. *Journal of Neuroscience*, *30*(34), 11379-11387.

Staresina, B. P., Bergmann, T. O., Bonnefond, M., van der Meij, R., Jensen, O., Deuker, L., Elger, C. E., Axmacher, N., & Fell, J. (2015). Hierarchical nesting of slow oscillations, spindles and ripples in the human hippocampus during sleep. *Nature Neuroscience*, *18*(11), 1679-1686.

Staresina, B. P., Niediek, J., Borger, V., Surges, R., & Mormann, F. (2023). How coupled slow oscillations, spindles and ripples coordinate neuronal processing and communication during human sleep. *Nature Neuroscience*, 1-9.

Steriade, M., McCormick, D. A., & Sejnowski, T. J. (1993). Thalamocortical oscillations in the sleeping and aroused brain. *Science*, *262*(5134), 679-685.

Tagliazucchi, E., von Wegner, F., Morzelewski, A., Brodbeck, V., Jahnke, K., & Laufs, H. (2013). Breakdown of long-range temporal dependence in default mode and attention networks during deep sleep. *Proceedings of the National Academy of Sciences*, *110*(38), 15419-15424.

Tustison, N. J., Avants, B. B., Cook, P. A., Zheng, Y., Egan, A., Yushkevich, P. A., & Gee, J. C. (2010). N4ITK: improved N3 bias correction. *IEEE transactions on medical imaging*, *29*(6), 1310-1320.

Vallat, R., & Walker, M. P. (2021). An open-source, high-performance tool for automated sleep staging. *Elife*, *10*.

Xia, T., Antony, J. W., Paller, K. A., & Hu, X. (2023). Targeted memory reactivation during sleep influences social bias as a function of slow-oscillation phase and delta power. *Psychophysiology*, *60*(5), e14224.

Yarkoni, T., Poldrack, R. A., Nichols, T. E., Van Essen, D. C., & Wager, T. D. (2011). Large-scale automated synthesis of human functional neuroimaging data. *Nature methods*, *8*(8), 665-670.

Yeshurun, Y., Nguyen, M., & Hasson, U. (2021). The default mode network: where the idiosyncratic self meets the shared social world. *Nature Reviews Neuroscience*, 1-12.

Zhang, Y., Brady, M., & Smith, S. (2001). Segmentation of brain MR images through a hidden Markov random field model and the expectation-maximization algorithm. *IEEE transactions on medical imaging*, *20*(1), 45-57.

**Acknowledgment**

Author contributions: Conceptualization, Y.L., H.W., J.G.; Investigation, H.W., Q.Z., J.Z., Y.L., J.G.; Writing – Original Draft, H.W., Y.L.; Writing – Review & Editing, H.W., Y.L., J.Z., Q.Z., J.G. Funding: This study is supported by the National Science and Technology Innovation 2030 Major Program (2022ZD0205500), the National Natural Science Foundation of China (32271093), the Beijing Municipal Science and Technology Commission (Z230010, L222033), Beijing United Imaging Research Institute of Intelligent Imaging Foundation (CRIBJZD202101), the Open Research Fund of the National Center for Protein Sciences at Peking University, and the Fundamental Research Funds for the Central Universities.

**Competing interests**

The authors declare that they have no competing interests.

**Data and materials availability**

The data that support the findings can be provided by the corresponding author Y.L. pending scientific review and a completed material transfer agreement. Requests for the data should be submitted to: https://datadryad.org/stash/dataset/doi:10.5061/dryad.2fqz612x0. The analysis code will be publicly available on https://gitlab.com/liu_lab/eeg-fmri-sleep upon publication.

**Supplementary Materials**

**Fig. S1 | Removal of MRI gradient noise from simultaneous collected EEG data. a**, Time series of both raw and preprocessed EEG data. The top row depicts the raw EEG data, which contains noise primarily from the MRI gradient magnetic field and electrocardiographic artifacts. The bottom row showcases the preprocessed EEG data (detailed in Methods). **b**, Power spectral density of the raw and preprocessed EEG data estimated by the fast Fourier transform. The raw EEG data is shown in the top row, while the preprocessed EEG data is in the bottom row. **c**, Time-frequency spectrogram of the raw and preprocessed EEG data, by the short-time Fourier transform. The top row represents the raw EEG data, and the bottom row displays the preprocessed EEG data.

**Fig. S2 | ERPs of SOs and spindles coupling during different sleep stages across all 107 subjects.**
**a.** ERP of SOs in different sleep stages using the broadband (0.1–30 Hz) EEG data. We align the trough of the DOWN-state of each SO at time zero (see Methods for details). The orange line represents the SO ERP in the N1 stage, the black line represents the SO ERP in the N2&N3 stage, and the green line represents the SO ERP in the REM stage. **b.** ERP of spindles in different sleep stages using the broadband (0.1–30 Hz) EEG data. We align the peak of each spindle at time zero (see Methods for details). The color scheme is the same as in panel a.

Fig. S3 | ERP and time-frequency patterns of SO-spindle coupling in the N1 stage. The averaged temporal frequency pattern and ERP across all instances of SO-spindle coupling, computed over all subjects, following the same procedure as in Fig. 2a, but for N1 stage.

**Fig. S4 | ERP and time-frequency patterns of SO-spindle coupling in the REM stage.** The averaged

temporal frequency pattern and ERP across all instances of SO-spindle coupling, computed over all

subjects, again following the same procedure as in **Fig. 2a**, but for REM stage.

**Fig. S5 | Brain-wide activity between SO UP-state (peak) and DOWN-state (trough). a**, Brain activity with SO DOWN-state (trough) modelled as event onset, whole-brain FWE corrected at the cluster level ($p < 0.05$) with a cluster-forming voxel threshold of $p_{unc.} < 0.001$. **b**, Brain activity with SO UP-state (peak) modelled as event onset. **c**, Differences in brain activity corresponding to SO UP-state and SO DOWN-state. The whole-brain results were displayed at an uncorrected threshold of $p < 0.01$ for visualization purpose only. No brain region was found significant in this contrast.

**Fig. S6 | Influence of the percentile threshold for SO detection on hippocampal activation (ROI)**

**during SO-spindle coupling.** We changed the percentile threshold for SO event detection in the EEG

data analysis and then reconstructed the GLM design matrix based on the SO events detected at each

threshold. The brain-wide activation pattern of SO-spindle couplings in the N2/3 stage was extracted

using the same method as shown in **Fig. 3**. The gray horizontal line represents the significant range

(71%–80%). * $p < 0.05$.

**Fig. S7 | Functional decoding using the ROI association method in Neurosynth. a**, Decoding results using positive activation. **b**, Decoding results using negative activation. Each row corresponds to the brain-wide activation patterns for sleep rhythms shown in **Fig. 3b-d**, while each column corresponds to topics in the Neurosynth database (detailed in Methods). Only topics with a decoded significance level of $p < 0.05$ are displayed.

| Descriptive statistics of demographic and sleep | Mean ± SE |
| --- | --- |
| | (n = 107) |
| Age | 22.40 ± 0.34 |
| Total Recording time (min) | 180.60 ± 8.40 |
| Wake time (min) | 84.60 ± 5.00 |
| Total Sleep time (min) | 95.98 ± 5.67 |
| N1 time (min) | 13.99 ± 1.19 |
| N2 time (min) | 56.21 ± 4.05 |
| N3 time (min) | 16.53 ± 1.35 |
| REM time (min) | 9.27 ± 0.86 |
| N1 ratio (%) | 14.57 ± 1.06 |
| N2 ratio (%) | 58.55 ± 1.70 |
| N3 ratio (%) | 17.22 ± 1.21 |
| REM ratio (%) | 9.66 ± 0.90 |
| Sleep Onset Latency (min) | 9.75 ± 0.85 |
| N2 latency (min) | 14.58 ± 1.15 |
| N3 latency (min) | 33.65 ± 3.02 |
| Sleep Efficiency (%) | 52.96 ± 1.60 |

**Table S1 | Descriptive results of demographic information and sleep characteristics**. Note: The total recorded time is equal to the awake time plus the total sleep time. The sleep onset latency is the time taken to reach the first sleep epoch. The Sleep Efficiency is the ratio of actual sleep time to total recording time.

| Subject ID | N1 Sleep Stage | | | | | | |
|---|---|---|---|---|---|---|---|
| | Time (min) | Number of SO | Number of spindle | Number of coupling | Density of SO (times/min) | Density of spindle (times/min) | Density of coupling (times/min) |
| 1 | 8.38 | 60 | 31 | 14 | 7.16 | 3.70 | 1.67 |
| 2 | 5.21 | 49 | 22 | 14 | 9.40 | 4.22 | 2.69 |
| 3 | 3.24 | 28 | 15 | 9 | 8.63 | 4.62 | 2.77 |
| 4 | 16.61 | 197 | 60 | 55 | 11.86 | 3.61 | 3.31 |
| 5 | 3.38 | 40 | 10 | 5 | 11.85 | 2.96 | 1.48 |
| 6 | 9.74 | 104 | 35 | 28 | 10.67 | 3.59 | 2.87 |
| 7 | 8.84 | 46 | 32 | 20 | 5.20 | 3.62 | 2.26 |
| 8 | 19.51 | 147 | 38 | 13 | 7.53 | 1.95 | 0.67 |
| 9 | 2.28 | 17 | 2 | 0 | 7.47 | 0.88 | 0.00 |
| 10 | 25.71 | 234 | 95 | 57 | 9.10 | 3.70 | 2.22 |
| 11 | 8.04 | 35 | 18 | 7 | 4.35 | 2.24 | 0.87 |
| 12 | 4.54 | 10 | 6 | 0 | 2.20 | 1.32 | 0.00 |
| 13 | 4.91 | 36 | 23 | 15 | 7.33 | 4.68 | 3.05 |
| 14 | 3.41 | 22 | 14 | 9 | 6.45 | 4.11 | 2.64 |
| 15 | 4.91 | 52 | 8 | 5 | 10.59 | 1.63 | 1.02 |
| 16 | 21.98 | 58 | 34 | 17 | 2.64 | 1.55 | 0.77 |
| 17 | 1.31 | 0 | 0 | 0 | 0.00 | 0.00 | 0.00 |
| 18 | 0.91 | 0 | 0 | 0 | 0.00 | 0.00 | 0.00 |
| 19 | 2.54 | 11 | 4 | 4 | 4.33 | 1.57 | 1.57 |
| 20 | 23.71 | 266 | 100 | 54 | 11.22 | 4.22 | 2.28 |
| 21 | 35.81 | 322 | 151 | 109 | 8.99 | 4.22 | 3.04 |
| 22 | 25.58 | 146 | 61 | 41 | 5.71 | 2.38 | 1.60 |
| 23 | 3.41 | 21 | 5 | 3 | 6.16 | 1.47 | 0.88 |
| 24 | 20.68 | 224 | 73 | 59 | 10.83 | 3.53 | 2.85 |
| 25 | 47.74 | 543 | 224 | 124 | 11.37 | 4.69 | 2.60 |

| 26 | 20.91 | 130 | 63  | 42  | 6.22  | 3.01 | 2.01 |
|----|-------|-----|-----|-----|-------|------|------|
| 27 | 7.91  | 92  | 46  | 21  | 11.63 | 5.82 | 2.65 |
| 28 | 10.71 | 94  | 49  | 38  | 8.78  | 4.58 | 3.55 |
| 29 | 22.71 | 229 | 100 | 63  | 10.08 | 4.40 | 2.77 |
| 30 | 2.84  | 28  | 12  | 7   | 9.85  | 4.22 | 2.46 |
| 31 | 7.41  | 60  | 20  | 17  | 8.10  | 2.70 | 2.29 |
| 32 | 0.98  | 6   | 2   | 1   | 6.14  | 2.05 | 1.02 |
| 33 | 3.21  | 38  | 18  | 12  | 11.84 | 5.61 | 3.74 |
| 34 | 7.81  | 67  | 29  | 16  | 8.58  | 3.71 | 2.05 |
| 35 | 4.44  | 31  | 13  | 8   | 6.98  | 2.93 | 1.80 |
| 36 | 41.34 | 423 | 145 | 107 | 10.23 | 3.51 | 2.59 |
| 37 | 5.31  | 35  | 18  | 8   | 6.59  | 3.39 | 1.51 |
| 38 | 13.34 | 140 | 53  | 46  | 10.49 | 3.97 | 3.45 |
| 39 | 20.34 | 217 | 61  | 40  | 10.67 | 3.00 | 1.97 |
| 40 | 5.74  | 51  | 19  | 14  | 8.88  | 3.31 | 2.44 |
| 41 | 8.11  | 113 | 24  | 20  | 13.93 | 2.96 | 2.47 |
| 42 | 25.68 | 271 | 108 | 70  | 10.55 | 4.21 | 2.73 |
| 43 | 23.21 | 204 | 58  | 35  | 8.79  | 2.50 | 1.51 |
| 44 | 16.94 | 195 | 69  | 39  | 11.51 | 4.07 | 2.30 |
| 45 | 6.31  | 58  | 32  | 18  | 9.19  | 5.07 | 2.85 |
| 46 | 7.24  | 80  | 23  | 13  | 11.04 | 3.18 | 1.79 |
| 47 | 9.14  | 78  | 36  | 24  | 8.53  | 3.94 | 2.62 |
| 48 | 35.71 | 361 | 187 | 106 | 10.11 | 5.24 | 2.97 |
| 49 | 7.01  | 71  | 23  | 19  | 10.13 | 3.28 | 2.71 |
| 50 | 27.18 | 174 | 71  | 52  | 6.40  | 2.61 | 1.91 |
| 51 | 13.38 | 129 | 50  | 34  | 9.64  | 3.74 | 2.54 |
| 52 | 17.54 | 71  | 28  | 5   | 4.05  | 1.60 | 0.29 |
| 53 | 24.31 | 269 | 104 | 83  | 11.07 | 4.28 | 3.41 |
| 54 | 11.54 | 131 | 60  | 42  | 11.35 | 5.20 | 3.64 |
| 55 | 10.14 | 93  | 27  | 14  | 9.17  | 2.66 | 1.38 |
| 56 | 4.41  | 49  | 21  | 16  | 11.11 | 4.76 | 3.63 |

| 57 | 5.01 | 34 | 21 | 11 | 6.79 | 4.19 | 2.20 |
|----|------|-----|-----|-----|-------|------|------|
| 58 | 6.01 | 56 | 28 | 13 | 9.32 | 4.66 | 2.16 |
| 59 | 37.38 | 404 | 139 | 103 | 10.81 | 3.72 | 2.76 |
| 60 | 7.08 | 68 | 27 | 13 | 9.61 | 3.82 | 1.84 |
| 61 | 8.74 | 86 | 43 | 23 | 9.84 | 4.92 | 2.63 |
| 62 | 17.61 | 181 | 56 | 48 | 10.28 | 3.18 | 2.73 |
| 63 | 48.61 | 388 | 187 | 117 | 7.98 | 3.85 | 2.41 |
| 64 | 9.74 | 51 | 16 | 9 | 5.23 | 1.64 | 0.92 |
| 65 | 22.21 | 215 | 96 | 80 | 9.68 | 4.32 | 3.60 |
| 66 | 36.78 | 312 | 149 | 98 | 8.48 | 4.05 | 2.66 |
| 67 | 33.51 | 263 | 110 | 67 | 7.85 | 3.28 | 2.00 |
| 68 | 10.71 | 112 | 36 | 19 | 10.46 | 3.36 | 1.77 |
| 69 | 58.34 | 608 | 213 | 161 | 10.42 | 3.65 | 2.76 |
| 70 | 14.21 | 97 | 51 | 31 | 6.83 | 3.59 | 2.18 |
| 71 | 21.98 | 170 | 85 | 61 | 7.74 | 3.87 | 2.78 |
| 72 | 55.31 | 600 | 203 | 120 | 10.85 | 3.67 | 2.17 |
| 73 | 24.61 | 207 | 89 | 81 | 8.41 | 3.62 | 3.29 |
| 74 | 23.31 | 252 | 96 | 81 | 10.81 | 4.12 | 3.47 |
| 75 | 4.74 | 34 | 17 | 9 | 7.17 | 3.58 | 1.90 |
| 76 | 37.11 | 360 | 145 | 97 | 9.70 | 3.91 | 2.61 |
| 77 | 14.41 | 146 | 64 | 53 | 10.13 | 4.44 | 3.68 |
| 78 | 12.21 | 134 | 54 | 33 | 10.97 | 4.42 | 2.70 |
| 79 | 6.18 | 66 | 14 | 8 | 10.69 | 2.27 | 1.30 |
| 80 | 8.18 | 76 | 27 | 15 | 9.29 | 3.30 | 1.83 |
| 81 | 2.84 | 29 | 8 | 3 | 10.20 | 2.81 | 1.06 |
| 82 | 7.41 | 75 | 29 | 21 | 10.12 | 3.91 | 2.83 |
| 83 | 1.91 | 18 | 7 | 3 | 9.42 | 3.66 | 1.57 |
| 84 | 2.14 | 21 | 7 | 3 | 9.80 | 3.27 | 1.40 |
| 85 | 10.38 | 47 | 37 | 12 | 4.53 | 3.57 | 1.16 |
| 86 | 10.34 | 89 | 53 | 28 | 8.60 | 5.12 | 2.71 |
| 87 | 5.98 | 60 | 19 | 11 | 10.04 | 3.18 | 1.84 |

| | | | | | | |
|------|-------|--------|-------|-------|-------|------|------|
| 88 | 0.41 | 0 | 0 | 0 | 0.00 | 0.00 | 0.00 |
| 89 | 10.91 | 113 | 46 | 29 | 10.36 | 4.22 | 2.66 |
| 90 | 9.54 | 100 | 37 | 28 | 10.48 | 3.88 | 2.93 |
| 91 | 4.18 | 49 | 18 | 12 | 11.73 | 4.31 | 2.87 |
| 92 | 2.14 | 19 | 5 | 2 | 8.86 | 2.33 | 0.93 |
| 93 | 10.11 | 116 | 38 | 38 | 11.47 | 3.76 | 3.76 |
| 94 | 1.91 | 3 | 0 | 0 | 1.57 | 0.00 | 0.00 |
| 95 | 8.68 | 76 | 39 | 22 | 8.76 | 4.49 | 2.54 |
| 96 | 30.71 | 344 | 123 | 87 | 11.20 | 4.01 | 2.83 |
| 97 | 11.71 | 135 | 40 | 36 | 11.53 | 3.42 | 3.07 |
| 98 | 11.91 | 108 | 55 | 37 | 9.07 | 4.62 | 3.11 |
| 99 | 16.41 | 133 | 84 | 51 | 8.10 | 5.12 | 3.11 |
| 100 | 7.34 | 80 | 26 | 15 | 10.89 | 3.54 | 2.04 |
| 101 | 6.41 | 39 | 24 | 11 | 6.08 | 3.74 | 1.72 |
| 102 | 9.24 | 100 | 45 | 31 | 10.82 | 4.87 | 3.35 |
| 103 | 3.34 | 15 | 8 | 3 | 4.49 | 2.39 | 0.90 |
| 104 | 13.41 | 142 | 64 | 45 | 10.59 | 4.77 | 3.36 |
| 105 | 10.58 | 105 | 44 | 39 | 9.93 | 4.16 | 3.69 |
| 106 | 6.48 | 63 | 25 | 25 | 9.73 | 3.86 | 3.86 |
| 107 | 21.51 | 217 | 75 | 50 | 10.09 | 3.49 | 2.32 |
| **Mean** | 13.99 | 129.64 | 51.89 | 34.44 | 8.70 | 3.48 | 2.23 |
| **SE** | 1.19 | 12.18 | 4.78 | 3.27 | 0.26 | 0.12 | 0.09 |

**Table S2 | Statistics of sleep duration, SO, spindle and coupling event numbers and densities for all 107 subjects during N1 stage.**

| Subject ID | N2&N3 Sleep Stage | | | | | | |
|---|---|---|---|---|---|---|---|
| | Time (min) | Number of SO | Number of spindle | Number of coupling | Density of SO (times/min) | Density of spindle (times/min) | Density of coupling (times/min) |
| 1 | 89.00 | 820 | 401 | 242 | 9.21 | 4.51 | 2.72 |
| 2 | 125.00 | 1417 | 659 | 298 | 11.34 | 5.27 | 2.38 |
| 3 | 22.50 | 225 | 115 | 75 | 10.00 | 5.11 | 3.33 |
| 4 | 87.00 | 1059 | 326 | 175 | 12.17 | 3.75 | 2.01 |
| 5 | 44.50 | 425 | 190 | 115 | 9.55 | 4.27 | 2.58 |
| 6 | 21.00 | 222 | 78 | 57 | 10.57 | 3.71 | 2.71 |
| 7 | 42.50 | 185 | 160 | 91 | 4.35 | 3.76 | 2.14 |
| 8 | 167.50 | 1865 | 882 | 502 | 11.13 | 5.27 | 3.00 |
| 9 | 25.50 | 112 | 28 | 24 | 4.39 | 1.10 | 0.94 |
| 10 | 65.00 | 715 | 322 | 197 | 11.00 | 4.95 | 3.03 |
| 11 | 69.00 | 511 | 302 | 155 | 7.41 | 4.38 | 2.25 |
| 12 | 62.50 | 413 | 316 | 162 | 6.61 | 5.06 | 2.59 |
| 13 | 89.00 | 826 | 449 | 283 | 9.28 | 5.04 | 3.18 |
| 14 | 41.00 | 390 | 191 | 93 | 9.51 | 4.66 | 2.27 |
| 15 | 105.00 | 359 | 96 | 51 | 3.42 | 0.91 | 0.49 |
| 16 | 184.50 | 708 | 425 | 228 | 3.84 | 2.30 | 1.24 |
| 17 | 97.50 | 395 | 306 | 155 | 4.05 | 3.14 | 1.59 |
| 18 | 23.50 | 212 | 86 | 75 | 9.02 | 3.66 | 3.19 |
| 19 | 17.50 | 137 | 48 | 39 | 7.83 | 2.74 | 2.23 |
| 20 | 102.00 | 1196 | 462 | 290 | 11.73 | 4.53 | 2.84 |
| 21 | 139.00 | 1236 | 671 | 359 | 8.89 | 4.83 | 2.58 |
| 22 | 56.50 | 684 | 284 | 166 | 12.11 | 5.03 | 2.94 |
| 23 | 43.00 | 508 | 182 | 117 | 11.81 | 4.23 | 2.72 |
| 24 | 75.50 | 872 | 313 | 194 | 11.55 | 4.15 | 2.57 |
| 25 | 136.00 | 1616 | 792 | 408 | 11.88 | 5.82 | 3.00 |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| 26 | 29.00 | 190 | 118 | 63 | 6.55 | 4.07 | 2.17 |
| 27 | 73.50 | 849 | 408 | 229 | 11.55 | 5.55 | 3.12 |
| 28 | 249.50 | 2355 | 1170 | 806 | 9.44 | 4.69 | 3.23 |
| 29 | 54.00 | 603 | 224 | 137 | 11.17 | 4.15 | 2.54 |
| 30 | 7.50 | 34 | 16 | 13 | 4.53 | 2.13 | 1.73 |
| 31 | 30.50 | 269 | 139 | 91 | 8.82 | 4.56 | 2.98 |
| 32 | 53.00 | 550 | 258 | 128 | 10.38 | 4.87 | 2.42 |
| 33 | 24.00 | 290 | 128 | 68 | 12.08 | 5.33 | 2.83 |
| 34 | 44.00 | 401 | 182 | 111 | 9.11 | 4.14 | 2.52 |
| 35 | 14.00 | 120 | 62 | 36 | 8.57 | 4.43 | 2.57 |
| 36 | 177.50 | 1925 | 782 | 518 | 10.85 | 4.41 | 2.92 |
| 37 | 113.50 | 1290 | 569 | 345 | 11.37 | 5.01 | 3.04 |
| 38 | 113.50 | 1286 | 570 | 303 | 11.33 | 5.02 | 2.67 |
| 39 | 72.50 | 660 | 230 | 134 | 9.10 | 3.17 | 1.85 |
| 40 | 119.00 | 827 | 284 | 176 | 6.95 | 2.39 | 1.48 |
| 41 | 43.00 | 438 | 176 | 101 | 10.19 | 4.09 | 2.35 |
| 42 | 79.50 | 917 | 324 | 154 | 11.53 | 4.08 | 1.94 |
| 43 | 197.00 | 1166 | 559 | 369 | 5.92 | 2.84 | 1.87 |
| 44 | 93.00 | 1136 | 362 | 205 | 12.22 | 3.89 | 2.20 |
| 45 | 159.00 | 696 | 411 | 227 | 4.38 | 2.58 | 1.43 |
| 46 | 53.00 | 575 | 165 | 102 | 10.85 | 3.11 | 1.92 |
| 47 | 143.00 | 596 | 317 | 207 | 4.17 | 2.22 | 1.45 |
| 48 | 149.00 | 1514 | 850 | 515 | 10.16 | 5.70 | 3.46 |
| 49 | 27.00 | 312 | 102 | 61 | 11.56 | 3.78 | 2.26 |
| 50 | 110.00 | 707 | 339 | 238 | 6.43 | 3.08 | 2.16 |
| 51 | 42.50 | 439 | 144 | 71 | 10.33 | 3.39 | 1.67 |
| 52 | 52.00 | 275 | 171 | 74 | 5.29 | 3.29 | 1.42 |
| 53 | 34.00 | 343 | 136 | 91 | 10.09 | 4.00 | 2.68 |
| 54 | 69.00 | 820 | 359 | 198 | 11.88 | 5.20 | 2.87 |
| 55 | 81.00 | 940 | 327 | 197 | 11.60 | 4.04 | 2.43 |
| 56 | 39.50 | 448 | 209 | 124 | 11.34 | 5.29 | 3.14 |

| | | | | | | |
|---|---|---|---|---|---|---|
| 57 | 32.50 | 301 | 140 | 76 | 9.26 | 4.31 | 2.34 |
| 58 | 51.50 | 525 | 218 | 119 | 10.19 | 4.23 | 2.31 |
| 59 | 55.50 | 561 | 212 | 150 | 10.11 | 3.82 | 2.70 |
| 60 | 114.00 | 1188 | 485 | 279 | 10.42 | 4.25 | 2.45 |
| 61 | 59.50 | 713 | 373 | 194 | 11.98 | 6.27 | 3.26 |
| 62 | 70.50 | 795 | 250 | 204 | 11.28 | 3.55 | 2.89 |
| 63 | 105.00 | 693 | 432 | 231 | 6.60 | 4.11 | 2.20 |
| 64 | 57.00 | 653 | 347 | 172 | 11.46 | 6.09 | 3.02 |
| 65 | 60.50 | 577 | 268 | 151 | 9.54 | 4.43 | 2.50 |
| 66 | 120.50 | 1203 | 622 | 390 | 9.98 | 5.16 | 3.24 |
| 67 | 199.00 | 915 | 521 | 274 | 4.60 | 2.62 | 1.38 |
| 68 | 50.00 | 595 | 183 | 118 | 11.90 | 3.66 | 2.36 |
| 69 | 127.50 | 1458 | 592 | 366 | 11.44 | 4.64 | 2.87 |
| 70 | 142.50 | 323 | 89 | 46 | 2.27 | 0.62 | 0.32 |
| 71 | 123.50 | 1261 | 460 | 324 | 10.21 | 3.72 | 2.62 |
| 72 | 65.00 | 195 | 80 | 55 | 3.00 | 1.23 | 0.85 |
| 73 | 92.00 | 1035 | 413 | 238 | 11.25 | 4.49 | 2.59 |
| 74 | 128.00 | 1500 | 638 | 307 | 11.72 | 4.98 | 2.40 |
| 75 | 31.50 | 289 | 153 | 81 | 9.17 | 4.86 | 2.57 |
| 76 | 149.00 | 1302 | 631 | 331 | 8.74 | 4.23 | 2.22 |
| 77 | 57.00 | 605 | 271 | 166 | 10.61 | 4.75 | 2.91 |
| 78 | 94.50 | 1097 | 448 | 247 | 11.61 | 4.74 | 2.61 |
| 79 | 58.50 | 704 | 259 | 126 | 12.03 | 4.43 | 2.15 |
| 80 | 131.00 | 709 | 625 | 236 | 5.41 | 4.77 | 1.80 |
| 81 | 42.50 | 489 | 183 | 126 | 11.51 | 4.31 | 2.96 |
| 82 | 25.50 | 284 | 124 | 89 | 11.14 | 4.86 | 3.49 |
| 83 | 32.50 | 383 | 166 | 110 | 11.78 | 5.11 | 3.38 |
| 84 | 48.00 | 488 | 173 | 117 | 10.17 | 3.60 | 2.44 |
| 85 | 48.00 | 376 | 194 | 105 | 7.83 | 4.04 | 2.19 |
| 86 | 56.00 | 499 | 300 | 154 | 8.91 | 5.36 | 2.75 |
| 87 | 49.00 | 583 | 214 | 106 | 11.90 | 4.37 | 2.16 |

| | | | | | | |
|---|---|---|---|---|---|---|
| 88 | 35.50 | 445 | 143 | 73 | 12.54 | 4.03 | 2.06 |
| 89 | 44.00 | 454 | 207 | 121 | 10.32 | 4.70 | 2.75 |
| 90 | 24.00 | 258 | 115 | 59 | 10.75 | 4.79 | 2.46 |
| 91 | 30.00 | 369 | 149 | 83 | 12.30 | 4.97 | 2.77 |
| 92 | 49.00 | 592 | 176 | 93 | 12.08 | 3.59 | 1.90 |
| 93 | 7.00 | 82 | 30 | 26 | 11.71 | 4.29 | 3.71 |
| 94 | 101.50 | 1235 | 503 | 324 | 12.17 | 4.96 | 3.19 |
| 95 | 26.00 | 322 | 134 | 76 | 12.38 | 5.15 | 2.92 |
| 96 | 21.00 | 239 | 87 | 71 | 11.38 | 4.14 | 3.38 |
| 97 | 40.50 | 481 | 177 | 108 | 11.88 | 4.37 | 2.67 |
| 98 | 90.50 | 886 | 410 | 210 | 9.79 | 4.53 | 2.32 |
| 99 | 29.00 | 243 | 138 | 81 | 8.38 | 4.76 | 2.79 |
| 100 | 26.00 | 309 | 96 | 48 | 11.88 | 3.69 | 1.85 |
| 101 | 79.00 | 609 | 310 | 173 | 7.71 | 3.92 | 2.19 |
| 102 | 21.50 | 234 | 117 | 58 | 10.88 | 5.44 | 2.70 |
| 103 | 71.00 | 871 | 309 | 165 | 12.27 | 4.35 | 2.32 |
| 104 | 41.50 | 428 | 202 | 130 | 10.31 | 4.87 | 3.13 |
| 105 | 30.50 | 390 | 131 | 81 | 12.79 | 4.30 | 2.66 |
| 106 | 52.50 | 590 | 278 | 160 | 11.24 | 5.30 | 3.05 |
| 107 | 8.50 | 72 | 31 | 19 | 8.47 | 3.65 | 2.24 |
| **Mean** | 72.74 | 668.10 | 298.90 | 172.80 | 9.64 | 4.19 | 2.46 |
| **SE** | 4.68 | 42.64 | 20.40 | 12.18 | 0.25 | 0.10 | 0.06 |

**Table S3 | Statistics of sleep duration, SO, spindle and coupling event numbers and densities for all 107 subjects during N2&N3 stage.**

| Subject ID | REM Sleep Stage | | | | | | |
|---|---|---|---|---|---|---|---|
| | Time (min) | Number of SO | Number of spindle | Number of coupling | Density of SO (times/min) | Density of spindle (times/min) | Density of coupling (times/min) |
| 1 | 2.00 | 13 | 7 | 6 | 6.50 | 3.50 | 3.00 |
| 2 | 4.00 | 26 | 12 | 9 | 6.50 | 3.00 | 2.25 |
| 3 | 4.50 | 39 | 17 | 12 | 8.67 | 3.78 | 2.67 |
| 4 | 7.00 | 48 | 14 | 11 | 6.86 | 2.00 | 1.57 |
| 5 | 3.50 | 35 | 13 | 7 | 10.00 | 3.71 | 2.00 |
| 6 | 24.50 | 264 | 48 | 40 | 10.78 | 1.96 | 1.63 |
| 7 | 5.00 | 18 | 21 | 3 | 3.60 | 4.20 | 0.60 |
| 8 | 21.00 | 215 | 78 | 61 | 10.24 | 3.71 | 2.90 |
| 9 | 4.50 | 27 | 20 | 13 | 6.00 | 4.44 | 2.89 |
| 10 | 15.00 | 153 | 57 | 51 | 10.20 | 3.80 | 3.40 |
| 11 | 4.50 | 33 | 18 | 11 | 7.33 | 4.00 | 2.44 |
| 12 | 4.50 | 22 | 10 | 6 | 4.89 | 2.22 | 1.33 |
| 13 | 2.00 | 10 | 7 | 7 | 5.00 | 3.50 | 3.50 |
| 14 | 0.50 | 0 | 0 | 0 | 0.00 | 0.00 | 0.00 |
| 15 | 16.00 | 186 | 39 | 30 | 11.63 | 2.44 | 1.88 |
| 16 | 11.50 | 51 | 26 | 23 | 4.43 | 2.26 | 2.00 |
| 17 | 12.50 | 51 | 30 | 14 | 4.08 | 2.40 | 1.12 |
| 18 | 4.00 | 35 | 9 | 9 | 8.75 | 2.25 | 2.25 |
| 19 | 11.50 | 108 | 40 | 38 | 9.39 | 3.48 | 3.30 |
| 20 | 5.50 | 55 | 20 | 14 | 10.00 | 3.64 | 2.55 |
| 21 | 12.00 | 105 | 59 | 34 | 8.75 | 4.92 | 2.83 |
| 22 | 0.50 | 0 | 0 | 0 | 0.00 | 0.00 | 0.00 |
| 23 | 11.50 | 73 | 42 | 22 | 6.35 | 3.65 | 1.91 |
| 24 | 8.00 | 81 | 27 | 18 | 10.13 | 3.38 | 2.25 |
| 25 | 32.50 | 381 | 120 | 79 | 11.72 | 3.69 | 2.43 |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| 26 | 2.50 | 10 | 9 | 6 | 4.00 | 3.60 | 2.40 |
| 27 | 7.50 | 94 | 32 | 26 | 12.53 | 4.27 | 3.47 |
| 28 | 5.50 | 42 | 17 | 14 | 7.64 | 3.09 | 2.55 |
| 29 | 17.50 | 178 | 71 | 43 | 10.17 | 4.06 | 2.46 |
| 30 | 6.00 | 60 | 24 | 23 | 10.00 | 4.00 | 3.83 |
| 31 | 13.00 | 85 | 37 | 33 | 6.54 | 2.85 | 2.54 |
| 32 | 0.50 | 0 | 0 | 0 | 0.00 | 0.00 | 0.00 |
| 33 | 1.00 | 9 | 5 | 2 | 9.00 | 5.00 | 2.00 |
| 34 | 9.50 | 85 | 41 | 27 | 8.95 | 4.32 | 2.84 |
| 35 | 2.50 | 16 | 9 | 8 | 6.40 | 3.60 | 3.20 |
| 36 | 9.50 | 83 | 32 | 26 | 8.74 | 3.37 | 2.74 |
| 37 | 24.00 | 260 | 137 | 69 | 10.83 | 5.71 | 2.88 |
| 38 | 4.00 | 38 | 17 | 16 | 9.50 | 4.25 | 4.00 |
| 39 | 13.00 | 137 | 49 | 30 | 10.54 | 3.77 | 2.31 |
| 40 | 8.50 | 81 | 28 | 23 | 9.53 | 3.29 | 2.71 |
| 41 | 7.00 | 74 | 20 | 13 | 10.57 | 2.86 | 1.86 |
| 42 | 17.00 | 178 | 82 | 63 | 10.47 | 4.82 | 3.71 |
| 43 | 25.00 | 165 | 93 | 68 | 6.60 | 3.72 | 2.72 |
| 44 | 15.50 | 170 | 68 | 43 | 10.97 | 4.39 | 2.77 |
| 45 | 7.50 | 67 | 40 | 21 | 8.93 | 5.33 | 2.80 |
| 46 | 0.50 | 0 | 0 | 0 | 0.00 | 0.00 | 0.00 |
| 47 | 1.00 | 7 | 5 | 3 | 7.00 | 5.00 | 3.00 |
| 48 | 46.00 | 428 | 286 | 99 | 9.30 | 6.22 | 2.15 |
| 49 | 0.50 | 0 | 0 | 0 | 0.00 | 0.00 | 0.00 |
| 50 | 29.50 | 223 | 104 | 64 | 7.56 | 3.53 | 2.17 |
| 51 | 19.50 | 204 | 59 | 46 | 10.46 | 3.03 | 2.36 |
| 52 | 8.00 | 18 | 10 | 9 | 2.25 | 1.25 | 1.13 |
| 53 | 6.00 | 62 | 21 | 14 | 10.33 | 3.50 | 2.33 |
| 54 | 6.00 | 42 | 21 | 11 | 7.00 | 3.50 | 1.83 |
| 55 | 1.00 | 4 | 2 | 1 | 4.00 | 2.00 | 1.00 |
| 56 | 20.00 | 243 | 73 | 51 | 12.15 | 3.65 | 2.55 |

| 57 | 0.50 | 0 | 0 | 0 | 0.00 | 0.00 | 0.00 |
|----|------|---|---|---|------|------|------|
| 58 | 6.00 | 51 | 22 | 17 | 8.50 | 3.67 | 2.83 |
| 59 | 45.00 | 501 | 212 | 91 | 11.13 | 4.71 | 2.02 |
| 60 | 0.50 | 0 | 0 | 0 | 0.00 | 0.00 | 0.00 |
| 61 | 8.00 | 72 | 42 | 26 | 9.00 | 5.25 | 3.25 |
| 62 | 10.50 | 116 | 34 | 34 | 11.05 | 3.24 | 3.24 |
| 63 | 5.50 | 40 | 24 | 14 | 7.27 | 4.36 | 2.55 |
| 64 | 7.50 | 78 | 49 | 26 | 10.40 | 6.53 | 3.47 |
| 65 | 9.00 | 87 | 61 | 24 | 9.67 | 6.78 | 2.67 |
| 66 | 22.00 | 200 | 98 | 50 | 9.09 | 4.45 | 2.27 |
| 67 | 22.00 | 199 | 88 | 54 | 9.05 | 4.00 | 2.45 |
| 68 | 18.50 | 206 | 62 | 46 | 11.14 | 3.35 | 2.49 |
| 69 | 5.50 | 43 | 13 | 11 | 7.82 | 2.36 | 2.00 |
| 70 | 10.00 | 88 | 22 | 21 | 8.80 | 2.20 | 2.10 |
| 71 | 29.50 | 255 | 112 | 79 | 8.64 | 3.80 | 2.68 |
| 72 | 0.00 | 0 | 0 | 0 | / | / | / |
| 73 | 14.50 | 108 | 53 | 38 | 7.45 | 3.66 | 2.62 |
| 74 | 13.50 | 139 | 57 | 37 | 10.30 | 4.22 | 2.74 |
| 75 | 3.50 | 25 | 13 | 8 | 7.14 | 3.71 | 2.29 |
| 76 | 4.50 | 35 | 16 | 12 | 7.78 | 3.56 | 2.67 |
| 77 | 14.00 | 139 | 70 | 43 | 9.93 | 5.00 | 3.07 |
| 78 | 26.00 | 297 | 150 | 72 | 11.42 | 5.77 | 2.77 |
| 79 | 2.50 | 19 | 7 | 6 | 7.60 | 2.80 | 2.40 |
| 80 | 6.50 | 27 | 21 | 11 | 4.15 | 3.23 | 1.69 |
| 81 | 4.00 | 35 | 14 | 8 | 8.75 | 3.50 | 2.00 |
| 82 | 3.00 | 30 | 11 | 9 | 10.00 | 3.67 | 3.00 |
| 83 | 10.50 | 113 | 43 | 31 | 10.76 | 4.10 | 2.95 |
| 84 | 1.00 | 6 | 1 | 1 | 6.00 | 1.00 | 1.00 |
| 85 | 8.00 | 88 | 33 | 20 | 11.00 | 4.13 | 2.50 |
| 86 | 0.00 | 0 | 0 | 0 | / | / | / |
| 87 | 3.50 | 31 | 11 | 11 | 8.86 | 3.14 | 3.14 |

| | | | | | | | |
|---|---|---|---|---|---|---|---|
| 88 | 1.50 | 13 | 3 | 2 | 8.67 | 2.00 | 1.33 |
| 89 | 8.50 | 75 | 42 | 27 | 8.82 | 4.94 | 3.18 |
| 90 | 4.00 | 38 | 16 | 9 | 9.50 | 4.00 | 2.25 |
| 91 | 1.50 | 12 | 6 | 3 | 8.00 | 4.00 | 2.00 |
| 92 | 2.00 | 15 | 8 | 7 | 7.50 | 4.00 | 3.50 |
| 93 | 1.50 | 13 | 5 | 3 | 8.67 | 3.33 | 2.00 |
| 94 | 5.50 | 53 | 26 | 13 | 9.64 | 4.73 | 2.36 |
| 95 | 6.00 | 53 | 20 | 17 | 8.83 | 3.33 | 2.83 |
| 96 | 7.00 | 75 | 26 | 18 | 10.71 | 3.71 | 2.57 |
| 97 | 0.00 | 0 | 0 | 0 | / | / | / |
| 98 | 11.00 | 95 | 52 | 33 | 8.64 | 4.73 | 3.00 |
| 99 | 15.00 | 117 | 60 | 47 | 7.80 | 4.00 | 3.13 |
| 100 | 4.50 | 49 | 18 | 13 | 10.89 | 4.00 | 2.89 |
| 101 | 3.50 | 19 | 12 | 8 | 5.43 | 3.43 | 2.29 |
| 102 | 7.50 | 62 | 30 | 21 | 8.27 | 4.00 | 2.80 |
| 103 | 1.00 | 5 | 3 | 1 | 5.00 | 3.00 | 1.00 |
| 104 | 8.50 | 77 | 36 | 23 | 9.06 | 4.24 | 2.71 |
| 105 | 6.50 | 50 | 22 | 16 | 7.69 | 3.38 | 2.46 |
| 106 | 2.50 | 17 | 12 | 7 | 6.80 | 4.80 | 2.80 |
| 107 | 16.50 | 155 | 58 | 38 | 9.39 | 3.52 | 2.30 |
| **Mean** | 9.27 | 85.17 | 36.64 | 23.14 | 7.94 | 3.49 | 2.32 |
| **SE** | 0.86 | 9.03 | 4.17 | 2.13 | 0.29 | 0.13 | 0.09 |

**Table S4 | Statistics of sleep duration, SO, spindle and coupling event numbers and densities for all 107 subjects during REM stage.**

| Brain areas | Laterality | Peak T-value | Cluster size | Peak coordinate (MNI) | | |
|---|---|---|---|---|---|---|
| | | | | x | y | z |
| Anterior Supramarginal Gyrus | L | -3.93 | 303 | -52 | -29 | 40 |
| Anterior Supramarginal Gyrus | R | -2.66 | 18 | 67 | -29 | 27 |
| Superior Frontal Gyrus | L | -4.53 | 2766 | -12 | 4 | 66 |
| Superior Frontal Gyrus | R | -4.86 | 2122 | 12 | 21 | 59 |
| Middle Frontal Gyrus | L | -5.57 | 3320 | -44 | 26 | 33 |
| Middle Frontal Gyrus | R | -5.56 | 3312 | 40 | 25 | 37 |
| Anterior Paracingulate Gyrus | L | -7.28 | 4153 | -5 | 38 | -10 |
| Anterior Paracingulate Gyrus | R | -7.13 | 3918 | 4 | 45 | -8 |
| Angular Gyrus | L | -3.52 | 519 | -62 | -54 | 26 |
| Angular Gyrus | R | -2.66 | 401 | 50 | -52 | 49 |
| Frontal Orbital Cortex | L | -6.46 | 6392 | -38 | 33 | -17 |
| Frontal Orbital Cortex | R | -7.26 | 5637 | 28 | 28 | -20 |
| Superior Parietal Lobule | L | -5.10 | 1774 | -35 | -44 | 58 |
| Superior Parietal Lobule | R | -5.96 | 1457 | 35 | -41 | 59 |
| Insular Cortex | L | -4.89 | 1004 | -32 | -26 | 16 |
| Insular Cortex | R | -6.34 | 1028 | 34 | -24 | 9 |
| Posterior Cingulate Gyrus | L | -10.83 | 4525 | -5 | -47 | 19 |
| Posterior Cingulate Gyrus | R | -10.47 | 5838 | 5 | -43 | 27 |
| Posterior Supramarginal Gyrus | L | -3.24 | 102 | -61 | -51 | 26 |
| Posterior Supramarginal Gyrus | R | -2.41 | 7 | 67 | -38 | 21 |

**Table S5 | Peak and significant cluster of fMRI activity during SO main effect.** We used the SO main effect whole-brain activation patterns in **Fig. 3b.** ROIs

were defined anatomically (see Methods). Cluster sizes are reported $p_{unc.} < 0.001$.

| Brain areas | Laterality | Peak T-value | Cluster size | Peak coordinate (MNI) | | |
|---|---|---|---|---|---|---|
| | | | | x | y | z |
| Anterior Supramarginal Gyrus | L | -6.36 | 1357 | -55 | -29 | 45 |
| Anterior Supramarginal Gyrus | R | -6.53 | 1671 | 55 | -26 | 49 |
| Superior Frontal Gyrus | L | -6.83 | 2274 | -20 | 33 | 38 |
| Superior Frontal Gyrus | R | 5.52 | 2320 | 12 | -4 | 75 |
| Middle Frontal Gyrus | L | -8.03 | 5350 | -42 | 30 | 30 |
| Middle Frontal Gyrus | R | -5.77 | 4194 | 31 | 28 | 41 |
| Anterior Paracingulate Gyrus | L | 7.15 | 1086 | -7 | 11 | 42 |
| Anterior Paracingulate Gyrus | R | 6.68 | 2222 | 2 | 15 | 41 |
| Angular Gyrus | L | -7.55 | 1240 | -48 | -58 | 50 |
| Angular Gyrus | R | -7.33 | 4588 | 53 | -53 | 34 |
| Frontal Orbital Cortex | L | -4.61 | 1149 | -41 | 21 | -11 |
| Frontal Orbital Cortex | R | -3.98 | 645 | 40 | 20 | -16 |
| Superior Parietal Lobule | L | -6.91 | 1687 | -36 | -56 | 61 |
| Superior Parietal Lobule | R | -7.79 | 3091 | 42 | -43 | 62 |
| Insular Cortex | L | -6.99 | 1103 | -37 | -20 | 15 |
| Insular Cortex | R | -7.26 | 2169 | 36 | -22 | 12 |
| Posterior Cingulate Gyrus | L | -7.85 | 3208 | -5 | -46 | 38 |
| Posterior Cingulate Gyrus | R | -7.10 | 3352 | 0 | -45 | 36 |
| Posterior Supramarginal Gyrus | L | -6.71 | 1263 | -54 | -49 | 39 |
| Posterior Supramarginal Gyrus | R | -6.61 | 2376 | 48 | -38 | 54 |

**Table S6 | Peak and significant cluster of fMRI activity during spindle main effect.** We used the Spindle main effect whole-brain activation patterns in **Fig.**

**3c.** ROIs were defined anatomically (see Methods). Cluster sizes are reported $p_{unc.} < 0.001$.

| Brain areas | Laterality | Peak T-value | Cluster size | Peak coordinate (MNI) | | |
|---|---|---|---|---|---|---|
| | | | | x | y | z |
| Anterior Supramarginal Gyrus | L | -4.95 | 1327 | -56 | -33 | 38 |
| Anterior Supramarginal Gyrus | R | -2.42 | 10 | 60 | -23 | 27 |
| Superior Frontal Gyrus | L | -3.50 | 398 | -14 | 10 | 64 |
| Superior Frontal Gyrus | R | -4.44 | 596 | 6 | 23 | 49 |
| Middle Frontal Gyrus | L | -4.74 | 2455 | -50 | 11 | 40 |
| Middle Frontal Gyrus | R | -6.01 | 1694 | 53 | 17 | 37 |
| Anterior Paracingulate Gyrus | L | -3.56 | 381 | -1 | 21 | 47 |
| Anterior Paracingulate Gyrus | R | -4.98 | 725 | 7 | 21 | 47 |
| Angular Gyrus | L | 1.71 | 0 | -62 | -54 | 26 |
| Angular Gyrus | R | 2.63 | 149 | 55 | -58 | 19 |
| Frontal Orbital Cortex | L | -4.72 | 1492 | -47 | 26 | -6 |
| Frontal Orbital Cortex | R | -4.24 | 872 | 40 | 27 | -2 |
| Superior Parietal Lobule | L | -4.83 | 1177 | -37 | -46 | 56 |
| Superior Parietal Lobule | R | -4.02 | 584 | 28 | -50 | 54 |
| Insular Cortex | L | -4.01 | 1385 | -38 | 19 | -1 |
| Insular Cortex | R | -4.36 | 1374 | 32 | 21 | 0 |
| Posterior Cingulate Gyrus | L | -2.93 | 83 | -9 | -32 | 40 |
| Posterior Cingulate Gyrus | R | -2.91 | 130 | 7 | -30 | 42 |
| Posterior Supramarginal Gyrus | L | -1.96 | 0 | -48 | -44 | 46 |
| Posterior Supramarginal Gyrus | R | -2.02 | 0 | 49 | -37 | 51 |

**Table S7 | Peak and significant cluster of fMRI activity during SO-spindle interaction.** We used the SO-spindle interaction effect whole-brain activation patterns in **Fig. 3d.** ROIs were defined anatomically (see Methods). Cluster sizes are reported $p_{unc.} < 0.001$.