Bacteroides thetaiotaomicron enhances H$_2$S production in *Bilophila wadsworthia*

Jade Davies[1], Melinda J. Mayer[1], Nathalie Juge[1], Arjan Narbad[1,2], Lizbeth Sayavedra[1,2,3]

[1]Quadram Institute Bioscience, Norwich, UK

[2]Joint last author

[3]Corresponding author: lizbeth.sayavedra@quadram.ac.uk

Abstract

Sulfate- and sulfite-reducing bacteria (SRB) are a group of strict anaerobes found within the human gut. *Bilophila wadsworthia*, a sulfite-reducing bacterium which produces hydrogen sulfide (H$_2$S) from taurine and isethionate respiration is a common member of the healthy commensal human gut microbiota, but has been implicated in several disease states including inflammatory bowel disease and colorectal cancer. *Bacteroides thetaiotaomicron*, one of the most prominent gut bacteria, has sulfatases which release sulfate, serving as a potential substrate for sulfate-reducing bacteria. Here, we showed that when *B. thetaiotaomicron* and *B. wadsworthia* were in co-culture, there was a significant increase in *B. thetaiotaomicron*'s growth and in H$_2$S production by *B. wadsworthia*. Differential gene expression analysis revealed increased expression of *B. wadsworthia*'s *dsrMKJOP* complex in co-culture, which delivers electrons for sulfite reduction to H$_2$S. This was accompanied by a decreased expression of genes associated with taurine, sulfolactate and thiosulfate respiration, indicating that *B. thetaiotaomicron* may provide an alternative source of sulfite to *B. wadsworthia*. We hypothesised adenosine 5'-phosphosulfate (APS) to be this intermediate. Indeed, *B. wadsworthia* was able to grow using APS or sulfite as electron acceptors. Endometabolomic and transcriptomic analyses revealed decreased production of indole by *B. thetaiotaomicron* in co-culture with *B. wadsworthia* due to enhanced tryptophan utilisation by *B. wadsworthia*. The results of this microbe-microbe interaction could have significant pro-inflammatory effects in the human gut environment.

Introduction

Sulfate- and sulfite-reducing bacteria (SRB) comprise a group of strict anaerobes and are found within the colonic mucosa as part of the commensal gut microbiota in at least 50% of humans [1, 2]. SRB utilise inorganic sulfate (SO_4^{2-}) or sulfite (SO_3^{2-}) as a terminal electron acceptor during energy metabolism, a process which occurs concomitantly with oxidation of molecular hydrogen or organic compounds. Hydrogen sulfide (H_2S) is the final product of dissimilatory sulfate reduction by SRB [3] and can freely diffuse across cell membranes [4]. H_2S is recognised as a hazardous product as it is both corrosive and toxic to many organs even at low concentrations [5]. Classical sulfate-reducing bacteria reduce sulfate to sulfite via sulfate adenylyltransferase (Sat) and adenylylsulfate reductase (AprAB) [6] (Figure S1). Sulfite then enters the dissimilatory sulfite reduction pathway. *Bilophila wadsworthia* is a Gram-negative member of the SRB, first identified in 1989 from gangrenous and perforated appendicitis samples [8].Unlike other SRB, *B. wadsworthia* cannot utilise sulfate [7-9], and instead degrades organosulfate compounds such as taurine, isethionate, and sulfoquinovase, generating H_2S as a by-product [10, 11]. In contrast, the majority of living organisms have the assimilatory sulfate reduction pathway [12]. This involves the reduction of sulfate to APS via sulfate adenylyltransferase (Sat), which is then converted to phosphoadenosine phosphosulfate (PAPS) via adenylylsulfate kinase / APS kinase. PAPS is then reduced to sulfite by phosphoadenosine phosphosulfate (PAPS) reductase, and the sulfite is then converted to H_2S by assimilatory sulfite reductase [12] (Figure S1).

B. wadsworthia is present in the faeces of 50-60% of healthy individuals [13], and can also be isolated from buccal and vaginal samples [14]. *B. wadsworthia* is considered to be virulent, as it is the third most common anaerobic isolate from appendicitis samples and appears to be clinically important in a variety of anaerobic infections [15]. Furthermore, it exhibits endotoxic activity [16], and is adherent to human embryonic intestinal cells *in vitro* [17]. Association studies in humans have linked *B. wadsworthia* enrichment in the gut with many diseases including colorectal cancer [18], multiple sclerosis [19], Parkinson's disease [20], dementia [21], non-alcoholic steatohepatitis [22], intrahepatic cholestasis in pregnancy [23], diabetic kidney disease [24] and schizophrenia [25]. Additionally, it was demonstrated that a human stool-derived *B. wadsworthia* strain was able to induce systemic inflammation in specific-pathogen-free mice [26]. Given *B. wadsworthia*'s status as a potential pathobiont in the human gut, it is important to investigate the factors that influence its abundance and function.

Bacteroides thetaiotaomicron is a Gram-negative obligate anaerobe within the
Bacteroidaceae family [27]. Originally isolated from the faeces of a healthy adult [27], *B.*
thetaiotaomicron is highly abundant constituting 1-6% of the total bacteria [28] with a 46%
prevalence [29], and is a key commensal member of the human gut microbiota. *B.*
thetaiotaomicron plays a major role in the human gut, including modulation of the host
mucosal immune system [30], and utilisation of a wide range of polysaccharides [31]. Some of
these polysaccharides can be host-derived, and include glycosaminoglycans such as
chondroitin sulfate, mucin, hyaluronate and heparan sulfate [32]. The degradation of
polysaccharides yields simple sugars for fermentation, resulting in the production of
beneficial short-chain organic acids including acetate, lactate, succinate and propionate [33];
B. thetaiotaomicron has also been shown to support the growth of butyrate-producing
Anaerostipes caccae in vitro [34]. Short-chain fatty acids (SCFAs) are a valuable source of
energy for human colonocytes and aid in maintaining healthy barrier function [35]. *B.*
thetaiotaomicron encodes 28 sulfatases which can cleave sulfated residues of the host
glycosaminoglycans to yield free sulfate in the mouse gut [36]. By increasing the availability
of free sulfate *in vivo*, *B. thetaiotaomicron* sulfatase activity could permit increased sulfate
reduction and H_2S production by SRB [37]. Indeed, *B. thetaiotaomicron* and *Desulfovibrio*
piger, a SRB, can co-colonise the gut, with *B. thetaiotaomicron* providing free sulfate to *D.*
piger and promoting H_2S production both *in vitro* and *in vivo* [2]. Additionally, *B.*
thetaiotaomicron generates hydrogen during polysaccharide fermentation and molecular
hydrogen, which is used by hydrogenotrophs, including SRB, as an electron donor [38, 39].
Given these established interactions with other SRBs, we aimed to further investigate the
dynamics between *B. thetaiotaomicron* and *B. wadsworthia*. Here, we used
transcriptomics and metabolomics to explore the mechanisms underpinning the interaction
of *B. wadsworthia* and *B. thetaiotaomicron* during anaerobic co-culture.

Results and Discussion

Co-culture of *B. wadsworthia* and *B. thetaiotaomicron* boosts H_2S production

B. wadsworthia (QI0013) and *B. thetaiotaomicron* (QI0072) were grown anaerobically in
monocultures and co-cultures on synthetic (BPM, see methods) media supplemented with
10 mM taurine at a 1:1 ratio (10^6 colony forming units (CFU)/mL inoculation density). At 8
hours post-inoculation, a significant increase in H_2S concentration was observed in the co-
culture compared to *B. wadsworthia* monoculture (Figure 1a) while *B. wadsworthia*

abundance remained unaffected by the co-culture (Figure 1b). The elevated H_2S levels
were attributed to a significantly increased H_2S production per *B. wadsworthia* cell when in
co-culture with *B. thetaiotaomicron* (Figure 1d). Interestingly, the interaction also appeared
to benefit *B. thetaiotaomicron*, which grew to significantly higher abundance in the co-
culture (Figure 1c).
To gain further insights into the potential mechanism that increases H_2S production in *B.*
wadsworthia, we performed transcriptomic and metabolomic analysis of the mono and co-
cultures at 8 hours. The most impacted pathways are discussed below.
113

Respiration of sulfite via hydrogen or lactate by *B. wadsworthia* was
increased in co-culture
In the co-culture, four genes associated with sulfur metabolism leading to H_2S production
in *B. wadsworthia* were overexpressed (Figure 2a, Table S1). These genes are part of the
dissimilatory sulfite reductase protein complex (*dsrMKJOP*), including *dsrM*
(WCP94_001671), *dsrK* (WCP94_001670), *dsrO* (WCP94_001668), and *dsrP*
(WCP94_001667) (Table S1). When *B. wadsworthia* respires taurine, sulfite (SO_3^{2-})
generated enters the dissimilatory sulfite reduction pathway and is further reduced by
DsrAB to produce H_2S (Figure S1). Expression of *dsrMKJOP* can be increased under H_2-
rich growth conditions in *D. vulgaris* [40]; the observed up-regulation in co-culture may
suggest a higher bioavailability of H_2 to *B. wadsworthia*. *B. thetaiotaomicron* can generate
hydrogen (H_2) during polysaccharide fermentation [41]; therefore, *B. wadsworthia* could
utilise this directly as an electron donor during dissimilatory sulfite reduction [38, 42]. The
increased expression of *dsrMKOP* in co-culture alludes to sulfite utilisation as an electron
acceptor, potentially leading to increased H_2S production.
129

In addition to hydrogen, lactate can also be utilised as an electron donor for the respiration
of sulfite [10, 43, 44] (Figure 3). Interestingly, two genes encoding lactate permeases were
overexpressed in co-culture (WCP94_002306, WCP94_000357) in *B. wadsworthia* (Table
S1), suggesting enhanced utilisation of lactate. The presence of different electron donors
might contribute to the increased expression of *dsrMKJOP* in *B. wadsworthia* under co-
culture conditions, leading to a higher H_2S production as a by-product. Taken together, the
transcriptomic data revealed an increased capacity for uptake and utilisation of hydrogen
and lactate in *B. wadsworthia* in co-culture with *B. thetaiotaomicron*; both compounds are
key electron donors for dissimilatory sulfite reduction. This is in line with increased H_2S
concentration by *B. wadsworthia* in co-culture (Figure 1a).
140

Surprisingly, the gene cluster associated with taurine metabolism exhibited reduced expression in the co-culture, despite the observed increase in H_2S concentration. This gene cluster includes taurine pyruvate dehydrogenase (*tpa*, WCP94_00949), alanine dehydrogenase (*ald*, WCP94_00950) and sulfoacetaldehyde reductase (*sarD*, WCP94_00947) [10] (Figure S1, Figure 3, Table S1). It has been shown that expression of these enzymes is elevated in *B. wadsworthia* cells grown in taurine, but not in isethionate-grown cells [10]. Isethionate sulfite-lyase *islA* and isethionate sulfite-lyase activating protein *isiB* were not among the differentially expressed genes, but acetaldehyde dehydrogenase (*adhE*, WCP94_001941), which encodes an enzyme able to convert acetaldehyde (a product of isethionate degradation) to acetyl coA was decreased in expression in co-culture (Table S1). The downregulation of *tpa* and *sarD,* combined with the increased expression of *dsrMKJOP* suggests that *B. wadsworthia* can also utilise the respiration pathway from sulfite or a precursor molecule in the presence of *B. thetaiotaomicron* (Figure 3). Sulfite could be produced through different metabolic pathways in *B. wadsworthia*: i) from sulfolactate, via the activity of a sulfo-lyase (WCP94_000771-772) which converts sulfolactate to sulfite and pyruvate; this enzyme showed decreased expression in co-culture; or ii) through the inner membrane protein YgaP (WCP94_002882)*,* a rhodanese-domain-containing protein that generates sulfite from thiosulfate, and which also showed decreased expression in co-culture (Table S1). Therefore, it is likely that in co-culture with *B. thetaiotaomicron*, *B. wadsworthia* utilises an alternative source of sulfite (Figure 3).

Modulation of sulfur metabolism in *B. wadsworthia* and *B. thetaiotaomicron* co-culture

As taurine utilisation by *B. wadsworthia* as a sulfite source in co-culture appeared to be decreased, alongside an increased expression of genes associated with the dissimilatory sulfite reductase pathway and elevated H_2S concentration, we investigated other possible sources of sulfite for *B. wadsworthia* in the co-culture. Increased expression of sulfate permease (WCP94_001830) and sulfatase (WCP94_001848) genes was observed in *B. wadsworthia* in co-culture (Table S1), suggesting an increased bioavailability of sulfate; however, whether these transporters are promiscuous for other compounds such as sulfite is unknown.

Given that *B. wadsworthia* cannot utilise sulfate, it was previously assumed that the genes for sulfate utilisation were absent from the genome [7, 8, 45]. The *B. wadsworthia* QI0013

genome does not contain genes encoding the Sat enzyme necessary for sulfate reduction. However, our analysis revealed that *B. wadsworthia* QI0013 encodes two gene clusters of the alpha and beta subunits of the adenylylsulfate reductase (WCP94_000309-310 and WCP94_000741-742) (Table S3, Figure S1). Despite low amino acid sequence similarity to AprAB proteins from *Desulfovibrio* strains (Figure S2, Table S4), we identified conserved protein domains, including FAD-dependent oxidoreductase 2 and succinate dehydrogenase/fumarate reductase flavoprotein, supporting the prediction of AprAB activity (Table S5). Furthermore, we found that the putative AprAB genes were widely distributed in publicly available *B. wadsworthia* genomes (Table S6). This suggests a potential role for *B. wadsworthia* in APS reduction via adenylylsulfate reductase. The putative proteins encoded by these genes could facilitate the conversion of APS to sulfite, offering an alternative precursor molecule for the dissimilatory sulfite reduction pathway. Interestingly, the *B. thetaiotaomicron* QI0072 genome encodes some enzymes involved in assimilatory sulfite reduction, including sulfate adenylyltransferase (Sat) (QI0072_1554, QI0072_1555) and adenylylsulfate kinase / APS kinase (QI0072_1556) (Table S3), which would catalyse the conversion of sulfate to both APS and PAPS. However, *B. thetaiotaomicron* QI0072 does not encode the enzymes for further utilisation of PAPS and APS. *B. wadsworthia* QI0013 expresses both phosphoadenosine phosphosulfate (PAPS) reductase (WCP94_00194), which reduces PAPS to sulfite, in addition to the putative AprAB which reduces APS to sulfite (Table S3).

We hypothesise that *B. thetaiotaomicron* QI0072 produces sulfate from sulfated saccharides via sulfatases [46], an activity which has been previously described to support the growth of *D. piger* via cross-feeding [47], and then reduces the sulfate to APS and/or PAPS. In *B. thetaiotaomicron*, genes encoding sulfate adenylyltransferase subunits 1 and 2 and adenylylsulfate kinase were consistently expressed at similar levels in both co-culture and monoculture (Table S3), suggesting the production of APS and PAPS. *B. thetaiotaomicron*-derived APS and PAPS could then be utilised directly by *B. wadsworthia*, where either AprAB or PAPS reductase yield sulfite, which then enters the dissimilatory sulfate reduction pathway (Figure 3). In this way, *B. thetaiotaomicron* may provide *B. wadsworthia* with an alternative source of sulfite, which may be energetically favourable over taurine degradation, or utilised when taurine or other organosulfur compounds are depleted. Indeed, we observed a decreased expression of genes associated with sulfite generation via taurine, thiosulfate and sulfolactate under co-culture conditions, with the phenotypic observation of increased H_2S concentration (Figure 1). The adenylylsulfate

reductase and PAPS reductase genes were expressed in *B. wadsworthia* in both co-culture and monoculture, without significant differences in expression levels (Table S3). Next, we investigated if *B. wadsworthia* would grow using sulfite or APS as electron acceptors instead of taurine. *B. wadsworthia* was able to show a significantly higher growth in media supplemented with 4 mM sulfite, 10 mM APS, or 10 mM taurine, compared to only anaerobe basal broth (ABB) (Figure 4).

Next, we tested the capacity of *B. thetaiotaomicron* to promote the sulfidogenic activity of other *B. wadsworthia* strains i.e *B. wadsworthia* QI0012, QI0014, and QI0015 strains. Co-culture with *B. thetaiotaomicron* resulted in significantly increased H_2S concentration in all *B. wadsworthia* strains tested compared to their respective monocultures (Figure 5a), suggesting that the sulfidogenic potential of *B. thetaiotaomicron* is not strain specific. Indeed, QI0014 did not produce detectable H_2S in monoculture at 8 h, whereas 195.3 ± 18.60 µM was produced in co-culture with *B. thetaiotaomicron* (Figure 5a). Interestingly, qPCR analysis of *B. wadsworthia* cell abundance revealed differences between strains with QI0012 and QI0013 showing slight increases in abundance in co-culture with *B. thetaiotaomicron* although this effect was not significant with QI0013 (Figure 5b), as shown above for this strain (Figure 1). A decreased abundance was observed in co-culture *B. wadsworthia* QI0014 and QI0015 strains with *B. thetaiotaomicron* (Figure 5b). Upon standardizing H_2S concentrations relative to *B. wadsworthia* cell numbers, it was observed that *Bacteroides thetaiotaomicron* consistently enhanced H_2S production across the various *B. wadsworthia* strains. Nonetheless, the specific mechanisms behind this increase may differ among strains (Figure 5).

The increased H_2S levels in QI0012 and QI0013 co-cultures are attributable to a combination of higher *B. wadsworthia* abundance and elevated H_2S production per cell. In contrast, the rise in H_2S observed with QI0014 and QI0015 is predominantly due to a substantial increase in H_2S production per *B. wadsworthia* cell (Figure 5d). Additionally, *B. thetaiotaomicron* exhibited a slight increase in abundance in all co-cultures compared to its monoculture (Figure 5c). Overall, *B. thetaiotaomicron* showed an ability to increase H_2S production in all four *B. wadsworthia* strains that were tested.

H$_2$S-utilising amino acid biosynthetic pathways were reduced in expression in *B. thetaiotaomicron*

Based on our genome analyses, *B. thetaiotaomicron* QI0072 does not encode known genes for H$_2$S production, but it does have the capacity to utilise H$_2$S during the biosynthesis of amino acids including cysteine and homocysteine. *B. thetaiotaomicron* QI0072 over-expressed twelve genes and under-expressed nine genes in co-culture associated with amino acid metabolism, compared to monoculture (Figure 2b). Interestingly, this strain under-expressed O-acetylserine sulfhydrolase (QI0072_3844) in co-culture (Table S2); this enzyme catalyses the conversion of O3-acetyl-L-serine and H$_2$S to L-cysteine and acetate, utilising pyridoxal-5'-phosphate (vitamin B6) as a co-factor [48]. This is relevant as this indicates that although *B. thetaiotaomicron* can utilise *B. wadsworthia*-derived H$_2$S, the gene encoding this enzyme is heavily decreased in expression in co-culture with *B. wadsworthia*, which could indirectly contribute to the higher H$_2$S concentration observed in the co-culture. Similarly, two genes involved in H$_2$S-utilising homocysteine biosynthesis were also differentially expressed; one gene encoding O-acetylhomoserine sulfhydrylase (EC 2.5.1.49) was notably decreased in expression in co-culture (QI0072_3143, -5.12 logFC), whereas another gene encoding this enzyme was increased in expression in co-culture (QI0072_2658, 1.24 logFC) (Table S2). This enzyme catalyses the conversion of O-acylhomoserine and H$_2$S to homocysteine [49, 50]. Overall, decreased homocysteine and cysteine biosynthesis by *B. thetaiotaomicron* in co-culture with *B. wadsworthia* results in lower H$_2$S utilisation by *B. thetaiotaomicron*, which indirectly contributes towards the overall phenotype of high H$_2$S concentration in the co-culture.

Alterations to the endometabolome in *B. wadsworthia* and *B. thetaiotaomicron* co-culture

The endometabolome of the *B. wadsworthia* and *B. thetaiotaomicron* co-culture was analysed at 8 hours post-inoculation, alongside monocultures. The *B. thetaiotaomicron* monoculture metabolite abundance clustered distinctly from the *B. wadsworthia* monoculture and co-culture conditions based on partial least squares discriminant analysis (PLS-DA) (Figure S3a). The heatmap displaying the top 50 differentially abundant metabolites showed clear differences between the conditions, most notably when the *B. thetaiotaomicron* monoculture is compared to the co-culture conditions (Figure S3b). The variable importance in projection (VIP) scores showed that the top 15 compounds contributing to differences between conditions were relatively high in abundance in the *B. thetaiotaomicron* monoculture and in relatively low abundance in the co-culture (Figure 6a). This suggests that these compounds are produced by *B. thetaiotaomicron* and either

not produced by *B. thetaiotaomicron* in co-culture with *B. wadsworthia* or consumed by *B.*

wadsworthia. The highest-scoring compound was Ala-Val, followed by the ester phenethyl

acetate, sphinganine and nicotinic compound 1-Methylnicotinamide (Figure 6a). Kynurenic

acid also had a significantly lower abundance in co-culture compared to *B.*

thetaiotaomicron monoculture (Figure 6a,b), suggesting a reduction in tryptophan

metabolism via this pathway. Kynurenic acid is a product of tryptophan degradation in

mammals [51], however, *Escherichia coli* has been demonstrated to produce kynurenic acid

from L-kynurenine in the rat small intestine [52] and the human gut microbiota may also

produce this compound, since kynurenic acid concentration is high in the distal colon

relative to other body sites [53]. Sphinganine can be produced by *Bacteroides* strains in the

gut and has the potential to control the levels of bioactive lipids in the liver [54], underscoring

its potential role in host-microbial interactions.

The transcriptomic data identified branched-chain amino acids (BCAAs) as significant

metabolites of interest, showcasing differential expression in Liv system genes responsible

for ABC transporters specific to BCAAs in *B. wadsworthia* (Figure 2a, supplementary

Table S1). This may signify a modified bioavailability of leucine, isoleucine, and valine for

B. wadsworthia in the co-culture environment. Indeed, valine and isoleucine were detected

in the endometabolome in all culture conditions, in addition to dipeptides of leucine

including threonylleucine, prolylleucine, N-leucyl-leucine and leucyl-leucine (Figure 6b). In

all cases, relative abundance was high in *B. thetaiotaomicron* monoculture, lower in *B.*

wadsworthia monoculture and the lowest abundance was in the co-culture condition

(Figure 6b). Taken together, this suggests lower bioavailability of BCAAs in the

endometabolome of *B. wadsworthia* and *B. thetaiotaomicron* in co-culture at 8 hours

compared to monocultures. This interplay could have implications for human health, since

increased concentrations of aromatic and BCAAs by the gut microbiome have been linked

to insulin resistance and type 2 diabetes mellitus [55].

Impact on tryptophan metabolism between *B. thetaiotaomicron* and *B.*
***wadsworthia*: insights from combined metabolomic and transcriptomic**
analyses

The endometabolome revealed differences in the relative concentrations of tryptophan,

indole, and indole-3-acetamide between *B. thetaiotaomicron* and *B. wadsworthia*

monocultures and their co-culture; the abundance of these metabolites was higher in the

B. thetaiotaomicron monoculture compared to the other culture conditions (Figure 6b and

6c). Bacterial produced indole has been shown to modulate intestinal inflammation in
animals, and inhibit quorum sensing and virulence factor production [56]. Indole-producing
members of the human gut microbiota include *E. coli*, *Proteus vulgaris*, *Paracolobactrum*
coliforme, *Achromobacter liquefaciens* and *Bacteroides spp.* [57]. These strains can degrade
tryptophan to indole, pyruvate and ammonia via the enzyme tryptophanase (TnaA), the
activity of which is induced by tryptophan [58] and inhibited by glucose [57]. Interestingly,
indole production from tryptophan is commonly observed in *B. thetaiotaomicron* [59], and *B.*
thetaiotaomicron-derived indole has been shown to inhibit virulence in enteropathogenic *E.*
coli (EPEC) and *Vibrio cholerae* by inhibiting T3SS expression [59]. Given that *B.*
wadsworthia cannot produce indole [60], it is likely that indole and its by-products are
derived from *B. thetaiotaomicron*. In co-culture, the relative abundance of these
metabolites decreased compared to the *B. thetaiotaomicron* monoculture (Figure 6b). To
confirm that this effect was not due to the concentrations being standardised to the total
cell count, the indole and tryptophan relative abundances were determined in the *B.*
thetaiotaomicron monoculture and co-culture, normalized to 10^9 *B. thetaiotaomicron* cells
only as determined by qPCR. This showed a decreased relative abundance of indole and
tryptophan in the endometabolome of the co-culture compared to *B. thetaiotaomicron*
monoculture (Figure 6c). This suggests that *B. wadsworthia* may either utilise indole
derived from *B. thetaiotaomicron* or directly uses tryptophan, thereby limiting the
availability of tryptophan for *B. thetaiotaomicron*. Transcriptomic data supported this
observation, showing increased expression of a tryptophan-specific transport protein
(WCP94_002980) and tryptophanyl-tRNA synthetase gene (WCP94_002967) in *B.*
wadsworthia during co-culture, indicating enhanced tryptophan uptake and incorporation
into new proteins. These combined metabolomic and transcriptomic findings suggest that,
under co-culture conditions, *B. wadsworthia* intensifies its tryptophan uptake and protein
incorporation, potentially diminishing tryptophan availability for *B. thetaiotaomicron*'s indole
and kynurenic acid production. These observations underscore a dynamic interplay
between the two bacteria in the utilisation of tryptophan and its downstream metabolites in
the co-culture environment.

Conclusions

The increased H_2S production by *B. wadsworthia* when co-cultured with *B.*
thetaiotaomicron could have implications for human gut health. Both H_2S and *B.*
wadsworthia have been implicated in gut inflammation and pathogenesis, making it crucial
to understand the change in H_2S production. Our co-culture experiments demonstrated
enhanced growth of *B. thetaiotaomicron*, indicating a potentially beneficial syntrophic

relationship. Transcriptomic analysis of *B. wadsworthia* revealed reduced expression of

genes involved in sulfite production from taurine, thiosulfate, and sulfolactate, suggesting

an alternative sulfite source in the co-culture. Our integrated transcriptomic and genomic

data indicate a putative cross-feeding mechanism, with *B. thetaiotaomicron* potentially

providing APS to *B. wadsworthia* for sulfite production; we next demonstrated that *B.*

wadsworthia can grow to high densities using APS as an electron acceptor, substantiating

this proposed cross-feeding interaction. Gene expression data suggested an increased

uptake of lactate in *B. wadsworthia*, which could be used as an electron donor in

dissimilatory sulfite reduction. Beyond sulfur metabolism, we observed alterations in

tryptophan metabolism in co-culture. *B. wadsworthia*'s increased expression of genes

associated with tryptophan uptake suggests a competition for tryptophan, potentially

impacting *B. thetaiotaomicron*'s indole production. Reduced indole production by *B.*

thetaiotaomicron in co-culture aligns with increased tryptophan utilisation by *B.*

wadsworthia, hinting at potential implications for gut health and inflammation [61, 62]. In

conclusion, the increased H_2S production by *B. wadsworthia* in co-culture, coupled with

the reduced H_2S utilisation by *B. thetaiotaomicron*, underscores a complex microbial

interaction. This proposed APS-mediated cross-feeding mechanism opens avenues for

further exploration into sulfate metabolism in the human gut. The overall shift towards

decreased indole and increased H_2S production in co-culture suggests potential pro-

inflammatory effects in the human gut environment, warranting further investigation.

Materials and Methods
B. thetaiotaomicron and *B. wadsworthia* isolation and growth

B. thetaiotaomicron QI0072 was isolated from a faecal sample donated by a healthy adult

between 50-80 years old recruited via the COMBAT study (ClinicalTrials.gov Identifier:

NCT03679533). An initial enrichment was obtained in anaerobic anaerobe basal broth

(ABB) liquid broth media (Oxoid, Thermo Fisher Scientific) supplemented with 10 mM

taurine, and the isolate was purified on Brain Heart Infusion agar (BHI, Oxoid) prepared

according to the manufacturer's instructions then supplemented with 0.5% v/v vitamin K

solution in ethanol (10 μL/L), resazurin (1 mg/L), hemin (5 mg/L) and L-cysteine

hydrochloride (0.5 g/L) (further referred as BHI+C) with 1.5% (w/v) agar. Additional

chemicals were from Merck. *B. wadsworthia* strains QI0012, QI0013, QI0014 and QI0015

were isolated from stool samples from healthy human donors recruited via the QIB Colon

Model study (ClinicalTrials.gov Identifier: NCT02653001) using ABB supplemented with 10

mM taurine as described by Sayavedra *et al.* [63]. Cell pellets were cryopreserved at -80°C

using Protect Select Anaerobe Cryopreservation tubes (Technical Service Consultants,

UK).

Bacterial co-culture experiments

BHI, de Man, Rogosa and Sharpe (MRS, Oxoid) and ABB liquid broth media were

prepared according to the manufacturer's instructions. BHI + supplements (BHI+S) was

prepared by adding hemin (10 mg/L) and yeast extract (5 g/L) to BHI broth before

autoclaving. Where required, solid media was made by the addition of 1.5% (w/v) agar to

the media before autoclaving. All media and culture vessels were maintained under

anaerobic conditions using an anaerobic cabinet (Don Whitley, UK) with materials pre-

reduced before use for at least 18 hours in an atmosphere of 5% CO_2, 10% H_2 in N_2 at

37°C. For co-culture experiments, *B. thetaiotaomicron* was grown on BHI+C, and *B.*

wadsworthia was grown on ABB supplemented with 10 mM taurine. After overnight

incubation, the second passage cultures were diluted to ~10^8 CFU/mL using estimated cell

count per optical density at 600 nm (OD_{600}) factor. OD_{600} was quantified using a

SPECTROstar Nano instrument (BMG Labtech). Co-culture experiments used anaerobic

synthetic media containing a mix of 1:1:1 (v/v) of BHI+C, MRS and Postgate C,

supplemented with 10 mM taurine (further referred to as BPM media). Postgate C

contained per litre of distilled water: sodium lactate (6 g), sodium sulphate (4.5 g),

ammonium chloride (1 g), yeast extract (1 g) potassium dihydrogen phosphate (0.5 g),

sodium citrate tri basic (0.3 g), magnesium sulphate 7-hydrate (0.06 g), iron sulphate 7-

hydrate (4 mg), calcium chloride (0.04 g), L-cysteine hydrochloride (0.5 g), and resazurin

(0.8 mg) [64].

Co-cultures were prepared by inoculating ~10^6 CFU/mL anaerobic BPM. The CFU used

for inoculum were confirmed by plating on ABB supplemented with 10 mM taurine for *B.*

wadsworthia, and BHI+C media for *B. thetaiotaomicron*. Experimental conditions included

negative control (no inoculum), monocultures and co-culture with final culture volumes of

10 mL. All cultures were performed in at least triplicate. Sub-samples were taken at 0, 2, 4,

6 and 8 h post-inoculation.

Growth of *B. wadsworthia* with different electron acceptors

415   The growth of *B. wadsworthia* under various conditions was assessed by monitoring its

416   OD_{600} after 48 hours of incubation on ABB media enriched with different electron

417   acceptors at a 1:100 dilution. These included 4 mM sodium sulfite, 10 mM adenosine 5′-

phosphosulfate (APS) sodium salt (Merck, UK), and 10 mM taurine, alongside a control containing ABB only. We used at least seven technical replicates. Differences in growth were tested using a linear model with the glmmTMB package.

Colourimetric determination of H$_2$S concentration

Determination of H$_2$S concentration in samples was performed using the methylene blue assay modified from Cline [65]. Briefly, 500 µL of bacterial cultures were taken and immediately fixed 1:1 with 5% zinc acetate and stored at -20°C. For calibration, zinc sulfide solutions were prepared in media diluted 1:200 within concentration range 0 – 40 µM. For analysis, fixed samples were diluted 1:100 in water to a final volume of 1 mL, producing a final sample dilution of 1:200. 80 µL of diamine reagent (250 mL 6M HCl, 1g N,N-dimethyl-1,4-phenylendiamine sulfate, 1.5 g FeCl$_3$.6H$_2$0) was added to all samples and standards which were then stored in the dark to allow methylene blue colour development. After 30 min, the samples were centrifuged at 13,000 x g for 5 min to pellet biomass. 300 µL of supernatant was taken for spectrophotometric absorbance measurement at 670 nm. H$_2$S concentration in µM was determined using the calibration curve as a standard. Statistical significance between culture conditions was established using unpaired t-test pair-wise comparisons and displayed graphically using GraphPad Prism 7 (GraphPad Software, Boston, USA). Results of unpaired t tests are shown, where * = $p \leq 0.05$, ** = $p \leq 0.01$, *** = $p \leq 0.001$, **** = $p \leq 0.0001$, ns = not significant ($p > 0.05$).

Absolute quantification of bacterial cells via qPCR

For quantification of bacterial cells in experimental cultures, DNA extraction was performed using the Maxwell® RSC Blood DNA kit (Promega) according to the manufacturer's protocol. Prior to extraction, 200 µL samples were boiled at 90°C for 10 min and 150 µL was transferred to a sterile microcentrifuge tube containing 30 µL of proteinase K solution and 300 µL lysis buffer. Each sample was vortexed for 10 s and incubated at 56°C for 20 min. Quantification of *B. wadsworthia* and *B. thetaiotaomicron* was performed via qPCR using KiCqStart® SYBR® Green qPCR ReadyMix™ with ROX (Sigma-Aldrich) on a StepOnePlus™ Real-Time PCR System (Thermo Fisher Scientific). Absolute quantification of gene copy numbers was performed by comparing samples to calibration standards prepared with known gene copy numbers in the range of 10 – 1 x 10^9 copies. The number of gene copies per cell was used to calculate the absolute cell counts per mL of culture. All samples and standards were assayed in triplicate. The reaction conditions and primers

452 used are shown in Table 1. Statistical significance between culture conditions was

453 established using unpaired t-test pair-wise comparisons and displayed graphically using

454 GraphPad Prism 7.

456 **Table 1**: Primers and reaction conditions used for absolute quantification of *B.*

457 *wadsworthia* and *B. thetaiotaomicron* via qPCR

Target bacterium	Primer name	Product size (bp)	Primer sequence (5' to 3') (W=A+T)	Primer conc. (nM)	Reaction conditions	Ref
B. wadsworthia	TPA-F	150	CAACGTCCCCACCATCAAGTTCTCTG	100	95°C 2 min, 40 cycles of 95°C 15 s, 62°C 30 s.	66
	TPA-R		TGAATTCGCGGAAGGAGCGAGAGGTC	100		
B. thetaiotao-micron	F_Bacter11	131	CCTWCGATGGATAGGGGTT	200	95°C 10 min, 40 cycles of 95°C 30 s, 60°C 1 min.	67
	R_Bacter08		CACGCTACTTGGCTGGTTCAG	200		

Transcriptomic sequencing and analysis

5 mL of cultures were taken at 8 h post-inoculation and cells were immediately pelleted at

15,000 x g for 2 min. The supernatant was removed, cell pellets were snap frozen on dry

ice, and stored at -80°C. RNA was extracted using the RNeasy® Mini Kit (Qiagen,

Germany) according to manufacturer's protocol with on-column DNase digestion. RNA

concentration was determined via Qubit RNA High Sensitivity kit (Thermo Fisher Scientific,

Massachusetts, USA) and Qubit 3.0 fluorometer (Life Technologies, Massachusetts, USA).

Total RNA was sequenced at Azenta Genewiz. rRNA was depleted using NEBNext rRNA

depletion kits for human/mouse/rat and bacteria species (E6310 and E7850). Samples

were sequenced using an Illumina NovaSeq 6000 instrument with 2x150 bp configuration

using a sequencing depth of 20 M paired-end reads per sample.

For differential expression analysis, raw data was analysed as previously described [68].

Briefly, reads were cleaned to remove sequencing adapters and residual ribosomal RNA

sequences using BBDuk (v 38.06, Bushnell B. - sourceforge.net/projects/bbmap/).

Cleaned reads were mapped to the QI0013 and QI0072 reference genomes using BBSplit

(v 38.06). The number of transcripts per gene was estimated using featureCounts (v.2.0)

[69]. Reference genomes were annotated with the BV-BRC comprehensive genome analysis

tool [70]. Differentially expressed genes were identified using edgeR with TMM normalisation

[71] using the Trinity RNASeq package [72], with a P value cut-off of 0.01 and a log2FC of 1

equating to a 2-fold change in gene expression. In the case of specific genes of interest

where there was no functional annotation, InterProScan [73] was used to assign predicted

protein functions based on domains where possible using the predicted amino acid

sequence. Additionally, Pathway Tools v23.0 [74] was used for preliminary pathway

enrichment analysis and pathway predictions using the reference genomes.

Endometabolome analysis via LC-MS

5 mL of cultures were taken at 8 h post-inoculation and cells were immediately pelleted at

4,000 x g for 10 min. All supernatant was removed, and cell pellets were snap-frozen on

dry ice. All LC-MS analysis was performed by Creative Proteomics, New York, USA. Prior

to analysis, bacterial pellets were thawed and 240 μL methanol added for metabolite

extraction. Samples were vortexed for 60 s, sonicated for 30 min at 4°C and stored at -

20°C for 1 h. Samples were pelleted at 12,000 x g for 15 min at 4°C. Finally, 200 μL of

supernatant and 5 μL of DL-o-Chlorophenylalanine (0.2 mg/mL) was transferred to a vial

for LC-MS analysis. QC samples were prepared by pooling all the samples in triplicate. All

samples were injected in triplicate. Separation was performed by ACQUITY UPLC

(Waters) combined with Q Exactive MS (Thermo) and screened with ESI-MS. The LC

system was comprised of ACQUITY UPLC HSS T3 (100×2.1 mm×1.8 μm) with ACQUITY

UPLC (Waters). The mobile phase was composed of solvent A (0.05% formic acid water)

and solvent B (acetonitrile) with a gradient elution (0-1 min, 5% B; 1-12.5 min, 5%-95% B;

12.5-13.5 min, 95% B; 13.5-13.6 min, 95%-5% B; 13.6-16 min, 5% B). The flow rate of the

mobile phase was 0.3 mL/min. The column temperature was maintained at 40°C, and the

sample manager temperature set at 4°C. Mass spectrometry parameters in ESI+ and ESI-

mode were as follows: ESI+: Heater Temp 300°C; Sheath Gas Flow rate, 45 arb; Aux Gas

Flow Rate, 15 arb; Sweep Gas Flow Rate, 1 arb; spray voltage, 3.0 KV; Capillary Temp,

350°C; S-Lens RF Level, 30%. ESI-: Heater Temp 300°C, Sheath Gas Flow rate, 45 arb;

Aux Gas Flow Rate, 15 arb; Sweep Gas Flow Rate, 1 arb; spray voltage, 3.2 KV; Capillary

Temp, 350°C; S-Lens RF Level, 60%.

Metabolites were identified using Compound Discoverer 3.0 (Thermo Fisher Scientific). Progenesis QI v 2.1 (Waters) was used for manual screening of the identified compounds, in order to minimise false positive identification results. Data was normalised using the Total Ion Count (TIC) method where the peak area of each metabolite was divided by the SUM of all metabolites area and then multiplied by one million.

To remove compounds with high analytical variability, compounds with RSD_{QC} >20% were discarded [75]. Endometabolomic data was standardised by calculating the concentration (µM) per 10^9 bacterial cells present as measured via qPCR, in order to account for cell density differences in monocultures and co-cultures [76]. Compounds at low concentrations across all samples (<2 µM per 10^9 cells) were removed. In positive ion mode, 65% of the identified metabolites remained after quality control (320 left from 490), and in negative mode 70% remained (267 left from 377). Given the higher peak intensity and number of reported compounds, positive mode was used for further analysis. The metabolomic data was auto-scaled and analysed using Metaboanalyst 5.0 [77], to obtain the PLS-DA for the global profile changes and Variable Importance in Projection (VIP) compounds that contribute highly to inter-condition differences, and the heatmaps showing feature clustering and inter-condition differences in relative abundance of compounds.

DNA extraction from isolates for whole genome sequencing

For whole genome sequencing, 1 mL of overnight culture was taken and centrifuged at 13,000 g for 2 min. Pellets were stored at -20°C until extraction. For extraction of high molecular weight DNA, Fire Monkey High Molecular Weight DNA (HMW-DNA) extraction kit was used according to manufacturer's protocol (Revolugen, UK). The additional overnight elution step recommended by the manufacturer was used to maximise DNA yield. DNA was quantified using Qubit High Sensitivity DNA Quantification Kit (Thermo Scientific, UK) and Qubit 3.0 fluorometer.

Whole genome sequencing

For whole genome sequencing of *B. thetaiotaomicron*, a hybrid approach using both short- and long-read sequencing was performed in-house at Quadram Institute Bioscience. For short-read sequencing, genomic DNA was normalised to 5 ng/µl with EB buffer (10 mM Tris-HCl) and sequencing was performed using an Illumina NextSeq 500 system with 2x150 bp paired-end reads. Libraries were prepared using Bead Linked Transposomes (BLT) (Illumina) and P7 and P5 Illumina 9 bp barcodes. Paired-end sequencing reads

were received as fastq files. For long-read sequencing, a minION was used (Oxford

Nanopore Technologies, Oxford, UK).

For genome assembly, short reads were cleaned using BBDuk (v 38.79) to trim the reads

and remove sequencing adaptors. Long reads were trimmed using Porechop (v 0.2.3) [78]

and a hybrid assembly was reconstructed using Unicycler (v 0.4.9) [79]. The completeness

and contamination of the assembly was checked using CheckM (v 1.0.18) [80]. Genomes

were annotated using Prokka (v 1.14.6) and BV-BRC [81, 82].

Sequence comparison of sulfur metabolism genes

For genes of interest, amino acid sequences were obtained from *B. wadsworthia* (QI0013)

and *B. thetaiotaomicron* strain 1 (QI0072) genomes in addition to those from reference

genomes *D. desulfuricans* subsp. *desulfuricans* DSM 642, *D. gigas* DSM 1382 and *D.*

alaskensis G20 via BV-BRC [70]. InterProScan [73] was used to assign predicted protein

function based on domains. Protein-protein comparison was performed using blastp on

NCBI-BLAST using default settings [83]. Amino acid alignments were performed using

Clustal omega v1.2.2 in Geneious prime v 2022.1.1 (Biomatters Ltd) and visualised with

GeneDoc v 2.7.000 (National Resource for Biomedical Supercomputing).

Ethics statement

The colon model study was approved by the Quadram Institute Bioscience Human

Research Governance Committee (IFR01/2015), and London - Westminster Research

Ethics Committee (15/LO/2169). The COMBAT study was approved by the University of

East Anglia's Faculty of Medicine and Health Sciences Ethical Review Committee

(Reference: 201819–039) and the Health Research Authority (IRAS number: 237251). The

colon model study was registered under the ClinicalTrials.gov Identifier NCT02653001 and

the COMBAT was registered under the NCT03679533.

Acknowledgements
571

572 The authors thank David Baker (Quadram Institute Bioscience) for library preparation and

573 whole genome sequencing.

574 # Funding
575

This work was supported by UKRI-BBSRC via the Norwich Research Park Doctoral
Training Partnership (grant no. BB/M011216/1) and institute strategic programme grants
Gut Microbes and Health BB/R012490/1 theme BBS/E/F/000PR10356 and Food,
Microbiome and Health BB/X011054/1 theme BBS/E/F/000PR13633. UKRI-BBSRC had
no role in the manner of conduct or outcome of the research project. Transcriptomic
sequencing was funded through a Quadram Institute Bioscience Institute Development
Grant (20414000X). LS was supported by a BBSRC Discovery Fellowship
(BB/Z514445/10).

Data availability

586 Transcriptomic sequencing data was submitted to NCBI under PRJNA1115121. The
587 genome of *Bilophila wadsworthia* QI0013 is available under project PRJNA1085689. The
588 genome of *Bacteroides thetaiotaomicron* was submitted under project PRJNA1151643.

Authors' contributions

591 JD, LS and AN conceived and designed the project. JD and LS designed the experiments.
592 JD performed experiments and analysed the data. LS supervised transcriptomic analyses.
593 JD and LS wrote the manuscript with contributions from all co-authors.

Conflict of interest
596 The authors declare that they have no conflicts of interest.

References

1. Stewart J, Chadwick V, Murray A. Carriage, quantification, and predominance of
methanogens and sulfate-reducing bacteria in faecal samples. Letters in Applied
Microbiology 2006; 43:58-63.
2. Rey Federico E, Gonzalez Mark D, Cheng J, Wu M, Ahern Philip P, Gordon Jeffrey
I. Metabolic niche of a prominent sulfate-reducing human gut bacterium.
Proceedings of the National Academy of Sciences 2013; 110:13582-7.
3. Santos A, Venceslau S, Grein F, Leavitt W, Dahl C, Johnston D, et al. A protein
trisulfide couples dissimilatory sulfate reduction to energy conservation. Science
2015; 350:1541.
4. Mathai JC, Missner A, Kügler P, Saparov SM, Zeidel ML, Lee JK, et al. No facilitator
required for membrane transport of hydrogen sulfide. Proceedings of the National
Academy of Sciences 2009; 106:16633-8.
5. Guidotti T. Hydrogen Sulphide. Occupational Medicine 1996; 46:367-71.

6. Parey K, Demmer U, Warkentin E, Wynen A, Ermler U, Dahl C. Structural,
biochemical and genetic characterization of dissimilatory ATP sulfurylase from
Allochromatium vinosum. PLoS One 2013; 8:e74707.
7. Denger K, Laue H, Cook A. Anaerobic taurine oxidation: a novel reaction by a
nitrate-reducing *Alcaligenes* sp. Microbiology 1997; 143:1919-24.
8. Baron E, Summanen P, Downes J, Roberts M, Wexler H, Finegold S. *Bilophila*
wadsworthia, gen. nov. and sp. nov., a unique Gram-negative anaerobic rod
recovered from appendicitis specimens and human faeces. Microbiology 1989;
135:3405-11.
9. Baron E. *Bilophila wadsworthia*: a Unique Gram-negative Anaerobic Rod. Anaerobe
1997; 3:83-6.
10. Peck S, Denger K, Burrichter A, Irwin S, Balskus E, Schleheck D. A glycyl radical
enzyme enables hydrogen sulfide production by the human intestinal bacterium
Bilophila wadsworthia. Proceedings of the National Academy of Sciences 2019;
116:3171.
11. Hanson B, Kits K, Löffler J, Burrichter A, Fiedler A, Denger K, et al. Sulfoquinovose
is a select nutrient of prominent bacteria and a source of hydrogen sulfide in the
human gut. The ISME Journal 2020.
12. Kushkevych I, Cejnar J, Treml J, Dordević D, Kollar P, Vítězová M. Recent
advances in metabolic pathways of sulfate reduction in intestinal bacteria. Cells
2020; 9.
13. Scanlan P, Shanahan F, Marchesi J. Culture-independent analysis of desulfovibrios
in the human distal colon of healthy, colorectal cancer and polypectomized
individuals. FEMS Microbiology Ecology 2009; 69:213-21.
14. Baron E, Curren M, Henderson G, Jousimies-Somer H, Lee K, Lechowitz K, et al.
Bilophila wadsworthia isolates from clinical specimens. Journal of Clinical
Microbiology 1992; 30:1882-4.
15. Finegold S, P. S, Hunt Gerardo S., E. B. Clinical Importance of *Bilophila*
wadsworthia. European Journal of Clinical Microbiology and Infectious Diseases
1992; 11:1058-63.
16. Mosca A, D'Alagni M, del Prete R, de Michele G, Summanen P, Finegold S, et al.
Preliminary evidence of endotoxic activity of *Bilophila wadsworthia*. Anaerobe 1995;
1:21-4.
17. Hunt Gerardo S, Garcia M, Wexler H, Finegold S. Adherence of *Bilophila*
wadsworthia to cultured human embryonic intestinal cells. Anaerobe 1998; 4:19-27.
18. Yazici C, Wolf P, Carroll T, Mutlu E, Xicola R, Llor X, et al. *Bilophila wadsworthia* Is
more abundant in the colonic microbiome of colorectal cancer cases compared to
healthy controls. Gastroenterology 2015; 148:S-100.
19. Tremlett H, Fadrosh D, Faruqi A, Zhu F, Hart J, Roalstad S, et al. Gut microbiota in
early pediatric multiple sclerosis: a case-control study. Eur J Neurol 2016; 23:1308-
21.
20. Baldini F, Hertel J, Sandt E, Thinnes CC, Neuberger-Castillo L, Pavelka L, et al.
Parkinson's disease-associated alterations of the gut microbiome predict disease-
relevant changes in metabolic functions. BMC Biology 2020; 18:62.
21. Sheng C, Yang K, He B, Du W, Cai Y, Han Y. Combination of gut microbiota and
plasma amyloid-β as a potential index for identifying preclinical Alzheimer's
disease: a cross-sectional analysis from the SILCODE study. Alzheimer's Research
& Therapy 2022; 14:35.
22. Jiao N, Baker SS, Chapa-Rodriguez A, Liu W, Nugent CA, Tsompana M, et al.
Suppressed hepatic bile acid signalling despite elevated production of primary and
secondary bile acids in NAFLD. Gut 2018; 67:1881.

23. Li G-H, Huang S-J, Li X, Liu X-S, Du Q-L. Response of gut microbiota to serum
metabolome changes in intrahepatic cholestasis of pregnant patients. World journal
of gastroenterology 2020; 26:7338-51.
24. He X, Sun J, Liu C, Yu X, Li H, Zhang W, et al. Compositional alterations of gut
microbiota in patients with diabetic kidney disease and type 2 diabetes mellitus.
Diabetes, Metabolic Syndrome and Obesity 2022; 15:755-65.
25. Thirion F, Speyer H, Hansen TH, Nielsen T, Fan Y, Le Chatelier E, et al. Alteration
of gut microbiome in patients with schizophrenia indicates links between bacterial
tyrosine biosynthesis and cognitive dysfunction. Biological Psychiatry Global Open
Science 2022.
26. Feng Z, Long W, Hao B, Ding D, Ma X, Zhao L, et al. A human stool-derived
Bilophila wadsworthia strain caused systemic inflammation in specific-pathogen-
free mice. Gut Pathogens 2017; 9:59-.
27. Xu J, Bjursell Magnus K, Himrod J, Deng S, Carmichael Lynn K, Chiang Herbert C,
et al. A genomic view of the human-*Bacteroides thetaiotaomicron* symbiosis.
Science 2003; 299:2074-6.
28. Hong P-Y, Wu J-H, Liu W-T. Relative abundance of *Bacteroides* spp. in stools and
wastewaters as determined by hierarchical oligonucleotide primer extension.
Applied and Environmental Microbiology 2008; 74:2882-93.
29. Human Microbiome Project C. Structure, function and diversity of the healthy
human microbiome. Nature 2012; 486:207-14.
30. Garber JM, Hennet T, Szymanski CM. Significance of fucose in intestinal health and
disease. Molecular Microbiology 2021; 115:1086-93.
31. Freitas M, Axelsson L-G, Cayuela C, Midtvedt T, Trugnan G. Indigenous microbes
and their soluble factors differentially modulate intestinal glycosylation steps in vivo.
Histochemistry and Cell Biology 2005; 124:423-33.
32. Hooper LV, Gordon JI. Glycans as legislators of host–microbial interactions:
spanning the spectrum from symbiosis to pathogenicity. Glycobiology 2001; 11:1R-
10R.
33. Chia LW, Mank M, Blijenberg B, Aalvink S, Bongers RS, Stahl B, et al. *Bacteroides
thetaiotaomicron* fosters the growth of butyrate-producing *Anaerostipes caccae* in
the presence of lactose and total human milk carbohydrates. Microorganisms 2020;
8.
34. Chia LW, Mank M, Blijenberg B, Aalvink S, Bongers RS, Stahl B, et al. Bacteroides
thetaiotaomicron Fosters the Growth of Butyrate-Producing Anaerostipes caccae in
the Presence of Lactose and Total Human Milk Carbohydrates. Microorganisms,
2020.
35. Porter Nathan T, Larsbrink J, Avci Fikri Y. Investigation and alteration of organic
acid synthesis pathways in the mammalian gut symbiont *Bacteroides
thetaiotaomicron*. Microbiology Spectrum; 10:e02312-21.
36. Ye M, Yu J, Shi X, Zhu J, Gao X, Liu W. Polysaccharides catabolism by the human
gut bacterium *Bacteroides thetaiotaomicron*: advances and perspectives. Critical
Reviews in Food Science and Nutrition 2021; 61:3569-88.
37. Dordević D, Jančíková S, Vítězová M, Kushkevych I. Hydrogen sulfide toxicity in the
gut environment: Meta-analysis of sulfate-reducing and lactic acid bacteria in
inflammatory processes. Journal of Advanced Research 2021; 27:55-69.
38. da Silva S, Venceslau S, Fernandes C, Valente F, Pereira I. Hydrogen as an energy
source for the human pathogen *Bilophila wadsworthia*. Antonie van Leeuwenhoek
2008; 93:381-90.
39. Khelaifia S, Lagier JC, Nkamga VD, Guilhot E, Drancourt M, Raoult D. Aerobic
culture of methanogenic archaea without an external source of hydrogen. European
Journal of Clinical Microbiology & Infectious Diseases 2016; 35:985-91.

40. Keon RG, Fu R, Voordouw G. Deletion of two downstream genes alters expression
of the hmc operon of Desulfovibrio vulgaris subsp. vulgaris Hildenborough. Archives
of Microbiology 1997; 167:376-83.
41. Ichikawa Y, Yamamoto H, Hirano S-i, Sato B, Takefuji Y, Satoh F. The overlooked
benefits of hydrogen-producing bacteria. Medical Gas Research 2023; 13:108-11.
42. Muyzer G, Stams A. The ecology and biotechnology of sulphate-reducing bacteria.
Nature Reviews Microbiology 2008; 6:441-54.
43. Qian Z, Tianwei H, Mackey HR, van Loosdrecht MCM, Guanghao C. Recent
advances in dissimilatory sulfate reduction: from metabolic study to application.
Water Research 2019; 150:162-81.
44. Duarte AG, Barbosa ACC, Ferreira D, Manteigas G, Domingos RM, Pereira IAC.
Redox loops in anaerobic respiration - The role of the widespread NrfD protein
family and associated dimeric redox module. Biochimica et Biophysica Acta (BBA) -
Bioenergetics 2021; 1862:148416.
45. Burrichter AG, Dörr S, Bergmann P, Haiß S, Keller A, Fournier C, et al. Bacterial
microcompartments for isethionate desulfonation in the taurine-degrading human-
gut bacterium *Bilophila wadsworthia*. BMC Microbiology 2021; 21:340.
46. Luis AS, Jin C, Pereira GV, Glowacki RWP, Gugel SR, Singh S, et al. A single
sulfatase is required to access colonic mucin by a gut bacterium. Nature 2021;
598:332-7.
47. Rey FE, Gonzalez MD, Cheng JY, Wu M, Ahern PP, Gordon JI. Metabolic niche of
a prominent sulfate-reducing human gut bacterium. Proceedings of the National
Academy of Sciences 2013; 110:13582-7.
48. Benoni R, De Bei O, Paredi G, Hayes CS, Franko N, Mozzarelli A, et al. Modulation
of *Escherichia coli* serine acetyltransferase catalytic activity in the cysteine synthase
complex. FEBS Letters 2017; 591:1212-24.
49. Guédon E, Martin-Verstraete I. Cysteine metabolism and its regulation in bacteria.
Amino acid biosynthesis~ pathways, regulation and metabolic engineering:
Springer, 2006:195-218.
50. Kulikova VV, Revtovich SV, Bazhulina NP, Anufrieva NV, Kotlov MI, Koval VS, et al.
Identification of O-acetylhomoserine sulfhydrylase, a putative enzyme responsible
for methionine biosynthesis in *Clostridioides difficile*: gene cloning and biochemical
characterizations. IUBMB Life 2019; 71:1815-23.
51. Dalgliesh C. Biological degradation of tryptophan. Quarterly Reviews, Chemical
Society 1951; 5:227-44.
52. Kuc D, Zgrajka W, Parada-Turska J, Urbanik-Sypniewska T, Turski WA. Micromolar
concentration of kynurenic acid in rat small intestine. Amino Acids 2008; 35:503-5.
53. Turski MP, Turska M, Paluszkiewicz P, Parada-Turska J, Oxenkrug GF. Kynurenic
acid in the digestive system–-new facts, new challenges. International Journal of
Tryptophan Research 2013; 6:IJTR.S12536.
54. Johnson EL, Heaver SL, Waters JL, Kim BI, Bretin A, Goodman AL, et al.
Sphingolipids produced by gut bacteria enter host metabolic pathways impacting
ceramide levels. Nature Communications 2020; 11:2471.
55. Neis EP, Dejong CH, Rensen SS. The role of microbial amino acid metabolism in
host metabolism. Nutrients 2015; 7:2930-46.
56. Lee J-H, Wood TK, Lee J. Roles of indole as an interspecies and interkingdom
signaling molecule. Trends in microbiology 2015; 23 11:707-18.
57. Smith EA, Macfarlane GT. Formation of phenolic and indolic compounds by
anaerobic bacteria in the human large intestine. Microbial Ecology 1997; 33:180-8.
58. Li G, Young KD. Indole production by the tryptophanase TnaA in *Escherichia coli* is
determined by the amount of exogenous tryptophan. Microbiology 2013; 159:402-
10.

59.   Gorelik O, Rogad A, Holoidovsky L, Meijler MM, Sal-Man N. Indole intercepts the
769   communication between enteropathogenic *E. coli* and *Vibrio cholerae*. Gut
770   Microbes 2022; 14:2138677.

771   60.   Claros MC, Schumacher U, Jacob M, Hunt Gerardo S, Kleinkauf N, Goldstein EJC,
772   et al. Characterization of *Bilophila wadsworthia* isolates using PCR fingerprinting.
773   Anaerobe 1999; 5:589-93.

774   61.   Bansal T, Alaniz RC, Wood TK, Jayaraman A. The bacterial signal indole increases
775   epithelial-cell tight-junction resistance and attenuates indicators of inflammation.
776   Proceedings of the national academy of sciences 2010; 107:228-33.

777   62.   Whitfield-Cargile CM, Cohen ND, Chapkin RS, Weeks BR, Davidson LA, Goldsby
778   JS, et al. The microbiota-derived metabolite indole decreases mucosal inflammation
779   and injury in a murine model of NSAID enteropathy. Gut microbes 2016; 7:246-61.

780   63.   Sayavedra L, Yasir M, Goldson A, Brion A, Le Gall G, Moreno-Gonzalez M, et al.
781   Energy generation drives gut colonization by *Bilophila wadsworthia*. Research
782   Square 2024.

783   64.   Postgate John R. Versatile medium for the enumeration of sulfate-reducing
784   bacteria. Applied Microbiology 1963; 11:265-7.

785   65.   Cline J. Spectrophotometric determination of hydrogen sulphide in natural waters.
786   Limnology and Oceanography 1969; 14:454-8.

787   66.   Natividad J, Lamas B, Pham H, Michel M, Rainteau D, Bridonneau C, et al.
788   *Bilophila wadsworthia* aggravates high fat diet induced metabolic dysfunctions in
789   mice. Nature Communications 2018; 9:2802-.

790   67.   Furet J-P, Firmesse O, Gourmelon M, Bridonneau C, Tap J, Mondot S, et al.
791   Comparative assessment of human and farm animal faecal microbiota using real-
792   time quantitative PCR. FEMS Microbiology Ecology 2009; 68:351-62.

793   68.   Sayavedra L, Li T, Bueno Batista M, Seah BKB, Booth C, Zhai Q, et al.
794   *Desulfovibrio diazotrophicus* sp. nov., a sulfate-reducing bacterium from the human
795   gut capable of nitrogen fixation. Environmental Microbiology 2021; 23:3164-81.

796   69.   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for
797   assigning sequence reads to genomic features. Bioinformatics 2014; 30:923-30.

798   70.   Olson RD, Assaf R, Brettin T, Conrad N, Cucinell C, Davis JJ, et al. Introducing the
799   bacterial and viral bioinformatics resource center (BV-BRC): a resource combining
800   PATRIC, IRD and ViPR. Nucleic acids research 2023; 51:D678-D89.

801   71.   Robinson MD, Oshlack A. A scaling normalization method for differential expression
802   analysis of RNA-seq data. Genome Biology 2010; 11:R25.

803   72.   Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Full-
804   length transcriptome assembly from RNA-Seq data without a reference genome.
805   Nature Biotechnology 2011; 29:644-52.

806   73.   Blum M, Chang H-Y, Chuguransky S, Grego T, Kandasaamy S, Mitchell A, et al.
807   The InterPro protein families and domains database: 20 years on. Nucleic Acids
808   Research 2021; 49:D344-D54.

809   74.   Karp PD, Midford PE, Billington R, Kothari A, Krummenacker M, Latendresse M, et
810   al. Pathway Tools version 23.0 update: software for pathway/genome informatics
811   and systems biology. Briefings in Bioinformatics 2021; 22:109-26.

812   75.   Sánchez-Illana Á, Piñeiro-Ramos JD, Sanjuan-Herráez JD, Vento M, Quintás G,
813   Kuligowski J. Evaluation of batch effect elimination using quality control replicates in
814   LC-MS metabolite profiling. Analytica Chimica Acta 2018; 1019:38-48.

815   76.   Zhang J, Luo W, Wang Z, Chen X, Lv P, Xu J. A novel strategy for D-psicose and
816   lipase co-production using a co-culture system of engineered *Bacillus subtilis* and
817   *Escherichia coli* and bioprocess analysis using metabolomics. Bioresources and
818   Bioprocessing 2021; 8:77.

819   77.   Pang Z, Chong J, Zhou G, de Lima Morais DA, Chang L, Barrette M, et al.
820   MetaboAnalyst 5.0: narrowing the gap between raw spectra and functional insights.
821   Nucleic Acids Research 2021; 49:W388-W96.
822   78.   Wick RR, Judd LM, Gorrie CL, Holt KE. Completing bacterial genome assemblies
823   with multiplex MinION sequencing. Microbial genomics 2017; 3.
824   79.   Wick RR, Judd LM, Gorrie CL, Holt KE. Unicycler: resolving bacterial genome
825   assemblies from short and long sequencing reads. PLOS Computational Biology
826   2017; 13:e1005595.
827   80.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM:
828   assessing the quality of microbial genomes recovered from isolates, single cells,
829   and metagenomes. Genome Research 2015; 25:1043-55.
830   81.   Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics 2014;
831   30:2068-9.
832   82.   Olson RD, Assaf R, Brettin T, Conrad N, Cucinell C, Davis JJ, et al. Introducing the
833   Bacterial and Viral Bioinformatics Resource Center (BV-BRC): a resource
834   combining PATRIC, IRD and ViPR. Nucleic Acids Research 2023; 51:D678-d89.
835   83.   Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, et al. Gapped
836   BLAST and PSI-BLAST: A new generation of protein database search programs.
837   Nucleic Acids Research 1997; 25:3389-402.
838

Figure 1: Co-culture of *B. wadsworthia* (Bw) QI0013 and *B. thetaiotaomicron* (Bt) QI0072. a) H_2S concentrations (µM) at 8 h. b-c) Cell counts of *B. wadsworthia* or *B. thetaiotaomicron* during culture measured via qPCR. d) H_2S concentration (µM) per 10^6 *B. wadsworthia* cells at 8 h. Each point represents a culture replicate (n=7). In the negative control bacterial cells were not added. Horizontal lines represent average, and error bars represent SD. Results of unpaired t-tests are shown where ** = p ≤ 0.01, *** = p ≤ 0.001, **** = p ≤ 0.0001.

a) *Bilophila wadsworthia*

b) *Bacteroides thethaiotamicron*

Figure 2: Differentially expressed genes (DEGs) of *B. wadsworthia* and *B. thetaiotaomicron* in co-culture versus monoculture. a) DEGs in *B. wadsworthia*. b) DEGs in *B. thetaiotaomicron*. Bar charts show numbers of DEGs increased (green) and decreased (red) in expression in co-culture relative to the respective monoculture in each functional gene class as annotated in BV-BRC.

Figure 3: Model of cross-feeding interaction between *B. wadsworthia* and *B. thetaiotaomicron*. Enzymes are shown in colours corresponding to transcriptomic data: green showed increased gene expression in co-culture; red showed decreased expression in co-culture; grey showed no change in expression between co-culture and monoculture. The blue star represents a putative enzyme where functionality is unproven. The dotted line represents a putative cross-feeding mechanism. Dashed lines represent electron transfer.

Figure 4: Growth (OD$_{600}$) of *B. wadsworthia* in ABB media supplemented with 10 mM adenosine 5'-phosphosulfate (APS), 4 mM sulfite or 10 mM taurine. Growth was compared to *B. wadsworthia* grown in ABB media. Each box plot represents the median and interquartile range of the distribution of 7 culture replicate. Results of mixed linear model analyses are shown where $* = p \leq 0.01$, $** = p \leq 0.001$.

Figure 5: Pairwise co-culture of Bt strain 1 with four *B. wadsworthia* strains (QI0012, QI0013, QI0014, QI0015). a) H_2S concentrations (µM) at 8 h. b) qPCR-determined *B. wadsworthia* cell counts at 8 h. c) qPCR-determined *B. thetaiotaomicron* cell counts at 8 h. d) H_2S concentration (µM) per 10^6 *B. wadsworthia* cells. Each point represents a culture replicate (n=3). Horizontal lines represent average, and error bars represent SD. Statistical significance between culture conditions was established using one-way analysis of variance (ANOVA) with Tukey's multiple comparison tests with a significance level set at α = 0.05. Results show * = p ≤ 0.05, ** = p ≤ 0.01, *** = p ≤ 0.001, **** = p ≤ 0.0001, ns = not significant (p > 0.05).

a)

b)

c)

Relative concentration (µM) per 10⁹ *Bacteroides thehtaiotamicron* cells

Figure 6: Comparisons of the endometabolome of *B. wadsworthia* and *B. thetaiotaomicron* in co-culture (Bw + Bt) with monocultures (Bw_mono, Bt_mono). a) The top compounds ranked based on the Variable Importance in Projection (VIP) scores. The coloured boxes on the right indicate the relative concentrations of the corresponding metabolite in each group. b) Specific metabolites of interest manually curated from the compound list detected via untargeted LC-MS. c) Relative intracellular concentrations of compounds related to tryptophan metabolism in *B. thetaiotaomicron* (Bt) in monoculture and co-culture with *B. wadsworthia* (Bw + Bt). Concentration is standardised to *B. thetaiotaomicron* cell counts. Each point represents a culture replicate. Box and whisker plots show line at mean, box at 25th – 75th percentile, whiskers to minimum and maximum.

Supplementary data

Figure S1: Metabolic pathways for sulfate reduction Dissimilatory sulfate reduction is a strictly anaerobic process performed by sulfate-reducing bacteria (SRB). Toxic isethionate intermediates are encapsulated within microcompartments in *B. wadsworthia* [1].

Figure S2: Alignment of amino acid sequences of A) AprA genes and B) AprB genes from *B. wadsworthia* QI0013 (Bwadsw), *Desulfovibrio gigas* DSM 1382 (Dgigas), *Desulfovibrio alaskensis* G20 (Dalask) and *Desulfovibrio desulfuricans* subsp. desulfuricans DSM 642 (Ddesul).

Figure S3: Comparisons of the endometabolome of *B. wadsworthia* and *B. thetaiotaomicron* in co-culture (Bw + Bt) with monocultures (Bw_mono, Bt_mono). a) PLS-DA plot of metabolites acquired from samples via untargeted LC-MS in positive ion mode. b) Heatmap displaying relative abundance of top 50 differentially abundant metabolites in the culture conditions.

References:

1   Burrichter, A. G. *et al.* Bacterial microcompartments for isethionate desulfonation in the taurine-degrading human-gut bacterium *Bilophila wadsworthia*. *BMC Microbiol.* **21**, 340 (2021). https://doi.org:10.1186/s12866-021-02386-w