Title: *In vitro* inhibition of the CFTR ion channel in the cervix thickens cervical mucus

Running Title: CFTR inhibition thickens cervical mucus in vitro

Authorship: Rachel J. Looney[1], Mackenzie Roberts MPH[1], Matthew Markovetz PhD[2], Rachelle Godiah

MPH[1], Shan Yao M.D[1], Kirsti Golgotiu[3], Shuhao Wei[1], Chris Cellucci[1], Leo Han M.D. MPH[1,4]

[1]Division of Reproductive and Developmental Sciences, Oregon National Primate Research Center,

Beaverton, OR, USA

[2]Marsico Lung Institute, University of North Carolina, Chapel Hill, NC, USA

[3]Department of Biomedical Engineering, Oregon Health and Science University, Portland, OR, USA

[4]Department of Obstetrics and Gynecology, Oregon Health and Science University, Portland, OR, USA

Corresponding Author:

Leo Han

Oregon National Primate Research Center

505 NW 185[th] Ave Rm 2116 Beaverton OR 97006

hanl@ohsu.edu

Key Words: CFTR, Cervix, Mucus, Fertility, Inh-172, ASL, PTMR

Word Count: 4809

Abstract

Study question: Does the inhibition of Cystic Fibrosis Transmembrane Conductance Regulator (CFTR)

alter cervical mucus produced by endocervical cells *in vitro*?

Summary answer: Inhibition of CFTR dehydrates and thickens *in vitro* endocervical secretions.

What is known already: Cervical mucus changes throughout the menstrual cycle in response to

hormonal fluctuations, helping to regulate fertility and access of pathogens to the reproductive tract.

CFTR is an anion channel that plays a critical role in mediating epithelial mucus secretions.

Study design, size, duration: *In vitro* experiments conducted with control and experimental arms.

Participants/materials, setting, methods: Primary endocervical cells obtained from rhesus macaques

Macaca mulatta were treated with vehicle controls or CFTR inhibitor Inh-172. Hydration and viscosity of

secreted mucus was measured through air surface liquid and particle tracking microrheology.

Main results and the role of chance: Endocervical cells treated with CFTR inhibitor Inh-172

demonstrated dehydrated, thicker mucus secretions compared to control ($p < .01$).

Limitations, reasons for caution: *In vitro* models do not fully recapitulate cellular mucus secretions.

Wider implications of the findings: Cervical mucus plays an important role in fertility regulation. Our

studies suggest that CFTR may be an important mediator of those changes and provide experimental

evidence for the infertility phenotype seen in women with cystic fibrosis. Additionally, assays used in

these studies are useful endpoints for studying cervical mucus regulation *in vitro*.

Study funding/competing interest(s): The Bill and Melinda Gates Foundation, The Cystic Fibrosis

Foundation, Oregon National Primate Research Center, open access

Introduction

Epithelial cells, lining the endocervical canal, control sperm and pathogen entry into the female

reproductive tract through cervical mucus secretion. Over the course of the menstrual cycle, fluctuations

in ovarian sex-steroids, estradiol (E2) and progesterone (P4), regulate the secreted mucus (Moghissi *et*

al., 1972; Najmabadi *et al.*, 2021). When E2 levels peak during the ovulation window, cervical mucus

increases in quantity and hydration, and decreases in viscosity, creating a permeable gel that allows

sperm to move through to the upper tract. Under the influence of P4, following ovulation during the

luteal phase, cervical mucus increases in concentration, becomes more viscous, and blocks sperm from

entering the reproductive tract (Moghissi, 1972).

Cervical mucus is a heterogenous mixture of water, mucin and non-mucin proteins, lipids,

carbohydrates, and ions. Outside the cervix, other epithelia, including the lung, GI tract, and eye, also

produce mucus that maintain organ health and functions. Mucins and proteins found in mucus are

retained across organ systems (Thornton *et al.*, 2008). The Cystic Fibrosis Transmembrane Conductance

Regulator (CFTR) ion channel has been shown to be a critical regulator of mucus characteristics in all

mucus secreting epithelia. Mutations in the CFTR gene lead to dysfunctional channel activity and the

autosomal recessive disease, cystic fibrosis (CF). CF is characterized by systemically thickened secretions

that can cause multi-organ dysfunction in the airway, GI tract, and reproductive tract. In the female

reproductive tract, CF is notable for thickened endocervical mucus as well as sub-fertile and infertile

phenotypes (Anthony *et al.*, 1970; Kopito *et al.*, 1973).

The CFTR channel moves chloride and bicarbonate (ions) into the lumen, promoting hydration

and the thinning of mucus through several mechanisms. In conjunction with the epithelial sodium

channel (ENaC) and other apical ion channels, CFTR maintains an ionic gradient for water to move

osmotically (Saint-Criq and Gray, 2017). CFTR has also been shown to play a critical role in hydration of

mucins. Intracellularly, large mucin glycoproteins are tightly packaged, with cations shielding their

negatively charged carbohydrate-rich ends. Upon release, luminal anions remove the cations resulting in

the expansion and unfolding of the mucins. In their expanded form, mucins readily bind water

(Muchekehu and Quinton, 2010). When CFTR is dysfunctional, mucins are unable to fully expand and

bind water, leading to the dehydrated, thickened mucus seen in CF (Clunes and Boucher, 2007).

Previous studies have demonstrated that CFTR in the cervix is both functionally and hormonally

regulated (Han *et al.*, 2022). While these findings may suggest that CFTR plays a role in regulating mucus

changes in the cervix, it remains unclear if CFTR directly alters mucus consistency and hydration. CFTR

inhibition of endocervical cells and its effect on mucosal thickness and hydration has yet to be

demonstrated in an experimental model. Elucidating the function of CFTR in cervical mucus regulation

would identify its role as a target for both fertility and contraceptive studies in the cervix.

Prior studies using 2D cultures of the endocervix lack experimental endpoints for measuring

mucus changes in response to perturbation experiments. To overcome this limitation, we adapted from

airway studies of mucus, airway surface liquid (ASL) as a method for measuring mucus hydration *in vitro*.

ASL measurements correlate tightly to interfacial hydration *in vivo* (Tarran and Boucher, 2002; Harvey *et*

al., 2011; Haq *et al.*, 2016). ASL hydration in normal airway epithelia (Tarran and Boucher, 2002;

Myerburg *et al.*, 2006) is lost in CF cells due to mucosal dehydration (Matsui *et al.*, 1998) and can be

restored in mutant cell lines with novel CFTR modulator therapies (Haq *et al.*, 2016; Berg *et al.*, 2019).

While ASL can be directly measured using confocal microscopy in the XZ axis (Tarran and Boucher, 2002),

ASL can also be assessed by measuring the edge-meniscus formed under brightfield microscopy that

allows 96-well experimental measurement and replication (Harvey *et al.*, 2011). Brightfield meniscus

width is a valid measurement of ASL and can be used for high throughput studies of CFTR modulator

therapies (Berg *et al.*, 2019).

While ASL is a measure of liquid-ion transport functionality, it does not provide a direct readout

of the biophysical status of the mucus in the lumen. Multiple studies have shown that mucus

concentration is the primary driver of aberrant biophysical properties (i.e. viscoelasticity) observed in CF

(Button *et al.*, 2012; Hill *et al.*, 2018). This has been demonstrated with classical rheometric and

microscale techniques relevant to the biophysics experienced by sperm within mucus, generally termed

particle tracking microrheology (PTMR). PTMR exploits the thermal diffusion of small (~1 μm)

fluorescent probes to sense the viscoelasticity of the mucus network.

In this investigation, we utilized both ASL measurements of mucus hydration and PTMR

measurements of viscosity in response to hormonal changes and CFTR inhibition. We hypothesized that

selectively inhibiting CFTR with CFTR Inhibtor-172 (Inh-172) would result in decreased mucosal

hydration, and increased mucus viscosity in vitro.

Methods

Primary cell culture

Rhesus macaque cervical tissue was obtained from the Oregon National Primate Research

Center tissue distribution program. Endocervical epithelia were scalpel separated from the cervical

stroma, enzymatically digested, and cultured using conditionally reprogramming conditions as

previously described (CRECs) (Han *et al.*, 2020). Expanded cells were passaged at 80% and placed onto

HTS Transwell® 96-well and 12-well Permeable Supports (0. 4 μm)(Corning, New York) in a serum-free

media (ReproLife CX Lifeline Cell Technology, Frederick, Maryland), supplemented with 1M calcium

chloride and 10nM E2 and taken to an air-liquid interface (ALI).

Airway surface Liquid

After 10-14 days at an ALI (when transepithelial electrical resistance >300 Ω), we performed the

air-surface liquid assay on treatment naïve cells. For ASL, we adapted an assay variation used for high-

throughput screening in human bronchial epithelial (HBE) cultures that measures the resorption of

added saline to the apical (mucus) side of ALI culture (Berg *et al.*, 2019). Cultures that create more

dehydrated mucus resorb more saline over time. To do this, we used two different versions of the

resorption assay. In a version similar to the previously published HTS, phosphate-buffered saline (PBS; 8

μL per well) (Thermo-Fisher Scientific) was added to the apical surface and images were taken using

brightfield microscopy (2x PlanApoλ NA 0.10 magnification, BZ-X Series, Keyence, Itastca, Illinois).

Images were taken pre-treatment, immediately post-treatment (0 hours), 24, 48, 72, and 96 hours.

We additionally assessed these changes using an apical addition of 8 μL per well of 100 mM

mannitol in PBS. Mannitol is a sugar alcohol that demonstrates osmotic based expansion of ASL in HBE

cells and has been studied in clinical trials as potential treatment for CF lung disease (Jaques *et al.*, 2008;

Keyrouz *et al.*, 2008). We used this combined method of assessing ASL (mannitol+ PBS) as we found that

it more robustly defined changes between controls and treatment in ASL studies (Locke *et al.*, 2014).

Following image acquisition, we used custom FIJI (Schindelin *et al.*, 2012) scripts to automatically

calculate the change in fluid meniscus. The scripts use pixel thresholding to determine the size of the

meniscus by measuring the annulus; the annulus is the area of dark fluid meniscus that emanates from

the edge of the well to the lighter center region (Figure 1A,B). This area increases with greater apical

hydration and decreases with reduced hydration. The metrics were reported as magnitude change from

baseline (time=0 hours) in downstream analysis and statistics. More negative values correspond to

greater reduction in the meniscus from the 0Hr time point (time of apical application of PBS). Thus,

larger change in meniscal reduction indicates increased dehydration of the cell. As part of our custom

scripts, we additionally adjusted for overlying artifacts (i.e. overlaying cellular mucus or debris) in the cell

cultures using thresholding automatic identification of outliers based on a pre-determined 95%

confidence interval. Using the 96-well plate, treatment conditions were repeated using two different

plates to account for edge effects with a total of either 48 or 96 wells per condition. Experiments

represent at least three or more separate primary cell culture experiments.

Air Surface Liquid Baseline Assays

We began by demonstrating the ability of our assay to measure apical hydration using mannitol

only treatments (Myerburg *et al.*, 2010). Cells were treated with vehicle or mannitol (100mM and

300mM) in PBS (8 μL) applied apically on day 1. We also sought to demonstrate that ASL measurements

responded to hormonal regulation of mucus. Clinically, E2 peaks during ovulation and results in thinner,

hydrated mucus. During the luteal phase, P4 dominates, and mucus becomes thicker and more

dehydrated. Thus, we would expect E2 to increase ASL relative to control, and P4 to decrease ASL

relative to control (Moghissi, 1972). To do this, we treated differentiated cells with E2 (10 nM) or E2/P4

(10 – 14 days of 10 nm E2 priming followed by 1nM E2 and 100 nM P4) basolaterally and compared the

responses to the vehicle only controls (Nussey and Whitehead, 2001; Hombach-Klonisch *et al.*, 2005;

Buckner *et al.*, 2011).

ASL Measurement of CFTR Inhibition

CFTR was inhibited with two small molecule CFTR inhibitors: Inh-172 (5 or 10μM) or GlyH-101

(5μM) (MilliporeSigma, Burlington, Vermont)(Verkman *et al.*, 2013). The cells were dosed in the

basolateral compartment of the Transwell® every other day.

Microrheology

To use PTMR for measuring viscosity, 1 µm diameter carboxylated FluoSpheres™ (ThermoFisher,

Fremont, CA), henceforth referred to as "beads", were added to the lumen of cell cultures. The thermally

driven motion of the beads was tracked to determine the viscoelastic properties of the luminal mucus.

Bead motion was recorded for 10s at a rate of 60 frames/s using a 40x air objective on a Zeiss AxioVert

A1 Fluorescent microscope (Zeiss, Jena, Germany). Individual bead trajectories were measured

automatically using a custom Python program that uses TrackPy (Markovetz *et al.*, 2019) for bead

localization and tracking. Bead motion was converted into mean squared displacements (MSD) and

complex viscosity (η*) values in accordance with the mathematics described previously (Hill *et al.*, 2014;

Markovetz *et al.*, 2019).

Statistics

One-way ANOVA was administered using SAS statistical software (Cary, NC). A general linear

model testing least squares means for effect was generated and p-values less than 0.05 were considered

significant. "N" was denoted as the number of primary cell cultures used for 96-well experiments.

Magnitude or reduction in the meniscus from baseline are expressed at means $\pm$ SEM.

Results

Air-surface liquid response to hormonal manipulation

When treated with mannitol (100mM and 300mM), compared to vehicle-only controls, cultures

demonstrated a dose-dependent increase in apical hydration at 24 and 48 hours post treatment (P<0.05)

(Figure 2). Treatment with 10 nM E2 resulted in greater ASL hydration when compared to the control

condition (p<0.001) (Figure 3A). Similarly, treatment with 100nM P4 resulted in greater reduction in ASL

than 10 nM E2 at every time point (P<0.05) (Figure 3B).

ASL response to CFTR Inhibition

10μM Inh-172 resulted in statistically significant ASL dehydration by 48, 72, and 96 hours

(p<0.01), while 5μM GlyH-101 had a smaller but still significant dehydration effect (p<0.05), compared to

the control condition (Figure 4A,B). Using mannitol in conjunction with PBS or the inhibitors amplified

the dehydration response (Figure 4B). The reduction of ASL volume by Inh-172 was visually notable

(Figure 5), and the resultant increase in mucus viscoelasticity was quantified by PTMR in n=6 cultures

and found to be significant (p<0.01)(Figure 6).

Discussion

Using two different phenotypic assays measuring mucus properties, we demonstrate that

inhibition of CFTR *in vitro* changes secreted mucus by endocervical epithelial cells. Specifically, our

results are consistent with the *clinical* findings that mucus is more hydrated in response to E2,

dehydrated when exposed to P4, and that CFTR inhibition results in abnormal, dehydrated, thickened

mucus secretions similar to that seen in cystic fibrosis (Kopito *et al.*, 1973). Combined with prior studies

that demonstrate the hormonal regulation of CFTR during the menstrual cycle, these findings further

support the integral role of CFTR in endocervical mucus changes.

Additionally, this study translates two assays, ASL and PTMR, into endocervical mucus studies

(Harvey *et al.*, 2011; Markovetz *et al.*, 2019). New to endocervical mucus studies, these assays provide

experimental endpoints that directly reflect mucus changes secreted by cultures, correlated to

biophysical properties of mucus itself. ASL, in particular, has potentially high experimental utility as it has

been both well validated in mechanistic studies of mucus regulation and used in a high-throughput

fashion for drug screening (Berg *et al.*, 2019). PTMR is a useful technique for this study for two reasons:

it operates and measures forces at the same scale as sperm in the cervical mucus environment, and it

has been shown that there is a strong correlation between *in vitro* mucus models and *in vivo* mucus in

the airway (Markovetz *et al.*, 2019). Uniquely, the reduced ciliary activity/presence in the CREC cultures

allows for PTMR to be performed *in situ*, further increasing its utility here.

Both assays provide potential endpoints for future studies of mucus *in vitro*. One challenging

characteristic of studying endocervical mucus is the lack of valid small animal models. Classic

mammalian models, such as rodents, are poor options for studying cervical mucus changes or lower

tract fertility as they lack a mucus secreting endocervix and sperm is ejaculated directly into the uterus

(Cunha *et al.*, 2019). In larger mammalian models, experimental rigor is challenging due to biological

variability, semi-quantitative nature of mucus appraisal, and invasiveness for mucus collection. An *in*

vitro model provides a tool for mechanistic studies to determine the key regulators of mucus changes

such as CFTR, but also a platform for drug discovery. With recent advances in primary cell culture such

as conditional reprogramming, there is further potential for study of genetic differences and

personalization of drugs response (Liu *et al.*, 2017).

Our previous studies demonstrate that CFTR is a hormonally regulated ion channel with E2 (mid

cycle) conditions increasing CFTR expression and, conversely, P4 (luteal) conditions decreasing CFTR

expression (Han *et al.*, 2022). In CF, the CFTR gene mutation produces a defective channel, resulting in

thick, viscous cervical mucus. Prior to the drug discovery and development of small molecules that

modulate mutant CFTR function, women with CF had known infertility/subfertility phenotypes that were

largely attributed to these mucus changes, preventing sperm penetration (Hodges *et al.*, 2008). In our

study, we used inhibitors to mimic CFTR mutation, recapitulating this phenotype *in vitro*. Thus, our study

is one of the first to experimentally confirm the relationship of CFTR inhibition and mucus changes using

loss of function experiments *in vitro*. Additionally, it provides evidence that CFTR is likely a critical

mediator of mucus changes in normal menstrual cycles.

Our *in vitro* study does not fully recapitulate either tissue level characteristics of epithelial

biology nor the full content of mucus secretions produced *in vivo*. However, the conditionally

reprogrammed platform used in this investigation provides a robust model where we can investigate

epithelial changes under multiple treatment conditions, while evaluating mucus production and quality.

We have previously demonstrated in proteomic studies, high similarity between the mucus produced in

vitro using this cellular system to *in vivo* mucus (Han *et al.*, 2020, 2022; Park *et al.*, 2022), and epithelial

only cell models have been used to high success in disease investigations and drug discovery in other

mucosal systems (Gentzsch *et al.*, 2017; Liu *et al.*, 2017).

The extent to which CFTR is responsible for mucus changes in the cervix is not fully quantified by

our studies. CFTR, while a critical channel with a well-known mutated human phenotype, is part of a

large ecosystem of ion channels. There are other ion channels, such as the epithelial sodium channel

(ENaC) that are hormonally active and may be involved in mucus hydration (Roberts *et al.*, 2023).

Investigating the relative contributions of different ion channels in mucosal hydration could provide a

more comprehensive understanding of female fertility and the development of nonhormonal drug

therapies. Additionally, future studies might develop our findings further by examining the effect of CFTR

inhibition on fertility and mucus consistency and hydration in *vivo* using a whole animal model.

Conclusion

Cervical mucus produced by the epithelia in the endocervix is a hormonally induced regulator of

female fertility. In this investigation we validate and employ two different techniques, ASL and PTMR, to

measure the viscosity and hydration of endocervical cell mucus. We determined that the inhibition of

CFTR using small molecule inhibitors results in significant dehydration and increased viscosity of cervical

mucus *in vitro*. These findings are consistent with the physiology seen clinically and predicted by

molecular studies of CFTR gene expression.

Author contribution statement

Rachel Looney: performed experiments, analyzed data, wrote paper

Mackenzie Roberts: performed experiments, analyzed data, wrote paper

Matthew Markovetz: performed experiments, analyzed data, wrote paper

Rachelle Godiah: performed experiments, analyzed data, wrote paper

Shan Yao performed experiments, analyzed data

Kirsti Golgotiu: performed experiments, analyzed data, wrote paper

Shuhao Wei: performed experiments, analyzed data

Chris Cellucci: performed experiments, analyzed data

Leo Han: conceived the study, performed experiments, analyzed data and wrote the paper

Acknowledgements

This work was supported, in whole or in part, by the Bill & Melinda Gates Foundation [INV-

024195]. Under the grant conditions of the Foundation, a Creative Commons Attribution 4.0 Generic

License has already been assigned to the Author Accepted Manuscript version that might arise from this

submission.

Funding

This work was funded by the Bill & Melinda Gates Foundation [INV-024195], Cystic Fibrosis

Foundation [MARKOV22I0], and Oregon National Primate Research Center [P51OD011092].

Declaration of interest

There is no conflict of interest to report.

Data Availability Statement

The data underlying this article will be shared on reasonable request to the corresponding

author.

References

Anthony E, Oppenheimer, Case AL, Esterly JR, Rothberg RM. Cervical mucus in cystic fibrosis: A possible

cause of infertility. *American Journal of Obstetrics and Gynecology* 1970;**108**:673–674.

Berg A, Hallowell S, Tibbetts M, Beasley C, Brown-Phillips T, Healy A, Pustilnik L, Doyonnas R, Pregel M.

High-Throughput Surface Liquid Absorption and Secretion Assays to Identify F508del CFTR

Correctors Using Patient Primary Airway Epithelial Cultures. *SLAS Discovery* 2019;**24**:724–737.

Buckner LR, Schust DJ, Ding J, Nagamatsu T, Beatty W, Chang TL, Greene SJ, Lewis ME, Ruiz B, Holman SL,

et al. Innate immune mediator profiles and their regulation in a novel polarized immortalized

epithelial cell model derived from human endocervix. *Journal of Reproductive Immunology*

2011;**92**:8–20.

Button B, Cai L-H, Ehre C, Kesimer M, Hill DB, Sheehan JK, Boucher RC, Rubinstein M. A periciliary brush

promotes the lung health by separating the mucus layer from airway epithelia. *Science*

2012;**337**:937–941.

Clunes MT, Boucher RC. Cystic fibrosis: the mechanisms of pathogenesis of an inherited lung disorder.

Drug Discovery Today: Disease Mechanisms 2007;**4**:63–72.

Cunha GR, Sinclair A, Ricke WA, Robboy SJ, Cao M, Baskin LS. Reproductive tract biology: Of mice and

men. *Differentiation* 2019;**110**:49–63.

Gentzsch M, Boyles SE, Cheluvaraju C, Chaudhry IG, Quinney NL, Cho C, Dang H, Liu X, Schlegel R, Randell

SH. Pharmacological Rescue of Conditionally Reprogrammed Cystic Fibrosis Bronchial Epithelial

Cells. *Am J Respir Cell Mol Biol* 2017;**56**:568–574.

Han L, Andrews W, Wong K, Jensen JT. Conditionally reprogrammed macaque endocervical cells retain

steroid receptor expression and produce mucus. *Biology of Reproduction* 2020;**102**:1191–1202.

Han L, Roberts M, Luo A, Wei S, Slayden OD, Macdonald KD. Functional evaluation of the cystic fibrosis

transmembrane conductance regulator in the endocervix. *Biology of Reproduction*

2022;**107**:732–740.

Haq IJ, Gray MA, Garnett JP, Ward C, Brodlie M. Airway surface liquid homeostasis in cystic fibrosis:

pathophysiology and therapeutic targets. *Thorax* 2016;**71**:284–287.

Harvey PR, Tarran R, Garoff S, Myerburg MM. Measurement of the Airway Surface Liquid Volume with

Simple Light Refraction Microscopy. *Am J Respir Cell Mol Biol* 2011;**45**:592–599.

Hill DB, Long RF, Kissner WJ, Atieh E, Garbarine IC, Markovetz MR, Fontana NC, Christy M, Habibpour M,

Tarran R, *et al.* Pathological mucus and impaired mucus clearance in cystic fibrosis patients result

from increased concentration, not altered pH. *Eur Respir J* 2018;**52**:1801297.

Hill DB, Vasquez PA, Mellnik J, McKinley SA, Vose A, Mu F, Henderson AG, Donaldson SH, Alexis NE,

Boucher RC, *et al.* A Biophysical Basis for Mucus Solids Concentration as a Candidate Biomarker

for Airways Disease. *PLOS ONE* 2014;**9**:e87681. Public Library of Science.

Hodges CA, Palmert MR, Drumm ML. Infertility in Females with Cystic Fibrosis Is Multifactorial: Evidence

from Mouse Models. *Endocrinology* 2008;**149**:2790–2797.

Hombach-Klonisch S, Kehlen A, Fowler PA, Huppertz B, Jugert JF, Bischoff G, Schlüter E, Buchmann J,

Klonisch T. Regulation of functional steroid receptors and ligand-induced responses in

telomerase-immortalized human endometrial epithelial cells. *Journal of Molecular*

Endocrinology 2005;**34**:517–534.

Jaques A, Daviskas E, Turton JA, McKay K, Cooper P, Stirling RG, Robertson CF, Bye PTP, LeSouëf PN,

Shadbolt B, *et al.* Inhaled Mannitol Improves Lung Function in Cystic Fibrosis. *Chest*

2008;**133**:1388–1396.

Keyrouz SG, Dhar R, Diringer MN. Variation in Osmotic Response to Sustained Mannitol Administration.

Neurocrit Care 2008;**9**:204–209.

Kopito LE, Kosasky HJ, Shwachman H. Water and electrolytes in cervical mucus from patients with cystic

fibrosis. *Fertil Steril* 1973;**24**:512–516.

Liu X, Krawczyk E, Suprynowicz FA, Palechor-Ceron N, Yuan H, Dakic A, Simic V, Zheng Y-L, Sripadhan P,

Chen C, *et al.* Conditional reprogramming and long-term expansion of normal and tumor cells

from human biospecimens. *Nat Protoc* 2017;**12**:439–451.

Locke LW, Myerburg MM, Markovetz MR, Parker RS, Weber L, Czachowski MR, Harding TJ, Brown SL,

Nero JA, Pilewski JM, *et al.* Quantitative imaging of airway liquid absorption in cystic fibrosis. *Eur*

Respir J 2014;**44**:675–684.

Markovetz MR, Subramani DB, Kissner WJ, Morrison CB, Garbarine IC, Ghio A, Ramsey KA, Arora H,

Kumar P, Nix DB, *et al.* Endotracheal tube mucus as a source of airway mucus for rheological

study. *Am J Physiol Lung Cell Mol Physiol* 2019;**317**:L498–L509.

Matsui H, Grubb BR, Tarran R, Randell SH, Gatzy JT, Davis CW, Boucher RC. Evidence for periciliary liquid

layer depletion, not abnormal ion composition, in the pathogenesis of cystic fibrosis airways

disease. *Cell* 1998;**95**:1005–1015.

Moghissi KS. The function of the cervix in fertility. *Fertil Steril* 1972;**23**:295–306.

Moghissi KS, Syner FN, Evans TN. A composite picture of the menstrual cycle. *Am J Obstet Gynecol*

1972;**114**:405–418.

Muchekehu RW, Quinton PM. A new role for bicarbonate secretion in cervico-uterine mucus release. *J*

Physiol 2010;**588**:2329–2342.

Myerburg MM, Butterworth MB, McKenna EE, Peters KW, Frizzell RA, Kleyman TR, Pilewski JM. Airway

surface liquid volume regulates ENaC by altering the serine protease-protease inhibitor balance:

a mechanism for sodium hyperabsorption in cystic fibrosis. *J Biol Chem* 2006;**281**:27942–27949.

Myerburg MM, Harvey PR, Heidrich EM, Pilewski JM, Butterworth MB. Acute regulation of the epithelial

sodium channel in airway epithelia by proteases and trafficking. *Am J Respir Cell Mol Biol*

2010;**43**:712–719.

Najmabadi S, Schliep KC, Simonsen SE, Porucznik CA, Egger MJ, Stanford JB. Cervical mucus patterns and

the fertile window in women without known subfertility: a pooled analysis of three cohorts.

Hum Reprod 2021;**36**:1784–1795.

Nussey S, Whitehead S. *Endocrinology: An Integrated Approach* [Internet]. 2001; BIOS Scientific

Publishers: OxfordAvailable from: http://www.ncbi.nlm.nih.gov/books/NBK22/.

Park D, Reddy AP, Wilmarth PA, Jensen JT, Han L. Mucus secretions from a conditionally reprogrammed

primary endocervical cell culture. *F S Sci* 2022;**3**:159–165.

Roberts M, Yao S, Wei S, Jensen JT, Han L. Hormonal regulation of non-cystic fibrosis transmembrane

conductance regulator ion channels in the endocervix. *F S Sci* 2023;**4**:163–171.

Saint-Criq V, Gray MA. Role of CFTR in epithelial physiology. *Cell Mol Life Sci* 2017;**74**:93–115.

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S,

Schmid B, *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods*

2012;**9**:676–682.

Tarran R, Boucher RC. Thin-film measurements of airway surface liquid volume/composition and mucus

transport rates in vitro. *Methods Mol Med* 2002;**70**:479–492.

Thornton DJ, Rousseau K, McGuckin MA. Structure and Function of the Polymeric Mucins in Airways

Mucus. *Annu Rev Physiol* 2008;**70**:459–486.

Verkman AS, Synder D, Tradtrantip L, Thiagarajah JR, Anderson MO. CFTR inhibitors. *Curr Pharm Des*

2013;**19**:3529–3541.

Figure Legends

Figure 1: (A) This figure illustrates the XZ- plane of a Transwell® during an experiment. The apical

meniscus view shows the change in meniscus corresponding to CFTR function. (B) This figure

demonstrates the image obtained from brightfield microscopy used in ASL studies. The cells can be seen

in the center of the field forming a monolayer. The darker outer ring is the meniscus or annulus formed

by mucus and PBS on the apical surface of the Transwell®.

Alt text: (A) A cross-sectional diagram showing an experimental setup of endocervical epithelial cells)

within a liquid-filled container. To the right of the container, there are two circular illustrations labeled

"Apical Meniscus View." One circle represents a hydrated, "unimpaired" meniscus, while the other

shows a dehydrated, "impaired" meniscus. (B) A grayscale circular image resembling a microscope view,

featuring concentric rings with distinct boundaries outlined in white. The distance between edge of the

well and the lighter region is the annulus.

Figure 2: Effect of mannitol on ASL hydration. Across all time points (24, 48 hours from administration)

mannitol (100 mM and 300 mM) reduced the measured change in meniscus from baseline compared to

the control (P<0.05).

Alt text: Bar chart showing reduction in meniscus from baseline for various treatment groups over time.

Figure 3: (A). Effect of E2 on ASL hydration. Across all time points (24, 48, 72, 96 hours from

administration) E2 (10 nM) reduced the measured change in meniscus from baseline compared to the

control (P<0.001). (B). Effect of E2 and P4/E2 on ASL hydration. Across all time points (24, 48, 72, 96

hours from administration) 100 nM P4 (after E2 priming) increased the measured change in meniscus

from baseline as compared E2 only (P<0.05).

Alt text: Bar charts showing reduction in meniscus from baseline for various treatment groups over time.

Figure 4: (A) Effect of InH-172 and GlyH-101 on ASL Hydration. Across time points 48, 72, and 96 hours

from administration, 10 μM Inh-172 and 5 μM GlyH-101 increased the measured change in meniscus

from baseline compared to the control (P<0.01, P<0.05 respectively). (B) Effect of Inh-172 and Mannitol

on ASL Hydration. Across time points 48, 72 and 96 hours from administration, 5 μM Inh-172 with 100

mM mannitol increased the measured change in meniscus from baseline compared to the control

(P<0.05, P<0.01, P<0.01 respectively).

Alt text: Bar charts showing reduction in meniscus from baseline for various treatment groups over time.

Chart A compares three groups, and chart B compares two groups.

Figure 5: Representative images obtained at 0 hr and 96 hr with control and Inh-172. Visually greater

reduction in meniscus in Inh-172 condition is seen.

Alt text: A collection of four images depicting a single culture well. The dark region represents the

meniscus which decreases over time in both conditions.

Figure 6: Effect of 10 μM Inh-172 on mucus complex viscosity (η*). Treatment with Inh-172 for 72 hours

increased mucus secretion η* (p<0.01).

Alt text: a graph demonstrating the difference between DMSO and Inh-172 on complex viscosity.

A

Apical Meniscus View

Hydrated, "Unimpaired"

Dehydrated, "Impaired"

Meniscus

CREC Endocervical Epithelial Cells

XZ-Plane View

B

	0Hr	96Hr
Control		
Inh-172		