Global transcription factors analyses reveal hierarchy and synergism of

regulatory networks and master virulence regulators in *Pseudomonas aeruginosa*

Jiadai Huang[1#], Yue Sun[1#], Fang Chen[1#], Shumin Li[2], Xiangkai You[1], Liangliang Han[1],

Jingwei Li[1], Zhe He[1], Canfeng Hua[1], Chunyan Yao[1], Tianmin Li[1], Beifang Lu[1], Yung-

Fu Chang[3], Xin Deng[1,4,5*]

[1] Department of Biomedical Sciences, City University of Hong Kong, 83 Tat Chee Rd,

Kowloon Tong, Hong Kong, China

[2] Department of Computer Science, The University of Hong Kong, Pokfulam, Hong

Kong, China

[3] Department of Population Medicine and Diagnostic Sciences, College of Veterinary

Medicine, Cornell University, Ithaca, NY 14853, United States.

[4] Shenzhen Research Institute, City University of Hong Kong, Shenzhen, Guangdong,

China

[5] Lead contact

[*]To whom correspondence should be addressed: Dr. Xin Deng. Tel: +852 3442 5693;

Fax: +852 3442 0549; E-mail: xindeng@cityu.edu.hk.

[#] These authors contributed equally to the paper as the first authors.

Abstract

The transcription factor (TF) regulatory network in *Pseudomonas aeruginosa* is

complex and involves multiple regulators that respond to various environmental

signals and physiological cues by regulating gene expression. However, the biological

functions of at least half of its 373 putative TFs remain uncharacterised. Herein,

chromatin immunoprecipitation sequencing (ChIP-seq) was used to investigate the

binding sites of 172 TFs in the *P. aeruginosa* PAO1 strain. The results revealed 81,009

significant binding peaks in the genome, more than half of which were located in the

promoter regions. To further decode the diverse regulatory relationships among TFs, a

hierarchical network was assembled into three levels: top, middle, and bottom.

Thirteen ternary regulatory motifs revealed flexible relationships among TFs in small

hubs, and a comprehensive co-association atlas was established, showing the

enrichment of seven core associated clusters. Twenty-four TFs were identified as the

master regulators of virulence-related pathways. The pan-genome analysis revealed

the conservation and evolution of TFs in *P. aeruginosa* complex and other species. A

Web-based database combining existing and new data from ChIP-seq and the high-

throughput systematic evolution of ligands by exponential enrichment was established

for searching TF-binding sites. This study provides important insights into the

pathogenic mechanisms of *P. aeruginosa* and related bacteria and is expected to

contribute to the development of effective therapies for infectious diseases caused by

this pathogen.

Keywords: *Pseudomonas aeruginosa*, transcription factor, ChIP-seq

Introduction

Transcription factors (TFs) are key regulators of transcription and thus play crucial roles in mediating multiple biological pathways and events in eukaryotes and prokaryotes. Based on their DNA-binding domains (DBDs), TFs can be classified into different families, such as the LysR, AraC, and LuxR families[1]. TFs can activate and repress the expression of a set of genes in response to various environmental and/or specific signal triggers[2,3]. Several approaches can be used to study TFs, including chromatin immunoprecipitation sequencing (ChIP-seq)[4] and ChIP-exo[5] in *in vivo* research and DNA affinity purification sequencing (DAP-seq)[6] and high-throughput systematic evolution of ligands by exponential enrichment (HT-SELEX)[7] in *in vitro* research. ChIP-seq is a powerful *in vivo* technique that enables more accurate genome-wide mapping of TF–DNA interactions than *in vitro* methods because TFs may interact with other co-regulators in an environment-specific pattern, thereby altering binding preferences *in vivo*[8,9].

These sequencing approaches have been used to illustrate the biological functions of TFs in several bacterial species, including *Mycobacterium tuberculosis*, *Vibrio cholerae*, *Salmonella enterica*, *Escherichia coli*, and *Clostridium thermocellum*[10-15]. In addition, several databases and models have been constructed based on experimental validation or machine learning for studying TFs in bacteria, such as CollecTF, PredicTF, PRODORIC, DBTBS, and RegulomePA[16-20]. However, the functions and spatiotemporal interactions of most bacterial TFs remain to be clarified.

The gram-negative bacterium *Pseudomonas aeruginosa* is a major human opportunistic pathogen that grows ubiquitously. *P. aeruginosa* tends to cause infection in burn victims, patients with cystic fibrosis (CF), and individuals hospitalised for long periods[21,22]. It utilises versatile pathways to exert its virulence, such as biofilm formation, quorum sensing (QS), type III and type VI secretion systems (T3SS and T6SS, respectively), motility, siderophore production, oxidative stress resistance, and antibiotic resistance, all of which are under the control of a complicated TF regulatory network (TRN). The metabolic changes regulated by the TRN enhance the pathogen's

adaptability under different infection and survival conditions.

P. aeruginosa employs diverse virulence pathways to establish successful infection,

with QS being one of the major mechanisms involving the expression of many

virulence genes. Its regulation occurs hierarchically, comprising the interconnected

las, *rhl*, and *pqs* systems[23,24]. Through its transcriptional regulator, LasR, and in

collaboration with its corresponding autoinducer, 3-oxo-C_{12}-HSL (encoded by *lasI*),

the *las* system triggers the activation of the *rhl* and *pqs* systems[23,25]. Additionally, *P.*

aeruginosa utilizes motility systems for surface attachment and colonization,

including: (i) swarming and swimming powered by flagella, which are multicellular

and individual cell movements, respectively, and (ii) twitching powered by type 4 pili

(T4P)[26], wherein T4P promote surface attachment by twitching to pull the cell closer

to the attachment sites[27]. The motility of *P. aeruginosa* is thus the initial step in its

pathogenic process and is associated with the upregulation of virulence and the

induction of host defence. Both QS and motility pathways are precisely controlled by

the sophisticated TRN in *P. aeruginosa*. Understanding how these virulence pathways

are integrated and regulated is crucial for developing novel therapeutic strategies

against *P. aeruginosa* infections.

Despite the TRN being essential for understanding how bacterial genes are regulated

synergistically, only a limited number of binding targets of TFs have been identified

in *P. aeruginosa* over the past decades. For example, ChIP-seq and RNA sequencing

(RNA-seq) co-analysis has been used to map a *P. aeruginosa* genomic regulatory

network (PAGnet) of 20 key virulence-related TFs, while HT-SELEX has been used

to characterise the DNA-binding specificities of 182 TFs[28]. DAP-seq has been used to

reveal a regulatory network of 55 response regulators (RRs) of two-component

systems in *P. aeruginosa*, 51 of which are from the PAO1 strain[29]. However, almost

half of the TFs remain functionally uncharacterized, and their downstream targets and

upstream regulators are unknown. To fill this knowledge gap, we performed ChIP-seq

of 172 TFs, representing the majority of TFs not covered by HT-SELEX, to map their

genomic binding sites *in vivo* (**Table S1**). This comprehensive dataset revealed a

hierarchical and co-association regulatory network among TFs and newly identified

virulence-related regulators. We integrated both the ChIP-seq and HT-SELEX data

into a web-based database for querying TF-binding patterns, which is publicly

available. Our findings and tools will significantly contribute to future mechanistic

studies on and a comprehensive understanding of the TRN in *P. aeruginosa* and other

bacterial species.

Results

ChIP-seq of 172 uncharacterised TFs reveals a global transcriptional atlas of *P.*

aeruginosa.

In this study, 373 proteins in *P. aeruginosa* were annotated as TFs with DNA-binding

activity based on the existing annotations in the *Pseudomonas* Genome Database[30]

(Fig. 1A). To profile the whole TF–DNA binding landscape and construct a

comprehensive regulatory network of *P. aeruginosa*, ChIP-seq experiments were

performed on 172 TFs to determine the TF-binding sites (TFBSs) **(Fig. 1A)**. Libraries

were constructed using vesicular stomatitis virus glycoprotein tagging, and raw

sequence reads were mapped to the PAO1 genome using bowtie2 (Version 2.3.4.1)[31].

Using MACS2[32], 81,009 significant binding peaks with a cut-off p-value of 0.001

were identified. Subsequently, the R package ChIPpeakAnno[33] was used to define the

peak locations and find the nearest genes. The peak location can be divided into six

features: upstream, overlapStart, inside, overlapEnd, downstream, and includeFeature.

The percentages of binding peak locations for each TF indicated that more than half

of the TFs preferentially bind to the upstream (upstream intergenic regions) and

overlapStart (overlapped with the transcription start sites [TSSs] of genes) regions

(Fig. 1B). Apart from these regions, some TFs bind to the inside regions (coding

regions of genes). Few TFs were found to preferentially bind to the rest of the peak

location features.

To further provide insights into the functional roles of different TFs in gene regulation,

we analysed the peak distance to the TSS, which revealed several patterns **(Fig. 1C)**.

First, most of the TFs exhibited peak distributions that were concentrated in the

proximal region around the TSS. Some TFs, such as Cor_CI, showed a broader

distribution of peak distance, indicating that these TFs might also be active at

positions farther from the TSS. Furthermore, certain TFs, including RpiR and DeoR,

displayed distinct bimodal or multimodal distributions, while other TFs, such as LysR

and AraC, showed smoother and more narrowly focused distributions, suggesting

their different binding preferences. The number of peaks of each TF and its DBD

types are summarised in a treemap in **Fig. 1D**. TFs in the LysR and AraC families accounted for almost half of all of the peaks. PA2718, classified in the MerR family, yielded the highest number (2,480) of binding peaks, followed by the TF PA0756 (2,080 binding peaks) from the OmpR family. Furthermore, eight TFs showed no more than 10 peaks, and the TFs PA4074 and PA4806 had only two peaks each. To preliminarily identify the different regulatory functions of TFs via their different binding preferences, we performed Gene Ontology (GO) functional enrichment analysis for each TF and presented the top 10 GO terms **(Fig. 1E)**. The significantly enriched GO terms (BH-adjusted $P < 0.05$) for all of the 172 TFs analysed by ChIP-seq are listed in **Table S2**, and these terms suggest divergent functions of the TFs. Of these functional categories, most were related to metabolism, such as transcription, translation, ribosome structure, and GTPase activity, while some of the functional categories were associated with virulence, including T4P, O antigen biosynthesis, and biofilm formation **(Fig. 1E)**.

Hierarchical networks of TFs based on pairwise interactions.

TF-coding genes can also serve as targets for other TFs, leading to interactions between the DNA-binding profiles of TFs. To clarify these relationships, lists of binding targets of the TFs obtained using ChIP-seq and HT-SELEX were used to assemble a system-level hierarchical regulatory network. After filtering by TFs that bind to promoter regions, 947 unique pairwise patterns between TFs were identified from a total of 13,375 promoter interactions.

We next defined a statistic index, h, based on the out-degree (O) and in-degree (I) of interactions to measure the direction and hierarchical level of TFs [$h = (O-I)/(O+I)$] in our established hierarchical regulatory network and thus identify hubs and information-flow bottlenecks[34]. The h value of each TF can reflect the direction and extent of information flow. A positive h indicates that the TF acts 'upstream', whereas a negative value suggests that it acts 'downstream'. The higher the absolute h value, the stronger the implication of the direction. A density plot of all of the TFs with their h values revealed that they can be approximately divided into three groups: top (h □

[0.75, 1]), middle (h □ [-0.75, 0.75]), and bottom (h □ [-1, 0.75]) levels **(Fig. S2A)**.

The top-level TFs are those that tend to control many other TFs, while the bottom-

level TFs are those that are more regulated by other TFs than acting as regulators

themselves. The middle-level TFs are those that connect the top- and bottom-level

TFs. Of the 231 TFs in this hierarchy, 100, 46, and 85 TFs were defined as top-,

middle-, and bottom-level TFs, respectively. The complete hierarchical information of

all of the TFs is provided in **Table S3**. To avoid extreme cases and aid visualisation,

we summarised the hierarchical regulatory network by removing TFs for which ($O+I$)

was more than 10 **(Fig. 2A and Fig. S2B)**. The network was visualised using

Cytoscape software[35].

Additionally, we investigated the factors influencing the hierarchical level of TFs,

such as DBD types (e.g., DBD families containing more TFs). TFs from the AraC,

GntR, and TetR families tended to cluster at the top level. TFs from the MarR family

and two-DBD-type TFs (i.e., TFs with two DBDs) were mostly found in the bottom

level. Additionally, among the characterised TFs, 58% were clustered at the bottom

level, probably because these TFs directly regulated the downstream genes involved

in important phenotypes, such as AlgR[36], AmrZ[37], MvfR[38], FleQ[39], and VqsM[40].

Overall, the organised hierarchical regulatory network profiling at the three levels

showed complex pairwise interactions among TFs in *P. aeruginosa*.

Ternary regulatory motifs show flexible relationships among TFs in *P. aeruginosa*.

Apart from their global hierarchical regulatory structure, we investigated the

constituent regulatory network motifs of TFs, which revealed small connectivity

patterns associated with canonical functions[41]. Positive and negative feedback loops

are prevalent in bacterial regulatory networks, which generally involve combinations

of several genes to appropriately respond to stimuli. One of the regulatory patterns

was a single auto-regulator that can self-regulate its expression or activity. Through

analysis, nine auto-regulator TFs were identified **(Fig. 2B)**. This number likely

represents a conservative estimate, as experiments may not optimally capture auto-

regulatory events that depend on native expression levels or specific physiological

conditions. Compared with non-auto-regulators, auto-regulators usually tend to be

repressors, which play important roles in maintaining a steady state[42,43]. For example,

the key T3SS activators PsrA and HrpL can repress their own expression in *P.*

aeruginosa and *P. savastanoi*, respectively[44-46].

In addition, the most fundamental regulatory pattern was the ternary TF motif. We

computed all possible motifs, which are listed in **Fig. 2C** and **Table S4**. The basic

triangular motifs were monodirectional regulatory structures with five different motifs.

The most frequently observed motif (6,535 occurrences) was motif_2, wherein two

different TFs co-regulate another TF. The remaining eight types of motifs, which

contained toggle switches, occurred much less frequently than the basic motifs. The

more complex the relationships among TFs, the lesser their occurrence. In particular,

motif_10, motif_12, and motif_13, characterised by two or three mutually regulating

TFs, were not found even once **(Fig. 2C)**. The high occurrence of different TFs in

diverse motifs revealed a complex and flexible regulatory network in *P. aeruginosa*.

Next, we constructed a basic motif_3 regulatory network based on potential TF–TF

interactions to detect the regulatory preferences based on DBD types **(Fig. 2D).** Basic

motif_3 was a typical hierarchical regulatory network in which TF1 regulates TF2,

which in turn regulates TF3. We found that the AraC, Fis, and GntR families tended to

harbour top-level TFs rather than middle- and bottom-level TFs, while the OmpR

family tended to harbour TFs from middle- and bottom-levels more than those from

the top level. Furthermore, the LysR and LuxR families revealed non-significant

differences among these three TF levels **(Fig. 2D)**. Taken together, our integrated

hierarchical regulatory profiling revealed a multifaceted TF–TF connection, reflecting

the regulatory preferences of TFs and DBD types in *P. aeruginos*a.

Clustering of 103 TF-binding motifs.

According to their binding sites, a total of 103 binding motifs were determined. The

TFs were clustered based on the positional weight matrix similarity of their motifs. A

circular phylogenetic tree was constructed using hierarchical clustering of the

pairwise similarity matrix **(Fig. S3A and Table S5)**. Each node in the tree represents a

TF, colour-highlighted according to its DBD family classification. The clustering

analysis revealed four distinct clusters of TFs, one each coloured light green (Cluster

1), light blue (Cluster 2), purple (Cluster 3), and pink (Cluster 4). These clusters

indicate groups of TFs with similar binding properties or regulatory functions. To

validate the identified motifs, we compared them against existing motifs in the

RegPrecise database[47], and we found the motif of PA3587 exhibited similarity to

motifs of its orthologs in other *Pseudomonadaceae* species **(Fig. S3B)**.

Within each identified cluster, TFs tend to share conserved motif patterns, indicating

potential functional relationships. For instance, it was found that the TFs in cluster 1

were enriched in adenine (A) and thymine (T) bases. Further, cluster 4 was more

likely to have an inverted repeat (IR) sequence, such as PA0367 and PA1015.

Clustering of TF binding motifs identified groups of TFs with similar intrinsic DNA-

binding specificities. As expected, many clusters contained TFs from the same DBD

families, reflecting evolutionary conservation and potential functional redundancy or

competitive binding at shared regulatory elements. Notably, the clustering also

uncovered associations between TFs from different DBD families, suggesting

convergent evolution of binding specificity or novel regulatory interactions that

warrant further investigation.

Genome-wide synergistic co-association of TFs in *P. aeruginosa*.

To further investigate the crosstalk among TFs in *P. aeruginosa*, we computed the co-

association scores based on the overlaps between the peaks of all pairs of TFs and

ChIP-seq data **(Fig. S4A and Table S6)**. Among a total of 16,175 TF pairs, we found

5,716 significant TF interactions with a co-association score of up to 0.1 based on an

elbow statistic **(Fig. S4B)**. The relationships among TFs can be further determined as

core clusters if their co-association scores are more than 0.4. The clustering analysis

of core networks using Glay clustering[48] resulted in seven associated modules, each

containing a diverse number of TFs with the DBD families highlighted in different

colours **(Fig. 3A)**.

Next, we summarised the potential functional crosstalk among the middle-scale co-

associated five TFs (PA2417, PA2718, PA0756, PmiR, and AgtR) in cluster 3. Among

a total of 1,241 target genes, 372 were co-bound by at least three TFs, and 35 were co-

bound by all five TFs, indicating that these TFs have a high level of collaboration in

regulatory patterns **(Fig. 3B)**. There were smaller intersections among different

combinations of TFs, highlighting potential co-regulation or shared binding sites.

Binding targets co-bound by more than four TFs are summarised in **Fig. 3C**. The

dense network and connectivity indicate a complex regulatory landscape, where these

TFs may orchestrate a coordinated regulation of gene expression. For example, peak

visualization of PA2504 revealed distinct binding patterns of these five TFs compared

with the control group **(Fig. 3D)**. PA2504 is the sole partner of the PppH RNA

hydrolase related to several important cellular factors[49]. In addition, four of the five

TFs (barring PA2718) can bind to the promoter region of *pqsH*, which is involved in

the *pqs* pathway **(Fig. 3E)**. Notably, PmiR has previously been verified to bind to the

pqsH promoter region and thereby regulate virulence[50]. PA2504 and *pqsH* are

highlighted in blue and green, respectively, in **Fig. 3C**.

Identified potential master regulators are associated with virulence.

After elucidating the binding targets of 172 TFs, we investigated their potential

functions in the virulence-related pathways and pathogenesis of *P. aeruginosa*. For

this purpose, we performed an enrichment analysis using a hypergeometric test to

define the master regulators (BH-adjusted $P < 0.05$) that might play important roles in

regulating specific pathways. We primarily focused on TF-binding profiles involving

the promoter regions of genes involved in nine pathways, namely QS, motility,

biofilm production, antibiotic resistance, T6SS, T3SS, reactive oxygen species (ROS)

resistance, pyocyanin production, and siderophore production. Accordingly, 134 of

the 172 TFs were found to bind to at least one target gene associated with the

abovementioned virulent pathways, and 24 were identified as master regulators of

genes involved in six pathways, namely siderophore production, biofilm production,

pyocyanin production, QS, motility, and ROS resistance pathways **(Fig. 4A and Table**

S7).

To experimentally validate the regulatory interactions identified by ChIP-seq, we

performed biochemical and genetic analyses on selected TFs. First, we conducted

Electrophoretic Mobility Shift Assays (EMSA) for four TFs, including PA0167,

PA0815, PA1380, and PA3094, using DNA fragments containing their predicted

binding sites from target gene promoters. These TFs showed specific binding to their

cognate DNA sequences (Figures S5A-D), confirming the direct binding of the ChIP-

seq-identified interactions.

To further validate the functional regulatory roles of these TFs, we constructed clean

deletion mutants of PA1380 and PA3094 (ΔPA1380 and ΔPA3094) along with their

complemented strains (ΔPA1380/p and ΔPA3094/p). RT-qPCR analysis revealed that

PA1380 positively regulates the expression of *cupB1* and *cupB3* (Figure S5E), two

genes within the CupB fimbrial cluster identified as ChIP-seq targets. Similarly,

PA3094 was confirmed to positively regulate *lecA* expression (Figure S5F), which

encodes a lectin involved in biofilm formation and host interactions[51]. Expression of

these target genes was restored to wild-type (WT) levels in the complemented strains,

validating the regulatory relationships predicted by ChIP-seq. These combined

biochemical and genetic validations demonstrate the accuracy and biological

relevance of our TF binding data.

The intersection of master regulators across these virulent pathways revealed that six

TFs (PA0167, PyrR, PA0707, PA0877, PA1504, and PA1484) only played an

important role in regulating the motility pathway, and four TFs (PA1380, PA0815,

PA5428, and PA3973) jointly regulated the motility and other pathways, including

biofilm production, pyocyanin production, and QS pathways **(Fig. 4B)**. Specifically,

PA0815 co-regulated motility, biofilm production, and QS pathways. PA1380 jointly

regulated motility and pyocyanin production pathways. PA3973 and PA5428 jointly

regulated motility and biofilm production pathways. The regulatory network of these

four master regulators and their target genes is summarised in **Fig. S5G**, represented

by lines connecting the TFs to the targets involved in specific pathways.

Predicted functions of regulators characterised in two metabolic pathways.

P. aeruginosa uses different global regulators associated with metabolism that ensure its survival and enhance its adaptability under fluctuating environmental conditions. For instance, it uses different regulatory mechanisms to respond to nutrient changes, oxygen limitation, or CF-associated lung infection conditions[52-55].

To define the core TFs that regulate metabolism, we performed master regulator analysis in metabolic pathways (*p.adjust* < 0.05) based on the TFBSs located in the promoter regions revealed by ChIP-seq for all TFs. We identified 14 TFs potentially involved in two pathways, namely the tricarboxylic acid (TCA) cycle and ribosome function pathways **(Fig. S6A-B)**. The radial plot highlights the master regulators within the TCA cycle and ribosome function pathways, and their joint regulation with their targets is presented in the regulatory network below. PA2957 and PA5438 were found to play putative roles in the TCA cycle by potentially binding to the promoter of certain genes, including *icd*, *sucA*, and *sdhC*, indicating that they coordinate the expression of genes crucial for central metabolism and energy production **(Fig. S6B)**. Furthermore, certain TFs, such as PA0611, PA0403 (PyrR), and PA1283, were found to be involved in regulating a broad array of ribosomal genes, such as *rpsB*, *rpsA*, and *rplL*. Notably, TF PyrR was found to play a putative role in regulating motility in *P. aeruginosa*. Overall, these regulators, which are related to the core metabolic pathways that enhance the adaptation of *P. aeruginosa* to different infection conditions and contribute to its pathogenicity, could offer potential drug targets for tackling *P. aeruginosa* infection in the future.

A global summary of transcription regulatory networks and functions in *P. aeruginosa*.

After studying the TFBSs of most of the TFs in *P. aeruginosa* both *in vivo* and *in vitro*, we organised these datasets to yield a comprehensive connection of virulence-related master regulators. The QS and motility pathways consisting of 18 and 20 TFs, respectively, were the key pathways in the virulent landscape and showed a close

collaboration with the other five pathways **(Fig. 5A)**. Four TFs (ExsA, CprR, AlpR,

and PA2449) were involved in T3SS, a direct host infection mechanism used by *P.*

aeruginosa. Additionally, six TFs (PA4436, PA0929, PA3771, PA2534, PA3699, and

PA5218) were involved in siderophore production, as identified using HT-SELEX,

and showed correlations with only one TF, PA2534, involved in the ROS resistance

pathway. Four TFs (CprR, PA0708, PA1520, and PA2479) were characterised in the

stringent response and persister cell pathways and were correlated with other

virulence-related pathways, namely QS, ROS resistance, T3SS, and antibiotic

resistance pathways.

To investigate the phylogenetic classification of the binding targets of the identified

TFs on a genome-wide scale, we derived a global atlas revealing the relationships

between the protein functions of target genes and the family categories of TFs **(Fig.**

5B). The Clusters of Orthologous Groups of proteins (COGs) database was used to

classify proteins based on the orthology concept[56]. Based on the annotated COGs of *P.*

aeruginosa, we found that most of the TF targets were involved in metabolism,

followed by poorly characterised categories. The *P. aeruginosa* PAO1 genome is 6.3

Mbp long, and approximately 40% of it remains uncharacterised but is believed to

encode well-conserved hypothetical proteins (HPs), which might have indispensable

and similar functions. Given that most of the TF targets are poorly characterised, it is

crucial to study these HPs in the future to provide more insights into *P. aeruginosa*

pathogenicity. Altogether, our results provide a resource of TFs and their putative

target genes in *P. aeruginosa* and, consequently, a basis for understanding the

regulatory network of virulence and orthological functions.

A Web-based database of all TFBSs in *P. aeruginosa*.

To enable a quick search of the TFBSs in *P. aeruginosa*, we developed a Web-based

database containing our ChIP-seq and HT-SELEX results at

https://jiadhuang0417.shinyapps.io/PATF_Net/, which is available to the public. It has

four parts: homepage, data page, upload, and contact. The homepage briefly describes

the background information and schematic workflow of ChIP-seq and HT-SELEX.

The data page provides two types of data: a network plot of TFs and targets, and

detailed TF–target information in a table format. The network plot illustrates the top

50 peaks for ease of visualisation. The code was generated using R, and the figures

were created using BioRender (https://www.biorender.com/).

The database offers multiple search modalities to facilitate data exploration: users can

perform TF-centric searches to query binding sites, target genes, and regulatory

networks for individual TFs, or utilize the target gene search function to identify all

TFs that regulate any gene of interest by entering its locus tag. To connect regulatory

data with biological function, we have implemented a virulence pathway browser that

allows users to explore TF binding patterns across curated gene sets for major *P.*

aeruginosa virulence pathways. Interactive visualization tools, including network

graphs and binding profile plots, facilitate intuitive exploration of regulatory

relationships. The primary purpose of PATF_Net is to store, search, and mine valuable

information on *P. aeruginosa* TFs for researchers investigating *P. aeruginosa*

infection. The current resource is based on the reference strain PAO1, which serves as

the foundation for most *P. aeruginosa* molecular studies and allows direct integration

with existing genomic annotations and functional data. However, *P. aeruginosa*

exhibits substantial genomic diversity across clinical isolates, and strain-specific

differences in TF binding patterns may contribute to phenotypic variation in virulence,

antibiotic resistance, and host adaptation. Extension of this resource to include strain-

specific regulatory maps from diverse clinical isolates would provide valuable

insights into the regulatory basis and represents an important direction for future

investigation.

Conservation and evolution of TFs in *P. aeruginosa*.

Differences between species are mainly due to differences in genes, whereas essential

genes are usually well conserved, especially in subspecies. To investigate the

conservation of TFs in *P. aeruginosa*, we performed a pan-genome analysis of 100

strains containing model strains and clinical strains using the Roary software[57]. A

total of 21,432 genes were identified across the 100 *P. aeruginosa* strains, of which

only 4,544 genes were classified as core genes (existing in more than 99% of strains),

revealing that most of the genes were cloud genes (existing in less than 15% of strains)

(Fig. 6A). However, when focusing on the TFs among these 100 strains, almost 89%

of TFs (331 of 373) were found to be well-conserved **(Fig. 6B)**, which implied that

these recognised TFs from PAO1 play essential roles in biological activities in all of

the *P. aeruginosa* subspecies.

The bar plot provides detailed information on the rest of the non-core TFs in PAO1, in

addition to several important TFs, such as LasR and ExsA, that are lacking in some

strains **(Fig. 7C)**. As is known, the *lasR* mutant of *P. aeruginosa* is associated with

CF lung disease progression and is considered a marker of an early CF-adaptive

phenotype[58]. ExsA is considered an important activator of the T3SS pathway in *P.*

aeruginosa, which is characterised by an acute infection phenotype. Additionally,

VqsM might have a relatively special function in PAO1 among *P. aeruginosa*

subspecies in addition to playing an important role in the QS and antibiotic resistance

pathways. Taken together, the core- and non-core TFs were both vital and worth

studying for understanding the metabolism and virulence-related regulatory

mechanisms in *P. aeruginosa*.

To further investigate the conservation of TFs in *P. aeruginosa*, we re-analysed the

ChIP-seq data of PhoB and RpoN in the PA14 strain[59,60] and compared them with

those in the PAO1 strain. **Fig. S7A** shows genome-wide binding peaks of PhoB and

RpoN throughout the genomes of PAO1 and PA14. We found that in PA14 and PAO1,

PhoB had 725 and 63 peaks, while RpoN yielded 363 and 1,238 peaks, respectively.

Next, we used MEME[61] to compare the motifs of PhoB and RpoN in PAO1 and PA14

based on their binding sequences from ChIP-seq **(Fig. S7B)**. PhoB and RpoN shared a

well-conserved motif in the two strains even though they showed different peak

distributions in the genomes. They had different numbers of peaks, and of the 690 and

345 respective annotation targets of PhoB and RpoN in PA14, 114 and 66 targets,

respectively, were unique to PA14. The rest of the targets were homologous between

PAO1 and PA14. We next performed target annotation and observed that the genes

regulated by these two TFs were significantly (Fisher's test) overrepresented in the

two tested strains **(Fig. S7C)**. For example, RpoN has been found to play an important

role in regulating T6SS in both PAO1 and PA14[60,62]. Overall, the conservation of the

representative TFs PhoB and RpoN in PAO1 and PA14 indicates that conserved TFs

may share similar functions in different species. This knowledge is expected to guide

drug development for different infectious strains.

TFs play a pivotal role in regulating gene expression, and studying their conservation

and evolution could provide insights into how gene expression patterns are conserved

or modified across different species and help to predict gene regulatory networks in

different species. For example, the conservation of a TF across multiple species

suggests that its targets and regulatory interactions are likely to be conserved as well.

Therefore, to study the conservation and evolution of *P. aeruginosa* TFs across other

species, we conducted phylogenetic analyses of the 373 TFs of *P. aeruginosa* across

other bacteria, fungi, archaea, plants, and animals. We found that three *P. aeruginosa*

species showed the most conserved TFs, followed by *P. savastanoi*, a phytopathogen

(Fig. 6D). Compared with bacteria, the conservation values of the TFs were lower in

fungi, archaea, plants, and animals. However, several orthologues of TFs were still

noted among these organisms. For instance, PA2032 had more than 25% and

approximately 30% identity with aminoadipate aminotransferase in humans and mice,

respectively. The phylogenetic tree of PA2032 across bacteria, archaea, fungi, plants,

and animals, with PAO1 as the root revealed that the bacterial TFs (purple) indicates a

high degree of conservation within prokaryotes, suggesting a fundamental role in core

regulatory processes **(Fig. S7D)**. In contrast, eukaryotic TFs (fungi, plants, and

animals) form distinct clades with longer branch lengths, indicating significant

divergence and specialization during eukaryotic evolution. These findings suggest that

while TF is conserved across domains of life, its functional roles and regulatory

mechanisms have undergone substantial diversification in eukaryotes. Altogether, the

high-level evolutionary conservation of TFs in *P. aeruginosa* strongly suggests that

the regulatory mechanisms are common to a wide range of bacteria.

Discussion

Bacterial pathogens use many mechanisms to establish infection and cause diseases in human hosts, making them a major public health concern worldwide. These bacteria can secrete a wide range of molecular particles that recognise and bind to host cell targets, thus damaging or preventing host responses[63]. Pathogenic bacteria use strategies to adapt to environmental stressors such as nutrient limitation or antibiotic treatment[64]. The underlying mechanism of these strategies involves sensing external signals and then responding to different environmental conditions. TFs are key molecules involved in responding to host or environmental signals by precisely modulating transcription levels; hence, their pivotal roles in infection conditions cannot be overstated. Nevertheless, our comprehension of bacterial TFs remains considerably limited, even in extensively examined model organisms such as *P. aeruginosa*. To address this knowledge gap and shed more light on TFs in *P. aeruginosa*, we conducted ChIP-seq experiments aimed at elucidating the TFBSs for 172 TFs with relatively unexplored biological functions.

ChIP-seq is a valuable technique to obtain high-resolution DNA–protein interaction mapping on a genome-wide scale to study the potential binding targets of TFs. The technique is highly sensitive and can detect low-abundance DNA-binding events, allowing the detection of weak TF–DNA interactions. To enhance the visualisation of more subtle binding peaks, particularly when investigating co-associations and TFs lacking well-defined signals, we devised a strategy involving TF-overexpressing plasmids. *In vitro* assays, such as DAP-seq or HT-SELEX, offer superior scalability to ChIP-seq but present their own challenges[29,65]. Our analysis revealed that TF binding events occur within coding regions, which is consistent with previous study demonstrating that bacterial TFs possess binding capabilities for coding regions and can regulate transcription through multiple mechanisms[66]. Besides, it may also regulate RNA stability, warranting integration with translatomics or epigenomics data. These findings extend the basic function of TFs is to recognise specific DNA sequences (i.e., motifs) at the promoter region and then upregulate or downregulate

the transcription of the target gene[67]. Notably, the literature underscores the

collaborative nature of TFs, which often exhibit interactions with other auxiliary

proteins at multiple promoter sites[68]. As shown in **Figures 3 and S3**, we present a TF-

binding motifs clustering tree and a TF co-association regulatory network based on

ChIP-seq data and core clusters of TFs with high co-association scores. Overall, our

above-described analysis has illustrated a regulatory atlas to study the biological

functions of *P. aeruginosa* TFs in specific contexts.

However, several limitations of the ChIP-seq approach should be acknowledged.

Firstly, TF overexpression ensures sufficient protein levels for ChIP-seq signal

detection but does not guarantee that all TFs are in their active conformational states,

as many bacterial TFs require allosteric activation by metabolites, cofactors, or post-

translational modifications. The cells under standard laboratory conditions which may

not activate all TFs to their maximal regulatory states, potentially leading to

underestimation of condition-specific binding peaks. Secondly, while we observed TF

binding at thousands of genomic sites, binding per se does not equate to functional

regulation, as chromatin context, cofactor availability, and competitive binding all

influence regulatory outcomes.

P. aeruginosa uses TFs to control biological functions, including metabolism and

virulence. Given the public health burden of *P. aeruginosa* infection, many studies

have investigated the roles of TFs in this pathogen. These studies have characterised

the regulatory mechanisms of individual TFs, such as FleQ, AlgR, LasR, VqsM, and

PvrA[40,69-72]. Additionally, a TRN comprising 690 genes and 1,020 regulatory

interactions with six biological modules and main motifs was established in 2011

using published data[73]. Furthermore, we developed a PAGnet containing 20 key

virulence-associated TFs based on ChIP-seq and RNA-seq data[74]. A previous study

used machine learning to identify modules of a group of genes related to known TFs

from published data[75]. Another study used DAP-seq to determine a regulatory

network of 55 RRs in *P. aeruginosa*[29]. To further identify and characterise unknown

TFs, we used HT-SELEX to describe the binding specificities of 182 TFs[28]. Compared

with the abovementioned studies, the present work profiled a comprehensive regulatory network of *P. aeruginosa*, combining existing and new data with experimental verification.

Additionally, although the TRN analysis revealed organizational patterns in *P. aeruginosa* regulatory network, the functional significance these topological features, including their specific contributions to pathogenicity, metabolic adaptation, and antibiotic resistance remains to be experimentally determined in the future work. The hierarchical structure and regulatory motifs we identified represent objective network properties derived from our binding data, but translating these structural observations into mechanistic understanding will require condition-specific functional studies, genetic validation, and phenotypic characterization. Our analysis provided a systematic framework and generating testable hypotheses rather than definitive functional conclusions. Nevertheless, these network-level organizational principles provided value to the community as a foundational reference, similar to other regulatory network maps[73] that were useful even before comprehensive validation.

We confirmed the functions and provided potential molecular mechanisms of a number of previously identified TFs to explain their phenotypes. PA0797 is known to regulate the *pqs* system and pyocyanin production[76]. In the present study, it was also found to bind to the *pqsH* promoter region and its motif was visualised. PA5428 was found to bind to the promoter regions of *aceA* and *glcB* genes[77], which was also demonstrated in our ChIP-seq results. PA4381 (CloR) was found to be associated with polymyxin resistance in a previous study[78] and to be possibly related to ROS resistance in the present study. Furthermore, PA5032 plays a putative role in biofilm regulation and also forms an operon with PA5033, an HP associated with biofilm formation[79].

While our phylogenetic analysis reveals varying degrees of TF conservation across bacterial species, the functional implications of this conservation remain to be fully explored. Many *P. aeruginosa* TFs have clear orthologs in both Gram-negative (e.g., *Klebsiella pneumoniae*) and Gram-positive pathogens (e.g., *Bacillus cereus*), yet

whether these orthologs regulate similar target genes and biological processes is largely unknown. Future comparative profiling of orthologous TFs could reveal the extent to which regulatory network architecture is conserved versus rewired during bacterial evolution, potentially identifying core regulatory modules governing universal bacterial strategies versus species-specific innovations. Such cross-species comparisons would enhance our understanding of regulatory network evolution and enable functional prediction in less well-characterized pathogens based on homology to experimentally validated *P. aeruginosa* regulators.

Our study is the first to illustrate the hierarchical regulatory network and ternary motifs of and genome-wide co-association relationships among TFs in *P. aeruginosa*, contributing to a better understanding of the fundamental traits of pathogenicity of this species. Traditional bacterial eradication strategies involving antibiotics often focus on pathogen extermination, yet this approach can inadvertently foster selective growth advantages and precipitate the emergence of drug-resistant strains. Therefore, prioritising TF-targeted drugs could potentially alleviate selective growth pressure and enhance future preventative measures against *P. aeruginosa* infection. The extensive datasets generated in this study offer valuable insights into understanding and targeting *P. aeruginosa* pathogenicity. The genome-wide binding profiles can be systematically analyzed through our hierarchical regulatory network framework to decode complex virulence mechanisms. The virulence-related master regulators and core regulatory clusters identified in this study highlighted key nodes of transcriptional control. Understanding these regulatory relationships is particularly valuable for identifying targets whose modulation would significantly impact virulence while accounting for potential compensatory mechanisms. This knowledge base thus provides a foundation for developing targeted approaches to combat *P. aeruginosa* infections, moving beyond traditional antibiotic strategies toward more sophisticated interventions based on regulatory network manipulation.

Methods

Strains, plasmids, and primers

The bacterial strains, plasmids, and primers used in the present study are listed in **Table S8**. The *P. aeruginosa* PAO1 strain, and its derivatives were grown at 37 °C in LB (Luria-Bertani) broth with shaking at 220 rpm or on LB agar plates. Antibiotics were used for *Escherichia coli* at the following concentrations: kanamycin at 50 µg/ml and carbenicillin at 60 µg/ml.

ChIP-seq and analysis

The chromatin immunoprecipitation (ChIP) procedures were modified from a previous study[80]. Briefly, for the VSVG-tagged, the ORF was amplified by PCR from the PAO1 genome and cloned into pAK1900 plasmid by HindIII/BamHI for the overexpressed TFs through HindIII site by using ClonEx- press MultiS One Step Cloning Kit (Vazyme, China). Wild-type *P. aeruginosa* containing empty pAK1900 or pAK1900-TF-VSV-G was cultured in LB medium supplemented at 37°C with shaking until mid-log phase ($OD_{600} = 0.6$), then we cross-linked the samples with 1% formaldehyde for 10 min. Subsequently, cross-linking was stopped by the addition of 125 mM glycine. Samples were centrifuged and washed thrice with a Tris buffer (20 mM Tris-HCl, pH 7.5, 150 mM NaCl). The resulting pellets were resuspended in 500 µl IP buffer (50 mM HEPES–KOH [pH 7.5], 150 mM NaCl, 1 mM EDTA, 1 % Triton X- 100, 0.1% sodium deoxycholate, 0.1% SDS, and mini-protease inhibitor cocktail (Roche), and then the DNA was broken into pieces (100 to 300 bp) with an ultrasonic processor. Insoluble cellular debris was removed by centrifugation at 4 °C, and the supernatant was used as the input sample in IP experiments. We next added 25 µl agarose-conjugated anti-VSV antibodies (Sigma) in the IP buffer. Washing, reverse cross-linked, and purification of the ChIP DNA were conducted. We used the NEXTflex ChIP-Seq kit (Bio Scientific) to construct the DNA fragment library, and agarose gel was used to cut DNA fragments between 150 and 250 bp. After sequencing, the raw reads were trimmed by Trim Galore[81] with default parameters. Then, the filtered reads were aligned to the PAO1 genome (NC_002516) using bowtie2. Only the unique reads after alignment will be used to perform peakcalling

601 with MACS2 ($P < 0.001$). MEME-ChIP was used to identify consensus motifs with

602 all peaks. TF target genes were then annotated by the R package ChIPpeakAnno.

Identification of virulence-related master regulators

We adopted the definition of "master regulator" from developmental biology[82], where

it refers to TFs that control the expression of multiple downstream genes governing a

specific biological process or lineage commitment. In our context, we use "master

regulator" to designate TFs that coordinately regulate multiple genes within specific

virulence-related pathways based on statistical enrichment criteria as described[83].

Briefly, we first generated gene lists associated with nine pathways[84], including QS,

motility, biofilm production, antibiotic resistance, T6SS, T3SS, ROS resistance,

pyocyanin, and siderophores. These gene lists are provided in full detail at our

PA_TFNet database. Then, we calculated the number of overlap genes between target

genes of TF and genes involved in the above eight pathways. The statistical

significance of the master regulator was identified using the Hypergeometric test

(BH-adjusted $P < 0.05$). R package ggplot2[85] was used to visualize the result.

GO analysis

617 GO enrichment analysis of TFs target genes was identified using R package

618 clusterProfile[86]. The enriched GO terms with BH-adjusted $P < 0.05$ were defined as

619 significantly enriched.

Pan-genome analysis

621 The information on the model and clinical *P. aeruginosa* strains[87,88] can be found in

622 **Table S9**. Briefly, the genome sequence of all strains were annotated by prokka[89] and

623 the outputs were then used as input files for roary.

Statistical analysis

625 Student's t-test was performed to analyze the RT-qPCR and lux production results

626 using Microsoft Office Excel 2019. $^{*}P < 0.05$, $^{**}P < 0.01$, and $^{***}P < 0.001$,

627 and results represent means □±□SD. All experiments were repeated at least twice.

Electrophoretic Mobility Shift Assay (EMSA)

DNA probes corresponding to predicted TF binding regions were PCR-amplified from
PAO1 genomic DNA. For binding reactions, 20 ng of purified DNA probe was
incubated with varying concentrations of recombinant TF protein in 20 μL binding
buffer containing 10 mM Tris-HCl (pH 7.4), 50 mM KCl, 5 mM MgCl□, and 10%
(v/v) glycerol. Reactions were incubated at room temperature for 20 min to allow
protein-DNA complex formation. Samples were then resolved on 6% native
polyacrylamide gels by electrophoresis at 90 V for 90 min in TBE buffer. Following
electrophoresis, gels were stained with GelRed for 5 min and visualized using a
ChemiDoc imaging system (Bio-Rad).

Reverse Transcription Quantitative PCR (RT-qPCR)

Bacterial cultures were grown in LB medium to mid-log phase (OD□ □ □ = 0.6), and
total RNA was extracted using the Bacterial Total RNA Isolation Kit (Sangon
Biotech). RNA concentration and purity were assessed using a NanoDrop 2000
spectrophotometer (Thermo Fisher Scientific). cDNA synthesis was performed using
600 ng of total RNA with HiScript III RT SuperMix (Vazyme). Quantitative PCR
reactions were carried out using ChamQ Universal SYBR qPCR Master Mix (Vazyme)
on a QuantStudio Real-Time PCR System (Applied Biosystems). Relative gene
expression levels were calculated using the $2^{(-\Delta\Delta Ct)}$ method, with expression in wild-
type strain set as 1.0. All RT-qPCR experiments were performed with three biological
replicates, each with three technical replicates.

Data availability

650 The ChIP-seq data was uploaded to National Center for Biotechnology Information
651 Gene Expression Omnibus database under accession GSE241603 and GSE271817.
652 Analysis codes have been deposited at https://github.com/dengxinb2315/PS-PATRnet-
653 code.

Acknowledgment

This study was funded by the Guangdong Major Project of Basic and Applied Basic Research (2020B0301030005), Health and Medical Research Fund (20190942), Shenzhen Science and Technology Fund (JCYJ20210324134000002), National Natural Science Foundation of China grants (32172358), the General Research Funds of Hong Kong (11103221, 11101722, and 11102223), and the Theme-based Research Scheme (T11-104/22-R). The funders had no role in study design, data collection, interpretation, or the decision to submit the work for publication.

Contributions

X.D. designed the project and obtained funding, material support and study supervision. X.D., J.H. wrote the manuscript and generated the figures and tables. J.H. performed bioinformatic analysis. J.H., Y.S., F.C., X.Y., and J.L performed the experiments. S.L. helped to write the code. Y.S. and L.H. helped to edit the manuscript. All authors reviewed the manuscript.

Declare of interest

The authors declare no competing interests.

    Figure legend

Figure 1. Overview of ChIP-seq results. (**A).** Density of all TFs (green) and ChIPed

TFs (orange) in this study throughout the *P. aeruginosa* genome. (**B).** Annotation

heatmap of all peak distribution with 6 locations: Upstream, where the peak is located

entirely upstream of the gene; Downstream, where the peak is positioned completely

downstream of the gene; Inside, where the peak is entirely contained within the gene

body; OverlapStart, where the peak overlaps with the 5' end of the gene; OverlapEnd,

where the peak overlaps with the 3' end of the gene; and IncludeFeature, where the

peak completely encompasses the gene. **(C).** Peak distance to the translational start

site (TSS) of each DBD family. **(D).** Treemap of the 172 TFs peak numbers based on

DBD family. Each box's size represents the family's size (number of peaks), and the

explained variance of each DBD type means the color shades of each box. DBD

families of ChIPed TFs are classified into 20 different categories: LuxR, LysR, Two

DBDs (Two DNA binding domains), AraC, TetR, ArsR, CRP, OmpR, GntR, MarR,

AsnC, Cro/CI, TyrR, Rrf2, MerR, IclR, Fis, RpiR, DeoR, and undetectable. **(E).** The

dot plot shows the top 10 gene ontology (GO) terms from the PseudoCAP annotation

of ChIPed TFs. The size of the dots indicates the significance of each functional

category, quantified by −log10 (p. adjust).

Figure 2. Hierarchical networks and network motifs. (**A).** Overview of hierarchy

regulatory network after removing TFs with degree O+I <= 10. Nodes indicate TFs,

and the color represents the hierarchical level. The top level was highlighted in red,

the middle in yellow, and the bottom highlighted in blue. The edges with arrows

indicate the regulatory direction. Gray means downward-pointing, and red means

upward-pointing. **(B).** The auto-regulator motif of 9 TFs. **(C).** Three transcription-

factor motifs occurrences with five basic triangular motifs and eight toggle switch

motifs. The circle presents TF, and the arrow indicates the regulatory direction**. (D).**

Alluvial diagram reveals basic triangle motif 3 (n = 1956) depending on DBD types.

The color of splines is highlighted in different DBD families, and the name of DBD

families are labeled.

Figure 3. Core co-association regulatory networks. (A). Core clusters of significant

co-binding patterns of TFs. Each TF is highlighted in different colors based on DBD

types. The co-association score by pair of TFs was calculated by Jaccard statistics,

which measures the ratio of the number of base pairs in overlapped binding peaks on

both TFs to the number of base pairs in their union. **(B).** The histogram's overlapped

target genes of TFs in cluster 3 represent the number of target peaks in the

individual/overlapped set. **(C).** The network of co-regulation of TFs in cluster 3 with

co-bound targets of more than 4. **(D-E).** Genome browser view of TFs in cluster 3

binding intensities at the PA2504 and *pqsH* locus.

Figure 4. Newly identified virulence-related master regulators. (A). Overview of

all identified master regulators related to 6 pathways, including QS, motility, biofilm,

siderophore, pyocyanin, and ROS. Each circle represents one TF, with the height

indicating the significance, quantified by -log10 (p.adjust), and the size indicates the

number of targets associated with the virulence pathway. **(B).** Intersection of master

regulators in 6 virulence pathways. The bar chart on the top shows the number of

intersections, while the matrix below indicates which TFs are involved in specific

biological processes such as siderophore production, pyocyanin production, biofilm

formation, ROS response, QS, and motility.

Figure 5. Overview of transcriptional network of TFs in *P. aeruginosa*. (A). The

interaction network of virulence-related master regulators in *P. aeruginosa*. The ten

virulence pathways are highlighted in different colors. The size of the circle represents

the degree, and the width of the edges indicates the overlapped number of TFs

between two pathways. **(B).** The target annotated using the COG database shows four

orthology classification profiling from different DBD types TFs. The size of rectangle

indicates the number of targets.

Figure 6. Conservation and variability of TFs in PAO1. (A). The pie chart shows

the proportions of genes categorized by their presence across *P. aeruginosa* strains for

all genes. **(B).** The pie chart shows the distribution of TFs identified from PAO1

across different conservation categories. **(C).** The bar plot of the proportion for non-

core TFs. Genes are categorized based on their presence frequency across P.

aeruginosa strains: Core genes (present in 99% ~ 100% strains), Soft core genes

(present in 95% ~ 99% strains), Shell genes (present in 15% ~ 95% strains), and

Cloud genes (present in 0% ~ 15% strains). **(D).** The conservation and evolutionary

relationship of all 373 TFs in PAO1 among bacteria, archaebacteria, fungi, plants, and

animals. The conservation value was normalized after blastp alignment[90]. The

phylogenetic trees were constructed using MEGA11[91] and plotted via R package

ggtree[92].

Supplementary information

Fig. S1. RNA-seq for all TFs and ChIPed TFs. Previous RNA-seq data shows the

expression level of all TFs and ChIPed TFs. The circle size indicates the log2-

transformation raw counts, and the color highlighted for ChIPed TFs represented

different DBD families.

Fig. S2. Distribution of Hierarchy Height h. $h = (O-I)/(O+I)$, O presents the extent

they regulate other TFs, and I indicates the level other TFs regulate them. **(A).** TFs

can be divided into almost equal three levels depending on index h: Bottom level (-1.0

< h < 0.75), Middle level (-0.75 < h < 0.75), and Top level (0.75 < h < 1.0). **(B).** The

number of TFs resided in three levels after removing TFs with $O+I < 10$.

Fig. S3 Circular phylogenetic tree of TF binding motifs. (A). Circular phylogenetic

tree representing the clustering of TFs based on their motif similarity. The tree was

constructed using hierarchical clustering with the Ward.D2 method applied to the

pairwise similarity matrix of motifs. The tree were visualized using ggtree[93]. Each

node represents a TF, color-coded by its DBD family classification as indicated in the

legend on the right. The motif logos are aligned with the respective TFs, with their

orientations adjusted to maintain legibility around the circular layout. **(B).** PA3587

and MetR display similar DNA binding motifs.

Fig. S4. Co-association network of TFs. (A). Heatmap reveals a full co-association
pattern of all TFs. **(B).** The density of co-association score for all TF pairs.
Determining co-association score as 0.1 (dashed in red) of significant TF co-
associations based on an elbow statistic.

Fig. S5. Validation and co-regulation of virulence-related master regulators. (A).
The validation of the binding sites of PA0167 by EMSA. **(B).** The validation of the
binding sites of PA0815 by EMSA. **(C).** The validation of the binding sites of PA1380
by EMSA. **(D).** The validation of the binding sites of PA3094 by EMSA. **(E).** The
detection of expression of target genes PA1380, *cupB1*, and *cupB3* in WT, ΔPA1380
and complementary strain by RT-qPCR. **(F).** The detection of expression of target
genes PA3094 and *lecA* in WT, ΔPA3094 and complementary strain by RT-qPCR. **(G).**
Graph diagram of interactions involving target genes of 4 TFs, including PA1380,
PA0815, PA5428, and PA3973. NC indicates negative control. Student's *t* test, $*P \leq$
0.05.

Fig. S6. Regulators involved in TCA and Ribosome pathways. (A). Radar plots
show the putative regulator in TCA and ribosome pathways. **(B).** Graph diagram of
interactions involving target genes of 14 TFs, including PA0611, PA5218, PA5431,
PA1759, PA0403 (PyrR), PA 1283, PA4381, PA5255 (AlgQ), PA4784, PA0475,
PA0893, PA0448, PA2957 and PA5438.

Fig. S7. Conserved TFs in *P. aeruginosa*. (A). The coverage of PhoB and RpoN
peak regions over PAO1 and PA14 chromosomes. Each line shows the location and
log2 Fold Enrichment of peaks signal in the chromosome. **(B).** The binding motifs of
PhoB and RpoN were analyzed via MEME-ChIP. All peaks were used to define the
binding motif. **(C).** Comparison of overlapped targets enriched in PAO1 and PA14 of
PhoB and RpoN. The Fisher test made the significance of the overlap ratio. **(D).**
Phylogenetic tree of PA2032 across different domains of life. The phylogenetic tree
was constructed using PA2032 sequence from PAO1 as the root.

Table S1. List of 172 ChIPed TFs.

Table S2. Enriched GO terms of 172 ChIPed TFs.

Table S3. Hierarchical regulatory network.

Table S4. Ternary regulatory motifs.

Table S5. PWM pairwise similarity scores.

Table S6. Co-association network.

Table S7. Virulence-related master regulators.

Table S8. Strains and primers used in this study.

Table S9. Reference list of strains for pan-genome analysis.

Reference

1   El-Gebali, S., Mistry, J., Bateman, A., Eddy, S. R., Luciani, A., Potter, S. C. *et al.* The
Pfam protein families database in 2019. *Nucleic Acids Res* **47**, D427-D432 (2019).
2   Salgado, H., Martinez-Antonio, A. & Janga, S. C. Conservation of transcriptional
sensing systems in prokaryotes: a perspective from *Escherichia coli*. *FEBS Lett* **581**,
3499-3506 (2007).
3   Seshasayee, A. S., Sivaraman, K. & Luscombe, N. M. An overview of prokaryotic
transcription factors : a summary of function and occurrence in bacterial genomes.
Subcell Biochem **52**, 7-23 (2011).
4   Johnson, D. S., Mortazavi, A., Myers, R. M. & Wold, B. Genome-wide mapping of in
vivo protein-DNA interactions. *Science* **316**, 1497-1502 (2007).
5   Rhee, H. S. & Pugh, B. F. Comprehensive Genome-wide Protein-DNA Interactions
Detected at Single-Nucleotide Resolution. *Cell* **147**, 1408-1419 (2011).
6   Bartlett, A., O'Malley, R. C., Huang, S. S. C., Galli, M., Nery, J. R., Gallavotti, A. *et*
al. Mapping genome-wide transcription-factor binding sites using DAP-seq. *Nat*
Protoc **12**, 1659-1672 (2017).
7   Jolma, A., Yan, J., Whitington, T., Toivonen, J., Nitta, K. R., Rastas, P. *et al.* DNA-
Binding Specificities of Human Transcription Factors. *Cell* **152**, 327-339 (2013).
8   Furey, T. S. ChIP-seq and beyond: new and improved methodologies to detect and
characterize protein-DNA interactions. *Nat Rev Genet* **13**, 840-852 (2012).
9   Fu, L. Y., Zhu, T., Zhou, X. K., Yu, R. R., He, Z. H., Zhang, P. J. *et al.* ChIP-Hub
provides an integrative platform for exploring plant regulome. *Nat Commun* **13**
(2022).
10   Minch, K. J., Rustad, T. R., Peterson, E. J., Winkler, J., Reiss, D. J., Ma, S. *et al.* The
DNA-binding network of *Mycobacterium tuberculosis*. *Nat Commun* **6**, 5829 (2015).
11   Ayala, J. C., Silva, A. J. & Benitez, J. A. H-NS: an overarching regulator of the *Vibrio*
cholerae life cycle. *Res Microbiol* **168**, 16-25 (2017).

12   Garcia-Pastor, L., Sanchez-Romero, M. A., Jakomin, M., Puerta-Fernandez, E. &
819   Casadesus, J. Regulation of bistability in the std fimbrial operon of *Salmonella*
820   *enterica* by DNA adenine methylation and transcription factors HdfR, StdE and StdF.
821   *Nucleic Acids Res* **47**, 7929-7941 (2019).

822   13   Seo, S. W., Kim, D., Latif, H., O'Brien, E. J., Szubin, R. & Palsson, B. O.
823   Deciphering Fur transcriptional regulatory network highlights its complex role
824   beyond iron metabolism in *Escherichia coli*. *Nat Commun* **5**, 4910 (2014).

825   14   Ishihama, A., Shimada, T. & Yamazaki, Y. Transcription profile of *Escherichia coli*:
826   genomic SELEX search for regulatory targets of transcription factors. *Nucleic Acids*
827   *Res* **44**, 2058-2074 (2016).

828   15   Hebdon, S. D., Gerritsen, A. T., Chen, Y. P., Marcano, J. G. & Chou, K. J. Genome-
829   Wide Transcription Factor DNA Binding Sites and Gene Regulatory Networks in
830   *Clostridium thermocellum*. *Front Microbiol* **12**, 695517 (2021).

831   16   Kilic, S., White, E. R., Sagitova, D. M., Cornish, J. P. & Erill, I. CollecTF: a database
832   of experimentally validated transcription factor-binding sites in Bacteria. *Nucleic*
833   *Acids Res* **42**, D156-160 (2014).

834   17   Oliveira Monteiro, L. M., Saraiva, J. P., Brizola Toscan, R., Stadler, P. F., Silva-Rocha,
835   R. & Nunes da Rocha, U. PredicTF: prediction of bacterial transcription factors in
836   complex microbial communities using deep learning. *Environ Microbiome* **17**, 7
837   (2022).

838   18   Dudek, C. A. & Jahn, D. PRODORIC: state-of-the-art database of prokaryotic gene
839   regulation. *Nucleic Acids Res* **50**, D295-D302 (2022).

840   19   Makita, Y., Nakao, M., Ogasawara, N. & Nakai, K. DBTBS: database of
841   transcriptional regulation in *Bacillus subtilis* and its contribution to comparative
842   genomics. *Nucleic Acids Res* **32**, D75-77 (2004).

843   20   Galan-Vasquez, E., Luna-Olivera, B. C., Ramirez-Ibanez, M. & Martinez-Antonio, A.
844   RegulomePA: a database of transcriptional regulatory interactions in *Pseudomonas*
845   *aeruginosa* PAO1. *Database-Oxford* (2020).

846   21   Bodey, G. P., Bolivar, R., Fainstein, V. & Jadeja, L. Infections caused by
847   *Pseudomonas aeruginosa*. *Rev Infect Dis* **5**, 279-313 (1983).

848   22   Stover, C. K., Pham, X. Q., Erwin, A. L., Mizoguchi, S. D., Warrener, P., Hickey, M. J.
849   *et al.* Complete genome sequence of *Pseudomonas aeruginosa* PAO1, an
850   opportunistic pathogen. *Nature* **406**, 959-964 (2000).

851   23   Lee, J. & Zhang, L. H. The hierarchy quorum sensing network in *Pseudomonas*
852   *aeruginosa*. *Protein Cell* **6**, 26-41 (2015).

853   24   Girard, G. & Bloemberg, G. V. Central role of quorum sensing in regulating the
854   production of pathogenicity factors in *Pseudomonas aeruginosa*. *Future Microbiol* **3**,
855   97-106 (2008).

856   25   Schuster, M. & Greenberg, E. P. Early activation of quorum sensing in *Pseudomonas*
857   *aeruginosa* reveals the architecture of a complex regulon. *Bmc Genomics* **8** (2007).

858   26   Kazmierczak, B. I., Schniederberend, M. & Jain, R. Cross-regulation of
859   *Pseudomonas* motility systems: the intimate relationship between flagella, pili and
860   virulence. *Curr Opin Microbiol* **28**, 78-82 (2015).

27   Kearns, D. B. A field guide to bacterial swarming motility. *Nat Rev Microbiol* **8**, 634-

644 (2010).

28   Wang, T. T., Sun, W. J., Fan, L. G., Hua, C. F., Wu, N., Fan, S. R. *et al.* An atlas of the binding specificities of transcription factors in *Pseudomonas aeruginosa* directs prediction of novel regulators in virulence. *Elife* **10** (2021).

29   Trouillon, J., Imbert, L., Villard, A. M., Vernet, T., Attree, I. & Elsen, S. Determination of the two-component systems regulatory network reveals core and accessory regulations across *Pseudomonas aeruginosa* lineages. *Nucleic Acids Research* **49**, 11476-11490 (2021).

30   Winsor, G. L., Griffiths, E. J., Lo, R., Dhillon, B. K., Shay, J. A. & Brinkman, F. S. Enhanced annotations and features for comparing thousands of *Pseudomonas* genomes in the *Pseudomonas* genome database. *Nucleic Acids Res* **44**, D646-653 (2016).

31   Langmead, B., Trapnell, C., Pop, M. & Salzberg, S. L. Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol* **10**, R25 (2009).

32   Zhang, Y., Liu, T., Meyer, C. A., Eeckhoute, J., Johnson, D. S., Bernstein, B. E. *et al.* Model-based analysis of ChIP-Seq (MACS). *Genome Biol* **9**, R137 (2008).

33   Zhu, L. J., Gazin, C., Lawson, N. D., Pages, H., Lin, S. M., Lapointe, D. S. *et al.* ChIPpeakAnno: a Bioconductor package to annotate ChIP-seq and ChIP-chip data. *BMC Bioinformatics* **11**, 237 (2010).

34   Gerstein, M. B., Kundaje, A., Hariharan, M., Landt, S. G., Yan, K. K., Cheng, C. *et al.* Architecture of the human regulatory network derived from ENCODE data. *Nature* **489**, 91-100 (2012).

35   Shannon, P., Markiel, A., Ozier, O., Baliga, N. S., Wang, J. T., Ramage, D. *et al.* Cytoscape: a software environment for integrated models of biomolecular interaction networks. *Genome research* **13**, 2498-2504 (2003).

36   Lizewski, S. E., Lundberg, D. S. & Schurr, M. J. The transcriptional regulator AlgR is essential for *Pseudomonas aeruginosa* pathogenesis. *Infect Immun* **70**, 6083-6093 (2002).

37   Jones, C. J., Ryder, C. R., Mann, E. E. & Wozniak, D. J. AmrZ modulates *Pseudomonas aeruginosa* biofilm architecture by directly repressing transcription of the psl operon. *J Bacteriol* **195**, 1637-1644 (2013).

38   Déziel, E., Gopalan, S., Tampakaki, A. P., Lépine, F., Padfield, K. E., Saucier, M. *et al.* The contribution of MvfR to *Pseudomonas aeruginosa* pathogenesis and quorum sensing circuitry regulation: multiple quorum sensing□regulated genes are modulated without affecting *lasRI*, *rhlRI* or the production of N□acyl□L□homoserine lactones. *Mol Microbiol* **55**, 998-1014 (2005).

39   Hickman, J. W. & Harwood, C. S. Identification of FleQ from *Pseudomonas aeruginosa* as ac□di□GMP□responsive transcription factor. *Mol Microbiol* **69**, 376-389 (2008).

40   Liang, H., Deng, X., Li, X., Ye, Y. & Wu, M. Molecular mechanisms of master regulator VqsM mediating quorum-sensing and antibiotic resistance in *Pseudomonas aeruginosa*. *Nucleic acids research* **42**, 10307-10320 (2014).

41   Cheng, C., Yan, K. K., Hwang, W., Qian, J., Bhardwaj, N., Rozowsky, J. *et al.*

Construction and analysis of an integrated regulatory network derived from high-
throughput sequencing data. *PLoS Comput Biol* **7**, e1002190 (2011).

42   Alon, U. Network motifs: theory and experimental approaches. *Nat Rev Genet* **8**, 450-
461 (2007).

43   Burda, Z., Krzywicki, A., Martin, O. C. & Zagorski, M. Motifs emerge from function
in model gene regulatory networks. *Proceedings of the National Academy of Sciences*
108, 17263-17268 (2011).

44   Huang, J., Yao, C., Sun, Y., Ji, Q. & Deng, X. Virulence-related regulatory network of
Pseudomonas syringae. *Comput Struct Biotechnol J* **20**, 6259-6270 (2022).

45   Shen, D. K., Filopon, D., Kuhn, L., Polack, B. & Toussaint, B. PsrA is a positive
transcriptional regulator of the type III secretion system in *Pseudomonas aeruginosa*.
Infect Immun **74**, 1121-1129 (2006).

46   Kojic, M., Jovcic, B., Vindigni, A., Odreman, F. & Venturi, V. Novel target genes of
PsrA transcriptional regulator of *Pseudomonas aeruginosa*. *Fems Microbiol Lett* **246**,
175-181 (2005).

47   Novichkov, P. S., Kazakov, A. E., Ravcheev, D. A., Leyn, S. A., Kovaleva, G. Y.,
Sutormin, R. A. *et al.* RegPrecise 3.0–a resource for genome-scale exploration of
transcriptional regulation in bacteria. *Bmc Genomics* **14**, 745 (2013).

48   Su, G., Kuchinsky, A., Morris, J. H., States, D. J. & Meng, F. GLay: community
structure analysis of biological networks. *Bioinformatics* **26**, 3135-3137 (2010).

49   Drabinska, J., Steczkiewicz, K., Kujawa, M. & Kraszewska, E. Searching for
biological function of the mysterious PA2504 protein from *Pseudomonas aeruginosa*.
International Journal of Molecular Sciences **22**, 9833 (2021).

50   Cui, G., Zhang, Y., Xu, X., Liu, Y., Li, Z., Wu, M. *et al.* PmiR senses 2-
methylisocitrate levels to regulate bacterial virulence in *Pseudomonas aeruginosa*.
Sci Adv **8**, eadd4220 (2022).

51   Chemani, C., Imberty, A., de Bentzmann, S., Pierre, M., Wimmerová, M., Guery, B. P.
et al. Role of LecA and LecB lectins in *Pseudomonas aeruginosa*-induced lung injury
and effect of carbohydrate ligands. *Infect Immun* **77**, 2065-2075 (2009).

52   Moreno, R. & Rojo, F. The importance of understanding the regulation of bacterial
metabolism. *Environ Microbiol* **25**, 54-58 (2023).

53   Dolan, S. K., Kohlstedt, M., Trigg, S., Ramirez, P. V., Wittmann, C., Kaminski, C. *et*
al. Contextual Flexibility in *Pseudomonas aeruginosa* Central Carbon Metabolism
during Growth in Single Carbon Sources. *Mbio* **11** (2020).

54   Oberhardt, M. A., Goldberg, J. B., Hogardt, M. & Papin, J. A. Metabolic Network
Analysis of *Pseudomonas aeruginosa* during Chronic Cystic Fibrosis Lung Infection.
J Bacteriol **192**, 5534-5548 (2010).

55   Korgaonkar, A. K. & Whiteley, M. *Pseudomonas aeruginosa* Enhances Production of
an Antimicrobial in Response to N-Acetylglucosamine and Peptidoglycan. *J*
Bacteriol **193**, 909-917 (2011).

56   Tatusov, R. L., Galperin, M. Y., Natale, D. A. & Koonin, E. V. The COG database: a
tool for genome-scale analysis of protein functions and evolution. *Nucleic Acids Res*
28, 33-36 (2000).

57   Page, A. J., Cummins, C. A., Hunt, M., Wong, V. K., Reuter, S., Holden, M. T. *et al.*

Roary: rapid large-scale prokaryote pan genome analysis. *Bioinformatics* **31**, 3691-
3693 (2015).

58   Hoffman, L. R., Kulasekara, H. D., Emerson, J., Houston, L. S., Burns, J. L., Ramsey,
B. W. *et al. Pseudomonas aeruginosa lasR* mutants are associated with cystic fibrosis
lung disease progression. *Journal of Cystic Fibrosis* **8**, 66-70 (2009).

59   Bielecki, P., Jensen, V., Schulze, W., Gödeke, J., Strehmel, J., Eckweiler, D. *et al.*
Cross talk between the response regulators PhoB and TctD allows for the integration
of diverse environmental signals in *Pseudomonas aeruginosa*. *Nucleic Acids*
Research **43**, 6413-6425 (2015).

60   Allsopp, L. P., Collins, A. C. Z., Hawkins, E., Wood, T. E. & Filloux, A. RpoN/Sfa2-
dependent activation of the *Pseudomonas aeruginosa* H2-T6SS and its cognate
arsenal of antibacterial toxins. *Nucleic Acids Res* **50**, 227-243 (2022).

61   Bailey, T. L., Boden, M., Buske, F. A., Frith, M., Grant, C. E., Clementi, L. *et al.*
MEME SUITE: tools for motif discovery and searching. *Nucleic Acids Res* **37**,
W202-208 (2009).

62   Shao, X., Zhang, X., Zhang, Y., Zhu, M., Yang, P., Yuan, J. *et al.* RpoN-Dependent
Direct Regulation of Quorum Sensing and the Type VI Secretion System in
Pseudomonas aeruginosa PAO1. *J Bacteriol* **200** (2018).

63   Wilson, J. W., Schurr, M. J., LeBlanc, C. L., Ramamurthy, R., Buchanan, K. L. &
Nickerson, C. A. Mechanisms of bacterial pathogenicity. *Postgrad Med J* **78**, 216-224
(2002).

64   Blair, J. M., Webber, M. A., Baylay, A. J., Ogbolu, D. O. & Piddock, L. J. Molecular
mechanisms of antibiotic resistance. *Nat Rev Microbiol* **13**, 42-51 (2015).

65   Baumgart, L. A., Lee, J. E., Salamov, A., Dilworth, D. J., Na, H., Mingay, M. *et al.*
Persistence and plasticity in bacterial gene regulation. *Nat Methods* **18**, 1499-1505
(2021).

66   Hua, C., Huang, J., Wang, T., Sun, Y., Liu, J., Huang, L. *et al.* Bacterial Transcription
Factors Bind to Coding Regions and Regulate Internal Cryptic Promoters. *Mbio* **13**,
e0164322 (2022).

67   Perez-Rueda, E., Hernandez-Guerrero, R., Martinez-Nunez, M. A., Armenta-Medina,
D., Sanchez, I. & Ibarra, J. A. Abundance, diversity and domain architecture
variability in prokaryotic DNA-binding transcription factors. *PLoS One* **13**, e0195332
(2018).

68   Browning, D. F., Butala, M. & Busby, S. J. W. Bacterial Transcription Factors:
Regulation by Pick "N" Mix. *J Mol Biol* **431**, 4067-4077 (2019).

69   Arora, S. K., Ritchings, B. W., Almira, E. C., Lory, S. & Ramphal, R. A
transcriptional activator, FleQ, regulates mucin adhesion and flagellar gene
expression in *Pseudomonas aeruginosa* in a cascade manner. *J Bacteriol* **179**, 5574-
5581 (1997).

70   Kong, W., Zhao, J., Kang, H., Zhu, M., Zhou, T., Deng, X. *et al.* ChIP-seq reveals the
global regulator AlgR mediating cyclic di-GMP synthesis in *Pseudomonas*
aeruginosa. *Nucleic acids research* **43**, 8268-8282 (2015).

71   Gilbert, K. B., Kim, T. H., Gupta, R., Greenberg, E. P. & Schuster, M. Global position
analysis of the *Pseudomonas aeruginosa* quorum☐sensing transcription factor LasR.

Mol Microbiol **73**, 1072-1085 (2009).

72 Pan, X., Fan, Z., Chen, L., Liu, C., Bai, F., Wei, Y. *et al.* PvrA is a novel regulator that
contributes to *Pseudomonas aeruginosa* pathogenesis by controlling bacterial
utilization of long chain fatty acids. *Nucleic Acids Research* **48**, 5967-5985 (2020).

73 Galan-Vasquez, E., Luna, B. & Martinez-Antonio, A. The Regulatory Network of
Pseudomonas aeruginosa. Microb Inform Exp **1**, 3 (2011).

74 Huang, H., Shao, X. L., Xie, Y. P., Wang, T. T., Zhang, Y. C., Wang, X. *et al.* An
integrated genomic regulatory network of virulence-related transcriptional factors in
Pseudomonas aeruginosa. Nat Commun **10** (2019).

75 Rajput, A., Tsunemoto, H., Sastry, A. V., Szubin, R., Rychel, K., Sugie, J. *et al.*
Machine learning from *Pseudomonas aeruginosa* transcriptomes identifies
independently modulated sets of genes associated with known transcriptional
regulators. *Nucleic Acids Research* **50**, 3658-3672 (2022).

76 Cui, G. Y., Zhang, Y. X., Xu, X. J., Liu, Y. Y., Li, Z., Wu, M. *et al.* PmiR senses 2-
methylisocitrate levels to regulate bacterial virulence in *Pseudomonas aeruginosa.*
Sci Adv **8** (2022).

77 Hwang, W., Yong, J. H., Min, K. B., Lee, K.-M., Pascoe, B., Sheppard, S. K. *et al.*
Genome-wide association study of signature genetic alterations among *pseudomonas*
aeruginosa cystic fibrosis isolates. *Plos Pathog* **17**, e1009681 (2021).

78 Gutu, A. D., Sgambati, N., Strasbourger, P., Brannon, M. K., Jacobs, M. A., Haugen,
E. *et al.* Polymyxin resistance of *Pseudomonas aeruginosa phoQ* mutants is
dependent on additional two-component regulatory systems. *Antimicrob Agents*
Chemother **57**, 2204-2215 (2013).

79 Zhang, L., Fritsch, M., Hammond, L., Landreville, R., Slatculescu, C., Colavita, A. *et*
al. Identification of genes involved in Pseudomonas aeruginosa biofilm-specific
resistance to antibiotics. *PLoS One* **8**, e61625 (2013).

80 Blasco, B., Chen, J. M., Hartkoorn, R., Sala, C., Uplekar, S., Rougemont, J. *et al.*
Virulence Regulator EspR of *Mycobacterium tuberculosis* Is a Nucleoid-Associated
Protein. *Plos Pathog* **8** (2012).

81 Felix Krueger, F. J., Phil Ewels, Ebrahim Afyounian, Michael Weinstein, Benjamin
Schuster-Boeckler, Gert Hulselmans, & sclamons. FelixKrueger/TrimGalore: v0.6.10
- add default decompression path (0.6.10). (2023).

82 Chan, S. S.-K. & Kyba, M. What is a master regulator? *Journal of stem cell research*
& therapy **3**, 114 (2013).

83 Fan, L. G., Wang, T. T., Hua, C. F., Sun, W. J., Li, X. Y., Grunwald, L. *et al.* A
compendium of DNA-binding specificities of transcription factors in *Pseudomonas*
syringae. Nat Commun **11** (2020).

84 Shao, X., Yao, C., Ding, Y., Hu, H., Qian, G., He, M. *et al.* The transcriptional
regulators of virulence for *Pseudomonas aeruginosa*: Therapeutic opportunity and
preventive potential of its clinical infections. *Genes Dis* **10**, 2049-2063 (2023).

85 Wickham, H. & Wickham, H. Data transformation. In: *ggplot2*: Elegant graphics for
data analysis. 203-220 (2016).

86 Yu, G., Wang, L.-G., Han, Y. & He, Q.-Y. clusterProfiler: an R package for comparing
biological themes among gene clusters. *Omics: a journal of integrative biology* **16**,

1038   284-287 (2012).

1039 87 Hu, Y. Y., Wang, Q., Sun, Q. L., Chen, G. X. & Zhang, R. A novel plasmid carrying
1040   carbapenem-resistant gene blaKPC-2 in *Pseudomonas aeruginosa*. *Infect Drug Resist*
1041   **12**, 1285-1288 (2019).

1042 88 Hu, Y., Liu, C., Wang, Q., Zeng, Y., Sun, Q., Shu, L. *et al.* Emergence and expansion
1043   of a carbapenem-resistant *Pseudomonas aeruginosa* clone are associated with
1044   plasmid-borne bla KPC-2 and virulence-related genes. *Msystems* **6**,
1045   10.1128/msystems. 00154-00121 (2021).

1046 89 Seemann, T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068-
1047   2069 (2014).

1048 90 Coordinators, N. R. Database resources of the National Center for Biotechnology
1049   Information. *Nucleic Acids Res* **41**, D8-D20 (2013).

1050 91 Tamura, K., Stecher, G. & Kumar, S. MEGA11: Molecular Evolutionary Genetics
1051   Analysis Version 11. *Mol Biol Evol* **38**, 3022-3027 (2021).

1052 92 Yu, G., Smith, D. K., Zhu, H., Guan, Y. & Lam, T. T. Y. ggtree: an R package for
1053   visualization and annotation of phylogenetic trees with their covariates and other
1054   associated data. *Methods in Ecology and Evolution* **8**, 28-36 (2017).

1055 93 Yu, G. Using ggtree to Visualize Data on Tree-Like Structures. *Curr Protoc*
1056   *Bioinformatics* **69**, e96 (2020).

1057

A

B

Auto-Regulators

4074	4165	4806
5360	1196	5483
1315	2258	3599

Top ●

Middle ●

Bottom ●

Downward →

Upward →

C

Basic Triangular Motifs

M1 n=5530
M2 n=6535
M3 n=1956

M4 n=4
M5 n=220

Toggle Switch Motifs

M6 n=41
M10 n=0
M7 n=91
M11 n=10
M8 n=10
M12 n=0
M9 n=2
M13 n=0

D

Basic Motif_3
(n =1956)

AraC
Arc~like
Fis
GntR
LuxR
LysR
OmpR
Sigma factor 54
TetR

AraC
Arc~like
Cro/CI
DeoR
Fis
LcrR
LuxR
LysR
LytTr
MarR
OmpR
TetR
Two DBDs

AraC
AsnC
Cro/CI
Fis
GntR
LuxR
LysR
OmpR
TetR
Two DBDs

A

B

Figure S1

Figure S2

Figure S3

Figure S4

Figure S5

A

B

Figure S7