Metabolic modeling of microbial communities in the chicken ceca reveals a landscape of competition and co-operation

Irina Utkina[1,2], Yi Fan[,3], Benjamin P. Willing[,3], and John Parkinson[1,2,4,*]

[1] Department of Molecular Genetics, University of Toronto, Toronto ON, Canada

[2] Program in Molecular Medicine, Hospital for Sick Children, Toronto ON, Canada

[3] Department of Agricultural, Food and Nutritional Science, University of Alberta, Edmonton, Canada.

[4] Department of Biochemistry, University of Toronto, Toronto ON, Canada.

* To whom correspondence should be addressed.
jparkin@sickkids.ca

Keywords: metabolic modeling, keystone species, community assembly, poultry microbiome, cross-feeding

ABSTRACT

With their ability to degrade dietary fibers to liberate otherwise unavailable substrates, members of the Bacteroidales exert a substantial influence on the microbiome of the lower intestine. Currently our knowledge of how this influence translates to the metabolic interactions that support community structure is limited. Here we applied constraints-based modeling to chicken cecal communities to investigate metabolic interactions in the presence and absence of *Bacteroides*. From metagenomic datasets previously generated from 30 chicken ceca, we constructed 233 metagenome-assembled genomes. Metabolic modeling of communities built from these genomes generated profiles of short chain fatty acids largely consistent with experimental assays and confirmed the role of *B. fragilis* as a metabolic hub, central to the production of metabolites consumed by other taxa. In its absence, communities undergo significant functional reconfiguration, with metabolic roles typically fulfilled by *B. fragilis* assumed by multiple taxa. Beyond *B. fragilis*, we found *Escherichia coli* and *Lactobacillus crispatus* also mediate influential metabolic roles that vary in the presence or absence of *B. fragilis*. Compensatory adaptations adopted by the microbiome in the absence of *B. fragilis* resulted in metabolic profiles previously associated with inflammatory bowel disease in humans, including energy deficiency, increased lactate production and altered amino acid metabolism. This work demonstrates the potential of chicken cecal microbiomes to investigate the complex metabolic interactions and key contributions that drive community dynamics.

INTRODUCTION

The gut microbiome consists of a complex, yet stable, community of microbes that confer beneficial roles to their host including the extraction of key nutrients, promoting gut health and limiting colonization by pathogens. Helping organize these communities are metabolic interactions involving the sharing (cross-feeding) of metabolites between taxa[1]. Many of these interactions are thought to be mediated by so-called keystone taxa which play a crucial role in shaping microbial community structure and function[2–7]. Within gut communities, keystone taxa are represented by genera such as *Bacteroides* and *Lactobacillus* that have been shown to exert considerable influence within their communities[8,9]. For example, while *Bacteroides* spp. play an important role in the degradation of complex carbohydrates and production of short chain fatty acids (SCFAs)[10,11], *Lactobacillus* spp. have been identified as divisive members, characterized by both positive and negative interactions with other microbes[12]. Currently our understanding of the cross-feeding relationships that drive community dynamics and the contribution of keystone taxa is limited.

Community metabolic modeling offers a powerful approach to unravel the complex interactions within the gut microbiome. Integration of genome-scale metabolic models from multiple microbes within a community-scale framework provides a mechanistic basis to dissect the metabolic landscape of microbial communities and the contributions of individual taxa[13–15]. Recent advances in community-scale metabolic modeling have enabled predictions of the production of key metabolites such as SCFAs, which help support gut health[16,17] as well as identification of potential cross-feeding interactions between bacterial species[13–15] and changes in the concentrations of metabolites associated with a variety of human diseases[18,19]. Beyond human health, the gut microbiome has attracted considerable interest from the livestock industry. For example, with global bans on the use of antibiotic growth promoters (AGPs)[20] attention is turning to manipulating gut communities in chickens to address issues of food insecurity associated with a surge in enteric infections[20].

In this study, we leverage community metabolic modeling to investigate the influence of keystone taxa on microbial communities associated with the chicken cecum, a major site of digestion of insoluble fibers and fermentation and host to the most diverse community in the poultry gut.

Applying *in silico* modelling based on previously published metagenomic data, we investigate the

impact of *Bacteroides fragilis* on community and metabolic dynamics. Our models reveal that *B.*

fragilis significantly impacts microbial diversity and structure of metabolic interactions. In

Bacteroides-depleted communities, we observe an increase in the influence of other taxa including

Escherichia coli, *Anaerostipes butyraticus*, and *Lactobacillus crispatus,* leading to a functional

reorganization of the microbiome. Our findings highlight the pivotal role of the unique metabolic

capabilities of *B. fragilis* and the potential implications for host health and productivity in its

absence.

RESULTS

Bacteroides species influence the microbiome composition and functional capacity

To examine the metabolic influence of keystone taxa within cecal microbial communities we divided 33 samples from a recent study of 7-day old chickens[11] into two groups: one characterized by the presence of *Bacteroides* species (HB, n=15), specifically *B. fragilis*, and the other without *Bacteroides* species (NB, n=18)[11]. Metagenomic datasets previously generated from these samples were used to generate metagenome-assembled genomes (MAGs). To enhance MAG quality and consistency across samples, we conducted both individual assemblies for each sample and a co-assembly of all samples. This combined approach allowed us to capture species present in individual samples and improve assembly quality for shared taxa by leveraging the combined data. Additionally, we re-assembled common taxa by pooling the corresponding reads from multiple samples (see Methods). This process yielded 237 high- and medium-quality MAGs (defined as having ≥50% completeness and ≤10% contamination) representing 227 taxa. Each MAG represented at least 0.5% relative abundance in at least one of the 33 samples (**Figure 1A**). Communities were largely dominated by representatives of families such as Lachnospiraceae, UBA660, and taxa belonging to Oscillospirales order, with significant variations in their relative abundances across samples (**Figure S1**). Next, we generated genome-scale metabolic models for each MAG, refining models through gap-filling and manual curation to ensure their ability to support growth on a corn feed diet and incorporate missing carbohydrate-active enzymes and fermentation pathways (see Methods). Subsequently, models were combined to reflect the cecal samples from which they were derived resulting in communities represented by 5 to 33 taxa with an average of 1,096±98 unique enzymes (as defined by enzyme commission numbers; ECs; **Figure 1A**). Further, each community contained from 2,187 to 3,368 (mean 3,001±280) unique reactions encompassing from 143 to 598 (mean 495±83) unique pathways (defined in MetaCyc[21]), with the number of unique pathways being significantly higher in NB communities compared to HB (p=0.0045, Wilcoxon Rank-Sum test; **Figure 1B**).

Our reconstructed metabolic models revealed distinct metabolic profiles among taxa. *E. coli* had the largest number of enzymes, with broad representation across all pathways suggesting high functional diversity (**Figure S2; Table S1**), although we acknowledge this may also reflect its well

annotated genome. *Bacteroides*, *Alistipes*, *Helicobacter*, *Barnesiella*, and *Coprobacter* spp., found exclusively in HB communities, exhibited expanded functionality in glycan and vitamin metabolism, suggesting they may play a dominant role in the breakdown of dietary fibers and vitamin synthesis. While most taxa possessed core functionalities across the major pathways, species within UBA1242 (e.g., *Caccovivens* spp.), UBA660 (e.g., *Faecenecus* and *Faecimonas* spp.), and CAG-508 (e.g., *Merdicola* spp.) families displayed sparser coverage of functions within carbohydrate metabolism. These microbial groups have been recently identified in diverse animal and human gut environments and are characterized by limited metabolic capabilities[22–27]. Despite their distant phylogenetic relationships, they each exhibit reduced genome sizes, diminished genetic repertoires, and numerous auxotrophies[28,29]. Further, these taxa exhibited minimal overlap with other species in amino acid metabolism capabilities, with distinct sets of enzymes involved in amino acid metabolism and xenobiotic biodegradation, suggesting they occupy a unique niche (e.g. mitigating environmental stressors) within the gut microbiome (**Figure S2**).

Figure 1. Taxonomic composition and functional analysis of chicken cecal microbial communities. A. Relative abundances of metagenome-assembled genomes (MAGs) consolidated by taxonomic family across 33 chicken cecal samples. The bar plots represent the numbers of

unique species, ECs, unique reactions, and unique pathways based on metabolic model's content.
Samples are categorized into two groups: those with high Bacteroides abundance (HB, gold) and
those without Bacteroides (NB, green). **B**. Distribution of numbers of unique metabolic reactions
and pathways in NB and HB samples. Significance determined using Wilcoxon rank-sum test. **C,**
E. PCoA of samples based on abundances of pathways (**C**) or reactions (**E**) content (i.e., reaction
and pathway counts within each sample were summarized by weighing them by the relative
abundance of the species contributing to that reaction or pathway), with dissimilarity matrices
calculated using the Bray-Curtis index. **D, F**. PCoA of samples based on the presence or absence
of pathways (**D**) or reactions (**F**) using Jaccard distances.
130

Interestingly, while the prevalence of *Lactobacillus crispatus, Anaerostipes butyraticus* and *E. coli*

in NB samples suggests that these taxa may become more dominant in the absence of *B. fragilis*,

members of UBA660 family, specifically, *Faecenecus gallistercoris* and *Faecimonas*

intestinavium, exhibit relatively high abundance (15-68%) in samples where the former three taxa

are absent or exhibit low abundance (**Figure S1**). This may indicate competitive relationships,

where less metabolically versatile taxa exploit specific substrates or metabolic niches that are less

accessible to other bacteria. Members of the Oscillospirales order exhibited enhanced functionality

in the metabolism of cofactors and vitamins, as well as carbohydrate metabolism, aligning with

their roles in fiber degradation and SCFA production (**Figure S2**). These diverse metabolic

capacities and interdependent roles of bacteria in the chicken cecal microbiome illustrate the

complex relationships driving microbial community structure.

Principal Coordinate Analysis revealed that the presence of *B. fragilis* significantly influences both

the taxonomic composition and metabolic potential of microbial communities (**Figures 1C-F**). NB

and HB samples segregate on the basis of reaction and pathway content, regardless of similarity

metric used. The high percentage of variance explained by the first two principal coordinates (70%

for Bray-Curtis dissimilarity based on pathway abundance and 67% for Jaccard distance) together

with a significant difference in metabolic pathway profiles (PERMANOVA $p=0.001$) highlights

the impact of *B. fragilis* on community metabolism. However, the differences in within-group

variability are not significant (TukeyHSD test $p=0.13$, permutest $p=0.11$) between the HB and NB

groups (**Figure S3C**). Moreover, we observed distinct pathway and reaction abundance profiles

between HB and NB groups (PERMANOVA $p=0.001$ for both), with greater variability within the

NB communities (TukeyHSD $p=0.015$ and $p=0.006$, permutest $p=0.009$ and $p=0.004$, for

pathways and reactions, respectively; **Figure S3B-C**). Overall, these findings show that *B. fragilis*

has a major influence on metabolic interactions within the gut microbiome.

Community metabolic modeling accurately predicts the production of acetate and

propionate

To further dissect the metabolic contributions of *B. fragilis* within their communities we

constructed *in silico* models for each of the 33 samples using the BacArena framework[30] in which

the relative abundance of each taxon was defined from the metagenomic datasets. Our *in silico*

model was designed to mimic the physiochemical conditions of the chicken cecum, including low

oxygen, neutral pH, peristalsis and release of cecal contents[29–32](see Methods). To assess the

accuracy of our *in silico* ceca models, simulations were performed for 16 hours (representing a

cycle of feeding), and predicted SCFA concentrations were compared with those previously

measured experimentally[11]. We found a significant correlation between the predicted and

experimentally derived concentrations for both acetate (spearman rho=0.53, p-value=0.0015;

Figure 2A) and propionate (spearman rho=0.52, p-value=0.0021; **Figure 2A**). Moreover, the

simulations reflected the experimentally measured differences for the NB and HB communities,

although we note the model over-predicted the production of propionate by *Bacteroides* species

(**Figure 2B**, **Figure S4**). The model was less predictive of butyrate production, potentially as a

result of the model missing the impact of host epithelial cells which are known to uptake

microbially derived butyrate[31,32]. Further, the discrepancy in the prediction of butyrate flux arises

from the presence of butyrate-producing organisms (e.g. *Butyricicoccus* and *Faecalibacterium*

spp.) in samples where the measured concentration was nearly zero. Previous studies have

demonstrated cross-feeding interactions between *Bifidobacterium* spp. and *Faecalibacterium* spp.,

with acetate produced by the former enhancing butyrate production in the latter[33]. In our

simulations, acetate produced by *Bifidobacterium pullorum* was utilized by other taxa, including

Butyricicoccus and *Faecalibacterium* spp.[34–36], to generate butyrate (**Figure S4**). This highlights

the importance of studying metabolic interdependencies to better understand SCFA production in

the gut microbiome. Overall, our models were able to accurately capture general trends and known

dynamics of SCFA production, correctly predicting interactions between key gut microbes.

Figure 2. **Comparison of predicted and measured SCFA concentrations in chicken cecal samples. A**. Spearman rank correlation between *in silico* predicted and experimentally measured concentrations of acetate, propionate, and butyrate, with correlation coefficients (Spearman rho) and p-values indicated for each SCFA. Lines are fitted linear regressions, and colored areas around them indicate the 95% confidence interval. Each data point represents a sample. **B**. Bar plots comparing the average predicted and measured SCFA concentrations in samples with no Bacteroides (NB) and high (HB) Bacteroides abundance. The plots indicate the mean concentrations and standard errors for acetate, propionate, and butyrate with asterisks corresponding to the significance levels (Wilcoxon Rank-Sum test): * - p-value < 0.05, ** - p-value <0.01, *** - p-value < 0.001.

E. coli and *B. fragilis* **drive metabolic dynamics in the chicken cecal microbiome**

Our analysis of metabolic interactions within chicken cecal microbial communities revealed distinct structural differences between HB and NB communities. To further investigate these differences, we used mutual information (MI) scores to construct a network based on metabolic

overlap, where metabolic similarity refers to the extent of shared metabolic reactions between
communities. We found that NB communities exhibited higher metabolic similarity than HB
communities (pairwise MI scores 0.134±0.076 for NB and 0.123±0.068 for HB; **Figure 3A**).
Although the difference in MI scores between NB and HB communities was not statistically
significant (p=0.065, Wilcoxon Rank-Sum test), the trend suggests that the absence of *B. fragilis*
leads to greater metabolic similarity. Since *Bacteroides* spp. play a fundamental role as primary
fermenters and are often considered keystone taxa in their communities[37–40], their potential
monopoly of key resources may drive the reduction in community interactions and metabolic
diversity. Moreover, the presence of both *B. fragilis* and *E. coli* resulted in further
compartmentalization of metabolic functions, as evidenced by significantly lower overlap in
metabolic functions (MI score 0.128±0.061) compared to NB communities with *E. coli* (MI score
0.154±0.08; p=0.022, Wilcoxon Rank-Sum test) and without *E. coli* (MI score 0.21±0.11; p=1e-
05, Wilcoxon Rank-Sum test) (**Figure 3A**). Interestingly, samples lacking *E. coli* species exhibited
highest pairwise MI scores (0.213±0.111 and 0.21±0.11, for HB and NB communities
respectively**)** indicating that absence of *E. coli* promotes higher metabolic similarity irrespective
of the presence or absence of *B. fragilis.* These findings suggest that consistent with its wide range
of metabolic functions, *E. coli* acts as a metabolic generalist – a species capable of performing a
broad range of metabolic activities, in contrast to specialists that have more limited, niche-specific
roles – to fulfill roles vacated by other keystone taxa to support the metabolic versatility of the
cecal microbiome (**Figure S2; Table S1**).
218
While MI scores did not show a significant difference in metabolic similarity in NB and HB
communities, the influence of additional taxa such as *E. coli* led us to investigate whether the
absence of *B. fragilis* results in communities exhibiting a greater diversity of less dominant taxa,
with increased competition and metabolic interactions. We applied three complementary metrics
(see Methods): Metabolic Resource Overlap (MRO) score, a metric that quantifies both
competition for shared resources as well as complementarity in metabolic capabilities[41];
Community Metabolic Dissimilarity (CMD) score, a metric that quantifies the diversity of
metabolic functions in a community; and Cross-Feeding Coefficient (CFC) based on the number
of metabolite exchanges. We found that NB communities had significantly higher MRO (p=1.2e-
05, Wilcoxon Rank-Sum test) and CFC (p=0.025, Wilcoxon Rank-Sum test) scores than HB

communities (**Figure 3B-C**). While NB communities also featured higher CMD scores, consistent
with reports showing that metabolic dissimilarities enhance cross-feeding between species[42], this
difference was not significant. Overall, these findings show that NB communities exhibit greater
metabolic similarity between communities, along with enhanced cross-feeding relationships and
higher metabolic diversity within individual communities, supporting the hypothesis that the
presence of *B. fragilis* reduces metabolic diversity and interactions.

**Figure 3. *E. coli* and *B. fragilis* influence metabolic similarity and cross-feeding in chicken
cecal microbiomes. A.** Mutual information-based network of chicken cecal microbial
communities, highlighting metabolic similarities based on metabolic reaction content. Nodes
represent samples, with sizes corresponding to the number of unique reactions in a sample, colored
by category: no Bacteroides (NB, green) and high Bacteroides (HB, gold). Edges are colored and
thickened according to the mutual information score, with thicker and darker edges indicating
higher similarity. The top region outlined in red indicates samples lacking E. coli. Sizes of the
smaller nodes within a sample correspond to the number of reactions covered by the metabolic
models of species belonging to a respective taxonomic family (grouped by taxonomic order
below). **B.** Bar plots showing community sizes (top) and box plots showing Metabolic Resource
Overlap (MRO) within each sample (summarizing five *in silico* replicates). NB communities
exhibit significantly higher MRO (p=1.203e-05, Wilcoxon Rank-Sum test) **C.** Cross-feeding
coefficient (left) and Community Metabolic Dissimilarity (CMD) score (right) between NB and
HB communities. NB communities show a significantly higher cross-feeding coefficient
(*p=0.025) and a higher average CMD score.

Functional reorganization and emergence of alternate hubs in response to the absence of keystone taxa in the gut

Building upon the finding that the presence or absence of *B. fragilis* influences cross-feeding relationships among other taxa, we extended our analyses to examine changes in metabolite exchange dynamics in a larger set of taxa that could be considered metabolic hubs (defined here as taxa exchanging at least 12 metabolites with at least 10 other taxa across samples within each category; 87 taxa total - 40, 22 and 25 taxa associated with NB only, HB only and both, respectively). These taxa were used to generate profiles of interactions with other taxa and subsequently clustered to identify cliques of cross-feeding taxa (**Figure 4A; Table S2**). Consistent with our previous findings *B. fragilis* represents the central hub in the HB communities, while in its absence, *E. coli, A. butyraticus* and *L. crispatus,* adopt a greater range of metabolic interactions. The shift towards a more partitioned network observed in NB communities was further characterized by the formation of distinct cliques, representing groups of taxa engaging in mutual cross-feeding interactions. Within each clique, a single taxon (e.g. *Mediterraneibacter sp904377845, Lactobacillus johnsonii* and *Borkfalkia sp944368795*) appears to dominate interactions acting as a metabolic hub for each group of taxa.

To further clarify how 'metabolic hub' taxa exert their influence within their communities, we examined their interactions with other taxa in the context of prevalence across samples, relative abundance, and provision of unique functionality (**Figure 4B, Figure S5**). Surprisingly, a taxon's unique metabolic potential within its community (as estimated by the number of unique enzymes) did not correlate with its number of metabolic interactions (**Figure S5A**) although we did observe a positive correlation with prevalence (**Figure S5B**). These findings suggest that while the number of unique functions a taxon brings to a community might increase its ability to engraft, the community does not exploit that additional functionality. Indeed, *E. coli*, which is both prevalent and possesses the greatest set of unique functions, appears to benefit in its community from the consumption of many metabolites from multiple species with only modest contributions to production (**Figure 4B**). Other prevalent taxa exhibiting a broad range of interactions, such as *L. crispatus* and *A. butyraticus,* with fewer unique enzymes did not exhibit the same disparity in consumption and production. At the other end of the spectrum, we found some less prevalent taxa, present in only a limited number of samples (e.g., *L. johnsonii* and *M. sp904377845*) exert their

influence by specializing in the production of a wide array of metabolites utilized by a large number of other taxa (**Figure 4B**).

Figure 4. Major metabolic hubs and cross-feeding dynamics in HB and NB microbial communities. A. Heatmap of interactions between species, highlighting top cross-feeders defined as those forming at least 10 cross-feeding pairs with the exchange of a minimum of 12 metabolites across samples within each category, HB or NB. The interactions observed across multiple samples within each category were normalized by the number of samples in which each cross-feeding pair was present. Key taxa are labeled in this heatmap, including taxa present in at least two samples, forming an average of at least 230 interactions per sample and possessing on average at least 5 unique EC per sample. The tiles in the heatmap are colored based on the normalized total number of interactions for each species pair, with a color gradient indicating the interaction intensity.

Blocks of taxon interactions represent 'cliques', indicating groups of species with frequent metabolic exchanges. For example, the green-highlighted clique shows *Lactobacillus johnsonii* acting as a 'metabolic hub,' engaging in extensive interactions with taxa from the families Lachnospiraceae (*Lachnoclostridium B sp000765215*, *Fusicatenibacter sp900543115*, *Blautia ornithocaccae*, *Eisenbergiella sp904392525*, *Choladocola avistercoris*, *Merdisoma faecalis*), Acutalibacteriaceae (*Eubacterium R faecavium*, *Acutalibacter ornithocaccae*, *Anaeromassilibacillus stercoravium*), Borkfalkiaceae (*Coproplasma stercoravium*, *Coproplasma stercorigallinarum*), Anaerotignaceae (*Anaerotignum merdipullorum*), Butyricicoccaceae (*Butyricicoccus avicola*), Ruminococcaceae (*Gemmiger avicola*), and Oscillospiraceae (*Dysosmobacter pullicola*), as well as with each other. Taxa mentioned in the main text are highlighted in bold. **B**. Stacked bar charts representing the intensity of cross-feeding per taxon. The bars indicate the average number of interacting species for each taxon per sample, divided into metabolite consumption (left) and production (right). Colors correspond to the number of metabolites exchanged, with categories ranging from 1-5, 6-10, 11-15, and >15 metabolites. The bar plots to the right represent the sample count (number of samples in which each species is present, blue), relative abundance (grey), and the average number of unique enzymes (ECs) taxon possesses per sample (purple). Subset of taxa displayed is present in at least two samples, exhibit an average of at least 230 total interactions per sample, and possess a minimum of 5 unique EC numbers per sample.

These findings reveal a hierarchy of taxa acting as metabolic hubs, where taxa such as *B. fragilis* and *E. coli* are responsible for driving the bulk of interactions, supplemented by additional hubs like *A. butyraticus* and *L. crispatus*. Further, within their communities, there exist cliques of taxa sharing metabolic interactions, organized around relatively rare taxa. This hierarchy highlights the varying abilities of each type of hub taxa to colonize within their communities, with *B. fragilis* and *E. coli* capable of colonizing many communities and establishing extensive cross-feeding networks. Others such as *M. sp904377845* and *L. johnsonii*, appear more limited in their ability to colonize communities and feature a smaller range of interactions with only a subset of taxa.

Role of key metabolic hubs in modulating cross-feeding interactions

To further assess how the presence/absence of different metabolic hubs may impact cross-feeding relationships within their communities, we constructed a linear model based on the abundance of *E. coli* and the presence/absence of three additional metabolic hub taxa (*A. butyraticus, L. crispatus* and *B. fragilis*; see Methods). A relatively high abundance (>5%) of *E. coli* abundance was associated with increased cross-feeding in *Bacteroides*-depleted communities, potentially indicating the ability of *E. coli* to replace community functions otherwise performed by *B. fragilis* (**Figure S5C; Table S3**). Interestingly, only one of the 15 HB samples exhibited such an abundance, indicating competition between these two taxa. Extending this analysis to two other

metabolic hub taxa (*A. butyraticus* and *L. crispatus*), we found their presence in the absence of *E.*
coli was associated with increased cross-feeding activity, suggesting their ability to provide an
alternative source of metabolic functions (**Figure S5D-E; Table S3**). Communities featuring both
E. coli together with *A.butyraticus* displayed lower cross-feeding activity, reflecting potential
competition between these taxa. In contrast, NB communities with *L. crispatus* showed higher
cross-feeding activity compared to HB communities, regardless of *E. coli* abundance, indicating
an antagonistic relationship between *L. crispatus* and *Bacteroides*.

Since our linear model focused only on taxa representing metabolic hubs, we were interested in
how other metabolic hub taxa, specifically those mediating interactions within a subset of taxa
(cliques), contribute to their communities. To avoid the confounding influence of dietary
metabolites, we further restricted these analyses to metabolites only produced by bacteria. As
before, *B. fragilis* represents the central hub in the HB communities, while *E. coli, A. butyraticus*
and L. crispatus assume additional interactions in NB communities (**Figure S6A; Table S4**).
Moreover, we also observed the formation of metabolic cliques, with greater numbers of
interactions in NB communities. Comparing the eight largest cliques specific to either NB or HB
communities (labelled A-H), we identified 48 unique microbially-derived metabolites exchanged
within these cliques (ranging from 24 in clique H to 37 in clique A). While taxa associated with
these cliques may not necessarily be present in the same samples, these cliques nonetheless
represent groups of taxa that occupy a similar metabolic niche – that is, they share specific sets of
metabolic functions and interactions they engage in, contributing to similar ecological roles across
different communities.

This aligns with previous research demonstrating enrichment of mutualistic interactions in co-
occurring microbial communities[41]. While there is a substantial overlap in metabolites exchanged
across the eight cliques investigated – 17 metabolites were common to all cliques, and an additional
16 were shared among at least five cliques (**Figure S6B**) – individual cliques exhibit unique
metabolic exchanges reflecting the specific taxa involved. This suggest that, despite a shared core
of metabolic pathways and intermediates, each clique possesses distinct metabolic interactions
driven by its constituent taxa. For instance, in the NB-specific clique B, *Bifidobacterium* spp. And
members of Lactobacillaceae uniquely produced and exchanged D-alanine, melibiose, and

spermidine (**Figure S6A-B**). Relative to other cliques, clique D, comprising members of
Oscillospirales, Lachnospirales, Christensenellales, Coriobacteriales, and RF39, exhibited a more
limited range of metabolite exchanges, lacking common fermentation products but uniquely
exchanging NAD, highlighting the role of redox reactions and energy metabolism within this
clique. These findings highlight how specific taxa contribute to functional diversity within
microbial communities by occupying distinct metabolic niches, thereby shaping the overall
metabolic landscape of their environments.

The changes in cross-feeding relationships associated with the absence of *B. fragilis* led us to
investigate how this translates in terms of exchange of specific metabolites. We therefore analyzed
the production and consumption of 18 metabolites selected on the basis of their relationship with
4 keystone taxa (see Methods; **Figure 5**). As expected, in HB communities *B. fragilis* is the
predominant producer, contributing to the majority of the production of 10 metabolites, with *E.*
coli, Mediterraneibacter glycyrrhizinilyticus, Faecalibacterium gallistercoris, and
Dysosmobacter pullicola largely responsible for the production of a more limited set of metabolites
including methanol, indole, uridine, formate and L-glutamine. In NB communities, the production
of these metabolites is redistributed to multiple other species. For example, consistent with
previous reports[33,36,43,44], in the absence of *B. fragilis*, *E. coli* assumes the production of multiple
metabolites otherwise produced by *B. fragilis* including acetate (along with members of
Clostridia), ammonium (with *A. butyraticus*), succinate, propionate, sorbitol and GABA. These
activities are supported through the production of lactate, uridine and L-glutamine by *L. crispatus*,
the latter consistent with previous reports of its production[45]. In the presence of *B. fragilis*, *E. coli*
alters its metabolic contributions to the generation of methanol, itself utilized by *B. fragilis.*
Interestingly, in NB communities *A. butyraticus* adopts the role of methanol production (**Figure**
5).

Together our predictions show that in the absence of a keystone taxon (here *B. fragilis*), the cecal
microbial community undergoes significant reorganization, characterized by the emergence of
alternate metabolic hubs as well as a shift in metabolite production and consumption profiles by
taxa present in both types of communities.

Figure 5. Shifts in cross-feeding networks: metabolic role redistribution in HB and NB communities. The Sankey diagram illustrates the cross-species interactions between primary producers and consumers of key cross-fed metabolites in the HB and NB communities. Producers are displayed on the left, cross-fed metabolites in the middle, and consumers on the right. Gold links represent contributions in HB communities, while green links represent those in NB communities. The thickness of each link is proportional to the contribution of the species (as a percentage) to the total production or consumption of each metabolite. To highlight the major players in cross-feeding dynamics, only the top three producers and consumers for each metabolite are shown, with a cutoff set at a minimum 5% contribution across samples within each category. Species highlighted in bold were previously described in Figure 4B. The diagram provides a visual summary of how key microbial taxa participate in the metabolic exchange, highlighting functional shifts between HB and NB communities.

Metabolic reorganization associated with the loss of *Bacteroides* in the chicken cecum reflects disease-associated metabolism in human IBD

Our comparisons of HB and NB communities, exhibit notable similarities to microbiome profiles associated with Inflammatory Bowel Disease (IBD) in humans (Ulcerative Colitis - UC and Crohn's Disease - CD). Previous studies have reported shifts in the abundance and roles of both *B. fragilis* and *E. coli* – major metabolic hubs in our cecal communities – that correlate with disease states. The former being associated with reduced abundance in IBD patients, while the latter is associated with increased abundance[46–50]. Additionally, we observed increased abundance of

Alistipes and *Faecalibacterium* species in HB samples (**Figure S1**), consistent with communities
associated with healthy individuals in contrast to IBD patients[49,51,52]. Similarities between our
study of HB and NB communities with healthy/IBD communities also extend to metabolic
profiles. For example, the heterolactic fermentation pathway associated with lactic acid bacteria is
more prevalent in NB communities, resulting in increased starch consumption and lactate
production (**Figure 6**), consistent with elevated lactate levels observed in IBD patients[53–55]. NB
communities are also associated with reduced production of propionate, a key source of energy for
gut epithelial cells and a hallmark of IBD[55–58]. While studies have reported varied metabolic
profiles in IBD, with both increased and decreased energy metabolism[55,59–61], our findings align
with the broader trend of altered energy metabolism in disease state. Specifically, we observed a
metabolic shift in *Bacteroides*-depleted communities characterized by increased consumption of
coenzyme A (CoA) and glycerol, alongside increased production of TCA cycle intermediates such
as 2-oxoglutarate and citrate, as well as GTP. CoA, essential for fatty acid metabolism and the
TCA cycle as an acyl group carrier, is being consumed more by the microbial community,
suggesting upregulated lipid and energy metabolism pathways. However, the concurrent
accumulation of TCA intermediates may indicate a bottleneck or dysregulation within the
microbial TCA cycle, potentially due to limited CoA availability. This imbalance could reflect a
shift towards fermentative metabolism in the absence of B. *fragilis*. Notably, reduced CoA levels
have been observed in the colonic tissues of pig models and human patients with ulcerative colitis
(UC), where they are associated with impaired fatty acid oxidation and energy metabolism in host
cells[55,62]. While our findings pertain to increased microbial consumption of CoA, this could
potentially decrease its availability in the gut lumen, impacting the host's energy metabolism.
432
Beyond changes in energy metabolism, our models also predict an increase in production of amino
acids, including L-phenylalanine, L-valine and L-lysine in NB communities relative to HB (**Figure**
6), consistent with recent studies reporting alterations in levels of these amino acids in IBD
patients[53,57,63,64]. Other notable metabolic similarities include reduced profiles of 5-Methylthio-D-
ribose, a metabolite involved in the methionine salvage pathway, in NB communities, indicative
of reduced sulfur cycle activity and consistent with deficiencies in sulfur-containing metabolites
in microbial communities of IBD patients[18,19]. Thiamin, typically deficient in CD patients[53,65], is
predominantly produced by *B. fragilis*, and consequently reduced in NB communities. Finally, the

increased consumption of sialic acid (Neu5Ac), a mucin-derived monosaccharide, by NB
microbiomes suggests enhanced sialidase activity, which has been linked to intestinal
inflammation and provides a growth advantage to bacteria like *E. coli*[66,67]. Thus, while the earlier
study reported reduced inflammation in HB chicken cecal microbiomes[11], the opposite may be
inferred for NB microbiomes.

Figure 6. Comparison of metabolite exchange patterns between HB and NB communities.
Each dot represents a metabolite being exchanged, with black dots indicating metabolites that do
not show statistically significant changes in production or consumption between NB and HB
groups. Statistically significant differences in total community fluxes (normalized to community
size) were identified using the Wilcoxon rank-sum test. Dashed red line indicates the threshold for
significance level, p-value < 0.05. Metabolites are color-coded based on the direction of their
exchange: green - increased consumption in HB; blue - increased production in HB; orange -
increased consumption in LB; purple - increased production in LB; red - switch from production
in LB to consumption in HB; pink - switch from consumption in LB to production in HB.
Metabolites with an absolute log2FoldChange greater than 0.7 are labeled and colored accordingly;
metabolites discussed in results are highlighted in bold.
457

Overall, our findings reveal parallels between the community and metabolic dynamics of chicken

cecal communities in the absence of *B. fragilis* and those associated with IBD, suggesting

Bacteroides-rich communities may confer a protective role against inflammation.

DISCUSSION

Previous studies suggest microbial communities in the gut display a hierarchical structure, with

so-called keystone taxa acting as influential members through their ability to provide their

communities with unique metabolic capacities[7,68–70]. The Bacteroidaceae family members are

considered to be one of the most influential taxa, exerting a significant impact on metabolic

interactions withinn their communities[37–39,71,72]. Previous studies have shown that poultry gut,

specifically the cecum, exhibits substantial fluctuations in the abundance of members of the

Bacteroidaceae family, including the genus *Bacteroides*[11,73,74]. Notably, *Bacteroides* spp. are less

abundant in intensively raised chickens compared to those raised in extensive systems (i.e. with

access to the outdoors)[75]. Shifts in the abundance of *Bacteroides* can impact the community

dynamics resulting in reduced feed efficiency, and heightened susceptibility to enteric diseases[76–

78]. In humans, such shifts have also been associated with IBD[79,80].

Here we leveraged a pre-existing dataset of metagenomic data from the cecal communities of

chickens displaying a range of *Bacteroides* abundance and performed community metabolic

modelling to dissect the influence of keystone taxa on metabolic interactions within their

communities. Notably we found that our sample-specific community models accurately predicted

the production SCFAs, particularly acetate and propionate, and recapitulated experimentally

observed differences between *Bacteroides*-rich (HB) and -depleted communities (NB). Our

findings demonstrate that the absence of *B. fragilis* results in more similar metabolic profiles across

different microbiomes, while simultaneously increasing the range of metabolic functions within

each individual community. However, this shift does not necessarily lead to an increase in overall

microbial diversity. Moreover, NB communities exhibited significantly higher metabolic resource

overlap within communities and increased cross-feeding interactions. This trend aligns with the

role of *Bacteroides* spp. as keystone taxa, which typically dominate and monopolize key metabolic

functions, thereby reducing metabolic diversity within their communities[81,82].

By defining metabolic hubs as microbial taxa with a central role in metabolite exchange and
distribution, we examined their influence on the overall metabolic landscape. While *B. fragilis* acts
as a central hub, its absence triggers a shift towards a decentralized network characterized by
distinct cross-feeding cliques and the emergence of new metabolic hubs, such as *E. coli*, *A.*
butyraticus, and *L. crispatus*. Contrary to the expectation that a larger enzymatic repertoire directly
translates to greater community influence, we observed a diverse range of strategies, from
generalists like *E. coli,* which can perform a broad range of metabolic activities, to specialists like
L. johnsonii, which focus on specific metabolic functions**.** Our analyses show that *E. coli* has a
significant impact on cross-feeding, with increased activity when *E. coli* abundance exceeds 5%
in *Bacteroides*-depleted communities. Interestingly, *L. johnsonii* and *L. crispatus,* previously
described as eliciting a Marmite effect on the gut microbiome[12], emerged as alternative metabolic
hubs in the absence of *Bacteroides*, reflecting the antagonistic relationship between *Lactobacilli*
and *Bacteroides* a observed in previous studies[11,83–85]. While *L. crispatus* had a positive effect on
cross-feeding in the absence of *E. coli*, its presence in samples where *E. coli* was abundant led to
a decrease in metabolic interactions. These observations highlight the shifting role of *L. crispatus*
as it can both contribute to and limit the community's metabolic interactions, depending on the
presence of keystone taxa (e.g. *E. coli* and *B. fragilis*).
505
In our study we found *B. fragilis* plays a central role in the production of metabolites utilized by
other taxa whereas NB communities were characterized by a greater distribution of metabolic
functions across a broader range of taxa. While this might be expected to contribute to community
stability, similar changes in distributions are thought to contribute to disease associated with the
human gut[86,87]. Indeed, our comparative analysis of HB and NB communities revealed striking
parallels with IBD-affected human gut; NB communities exhibit increased lactate production,
energy deficiency, and altered amino acid metabolism, alongside increased markers of
inflammation. In contrast, HB communities displayed metabolic signatures associated with a
healthy gut state including a higher production of SCFAs and essential vitamins, such as thiamin.
Further, consistent with our cecal models, a decreased abundance of *Bacteroides*, particularly *B.*
fragilis, which are typically abundant in healthy individuals, has been observed in patients with
IBD, allowing opportunistic pathogens such as *E. coli* to flourish[46–50]. These findings highlight the

potential of using models of chicken cecal communities to study gut microbial ecology and disease

mechanisms.

This work establishes a framework for utilizing community metabolic modeling to study the

poultry gut microbiome. Our approach facilitates the prediction of metabolic outputs, such as

SCFA production, as well as elucidation of cross-feeding relationships. Moreover, our framework

offers a mechanistic understanding of how keystone taxa, like *Bacteroides,* influence the

ecological dynamics and metabolic landscape of the gut microbiome. Despite its promising results,

we acknowledge our platform is not without some limitations. For example, community-level

metabolic models may risk overemphasizing the metabolic roles of well characterized taxa, such

as *E. coli*[88,89]. Further, the absence of host-microbiota interactions ignore the potential contribution

of the host in terms of uptake of metabolites, as well as the influence of the host immune system.

While the models accurately predict overall trends in metabolite production, they may

overestimate or underestimate specific metabolic fluxes. This can be attributed to limitations in

the functionality of some bacterial models, influenced in turn by the quality of the corresponding

MAGs. Finally, the lack of data from upstream gut compartments may also limit our understanding

of the precise metabolic environment in the cecum; here we assume that dietary components in

amounts adjusted for known estimates of upstream absorption are directly available to members

of the cecal community, ignoring the potential metabolic contributions from microbes present

upstream of the ceca. Future research could integrate host factors, explore other gut regions, and

assess the potential for targeted interventions to beneficially manipulate the microbiome.

MATERIALS AND METHODS

External data collection

The experimental methods and data collection procedures have been described previously[11].

Publicly available raw sequence data and associated metadata associated with the study were

downloaded from the sequence read archive (SRA).

Chicken cecal metagenomic samples analysis

Metagenomic reads were subjected to quality filtering using BBTools/BBMap (v39.03)[90]. Single-sample assemblies and co-assemblies of short reads were conducted with MEGAHIT (v1.2.9)[91], specifying a minimum contig length parameter of '–min-contig-len' set to 1000. The Bowtie2 (v2.2.3)[92] tool was employed to index assemblies and map filtered reads onto the contigs, creating abundance profiles. Following this, the metagenome assemblies underwent separate binning processes utilizing MetaBAT2 (v2.13)[93] and CONCOCT (v1.1.0)[94]. Initially, samtools (v1.9)[95] sorted the read mappings, and the read coverage was determined using the MetaBAT2 script 'jgi_summarize_bam_contig_depths'. Subsequently, the binning process involved the 'metabat2' command with default settings for each assembly, using contig coverage across samples as input. For CONCOCT, accessory scripts were employed to cut contigs into 10 kb fragments (cut_up_fasta.py), and the read coverage for these fragments was computed using CONCOCT_coverage_table.py. The fragment coverages were then utilized for binning the 10 kb fragments, and the clustered fragments underwent merging (merge_cutup_clustering.py) to generate the final CONCOCT bins (extra_fasta_bins.py). To dereplicate and consolidate the produced outputs from CONCOCT and MetaBAT2 binners, the metaWRAP (v1.2.3)[96] 'bin refinement' command was employed, utilizing parameters '-x 10 -c 50'. These parameters signify a maximum bin contamination threshold of 10% and a minimum bin completeness threshold of 50%, as determined by CheckM[97] estimates. Taxonomic annotation of generated bins was performed using GTDB-Tk (v2.3.2)[98] command 'gtdbtk classify_wf' with default settings. Finally, the abundances of the draft genomes (bins) across the samples were quantified using the metaWRAP 'quant_bins' module.

To improve the consistency and quality of models representing the same species across different samples, the bins for each common species (species that are not unique to one sample) were reassembled by pooling the reads corresponding to that species from all samples. These reassembled bins were then re-binned using the previously described procedure to generate a single MAG per species (**Figure S3A;Table S5**). The only exception was a pooled *E. coli* MAG that was 8.2Mb in size, which exceeds the typical size range for *E. coli* genomes, suggesting it was likely an artifact resulting from strain heterogeneity. To represent the multiple *E. coli* strains present across samples without retaining the 23 sample-specific *E. coli* individual assemblies with varying

size and completeness, the bins derived from the individual assemblies were clustered based on

average nucleotide identity using the fastANI software[99]. A distance matrix was constructed, and

the bins were then clustered using the *hclust* function from the R-package stats. Five clusters were

selected, and five cluster-specific *E. coli* strains were re-assembled, each representing a species

pooled from samples within the corresponding cluster, followed by re-binning as described earlier.

The resulting 5 MAGs vary in size from 4 to 5.8 Mb which is within the expected range for the *E.*

coli genome[100] (**Figure S3A**).

As a result of applying these steps to the co-assembly of all samples, we found that only five

species had MAGs from the co-assembly that were of higher quality - based on completeness,

contamination, and N50 metrics - than those obtained from the pooled or single-sample assemblies.

These species were *Bacteroides fragilis*, *Anaerostipes butyraticus*, *Alistipes excrementavium*,

Limihabitans stercoravium, and *Caccovivens* s*p944345705*. Consequently, for these five species,

the representative MAG was selected from the co-assembly. The characteristics and classification

of the 237 reconstructed MAGs are provided in the **Table S5**.

Model reconstruction and curation

The MAGs with a minimum relative abundance of 0.5% in each sample underwent reconstruction

using gapseq v1.2 (commit 3e2d9393, Sequence DB md5sum: 17e92c9) with default settings[101],

resulting in a total of 233 metabolic models. Each sample was represented by 5 to 33 unique

MAGs. Since the specific dietary composition used as chicken feed was not disclosed in the

original study, the nutritional input for the models was formulated based on a typical chicken corn-

based diet[102] by translating molecular constituents into corresponding metabolites using vmh.org,

foodb.ca and feedtables.com resources[103–105]. Due to limited information on fiber in the chicken

diet, cellulose was utilized as a primary source of fibre according to available literature[106,107]. The

absorption and digestion of metabolites in the upper gut compartments were taken into account

when calculating the dietary input based on available information from studies on dietary

absorption in poultry gut[108–113]. Additionally, urea and mucus were introduced to depict the cecal

intestinal setting. Subsequently, this resulting nutritional composition served as an input medium

for gapfilling step via gapseq. To ensure microbial degradation capabilities towards dietary

components, profiling of carbohydrate-active enzymes (CAZymes) encoded within metagenomes

used for model reconstruction was performed using dbCAN3 tool[114], which is established upon

CAZy database - an expert-curated and routinely updated resource. Predicted CAZymes were

filtered by e-value threshold (> 1e-50) and required minimum 90% coverage with enzyme(s)

assigned EC number(s). Predicted EC numbers along with substrates were mapped to respective

reactions within modelSEED database and incorporated into models through gapseq adapt.

Additionally, metabolic models of the species known as SCFA producers, were examined on the

presence of fermentation pathways documented in the literature; missing pathways were added to

the models.

Quantitative PCR

qPCR of cecal digesta was performed using PercfeCTa SYBR green Supermix (Quantabio) with

Primers used target the universal 16S rRNA gene: Forward CGGYCCAGACTCCTACGGG,

Reverse TTACCGCGGCTGCTGGCAC[115]. The copy numbers were calculated back to copy

numbers per gram of cecal digesta.

Set up of *in silico* ceca and simulation of sample-specific community models

The microbial communities were simulated using the BacArena software[30], which combines Flux

Balance Analysis and individual-based modeling. BacArena represents each organism individually

on a two-dimensional grid to model a spatial environment. Metabolites can be exchanged between

individuals and diffuse in the environment, allowing for the dynamic update of the state of each

microbe and its surroundings. The initial microbial community was seeded on a 100x100 cell grid.

To mimic the flow of luminal contents in and out of the cecum while maintaining a relatively

stable bacterial population, the following approach was implemented. Multiple studies have shown

that the cecal contents are released only 2-3 times per day [116,117], which is a key feature for realistic

simulation, considering the cecum's role as a major hub for bacterial fermentation and SCFA

production.

At each hourly iteration, metabolites representing the metabolic contents from the upstream

compartment were added to the grid and evenly distributed between the grid cells, simulating

environmental mixing and allowing access to dietary and bacterial-derived metabolites to all cells

within the simulated environment. Every fourth iteration, a fraction of the cecum, including both

metabolites and microbes, was randomly removed to mimic emptying of the cecum. The number

of grid cells selected for removal of contents was determined to be 25% of the total to maintain

realistic fluctuations in bacterial density and a generally consistent community size. Low oxygen

concentrations (1e-08 mM) were to establish a nearly anaerobic environment reflective of reported

conditions in chicken ceca, along with pH of 7[118,119].

The absolute abundances of microbial communities were measured via quantitative PCR (qPCR)

assay, yielding copy numbers per gram of digesta ranging from 1.62e+09 to 3.02e+11. To reduce

skewness and ensure smaller values were distinguishable while preventing larger values from

dominating, we log-transformed these abundance values. Next, the log-transformed values were

rescaled to fit between 1000 and 5000. These rescaled values were then used as the total

abundances for the microbial communities in each sample. This approach ensured that even

species with low abundances were represented by at least 5 individuals on the grid, while also

allowing space for the community to grow and move within a grid of 10,000 cells. The

communities were simulated in 5 replicates during 16 iterations, each representing an hour, to

mimic the day of feeding.

Comparison of functional capacities of community models

Sets of unique reactions and pathways within each sample-specific microbiome were extracted

from the corresponding metabolic models comprising the sample. Binary matrices indicating the

presence or absence of reactions and pathways in each sample were created for subsequent Jaccard

distance calculations. Additionally, to quantify the metabolic potential within each sample while

accounting for the varying contributions of different species, we summarized the reaction and

pathway counts within each sample. Each reaction or pathway count was weighted by the relative

abundance of the species contributing to that reaction or pathway. The resulting incidence matrices

provided detailed information on the pathway and reaction abundance in the samples. Dissimilarity

matrices were constructed from the incidence matrices described using *vegdist* function from the

vegan R-package, and Principal Coordinate Analysis was performed using *cmdscale* function from

stats R-package. The network showcasing functional similarity between sample-specific

microbiomes was constructed using mutual information score calculated via *mutinformation*

function from infotheo R-package based on the metabolic reaction content of the samples. This

network was visualized using Cytoscape v.3.10.3[120], where large nodes represent samples, with

their sizes corresponding to the number of unique reactions in each sample. The smaller nodes

within a sample, representing taxonomic families, have sizes corresponding to the number of

reactions covered by the metabolic models of the species belonging to the respective taxonomic

families.

Community metabolic dissimilarity (CMD) score was calculated as the average of pairwise

metabolic dissimilarities based on EC numbers within the community, adjusted for the community

size. Specifically, the Bray-Curtis dissimilarity matrix for a sample was calculated from a

presence/absence matrix where the columns represent species and the rows represent EC numbers,

using the vegdist function from the vegan R-package. CMD score was then computed as a mean

of the dissimilarity matrix, with further log-adjustment to account for the diminishing contribution

of additional species in larger communities.

Evaluation of cross-feeding and competition in *in silico* microbiomes

To evaluate the extent of metabolic competition within Bacteroides-rich and Bacteroides-depleted

microbial communities, the metabolic resource overlap (MRO) between species was quantified

using the approach described in the original study[41]. The metabolic resource overlap (MRO) is a

metric that estimates the extent of overlap between the growth requirements of two or more

metabolic models[41]. As such, it quantifies the potential for species to compete for a predetermined

set of compounds in the media.

Cross-feeding interactions within microbial communities were investigated using metabolic

reaction fluxes. We filtered negligible fluxes to retain absolute total flux values above 1e-06

millimoles per hour per gram of dry weight (mmol/gDW/hour) and categorized species as

metabolite producers or consumers. By merging producer and consumer datasets on shared

metabolites, cross-feeding interactions were summarized by counting and normalizing interaction

occurrences across replicates within each category (based on *Bacteroides* presence). Key cross-

feeders were identified using a threshold of at least 10 interactions involving the exchange of 12

or more distinct metabolites, focusing on species with substantial and consistent metabolic activity.

Cross-feeding was quantified by retrieving all metabolite exchanges (import and export fluxes)

during simulations and calculating the number of total interactions (cross-feeding pairs) in each

sample. A cross-feeding coefficient was introduced as a metric of cross-feeding activity within a

community, defined as the number of total cross-feeding pairs formed divided by all possible

combinations of species pairs in the community. Interaction data were visualized using

ComplexHeatmap R-package.

Next, the total production and consumption for each metabolite was calculated over the course of

the simulation. The contribution of each species to the production and consumption was

determined as a percentage of the total production/consumption. Exchanged metabolites were

filtered to include only non-dietary metabolites, which are exclusively produced by bacteria. The

pairwise interactions involving non-dietary metabolites summarized across samples are visualized

as a heatmap in **Figure S6A** with raw data used to plot heatmap provided in **Table S3**.

To focus on metabolic niche adjustments involving keystone taxa, metabolites were further filtered

based on the following criteria: 1) they are predominantly produced or consumed by keystone taxa,

defined as those where at least 30% of produced metabolite is attributed to the production or

consumption by a single keystone taxon; 2) to ensure the metabolite is involved in cross-feeding,

at least 1% of the produced amount must be consumed by other species; 3) metabolites with the

same top producer (taxon that produces the highest amount of the metabolite) in both HB and NB

communities were excluded to emphasize shifts in metabolic roles. This filtering process resulted

in the identification of 22 key cross-fed metabolites. Of these, 18 were selected for downstream

visualization. Simple sugars D-Gluconate, D-Ribose, and D-Arabinose were excluded due to a

very low production levels, and L-Lactate was excluded because its production and consumption

patterns significantly overlapped with those of D-Lactate, so only D-Lactate is shown. For each

key cross-fed metabolite, the top three producers and consumers were identified based on their

relative contributions. The flux data was then subset to include only these key producer and

consumer species. Sankey diagrams were generated using the networkD3 R-package. To highlight

major contributors, a minimum contribution threshold of 5% was used for species shown in **Figure**

5.

To investigate differences in metabolite production and consumption between HB and NB microbial communities, the fluxes of exchanged metabolites were summarized within each community at every hour of the simulation and normalized by the total community size at that time point. Subsequently, the community fluxes (normalized by abundance) were aggregated over the 16-hour simulation period, and differences between the categories were evaluated using the Wilcoxon rank-sum statistical test. For graphical representation, metabolic fluxes were log2-transformed.

Statistical analyses

All statistical analyses were performed in R v4.2.3. Significant differences for numbers of reactions and pathways, metabolite concentrations, metabolic fluxes, community metabolic diversity and cross-feeding activity coefficients were determined by non-parametric Mann–Whitney U-test (Wilcoxon rank-sum test). Spearman's rank correlation coefficients and associated p-values were calculated between measured and predicted SCFA concentrations. Given *E. coli*'s high degree of connectivity, we explored its influence on metabolic exchanges within microbial communities. To evaluate how the presence or absence of this pivotal taxon determines the degree of cross-feeding activity, we constructed a linear model incorporating community size and the abundance of *E. coli* alongside key metabolic hubs: *A. butyraticus, L. crispatus,* and *B. fragilis* using the *lm* function in R. The response variable was the cross-feeding coefficient (CrossFeeding_coef), which quantifies the degree of metabolic interactions within the community. Predictor variables included categorical levels of *E. coli* abundance (Ecoli_category: "NoEcoli", "LowEcoli" (<5% relative abundance), "HighEcoli" (>5% relative abundance)), *Bacteroides* presence or absence that defines the community category (Bacteroides_presence: "NB" or "HB"), the presence of key taxa such as *Anaerostipes butyraticus* (Abutyraticus_presence) and *Lactobacillus crispatus* (Lcrispatus_presence), and community size (comm_size). Interaction terms between *E. coli* abundance and the other predictors were included to capture potential synergistic or antagonistic effects on cross-feeding. The model was specified as follows:
 'CrossFeeding_coef ~ Ecoli_category:(Bacteroides_presence + Abutyraticus_presence + Lcrispatus_presence) + comm_size'. The model summary is provided in the **Table S3**. All statistical analyses employed a significance threshold of p-value < 0.05.

FUNDING SOURCES

This work was funded by grants from the Natural Sciences and Engineering Research Council of

Canada (RGPIN-2019-06852) and the Ontario Research Fund-Research Excellence. High

performance computing was provided by the SciNet HPC Consortium. SciNet is funded by

Innovation, Science and Economic Development Canada; the Digital Research Alliance of

Canada; the Ontario Research Fund: Research Excellence; and the University of Toronto. The

funders had no role in study design, data collection and analysis, decision to publish, or preparation

of the manuscript.

AUTHOR CONTRIBUTIONS

J.P. and I.U. conceived and designed the study. I.U. performed simulations and data analysis. Y.F.

performed qPCR experiments. B.P.W. provided critical reagents and contributed to data analysis. J.P.

and I.U wrote the manuscript. All authors reviewed and/or edited the manuscript.

COMPETING INTERESTS

The authors declare no competing interests.

DATA AVAILABILITY STATEMENT

Microbial metabolic models and the source code used for simulations and data analysis have been

deposited on GitHub (github.com/ParkinsonLab/Chicken_microbiome).

REFERENCES

1.   Culp, E. J. & Goodman, A. L. Cross-feeding in the gut microbiome: Ecology and

mechanisms. *Cell   Host   and   Microbe*   vol.   31   Preprint   at

https://doi.org/10.1016/j.chom.2023.03.016 (2023).

2.   Gibbons, S. M. Keystone taxa indispensable for microbiome recovery. *Nature Microbiology*

vol. 5 Preprint at https://doi.org/10.1038/s41564-020-0783-0 (2020).

3.   Banerjee, S., Schlaeppi, K. & van der Heijden, M. G. A. Keystone taxa as drivers of

microbiome structure and functioning. *Nature Reviews Microbiology* vol. 16 Preprint at

https://doi.org/10.1038/s41579-018-0024-1 (2018).

4.   Herren, C. M. & McMahon, K. D. Keystone taxa predict compositional change in microbial
communities. *Environ Microbiol* **20**, (2018).

5.   Garrett, W. S. *et al.* Enterobacteriaceae Act in concert with the gut microbiota to induce
spontaneous and maternally transmitted colitis. *Cell Host Microbe* **8**, (2010).

6.   Hajishengallis, G. *et al.* Low-abundance biofilm species orchestrates inflammatory
periodontal disease through the commensal microbiota and complement. *Cell Host Microbe*
10, (2011).

7.   Banerjee, S. *et al.* Network analysis reveals functional redundancy and keystone taxa
amongst bacterial and fungal communities during organic matter decomposition in an arable
soil. *Soil Biol Biochem* **97**, (2016).

8.   Lan, P. T. N., Hayashi, H., Sakamoto, M. & Benno, Y. Phylogenetic analysis of cecal
microbiota in chicken by the use of 16S rDNA clone libraries. *Microbiol Immunol* **46**,
(2002).

9.   Qi, Z., Shi, S., Tu, J. & Li, S. Comparative metagenomic sequencing analysis of cecum
microbiotal diversity and function in broilers and layers. *3 Biotech* **9**, (2019).

10.   Oakley, B. B. *et al.* Successional changes in the chicken cecal microbiome during 42 days
of growth are independent of organic acid feed additives. *BMC Vet Res* **10**, 1–8 (2014).

11.   Fan, Y., Ju, T., Bhardwaj, T., Korver, D. R. & Willing, B. P. Week-Old Chicks with High
Bacteroides Abundance Have Increased Short-Chain Fatty Acids and Reduced Markers of
Gut Inflammation . *Microbiol Spectr* **11**, (2023).

12.   Zou, A., Sharif, S. & Parkinson, J. Lactobacillus elicits a ' Marmite effect ' on the chicken
cecal microbiome. *NPJ Biofilms Microbiomes* **4**, 1–5 (2018).

13.   Melkonian, C. *et al.* Microbial interactions shape cheese flavour formation. *Nat Commun*
14, (2023).

14.   Giordano, N. *et al.* Genome-scale community modelling reveals conserved metabolic cross-
feedings in epipelagic bacterioplankton communities. *Nat Commun* **15**, 2721 (2024).

15.   Zampieri, G., Campanaro, S., Angione, C. & Treu, L. Metatranscriptomics-guided genome-
scale metabolic modeling of microbial communities. *Cell Reports Methods* **3**, (2023).

16.   Quinn-Bohmann, N. *et al.* Microbial community-scale metabolic modelling predicts
personalized short-chain fatty acid production profiles in the human gut. *Nat Microbiol* **9**,
1700–1712 (2024).

17. Bauer, E. & Thiele, I. From metagenomic data to personalized in silico microbiotas : predicting dietary supplements for Crohn ' s disease. *NPJ Syst Biol Appl* **4**, (2018).

18. Marcelino, V. R. *et al.* Disease-specific loss of microbial cross-feeding interactions in the human gut. *Nat Commun* **14**, (2023).

19. Heinken, A., Hertel, J. & Thiele, I. Metabolic modelling reveals broad changes in gut microbial metabolism in inflammatory bowel disease patients with dysbiosis. *NPJ Syst Biol Appl* **7**, (2021).

20. Kumar, S. *et al.* Effect of antibiotic withdrawal in feed on chicken gut microbial dynamics , immunity , growth performance and prevalence of foodborne pathogens. *PLoS One* **13**, 1–23 (2018).

21. Caspi, R. *et al.* The MetaCyc database of metabolic pathways and enzymes and the BioCyc collection of pathway/genome databases. *Nucleic Acids Res* **44**, (2016).

22. Humbel, F. *et al.* Association of Alterations in Intestinal Microbiota With Impaired Psychological Function in Patients With Inflammatory Bowel Diseases in Remission. *Clinical Gastroenterology and Hepatology* **18**, (2020).

23. Bowerman, K. L. *et al.* Disease-associated gut microbiome and metabolome changes in patients with chronic obstructive pulmonary disease. *Nat Commun* **11**, (2020).

24. Crossfield, M. *et al.* Archaeal and Bacterial Metagenome-Assembled Genome Sequences Derived from Pig Feces. *Microbiol Resour Announc* **11**, (2022).

25. Gilroy, R. *et al.* Metagenomic investigation of the equine faecal microbiome reveals extensive taxonomic diversity. *PeerJ* **10**, (2022).

26. Zhang, X.-X. *et al.* A Catalog of over 5,000 Metagenome-Assembled Microbial Genomes from the Caprinae Gut Microbiota. *Microbiol Spectr* **10**, (2022).

27. Gilroy, R. *et al.* Extensive microbial diversity within the chicken gut microbiome revealed by metagenomics and culture. *PeerJ* **9**, (2021).

28. Nayfach, S., Shi, Z. J., Seshadri, R., Pollard, K. S. & Kyrpides, N. C. New insights from uncultivated genomes of the global human gut microbiome. *Nature* **568**, (2019).

29. Wang, Y. *et al.* Phylogenomics of expanding uncultured environmental Tenericutes provides insights into their pathogenicity and evolutionary relationship with Bacilli. *BMC Genomics* **21**, (2020).

30.   Bauer, E., Zimmermann, J., Baldini, F., Thiele, I. & Kaleta, C. BacArena: Individual-based
848   metabolic modeling of heterogeneous microbes in complex communities. *PLoS Comput*
849   *Biol* **13**, (2017).

850   31.   Salvi, P. S. & Cowles, R. A. Butyrate and the intestinal epithelium: Modulation of
851   proliferation and inflammation in homeostasis and disease. *Cells* vol. 10 Preprint at
852   https://doi.org/10.3390/cells10071775 (2021).

853   32.   Van Immerseel, F. *et al.* Interactions of butyric acid- and acetic acid-treated Salmonella
854   with chicken primary cecal epithelial cells in vitro. *Avian Dis* **48**, (2004).

855   33.   Rios-Covian, D., Gueimonde, M., Duncan, S. H., Flint, H. J. & De Los Reyes-Gavilan, C.
856   G. Enhanced butyrate formation by cross-feeding between Faecalibacterium prausnitzii and
857   Bifidobacterium adolescentis. *FEMS Microbiol Lett* **362**, (2015).

858   34.   Reichardt, N. *et al.* Phylogenetic distribution of three pathways for propionate production
859   within the human gut microbiota. *ISME Journal* **8**, (2014).

860   35.   Geirnaert, A. *et al.* Butyricicoccus pullicaecorum, a butyrate producer with probiotic
861   potential, is intrinsically tolerant to stomach and small intestine conditions. *Anaerobe* **30**,
862   (2014).

863   36.   Louis, P. & Flint, H. J. Diversity, metabolism and microbial ecology of butyrate-producing
864   bacteria from the human large intestine. *FEMS Microbiol Lett* **294**, (2009).

865   37.   Feng, J. *et al.* Polysaccharide utilization loci in Bacteroides determine population fitness
866   and community-level interactions. *Cell Host Microbe* **30**, (2022).

867   38.   Liu, H. *et al.* Ecological dynamics of the gut microbiome in response to dietary fiber. *ISME*
868   *Journal* **16**, (2022).

869   39.   Shin, J. H. *et al.* Bacteroides and related species: The keystone taxa of the human gut
870   microbiota. *Anaerobe* vol. 85 Preprint at https://doi.org/10.1016/j.anaerobe.2024.102819
871   (2024).

872   40.   Weiss, A. S. *et al.* Nutritional and host environments determine community ecology and
873   keystone species in a synthetic gut bacterial community. *Nat Commun* **14**, (2023).

874   41.   Zelezniak, A. *et al.* Metabolic dependencies drive species co-occurrence in diverse
875   microbial communities. *Proc Natl Acad Sci U S A* **112**, (2015).

876   42.   Giri, S. *et al.* Metabolic dissimilarity determines the establishment of cross-feeding
877   interactions in bacteria. *Current Biology* **31**, (2021).

43.   Krautkramer, K. A., Fan, J. & Bäckhed, F. Gut microbial metabolites as multi-kingdom
intermediates. *Nature Reviews Microbiology* vol. 19 Preprint at
https://doi.org/10.1038/s41579-020-0438-4 (2021).

44.   Dabrock, B., Riedel, J., Bertram, J. & Gottschalk, G. Isopropylbenzene (cumene) - a new
substrate for the isolation of trichloroethene-degrading bacteria. *Arch Microbiol* **158**,
(1992).

45.   Liu, C. J. *et al.* Complete genome sequences and comparative genome analysis of
Lactobacillus plantarum strain 5-2 isolated from fermented soybean. *Genomics* **106**, (2015).

46.   Zhou, Y. & Zhi, F. Lower Level of Bacteroides in the Gut Microbiota Is Associated with
Inflammatory Bowel Disease: A Meta-Analysis. *BioMed Research International* vol. 2016
Preprint at https://doi.org/10.1155/2016/5828959 (2016).

47.   Vich Vila, A. *et al.* Gut microbiota composition and functional changes in inflammatory
bowel disease and irritable bowel syndrome. *Sci Transl Med* **10**, (2018).

48.   Xu, X. *et al.* The gut metagenomics and metabolomics signature in patients with
inflammatory bowel disease. *Gut Pathog* **14**, (2022).

49.   Lloyd-Price, J. *et al.* Multi-omics of the gut microbial ecosystem in inflammatory bowel
diseases. *Nature* **569**, (2019).

50.   Knights, D. *et al.* Complex host genetics influence the microbiome in inflammatory bowel
disease. *Genome Med* **6**, (2014).

51.   Pascal, V. *et al.* A microbial signature for Crohn's disease. *Gut* **66**, (2017).

52.   Willing, B. P. *et al.* A pyrosequencing study in twins shows that gastrointestinal microbial
profiles vary with inflammatory bowel disease phenotypes. *Gastroenterology* **139**, (2010).

53.   Franzosa, E. A. *et al.* Gut microbiome structure and metabolic activity in inflammatory
bowel disease. *Nat Microbiol* **4**, (2019).

54.   Hove, H. & Mortensen, P. B. Influence of intestinal inflammation (IBD) and small and large
bowel length on fecal short-chain fatty acids and lactate. *Dig Dis Sci* **40**, (1995).

55.   Roediger, W. E. W. THE COLONIC EPITHELIUM IN ULCERATIVE COLITIS: AN
ENERGY-DEFICIENCY DISEASE? *The Lancet* **316**, (1980).

56.   Kameyama, J. I., Narui, H., Inui, M. & Sato, T. Energy Level in Large Intestinal Mucosa in
Patients with Ulcerative Colitis. *Tohoku J Exp Med* **143**, (1984).

57.   Bjerrum, J. T. *et al.* Metabonomics of human fecal extracts characterize ulcerative colitis,
909   Crohn's disease and healthy individuals. *Metabolomics* **11**, (2015).

910   58.   Xu, H. M. *et al.* Characterization of short-chain fatty acids in patients with ulcerative colitis:
911   a meta-analysis. *BMC Gastroenterol* **22**, (2022).

912   59.   Braun, T. *et al.* Diet-omics in the Study of Urban and Rural Crohn disease Evolution
913   (SOURCE) cohort. *Nat Commun* **15**, (2024).

914   60.   Morgan, X. C. *et al.* Dysfunction of the intestinal microbiome in inflammatory bowel
915   disease and treatment. *Genome Biol* **13**, (2012).

916   61.   Lewis, J. D. *et al.* Inflammation, Antibiotics, and Diet as Environmental Stressors of the
917   Gut Microbiome in Pediatric Crohn's Disease. *Cell Host Microbe* **18**, (2015).

918   62.   Ellestad-Sayed, J. J., Nelson, R. A., Adson, M. A., Palmer, W. M. & Soule, E. H.
919   Pantothenic acid, coenzyme A, and human chronic ulcerative and granulomatous colitis.
920   *American Journal of Clinical Nutrition* **29**, (1976).

921   63.   Veseli, I. *et al.* Microbes with higher metabolic independence are enriched in human gut
922   microbiomes under stress. *bioRxiv* (2023) doi:10.1101/2023.05.10.540289.

923   64.   Bushman, F. D. *et al.* Multi-omic Analysis of the Interaction between Clostridioides
924   difficile Infection and Pediatric Inflammatory Bowel Disease. *Cell Host Microbe* **28**,
925   (2020).

926   65.   Filippi, J., Al-Jaouni, R., Wiroth, J. B., Hébuterne, X. & Schneider, S. M. Nutritional
927   deficiencies in patients with Crohn's disease in remission. *Inflamm Bowel Dis* **12**, (2006).

928   66.   Huang, Y. L., Chassard, C., Hausmann, M., Von Itzstein, M. & Hennet, T. Sialic acid
929   catabolism drives intestinal inflammation and microbial dysbiosis in mice. *Nat Commun* **6**,
930   (2015).

931   67.   Liang, Q. *et al.* Sialic acid plays a pivotal role in licensing Citrobacter rodentium's transition
932   from the intestinal lumen to a mucosal adherent niche. *Proc Natl Acad Sci U S A* **120**,
933   (2023).

934   68.   Xun, W. *et al.* Specialized metabolic functions of keystone taxa sustain soil microbiome
935   stability. *Microbiome* **9**, (2021).

936   69.   Ze, X., Duncan, S. H., Louis, P. & Flint, H. J. Ruminococcus bromii is a keystone species
937   for the degradation of resistant starch in the human colon. *ISME Journal* **6**, (2012).

70. Trosvik, P. & de Muinck, E. J. Ecology of bacteria in the human gastrointestinal tract--
identification of keystone and foundation taxa. *Microbiome* **3**, (2015).

71. Cartmell, A. *et al.* A surface endogalactanase in Bacteroides thetaiotaomicron confers
keystone status for arabinogalactan degradation. *Nat Microbiol* **3**, (2018).

72. Fisher, C. K. & Mehta, P. Identifying keystone species in the human gut microbiome from
metagenomic timeseries using sparse linear regression. *PLoS One* **9**, (2014).

73. Rychlik, I. Composition and function of chicken gut microbiota. *Animals* **10**, (2020).

74. Mancabelli, L. *et al.* Insights into the biodiversity of the gut microbiota of broiler chickens.
Environ Microbiol **18**, (2016).

75. Marcolla, C. S., Ju, T., Lantz, H. L. & Willing, B. P. Investigating the cecal microbiota of
broilers raised in extensive and intensive production systems. *Microbiol Spectr* **11**, (2023).

76. Roberts, T. *et al.* New issues and science in broiler chicken intestinal health: Intestinal
microbial composition, shifts, and impacts. *Worlds Poult Sci J* **71**, (2015).

77. Huang, Y. *et al.* Community composition of cecal microbiota in commercial yellow broilers
with high and low feed efficiencies. *Poult Sci* **100**, (2021).

78. Chen, Y. *et al.* Chicken cecal microbiota reduces abdominal fat deposition by regulating fat
metabolism. *NPJ Biofilms Microbiomes* **9**, (2023).

79. Lo Sasso, G. *et al.* Inflammatory Bowel Disease-Associated Changes in the Gut: Focus on
Kazan Patients. *Inflamm Bowel Dis* **27**, (2021).

80. Zheng, L. & Wen, X. L. Gut microbiota and inflammatory bowel disease: The current status
and perspectives. *World J Clin Cases* **9**, (2021).

81. Midani, F. S. & David, L. A. Tracking defined microbial communities by multicolor flow
cytometry reveals tradeoffs between productivity and diversity. *Front Microbiol* **13**, (2023).

82. Goodman, A. L. *et al.* Identifying Genetic Determinants Needed to Establish a Human Gut
Symbiont in Its Habitat. *Cell Host Microbe* **6**, (2009).

83. Ray, K. J., Santee, C., McCauley, K., Panzer, A. R. & Lynch, S. V. Gut Bifidobacteria
enrichment following oral Lactobacillus-supplementation is associated with clinical
improvements in children with cystic fibrosis. *BMC Pulm Med* **22**, (2022).

84. Ma, C. *et al.* The gut microbiome stability is altered by probiotic ingestion and improved
by the continuous supplementation of galactooligosaccharide. *Gut Microbes* (2020)
doi:10.1080/19490976.2020.1785252.

969 85. Armougom, F., Henry, M., Vialettes, B., Raccah, D. & Raoult, D. Monitoring bacterial
970   community of human gut microbiota reveals an increase in Lactobacillus in obese patients
971   and Methanogens in anorexic patients. *PLoS One* **4**, (2009).

972 86. Nieuwdorp, M., Gilijamse, P. W., Pai, N. & Kaplan, L. M. Role of the microbiome in energy
973   regulation and metabolism. *Gastroenterology* **146**, (2014).

974 87. Fan, Y. & Pedersen, O. Gut microbiota in human metabolic health and disease. *Nature*
975   *Reviews Microbiology* vol. 19 Preprint at https://doi.org/10.1038/s41579-020-0433-9
976   (2021).

977 88. Diener, C. & Gibbons, S. M. More is Different: Metabolic Modeling of Diverse Microbial
978   Communities. *mSystems* **8**, (2023).

979 89. Bosi, E., Bacci, G., Mengoni, A. & Fondi, M. Perspectives and Challenges in Microbial
980   Communities Metabolic Modeling. **8**, 1–9 (2017).

981 90. Bushnell, B. BBTools. *DOE Joint Genome Institute* Preprint at (2014).

982 91. Li, D., Liu, C. M., Luo, R., Sadakane, K. & Lam, T. W. MEGAHIT: An ultra-fast single-
983   node solution for large and complex metagenomics assembly via succinct de Bruijn graph.
984   *Bioinformatics* **31**, (2015).

985 92. Langmead. Bowtie2. *Nat Methods* **9**, (2013).

986 93. Kang, D. D. *et al.* MetaBAT 2: An adaptive binning algorithm for robust and efficient
987   genome reconstruction from metagenome assemblies. *PeerJ* **2019**, (2019).

988 94. Alneberg, J. *et al.* Binning metagenomic contigs by coverage and composition. *Nat Methods*
989   **11**, (2014).

990 95. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**,
991   (2009).

992 96. Uritskiy, G. V., DiRuggiero, J. & Taylor, J. MetaWRAP—a flexible pipeline for genome-
993   resolved metagenomic data analysis. *Microbiome* **6**, (2018).

994 97. Parks, D. H., Imelfort, M., Skennerton, C. T., Hugenholtz, P. & Tyson, G. W. CheckM:
995   Assessing the quality of microbial genomes recovered from isolates, single cells, and
996   metagenomes. *Genome Res* **25**, (2015).

997 98. Chaumeil, P. A., Mussig, A. J., Hugenholtz, P. & Parks, D. H. GTDB-Tk v2: memory
998   friendly classification with the genome taxonomy database. *Bioinformatics* **38**, (2022).

99.   Jain, C., Rodriguez-R, L. M., Phillippy, A. M., Konstantinidis, K. T. & Aluru, S. High
throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat*
Commun **9**, (2018).

100.   Abram, K. *et al.* Mash-based analyses of Escherichia coli genomes reveal 14 distinct
phylogroups. *Commun Biol* **4**, (2021).

101.   Zimmermann, J., Kaleta, C. & Waschina, S. gapseq: informed prediction of bacterial
metabolic pathways and reconstruction of accurate metabolic models. *Genome Biol* **22**,
(2021).

102.   Zou, A. *et al.* Systematic profiling of the chicken gut microbiome reveals dietary
supplementation with antibiotics alters expression of multiple microbial pathways with
minimal impact on community structure. *Microbiome* **10**, (2022).

103.   Noronha, A. *et al.* The Virtual Metabolic Human database : integrating human and gut
microbiome metabolism with nutrition and disease. *Nucleic Acids Res* **47**, 614–624 (2018).

104.   www.foodb.ca.

105.   AFZ. INRA-CIRAD-AFZ feed tables: composition and nutritive values of feeds for cattle,
sheep, goats, pigs, poultry, rabbits, horses, and salmonids. Feedtables.com, INRAE,
CIRAD, AFZ. (2017).

106.   Tejeda, O. J. & Kim, W. K. Role of dietary fiber in poultry nutrition. *Animals* vol. 11
Preprint at https://doi.org/10.3390/ani11020461 (2021).

107.   Erik, K. & Knudsen, B. Fiber and nonstarch polysaccharide content and variation in
common crops used in broiler diets 1. 2380–2393 (2012).

108.   Riesenfeld, G., Sklan, D., Bar, A., Eisner, U. & Hurwitz, S. Glucose absorption and starch
digestion in the intestine of the chicken. *Journal of Nutrition* **110**, 117–121 (1980).

109.   Khalil, M. M., Abdollahi, M. R., Zaefarian, F. & Ravindran, V. Measurement of ileal
endogenous energy losses and true ileal digestible energy of cereal grains for broiler
chickens. *Poult Sci* **99**, (2020).

110.   Bielorai, R. & Iosif, B. Amino acid absorption and endogenous amino acids in the lower
ileum and excreta of chicks. *Journal of Nutrition* **117**, 1459–1462 (1987).

111.   Bielorai, R., Tamir, M. & Hurwitz, S. Amino acid absorption along the intestinal tract of
chicks fed heated and raw soybean meal. *Journal of Nutrition* **107**, 1775–1778 (1977).

112.   Tancharoenrat, P., Ravindran, V., Zaefarian, F. & Ravindran, G. Digestion of fat and fatty
1030   acids along the gastrointestinal tract of broiler chickens. *Poult Sci* **93**, 371–379 (2014).

1031   113.   Rodriguez-Sanchez, R., Tres, A., Sala, R., Guardiola, F. & Barroeta, A. C. Evolution of
1032   lipid classes and fatty acid digestibility along the gastrointestinal tract of broiler chickens
1033   fed different fat sources at different ages. *Poult Sci* **98**, 1341–1353 (2019).

1034   114.   Zheng, J. *et al.* DbCAN3: Automated carbohydrate-Active enzyme and substrate
1035   annotation. *Nucleic Acids Res* **51**, (2023).

1036   115.   Forgie, A. J. *et al.* Pea polyphenolics and hydrolysis processing alter microbial community
1037   structure and early pathogen colonization in mice. *Journal of Nutritional Biochemistry* **67**,
1038   (2019).

1039   116.   Duke, G. E., Eccleston, E., Louis, C. F. & Bedbury, H. P. Cellulose digestion by domestic
1040   turkeys fed low or high fiber diets. *Journal of Nutrition* **114**, 95–102 (1984).

1041   117.   Duke, G. E. Relationship of cecal and colonic motility to diet, habitat, and cecal anatomy
1042   in several avian species. *Journal of Experimental Zoology* **252**, 38–47 (1989).

1043   118.   Ravindran, V. Feed enzymes: The science, practice, and metabolic realities. *Journal of
1044   Applied Poultry Research* **22**, 628–636 (2013).

1045   119.   Nguyen, H. T., Bedford, M. R., Wu, S. B. & Morgan, N. K. Soluble non-starch
1046   polysaccharide modulates broiler gastrointestinal tract environment. *Poult Sci* **100**, (2021).

1047   120.   Su, G., Morris, J. H., Demchak, B. & Bader, G. D. Biological Network Exploration with
1048   Cytoscape 3. *Curr Protoc Bioinformatics* **2014**, (2014).

1049

1050

1051
1052

1053

1054

1055

1056