Dominant contribution of Asgard archaea to eukaryogenesis

Victor Tobiasson[1*], Jacob Luo[1], Yuri I Wolf[1], Eugene V Koonin[1*]

1. Computational Biology Branch, Division of Intramural Research, National Library of Medicine, National Institutes of Health, Bethesda, MD 20894, USA

*Correspondence to: victor.tobiasson@nih.gov or koonin@ncbi.nlm.nih.gov

Abstract

The origin of eukaryotes is one of the key problems in evolutionary biology [1,2]. The
demonstration that the Last Eukaryotic Common Ancestor (LECA) already contained the
mitochondrion, an endosymbiotic organelle derived from an alphaproteobacterium, and the
discovery of Asgard archaea, the closest archaeal relatives of eukaryotes [3-7], inform and
constrain evolutionary scenarios of eukaryogenesis [8]. We undertook a comprehensive
analysis of the origins of core eukaryotic genes tracing to the LECA within a rigorous statistical
framework centered around evolutionary hypotheses testing using constrained phylogenetic
trees. The results reveal dominant contributions of Asgard archaea to the origin of most of
the conserved eukaryotic functional systems and pathways. A limited contribution from
Alphaproteobacteria was identified, primarily relating to the energy transformation systems
and Fe-S cluster biogenesis, whereas ancestry from other bacterial phyla was scattered across
the eukaryotic functional landscape, without clear consistent trends. These findings suggest a
model of eukaryogenesis in which key features of eukaryotic cell organization evolved in the
Asgard lineage leading to the LECA, followed by the capture of the Alphaproteobacterial
endosymbiont, and augmented by numerous but sporadic horizontal acquisition of genes
from other bacteria both before and after endosymbiosis.

Main

29
Eukaryotes drastically differ from archaea and bacteria (collectively, prokaryotes) by the highly complex
organization of the eukaryotic cells. The signature features of this organizational complexity include the
eponymous nucleus, the endomembrane system, the elaborate cytoskeleton, and the energy-converting
mitochondrion which evolved from an Alphaproteobacterial endosymbiont [9]. Early models of
eukaryogenesis featured a protoeukaryote lacking mitochondria that captured and domesticated an
alphaproteobacterium at a relatively late stage of evolution [10,11]. However, subsequent research
revealed no primary amitochondrial eukaryotes (although many have lost mitochondria through
reductive evolution) [12], suggesting that the Last Eukaryotic Common Ancestor (LECA) was already
endowed with mitochondria along with the other signatures of the eukaryotic cellular organization [1,2,8].
These findings lend credence to scenarios of eukaryogenesis in which mitochondrial endosymbiosis
triggered cellular reorganization, giving rise to the eukaryotic cellular complexity [1,2,8,13-15].

41
Phylogenomic analyses show that those conserved, core genes of eukaryotes that can be traced to the
LECA are a mix originating from archaea and various bacteria, and in early studies, the bacterial
contributions quantitatively exceeded the archaeal ones. The archaea-derived genes were found
primarily in information processing systems (replication, transcription and translation) whereas genes of
apparent bacterial origin comprised the operational component of the eukaryotic gene complement, in
particular encoding metabolic enzymes [16-19].

The study of eukaryogenesis was transformed by the discovery and exploration of Asgard archaea

(currently, phylum *Promethearchaeota* within kingdom *Promethearchaeati* [20, 21], hereafter Asgard)

which includes the closest known archaeal relatives of eukaryotes [3-7]. Phylogenetic analyses of highly

conserved genes have been pushing the eukaryotic branch progressively deeper within the Asgard tree.

The latest such analysis identified the order *Hodarchaeales*, within the Asgard class *Heimdallarchaeia*, as

the likely sister group of eukaryotes [4]. However, with the continued influx of Asgard sequence data, the

exact affinity of eukaryotes within Asgard remains a moving target [22,23]. Notably, it has been shown that

Asgard archaea encode, express and utilize homologs of a broad variety of eukaryote signature proteins

beyond the core information processing componentry; in particular, cytoskeletal proteins and proteins

involved in membrane remodeling [4,5,7,24,25]. Furthermore, these genes of Asgard archaea have undergone

extensive duplication and subfunctionalization pre-LECA, further highlighting their prominent role in

eukaryogenesis [26].

A variety of models for eukaryogenesis have been proposed, differing with respect to the timing,

topology and origin of the eukaryotic cellular organization and genetic repertoire [1,2,8]. The

endosymbiotic origin of the mitochondria from an alphaproteobacterium is indisputable, but the nature

of the host of the proto-mitochondrial endosymbiont remains a matter of debate. The most

straightforward models posit an Asgard host [2,8,27]. However, a major conundrum for such scenarios of

eukaryogenesis is the chemistry of membrane lipids and the enzymology of their biosynthesis, which are

unrelated in archaea and bacteria, with eukaryotes utilizing the bacterial type [28]. Accordingly, any model

of eukaryogenesis that includes an archaeal host would require a change in membrane composition.

More complex, alternative models of eukaryogenesis, underpinned by metabolic symbiosis, postulate

two endosymbiotic events whereby an Asgard archaeon was first engulfed by a bacterium, the archaeal

membrane being lost in the process, and then, a second endosymbiosis gave rise to the mitochondria

[15,29,30]. Even if seemingly less parsimonious than binary scenarios, such models of eukaryogenesis

account for the continuity of bacterial membranes (while requiring retargeting at least part of the

Archaeal membrane proteome) and appear compatible with the syntrophic lifestyle of at least some

Asgard archaea, found in consortia with bacteria, Myxococcota in particular (formerly

Deltaproteobacteria)[31,32]. Furthermore, a pivotal study by Pittis and Gabaldon [35] in which relative timing

of events in eukaryogenesis was inferred by comparing stem length from different ancestors in

phylogenetic trees of conserved eukaryotic genes, suggested a late capture of the mitochondrion,

apparently preceded by acquisition of many genes from different bacteria, possibly through one or

multiple earlier endosymbioses [33-35].

Here, we took advantage of the rapidly growing collection of archaeal, bacterial and eukaryotic genome

sequences to assess the origins of core eukaryotic genes within a rigorous statistical framework based

on evolutionary hypotheses testing using constrained phylogenetic trees. The results reveal a consistent,

major link between Asgard archaea and the origin of most functional classes of eukaryotic genes,

demonstrating their dominant contribution to eukaryogenesis. These findings are compatible with a

scenario of eukaryogenesis where many signature, complex features of eukaryotic cells evolved in the

Asgard ancestor of eukaryotes.

Phylogenomic analysis and global evolutionary associations of conserved eukaryotic proteins

To represent prokaryotes, a database of 75 million prokaryote protein sequences was curated from

47,545 complete prokaryotic genomes obtained from the NCBI GenBank in November 2023 (prok2311)

and supplemented with proteins extracted from 63 Asgard genome assemblies [4,5]. The initial eukaryotic

database consisted of 30 million sequences from 993 species present in EukProt v3 [73], cleaned using

mmseqs2 [75] to remove prokaryotic contaminants (Figure 1A). In order to accurately infer the origins of

genes in the LECA, it is essential to exclude genes present only within narrow subsets of species. These

possibly result from post-LECA horizontal gene transfer (HGT) between eukaryotes and prokaryotes, a

major phenomenon given the pervasive cohabitation of eukaryotes with prokaryotes [36-39]. To prevent

such genes from affecting our analysis, we reconstructed the "soft-core" pangenome for each of our

assigned prokaryotic and eukaryotic taxonomic classes (see Methods). These pangenomes included only

those genes that are present in at least 50% of the species within each class of bacteria and archaea in

prok2311 (20.7 million), and at least 20% of the species in eukaryotes (12.7 million), ensuring that our

inferences were based only on widespread protein families and would not reflect lineage-specific HGT

(supplementary Figure 1D).

To identify close connections between prokaryotic and eukaryotic protein families, separate hidden

Markov model (HMM) databases for prokaryotes and eukaryotes were constructed using a custom,

cascaded, sequence-to-profile clustering pipeline, implemented based on mmseqs2 clustering, followed

by a multistep data-reduction and multiple sequence alignment (MSA) procedure to generate HMM

profiles using diversified muscle5 alignments [43] and HH-suite [40] (see Methods). The resulting eukaryotic

HMMs were queried against the prokaryotic dataset using HHblits [40] to identify sets of homologous

protein sequences. Each eukaryotic cluster sequence and all its significant prokaryotic hits constituted

an individual set, hereinafter referred to as a Eukaryotic-Prokaryotic Orthologous Cluster (EPOC). Each

EPOC contains a unique set of eukaryotic proteins. These EPOCs were used for phylogenetic tree

construction, annotation, and evolutionary hypothesis testing (Figure 1A). The final EPOCs include 10.9

million prokaryotic and 1.74 million eukaryotic sequences, mapping to 52% and 14% of the respective

non-redundant pangenome datasets.

To infer the most likely prokaryotic ancestry of the eukaryotic proteins given the data in each EPOC,

rather than relying on tree topology directly, we employed a probabilistic approach for evolutionary

hypothesis testing using constraint trees (Figure 1A). Following the construction of an initial master tree,

we carried out further constrained tree calculations, exhaustively sampling all arrangements of the

prokaryotic sister clades nearest to the eukaryotic clade(s) and obtaining Expected Likelihood Weights

(ELW) for the set of possible sister clade models (supplementary Figure 2, supplementary Data 1) [41].

Given that the ELW metric is analogous to model selection confidence, here we take it to be

proportional to the probability that a sampled prokaryotic clade is the best estimate of the closest sister

clade of a eukaryotic clade. For each EPOC, our analysis dynamically accounts for long branch outliers

and is capable of resolving eukaryotic paraphyly, treating each resolved eukaryotic clade within a EPOC

individually for downstream analysis (see Methods). The resulting data included 14,400 EPOCs

annotated using profiles generated from KEGG Orthology Groups (KOGs) [42], each with an MSA
generated using muscle5 [43], a maximum likelihood tree inferred using IQtree2 [44] and associated ELW
values for all candidate prokaryotic sister clades. The analysis of prokaryotic ancestry was performed
only for those eukaryotic clades that included more than 5 distinct taxonomic labels, with at least one
coming from Amorphea and one from Diaphoretickes, the two expansive eukaryotic clades considered
to emit from either the first or the second bifurcation in the evolution of eukaryotes [45,46]. Thus, although
we did not attempt to explicitly reconstruct the LECA gene set, these clades represent genes which,
given their broad taxonomic distribution in our data, were likely to have had an ancestor in the LECA.
139
Considering the global distribution of ELW values across all 13,500 EPOCs covering 3950 unique KOGs,
the single greatest average ELW (aELW), here referred to as "association", was with Asgard archaea.
Further associations with Cyanobacteria, Actinomycetota, Betaproteobacteria and Alphaproteobacteria,
as well as trace association with many additional bacterial classes, were detected at lower levels (Figure
1B). Due to the scope of the analysis, MSA quality and topological diversity of the trees varied across the
EPOCs such that some EPOCs showed low ELW values for all clades, indicating that the data in these
cases was insufficient to discern their eukaryotic ancestries. Excluding such low-quality EPOCs with a
maximum ELW < 0.4 yielded a robust core set of 4290 EPOCs covering 2100 KOGs, improving the
reliability of the results. This core set covers a wide range of information processing systems, ubiquitous
metabolic pathways, and transporters, as well as regulatory and housekeeping proteins. Limiting the
analysis to this core subset of well assigned eukaryotic families with wide taxonomic coverage increased
the global Asgard association, which now accounted for 50% of all likelihood weight across more than
4,000 unique data points, covering more than 2,000 unique KOGs (Figure 1B).
153
To ensure these results were robust to the choice of the clustering approach, we performed additional
cluster annealing steps (see Methods, supplementary Figure 8A). In brief, starting with the mmseqs2
clusters, we performed HMM-HMM searches and clustered the results with a greedy set cover
approach. We then further partitioned these superclusters by constructing per-cluster trees and
identifying sets of leaves with mutually short pairwise distances, indicating the existence of separate
clades. These annealed clades, now strictly defined to be monophyletic, given significant sequence
similarity, were then used for EPOC construction. Additionally, we reconstructed our full dataset using
the Genome Taxonomy Database (GTDB) taxonomy to verify that the detected eukaryote-prokaryote
associations were not substantially affected by our use of the NCBI taxonomy [47]. We note that, while
relative contributions from specific taxa varied due to remapping of taxonomic labels (for example,
merging of Betaproteobacteria into Gammaproteobacteria), or due to the stricter annealed EPOC
definition limiting the scope of the data; our key observation of a dominant Asgard association for
eukaryotes held independent of clustering and taxonomy (supplementary Figure 1E, 8B, 8C). Given the
robustness of the results to both cluster specification and taxonomy, all subsequent analysis was based
on the core EPOC set described above. Finally, although our results suggest the presence of diverse
paralogs of apparent Asgard origin in the LECA [5,26,48], paralogy cannot account for this wide contribution
of Asgard to eukaryogenesis (supplementary Figure 1B, supplementary Figure 3A).
171

Averaging ELW scores across EPOCs sharing KEGG ontology supported strong Asgard association with,
among other functional systems and pathways, the ribosome, RNA and DNA polymerases, Ras-like
GTPases, as well as ubiquitin-mediated protein degradation and the proteasome. These findings
validated and expanded upon previous observations [2,4,5], extending Asgard associations to large parts of
the core metabolic network (Figure 1C). In contrast, prominent Alphaproteobacterial associations were
relatively limited, primarily including proteins involved in oxidative phosphorylation and Fe-S clusters
biogenesis, with weaker associations across the mitochondrial metabolic pathways. Thus, we observed a
far stronger overall association between Asgard archaea and eukaryotes across diverse biological
functions and pathways than previously described [4,5] although the association between eukaryotic core
metabolism and a wide range of bacterial phyla remained cumulatively substantial. We also observed a
consistent association with Cyanobacteria, a known endosymbiont of Archaeplastida and the ancestor of
plastids, with a strong impact on the evolution of Eukarya post LECA. However, we note that pathways
driving this Cyanobacterial association are unrelated to the photosynthetic capabilities of the
chloroplast, but rather exhibit a broad association with multiple metabolic pathways, without apparent
trends, similar to the observations for other bacterial taxa (supplementary Figure 4)[49-51]. Thus, we
conclude that this Cyanobacterial signal most likely stems from events pre-LECA and is distinct from the
later contributions from the chloroplast.
189

190
Figure 1. Phylogenetic associations of functional classes of eukaryotic proteins with major archaeal
and bacterial taxa.

A) Simplified EPOC generation and processing pipeline. From left to right, one eukaryotic and one
prokaryotic sequence database transformed into HMMs. The query of eukaryotic against prokaryotic
HMMs generated 13,500 annotated EPOCs. The detected prokaryotic clades were assessed for their
probability to be the closest eukaryotic sister phyla by permutation testing using IQTree2 (see
supplementary Figure 2). **B)** Global aELW for all EPOCs within unfiltered, core and non-paralogous
subsets and top-level breakdowns for metabolic enzymes and proteins of core information processing
pathways as defined by the KEGG ontology BRITE at the coarsest A level (KO:09100, KO:09120). Grey
markers group prokaryotic taxa with an aELW < 0.01. **C)** Further subdivision of aELW values across more
specific functional categories of genes BRITE level B, counting the total number of EPOCs and colored by
relative aELW of taxa per category.
203

Broad Asgard contributions to eukaryogenesis

In accord with a dominant contribution of Asgard archaea to eukaryogenesis, we observed consistent
and major associations of Asgard proteins with a wide array of cellular functions. In previous work, the
strongest Asgard traces have been noted across the information processing systems, with unambiguous
protein associations with DNA and RNA processing as well as protein expression, trafficking and
signaling [2,4,5]. Here, we confirmed and extended Asgard associations with genome replication and
transcription, and further detected pronounced Asgard traces for nucleotide excision repair, mismatch
repair and homologous recombination. Additional well-known associations such as ribosomal proteins
were expanded upon to include translation factors, components of the co-translational insertion
machinery, protein targeting and aminoacyl-tRNA biosynthesis (Figure 2A, supplementary data 1). Thus,
all groups of core eukaryotic proteins involved in information processing appear to be almost exclusively
of Asgard descent (Figure 1C).
216

We further detected Asgard associations extending far beyond the information processing systems,
including prominent contributions of Asgard archaea to the protein machinery involved in
nucleocytoplasmic transport as well as downstream protein sorting, glycosylation and targeting. Central
components of the ER-associated, N-linked glycan biosynthesis and transfer, including both cytoplasmic
and lumenal monoglycosyltransferases, as well as the core of the oligosaccharyltransferase complex
(OSTC) were strongly associated with Asgard (Figure 2A). Enzymes associated with glycosylation
maturation in the Golgi complex were not recovered in our core data, possibly, due to extensive
diversification of domain architecture in eukaryotes, limiting their taxonomic scope and apparent
presence in the LECA. The Asgard connections of the eukaryotic glycosylation machinery further
included the synthesis of GPI-anchors, which post-translationally tether targeted proteins to the
membrane [52], here detected as unambiguously Asgard-derived. We also detected an Asgard origin of the
7-subunit (UDP-GlcNAc)-transferring (GPI-GnT)-monoglycosyltransferase complex responsible for
initiating GPI-anchor synthesis, components required for the maturation of the GPI-anchor, as well as
the transamidase complex and factors responsible for transfer of the protein onto the mature GPI-
anchor (Figure 2B).
232

Of major importance to eukaryogenesis is the provenance of the pathways for the biosynthesis of bacterial-type lipids, given that all binary symbiogenesis scenarios with a Archaeal host require a transition from archaeal to bacterial lipids in the membranes [8]. Although Asgard associations were observed for large parts of the overall metabolic network, the pathways associated with fatty acid synthesis and decay showed a wide distribution of aELW values, suggesting a higher than average degree of mosaicism. As we enforce a minimum ELW of 0.4 for the most likely ancestors in our core data, such mosaicism is interpreted as separate, individually likely ancestries, averaged across EPOCs within a pathway, rather than a general uncertainty of the constituent EPOCs. The global aELW values favored Asgard origin for these pathways, but there were also associations with Actinomycetota and Alphaproteobacteria, further discussed below. However, higher Asgard aELW was observed within the adjoining ER localized pathways for sphingolipid metabolism, a broad class of derived plasma membrane lipids in eukaryotes. Previous studies have highlighted possible convergent origins of this pathway in bacteria and eukaryotes [53,54], but here we detected a global association with Asgard archaea (Figure 2A). Of further note are the ER-associated mevalonate pathway, converting Acetyl-CoA to mevalonate and further to Farnesyl and Geranyl diphosphate, which we also found to be associated with Asgard (Figure 2B). In eukaryotes, mevalonate is required for isoprenoid synthesis, in turn forming the precursor units for both sterols, carotenoids and terpenoids, synthesized in the ER lumen via the mevalonate pathway. However, in Archaea, isoprenoids are the precursors for the ether-linked membrane lipids [55]. In conclusion, we detected Asgard associations across a wide range of cellular functions and metabolic pathways and noted a relationship between Archaeal lipid synthesis and eukaryotic steroid precursor metabolism. A distinctly weaker Asgard signal was detected for pathways involved in bacterial lipid biosynthesis, suggesting a complex evolutionary history.

Figure 2. Asgard origins of eukaryotic pathways. A) Top 15 KEGG pathways with strongest Asgard aELW. KEGG maps with more than 20 EPOCs included. Taxonomy colored as per Figure 1. **B)** Further notable

259 Asgard systems, aELW calculated over either KEGG "map" number (ex. map0000X), reaction number
260 (R0000X), or module number (ex. M0000X).
261
Limited and specific Alphaproteobacterial contributions

In line with the central role of mitochondria in eukaryotic energy metabolism, and as recently reported
for central carbon metabolism [56], we primarily observe clear associations between Alphaproteobacteria
and mitochondrially localized metabolic pathways. Apart from the components of the mitochondrial
translation system, the most prominent Alphaproteobacterial associations were evident within
complexes involved in oxidative phosphorylation and the associated ubiquinone synthesis
(supplementary Figure 3B). Outside these central energy-transforming functions, we only detected
sparse contributions from core Alphaproteobacterial genes. One such prominent association was the
pathway for iron sulfur cluster (ISC) biogenesis. As previously reported [57], the ISC assembly machinery is
of Alphaproteobacterial origin, and accordingly, we found the 4Fe-4S ICSA platforms, as well as IBA57
and Fe-S cluster binding ferredoxin-1 and 2, to be associated with Alphaproteobacteria (Figure 3,
supplementary Figure 3B, supplementary Data 1). However, the 2Fe-2S precursor scaffold ISCU showed
a mosaic association with diverse prokaryotes (including Alphaproteobacteria), and a weaker but clearly
detectable Asgard association. Furthermore, the cysteine desulfurase NFS1 and the upstream pathways
for the biosynthesis of sulfur-containing amino acids, cysteine, and methionine, were Asgard associated.
The Alphaproteobacterial associations with Fe-S synthesis were therefore restricted primarily to
synthesis and integration of 4Fe-4S into the oxidative phosphorylation complexes. Outside the
mitochondria, Alphaproteobacterial associations with Fe-S metabolism cease. Here, ISC insertion
depends on the cytosolic targeting complex CIA, which consists of CIAO1, CIA2B and MMS19 [59]. For the
CIA components CIA1 and CIA2B, we observed clear association with Asgard whereas MMS19 is not
detected in our data. Taken together, these observations indicate that the contributions of
Alphaproteobacteria to eukaryogenesis are mostly centered around the mitochondrion.
284

286 **Figure 3. Ancestry of Eukaryotic Fe-S cluster synthesis pathways.** Schematics for Fe-S cluster synthesis
287 and assembly including downstream cytosolic Fe-S insertion are shown. Boxes indicate individual KOGs
288 covered by EPOCs. Full color indicates support for ancestry with an aELW > 0.5, boxes with mixed color

indicate unresolved ancestries with ELWs < 0.5 for the two most likely estimates. Mixed Prokaryotic ancestries represent all possible taxa, including Alphaproteobacteria.

Absence of functionally consistent contributions from other bacteria

Although our analysis greatly expands the Asgard contributions to eukaryogenesis, and reveals limited but consistent Alphaproteobacterial associations, contributions from other bacteria were detected as well. For some biological functions, this "diverse" bacterial component accounted for the majority of the aELW, and roughly one third of the analyzed KOGs (600 of 2,100) and EPOCs (1,580 of 4,300) had their strongest association neither with Asgard, nor with Alphaproteobacteria . However, in a sharp contrast to the limited but functionally related associations between Alphaproteobacterial and oxidative phosphorylation and Fe-S metabolism, EPOCs associated with diverse bacteria showed few functional trends, instead demonstrating low level association across most metabolic functions. Although Cyanobacteria and Actinomycetota exhibited stronger global aELW compared to Alphaproteobacteria, this association stemmed from multiple small contributions across disjoint pathways rather than specific functionally related contributions (supplementary Figure 4). To emphasize this trend, across the diverse set of bacterial taxa, only Alphaproteobacteria and Myxococcota were associated with a KEGG pathway containing more than 20 EPOCs and with a greater aELW than Asgard (Figure 4; see supplementary Figure 6A for alternate cutoffs). Specifically, oxidative phosphorylation shows a strong signal for Alphaproteobacteria (see supplementary Figure 3 for OXPHOS complexes) while Myxococcota (formerly, Deltaproteobacteria) showed a consistent associations with metabolic pathways involved in nicotinate, nucleotide, and nucleoside-sugar modification and metabolism. However, these nucleotide-related associations appear to be limited to phosphatases and phosphoribosyltransferases acting on nucleoside-sugars, specifically 5' and 3' nucleotidases and are as such much more limited in functional scope than pathways of Alphaproteobacterial associations. Previous studies highlighted a link between Myxococcota and steroid biosynthesis in Eukarya [60]. While we observed a clear, if narrow (6 EPOCs), association with Myxococcota for diterpenoid biosynthesis, downstream of the mevalonate pathway and upstream of steroid synthesis, our criteria on coverage and taxonomic scope limited the representation of downstream steroid pathways in the core data (supplementary Figure 5 and 6A).

Although few specific associations were observed with diverse bacteria on the level of pathways, the possibility remained that individual proteins with key functions could originate from such taxa. For this analysis, we considered a stricter subset of the core EPOCs, requiring a eukaryotic outgroup containing at least 15 taxonomic clades, to ensure functional centrality, and prokaryotic sister taxa with at least 20 sequences and an ELW > 0.7 to ensure strong association. Of the 130 unique EPOCs meeting these criteria, 33 were found to be associated with diverse bacterial lineages, mostly Actinomycetota and FCB group bacteria(supplementary Figure 6B). The remaining 97 EPOCs were found to be Asgard-derived. The 33 EPOCs of prominent bacterial origin covered a wide range of cellular functions from major facilitator superfamily transporters to lipases, to components of core sugar metabolism and cardiolipin synthesis. Once again, we observed no clear functional trend for these bacterial contributions. Thus, although single protein associations were individually significant, no pathways appeared to have been derived from any single bacterial taxon. Instead, we interpret these diffuse associations with diverse

bacteria as indications of multiple bacterial contributions, each with a limited functional scope, likely

originating from isolated HGT events rather than additional endosymbioses.

Figure 4. Strongest bacterial contributions to eukaryotic pathways compared to the contributions of

Asgard archaea. Each prokaryotic taxon is represented by the highest aELW of any KEGG BRITE pathway

with at least 20 EPOCs, relative to that of Asgard for the same pathway (background, pale blue). The

strongest prokaryotic association relative to Asgard was that of Myxococcota with Nicotinate

metabolism followed by Alphaproteobacteria with oxidative phosphorylation.

Ancestral stem lengths and the amount of evolution pre- and post-LECA

We attempted to use tree stem length analysis to infer the timing of acquisition of core eukaryotic

genes, as originally implemented by Pittis and Gabaldon [35], with the mitochondrion as a test case. As the

time elapsed post-LECA is the same for all genes, it has been proposed that the relative stem length in

gene trees is proportional to the time elapsed since the acquisition of the gene from its prokaryotic

donor to the LECA. Using this approach, Pittis and Gabaldon found that proteins of Alphaproteobacterial

descent had significantly shorter stems than proteins of archaeal descent, consistent with a

mitochondria-late scenario for eukaryogenesis [33-35]. The raw stem length is defined as the distance from

the Last Common Ancestor (LCA) of Eukarya and the most likely prokaryotic sister phylum, the First

Eukaryotic Common Ancestor (FECA)[48,61], to the LECA. To account for differences in evolutionary rates,

this raw stem length is normalized by dividing by the median eukaryotic branch length, measured from

the LECA (Figure 5A).

Across the entire set of eukaryotic stem lengths for our dataset (5,850 stems), we observed a wide

distribution with a sharp maximum close to 0.05, highly reminiscent of previous findings [35] (Figure 5B

5C). In our analysis, Alphaproteobacterial stems shorter than 0.3 (70% of all stems) were significantly
longer than those of Asgard origin, but Alphaproteobacterial stems longer than 0.35 were instead
shorter than Asgard on average, showing a non-obvious yet reproducible (supplementary figure 7B) and
statistically significant relationship (Figure 5B). Stems of Cyanobacterial origin followed a similar but
weaker trend. Thus, although the observation of globally shorter stem lengths for genes of
alphaproteobacterial descent applied only to stems longer than 0.35, we effectively recapitulate the
findings of Pittis and Gabaldon, who primarily analyzed stems within that range. When we limited our
analysis to ribosomal proteins and components of oxidative phosphorylation complexes, that is, genes
of well-known and easily identifiable Archaeal and Alphaproteobacterial ancestries, respectively, the
above trends were further accentuated (Figure 5C). Contrary to the notion of stem lengths as a measure
of the acquisition timing, we observed variances of more than two orders of magnitude in normalized
stem lengths for the oxidative phosphorylation complexes that are assumed to have been acquired
simultaneously during mitochondrial symbiogenesis (Figure 5B). Taken together, these findings suggest
that even normalized stem lengths are determined, primarily, by factors other than the gene acquisition
time, at least, across the long evolutionary timescales analyzed here.
370
Comparison of the stem length distributions across functional classes of genes (Figure 5A) prompted a
possible explanation for these observations. Shorter than average stems were observed for genetic
information processing, belying the expectation of these genes being the longest-residing genes in the
nascent eukaryotic lineage. We suggest that another major factor determining the relative stem length
of a gene might be the amount of evolution post-acquisition, necessary to adjust an acquired gene to a
new biological environment. The genes inherited from the Asgard ancestor would have been pre-
adapted to the cellular environment of the evolving protoeukaryotes, whereas genes acquired from
radically different bacterial sources had to substantially adapt post-acquisition, increasing the apparent
lengths of their pre-LECA stems, which is at least partially supported by our general observation of long
stems from diverse bacteria (supplementary Figure 7C).
381

Figure 5. Ancestral stem length distributions for different groups of prokaryotes contributing to eukaryogenesis and different functional classes of genes. A) Schematic of stem length calculation. Stem Lengths (SL) were calculated as the lengths of branches from the shared root of Eukarya and the inferred ancestral prokaryotic sister phylum to the root of the Eukarya clade. Normalized Stem Lengths (NSL) shown in A and B were calculated by dividing SL by median Branch Lengths from the root of Eukarya to individual Eukarya leaves (BLs). **B)** Left, Probability Density Function (PDF) of normalized stem lengths taken from samples of individual core eukaryotic genes associated with prokaryotic taxa, obtained through KDE. Thin bar indicates 5[th] percentile range, thick bar includes 25[th] percentile ranges with median indicated. Right, fractional Cumulative Distribution Functions (CDF) of normalized stem lengths together with 20 of 200 bootstrap replicates from EPOCs within each category. Second Y loglinear axis indicates value of the 2-sided Mann Whitney U test of bootstrap CDF distributions per stem length between Asgards and either Cyano- or Alphaproteobacteria. **C)** As in B with genes grouped by BRITE A Metabolism (KO:09100), Genetic Information Processing (KO:09120), Oxidative phosphorylation (map00190 excluding KOGs for V-type ATPases).

Discussion

Reconstruction of eukaryogenesis is a moving target, as strikingly demonstrated by the discovery and expanding exploration of Asgard archaea which revolutionized the field [2-7]. Nevertheless, recent progress in prokaryote and eukaryote genome sequencing and the advances of metagenomics have created unprecedented opportunities for understanding the phylogenomic origins of eukaryotes. In this work, we took advantage of this expanded genome collection, coupling it with maximally sensitive HMM profile-profile searches, and performed a comprehensive phylogenetic analysis of core eukaryotic genes tentatively mapped to the LECA, within a statistical framework focused on testing evolutionary hypotheses using constraint trees.

Many previous studies, especially early ones, demonstrated the apparent chimeric origins of the core eukaryotic gene set, with a bacterial contribution quantitatively exceeding the archaeal one, the latter being largely limited to information processing [17-19,62]. The discovery and exploration of Asgard archaea partially changed this notion by demonstrating that many genes involved in various cellular processes and systems, such as membrane remodeling and cytoskeleton, were of apparent Asgard origin [5,6,8]. Indeed, the presence of an elaborate cytoskeleton in Asgard archaea has been subsequently experimentally demonstrated [24]. The present work substantially extends the dominance of Asgard in the ancestry of the eukaryotes by demonstrating a likely Asgard origin for the majority of genes traced to the LECA, and a major Asgard association for nearly all functional systems and pathways of the eukaryotic cell. In only a handful of metabolic pathways did the detected bacterial contribution exceed that of Asgard archaea. The most conspicuous of the bacterial contributions, not unexpectedly, comes from Alphaproteobacteria, and accounts for the core components of the electron transport chain, iron-sulfur cluster metabolism, and the mitochondrial translation system. However, although prominent and functionally coherent, the Alphaproteobacterial contribution is relatively small, not comparable to that of Asgard, and notably, almost entirely restricted to mitochondrial functions. The contributions of other bacterial phyla, although cumulatively substantial, and in select cases strongly supported for individual EPOCs, failed to show any consistent functional trends, with the exception of notable but limited associations between Myxococcota and the synthesis of nucleotides, nucleoside-sugars and steroid precursors.

These findings are best compatible with a scenario of eukaryogenesis in which the Asgard FECA (the base node of the lineage leading from Asgard to the LECA) already possessed many signature features of eukaryotic cells, including the cytoskeleton and the endomembrane system as posited in some previous models [26,63]. Under this scenario, mitochondrial acquisition was a relatively late event that made a limited, but functionally crucial, contribution to the gene composition of the emerging eukaryote (Figure 6). Other bacterial contributions appear to be piecemeal, possibly, stemming from a continuous but non-specific capture of genes from diverse bacteria during the evolutionary path from the ancestral Asgard archaeon to the LECA (Figure 6). The syntrophic lifestyle of many Asgard archaea [32] is conducive to extensive gene influx from bacteria, enhancing the general trend of mosaic gene composition of prokaryotic genomes [71,72]. Much of the bacterial gene acquisition likely preceded the emergence of the proto-eukaryotic lineage and more occurred along the path from the FECA to the LECA (Figure 6). Thus,

438 some of the core eukaryotic genes, although proximally inherited from the Asgard ancestor, might
439 ultimately be of bacterial origin, having been carried via Asgards into the FECA. As such, we obtained no
440 indication of another symbiotic event contributing to eukaryogenesis.
441
How eukaryotes acquired the bacterial-type membranes remains an enigma. Whereas the eukaryotic
isoprenoid synthesis appears to have been derived from Asgard lipid synthesis, the ancestry of the
enzymes involved in membrane lipid biosynthesis appears complex and possibly mosaic, remaining
unresolved in the present analysis. Nevertheless, the Asgard affinity of adjoining systems and pathways,
such as the synthesis of GPI anchors, sphingolipids, and isoprenoids, suggests that a substantial
component of the lipid biosynthesis and modification capacity of eukaryotes persisted from the Asgard
ancestor. If an Asgard archaeal cell was the principal scaffold on which the eukaryotic cell evolved,
replacement of the archaeal membrane with a bacterial one must have occurred at an early stage of
eukaryogenesis, progressing through a mixed membrane stage. Viable bacteria with a mixed bacterial-
archaeal membrane have been experimentally engineered, albeit typically with a substantial fitness
decrease [64,65,79], demonstrating the possibility of such a scenario. Membrane replacement might have
even predated mitochondrial endosymbiosis, following early capture of the necessary bacterial enzymes
by the Asgard archaeon (Figure 6).
455
A major issue for understanding eukaryogenesis is the origin of the nucleus, the ubiquitous, eponymous
organelle of eukaryotic cells [66,67]. Our analysis supports the near complete origin of information
processing systems located in the nucleus from Asgard ancestors. However, given that the nuclear
envelope membrane is an extension of the ER and as such is of the bacterial type [68], whereas the nuclear
pore consists largely of repetitive proteins of uncertain provenance [69], the origin of the nucleus as a
membrane-bounded compartment remains an unresolved question that has to be addressed in
conjunction with the membrane replacement problem.
463

Figure 6. Hypothetical scenario of eukaryogenesis. The figure shows the gene flow along the succession
of events leading from a simple archaeal ancestor through the more complex Asgard intermediate
capturing the proto-mitochondrial Alphaproteobacterial endosymbiont to the LECA. The wide colored
shape denotes the dominant trend of the origin of eukaryotic genes from Asgard ancestors, with light
blue portions corresponding to archaeal evolution and the light brown portion to the evolution of
protoeukaryotes starting with the Alphaproteobacterial endosymbiosis. The purple arrows denote the
smaller, piecemeal contributions of gene influx from various bacteria that is posited to have been
occurring at all stages of the depicted evolutionary scenario.
473
The conclusions of this work are subject to caveats stemming, above all, from the biased and still limited
sampling of sequenced archaeal and bacterial genomes. In particular, almost all currently available
genomes of Asgard archaea are incomplete metagenomic assemblies, with the exception of only a few
closed circular genomes [22-24,31,70] which inevitably leads to uncertainty in the estimation of the Asgard
contribution to eukaryogenesis. The same pertains to Alphaproteobacteria, both because of the
sequencing bias whereby many of the sequenced alphaproteobacterial genomes come from symbionts
and parasites with reduced gene complements, and because the ancestor of the mitochondria
apparently belongs outside of the currently known diversity of Alphaproteobacteria [71]. Other bacterial
phyla are even more severely under-sampled, potentially, leading to underestimates of their
contributions to the evolution of eukaryotes. For example, given the limited genomic information, it is
difficult to rule out that we capture only part of a major signal from Myxococcota, proposed as one of
the partners in the syntrophic scenario of eukaryogenesis [15,22]. Our pangenome-based approach to
exclude late HGT events from ancestral inference potentially could overlook possible contributions from
narrow or severely under-sampled clades and might be insufficient to compensate for extensive and
continuous horizontal transfer across diverse eukaryotic taxa. Additionally, if future work indicates a
substantial mismatch between the representations of eukaryotic biology in KEGG and in a reconstructed
gene complement of the LECA, our results should be revisited. Expanding sampling of prokaryotic
genome diversity and further technical improvements in horizontal transfer detection might in the
future lead to a substantial modification or even complete revision of the model of eukaryogenesis. It
nevertheless appears that the dominant contribution of Asgard archaea, the main finding of this work, is
here to stay.
495

Methods

Database curation

For prokaryotes, 75 million sequences were curated from 47545 fully sequenced prokaryotic genomes obtained from the NCBI GenBank (https://ftp.ncbi.nlm.nih.gov/genomes/) in November 2023 (prok2311) and supplemented with 441150 proteins sequences from 63 Asgard metagenomes [4,5] (prok2311As), selected to represent all clades in these two publications where available. Protein sequences were either directly taken from GenBank annotations, or produced by Prodigal v2.6.3 [72] trained on the set of 12 complete or chromosome-level assembled Asgard genomes. To only include those sequences which are widely present within prokaryotic families, and to minimize the possibility of post-LECA horizontal transfer from eukaryotes, soft-core pangenomes were constructed for each of our 26 curated taxonomic groups, mostly conforming to the NCBI taxonomy rank "class" (see below and supplementary Data 1). To construct the soft-core pangenomes, all sequences from each taxonomic group were clustered individually using mmseqs2 [75] "mmseqs cluster -s 7 -c 0.9 --cov-mode 0" [75]. All clusters which did not contain sequences from at least 50% of all bacterial species (here defined by lowest rank taxonomic ID within the NCBI taxonomy), or 20% of eukaryotic sequences (see below), were rejected from the database. The effects of alternative pangenome definitions, such as 10, 24, 50, and 67% proteins per class, are shown in supplementary figure 1D.

The eukaryotic database was constructed starting from a curated set of 72 genomes available on the NCBI GenBank. In order to provide a more accurate representation of eukaryotic diversity, this dataset was supplemented with EukProt v3 [73], a curated database aiming to provide a sparse representation of eukaryotic biology, to create Euk72Ep containing 30.3 million sequences. As sequences included in EukProt v3 come from a diverse range of sources, some known to contain contaminant prokaryotic sequences, the database was screened for prokaryotic contamination. This screen was carried out by first constructing candidate prokaryotic and eukaryotic HMM databases as described below and querying both databases against the eukaryotic sequence database using hhblits [40]. All eukaryotic sequences with a top hit alignment score from a prokaryotic HMM were removed from the database prior to initial clustering.

Curation of taxonomic labels

In order to provide a compromise between taxonomic specificity and accuracy of clade assignment, we curated a set of taxonomic labels for both Euk72Ep and Prok2311As to act as representatives. To construct this set, we started with manually assigning all species in EukProt to their closest relative in the NCBI taxonomy. Then, each species from Prok2311As and Euk72Ep was assigned a taxonomic label corresponding to their "Class" rank (for example, Alphaproteobacteria, Thermococci, Mammalia) based on the NCBI Taxonomy of November 2023 using tools from the ete3 3.1.3. toolkit [74] as well as custom scripts. For species without a corresponding "Class" rank, the closest relevant rank was manually assigned. This procedure yielded an initial list of 317 unique taxonomic class identifiers. As the rank "Class" does not evenly partition biological diversity, identifiers were further manually mapped to a curated set of 45 eukaryotic and 26 prokaryotic taxonomic superclasses, each present in the NCBI

taxonomy (for example, Metazoa, Streptophyta, TACK archaea; see supplementary Data 1. Due to the
partially unresolved taxonomic relationship within Asgard archaea, all candidate Asgard sequences from
Prok2311 or from references 6,7 were assigned the class label "Asgard". Following the recent
reclassification of Deltaproteobacteria into Myxococcota, Desulfobacterota, Bdellovibrionota and
SAR324 species which could not be mapped into either of these clades using existing taxonomic
annotation were classified as Deltaproteobacteria (0.025% of total data).
543

Sequence clustering and profile database generation
To transform Euk72Ep and Prok2311As into profile databases suitable for sensitive HMM-HMM
searches, we implemented an unsupervised, cascaded sequence-profile clustering pipeline using the
tools available in the mmseqs2 software suite [75]. In brief, each sequence database was initially clustered
at 90% sequence identity and 80% pairwise coverage using "mmseqs linclust" and cluster
representatives were chosen using "mmseqs result2repseq". This procedure provides a non-redundant
set of 5.9 million prokaryotic and 12.7 million eukaryotic sequences for further analysis. The non-
redundant sequences are further collapsed into a set of initial profiles and consensus sequence pairs
using "mmseqs cluster", "mmseqs result2profile" and "mmseqs profile2consensus". All consensus
sequences are queried against their profiles using "mmseqs search" and results clustered using
"mmseqs clust". Clusters are mapped to their original non-redundant sequences using "mmseqs
mergeclusters" and profiles and consensus sequences are constructed once again. In order to grow
clusters based on their sequence-profile alignments, this procedure of "mmseqs profile2consensus -
mmseqs search - mmseqs clust - mmseqs result2profile" is iterated until cluster sizes converge. An 80%
pairwise coverage is maintained throughout all steps of the cascaded clustering protocol to avoid
homologous overextension. The final clustered databases contain 14.1 million and 91.000 clusters for
Euk72Ep and Prok2311As, respectively.
561

To transform the unsupervised clusters into a set of HMM profiles, we employed a mixed MSA
construction strategy coupled with automatic data reduction to create accurate alignments for a wide
diversity of cluster sizes and sequence diversities. To construct HMMs, all non-singleton sequence
clusters are first aligned using FAMSA 2.0.1. [76] to produce an initial alignment. Using the initial
alignment, we then reduce the sequence space by calculating an approximate tree using FastTree 2.1.1.
using "FastTree -gamma" [77], remove long branch outliers as described in "EPOC tree construction and
processing", and iteratively removing the closest pairwise leaves using ete3 [74], keeping the leaf closest
to the root, until the desired sequence set size is reached. For profile generation, we retain no more
than 300 sequences for both Prok2311As and Euk72Ep. This reduced sequence set is then realigned with
muscle5 5.1.linux64 [43] using "muscle -diversified" with 5 replicates, and the maximum column
confidence alignment is extracted using "muscle -maxcc". This alignment is then trimmed to keep
columns with more than 0.2 bits of Shannon information defined as the difference of log2(20) and the
Shannon entropy of the column amino acid distribution. This process of data
reduction/alignment/trimming is referred to as "prune-and-align" and employed later for downstream
analysis. Trimmed alignments are turned into an HMM database using HH-suite with "hhconsensus -M

50", hhmake and "cstranslate -f -x 0.3 -c 4 -I a3m". The final HH-suite databases contained 480.000 and
1.6 million profiles for prokaryotes and eukaryotes, respectively.
579

Cluster annealing

As the initial mmseqs2 based clustering was carried out in an unsupervised manner, we considered it
necessary to ensure that the results were not contingent on this unsupervised partitioning of the data.
To this end, a further cluster annealing step was developed which aims to redefine clusters based on
clades well separated monophyletic clades on phylogenetic trees, rather than by sequence similarity
alone. Starting with the existing clustered databases Euk72Ep and Prok2311As, HMM profiles were
calculated for each non-singleton cluster and all vs all HMM-HMM searches were performed using
Hhblits 3.3.0 (HHblits -n 1 -p 80), retaining hits with at least 80% HHblits probability and 50% pairwise
profile coverage. The resulting hits were then clustered using greedy set clustering, as described in
mmseqs2, implemented in python. These superclusters represent the limit of detection using out
sequence-based methods, but might involve overclustering, possibly merging multiple similar gene
families into one set. To mitigate the potential effect of such overclustering, cluster partitioning was
performed based on phylogenetic tree analysis. To this end, the sequence space of the superclusters
was first reduced using the "prune-and-align" approach described above, after which a master tree was
constructed using FastTree -gamma. Given the resulting set of trees, we sought to identify well
separated monophyletic sets of leaves, the presence of which would warrant partitioning of the
respective superclusters. To perform this tree partitioning in an unsupervised manner, the all-vs-all
pairwise leaf distances were calculated as a square matrix and embedded into 2 dimensions using UMAP
version 0.5.3 (n_neighbors=50, min_dist=0.3, distance='euclidean'). This embedding was clustered using
HDBSCAN as implemented in sklearn version 1.3.0 with default parameters. We explicitly allow single
cluster partitions to ensure that trees with uniform distributions of pairwise branch lengths remained as
single clusters. Mapping clusters back to the phylogenetic trees showed that UMAP embedded
HDBSCAN clusters were well separated in the original trees (Supplementary figure 8). These annealed
clades were used in downstream processing, representing a stricter definition of EPOCS than the initial
cascaded mmseqs2 clusters. The annealing process redistributed sequences from the 1.6 million
eukaryotic clusters into 1.5 million annealed clusters and from 460.000 prokaryotic sequences into
280.000 annealed clusters, highlighting the already dense representation of initial mmseqs2 clusters.
Because the embeddings were done per tree, the partitioning into subsets is independent of branch
lengths and merely relies on the internal distribution of data. Thus, well-formed subclades could be
identified in sequence sets independent on the internal diversity, an essential feature when applying
this procedure.
611

Search for prokaryotic homologs of eukaryotic proteins

Given that clade-specific protein families were not of direct relevance to this study, we reduced the
search space by excluding HMM profiles from eukaryotic clusters with less than 10 sequences and a
lowest common ancestor with a taxonomic rank below "Superkingdom" as calculated by "mmseqs lca".
This requires a sequence cluster to contain at least two sequences from different kingdoms, as defined
by the NCBI taxonomy (ex. Amoebozoa and Haptista, or Viridiplanta and Opistokonta), retaining 142,000

profiles of conserved eukaryotic proteins. To associate eukaryotic sequence clusters with homologous
prokaryotic clusters, we queried the Euk72Ep HMM database against all Prok2311As HMM profiles using
a single iteration of HHBlits 3.3.0 as "HHblits -n 1 -p 80" and retaining the hits with at least 80% HHblits
probability and 80% pairwise profile length. These filters resulted in a total of 20,700 accepted
eukaryotic query profiles targeting a total of 8,300 unique prokaryotic profiles. These clusters contained
a total of 5.7 million and 1.9 million sequences.
624

EPOC multiple sequence alignment construction

Unless otherwise stated, all subsequent steps were carried out using custom python 3.9 scripts, with
taxonomy and tree parsing carried out using ete3 [74]. All sequences from a eukaryotic cluster with at
least 10 sequences and a lowest common ancestor at the rank of "Superkingdom" (NCBI taxonomy),
together with all members of the homologous prokaryotic clusters identified in the HMM-HMM search
with at least 80% probability and pairwise sequence coverage, forms an EPOC. As EPOCs varied in size by
more than 5 orders of magnitude (10 to 100,000 sequences), robust data subsampling is necessary for
accurate downstream MSA generation and tree construction. We employed a variant of the "prune-and-
align" strategy described above for profile generation taking into account the taxonomic distribution of
the constituent sequences. Rather than iteratively pruning the closest leaves pairwise, we first label
each leaf with its corresponding taxonomic label and only prune monophyletic groups, maintaining a
count of all pruned leaves per clade. If the procedure fails to reach the target leaf number before
converging, all leaves are ordered by the number of pruned relatives and leaves with the lowest number
of relatives deleted (retaining the largest monophyletic groups) until the target size is reached. The
result approximates a maximum diversity representation, preferentially pruning isolated singleton
clades. Using this modified protocol, eukaryotic and prokaryotic sequences are cropped to a maximum
of 30 and 70 sequences, respectively. Each EPOC therefore consists of a maximum of 100 representative
sequences. All EPOCs are aligned using "muscle -diversified" with 5 replicates, and the maximum
confidence alignment is extracted using the --maxcc argument. This alignment is then trimmed to keep
columns with more than 0.15 bits of Shannon information content for tree reconstruction.
645

EPOC tree construction and processing

From the pruned and trimmed alignment, we created a maximum likelihood tree using IQtree2 2.3.5.
"IQtree2 -B 1000 -bnni -m MFP -mset LG,Q.pfam --cmin 4 --cmax 12" with model parameters estimated
by model finder plus [44]. Candidate models and rate category search ranges were selected based on a
test set of 200 randomly chosen EPOCs from the filtered core set analyzed with full model parameter
evaluation by Model finder [44] and 10 tree replicates. The LG and Q.pfam models and rate category
ranges were chosen as they consistently produced the highest ranking log-likelihoods for a random
sample of 200 EPOCs run with full model estimation in IQtree.
654

As the master trees are built from sequences obtained through sensitive cascaded sequence-profile
clustering we would in rare cases observe long-stemmed clade outliers, in addition to more common
long stem leaf outliers. In order to assess and remove such erroneous data, we first estimated the log-
normal distribution of all stem lengths using scipy.stats 1.11.1 and excluded any stems outside the 0-

99.5 % Probability Point Function interval. This simultaneously removes long terminal leaf branches as
well as highly diverged clades. In rare cases where this pruning would remove either the entire
eukaryotic clade or more than 30 % of all leaves, the EPOC was discarded (387 EPOCs). The resulting
trees were then re-rooted using a weighted midpoint approach so that the sum of all branch stem
lengths on either side of the root was equal. For clarity, no trees in our current study have been
"phylogenetically rooted" in the biological sense. All references to the "root" refer to the origin of the
data structure, not the biological concept.
666
To assign taxonomic clades, all leaf nodes were assigned taxonomic labels from our curated set and
monophyletic clades were collapsed, taking as the representative the lowest branching leaf. We found
that assigning clades purely based on a monophyletic definition of taxonomic purity severely hampered
the analysis as leaves can be taxonomically incongruous with their neighbors. This effect most likely
stems from cases of locally erroneous tree topology or local HGT, but severely complicates the
downstream analysis. We therefore adopted the notion of a "soft LCA" representing the root of a close-
to-monophyletic clade. To greedily rank all the best "soft LCAs", for each taxonomic label, the tree was
traversed from each monophyletic clade root to the tree root, calculating a score for each internal node:
675
676 (number of leaves with label X in clade / clade size) * (number of leaves with label X in clade / total
677 number of label X)
678
This metric balances taxonomic "purity" and "scope" for each possible clade of label X. All such nodes
are then ordered by this score. Because this score is sensitive to the total number of taxonomic labels in
an EPOC, it was not used as a filtering criterion at any point in the analysis, yet is present in the
metadata, mentioned here for completeness. To avoid overinterpreting small clades, clades were
considered valid if they represented at least 3 sequences with clade purity > 0.8 for prokaryotes, and at
least 5 sequences with purity > 0.8 for eukaryotes. Trees that fail to identify either any eukaryotic or any
prokaryotic clades under these constraints, primarily derived from small EPOCs, are discarded (411
EPOCs). Trees with more than 3 valid eukaryotic clades were considered to show high paraphyly and
likewise discarded (781 EPOCs).
688
Evolutionary hypothesis testing using constraint trees
In order to assess the relative probabilities of each sampled prokaryotic clades to represent the
eukaryotic sister clade, we employed evolutionary hypothesis testing using constraint trees as
implemented in IQtree2 [44]. For each EPOC and for each detected eukaryotic clade, we generated a set of
constraint trees, one for each possible prokaryotic sister. Each constraint tree had three defined clades
enforced throughout tree calculations: 1) the eukaryotic group as defined by all sequences below the
eukaryotic LCA, 2) one of the prokaryotic sister groups and 3) all other prokaryotic leaves. We then
construct a set of local trees from slices of the original MSA using IQtree2, forcing the constraint tree
topology using "iqtree2 -g", using the same evolutionary model as the original, unconstrained "master
tree". The set of constrained trees is then ranked using "iqtree2 -z" calculating the relative model
assignment confidence for each of the constrained topologies and resulting test metrics are saved for

downstream analysis. As the number of trees to evaluate scales by the number of eukaryotic clades

multiplied by the number of prokaryotic clades, we limit the sampling to a maximum of 3 clades per tree

for eukaryotes, discarding rare cases of high eukaryotic paraphyly, and only consider the 12 closest

prokaryotic clades to each individual eukaryotic clade, based on their topological distance calculated as

the number of non-root tree bifurcations from one node to another (supplementary Figure 2). The

procedure results in an Expected Likelihood Weight (ELW)[41] for each of the evaluated model trees, here

taken as being analogous to model selection confidence that an evaluated prokaryotic clade is the most

significant sister to a eukaryotic clade.

EPOC annotation

To functionally annotate protein families within EPOCs, we generated profiles based on protein

sequences in KEGG release 110 [42]. For each KOG, we extracted Kegg metadata including KEGG pathways

and BRITE labels via KEGG's API. At the time of parsing (January 2024), KEGG contained 26,695 KOGs.

Due to KEGG API constraints, protein sequences in KEGG were extracted from Uniprot, first by mapping

KEGG proteins to Uniprot IDs via Uniprot's ID mapping service (uniprot.org/id-mapping), then by

extracting the sequences for these Uniprot IDs corresponding to each KOG [78].

For each KOG, we generated HMM profiles using the aforementioned prune-and-align pipeline. To

increase specificity of labeling, we partitioned those KOGs which encompass both prokaryotic and

eukaryotic sequences into separate taxonomic groups and sequence alignments were generated

separately for prokaryotes and eukaryotes. Each eukaryotic HMM profile forming the basis for a EPOC

was queried against our KEGG profile database using "HHblits -n 3 -p 80". Results were subsequently

filtered to a pairwise profile coverage of 0.7 using custom scripts. As a result, 12,600 of the 13,500

EPOCs could be annotated by at least one KOG profile at 80% probability and 70% pairwise coverage.

The highest probability hits were used for annotation.

Data filtering and removal of cases of possible late horizontal gene transfer

Unless otherwise stated, all data is interpreted from the core set of EPOCs meeting the following

criteria: 1) the eukaryotic constituent cluster profile identifies at least one target profile in our KEGG

profile database at a probability of 80% and 70% coverage, 2) the eukaryotic clades include more than 5

distinct taxonomic labels, at least one from Amorphea and one from Diaphoretickes, encompassing

LECA, 3) only prokaryotic sister clades with $0.4 < ELW < 0.99$ were considered reliable and were included

in the analysis. We note the independence on aELW values on the choice og number of wukaryotic

clades as showin in Supplementary Figure 7A. Cases of ELW >= 0.99 were not considered as these are

found to be disproportionately eukaryotic-like, often branching from within the eukaryotic clade itself,

and thus likely derived from late horizontal transfer post LECA. Due to the presence of such high ELW

values within Alphaproteobacterial associations with Oxidative phosphorylation, in this case, as an

exception, ELW values of 1 were included.

Taxonomy remapping and EPOC construction under GTDB taxonomy

To ensure that the results were robust to taxonomy, the prok2311 database was reformatted under the

GTDB taxonomy (version 220) at the level of "phyla" [47]. For taxonomic mapping, first, genomes present

in both prok2311 and GTDB were directly annotated with the GTDB taxonomy. For the remaining

genomes, taxonomy was assigned through consensus voting by using the set of GTDB marker genes,

searched against our data using "mmseqs search -s 7 -c 0.5 -e 1e-10". A phyla level taxonomic label was

assigned to each genome based on the largest number of total hits. Genomes without significant hits to

GTDB marker genes were discarded for the purpose of this analysis (0.92% of all data). This procedure

mapped the prok2311 data to 96 GTDB phyla which were used as taxonomic assignment for EPOC

calculations as described above. All additional Asgard data taken from Ref 8 were directly assigned as

Asgardarchaeota.

Data visualization and plotting

All plots were prepared using python v3.9 and pandas v.2.0.3 with Altair v.4.2.2. Layout, annotation, and

vector editing was done using Inkscape v.1.1.1. All statistical tests were carried out using scipy.stats

1.11.1

Data availability
All final non-redundant databases with taxonomy annotation, clustering data, HMM profiles, EPOCs data
and final parsed and tabulated results and metadata is available at Zenodo
https://zenodo.org/records/15048010
762
763 **Code availability**
764 All code and scripts used to generate the final parsed data is available on Github at:
765 https://github.com/VictorTobiasson/eukgen
766
767 **Supplementary Data File 1**

768 Excel file with four sheets: 1) List of revised taxonomy mapping species without well curated taxonomy
769 to closest existing taxonomic ranks. 2) List of revised taxonomy merging sparsely populated classes. 3)
770 Manually curated list of genes involved in Fe-S assembly. 4) Tabular data of all tables included in figures.
771
772 **Author contributions**
773 E.V.K. initiated the study; V.T. and Y.I.W. developed the pipeline for data analysis; V.T. and J.L. collected
774 the data; V.T., J.L., Y.I.W and E.V.K. performed data analysis; V.T. and E.V.K wrote the manuscript which
775 was read, edited and approved by all authors.
776

777 **Acknowledgements**
778 The authors thank Toni Gabaldon and Koonin Group members for helpful discussions. The authors'
779 research is supported by the Intramural Research Program of the National Institutes of Health of the
780 USA (National Library of Medicine).

781
782 **Competing interests**
783 The authors declare no competing interests.
784
785

References

References

1   Embley, T. M. & Martin, W. Eukaryotic evolution, changes and challenges. *Nature* **440**, 623-630
(2006).
2   Vosseberg, J. *et al.* The emerging view on the origin and early evolution of eukaryotic cells.
Nature **633**, 295-305 (2024). https://doi.org:10.1038/s41586-024-07677-6
3   van der Giezen, M. Hydrogenosomes and mitosomes: conservation and evolution of functions. *J*
Eukaryot Microbiol **56**, 221-231 (2009). https://doi.org:JEU407 [pii] 10.1111/j.1550-
7408.2009.00407.x
4   Spang, A. *et al.* Complex archaea that bridge the gap between prokaryotes and eukaryotes.
Nature **521**, 173-179 (2015). https://doi.org:10.1038/nature14447
5   Zaremba-Niedzwiedzka, K. *et al.* Asgard archaea illuminate the origin of eukaryotic cellular
complexity. *Nature* **541**, 353-358 (2017). https://doi.org:10.1038/nature21031
6   Liu, Y. *et al.* Expanded diversity of Asgard archaea and their relationships with eukaryotes.
Nature **593**, 553-557 (2021). https://doi.org:10.1038/s41586-021-03494-3
7   Eme, L., Spang, A., Lombard, J., Stairs, C. W. & Ettema, T. J. G. Archaea and the origin of
eukaryotes. *Nat Rev Microbiol* **15**, 711-723 (2017). https://doi.org:10.1038/nrmicro.2017.133
8   Eme, L. *et al.* Inference and reconstruction of the heimdallarchaeial ancestry of eukaryotes.
Nature **618**, 992-999 (2023). https://doi.org:10.1038/s41586-023-06186-2
9   Lopez-Garcia, P. & Moreira, D. The symbiotic origin of the eukaryotic cell. *C R Biol* **346**, 55-73
(2023). https://doi.org:10.5802/crbiol.118
10   Alberts, B. *et al.* *Molecular Biology of the Cell*. 7 edn, (W.W. Norton & Co, 2022).
11   Cavalier-Smith, T. Eukaryotes with no mitochondria. *Nature* **326**, 332-333 (1987).
https://doi.org:10.1038/326332a0
12   Sogin, M. L., Gunderson, J. H., Elwood, H. J., Alonso, R. A. & Peattie, D. A. Phylogenetic meaning
of the kingdom concept: an unusual ribosomal RNA from Giardia lamblia. *Science* **243**, 75-77
(1989). https://doi.org:10.1126/science.2911720
13   Sagan, L. On the origin of mitosing cells. *J Theor Biol* **14**, 255-274 (1967).
14   Martin, W., Hoffmeister, M., Rotte, C. & Henze, K. An overview of endosymbiotic models for the
origins of eukaryotes, their ATP-producing organelles (mitochondria and hydrogenosomes), and
their heterotrophic lifestyle. *Biol Chem* **382**, 1521-1539 (2001).
https://doi.org:10.1515/BC.2001.187
15   Lopez-Garcia, P. & Moreira, D. The Syntrophy hypothesis for the origin of eukaryotes revisited.
Nat Microbiol **5**, 655-667 (2020). https://doi.org:10.1038/s41564-020-0710-4
16   Brown, J. R. & Doolittle, W. F. Archaea and the prokaryote-to-eukaryote transition. *Microbiol*
Mol Biol Rev **61**, 456-502 (1997). https://doi.org:10.1128/mmbr.61.4.456-502.1997
17   Rivera, M. C., Jain, R., Moore, J. E. & Lake, J. A. Genomic evidence for two functionally distinct
gene classes. *Proc Natl Acad Sci U S A* **95**, 6239-6244 (1998).
https://doi.org:10.1073/pnas.95.11.6239
18   Ribeiro, S. & Golding, G. B. The mosaic nature of the eukaryotic nucleus. *Mol Biol Evol* **15**, 779-
788 (1998). https://doi.org:10.1093/oxfordjournals.molbev.a025983
19   Koonin, E. V. *et al.* A comprehensive evolutionary classification of proteins encoded in complete
eukaryotic genomes. *Genome Biol* **5**, R7 (2004).
20   Imachi, H. *et al.* Promethearchaeum syntrophicum gen. nov., sp. nov., an anaerobic, obligately
syntrophic archaeon, the first isolate of the lineage 'Asgard' archaea, and proposal of the new
archaeal phylum Promethearchaeota phyl. nov. and kingdom Promethearchaeati regn. nov. *Int J*
Syst Evol Microbiol **74** (2024). https://doi.org:10.1099/ijsem.0.006435

21   Lombard, J., Lopez-Garcia, P. & Moreira, D. The early evolution of lipid membranes and the
three domains of life. *Nat Rev Microbiol* **10**, 507-515 (2012).
https://doi.org:10.1038/nrmicro2815

22   Moreira, D. & Lopez-Garcia, P. Symbiosis between methanogenic archaea and delta-
proteobacteria as the origin of eukaryotes: the syntrophic hypothesis. *J Mol Evol* **47**, 517-530
(1998). https://doi.org:10.1007/pl00006408

23   Krupovic, M., Dolja, V. V. & Koonin, E. V. The virome of the last eukaryotic common ancestor and
eukaryogenesis. *Nat Microbiol* **8**, 1008-1017 (2023). https://doi.org:10.1038/s41564-023-01378-
y

24   Imachi, H. *et al.* Isolation of an archaeon at the prokaryote-eukaryote interface. *Nature* **577**,
519-525 (2020). https://doi.org:10.1038/s41586-019-1916-6

25   Lopez-Garcia, P. & Moreira, D. Cultured Asgard Archaea Shed Light on Eukaryogenesis. *Cell* **181**,
232-235 (2020). https://doi.org:10.1016/j.cell.2020.03.058

26   Pittis, A. A. & Gabaldon, T. Late acquisition of mitochondria by a host with chimaeric prokaryotic
ancestry. *Nature* **531**, 101-104 (2016). https://doi.org:10.1038/nature16941

27   Gabaldon, T. Origin and Early Evolution of the Eukaryotic Cell. *Annu Rev Microbiol* **75**, 631-647
(2021). https://doi.org:10.1146/annurev-micro-090817-062213

28   Steinegger, M. & Soding, J. MMseqs2 enables sensitive protein sequence searching for the
analysis of massive data sets. *Nat Biotechnol* **35**, 1026-1028 (2017).
https://doi.org:10.1038/nbt.3988

29   Steinegger, M. *et al.* HH-suite3 for fast remote homology detection and deep protein
annotation. *BMC Bioinformatics* **20**, 473 (2019). https://doi.org:10.1186/s12859-019-3019-7

30   Richter, D. J. *et al.* EukProt: A database of genome-scale predicted proteins across the diversity
of eukaryotes *Peer Community J* **2**, e56 (2022).

31   Strimmer, K. & Rambaut, A. Inferring confidence sets of possibly misspecified gene trees. *Proc*
Biol Sci **269**, 137-142 (2002). https://doi.org:10.1098/rspb.2001.1862

32   Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y. & Morishima, K. KEGG: new perspectives on
genomes, pathways, diseases and drugs. *Nucleic Acids Res* **45**, D353-D361 (2017).
https://doi.org:10.1093/nar/gkw1092

33   Edgar, R. C. Muscle5: High-accuracy alignment ensembles enable unbiased assessments of
sequence homology and phylogeny. *Nat Commun* **13**, 6968 (2022).
https://doi.org:10.1038/s41467-022-34630-w

34   Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in
the Genomic Era. *Mol Biol Evol* **37**, 1530-1534 (2020). https://doi.org:10.1093/molbev/msaa015

35   Derelle, R. *et al.* Bacterial proteins pinpoint a single eukaryotic root. *Proc Natl Acad Sci U S A*
112, E693-699 (2015). https://doi.org:10.1073/pnas.1420657112

36   Al Jewari, C. & Baldauf, S. L. Conflict over the Eukaryote Root Resides in Strong Outliers, Mosaics
and Missing Data Sensitivity of Site-Specific (CAT) Mixture Models. *Syst Biol* **72**, 1-16 (2023).
https://doi.org:10.1093/sysbio/syac029

37   Paulick, M. G. & Bertozzi, C. R. The glycosylphosphatidylinositol anchor: a complex membrane-
anchoring structure for proteins. *Biochemistry* **47**, 6991-7000 (2008).
https://doi.org:10.1021/bi8006324

38   Stankeviciute, G. *et al.* Convergent evolution of bacterial ceramide synthesis. *Nat Chem Biol* **18**,
305-312 (2022). https://doi.org:10.1038/s41589-021-00948-7

39   Biran, A., Santos, T. C. B., Dingjan, T. & Futerman, A. H. The Sphinx and the egg: Evolutionary
enigmas of the (glyco)sphingolipid biosynthetic pathway. *Biochim Biophys Acta Mol Cell Biol*
Lipids **1869**, 159462 (2024). https://doi.org:10.1016/j.bbalip.2024.159462

40   Hoshino, Y. & Gaucher, E. A. On the Origin of Isoprenoid Biosynthesis. *Mol Biol Evol* **35**, 2185-
2197 (2018). https://doi.org:10.1093/molbev/msy120
41   Richards, T. A. & van der Giezen, M. Evolution of the Isd11-IscS complex reveals a single alpha-
proteobacterial endosymbiosis for all eukaryotes. *Mol Biol Evol* **23**, 1341-1344 (2006).
https://doi.org:10.1093/molbev/msl001
42   Cassier-Chauvat, C., Marceau, F., Farci, S., Ouchane, S. & Chauvat, F. The Glutathione System: A
Journey from Cyanobacteria to Higher Eukaryotes. *Antioxidants (Basel)* **12** (2023).
https://doi.org:10.3390/antiox12061199
43   Kassube, S. A. & Thoma, N. H. Structural insights into Fe-S protein biogenesis by the CIA
targeting complex. *Nat Struct Mol Biol* **27**, 735-742 (2020). https://doi.org:10.1038/s41594-020-
0454-0
44   Hedges, S. B. The origin and evolution of model organisms. *Nat Rev Genet* **3**, 838-849 (2002).
https://doi.org:10.1038/nrg929
45   Vosseberg, J. *et al.* Timing the origin of eukaryotic cellular complexity with ancient duplications.
Nat Ecol Evol **5**, 92-100 (2021). https://doi.org:10.1038/s41559-020-01320-z
46   Caforio, A. *et al.* Converting Escherichia coli into an archaebacterium with a hybrid heterochiral
membrane. *Proc Natl Acad Sci U S A* **115**, 3704-3709 (2018).
https://doi.org:10.1073/pnas.1721604115
47   Hoekzema, M., Jiang, J. & Driessen, A. J. M. Optimizing Archaeal Lipid Biosynthesis in Escherichia
coli. *ACS Synth Biol* **13**, 2470-2479 (2024). https://doi.org:10.1021/acssynbio.4c00235
48   Tamarit, D. *et al.* A closed Candidatus Odinarchaeum chromosome exposes Asgard archaeal
viruses. *Nat Microbiol* **7**, 948-952 (2022). https://doi.org:10.1038/s41564-022-01122-y
49   Rodrigues-Oliveira, T. *et al.* Actin cytoskeleton and complex cell architecture in an Asgard
archaeon. *Nature* **613**, 332-339 (2023). https://doi.org:10.1038/s41586-022-05550-y
50   Munoz-Gomez, S. A. *et al.* Site-and-branch-heterogeneous analyses of an expanded dataset
favour mitochondria as sister to known Alphaproteobacteria. *Nat Ecol Evol* **6**, 253-262 (2022).
https://doi.org:10.1038/s41559-021-01638-2
51   Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site
identification. *BMC Bioinformatics* **11**, 119 (2010). https://doi.org:10.1186/1471-2105-11-119
52   Huerta-Cepas, J., Serra, F. & Bork, P. ETE 3: Reconstruction, Analysis, and Visualization of
Phylogenomic Data. *Mol Biol Evol* **33**, 1635-1638 (2016).
https://doi.org:10.1093/molbev/msw046
53   Deorowicz, S., Debudaj-Grabysz, A. & Gudys, A. FAMSA: Fast and accurate multiple sequence
alignment of huge protein families. *Sci Rep* **6**, 33964 (2016). https://doi.org:10.1038/srep33964
54   Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2--approximately maximum-likelihood trees for
large alignments. *PLoS One* **5**, e9490 (2010). https://doi.org:10.1371/journal.pone.0009490
55   UniProt, C. UniProt: the universal protein knowledgebase in 2021. *Nucleic Acids Res* **49**, D480-
D489 (2021). https://doi.org:10.1093/nar/gkaa1100
917
918

921
Supplementary Figure 1. Global database statistics. A) Distribution of taxonomic labels present within non-redundant Euk72Ep and Prok2111As (see Methods). Euk72Ep limited to only display taxa with more than 100.000 sequences (99.69% of sequences). **B)** CDF fraction of cluster size distribution following cascaded mmseqs profile/sequence clustering. **C)** CDF fraction of N_{eff} values as calculated by HH-suite for resulting profile databases. **D)** The effect of different pangenome regimes on final aELW values. Pangenomes are reconstructed enforcing a protein presence across 0, 25, 50, or 67% of species within clades and data shown for the full dataset of 14400 EPOCs or core dataset, as described in methods. 50% criteria marked in bold as it is used as reference for the core data present throughout the paper. **E)** Global aELW for data remapped to the GTDB taxonomy at the level of Phylum using marker genes (see Methods). Displaying the top 26 strongest individual contributing phyla out of 92 accounting for 96.4% of all ELW.

933
Supplementary Figure 2. Evolutionary hypothesis testing constraint trees and Expected Likelihood
Weights. Overview of ELW calculation procedure. One unrooted master tree is constructed from all
sequences within a EPOC and the closest n sister clades to the eukaryotic outgroup identified. A set of
unrooted constraint trees are generated, each enforcing three clades, one eukaryotic, one sample sister
and one outgroup with remaining prokaryotic sequences, is present. The set of subtrees are then
constructed using IQtree, each forced to conform to a single guide tree. Log likelihood values for
resulting trees are compared and evaluated using IQtree -z to produce confidence sets and Expected
Likelihood Weight estimations.

Supplementary Figure 3. aELW breakdown of additional maps and pathways with prominent Asgard

and Alphaproteobacterial contributions. A) Overview of most paralogous KOGs estimated by number of

separate EPOCs traceable to the LECA. **B)** Oxidative phosphorylation (map00190) also contains entries

for V-type ATPases which are excluded here as they are not part of mitochondrial metabolism. Fe-S

scaffolds are not members of any map the manually curated set includes KOGs: K22063, K22072,

K22068, K22073, K22070, K22071, K04043, K04044 and K04082. The CIA complex is likewise not

grouped within BRITE and the manual set contains KOGs: K24730, K26403 and K15075.

Supplementary Figure 4. Strongest per taxa association of diverse bacteria to eukaryotes. Top
bacterial taxa associated taxa with aELW presented for their 15 most prominent pathways in KEGG Brite
B. Pathways with more than 20 EPOCs shown.

Supplementary Figure 5. Sparse Myxococcota association with eukaryotes. A) Top, aELW breakdown of enriched Myxococcota pathways showing primary association with nucleotide synthesis. Bottom, curated set of Myxococcota enriched EPOCs containing general nucleic acid metabolism and modification pathways corresponding to KEGG map00760, map01232, map00240 and map00230 with notable presence of Myxococcota. Further selection of only those KOGs within the above pathways which include Myxococcota. **B)** List of KOGs with Myxococcota presence within the curated set of nucleotide synthesis highlighting phosphatases and phosphotransferases.

Supplementary Figure 6. KOGs of strong bacterial association. A) Same as main figure 4 under less stringent selection criteria indicating most prominent prokaryotic pathways with a coverage of at least 10 and 5 EPOCs, aELW shown relative to that of Asgard for the same pathway in pale blue. **B)** Individual EPOCs strongly associated with diverse prokaryotic taxa under stringent selection criteria. EPOCs must contain a eukaryotic outgroup with at least 15 unique taxa encompassing the LECA, have a prokaryotic sister clade with an ELW of > 0.7 which is not derived from Asgard, Euryarchaeota, the TACK group, DPANN group, Cyanobacteriota or Alphaproteobacteria and contain more than 20 sequences. The EPOCs which meet these criteria are highlighted with respective KOGs indicated, aELW displayed.

973
Supplementary Figure 7: **Clade scope definition and stem length distribution for diverse bacteria**
compared to Asgard. **A)** Global aELW dependence on choice of clade size cutoff using core data.
Estimated aELW are independent on choice of eukaryotic clade scope. **B)** 20 full technical replicates,
including muscle5 alignment, IQtree2 calculations and constraint tree generation, of 175 randomly
sampled Alphaproteobacterial stems showing reproducibility of stem length distributions. Compared
with distributions drawn from Asgard and Cyanobacterial stems as per Figure 5. **C)** Left, Probability
Density Function (PDF) of normalized stem lengths taken from samples of individual core eukaryotic
genes associated with prokaryotic taxa, obtained through KDE. Thin bar indicates 5[th] percentile ranges,
thick bar includes 25[th] percentile ranges with median indicated. Right, fractional cumulative distribution
functions of normalized stem lengths.
984

985
Supplementary Figure 8: Annealed cluster EPOCS A) Examples of annealed clades defined based on
embedded and clustered pairwise distance matrices of phylogenetic trees. Top, embedding of pairwise
leaf distance of phylogenetic tree, bottom. Clustered by HDBSCAN allowing single cluster partitions as
shown, left. Partitions form monophyletic subsets of well partitioned branched. **B)** Resulting top level
aELW values derived from the core data clusters as well as annealed EPOCs showing broad correlation
between different core and annealed partitioning schemes. **C)** Correlation of aELW vectors calculated
for all Prokaryotic taxa across all KEGG maps using the annealed and core EPOCs indicating good
agreement for Asgard assignments but greater variance between prokaryotic taxa. Overlaid on global
aELW values. Taxa for which aELW is less than 0.01 are found to be associated only for very few
pathways and as such correlate strongly, notably DPANN has an aELW of 0 and do not contribute to any
pathways, and therefore with a correlation of 1.0 between the two reconstructions.
997

1   Embley, T. M. & Martin, W. Eukaryotic evolution, changes and challenges. *Nature* **440**, 623-630
(2006).
2   Vosseberg, J. *et al.* The emerging view on the origin and early evolution of eukaryotic cells.
Nature **633**, 295-305 (2024). https://doi.org:10.1038/s41586-024-07677-6
3   Eme, L., Spang, A., Lombard, J., Stairs, C. W. & Ettema, T. J. G. Archaea and the origin of
eukaryotes. *Nat Rev Microbiol* **15**, 711-723 (2017). https://doi.org:10.1038/nrmicro.2017.133
4   Eme, L. *et al.* Inference and reconstruction of the heimdallarchaeial ancestry of eukaryotes.
Nature **618**, 992-999 (2023). https://doi.org:10.1038/s41586-023-06186-2
5   Liu, Y. *et al.* Expanded diversity of Asgard archaea and their relationships with eukaryotes.
Nature **593**, 553-557 (2021). https://doi.org:10.1038/s41586-021-03494-3
6   Spang, A. *et al.* Complex archaea that bridge the gap between prokaryotes and eukaryotes.
Nature **521**, 173-179 (2015). https://doi.org:10.1038/nature14447
7   Zaremba-Niedzwiedzka, K. *et al.* Asgard archaea illuminate the origin of eukaryotic cellular
complexity. *Nature* **541**, 353-358 (2017). https://doi.org:10.1038/nature21031
8   Lopez-Garcia, P. & Moreira, D. The symbiotic origin of the eukaryotic cell. *C R Biol* **346**, 55-73
(2023). https://doi.org:10.5802/crbiol.118
9   Alberts, B. *et al. Molecular Biology of the Cell*. 7 edn, (W.W. Norton & Co, 2022).
10   Cavalier-Smith, T. Eukaryotes with no mitochondria. *Nature* **326**, 332-333 (1987).
https://doi.org:10.1038/326332a0
11   Sogin, M. L., Gunderson, J. H., Elwood, H. J., Alonso, R. A. & Peattie, D. A. Phylogenetic meaning
of the kingdom concept: an unusual ribosomal RNA from Giardia lamblia. *Science* **243**, 75-77
(1989). https://doi.org:10.1126/science.2911720
12   van der Giezen, M. Hydrogenosomes and mitosomes: conservation and evolution of functions. *J
Eukaryot Microbiol* **56**, 221-231 (2009). https://doi.org:JEU407 [pii] 10.1111/j.1550-
7408.2009.00407.x
13   Sagan, L. On the origin of mitosing cells. *J Theor Biol* **14**, 255-274 (1967).
14   Martin, W., Hoffmeister, M., Rotte, C. & Henze, K. An overview of endosymbiotic models for the
origins of eukaryotes, their ATP-producing organelles (mitochondria and hydrogenosomes), and
their heterotrophic lifestyle. *Biol Chem* **382**, 1521-1539 (2001).
https://doi.org:10.1515/BC.2001.187
15   Lopez-Garcia, P. & Moreira, D. The Syntrophy hypothesis for the origin of eukaryotes revisited.
Nat Microbiol **5**, 655-667 (2020). https://doi.org:10.1038/s41564-020-0710-4
16   Brown, J. R. & Doolittle, W. F. Archaea and the prokaryote-to-eukaryote transition. *Microbiol
Mol Biol Rev* **61**, 456-502 (1997). https://doi.org:10.1128/mmbr.61.4.456-502.1997
17   Ribeiro, S. & Golding, G. B. The mosaic nature of the eukaryotic nucleus. *Mol Biol Evol* **15**, 779-
788 (1998). https://doi.org:10.1093/oxfordjournals.molbev.a025983
18   Rivera, M. C., Jain, R., Moore, J. E. & Lake, J. A. Genomic evidence for two functionally distinct
gene classes. *Proc Natl Acad Sci U S A* **95**, 6239-6244 (1998).
https://doi.org:10.1073/pnas.95.11.6239
19   Koonin, E. V. *et al.* A comprehensive evolutionary classification of proteins encoded in complete
eukaryotic genomes. *Genome Biol* **5**, R7 (2004).
20   Imachi, H. *et al.* Promethearchaeum syntrophicum gen. nov., sp. nov., an anaerobic, obligately
syntrophic archaeon, the first isolate of the lineage 'Asgard' archaea, and proposal of the new
archaeal phylum Promethearchaeota phyl. nov. and kingdom Promethearchaeati regn. nov. *Int J
Syst Evol Microbiol* **74** (2024). https://doi.org:10.1099/ijsem.0.006435
21   Tamarit, D. *et al.* Description of Asgardarchaeum abyssi gen. nov. spec. nov., a novel species
within the class Asgardarchaeia and phylum Asgardarchaeota in accordance with the SeqCode.
Syst Appl Microbiol **47**, 126525 (2024). https://doi.org:10.1016/j.syapm.2024.126525

22　Wu, F. *et al.* Unique mobile elements and scalable gene flow at the prokaryote-eukaryote
boundary revealed by circularized Asgard archaea genomes. *Nat Microbiol* **7**, 200-212 (2022).
https://doi.org:10.1038/s41564-021-01039-y
23　Valentin-Alvarado, L. E. *et al.* Complete genomes of Asgard archaea reveal diverse integrated
and mobile genetic elements. *Genome Res* **34**, 1595-1609 (2024).
https://doi.org:10.1101/gr.279480.124
24　Rodrigues-Oliveira, T. *et al.* Actin cytoskeleton and complex cell architecture in an Asgard
archaeon. *Nature* **613**, 332-339 (2023). https://doi.org:10.1038/s41586-022-05550-y
25　Tran, L. T., Akil, C., Senju, Y. & Robinson, R. C. The eukaryotic-like characteristics of small GTPase,
roadblock and TRAPPC3 proteins from Asgard archaea. *Commun Biol* **7**, 273 (2024).
https://doi.org:10.1038/s42003-024-05888-1
26　Vosseberg, J. *et al.* Timing the origin of eukaryotic cellular complexity with ancient duplications.
Nat Ecol Evol **5**, 92-100 (2021). https://doi.org:10.1038/s41559-020-01320-z
27　Richards, T. A. *et al.* Reconstructing the last common ancestor of all eukaryotes. *PLoS Biol* **22**,
e3002917 (2024). https://doi.org:10.1371/journal.pbio.3002917
28　Lombard, J., Lopez-Garcia, P. & Moreira, D. The early evolution of lipid membranes and the
three domains of life. *Nat Rev Microbiol* **10**, 507-515 (2012).
https://doi.org:10.1038/nrmicro2815
29　Moreira, D. & Lopez-Garcia, P. Symbiosis between methanogenic archaea and delta-
proteobacteria as the origin of eukaryotes: the syntrophic hypothesis. *J Mol Evol* **47**, 517-530
(1998). https://doi.org:10.1007/pl00006408
30　Krupovic, M., Dolja, V. V. & Koonin, E. V. The virome of the last eukaryotic common ancestor and
eukaryogenesis. *Nat Microbiol* **8**, 1008-1017 (2023). https://doi.org:10.1038/s41564-023-01378-
y
31　Imachi, H. *et al.* Isolation of an archaeon at the prokaryote-eukaryote interface. *Nature* **577**,
519-525 (2020). https://doi.org:10.1038/s41586-019-1916-6
32　Lopez-Garcia, P. & Moreira, D. Cultured Asgard Archaea Shed Light on Eukaryogenesis. *Cell* **181**,
232-235 (2020). https://doi.org:10.1016/j.cell.2020.03.058
33　Gabaldon, T. Relative timing of mitochondrial endosymbiosis and the "pre-mitochondrial
symbioses" hypothesis. *IUBMB Life* **70**, 1188-1196 (2018). https://doi.org:10.1002/iub.1950
34　Gabaldon, T. Origin and Early Evolution of the Eukaryotic Cell. *Annu Rev Microbiol* **75**, 631-647
(2021). https://doi.org:10.1146/annurev-micro-090817-062213
35　Pittis, A. A. & Gabaldon, T. Late acquisition of mitochondria by a host with chimaeric prokaryotic
ancestry. *Nature* **531**, 101-104 (2016). https://doi.org:10.1038/nature16941
36　Soucy, S. M., Huang, J. & Gogarten, J. P. Horizontal gene transfer: building the web of life. *Nat
Rev Genet* **16**, 472-482 (2015). https://doi.org:10.1038/nrg3962
37　Artamonova, II, Lappi, T., Zudina, L. & Mushegian, A. R. Prokaryotic genes in eukaryotic genome
sequences: when to infer horizontal gene transfer and when to suspect an actual microbe.
Environ Microbiol **17**, 2203-2208 (2015). https://doi.org:10.1111/1462-2920.12854
38　Keeling, P. J. Horizontal gene transfer in eukaryotes: aligning theory with data. *Nat Rev Genet*
25, 416-430 (2024). https://doi.org:10.1038/s41576-023-00688-5
39　Wittmers, F. *et al.* Symbionts of predatory protists are widespread in the oceans and related to
animal pathogens. *Cell Host Microbe* **33**, 182-199 e187 (2025).
https://doi.org:10.1016/j.chom.2025.01.009
40　Steinegger, M. *et al.* HH-suite3 for fast remote homology detection and deep protein
annotation. *BMC Bioinformatics* **20**, 473 (2019). https://doi.org:10.1186/s12859-019-3019-7
41　Strimmer, K. & Rambaut, A. Inferring confidence sets of possibly misspecified gene trees. *Proc
Biol Sci* **269**, 137-142 (2002). https://doi.org:10.1098/rspb.2001.1862

42   Kanehisa, M., Furumichi, M., Tanabe, M., Sato, Y. & Morishima, K. KEGG: new perspectives on
genomes, pathways, diseases and drugs. *Nucleic Acids Res* **45**, D353-D361 (2017).
https://doi.org:10.1093/nar/gkw1092
43   Edgar, R. C. Muscle5: High-accuracy alignment ensembles enable unbiased assessments of
sequence homology and phylogeny. *Nat Commun* **13**, 6968 (2022).
https://doi.org:10.1038/s41467-022-34630-w
44   Minh, B. Q. *et al.* IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in
the Genomic Era. *Mol Biol Evol* **37**, 1530-1534 (2020). https://doi.org:10.1093/molbev/msaa015
45   Al Jewari, C. & Baldauf, S. L. Conflict over the Eukaryote Root Resides in Strong Outliers, Mosaics
and Missing Data Sensitivity of Site-Specific (CAT) Mixture Models. *Syst Biol* **72**, 1-16 (2023).
https://doi.org:10.1093/sysbio/syac029
46   Derelle, R. *et al.* Bacterial proteins pinpoint a single eukaryotic root. *Proc Natl Acad Sci U S A*
112, E693-699 (2015). https://doi.org:10.1073/pnas.1420657112
47   Parks, D. H. *et al.* GTDB: an ongoing census of bacterial and archaeal diversity through a
phylogenetically consistent, rank normalized and complete genome-based taxonomy. *Nucleic*
Acids Res **50**, D785-D794 (2022). https://doi.org:10.1093/nar/gkab776
48   Makarova, K. S., Wolf, Y. I., Mekhedov, S. L., Mirkin, B. G. & Koonin, E. V. Ancestral paralogs and
pseudoparalogs and their role in the emergence of the eukaryotic cell. *Nucleic Acids Res* **33**,
4626-4638 (2005). https://doi.org:33/14/4626 [pii]
10.1093/nar/gki775
49   Keeling, P. J. The number, speed, and impact of plastid endosymbioses in eukaryotic evolution.
Annu Rev Plant Biol **64**, 583-607 (2013). https://doi.org:10.1146/annurev-arplant-050312-
120144
50   Archibald, J. M. Genomic perspectives on the birth and spread of plastids. *Proc Natl Acad Sci U S*
A **112**, 10147-10153 (2015). https://doi.org:10.1073/pnas.1421374112
51   Ponce-Toledo, R. I., Lopez-Garcia, P. & Moreira, D. Horizontal and endosymbiotic gene transfer
in early plastid evolution. *New Phytol* **224**, 618-624 (2019). https://doi.org:10.1111/nph.15965
52   Paulick, M. G. & Bertozzi, C. R. The glycosylphosphatidylinositol anchor: a complex membrane-
anchoring structure for proteins. *Biochemistry* **47**, 6991-7000 (2008).
https://doi.org:10.1021/bi8006324
53   Biran, A., Santos, T. C. B., Dingjan, T. & Futerman, A. H. The Sphinx and the egg: Evolutionary
enigmas of the (glyco)sphingolipid biosynthetic pathway. *Biochim Biophys Acta Mol Cell Biol*
Lipids **1869**, 159462 (2024). https://doi.org:10.1016/j.bbalip.2024.159462
54   Stankeviciute, G. *et al.* Convergent evolution of bacterial ceramide synthesis. *Nat Chem Biol* **18**,
305-312 (2022). https://doi.org:10.1038/s41589-021-00948-7
55   Hoshino, Y. & Gaucher, E. A. On the Origin of Isoprenoid Biosynthesis. *Mol Biol Evol* **35**, 2185-
2197 (2018). https://doi.org:10.1093/molbev/msy120
56   Santana-Molina, C., Williams, T. A., Snel, B. & Spang, A. Chimeric origins and dynamic evolution
of central carbon metabolism in eukaryotes. *Nat Ecol Evol* (2025).
https://doi.org:10.1038/s41559-025-02648-0
57   Richards, T. A. & van der Giezen, M. Evolution of the Isd11-IscS complex reveals a single alpha-
proteobacterial endosymbiosis for all eukaryotes. *Mol Biol Evol* **23**, 1341-1344 (2006).
https://doi.org:10.1093/molbev/msl001
58   Cassier-Chauvat, C., Marceau, F., Farci, S., Ouchane, S. & Chauvat, F. The Glutathione System: A
Journey from Cyanobacteria to Higher Eukaryotes. *Antioxidants (Basel)* **12** (2023).
https://doi.org:10.3390/antiox12061199

59   Kassube, S. A. & Thoma, N. H. Structural insights into Fe-S protein biogenesis by the CIA
targeting complex. *Nat Struct Mol Biol* **27**, 735-742 (2020). https://doi.org:10.1038/s41594-020-
0454-0
60   Hoshino, Y. & Gaucher, E. A. Evolution of bacterial steroid biosynthesis and its impact on
eukaryogenesis. *Proc Natl Acad Sci U S A* **118** (2021). https://doi.org:10.1073/pnas.2101276118
61   Donoghue, P. C. J. *et al.* Defining eukaryotes to dissect eukaryogenesis. *Curr Biol* **33**, R919-R929
(2023). https://doi.org:10.1016/j.cub.2023.07.048
62   Hedges, S. B. The origin and evolution of model organisms. *Nat Rev Genet* **3**, 838-849 (2002).
https://doi.org:10.1038/nrg929
63   Koonin, E. V. & Yutin, N. Origin and evolution of eukaryotic large nucleo-cytoplasmic DNA
viruses. *Intervirology* **53**, 284-292 (2010). https://doi.org:000312913 [pii]
10.1159/000312913
64   Caforio, A. *et al.* Converting Escherichia coli into an archaebacterium with a hybrid heterochiral
membrane. *Proc Natl Acad Sci U S A* **115**, 3704-3709 (2018).
https://doi.org:10.1073/pnas.1721604115
65   Hoekzema, M., Jiang, J. & Driessen, A. J. M. Optimizing Archaeal Lipid Biosynthesis in Escherichia
coli. *ACS Synth Biol* **13**, 2470-2479 (2024). https://doi.org:10.1021/acssynbio.4c00235
66   Martin, W. & Koonin, E. V. Introns and the origin of nucleus-cytosol compartmentalization.
Nature **440**, 41-45 (2006). https://doi.org:nature04531 [pii]
10.1038/nature04531
67   Lopez-Garcia, P. & Moreira, D. Selective forces for the origin of the eukaryotic nucleus. *Bioessays*
28, 525-533 (2006). https://doi.org:10.1002/bies.20413
68   Padilla-Mejia, N. E., Makarov, A. A., Barlow, L. D., Butterfield, E. R. & Field, M. C. Evolution and
diversification of the nuclear envelope. *Nucleus* **12**, 21-41 (2021).
https://doi.org:10.1080/19491034.2021.1874135
69   Mans, B. J., Anantharaman, V., Aravind, L. & Koonin, E. V. Comparative genomics, evolution and
origins of the nuclear envelope and nuclear pore complex. *Cell Cycle* **3**, 1612-1637 (2004).
https://doi.org:1316 [pii]
70   Tamarit, D. *et al.* A closed Candidatus Odinarchaeum chromosome exposes Asgard archaeal
viruses. *Nat Microbiol* **7**, 948-952 (2022). https://doi.org:10.1038/s41564-022-01122-y
71   Munoz-Gomez, S. A. *et al.* Site-and-branch-heterogeneous analyses of an expanded dataset
favour mitochondria as sister to known Alphaproteobacteria. *Nat Ecol Evol* **6**, 253-262 (2022).
https://doi.org:10.1038/s41559-021-01638-2
72   Hyatt, D. *et al.* Prodigal: prokaryotic gene recognition and translation initiation site
identification. *BMC Bioinformatics* **11**, 119 (2010). https://doi.org:10.1186/1471-2105-11-119
73   Richter, D. J. *et al.* EukProt: A database of genome-scale predicted proteins across the diversity
of eukaryotes *Peer Community J* **2**, e56 (2022).
74   Huerta-Cepas, J., Serra, F. & Bork, P. ETE 3: Reconstruction, Analysis, and Visualization of
Phylogenomic Data. *Mol Biol Evol* **33**, 1635-1638 (2016).
https://doi.org:10.1093/molbev/msw046
75   Steinegger, M. & Soding, J. MMseqs2 enables sensitive protein sequence searching for the
analysis of massive data sets. *Nat Biotechnol* **35**, 1026-1028 (2017).
https://doi.org:10.1038/nbt.3988
76   Deorowicz, S., Debudaj-Grabysz, A. & Gudys, A. FAMSA: Fast and accurate multiple sequence
alignment of huge protein families. *Sci Rep* **6**, 33964 (2016). https://doi.org:10.1038/srep33964
77   Price, M. N., Dehal, P. S. & Arkin, A. P. FastTree 2--approximately maximum-likelihood trees for
large alignments. *PLoS One* **5**, e9490 (2010). https://doi.org:10.1371/journal.pone.0009490

78 UniProt, C. UniProt: the universal protein knowledgebase in 2021. *Nucleic Acids Res* **49**, D480-
D489 (2021). https://doi.org:10.1093/nar/gkaa1100
79 Chen, H., Zheng, F., Feng, X., et al. Engineering archaeal membrane-spanning lipid GDGT
biosynthesis in bacteria: Implications for early life membrane transformations. mLife. **4**:193-204
(2025) https://doi.org:10.1002/mlf2.70001
1192