The Prevalence of Killer Yeasts in the Gardens of Fungus-Growing Ants and the Discovery of a Novel Killer Toxin named Ksino.

Rodolfo Bizarria Jr.[a,b], Jack W. Creagh[a], Renato A. Corrêa dos Santos[c], Lily LGivens, Sarah A. Coss[a], Tanner J. Badigian[a], Adrian V. Chavez[a], Rim T. Tekle[a], Noah Fredstrom[d], F. Marty Ytreberg[e,f], Maitreya J. Dunham[d], Andre Rodrigues[b]*, Paul A. Rowley[a,f]*

[a]Department of Biological Sciences, University of Idaho, Moscow, Idaho, USA.

[b]Department of General and Applied Biology, São Paulo State University (UNESP), Institute of Biosciences, Rio Claro, São Paulo, Brazil.

[c]Laboratory of Computational, Evolutionary, and Systems Biology, Center for Nuclear Energy in Agriculture, University of São Paulo (USP), Piracicaba, São Paulo, Brazil

[d]Department of Genome Sciences, University of Washington, Seattle, WA 98195, USA.

[e]Department of Physics, University of Idaho, Moscow, ID, 83844, USA.

[f]Institute for Modeling Collaboration and Innovation, University of Idaho, Moscow, ID, 83844, USA.

*Corresponding authors

Abstract

Killer toxins are proteinaceous antifungal molecules produced by yeasts, with activity against a wide range of human and plant pathogenic fungi. Fungus gardens of attine ants in Brazil were surveyed to determine the presence of killer toxin-producing yeasts and to define their antifungal activities and ecological importance. Our results indicate that 10 out of 59 yeasts species isolated from fungal gardens are killer yeasts. Killer yeasts were less likely to inhibit the growth of yeasts isolated from the same environment but more effective at inhibiting yeast isolated from other environments, supporting a role for killer yeasts in shaping community composition. All killer yeasts harbored genome-encoded killer toxins lacked cytoplasmic

toxin-encoding elements (i.e., double-stranded RNA satellites and linear double-stranded
DNAs). Of all the killer yeasts associated with attine ants, *Candida sinolaborantium* (strain
LESF 1467) showed a broad spectrum of antifungal activities against 39 out of 69 57% of yeast
strains tested for toxin susceptibility. The complete genome sequence of *C. sinolaborantium*
LESF 1467 identified a new killer toxin, Ksino, with similarities in primary sequence and
tertiary structure to the *Saccharomyces cerevisiae* killer toxin named Klus. Surveys of publicly
available genome databases identified homologs of Ksino in the genomes of yeast strains of
Saccharomycetes and *Pichiomycetes*, as well as other species of Ascomycota and
Basidiomycota filamentous fungi. This demonstrates that killer yeasts can be widespread in
attine ant fungus gardens, possibly influencing fungal community composition and the
importance of these complex microbial communities for discovering novel antifungal
molecules.

Keywords: Killer yeasts, Killer toxins, Antifungals, Budding yeast, Fungus-growing ants

Importance
Attine ants perform essential ecosystem services through the harvesting of substrates for
fungiculture. The cultured fungi are a food source for attine ants. Characterizing antifungal
toxin-producing yeasts (killer yeasts) is vital to understanding how they might protect gardens
from invasion by unwanted fungal species. This study describes a new toxin named Ksino from
the yeast *Candida sinolaborantium*, a member of a new group of toxins found across many
different species of fungi. This work supports the role of killer yeasts in the ecology of
fungicultures and competition between fungi. The observed high prevalence of killer yeasts in
fungal gardens also enables the discovery of novel antifungal molecules with the potential to
be applied against disease-causing fungi.

1. Introduction

Killer yeasts were first described in *Saccharomyces cerevisiae* in the 1960s (Makower
and Bevan, 1963; Woods and Bevan, 1968), after which many strains and species of yeasts and
yeast-like fungi were also observed to secrete antifungal killer toxins (Woods and Bevan, 1968;

Golubev et al., 2002; Pfeiffer et al., 2004; Koltin and Day, 1976; Castillo and Cifuentes, 1994).

Killer toxins can have a range of antifungal activities ranging from narrow to broad

specificities, affecting cells from close or distantly related fungal species (Boynton, 2019;

Golubev, 1998; Golubev, 2006). The prevalence of killer yeasts strains in natural environments

is reported between 5% to 33% for positive killer phenotypes (Philliskirk and Young, 1975;

Stumm et al., 1977; Starmer et al., 1987; Abranches et al. 1997, Abranches et al. 1998; Trindade

et al. 2002; Pieczynska et al., 2013; Wojcik and Kordowska-Wiater, 2015) but can also be as

high as 59% in yeast associated with wine production (Hidalgo and Flores, 1994). Killer toxins

are known to be active against human and plant pathogenic yeasts (Middelbeek et al. 1980;

Hodgson et al. 1995; Buzzini et al., 2001; Giovati et al., 2018) and can control spoilage

organisms that are important for agriculture and food industry (Walker et al., 1995; Chessa et

al., 2017). The rise in acquired drug resistance and the emergence of drug-resistant fungal

pathogens justifies the search for novel antifungal killer toxins from diverse natural

environments, including insect-associated fungal communities.

Killer yeast communities have been previously reported in fungus-growing ant colonies

(Hymenoptera: tribe Attini: subtribe Attina, hereafter called 'attine ants'). Attine ants

established a long-term obligate mutualism with basidiomycete fungi in the *Agaricaceae* and

Pterulaceae families, using them as the primary source of food for their colonies (Branstetter

et al., 2017; Nygaard et al., 2016; Schultz and Brady, 2008, Möller, 1893; Weber, 1972).

According to their cultivars and foraging substrates, each species approaches fungiculture

differently, and these interactions are considered a model for studying the evolution of

mutualisms (Schultz and Brady, 2008, Schultz et al., 2015). Moreover, attine ants are key

ecological agents in the Neotropics as dominant herbivores, promoting nutrient cycling, soil

fertility, and seed dispersal (Corrêa et al., 2010; Meyer et al., 2011, Farji-brener and

Werenkraut, 2015; Sternberg et al., 2007; Leal and Oliveira, 1998). The foraging behavior of

attine ants also contributes to the presence of a large and complex community of environmental

yeasts in fungus gardens (Craven et al. 1970; Fisher et al., 1996; Pagnocca et al., 2008; Arcuri

et al., 2014; Bizarria Jr. et al., 2022). Fungus gardens are likely an important source of new

killer toxins due to the high concentration of simple sugars (Silva et al., 2003, Silva et al.,

2006), a large density of yeasts and filamentous fungi (Fisher et al., 1996; Craven et al., 1970;

Pagnocca et al., 1996; Rodrigues et al., 2008; Bizarria Jr. et al., 2022), and the hypothesized

role on garden "immunity" by yeasts (Rodrigues et al. 2009) – all key elements for species

competition (Boynton, 2019). Killer yeasts are primarily associated with the basidiomycete

fungus farmed by attine ants, the bodies of ants, and the leaves foraged by the ants (Carreiro et

al. 2002; Robledo-Leal et al. 2016; Bizarria Jr. et. al. 2023). In prior studies, killer positive

phenotypes have been identified in 5% to 11% of the interactions between yeasts derived from

attine ant fungus gardens (Carreiro et al. 2002; Bizarria Jr. et. al. 2023).

Killer toxins can modify the community composition of yeasts as an outcome of

competition or cooperation between killer yeasts, killer toxin-susceptible yeasts, and killer

toxin-resistant yeasts that are determined by spatial distribution, pH, and ploidy (Károlyi et al.,

2005; McBride et al. 2008; Wloch-Salamon et al. 2008). Killer toxin production also plays a

potential role in yeast dispersal, resource consumption among species, and invade or prevent

invasions from competitors (Boynton, 2019; Buser et al., 2021; Pintar and Starmer, 2003;

Ganter and Starmer, 1992; Travers-Cook et al., 2023). The coexistence of the different

phenotypes is known to be affected by environmental parameters, including nutrient

availability (Pintar and Starmer, 2003; Wloch-Salamon et al. 2008), cell density (Greig and

Travisano 2008; Vadasz et al. 2003), pH, and temperature (Bussey et al. 1988; McBride et al.

2008; Magliani et al. 1997), which can reflect the production and distribution of killer toxins

in spatially structured environments (Boynton, 2019; Giometto et al. 2021; Kerr et al. 2002;

Libberton et al. 2015; Wloch-Salamon et al. 2008). On the other hand, killer toxins are often

reported to play a role in interference competition in environments that enforce the interaction

between toxin-producing and neighboring susceptible cells (such as well-mixed habitats like

laboratory co-cultures and wine fermentation, changing the community structure and function,

as an outcome of competition or cooperation between killer yeasts, killer toxin-susceptible

yeasts, and killer toxin-resistant yeasts (Bussey et al., 1988; Hidalgo and Flores, 1994;

Pieczynska et al., 2016; Boynton, 2019).

Killer toxins are encoded by different genetic elements, including chromosomal genes,

extrachromosomal elements (such as non-autonomous double-stranded RNA (dsRNA)

satellites that are maintained by mycoviruses), and autonomous linear double-stranded DNAs

(dsDNA) (Bevan et al., 1973; Somers and Bevan, 1969; Magliani et al., 1997; Rowley, 2017).

DsRNA satellites have been previously found to be associated with *Hanseniaspora uvarum*,

Pichia kluyveri, *Saccharomyces* spp., *Torulaspora delbrueckii*, and *Zygosaccharomyces bailii*

(Ivannikova et al 2006, Ivannikova et al 2007; Naumov et al., 2009; Schmitt and Neuhausen,

1994; Ramírez et al., 2015; Rodríguez-Cousiño et al., 2017; Zorg et al., 1988), while dsDNA

plasmids have been identified in the genera *Babjevia, Debaryomyces*, *Kluyveromyces*,

Millerozyma, and *Pichia* (Klassen and Meinhardt 2007; Schaffrath et al. 2018; Klassen et al. 2017). The association of the killer phenotype with the presence of dsRNA satellites has been most thoroughly studied in *S. cerevisiae* (Crabtree et al., 2023; Vijayraghaven et al., 2023). For example, in a survey of 1,270 *S. cerevisiae* isolates, the frequency of killer yeasts was 50%, and 60% of these killer yeasts had dsRNA satellites (Crabtree et al., 2023). Killer toxin production by the remaining 40% of killer yeasts was closely correlated with the chromosomal killer toxin gene *KHS1*. Considering the many killer yeasts described, only a few studies have defined chromosomal genes that encode the toxins responsible for the observed antifungal effects. Chromosomally encoded killer toxins and their homologs, including K1-like (KKT killer toxins), KHR, KHS, SMKT, KP4-like, PMKT, and other *Pichia* toxins, have been described in many different species of fungi (Goto et al., 1990; Goto et al., 1991; Suzuki and Nikkuni, 1994; Belda et al., 2017, Fredericks et al. 2021, Brown 2011, Lu and Faris 2019). Despite the prevalence of genomic-encoded toxins, some dsRNA satellite-encoded toxins were observed to have genomic homologs in different Saccharomycotina lineages (Frank and Wolfe, 2009; Fredericks et al., 2021). However, the frequency of killer yeasts and the prevalence of dsRNA satellites and dsDNA extrachromosomal elements in different lineages of Saccharomycotina and yeast-like fungi is still understudied.

In this study, we explore the nature and prevalence of the killer phenotype in different Saccharomycotina lineages associated with attine ant fungus gardens, which are habitats that are rich in yeasts, shedding light on their putative ecological role in this environment (Bizarria Jr. et al. 2023). It was revealed that fungus gardens harbor diverse killer yeasts with narrow and broad antifungal activities. Unlike the well-studied killer yeasts of the *Pichia* and the *Saccharomyces* genera, attine ant-associated killer yeasts were devoid of cytoplasmic toxin-encoding elements such as linear dsDNA plasmids and dsRNA satellites (Magliani et al., 1997; Marquina et al., 2002). Therefore, to identify the gene(s) responsible for killer toxin production, a genome mining approach was used by determining the whole genome sequence of the killer yeast *Candida sinolaborantium* that displayed a broad spectrum of antifungal activities against many species of yeasts, including human pathogens. This approach identified a new killer toxin named Ksino. This novel killer toxin has homologs in both yeasts and filamentous fungi and shares a similar antifungal activity and predicted structure to an *S. cerevisiae* killer toxin named Klus. Collectively, it was found that fungus gardens of attine ants can be used to explore the ecological role of killer yeasts and discover new killer toxins for possible future application as natural antifungals.

2. Results

Fungus gardens of attine ants harbor killer yeasts. To assess the diversity of yeasts associated with fungus-growing ants and their potential for antifungal toxin production, fungus gardens of four ant fungiculture systems were surveyed from 28 nests from four cities in Brazil (Fig. 1A; Table S1). The ant species included the leaf-cutting ant *Acromyrmex coronatus*, the non-leaf-cutting ant *Mycetomoellerius tucumanus*, and the lower attines *Mycetophylax* aff. *auritus* and *Mycocepurus goeldii* that cultivate Agaricaceae fungi, and *Apterostigma goniodes* that cultivates Pterulaceae fungi (Fig. 1B). Ants were selected to broadly represent different fungiculture systems, foraging and preparation behaviors, and the yeast microbiota associated with fungus gardens. Yeasts were isolated from fungus gardens by culturing on synthetic growth media. Microsatellite amplification, sequencing of the D1/D2 domain of the large subunit ribosomal RNA gene, and phylogenetic reconstruction were used for taxonomy. The 180 isolated yeast strains comprised 59 species belonging to eight families and six orders from the Saccharomycotina subphylum previously characterized in a large study of yeasts and yeast-like fungi associated with fungus gardens of attine ants (Fig. 1C; Table S1) (Bizarria Jr. et al., 2023). Of the yeasts isolated, 91 were associated with *Acromyrmex coronatus*, 13 with *Mycetomoellerius tucumanus*, 42 with *Mycetophylax* aff. *auritus*, 25 with *Mycocepurus goeldii*, and nine with *Apterostigma goniodes* (Table S1) (Bizarria Jr. et al., 2023).

FIG 1. Fungus gardens of different attine ants harbor different killer yeasts. **A.** Images of fungus gardens of different ant species from which yeasts were isolated (arrowheads). The gardens of the leaf-cutting ant *Acromyrmex coronatus* (circle), the non-leaf-cutting ant *Mycetomoellerius tucumanus* (square), the lower attines *Mycetophylax* aff. *auritus* (triangle), and *Mycocepurus goeldii* (star). The fungus garden of *Apterostigma goniodes* is not represented. **B.** Representative images of attine ant species: *Acromyrmex coronatus* (CASENT0173791, April Nobile) (circle); *Mycetomoellerius tucumanus* (CASENT0909391, Will Ericson) (square); *Mycetophylax auritus* (CASENT0901666, Ryan Perry) (triangle), *Mycocepurus goeldii* (CASENT0173988, April Nobile) (star), and *Apterostigma goniodes* (CASENT0922040, Michele Esposito) (diamond). The scale bar on all images represents 0.2 mm. Photos of ants were downloaded from "www.antweb.org." **C.** Phylogenetic relationship of yeasts isolated from fungal gardens as inferred by maximum likelihood and the rRNA gene. Symbols in tree leaves indicate the isolation source of yeasts, as depicted in panels A and B. Red shading and skulls denote the species confirmed as killer yeasts. The phylogenetic clades were highlighted by order following the updated classification

From the 59 yeast species identified that were associated with attine ant gardens, ten
species had isolates that were able to produce killer toxins (10 out of 59; 17% of killer yeasts)
(Fig. 1C). Killer toxin production was scored by the absence of growth around a killer yeast or
the staining of the lawn yeast with the redox indicator methylene blue (as an indicator of cell
death) at pH 4.6. This pH was used as many previously described killer toxins are active in
these conditions and it is close to the pH observed in fungus gardens where these yeasts were
isolated (pH 5.1-5.4). The identified killer yeasts were effective at inhibiting strains that were
either previously determined to be susceptible to killer toxins (Fredericks et al., 2021), human
pathogenic yeasts, or yeasts randomly selected from fungus gardens and other sources (Fig. 2A
and S1; File S1). From the attine ant gardens, *Candida blattae*, *Candida sinolaborantium*,
Candida sp. (closest to *Candida temnochilae*), *Candida* sp. (closest to *Candida
membranifaciens*), *Diutina catenulata*, *Jamesozyma spencerorum*, *Kodamaea ohmeri*,
Schwanniomyces vanrijiae, *Wickerhamomyces ciferrii*, and *Wickerhamomyces* sp. (closest to
Wickerhamomyces rabaulensis) were identified as killer yeasts. These yeasts had unique
antifungal activities when compared to other previously identified *Saccharomyces* killer toxins
(K1, K1L, K2, K21, K28, K45, K62, K74, and Klus), as has been observed in a previous screen
for killer yeasts of *S. cerevisiae* (Crabtree et al., 2023). Killer yeasts were only absent from the
fungus gardens of *M. goeldii*, but were recovered from the fungus gardens of *Acromyrmex
coronatus* (7% killer yeasts, 6 out of 91 isolates), *Mycetomoellerius tucumanus* (46%, 6 out of
13), *Mycetophylax* aff. *auritus* (43%, 18 out of 42), and *Apterostigma goniodes* (22%, 2 out of
9) (Fig. 2B). Thirty-two out of 180 (18%) yeasts from attine ants were also capable of inhibiting
the growth of susceptible yeasts, including human pathogenic yeasts such as *Candida albicans*,
Candida glabrata, and *Candida auris* (Fig. S1). The attine ant-associated killer yeasts *J.
spencerorum* LESF 1468 and *S. vanrijiae* LESF 1521 could only inhibit the growth of
pathogenic yeasts (Fig. S1, File S1). Many killer yeasts from the same species had similar
antifungal activities as demonstrated by their phenotypic clustering (see *Candida* sp. closest to
C. temnochilae, *C. blattae*, *D. catenulata*, *S. vanrijiae*, *Wickerhamomyces* sp. closest to *W.
rabaulensis*, and *W. ciferrii*) (Fig. 2A). Specifically, *Candida blattae* killer yeasts had similar
antifungal activities even when isolated from the fungicultures of different ant species
(compare *Candida blattae* isolates from *Mycetophylax* aff. *auritus* and *Apterostigma goniodes*)

(Fig. 2A). However, there were isolates of *S. vanrijiae* killer yeasts from the fungiculture of
Mycetophylax aff. *auritus* with strikingly different spectra of antifungal activities, which
suggests that isolates of the same species were not always clonal. *Candida sinolaborantium*
(strain LESF 1467) had the broadest range of antifungal activity, inhibiting 57% of all 69 lawn
strains tested (Fig. 2A and S1, File S1). Comparing interactions between ant-associated killer
yeasts and susceptible lawn strains from fungus gardens and other environmental sources found
that growth inhibition was more prevalent for yeasts from different sources (Chi-squared test,
p-value < 0.01) (Fig. 2C; Table S2). Specifically, only 1% (6 of 647) of interactions between
ant-associated yeasts from the same type of fungiculture resulted in growth inhibition. In
contrast, 4% (301 of 8,353) of interactions between ant-associated killer yeasts and yeasts from
fungus gardens and other environmental sources isolation sources resulted in growth inhibition.
239

The ability to produce killer toxins is often associated with the presence of linear
dsDNA plasmids or mycovirus-associated dsRNA satellites. We assayed all attine ant-
associated killer yeasts for the presence of these genetic elements using solvent extraction of
nucleic acids, cellulose chromatography, and gel electrophoresis (Crabtree et al., 2019). In
particular, the use of cellulose chromatography selectively enriches dsRNAs derived from
mycoviruses and satellites, reducing the false negative rate compared to solvent extraction
alone and without the bias of reverse-transcriptase PCR (Crabtree et al., 2023). However, we
did not find that any of the isolated attine ant-associated strains possess dsRNAs (i.e., viruses
and satellites) or dsDNA cytoplasmic elements (i.e., plasmids). Double-stranded RNA viruses
have been found in many different species of yeasts without satellite dsRNAs (Crucitti et al.,
2021; Lee et al., 2022; Taylor et al., 2013); thus we broadened the survey to include all attine-
associated yeasts but again did not identify any dsRNA viruses. This suggested that the killer
toxins associated with yeasts from fungus gardens are genome-encoded and that dsRNA
viruses, dsRNA satellites, and DNA plasmids were not observed in yeasts associated with attine
ant species in the geographical area sampled.
255

FIG 2. Killer yeasts from fungus gardens have unique spectra of antifungal activity. A. Heatmap with the interactions between canonical *Saccharomyces* killer yeasts (K1, K1L, K2, Klus, K21, K28, K45, K62, K74), killer yeasts associated with ants, and killer toxin susceptible strains. Origins of the ant-associated killer yeasts; *Acromyrmex coronatus* (circle); *Mycetomoellerius tucumanus* (square); *Mycetophylax* aff. *auritus* (triangle), *Apterostigma goniodes* (diamond). Killer toxin activity was qualitatively assessed based on the presence and size of growth inhibition zones and/or methylene blue staining around killer yeasts (representative images of these phenotypes are shown). Darker colors on the heatmap represent a more prominent killer phenotype, with yellow indicating no observable killer phenotype. Clusters on the dendrograms connecting individual killers or susceptible yeasts indicate similar susceptibilities to killer toxins or antifungal activities. **B.** Number of killer yeasts and non-killers associated with the different attine ant fungicultures. **C**. Number of killer toxin-positive and negative interactions between yeast from attine ant environment and susceptible strains (Table S1.), growth inhibition by killer yeasts is different among the same or different fungicultures and locations.

274
FIG 3. The antifungal activities of *Candida sinolaborantium*. A. Growth phenotypes associated with the co-
culture of the killer yeast *C. sinolaborantium* with *K. ohmeri* at room temperature (RT) and pH 4.6. Cell
morphology of *K. ohmeri* inside (+Halo) and outside (-Halo) the white halo around *C. sinolaborantium* was
evaluated by microscopy. **B**. The antifungal activity of protein precipitates derived from the spent growth medium
of *C. sinolaborantium* and the heat stability of these activities at 60°C and 98°C. Killer yeast activity of *C.
sinolaborantium* under different pH and temperature conditions against **C.** *K. ohmeri* and *C. castelli*. For all pH
and temperature data, the means were significantly heterogeneous (one-way anova, P<0.01). Means with the same
letter are not significantly different from each other (Tukey–Kramer test, P>0.05). **E**. Proportion of elongated to
normal cells of *K. ohmeri* inside (+Halo) and outside (-Halo) the white halo around *C. sinolaborantium*, as
depicted in Fig. 3A. **F**. The same growth areas as shown in **E**. but colony forming units (CFU) were measured to
indicate *K. ohmeri* cell survival. Asterisks indicate different means for the Welch Two Sample t-test (p-value <
0.01). All error bars are standard error (n>3).

Characterization of the killer yeast *Candida sinolaborantium*. *C. sinolaborantium* was

identified as a killer yeast with a broad antifungal spectrum that could inhibit 39 of the 69 yeast

strains tested (Fig. 2 and S1). In contrast, the next most potent killer yeasts could only inhibit

11 strains tested. *C. sinolaborantium* was also able to cause large zones of growth inhibition in

many different species of yeasts that were resistant to other killer yeasts (Fig. 2A). After several

days of incubation, we observed that *K. ohmeri* produced a striking white halo between a zone of growth inhibition and methylene blue staining when challenged by the killer yeast *C. sinolaborantium* (Fig. 3A). This phenotype was unique to the pairing of *C. sinolaborantium* and *K. ohmeri*, and was distinct from the growth inhibition caused by the killer toxin K62 (the only other known killer toxin capable of inhibiting *K. ohmeri*) (Fig. 2A). The white halo also appeared to be raised from the surface of the agar plate (Fig. S2). When cells in this area (+halo) were observed under a microscope, they appeared to be elongated compared to cells of *K. ohmeri* outside of the halo (-halo) (Fig. 3A). The killer toxin (or toxins) secreted by *C. sinolaborantium* were precipitated by ethanol and remained active when incubated at room temperature, but were inactivated by heating to 60°C and 98°C (Fig. 3B). The inhibition of *K. ohmeri* by *C. sinolaborantium* was optimal at pH 5 (Fig. 3C, Table S3), with loss of killing at pH >5.5. The optimal temperature for killing *K. ohmeri* by *C. sinolaborantium* was 20°C. Using *C. castelli* as a sensitive lawn indicated a wider optimum temperature (17-25°C) and pH (3.5-5.0) (Fig. 3D, Table S3 and S4). In both cases the there was no killer activity at 35°C (Fig. 3C and 3D, Table S4). Further analysis of the white halo of *K. ohmeri* revealed an 8-fold increase in cell elongation in the white halos (+halo) when compared to untreated cultures of *K. ohmeri* (-halo) (Fig. 3E and Table S5). Cells of *K. ohmeri* from the white halo or untreated culture were seeded on agar to measure viability by colony-forming units. Cells isolated from the white halos showed a 2.1-fold reduction in colony-forming units compared to *K. ohmeri* isolated away from halos (Fig. 3F and Table S6). Ultrafiltration of culture media was used to determine the approximate molecular weight of the antifungal molecules produced by *C. sinolaborantium*. Antifungal activity was observed for fractions that enriched molecules >100 kDa and >30 kDa. Raw filtrates and concentrated ethanol-precipitated fractions were used to challenge a panel of yeasts previously shown to be susceptible to toxins produced by *C. sinolaborantium* (Fig. 2). The spent culture medium showed less antifungal activity compared to the precipitated medium, with the latter capable of inhibiting 21/29 susceptible yeasts that were inhibited in killer assays on agar (Fig. S3). Antifungal activities were similar before and after passage through a filter with a molecular weight cut-off (MWCO) of 100 kDa and were mostly retained in the MWCO 30 kDa filter. Only weak antifungal activity was detectable for the fraction that passed through the MWCO 30 kDa filter (Fig. S3). The appearance of white halos was observed using fractions captured by filters with MWCO 100 kDa and MWCO 30 kDa when tested against *K. ohmeri* but not with MWCO 5 kDa. Together, these data show that *C. sinolaborantium* is a killer yeast with a broad spectrum of antifungal activities, with similar temperature and pH optima of other previously described yeast killer toxins, but with the ability

to induce cell elongation in *K. ohmeri*. Fractions from ultrafiltration are consistent with the

antifungal activities of *C. sinolaborantium,* suggesting either the production of multiple

chromosomally encoded killer toxins with similar antifungal properties or perhaps large

oligomers or complexes of a single >30 kDa killer toxin.

Ksino: a novel killer toxin produced by *C. sinolaborantium* with structural homology to

the Klus killer toxin. To determine the gene or genes responsible for killer toxin production

by *C. sinolaborantium*, the genome sequence of the yeast was determined (Table S7). The

resulting 11.21 Mb assembly resulted in a BUSCO completeness (Complete and single-copy

BUSCOs) of around 98% using the saccharomycetes_odb10 dataset (Seppey et al., 2019), with

GC content of 49%, an N50 of 155,979 bp, and a total of 5,951 protein-coding genes (Table

S7). To identify genome-encoded killer toxins in the *C. sinolaborantium* proteome, BLASTp

was performed using a database of known killer toxins from yeasts (Table S8). This approach

identified 12 toxin-like candidates, all with a sequence alignment length greater than 100 amino

acids and an identity of at least 24% compared to any known killer toxin (Table S9). One *C.*

sinolaborantium protein had a 28% identity and 42% similarity over an alignment length of

151 amino acids with the *S. cerevisiae* killer toxin Klus and was named Ksino (Genbank

Accession: XAT76252) (Table S9) (Rodríguez-Cousiño et al. 2011).

FIG 4. Ksino from *C. sinolaborantium* defines a novel family of killer toxins. **A.** The phenotypes associated with the co-culture of the Klus-encoding killer yeast *S. cerevisiae* strain DSM-70459 with *K. ohmeri* (+Mlus). Treatment with cycloheximide (CHX) causes the loss of the Mlus satellite (-Mlus) and the loss of the white halo phenotype. The ratio of **B.** elongated and **C.** viable *K. ohmeri* cells in the white halo around Klus-producing *S. cerevisiae* (+Halo) and in axenic culture (-Halo). Asterisks indicate different means for Welch Two Sample t-test (p-value <0.01), ns = not significant. **D.** A maximum likelihood phylogeny of Ksino homologs identified in diverse fungi. The Ksino from *C. sinolaborantium* is highlighted in bold, and all Saccharomycotina yeasts are highlighted in blue. The dsRNA-encoded killer toxin Klus is represented as an outgroup. Bootstrap values higher than 70 are shown. All error bars are standard error (n>3).

The sequence similarity between Ksino and Klus prompted a more detailed investigation into whether these killer toxins share similar antifungal properties. Although we had previously failed to detect growth inhibition of *K. ohmeri* by *S. cerevisiae* killer toxin Klus (Fig. 3), we noted the appearance of white halos instead of a zone of growth inhibition or methylene blue staining after prolonged co-culture (Fig. 4A). The appearance of white halos was similar to those produced by *C. sinolaborantium* and was dependent on the presence of the Mlus satellite dsRNA that encodes the Klus killer toxin gene (Fig. 4A). Analysis of the cells within the white halo found a significant proportion of elongated cells, as previously observed upon exposure of *K. ohmeri* to Ksino (Fig. 4B). However, unlike the Ksino halos, Klus did not cause a significant loss of cell viability (Fig. 4C).

To identify other possible Ksino-homologous proteins in different organisms, we

performed a BLASTp search across the NCBI database, resulting in 84 protein sequences from

various Ascomycota and Basidiomycota fungi. Ksino homologs were found in

Saccharomycetes, Pichiomycetes yeasts, and other filamentous fungi with identity and

alignment lengths higher than 30% and 109 amino acids, respectively (Fig. 4D and Table S11).

Phylogenetic analysis of homologs, with Klus as an outgroup, indicates that proteins with the

highest identity of Ksino (35-42% identity) were all uncharacterized genes of the

Saccharomycotina yeasts from the lineages of *Debaryomycetaceae* and the

Saccharomycetaceae (Fig. 4D). Saccharomycotina species with Ksino-like genes included

Candida anglica, *Kazachstania africana*, *Scheffersomyces stipitis*, and *Suhomyces*

tanzawaensis, with both *K. africana* and *S. stipitis* (Basionym: *Pichia stipitis*) being previously

reported as killer yeasts (Antunes and Aguiar, 2012; Fredericks et al. 2021). Filamentous fungi

encode the majority of Ksino homologs (75 out of 84), including species belonging to the

genera *Fusarium*, *Mycena*, and *Aspergillus* (Fig. 4D and Table S11), known as plant and human

pathogens (*Fusarium* spp.), saprophytic and biotrophic plant associated fungi (*Mycena* spp.),

and fungi species that are ubiquitous in the environment (*Aspergillus* spp.) (Summerell et al.

2010, Harder et al. 2023; Thoen et al. 2020; Houbraken et al. 2020; Jurjevic et al. 2012;

Peterson et al. 2001).

The Ksino open reading frame was determined to be 202 amino acids long, comparable

to Klus at 242 amino acids. Linear representation of the secondary structure revealed a similar

pattern of alpha helices and beta sheets (Fig. 5A; Fig. S5). There was also evidence of potential

post-translational modification sites, including N-terminal signal sequence cleavage sites

predicted by psipred (Buchan et al., 2013) and dibasic motifs (i.e., KR) in the unstructured

region before helix 1 (Fig. S5). Dibasic motifs in killer toxins are known to be proteolytically

cleaved by the Golgi-specific protease Kex2 (Julius et al., 1984). As Ksino has no sequence

homology to a protein of known structure, AlphaFold2 was used to generate tertiary structure

models of Ksino and Klus (Jumper et al., 2021). Ksino and Klus had global LDDT scores of

63.6 and 65.1, respectively, and per residue, LDDT was greater than 80 in regions of secondary

structure for both proteins (Fig. 5A and S5). Molecular dynamics simulation improved the bond

angles of the alphafold model to Ramachandran favored regions in models of Ksino (77% to

91%) and Klus (73% to 89%) (Table S10, Fig. S6). Ksino and Klus simulations reached a stable

RMSD of around 1.6 nm after approximately 20 ns of simulation (Fig. S6). Molprobity scores,

analogous to structure resolution, were determined to be 1.54 for Ksino and 1.86 for Klus

(Studer et al., 2019). Overlay of the final energy-minimized structures of Ksino with Klus

compared to the original AlphaFold2 models yielded an RMSD of 2.6 Å (Fig. 5B). The

predicted organization of the Ksino structure consists of a discontinuous five-stranded

antiparallel beta-sheet packed against a pair of antiparallel alpha helices (Fig. 5B and S5). The

tertiary structure of Klus follows the same overall organization but with one less beta-turn.

Ksino has eight cysteine residues, compared to the seven in Klus (Fig. S5). The predicted

structures have three intramolecular disulfide bonds. In Ksino, C105-C127 is positioned to

crosslink the C-terminus of helix 1 to the beta-sheet 2, similar to C141-C162 in Klus (Fig. S5).

The remaining predicted disulfide bonds are unique to both structural models (Fig. S5). These

simulations collectively suggest that the Klus and Ksino proteins share significant structural

homology despite having low amino acid identity.

Figure 5. Molecular modeling of Ksino indicates structural homology to Klus. pLDDT confidence per residue

and predicted disulfide bond arrangement of AlphaFold2 models of Ksino and Klus. A. Secondary structure

representations aligned with per-residue pLDDT (predicted local distance difference test) scores from

AlphaFold2. Alpha helices are represented as blue rectangles, beta sheets as red arrows, and unstructured regions

as orange lines. Predicted disulfide linkages are indicated as horizontal dashed lines. Scissors with vertical dashed

lines represent predicted signal sequence cleavage sites. B. Ramachandran plots of general residues (non-

proline/glycine) generated by SWISS structure assessment tool demonstrating improved Ramachandran favored

after performing 100 ns simulation. The unrelaxed structure represents AlphaFold2's raw output (Ramachandran

et al., 1963). **B.** Full protein RMSD over 100 ns molecular dynamics simulation. GROMACS was used to generate

alignments of each snapshot back to the structure at 0 ns. C. Ksino and Klus AlphaFold2 models with cysteine

residues highlighted as yellow spheres to illustrate predicted disulfide linkage arrangement. The cartoon helices

and beta sheets are colored to match panel A.

The molecular weight of Ksino was predicted to be 21.5 kDa, which appeared to be
smaller than the predicted molecular weight determined by ultrafiltration (Figure S3). To
determine if Ksino was was expressed by *C. sinolaborantium*, total RNA was extracted from
laboratory cultured yeast cells in order to assay for the presence of Ksion transcript by reverse
transcriptase PCR (RT-PCR). This approach was successful in detecting the presence of Ksino
RNA transcripts (Figure S4). To confirm that Ksino is an active killer toxin, the gene was
amplified by PCR and cloned into an *S. cerevisiae* expression vector under the control of a
galactose-inducible promoter. *C. sinolaborantium* belongs to the Serinales order and uses the
CUG codon to encode serine (Groenewald et al 2023; Krassowski et al., 2018). Ksino has one
CUG codon that will introduce a leucine at position 186 when expressed by *S. cerevisiae*. This
mutation is predicted to have a small destabilizing effect on folding compared to the wild type
based on FoldX 5.0 and molecular dynamics simulation (estimated $\Delta\Delta G_{Folding}$ = 1.0 kcal/mol)
(Fig. 5). Therefore, residue 186 was recoded for serine in *S. cerevisiae* (L186S). When the
strains were grown on dextrose or galactose containing media, no growth inhibition of *K.
ohmeri* was observed during co-culture with a laboratory strain of *S. cerevisiae* transformed
with the wild type or L186S expression vectors (Fig. SXX). However, challenging the strain
C. castelli Y-17070 with Ksino wild type or L186S on galactose triggered large zones of
inhibition and methylene blue staining, indicating that Ksino is a novel killer toxin (Figure 6A).
Growth of *S. cerevisiae* under different temperatures and pH identified that Ksino has an
optimum antifungal activity at pH4-4.5 (Figure 6B and Table S13) and 20-25°C (Figure 6C
and Table S14), which was narrower than the killer phenotype of *C. sinolaborantium* (Figure
3). The inability of Ksino expressed by *S. cerevisiae* to prevent the growth of *K. ohmeri* and
differences in the optimal conditions for antifungal activity on *C. castelli*, indicated that Ksino
may be one of many antifungal molecules produced by *C. sinolaborantium*, or that expression
of Ksino by *S. cerevisiae* alters the toxins spectrum of activity (Fig. 4F and S8).

Figure 6. Ksino is an antifungal killer toxin when expressed by *S. cerevisiae*. A. Recombinant expression of Ksino (L186S) by *S. cerevisiae* using galactose induction with a lawn of *C. castelli* Y-17070 (pH 4.5). Dextrose was used as a non-induced control. n/a indicates an empty vector negative expression control. Zones of growth inhibition were compared to the killer phenotype of *C. sinolaborantium*. B. Killer yeast activity under different **B**. pH and **C**. temperature conditions as measured by zone of growth inhibition and methylene blue staining. Representative images of the killer phenotype are displayed under each histogram. The means were significantly heterogeneous (one-way anova, P<0.01) for both the pH and temperatures tested. Means with the same letter are not significantly different from each other (Tukey–Kramer test, P>0.05). All error bars are standard error (n=3).

3.   Discussion

The large number of killer yeasts in fungicultures suggests they might play a role in attine ant fungus gardens by suppressing fungal competitors or allelopathy effects on complex dynamics of microbial interactions. Different studies have revealed that yeasts are abundant in attine ant environments that likely represent differences in the foraging composition and

substrate preparation behaviors observed among the different fungicultures (Bizarria Jr. et al.,

2022, de Fine Licht and Boomsma, 2010; Ronque et al., 2019). Yeasts from fungus gardens of

the leaf-cutting ant *Atta texana* were observed to suppress the growth of fungal garden

contaminants, especially *Escovopsis* and *Syncephalastrum* (Rodrigues et al. 2009). These

antifungal properties were hypothesized to contribute to fungal garden "immunity," whereby

the resident fungi within a fungiculture prevent invasion by other fungal species. This would

require interplay between the cleaning behaviors of attine ants in their gardens to remove

unwanted fungi and other microbial interactions within attine ant colonies, i.e., allelopathy.

Our study indicates that fungus gardens of attine ants harbor killer yeasts with similar

prevalence to other surveys (Philliskirk and Young, 1975; Pieczynska et al., 2013; Starmer et

al., 1987; Stumm et al., 1977; Trindade et al. 2002; Wojcik and Kordowska-Wiater, 2015;

Abranches et al. 1997, Abranches et al. 1998). This expands the previous findings regarding

the prevalence of killer yeasts in laboratory-reared attine ant colonies of *Atta sexdens* (Carreiro

et al., 2002). Moreover, the prevalence of killer yeasts among the different ant species, such as

Mycetophylax aff. *auritus* (43%, eight nests sampled) and *Mycetomoellerius tucumanus* (46%,

four nests sampled) were considerably higher than in other surveyed fungicultures.

Killer yeasts from attine ant colonies were more likely to inhibit the growth of

susceptible strains from different fungicultures. This supports previous observations of killer

and non-killer yeast interactions whereby local selective pressure for killer toxin resistance

allows the coexistence of killer and non-killer yeasts in a given niche (Starmer et al. 1987;

Ganter and Starmer, 1992, Abranches et al. 1997, Trindade et al. 2002). The susceptibility of

yeast populations that have never been exposed to a specific killer toxin would greatly depend

on the standing genetic variation (perhaps in the cell wall and membrane toxin receptors)

among different lineages of yeasts dictated by the population's evolutionary history. Our

findings support the hypothesis that yeasts contribute to fungal garden "immunity" since killer

yeasts are expected to prevent the invasion of other fungal species into the fungus garden. The

coexistence of killer and non-killer yeast species could be an outcome of microhabitats in the

spatially structured environment of the fungus garden, which may harbor different populations

of yeasts and allow the colonization by killer yeasts (Giometto et al. 2021, Wloch-Salamon et

al. 2008).

Saccharomycotina yeast species isolated in attine ant gardens have been previously

found to secrete killer toxins. Among the yeast species, the ability to produce toxins was

observed for *Diutina catenulata* (Basionym: *Candida catenulata*; Stopiglia et al., 2014; Fuentefria et al., 2007), *Kodamaea ohmeri* (Synonym: *Pichia ohmeri*; Fuentefria et al., 2006, Fuentefria et al., 2008; Coelho et al., 2009), *Schwanniomyces vanrijiae* (Madbouly et al., 2020), and *Wickerhamomyces ciferrii* (Basionym: *Hansenula ciferrii*; Nomoto et al., 1984). The exception was *Jamesozyma spencerorum* (Synonym: *Kazachstania spencerorum* nom. inv.), which, to our knowledge, is the first report of a killer phenotype displayed by this species, although toxin production has been previously reported in the genus, specifically *K. africana*, *K. exigua* and *K. unispora* (Perez et al., 2016; Stopiglia et al., 2014; Fredericks et al., 2021). These yeast species have also been identified in substrates that are commonly foraged or associated with attine ants, including plants (*Schwanniomyces vanrijiae*, Madbouly et al., 2020; *Kodamaea ohmeri*, Fuentefria et al., 2006), insects and their environments (*Candida blattae*, Nguyen et al., 2007; *Candida sinolaborantium*, Suh et al., 2005; Guamán-Burneo et al., 2015; *Schwanniomyces vanrijiae*, Maksimova et al., 2016; *Kodamaea ohmeri*, Benda et al., 2008; Amos et al., 2019; Suh and Blackwell, 2005), and soil (*Jamesozyma spencerorum*, Synonym : *Kazachstania spencerorum* nom. inv.; Vaughan-Martini et al., 1995; Kurtzman, 2003; Kurtzman and Robnett, 2003; Liu et al., 2024).

Importantly, killer yeast species appear to be unique between colonies of the same ant species (Bizarria Jr. et al., 2023). This observed diversity could be due to the foraging behavior of ants, whereby sampling of the surrounding environment would introduce different fungal species to the fungiculture of the same species of ants in different geographical locations (Bizarria Jr. et al., 2023). However, despite the potential role of killer yeasts in garden immunity, whether the occurrence of killer yeasts in a fungiculture is acquired by chance, inherited from the environment, or inherited during garden propagation remains unknown (Bizarria Jr. et al., 2022). Niche construction by the fungal cultivar species could also play a role in yeast diversity in the fungus garden environment or by other features such as the pH of the fungus garden. Interestingly, the pH observed in the different fungus gardens (pH 5.1-5.4) is consistent with the optimal antifungal activities of *C. sinolaborantium* (pH 3.5-5.0) and Ksino (pH 4.0-4.5). However, the association between killer yeasts and fungicultures still requires robust empirical evidence to reject the stochastic nature of killer yeast colonization. Future yeast isolation will provide new directions to clarify whether killer yeast occurrence is modulated by ant foraging choice or fungus garden features such as pH and fungal cultivar species.

Our survey of the encoding elements of killer toxins indicated that the 180 isolated
strains of Saccharomycotina yeasts were devoid of cytoplasmic killer toxin-encoding elements
such as dsRNA satellites and dsDNA linear plasmids. The frequency of dsRNA elements
associated with *Saccharomyces* yeasts has been observed to range from 10% to 51% in wild
and domesticated strains (Pieczynska et al., 2013; Crabtree et al., 2023). It is expected that
more of these genetic elements will be found in other Saccharomycotina lineages, given their
discovery in non-*Saccharomyces* yeasts and the frequency of genome-integrated molecular
fossils of plasmid and virus sequences in different lineages (Frank and Wolfe, 2009; Liu et al.,
2010; Satwika et al., 2012; Myers et al., 2020; Villan Larios, et al. 2023; Taylor and Bruenn,
2009, Lee et al., 2022). Most of the yeast isolated from fungicultures belong to the Serinales,
where the CUG codon is translated into serine instead of leucine in the universal genetic code
(Groenewald et al 2023). The rewiring of the fungal genetic code has been suggested to prevent
the invasion of viruses and plasmids by stopping the faithful translation of viral proteins
(Krassowski et al., 2018). Alternatively, RNAi might also contribute to limiting viral infections
in these yeasts, even though, in some cases, yeast lineages maintain viruses that suppress RNAi
(Drinnenberg et al., 2009; Drinnenberg et al., 2011; Lee et al., 2022; Segers et al., 2006; Segers
et al., 2007; Rodriguez Coy et al., 2022). RNAi and genome recoding could be possible reasons
for the absence of cytoplasmic elements in the yeasts associated with attine ant fungiculture.
However, this still requires a more in-depth investigation to exclude the stochastic nature of
virus acquisition by fungi and sampling biases.
556
The lack of extrachromosomal elements in killer yeasts associated with attine ant
fungiculture suggests that killer toxin genes are genome-encoded. Our survey indicates that the
killer toxins from different Saccharomycotina lineages associated with fungus gardens differ
in their spectrum of antifungal activity from canonical *Saccharomyces* killer toxins (K1, K2,
K28, etc.). In particular, toxins secreted by *Candida sinolaborantium* had the broadest activity,
including toward infectious human pathogens (e.g., *Candida albicans*, *C. glabrata*, and *C.*
auris). A unique response to the killer activity of *C. sinolaborantium* was also observed in *K.*
ohmeri, where cells exposed to the killer yeast grew as filaments. Usually, dimorphism in
yeasts has been demonstrated to be triggered by nitrogen starvation (Gimeno et al., 1992),
different carbon sources, and temperature (Rupert and Rusche, 2022). However, to our
knowledge, this is the first case of this phenotype associated with exposure to a killer yeast . The
broad spectrum antifungal activities of *C. sinolaborantium* and the ability to fractionate
antifungal activities by ultrafiltration indicates the likely presence of multiple unidentified

chromosomally encoded toxins. After determining the genomic sequence of *C.*
sinolaborantium, the new killer toxin Ksino was discovered due to its sequence homology to
the canonical *Saccharomyces* killer toxin Klus. However, the killer activity of *C.*
sinolaborantium was not fully reproduced when Ksino was heterologously expressed by *S.*
cerevisiae and was unable to inhibit or cause filamentouse growth in *K. ohmeri*. Instead robust
growth inhibition was only observed after challenging *C. castelli.* These data also supports the
expression of multiple killer toxins by *C. sinolaborantium* or that ectopic heterologous
expression by *S. cerevisiae* is inadequate due to potential deficiencies in extracellular export,
maturation, or Ksino's innate toxicity.

Many proteins from different Ascomycete and Basidiomycete fungi lineages were
identified in our search for Ksino-like proteins, including filamentous fungi and yeast (from
Saccharomycetes and Pichiomycetes). The Ksino-like proteins from yeasts of the
Saccharomycotina subphylum (*Candida anglica*, *Kazachstania africana*, *Scheffersomyces*
stipitis, and *Suhomyces tanzawaensis*) share a closer evolutionary origin to Ksino and therefore
might represent bonafide killer toxins. Despite our approach to identifying killer toxins being
limited by primary sequence similarity to known toxins, it has successfully identified Ksino.
This combination of genome sequencing, sequence homology searches, and structural
modeling could be used more widely on the many killer yeasts that are devoid of cytoplasmic
extrachromosomal elements to identify new killer toxins (Boynton, 2019; Crabtree et al. 2023).
590

In summary, this study demonstrates that (i) Fungus gardens of attine ants can harbor
killer yeasts, an underappreciated environment for killer yeasts. (ii) DsRNA and dsDNA
extrachromosomal elements are lacking from yeasts associated with attine ant fungicultures.
(iii) Genome mining can discover new chromosomally encoded killer toxins, specifically Ksino
secreted by *Candida sinolaborantium*. (iv) Ksino has sequence and structural homology to the
Saccharomyces killer toxin Klus and is a member of a new family of killer toxins in
Ascomycete and Basidiomycete fungi.

Funding sources

This work was supported by Fundação de Amparo à Pesquisa do Estado de São Paulo
(FAPESP) [scholarships No. 2019/24412–2 and 2021/09980-4] to RBJ and [grant No.

2019/03746–0] to AR. The authors also thank the Conselho Nacional de Desenvolvimento

Científico e Tecnológico (CNPq) for a research fellowship [grant No. 305269/2018–6] to AR

and a scholarship [No. 142396/2019–2] to RBJ. This work was supported by the Institute for

Modeling Collaboration and Innovation at the University of Idaho (National Institutes of

Health grant P20GM104420), Idaho Institutional Development Awards Network of

Biomedical Research Excellence (INBRE) Program Core Technology Access Grant (NIH

Grant Nos. P20GM103408), and National Science Foundation Grant No. 2143405. The content

is solely the responsibility of the authors and does not necessarily represent the official views

of these funding agencies.

Data Availability

Data supporting the results in the paper are available in the Supplementary Material.

The GenBank accession number for the Open reading frames of the Ksino killer toxin of

Candida sinolaborantium LESF 1467 is PP790515. The sequenced genome of *Candida*

sinolaborantium can be accessed through BioProject PRJNA916362, BioSample

SAMN32422847, Assembly JAQRFW000000000. The raw sequence reads are also available

through the NCBI Sequence Read Archive (SRR23081998). Gene prediction and annotation

are provided in https://figshare.com/s/1f31cd262fc9de969d38

(10.6084/m9.figshare.25895278).

Acknowledgments

We would like to thank Conselho de Gestão do Patrimônio Genético (CGEn) for

providing permits for access to genetic heritage: #AE012D8, #A877F11 and #AA39A6D. We

also thank Amanda Viana and Luciana Simão Carneiro for preparing and preserving yeast

cultures. We also would like to thank the ARS Culture Collection (NRRL), Westerdijk Fungal

Biodiversity Institute (WI-KNAW), and Dr. Roland Klassen for providing yeast cultures used

in this study. We also thank Dr. Primrose Boynton for the critical reading of this manuscript.

Photos of ants in Fig. 1 were accessed from "www.antweb.org": CASENT0173791

Acromyrmex coronatus (Photographer: April Nobile); CASENT0909391 *Mycetomoellerius*

tucumanus (Photographer: Will Ericson); CASENT0901666 *Mycetophylax auritus*

(Photographer: Ryan Perry); CASENT0173988 *Mycocepurus goeldii* (Photographer: April

Nobile); CASENT0922040 *Apterostigma goniodes* (Photographer: Michele Esposito).

Authors' contributions

RB wrote the manuscript supervised by PAR and with inputs from AR, JWC, and MY;

RB conducted cross-interaction killer assays and dsRNA extractions. JWC performed protein

structure predictions and analysis. TJB conducted cross-interactions with pathogenic yeasts.

RACS performed genome assemblage and annotation. SAC performed killer assays for Ksino

characterization. SAC performed microscopic evaluations and killer assays. RB and RTT

performed dsDNA extractions. NF and MJD performed genome sequencing. PAR and AR

supervised and obtained funding for the project. AR is responsible for permit applications. All

authors discussed and reviewed the manuscript and approved the final version.

Supplementary figures

646

647 **FIG S1. Killer yeasts from attine ant gardens can inhibit the growth of opportunistic human pathogens.** A
648 heatmap showing interactions between killer yeasts associated with attine ants and opportunistic human
649 pathogens. Origins of the ant-associated killer yeasts; *Acromyrmex coronatus* (circle); *Mycetomoellerius*
650 *tucumanus* (square); *Mycetophylax* aff. *auritus* (triangle), *Apterostigma goniodes* (diamond). Killer toxin activity
651 was qualitatively assessed based on the presence and size of growth inhibition zones and/or methylene blue
652 staining around killer yeasts as diagrammed. Darker colors on the heatmap represent a more prominent killer
653 phenotype, with yellow indicating no observable killer phenotype. Clusters on the dendrograms connecting
654 individual killers or susceptible yeasts indicate similar susceptibilities to killer toxins or antifungal activities.
655 Darker colors on the cluster diagram represent a more prominent killer phenotype, with yellow indicating no
656 observable killer phenotype.

FIG S2. Elongated growth of *K. ohmeri* after coculture with the killer yeast *C. sinolaborantium*. A. Representative images showing the interaction between *K. ohmeri* and the killer yeast *C. sinolaborantium* on YPD agar. After extended incubation times at ~22°C (15 - 22 days), vertical growth from the surface of the agar was observed. B. Elongated growth phenotypes are not observed when *C. sinolaborantium* is grown in isolation on agar.

FIG S3. Antifungal activities exhibited by fractionated media after growth of *Candida sinolaborantium*. A. The workflow demonstrates the ultrafiltration of YPD-spent growth media to obtain fractions of antifungal molecules. Crude extracts were processed by ultrafiltration with molecular weight cut-off (MWCO) filters of 100 kDa, 30 kDa, and 5 kDa. The resulting protein preparations were used to challenge susceptible yeasts before (>) and after (<) ultrafiltration. Fractions were used **B.** before or **C.** after ethanol-precipitation **D.** Phenotypic examples of antifungal activity detectable in different fractions against representative susceptible yeast species.

672 **FIG S4. Detection of Ksino expression by RT-PCR.** XXXXXX

FIG S5. Phylogeny of Ksino homologs identified in diverse fungi. The Ksino-encoding yeast *C. sinolaborantium* is highlighted in bold, and Saccharomycotina in blue. The tree was inferred with maximum likelihood criteria, and numbers on branches are ultrafast bootstrap support values (only values higher than 70 are shown). The dsRNA-encoded killer toxin Klus is represented as an outgroup.

FIG S6. Computational predictions of protein stability with leucine and serine at position 186 in Ksino. A. AlphaFold2 models of Ksino wild type S186 and mutant L186 after 100 ns of molecular dynamics simulation. Residues 186 and P30 are shown as sticks. P30 forms a hydrogen bond (1.6 Å) with the hydroxyl of S186 but is too distant to create this bond in L186. FoldX 5.0 analysis revealed a $\Delta\Delta G_{Folding}$ of 1.0 kcal/mol for L186S. **B.** 100 ns MD simulation of Ksino L186 AlphaFold2 model. Unlike wild type and Klus, the RMSD stabilizes around 0.55 nm.

FIG S7. Recombinant expression of Ksino does not cause a white coloration of *S. cerevisiae*. Recombinant expression of Ksino by *S. cerevisiae* using galactose induction without a lawn of *K. ohmeri*. Dextrose was used as a non-induced control. n/a indicates an empty vector negative expression control.

Supplementary Tables

Table S1. Strains and species of yeasts used in the current study.

Table S2. The effect of isolation source on killer yeasts activity.

Table S3. The effect of pH on the growth inhibition of *Kodamaea ohmeri* LESF 1474 and *C. castelli* by *Candida sinolaborantium* LESF 1467

Table S4. The effect of temperature on the growth inhibition of *Kodamaea ohmeri* LESF 1474 and *C. castelli* by *Candida sinolaborantium* LESF 1467.

Table S5. The ratio between elongated and yeast morphology of *Kodamaea ohmeri* LESF 1474 cells challenged with killer toxins.

Table S6. Cell survival of *Kodamaea ohmeri* LESF 1474 cells against killer toxins.

Table S7. Statistics of the assembly, prediction, and annotation of the *Candida sinolaborantium* LESF1467 genome.

Table S8. Killer toxins were used as queries for the BLASTp search of the *Candida sinolaborantium* LESF1467 genome.

Table S9. Blastp hits of killer toxins towards *Candida sinolaborantium* LESF1467 genome.

Table S10. SWISS-MODEL structure assessment scores of Ksino top unrelaxed model and Klus top unrelaxed model compared to models after 100 ns of molecular dynamics simulation. Global pLDDT score is also included in the last row.

Table S11. Statistics of Blastp hits of Ksino homologs.

Table S12. Nucleotide sequences used in phylogenetic analysis of yeast species delimitation.

Table S13. The effect of pH on the growth inhibition of *Kodamaea ohmeri* LESF 1474 and *Candida castyelli* by Ksino expressed by *S. cerevisiae*.

Table S14. The effect of temperature on the growth inhibition of *Kodamaea ohmeri* LESF 1474 and *Candida castyelli* by Ksino expressed by *S. cerevisiae*.

Table S15. Yeasts used in killer assays as susceptible or killer positive standards.

Supplementary files

File S1 - All killer assay data.

File S2 - PDB coordinates for the molecular models of Ksino and Klus.

4. Materials and Methods

4.1. Yeast source and phylogenetic analysis

To investigate the prevalence of killer yeast in the fungus garden of attine ants, a representative collection of 180 yeasts previously characterized by Bizarria Jr. et al. (2023) was examined (Table S1). Briefly, yeasts were obtained by fungus garden suspension in saline solution (0.85% NaCl) supplemented with 0.05% Tween 80. Suspensions were ten-fold diluted and surface spread on nutrient media (Yeast-malt agar and Sabouraud dextrose agar, supplemented with 150 µg mL^{-1} chloramphenicol, and pH adjusted to 4.5; details in Bizarria

Jr. et al. 2023). Yeast characterization involved microsatellite amplification analysis (MSP-
PCR; Meyer et al. 1993) and sequencing of the D1/D2 region of the large subunit (LSU)
ribosomal RNA gene (Kurtzman and Robnett, 1998). Our selection of the 180
Saccharomycotina yeasts spanned from fungus gardens of the leaf-cutting ant *Acromyrmex*
coronatus (n = 91), the non-leaf-cutting ant *Mycetomoellerius tucumanus* (n = 13), the lower
attines *Mycetophylax* aff. *auritus* (n = 42), and *Mycocepurus goeldii* (n = 25), which cultivate
Agaricaceae fungi; and *Apterostigma goniodes* (n = 9), which cultivates *Pterulaceae* fungi.

To assess the phylogenetic relationship between the yeast strains, we performed a
phylogenetic analysis with representative species and their closest relatives (the accession
number of sequences used in phylogenetic analysis is listed in Table S12). Sequences were
aligned with MAFFT v.7. (Katoh et al. 2019), and nucleotide substitution models were
generated using the Bayesian information criterion (BIC) by the standard model generation in
IQ-TREE2. SYM+FQ+I+G4 was selected as the nucleotide substitution model. Maximum
likelihood (ML) phylogenetic trees were reconstructed with ultrafast bootstrap in IQTREE2
(Nguyen et al. 2015), with 10,000 replicates for ultrafast bootstrap (tree completed after 262
iterations, LogL: -14973.244). The final trees were edited in FigTree v.1.4.3 (Rambaut, 2016).

4.2. Killer yeast assays

Killer assays were performed to confirm killer toxin production by yeasts, as described
in Crabtree et al. (2019). The 180 yeast strains were seeded onto YPD 4.6 dextrose "killer
assay" agar plates (yeast extract 10 g L^{-1}, peptone 20 g L^{-1}, dextrose 20 g L^{-1}, sodium citrate
29.9 g L^{-1}, supplemented with 0.003% w/v methylene blue, and pH adjusted to 4.6 with HCl),
seeded 10^5 cells mL^{-1} with one of the 49 selected yeast susceptible strains (i.e. from fungus
gardens and sources not related to attine ant environment, Table S13). Nine *Saccharomyces*
spp. strains that encode known toxins were used as positive controls for killer toxin productio,
including K1, K1L, K21, K2, K28, K45, K62, K74, and Klus (Table S13). Yeasts were also
evaluated using 20 pathogenic yeasts as susceptible strains (Table S13). Toxin production was
recorded after five days of incubation at room temperature, either by an inhibition zone or a
methylene blue stain of susceptible strains. Killer activity severity was scored following
Fredericks et al. (2021) by zone of inhibition with or without methylene blue halo or by strong
or weak methylene blue halo without inhibition. We applied heatmaps to visualize the species
interactions, employing the function heatmap.2 from the gplots package, with the default

configuration for dendrogram display (Warnes et al., 2024). To evaluate the effect of the yeast sources (i.e., same or different fungicultures or sites) on killer toxin activity (i.e., killed or non-killed). We selected 9,000 interactions (toward 50 strains that showed susceptibility to killer toxins) and analyzed them with Pearson's Chi-squared test. All yeasts used in killer assays are listed in Table S15.

4.3.   Extraction of dsRNA encoding elements

To determine if the yeast strains host dsRNA elements, we extracted the dsRNA elements from all yeast strains following Okada et al. (2015), with modifications as stipulated in Crabtree et al. (2019). The K1 killer yeast *S. cerevisiae* BJH001 was used as a positive control for all assays. Specifically, yeast cultures were grown in yeast-peptone-dextrose broth (YPD: yeast extract 10 g L^{-1}, peptone 20 g L^{-1}, dextrose 20 g L^{-1}), centrifuged at 3000 × g for 5 min, and the broth was removed by aspiration. Cells were resuspended in water, centrifuged at 8,000 × g, and aspirated. A total of 450 µL of 2× sodium chloride-Tris-EDTA buffer (STE: 200 mM NaCl; 20 mM Tris-HCl, pH 8.0; 30 mM EDTA) and 0.5 µL β-mercaptoethanol were added, and cells were disrupted in a cell disruptor (Digital Disruptor Genie, Scientific Industries) for 3 min at 3000 rpm. Then, 50 µL of 10% (w/v) sodium dodecyl sulfate (SDS) solution and 350 µL phenol–chloroform–isoamyl alcohol (25:24:1) were added to crude extracts, which were homogenized and centrifuged at 10,000 × g for 5 min. Then, 500 µL of supernatant was mixed with absolute ethanol (5:1), and the final volume of 600 µL was transferred to cellulose columns (i.e., 0.05 g cellulose D in a 0.6 mL tube bottom punctured and placed in collecting tubes).

Column tubes were centrifuged at 10,000 × g for 30 seconds, and the flow-through was discarded. Then, 350 µL of wash buffer (1× STE containing 16% v/v EtOH) was added to the column, and the tubes were centrifuged at 10,000 × g for 30 seconds, followed by the flow-through discard. Columns were transferred to a collecting tube, adding 350 µL of 1× STE, and tubes were centrifuged at 10,000 × g for 30 seconds. The eluted fraction was recovered, and 1 mL of absolute ethanol with 40 µL of 3 M aqueous sodium acetate (pH 5.2) was added. Tubes were inverted ten times and centrifuged at 20,000 × g for 15 min. The ethanol mix was aspirated without disturbing the precipitated pellet, which was then air-dried and dissolved in 15 µL of molecular-grade water. The presence of dsRNAs was examined (loading 5 µL of each sample) by 0.8% TAE agarose gel electrophoresis, stained with ethidium bromide, at 130 v for 30 min

in 1% TAE buffer. As a molecular weight reference, a 1 kb DNA ladder was used. The

Saccharomyces cerevisiae strain YJM1307, known to host dsRNA totivirus and satellites, was

used as positive control.

4.4. Extraction of dsDNA linear plasmid

To determine if dsDNA linear plasmids encode the killer toxins, we extracted the

dsDNA elements from positive killer yeast, *Kluyveromyces lactis* AWJ137 with the linear

DNA plasmids pGKL2 (13.5 kb) and pGKL1 (8.9 kb) and *Pichia acaciae* NRRL Y-18665 with

the linear plasmids pPac1-1 (12.6 kb) and pPac1-2 (6.8 kb) (Worsham and Bolen, 1990; Gietz

et al., 1995). Cultures were grown in YPD overnight, 200 µL were centrifuged at 3000 × g for

5 min, and the broth was removed by aspiration. Yeast cells were resuspended in 10 µL

Zymolase solution (5% of 50 mM Tris-HCl, pH 8; 10% of 5 mM EDTA, and 85% water) and

7.2 µL of 10 mg mL^{-1} of Zymolase (United States Biological Life Sciences, Swampscott, MA),

and the suspension was incubated at 30°C for 1 h. After incubation, 1 µL of 10% SDS and 3

µL of proteinase K (New England Biolabs, Ipswich, MA) were added to the suspension,

followed by an incubation of 1 h at 65°C. After incubation, 5 µL of loading dye was added,

and the suspension was centrifuged at 3000 × g for 5 min. The presence of dsDNA plasmids

was examined (loading 10 µL of each sample) by 0.6% TAE agarose gel electrophoresis,

stained with ethidium bromide, at 130 v for 1 h in 1% TAE buffer.

4.5. Toxin precipitation and killer toxin activity at different temperatures and
pH

To investigate the stability of the toxins produced by *C. sinolaborantium* LESF 1467,

the isolate extracts on YPD medium (pH 4.6) were filtered, mixed with 1:1 absolute ethanol,

centrifuged at 20,000 × g for 20 min, followed by supernatant aspiration and suspension in 10

µL of YPD (pH 4.6). To confirm the killer activity of the precipitates, 4 µL of the suspended

proteins were used in killer assays against *Kodamaea ohmeri* LESF 1474. To evaluate the

thermostability, the precipitated toxins were screened in killer assays incubating at room

temperature or heat-inactivated at 60°C and 98°C for 2 min. For quantitative analysis of killer

toxin expression, 2 µl of 10^8 cells mL^{-1} were plated onto susceptible lawns. The killer activity

was also evaluated for activity at different pHs, adjusting the pH to 3.5, 4, 4.5, 5, 5.5, and 6

with HCl 12N, and at various temperatures, incubating at 17, 20, 25, 30, and 35°C. The area of

activity on killer assays (i.e., zone of inhibition and cell differentiation of the lawn strain) was

measured in cm² after four days with ImageJ 1.53t (Schneider et al., 2012). Since the area data

do not satisfy the assumptions of parametric tests, we analyzed variance with Kruskal-Wallis

followed by the Wilcoxon rank sum test with Bonferroni correction as a post-hoc for pairwise

multiple comparisons. Analyses were performed using the software RStudio v. 2023.12.0.369

(POSIT TEAM, 2023), and R v. 4.3.2 (R CORE TEAM, 2023). Figures were prepared using

ggplot2 (Wickham, 2016).

4.6.   **Production of elongated cells by *Kodamaea ohmeri* against killer toxins**

We observed that the yeast *K. ohmeri* LESF 1474 produces elongated cells in the

presence of *C. sinolaborantium* toxins and their precipitates. We suspended the cells in the

white and raised zones of killer assays and on axenic cultures in $1\times$ PBS. Cell morphology was

checked under a microscope (Zeiss Primo Star iLED) by counting budding (2 to 17×10^6 cells

mL^{-1}) and elongated cells (2 to 15×10^5 cells mL^{-1}) with a Neubauer chamber. Cell survival

was evaluated by adjusting the suspensions to 10^3 cells mL^{-1} and seeding 100 µL in YPD.

Colony-forming units (CFU) were counted after three days of incubation at room temperature.

A similar elongated cell phenotype was observed in the presence of *S. cerevisiae* DSM-70459,

which secretes the known killer toxin Klus (encoded by dsRNA elements). Microscopic and

survival evaluations were also conducted in the presence of *S. cerevisiae* DSM-70459, as

described above. To confirm that the elongated cell phenotype is related to the presence of the

dsRNA satellites, dsRNA satellites were cured from *S. cerevisiae* by growing on a YPD

medium supplemented with 5.0 µM cycloheximide. Loss of killer toxin production and the loss

of dsRNAs were assayed as described above. The ratio between normal and elongated cells

was compared between *K. ohmeri* in the white halo around *C. sinolaborantium* and *S.*

cerevisiae DSM-70459 with axenic cultures surrounding the halo, with ten independent

replicates. The data was checked for the assumption of parametric tests (performing Shapiro-

Wilk and Bartlett tests, respectively, for normality and homogeneity of variances). The ratio

data in both treatments, towards *C. sinolaborantium* and *S. cerevisiae* DSM-70459, were found

not to satisfy the assumptions of equal variance (Bartlett test, df = 1, p-value < 0.001), so we

conducted the Welch Two Sample t-test to compare means between treatments. The same

procedure was performed for cell survival data towards *C. sinolaborantium*, which also did not

satisfy the assumptions of equal variance (Bartlett's K-squared = 4.16, df = 1, p-value = 0.04).

Cell survival data towards *S. cerevisiae* DSM-70459 satisfied the assumptions for parametric

tests (Bartlett's K-squared = 0.52, df = 1, p-value = 0.47; Shapiro-Wilk p-value > 0.05), so we
conducted the Two Sample t-test to compare means between treatments.

4.7. Genomic DNA extraction, sequencing, and annotation

To extract genomic DNA from *C. sinolaborantium*, the yeast was grown in YPD broth;
cells were recovered by centrifugation (3000 × g for 5 min), washed, and suspended in 500 µL
TE buffer (10 mM Tris-Cl; 0.1 mM EDTA, pH 8). Recovered cells were transferred to a
microtube, centrifugation as previously described, and the supernatant was removed. The pellet
was disrupted by vortexing. Cells were suspended in a lysis buffer (2% Triton X-100, 1% SDS,
100 mM NaCl, 10 mM Tris-HCl, pH 8, 1 mM EDTA pH 8) with approximately 300 µL of
glass beads. A volume of 200 µL of phenol-chloroform-isoamyl alcohol solution was added,
and the sample was vortex for 3 min at 3000 rpm (Disruptor Genie), followed by the addition
of 200 µL of TE buffer and centrifugation for 5 min at 21000 × g. The upper layer was
transferred to a new microtube and mixed by inversion with 1 mL of absolute ethanol. The tube
was centrifuged for 3 min at 21000 × g, the supernatant removed, and the pellet suspended in
400 µL of TE buffer. Then, 30 µL of DNAse-free RNAse A was added, and the solution was
mixed and incubated for 5 min at 37°C. The DNA was precipitated by adding 10 µL of 4 M
ammonium acetate and 1 mL of absolute ethanol, mixing by inversion, and centrifuging for 3
min at 21000 × g. The supernatant was discarded and the pellet was dried for approximately
20 min at room temperature. Genomic DNA was suspended in 100 µL of TE buffer and stored
at -20°C.

The genome was sequenced by Illumina NextSeq 550 using the paired-end method.
Libraries were created with the Illumina DNA Prep kit for WGS
(https://www.illumina.com/products/by-type/sequencing-kits/library-prep-kits/nextera-dna-
flex.html). The initial quality analysis and cleaning steps were performed with fastp v. 0.20.1
(CHEN et al., 2018). The genome was assembled with SPAdes v. 3.15.5 (GUREVICH et al.,
2013) with multiple k-mers (21, 31, 41, 51, 61, 71, and 81). The blobtools v. 3.5.2 software
(LAETSCH; BLAXTER, 2017) was used to check contaminants. Briefly, Blastn 2.13.0+
(ALTSCHUL et al., 1990) was used to search for structures across the entire NCBI nt database;
minimap2 v. 2.24-r1122 (LI, 2018) and samtools v.1.13 (LI et al., 2009) were used to obtain
sequencing coverage. The blobtools visualization was used to ascertain the results. The rDNA
and ITS regions were annotated with ITSx v. 1.3 (BENGTSSON-PALME et al., 2013), which
was compared to an amplicon previously submitted to NCBI for this isolate (accession:

ON493969) as part of the contamination verification steps. Final assembly statistics were
obtained with QUAST v. 5.2.0 (BANKEVICH et al., 2012). Completeness was checked using
BUSCO v. 5.4.4 (SEPPEY et al., 2019) with the saccharomycetes_odb10 dataset (2,137
BUSCOs). The genetic prediction was performed using Braker v. 2.1.6 (BRŮNA et al., 2021)
[which includes Augustus v. 3.4.0 (STANKE; MORGENSTERN, 2005), ProtHint v. 2.6.0 and
GeneMark-ES v. 4.71 (LOMSADZE et al. 2005)] with OrthoDB v. 10 methods
(KRIVENTSEVA et al., 2019). InterProScan v. 5.60 (Jones et al., 2014) was used for
functional annotation of predicted proteins.

4.8.   Genome mining for toxin-encoding regions

To identify toxin-encoding chromosomal regions, a search using proteins predicted
from whole genome sequencing was performed with BLASTP 2.9.0, using a protein sequence
database of known killer toxins. Results with query/hit alignments greater than 100 were
selected. An open reading frame similar to the Klus toxin (28% identity, with 151 aa alignment
length and an e-value of 1.56E^{-05}) was found and selected for further analysis. Polymerase
chain reactions (PCR) were performed to amplify the region of interest using primers designed
with Primer3 software. 4.1.0 (https://primer3.ut.ee/). The primers were designed to target Ksino
untranslated regions (961F: 5'-TTAACGACTTTCGTCTTCGCTATCC-3' and 961R: 5'-
ATTGAGATCAGGTGGCCTGTGTAGC-3') followed by nested-PCR to amplify the open
reading frame using the previous PCR product as the template (961NF: 5'-
CGATCACCTAGCCCAAAATGC-3' and 961NR: 5'-AAAGTGTTGGCCAAGGACACG-
3'). Both of the amplification reactions used Phusion high fidelity master mix (New England
Biolabs, Ipswich, MA) with the amplification conditions of 98°C for 3 min, 30 cycles at 98°C
for 30 s, 63°C for 30 s, 72°C for 30 s, and a final extension step at 72°C for 5 min. Amplicons
were sequenced to confirm the identity of the region of interest.

4.9.   Prediction of tertiary structure and comparative analysis of proteins

After confirming the genomic coding region of Ksino was the correct size using PCR,
the sequence was used to predict protein structure via the AlphaFold2 software (Jumper et al.,
2021) with default parameters. The B factor columns of each unrelaxed structure were used to
create per residue pLDDT graphs. Secondary structure was assigned using PyMol to create a
linear representation of the protein and aligned to per residue graphs. The top-scoring relaxed

model from AlphaFold2 was used for molecular dynamics simulation using the GROMACS package (Spoel et al., 2005). Briefly, the system was set up using the AMBER99SB-ILDN force field with TIP3P water, a dodecahedral box with 1.0 nm padding, and 0.15 M NaCl. The system was then subjected to energy minimization by steepest descent, followed by NVT and NPT equilibration with a pressure of 1.0 atm and temperature of 300 K. Pressure and temperature were maintained constant using arrinello-Rahman with isotropic coupling and V-rescale with Protein Non-protein as the coupling groups, respectively. This setup was chosen to match standard cellular conditions closely: 26.85°C, 0.15 M NaCl, pH 7, and 1 atm. Simulations were run for 100 ns with a 2.0 fs timestep, and snapshots were saved every 2 fs. The RMSD was calculated as a function of time using the resulting trajectories. In addition, the final snapshot of each trajectory was saved as a PDB structure file and compared to relaxed and unrelaxed AlphaFold2 models using the SWISS structure assessment tool (Studer et al., 2019).

FoldX was used to predict the $\Delta\Delta G_{Folding}$ of L186S (Delgado et al., 2019). Amber relaxed Ksino L186 was used as the starting structure, and FoldX PDBrepair was carried out six times to optimize the structure. Then, L186 in the optimized structure was mutated to every other residue using PositionScan. $\Delta\Delta G_{Folding}$ was taken from the output file.

4.10. Bioinformatic discovery of Ksino homologs in fungi

A search for similar sequences in NCBI (https://www.ncbi.nlm.nih.gov/) was also carried out with Blastp, with the results filtered by alignment length (above 109), identity (above 30%), and e-value (less than 0.03). The protein sequences were aligned with MUSCLE (https://www.ebi.ac.uk/jdispatcher/msa/muscle), with the alignment consisting of 86 sequences and 426 positions. Amino acid substitution models were generated according to the Bayesian information criterion (BIC), with WAG+R4 being the chosen model. Maximum likelihood (ML) phylogenetic trees were reconstructed in IQTREE2 (Nguyen et al., 2015), with 10,000 ultrafast bootstrap replicates (tree completed after 163 iterations, LogL: -18525.485). The final trees were edited in FigTree v.1.4.3 (RAMBAUT, 2016).

4.11. Cloning and transformation in *Escherichia coli*

To study the coding region, a "TA" terminal was added to the amplicon in a reaction that included 1 μL Taq polymerase, 19 μL of PCR product, 1 μL of 10 μM dNTPs, 5 μL of 10×

buffer, and 24 µL of sterile ultrapure water. The reaction was incubated at 72°C for 20 min.

For cloning, the pCR™8/GW/TOPO™ TA Cloning kit (Invitrogen, Waltham, MA) was used,

adding 1 µL of amplicons with TA terminals, 0.25 µL of saline solution from the kit, and 0.25

µL of the pCR™8/GW/TOPO™ vector. The reaction was incubated at room temperature for 2

h, followed by adding 25 µL of competent *E. coli* cells (One-Shot TOP10 Competent Cells,

Invitrogen, Waltham, MA). The reaction was kept on ice for 30 min, followed by heat shock

at 42°C for 30 s and then ice bath for 2 min. Subsequently, 250 µL of Super Optimal broth

(SOC: 20 g L^{-1} Tryptone, 5 g L^{-1} Yeast extract, 0.58 g L^{-1} NaCl, 0.19 g L^{-1} KCl, 2, 03 g L^{-1}

MgCl$_2$, 1.20 g L^{-1} MgSO$_4$, and 3.6 g L^{-1} dextrose) preheated to 37°C was added, and incubated

at 37°C for 1 h. Then, 250 µL were plated on Luria bertani medium (LB: 10 g L^{-1} of tryptone,

5 g L^{-1} of yeast extract, 10 g L^{-1} of NaCl, and 15 g L^{-1} of agar) supplemented with 100 µg mL-

1 of spectinomycin. Plasmids were extracted using the High-Speed Mini Plasmid kit (IBI

Scientific, Dubuque, IA), following manufacturer recommendations.

Due to the transcriptional diversity of CUG-Leu codons in some yeasts (Krassowski et

al., 2018), the vectors containing the region of interest were subjected to targeted mutations

(SDM: Site-Directed Mutagenesis) in a CUG codon starting at position 556 of the DNA

sequence (position 186 in the protein sequence). Thus, the mutants consist of the alternatives

identified for CUG in other yeasts, being called L186A and L186S, with L being the original

amino acid, 186 the position, and "S" the mutation for Ser. To obtain the mutants, we design

primers   targeting   position   186   as   follows: 961GCG: 5'-

CAGCTGTGGCGCGAACCATAGCGGG-3',   961TCG: 5'-

CAGCTGTGGCTCGAACCATAGCGGG-3'   (forward),   and   961SNM: 5'-

AGTCCCGCAGGCCAATCC-3' (reverse). Amplification reactions included 12.5 µL of

Phusion High Fidelity Master Mix, 1 µL 10 µM forward primer, 1 µL 10 µM reverse primer,

9.5 µL sterile ultrapure water, 1 µL 1:9 diluted DNA (i.e., pCR™8 plasmid with the region of

interest). Amplification conditions were as follows: 98°C for 30 min, 25 cycles at 98°C for 5

s, 67°C for 10 s, 72°C for 50 s, and a final extension step at 72°C for 5 min. To remove the

template DNA from the reaction, a Dpnl treatment was performed by mixing 22 µL of PCR

product, 2 µL Dpnl, 3 µL CutSmart buffer, and 3 µL sterile ultrapure water. The reaction was

incubated at 37°C for 1 day. For the addition of 5'-P and 3'-OH, we performed a PNK treatment:

30 µL of the treated product, 5 µL PNK buffer, 1 µL PNK, 5 µL 10 mM ATP, 1 µL 0.1 M DTT,

8 µL of sterile ultrapure water. The reactions were incubated for 30 min at 37°C and inactivated

at 65°C for 10 min. The product was purified using the Monarch DNA & PCR Cleanup kit

(New England Biolabs, Ipswich, MA), following the manufacturer's protocols. To re-ligate the

plasmids, the following reaction was performed: 2 μL of purified product, 1 μL T4 DNA ligase,

2 μL 10× binding buffer, 15 μL sterile ultrapure water. The reaction was incubated for 2 h at

room temperature, and then 25 μL of competent *E. coli* cells (One-Shot TOP10 Competent

Cells, Invitrogen, Waltham, MA) were added. Transformed cultures were selected in LB

supplemented with 100 μg mL^{-1} spectinomycin, plasmids extracted with the High-Speed Mini

Plasmid kit (IBI Scientific, Dubuque, IA), and mutations checked by sequencing (ELIM

Biopharmaceuticals, Hayward, CA).

The vectors were subjected to a Gateway™ recombination from the vectors in pCR8 to

pAG306-GAL-ccdB. For recombination, the reaction was conducted with 0.5 μL of the source

vector, 0.5 μL of the destination vector, 1 μL of sterile ultrapure water, and 0.5 μL of LR

Clonase II (New England Biolabs, Ipswich, MA). The reaction was incubated for 4 h at room

temperature, followed by adding 0.25 μL of Proteinase K (New England Biolabs, Ipswich, MA)

and incubating at 37°C for 10 min. Finally, the *E. coli* transformation was performed as

previously described, and cultures were selected in LB supplemented with 100 μg mL^{-1} of

chloramphenicol and ampicillin.

1023

1024 **4.12. Transformation and heterologous expression of killer toxins**

1025 **by*Saccharomyces cerevisiae***

1026

For yeast transformation, the integrative vector containing a killer toxin gene was

linearized with the NcoI enzyme by cutting at the *URA3* locus, followed by inactivation of the

enzyme at 65°C for 30 min. The product was purified using the Monarch DNA & PCR Cleanup

kit (New England Biolabs, Ipswich, MA), following the manufacturer's recommendations. *S.*

cerevisiae CRY1 cells were grown in YPD until an optical density of 1.3 was obtained, then

the cells were washed with sterile water and suspended in 10 mL of TE buffer supplemented

with 100 mM LiAc, and incubated at 30°C for 30 min under 130 rpm agitation. Cells were

recovered by centrifugation at 1500 × g for 5 min at 4°C and suspended in 5 mL of ice-cold

1M sorbitol, followed by centrifugation and suspended in 550 μL of ice-cold 1M sorbitol. For

transformation, 80 μL of the suspended cells were added to 5 to 10 μg of the linearized plasmids

and transferred to electroporation cuvettes. The cuvettes were incubated on ice for 10 min and

pulsed with GenePulser (200 Ω, 1.5 kV, and 25 μF). To recover the transformed cultures, the

cells were grown at 30°C for 2 days in CM medium without uracil (complete media: 2.5 g L^{-1}

of the drop-out mix without uracil, 1.7 g L^{-1} of yeast nitrogen base YNB, 5 g L^{-1} ammonium

sulfate, 20 g L^{-1} dextrose). For killer toxin expression was induced by overnight growth in

YPGal (yeast extract 10 g L^{-1}, peptone 20 g L^{-1}, galactose 20 g L^{-1}) followed by plating on
either YPD or YPGal with citrate-phosphate 29.9 g L^{-1}, supplemented with 0.003% w/v
methylene blue and pH adjusted to 4.6 with HCl and 15 g L^{-1} agar seeded with 10^5 cells mL^{-1}
of a susceptible lawn strain. *C. sinolaborantium* and *S. cerevisiae* CRY1 were used as positive
and negative controls for killer activity, respectively.

1047

1048 **4.13. Detection of killer toxin expression by RT-PCR**

1049

An overnight culture of Ksino was grown in YPD and harvested at OD600 0.9-1.2.
Speroplasting was performed by incubating the harvested cells at 30°C for 30 min in buffer Y1
(1M sorbitol, 0.1 M EDTA, pH 7.4, B-ME 0.1%, and 0.1% v/v zymolase 20T (10 mg/mL)).
Spheroplasts were used as input for a Qiagen RNeasy procedure. The RNA quality was checked
using a spectrophotometer, and 10 μg of total RNA was treated with 1 μL of DNase I
(ThermoFisher Scientific) in the presence of 10 μL of DNase reaction buffer at 37°C for 10
min, and then inactivated by heat at 75°C for 10 min. Transcripts encoding Ksino or actin were
detected by RT-PCR (Superscript IV) following the manufacturer's instructions using the
primers 5'-GGCCTTGAGATACCCCATCG-3' and
5'-CCACGTGAGTAACACCGTCA-3' (for *ACT1*) and 5'-
GATCTACGACAGCAGCCTAGA-3' and
5'-GTGGTGTACATCGTTGACTCC-3' (for Ksino). Briefly, the RNA template is denatured
with a reverse primer mix at 65°C for 5 min and incubated on ice for at least 1 min. The reverse
transcription reaction mix was added to the denatured RNA and incubated for 10 minutes at
55°C and then inactivated at 80°C for 10 minutes. A 20 L PCR was performed with the cycling
instructions as follows (98°C for 30 s, then 30 cycles of 98°C for 10 s, 60°C for 30 s, 72°C for
10 s followed by one cycle of 72°C for 5 min). PCR products were analyzed by agarose gel
electrophoresis.

References

Abranches, J., Mendonça-Hagler, L. C., Hagler, A. N., Morais, P. B., Rosa, C. A. (1997). The
incidence of killer activity and extracellular proteases in tropical yeast communities. **Canadian**
Journal of Microbiology, 43(4), 328-336. doi: 10.1139/m97-046

1072

1073 Abranches, J., Valente, P., Nóbrega, H. N., Fernandez, F. A., Mendonça-Hagler, L. C., Hagler,
1074 A. N. (1998). Yeast diversity and killer activity dispersed in fecal pellets from marsupials and
1075 rodents in a Brazilian tropical habitat mosaic. **FEMS Microbiology Ecology**, 26(1), 27-33.
1076 doi: 10.1111/j.1574-6941.1998.tb01558.x

1077

1078 Altschul, S. F., Gish, W., Miller, W., Myers, E. W., Lipman, D. J. (1990). Basic local alignment
1079 search tool. **Journal of molecular biology**, 215(3), 403-410. doi: 10.1016/S0022-
1080 2836(05)80360-2

1081

1082 Amos, B. A., Hayes, R. A., Leemon, D. M., Furlong, M. J. (2019). Small hive beetle
1083 (Coleoptera: Nitidulidae) and the yeast, *Kodamaea ohmeri*: a facultative relationship under
1084 laboratory conditions. **Journal of Economic Entomology**, 112(2), 515-524. doi:
1085 10.1093/jee/toy378

1086

1087 Antunes, J., Aguiar, C. (2012). Search for killer phenotypes with potential for biological
1088 control. **Annals of microbiology**, 62, 427-433. doi: 10.1007/s13213-011-0256-z

1089

1090 Arcuri, S. L., Pagnocca, F. C., da Paixão Melo, W. G., Nagamoto, N. S., Komura, D. L.,
1091 Rodrigues, A. (2014). Yeasts found on an ephemeral reproductive caste of the leaf-cutting ant
1092 *Atta sexdens rubropilosa*. **Antonie Van Leeuwenhoek**, 106(3), 475–487. doi:
1093 10.1007/s10482-014-0216-2

1094

1095 Bankevich, A., et al. (2012). SPAdes: a new genome assembly algorithm and its applications
1096 to single-cell sequencing. **Journal of computational biology**, 19(5), 455-477. doi:
1097 10.1089/cmb.2012.0021

1098

1099 Belda, I., Ruiz, J., Alonso, A., Marquina, D., Santos, A. (2017). The biology of *Pichia
1100 membranifaciens* killer toxins. **Toxins**, 9(4), 112. doi: 10.3390/toxins9040112

1101

1102 Benda, N. D., Boucias, D., Torto, B., Teal, P. (2008). Detection and characterization of
1103 *Kodamaea ohmeri* associated with small hive beetle *Aethina tumida* infesting honey bee hives.
1104 **Journal of Apicultural Research**, 47(3), 194-201. doi: 10.1080/00218839.2008.11101459

1105

Bengtsson-Palme, J., et al. (2013). Improved software detection and extraction of ITS1 and ITS 2 from ribosomal ITS sequences of fungi and other eukaryotes for analysis of environmental sequencing data. **Methods in Ecology and Evolution**, 4(10), 914-919. doi: 10.1111/2041-210X.12073

Bevan, E. A., Herring, A. J., Mitchell, D. J. (1973). Preliminary characterization of two species of dsRNA in yeast and their relationship to the "killer" character. **Nature**, 245(5420), 81-86. doi: 10.1038/245081b0

Bizarria Jr, R., de Castro Pietrobon, T., Rodrigues, A. (2023). Uncovering the Yeast Communities in Fungus-Growing Ant Colonies. **Microbial Ecology**, 86(1), 624-635. doi: 10.1007/s00248-022-02099-1

Bizarria Jr, R., Pagnocca, F. C., Rodrigues, A. (2022). Yeasts in the attine ant–fungus mutualism: Diversity, functional roles, and putative biotechnological applications. **Yeast**, 39(1-2), 25-39. doi: 10.1002/yea.3667

Boynton, P. J. (2019). The ecology of killer yeasts: Interference competition in natural habitats. **Yeast**, 36(8), 473–485. doi: 10.1002/yea.3398

Branstetter, M. G., Ješovnik, A., Sosa-Calvo, J., Lloyd, M. W., Faircloth, B. C., Brady, S. G., and Schultz, T. R. (2017). Dry habitats were crucibles of domestication in the evolution of agriculture in ants. **Proceedings of the Royal Society B: Biological Sciences**, 284(1852), 20170095. doi: 10.1098/rspb.2017.0095

Brown, D. W. (2011). The KP4 killer protein gene family. **Current genetics**, 57, 51-62. doi: 10.1007/s00294-010-0326-y

Brůna, T., Hoff, K. J., Lomsadze, A., Stanke, M., Borodovsky, M. (2021). BRAKER2: automatic eukaryotic genome annotation with GeneMark-EP+ and AUGUSTUS supported by

a protein database. **NAR genomics and bioinformatics**, 3(1), lqaa108. doi:
10.1093/nargab/lqaa108

Buchan DWA, Minneci F, Nugent TCO, Bryson K, Jones DT (2013). Scalable web services
for the PSIPRED Protein Analysis Workbench. **Nucleic Acids Res.** 41(W1): W349–W357.
doi: 10.1093/nar/gkt381.

Buser, C. C., Jokela, J., Martin, O. Y. (2021). Scent of a killer: How could killer yeast boost its
dispersal?. **Ecology and Evolution**, 11(11), 5809-5814. doi: 10.1002/ece3.7534

Bussey, H., Vernet, T., Sdicu, A. M. (1988). Mutual antagonism among killer yeasts:
Competition between K1 and K2 killers and a novel cDNA-based K1-K2 killer strain of
Saccharomyces cerevisiae. **Canadian journal of microbiology**, 34(1), 38-44. doi:
10.1139/m88-007

Buzzini, P., Martini, A. (2001). Discrimination between *Candida albicans* and other
pathogenic species of the genus *Candida* by their differential sensitivities to toxins of a panel
of killer yeasts. **Journal of clinical microbiology**, 39(9), 3362-3364.
doi:10.1128/jcm.39.9.3362-3364.2001

Carreiro, S. C., Pagnocca, F. C., Bacci, M., Bueno, O. C., Hebling, M. J. A., Middelhoven, W.
J. (2002). Occurrence of killer yeasts in leaf-cutting ant nests. **Folia Microbiologica**, 47(3),
259–262. doi: 10.1007/BF02817648

Castillo, A., Cifuentes, V. (1994). Presence of double-stranded RNA and virus-like particles in
Phaffia rhodozyma. **Current genetics**, 26, 364-368. doi: 10.1007/BF00310502

Chen, S., Zhou, Y., Chen, Y., Gu, J. (2018). fastp: an ultra-fast all-in-one FASTQ preprocessor.
Bioinformatics, 34(17), i884-i890. doi: 10.1093/bioinformatics/bty560

1165

Chessa, R., et al. (2017). Biotechnological exploitation of *Tetrapisispora phaffii* killer toxin: heterologous production in *Komagataella phaffii* (*Pichia pastoris*). **Applied microbiology and biotechnology**, 101(7), 2931-2942. doi: 10.1007/s00253-016-8050-2

Coelho, A. R., Tachi, M., Pagnocca, F. C., Nobrega, G. M. A., Hoffmann, F. L., Harada, K. I., Hirooka, E. Y. (2009). Purification of *Candida guilliermondii* and *Pichia ohmeri* killer toxin as an active agent against *Penicillium expansum*. **Food Additives and Contaminants**, 26(1), 73-81. doi: 10.1080/02652030802227227

Corrêa, M. M., Silva, P. S., Wirth, R., Tabarelli, M., Leal, I. R. (2010). How leaf-cutting ants impact forests: drastic nest effects on light environment and plant assemblages. **Oecologia**, 162, 103-115. doi: 10.1007/s00442-009-1436-4

Crabtree, A. M., Kizer, E. A., Hunter, S. S., Van Leuven, J. T., New, D. D., Fagnan, M. W., Rowley, P. A. (2019). A rapid method for sequencing double-stranded RNAs purified from yeasts and the identification of a potent K1 killer toxin isolated from *Saccharomyces cerevisiae*. **Viruses**, 11(1), 70. doi: 10.3390/v11010070

Crabtree, A. M., Taggart, N. T., Lee, M. D., Boyer, J. M., Rowley, P. A. (2023). The prevalence of killer yeasts and double-stranded RNAs in the budding yeast *Saccharomyces cerevisiae*. **FEMS Yeast Research**, 23, foad046. doi: 10.1093/femsyr/foad046

Craven, S. E., Dix, M. W., Michaels, G. E. (1970). Attine fungus gardens contain yeasts. **Science**, 169(3941), 184–186. doi: 10.1126/science.169.3941.184

Crucitti, D., Chiapello, M., Oliva, D., Forgia, M., Turina, M., Carimi, F., La Bella, F., Pacifico, D. (2021). Identification and molecular characterization of novel mycoviruses in *Saccharomyces* and non-*Saccharomyces* yeasts of oenological interest. **Viruses**, 14(1), 52. doi: 10.3390/v14010052

De Fine Licht, H. H., Boomsma, J. J. (2010). Forage collection, substrate preparation, and diet composition in fungus-growing ants. **Ecological Entomology**, 35(3), 259-269. doi: 10.1111/j.1365-2311.2010.01193.x

1199

1200 Delgado J, Radusky LG, Cianferoni D, Serrano L (2019). FoldX 5.0: Working with RNA, small
1201 molecules and a new graphical interface. **Bioinformatics**. 35(20): btz184. doi:
1202 10.1093/bioinformatics/btz184.

1203

1204 Drinnenberg, I. A., Fink, G. R., Bartel, D. P. (2011). Compatibility with killer explains the rise
1205 of RNAi-deficient fungi. **Science**, 333(6049), 1592-1592. doi: 10.1126/science.1209575

1206

1207 Drinnenberg, I. A., Weinberg, D. E., Xie, K. T., Mower, J. P., Wolfe, K. H., Fink, G. R., Bartel,
1208 D. P. (2009). RNAi in budding yeast. **Science**, 326(5952), 544-550. doi:
1209 10.1126/science.1176945

1210

1211 Farji-brener, A. G., Werenkraut, V. (2015). A meta-analysis of leaf-cutting ant nest effects on
1212 soil fertility and plant performance. **Ecological Entomology**, 40(2), 150-158. doi:
1213 10.1111/een.12169

1214

1215 Fisher, P. J., Stradling, D. J., Sutton, B. C., Petrini, L. E. (1996). Microfungi in the fungus
1216 gardens of the leaf-cutting ant *Atta cephalotes*: a preliminary study. **Mycological Research**,
1217 100(5), 541–546. doi: 10.1016/S0953-7562(96)80006-2

1218

1219 Frank, A. C., Wolfe, K. H. (2009). Evolutionary capture of viral and plasmid DNA by yeast
1220 nuclear chromosomes. **Eukaryotic Cell**, 8(10), 1521-1531. doi: 10.1128/ec.00110-09

1221

1222 Fredericks, L. R., et al. (2021). The species-specific acquisition and diversification of a K1-
1223 like family of killer toxins in budding yeasts of the *Saccharomycotina*. **PLoS genetics**, 17(2),
1224 e1009341. doi: 10.1371/journal.pgen.1009341

1225

1226 Fuentefria, A. M., Faganello, J., Pazzini, F., Schrank, A., Valente, P., Vainstein, M. (2007).
1227 Typing and patterns of cellular morphological alterations in *Cryptococcus neoformans* and
1228 *Cryptococcus gattii* isolates exposed to a panel of killer yeasts. **Medical mycology**, 45(6), 503-
1229 512. doi: 10.1080/13693780701416580

1230

1231   Fuentefria, A. M., Franskoviaki, I. M., Mercado, L. W., Ramos, J. P., Vainstein, M. H., Valente,
1232   P. (2006). Inhibition of clinical and environmental *Cryptococcus neoformans* isolates by two
1233   Brasilian killer yeasts. **Journal of basic microbiology**, 46(2), 87-93. doi:
1234   10.1002/jobm.200510018

1235

1236   Fuentefria, A. M., Perez, L. R., Alves d'Azevedo, P., Pazzini, F., Schrank, A., Vainstein, M.
1237   H., Valente, P. (2008). Typing of *Staphylococcus epidermidis* clinical strains by a selected
1238   panel of Brasilian killer yeasts. **Journal of basic microbiology**, 48(1), 25-30. doi:
1239   10.1002/jobm.200700143

1240

1241   Ganter, P. F., Starmer, W. T. (1992). Killer Factor as a Mechanism of Interference Competition
1242   in Yeasts Associated with Cacti: Ecological Archives. **Ecology**, 73(1), 54-67. doi:
1243   10.2307/1938720

1244

1245   Gietz, R. D., Schiestl, R. H., Willems, A. R., Woods, R. A. (1995). Studies on the
1246   transformation of intact yeast cells by the LiAc/SS-DNA/PEG procedure. **Yeast**, 11(4), 355-
1247   360. doi: 10.1002/yea.320110408

1248

1249   Gimeno, C. J., Ljungdahl, P. O., Styles, C. A., Fink, G. R. (1992). Unipolar cell divisions in
1250   the yeast *S. cerevisiae* lead to filamentous growth: regulation by starvation and RAS. **Cell**,
1251   68(6), 1077-1090. doi: 10.1016/0092-8674(92)90079-R

1252

1253   Giometto, A., Nelson, D. R., Murray, A. W. (2021). Antagonism between killer yeast strains
1254   as an experimental model for biological nucleation dynamics. **Elife**, 10, e62932. doi:
1255   10.7554/eLife.62932

1256

1257   Giovati, L., et al. (2018). Candidacidal activity of a novel killer toxin from *Wickerhamomyces*
1258   *anomalus* against fluconazole-susceptible and-resistant strains. **Toxins**, 10(2), 68. doi:
1259   10.3390/toxins10020068

1260

1261   Golubev, W. I. (1998). Mycocins (killer toxins). In **The Yeasts** (pp. 55-62). Elsevier. doi:
1262   10.1016/B978-044481312-1/50011-3

1263

1264 Golubev, W. I. (2006). Antagonistic interactions among yeasts. In: Péter, G., Rosa, C. (eds)
1265 Biodiversity and Ecophysiology of Yeasts. **The Yeast Handbook**. Springer, Berlin,
1266 Heidelberg. doi: 10.1007/3-540-30985-3_10

1267

1268 Golubev, W. I., Pfeiffer, I., Golubeva, E. (2002). Mycocin production in *Trichosporon*
1269 *pullulans* populations colonizing tree exudates in the spring. **FEMS microbiology ecology**,
1270 40(2), 151-157. doi: 10.1111/j.1574-6941.2002.tb00947.x

1271

1272 Goto, K., Iwatuki, Y., Kitano, K., Obata, T., Hara, S. (1990). Cloning and nucleotide sequence
1273 of the KHR killer gene of *Saccharomyces cerevisiae*. **Agricultural and biological chemistry**,
1274 54(4), 979-984. doi: 10.1080/00021369.1990.10870081

1275

1276 Goto, K., Fukuda, H., Kichise, K., Kitano, K., Hara, S. (1991). Cloning and nucleotide
1277 sequence of the KHS killer gene of *Saccharomyces cerevisiae*. **Agricultural and biological**
1278 **chemistry**, 55(8), 1953-1958. doi: 10.1080/00021369.1991.10870924

1279

1280 Greig, D., Travisano, M. (2008). Density-dependent effects on allelopathic interactions in
1281 yeast. **Evolution**, 62(3), 521-527. doi: 10.1111/j.1558-5646.2007.00292.x

1282

1283 Groenewald, M., et al. (2023). A genome-informed higher rank classification of the
1284 biotechnologically important fungal subphylum Saccharomycotina. **Studies in Mycology**,
1285 105, 1–22. doi: 10.3114/sim.2023.105.01

1286

1287 Guamán-Burneo, M. C., et al. (2015). Xylitol production by yeasts isolated from rotting wood
1288 in the Galápagos Islands, Ecuador, and description of *Cyberlindnera galapagoensis* fa, sp. nov.
1289 **Antonie Van Leeuwenhoek**, 108, 919-931. doi: 10.1007/s10482-015-0546-8

1290

1291 Gurevich, A., Saveliev, V., Vyahhi, N., Tesler, G. (2013). QUAST: quality assessment tool for
1292 genome assemblies. **Bioinformatics**, 29(8), 1072-1075. doi: 10.1093/bioinformatics/btt086

1293

1294 Harder, C. B. et al. (2023). *Mycena* species can be opportunist-generalist plant root invaders.
1295 **Environmental Microbiology**, 25(10), 1875-1893. doi: 10.1111/1462-2920.16398

1296

1297 Hidalgo, P., Flores, M. (1994). Occurrence of the killer character in yeasts associated with
1298 Spanish wine production. **Food microbiology**, 11(2), 161-167. doi: 10.1006/fmic.1994.1019

1299

1300 Hodgson, V. J., Button, D., Walker, G. M. (1995). Anti-*Candida* activity of a novel killer toxin
1301 from the yeast *Williopsis mrakii*. **Microbiology**, 141(8), 2003-2012. doi:10.1099/13500872-
1302 141-8-2003

1303

1304 Houbraken, J. et al. (2020). Classification of *Aspergillus*, *Penicillium*, *Talaromyces* and related
1305 genera (Eurotiales): An overview of families, genera, subgenera, sections, series and species.
1306 **Studies in mycology**, 95, 5-169. doi: 10.1016/j.simyco.2020.05.002

1307

1308 Ivannikova, Y. V., Naumova, E. S., Naumov, G. I. (2006). Detection of viral dsRNA in the
1309 yeast *Saccharomyces bayanus*. **Doklady Biological Sciences**. Vol. 406. New York:
1310 Consultants Bureau, c1965-1992. doi: 10.1134/S0012496606010297

1311

1312 Ivannikova, Y. V., Naumova, E. S., Naumov, G. I. (2007). Viral dsRNA in the wine yeast
1313 *Saccharomyces bayanus* var. *uvarum*. **Research in microbiology**, 158(8-9), 638-643. doi:
1314 10.1016/j.resmic.2007.07.008

1315

1316 Jones, P., et al. (2014). InterProScan 5: genome-scale protein function classification.
1317 **Bioinformatics**, 30(9), 1236-1240. doi: 10.1093/bioinformatics/btu031

1318

1319 Julius D, Brake A, Blair L, Kunisawa R, Thorner J (1984). Isolation of the putative structural
1320 gene for the lysine-arginine-cleaving endopeptidase required for processing of yeast prepro-α-
1321 factor. **Cell**. 37(3): 1075–1089. doi: 10.1016/0092-8674(84)90442-2.

1322

1323 Jumper, J., et al. (2021). Highly accurate protein structure prediction with AlphaFold. **Nature**,
1324 596(7873), 583-589. doi: 10.1038/s41586-021-03819-2

1325

Jurjevic, Z., Peterson, S. W., Horn, B. W. (2012). *Aspergillus* section *Versicolores*: nine new species and multilocus DNA sequence based phylogeny. **IMA fungus**, 3(1), 59-79. doi:10.5598/imafungus.2012.03.01.07

Károlyi, G., Neufeld, Z., Scheuring, I. (2005). Rock-scissors-paper game in a chaotic flow: The effect of dispersion on the cyclic competition of microorganisms. **Journal of theoretical biology**, 236(1), 12-20. doi: 10.1016/j.jtbi.2005.02.012

Katoh, K., Rozewicki, J., Yamada, K. D. (2019). MAFFT online service: multiple sequence alignment, interactive sequence choice and visualization. **Briefings in Bioinformatics**, 20(4), 1160-1166. doi: 10.1093/bib/bbx108

Kerr, B., Riley, M. A., Feldman, M. W., & Bohannan, B. J. (2002). Local dispersal promotes biodiversity in a real-life game of rock–paper–scissors. **Nature**, 418(6894), 171-174. doi: 10.1038/nature00823

Klassen, R., Meinhardt, F. (2007). Linear protein-primed replicating plasmids in eukaryotic microbes. **Microbial Linear Plasmids**. Berlin, Heidelberg: Springer Berlin Heidelberg, 187-226. doi: 10.1007/7171_2007_095

Klassen, R., Schaffrath, R., Buzzini, P., Ganter, P. F. (2017). Antagonistic interactions and killer yeasts. **Yeasts in natural ecosystems: ecology**, 229-275. doi: 10.1007/978-3-319-61575-2_9

Koltin, Y., Day, P. R. (1976). Suppression of the killer phenotype in *Ustilago maydis*. **Genetics**, 82(4), 629-637. doi: 10.1093/genetics/82.4.629

Krassowski, T., et al. (2018). Evolutionary instability of CUG-Leu in the genetic code of budding yeasts. **Nature communications**, 9(1), 1887. doi: 10.1038/s41467-018-04374-7

1355

1356 Kriventseva, E. V., Kuznetsov, D., Tegenfeldt, F., Manni, M., Dias, R., Simão, F. A., Zdobnov,
1357 E. M. (2019). OrthoDB v10: sampling the diversity of animal, plant, fungal, protist, bacterial
1358 and viral genomes for evolutionary and functional annotations of orthologs. **Nucleic acids**
1359 **research**, 47(D1), D807-D811. doi: 10.1093/nar/gky1053

1360

1361 Kurtzman, C. P. (2003). Phylogenetic circumscription of *Saccharomyces*, *Kluyveromyces* and
1362 other members of the Saccharomycetaceae, and the proposal of the new genera *Lachancea*,
1363 *Nakaseomyces*, *Naumovia*, *Vanderwaltozyma* and *Zygotorulaspora*. **FEMS yeast research**,
1364 4(3), 233–245. doi: 10.1016/S1567-1356(03)00175-2.

1365

1366 Kurtzman C. P., Robnett, C. J. (1998) Identification and phylogeny of ascomycetous yeasts
1367 from analysis of nuclear large subunit (26S) ribosomal DNA partial sequences. **Antonie Van**
1368 **Leeuwenhoek,** 73(4):331–371. doi: 10.1023/a:1001761008817.

1369

1370 Kurtzman, C. P., Robnett, C. J. (2003). Phylogenetic relationships among yeasts of the
1371 'Saccharomyces complex' determined from multigene sequence analyses. **FEMS yeast**
1372 **research**, 3(4), 417-432. doi: 10.1016/S1567-1356(03)00012-6

1373

1374 Laetsch, D. R., Blaxter, M. L. (2017). BlobTools: Interrogation of genome assemblies.
1375 **F1000Research**, 6(1287), 1287. doi: 10.12688/f1000research.12232.1

1376

1377 Leal, I. R., Oliveira, P. S. (1998). Interactions between Fungus-Growing Ants (Attini), Fruits
1378 and Seeds in Cerrado Vegetation in Southeast Brasil. **Biotropica**, 30(2), 170-178. doi:
1379 10.1111/j.1744-7429.1998.tb00052.x

1380

1381 Lee, M. D., Creagh, J. W., Fredericks, L. R., Crabtree, A. M., Patel, J. S., Rowley, P. A. (2022).
1382 The Characterization of a Novel Virus Discovered in the Yeast *Pichia membranifaciens*.
1383 **Viruses**, 14(3), 594. doi: 10.3390/v14030594

1384

1385 Li, H. (2018). Minimap2: pairwise alignment for nucleotide sequences. **Bioinformatics**,
1386 34(18), 3094-3100. doi: 10.1093/bioinformatics/bty191

1387

1388 Li, H., et al. (2009). The sequence alignment/map format and SAMtools. **Bioinformatics**,
1389 25(16), 2078-2079. doi: 10.1093/bioinformatics/btp352

1390

1391 Libberton, B., Horsburgh, M. J., Brockhurst, M. A. (2015). The effects of spatial structure,
1392 frequency dependence and resistance evolution on the dynamics of toxin-mediated microbial
1393 invasions. **Evolutionary applications**, 8(7), 738-750. doi: 10.1111/eva.12284

1394

1395 Liu, H., Fu, Y., Jiang, D., Li, G., Xie, J., Cheng, J., Peng, Y., Ghabrial, S. A. Yi, X. (2010).
1396 Widespread horizontal gene transfer from double-stranded RNA viruses to eukaryotic nuclear
1397 genomes. **Journal of virology**, 84(22), 11876-11887. doi: 10.1128/jvi.00955-10

1398

1399 Liu, F., et al. (2024). *Saccharomycetaceae*: delineation of fungal genera based on
1400 phylogenomic analyses, genomic relatedness indices and genomics-based synapomorphies.
1401 **Persoonia**, 52(1), 1-21. doi: 10.3767/persoonia.2024.52.01.

1402

1403 Lomsadze, A., Ter-Hovhannisyan, V., Chernoff, Y. O., Borodovsky, M. (2005). Gene
1404 identification in novel eukaryotic genomes by self-training algorithm. **Nucleic acids research**,
1405 33(20), 6494-6506. doi: 10.1093/nar/gki937

1406

1407 Lu, S., Faris, J. D. (2019). *Fusarium graminearum* KP4-like proteins possess root growth-
1408 inhibiting activity against wheat and potentially contribute to fungal virulence in seedling rot.
1409 **Fungal genetics and biology**, 123, 1-13. doi: 10.1016/j.fgb.2018.11.002

1410

1411 Madbouly, A. K., Elyousr, K. A. A., Ismail, I. M. (2020). Biocontrol of *Monilinia fructigena*,
1412 causal agent of brown rot of apple fruit, by using endophytic yeasts. **Biological control**, 144,
1413 104239. doi: 10.1016/j.biocontrol.2020.104239

1414

1415 Magliani, W., Conti, S., Gerloni, M., Bertolotti, D., Polonelli, L. (1997). Yeast killer systems.
1416 **Clinical microbiology reviews**, 10(3), 369-400. doi: 10.1128/cmr.10.3.369

1417

1418 Makower, M., Bevan, E. A. (1963). The inheritance of a killer character in yeast
1419 (*Saccharomyces cerevisiae*). **Genetics today. Proceedings of the XI International Congress**
1420 **of Genetics**, v. 1, 202.

1421

1422 Maksimova, I. A., Glushakova, A. M., Kachalkin, A. V., Chernov, I. Y., Panteleeva, S. N.,
1423 Reznikova, Z. I. (2016). Yeast communities of Formica aquilonia colonies. **Microbiology**, 85,
1424 124-129. doi: 10.1134/S0026261716010045

1425

1426 Marquina, D., Santos, A., Peinado, J. (2002). Biology of killer yeasts. **International**
1427 **Microbiology**, 5, 65-71. doi: 10.1007/s10123-002-0066-z

1428

1429 McBride, R., Greig, D., Travisano, M. (2008). Fungal viral mutualism moderated by ploidy.
1430 **Evolution**, 62(9), 2372-2380. doi: 10.1111/j.1558-5646.2008.00443.x

1431

1432 Meyer, W., Mitchell, T.G., Freedman, E.Z., Vilgalys, R. (1993) Hybridization probes for
1433 conventional DNA fingerprinting used as single primers in the polymerase chain reaction to
1434 distinguish strains of *Cryptococcus neoformans*. **Journal of Clinical Microbiology**,
1435 31(9):2274–2280. doi: 10.1128/jcm.31.9.2274-2280.1993.

1436

1437 Meyer, S. T., Leal, I. R., Tabarelli, M., Wirth, R. (2011). Ecosystem engineering by leaf-cutting
1438 ants: nests of *Atta cephalotes* drastically alter forest structure and microclimate. **Ecological**
1439 **entomology**, 36(1), 14-24. doi: 10.1111/j.1365-2311.2010.01241.x

1440

1441 Middelbeek, E. J., Hermans, J. M., Stumm, C., Muytjens, H. L. (1980). High incidence of
1442 sensitivity to yeast killer toxins among *Candida* and *Torulopsis* isolates of human origin.
1443 **Antimicrobial Agents and Chemotherapy**, 17(3), 350-354. doi: 10.1128/aac.17.3.350

1444

1445 Möller, A. (1893). **Die Pilzgärten einiger südamerikanischer Ameisen** (No. 6). G. Fischer.

1446

1447 Myers, J. M., et al. (2020). Survey of early-diverging lineages of fungi reveals abundant and
1448 diverse mycoviruses. **mBio** 11: e02027-20. doi: 10.1128/mbio.02027-20

1449

Naumov, G. I., Ivannikova, Y. V., Chernov, I. Y., Naumova, E. S. (2009). Natural polymorphism of the plasmid double-stranded RNA of the *Saccharomyces* yeasts. **Microbiology**, 78, 208-213. doi: 10.1134/S0026261709020118

Nguyen, L. T., Schmidt, H. A., Von Haeseler, A., Minh, B. Q. (2015). IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. **Molecular Biology and Evolution**, 32(1), 268-274. doi: 10.1093/molbev/msu300

Nguyen, N. H., Suh, S. O., Blackwell, M. (2007). Five novel *Candida* species in insect-associated yeast clades isolated from Neuroptera and other insects. **Mycologia**, 99(6), 842–858. doi: 10.1080/15572536.2007.11832516

Nomoto, H., Kitano, K., Shimazaki, T., Kodama, K., Hara, S. (1984). Distribution of killer yeasts in the genus *Hansenula*. **Agricultural and biological chemistry**, 48(3), 807-809. doi: 10.1271/bbb1961.48.807

Nygaard, S., et al. (2016). Reciprocal genomic evolution in the ant–fungus agricultural symbiosis. **Nature Communications**, 7(1), 12233. doi: 10.1038/ncomms12233

Okada, R., Kiyota, E., Moriyama, H., Fukuhara, T., Natsuaki, T. (2015). A simple and rapid method to purify viral dsRNA from plant and fungal tissue. **Journal of General Plant Pathology**, 81, 103-107. doi: 10.1007/s10327-014-0575-6

Pagnocca, F. C., Carreiro, S. C., Bueno, O. C., Hebling, M. J., Da Silva, O. A. (1996). Microbiological changes in the nests of leaf-cutting ants fed on sesame leaves. **Journal of Applied Entomology**, 120(1-5), 317–320. doi: 10.1111/j.1439-0418.1996.tb01612.x

Pagnocca, F. C., Rodrigues, A., Nagamoto, N. S., Bacci, M. (2008). Yeasts and filamentous fungi carried by the gynes of leaf-cutting ants. **Antonie van Leeuwenhoek**, 94(4), 517–526. doi: 10.1007/s10482-008-9268-5

1480

1481 Perez, M. F., Contreras, L., Garnica, N. M., Fernández-Zenoff, M. V., Farías, M. E., Sepulveda,
1482 M., Ramallo, J., Dib, J. R. (2016). Native killer yeasts as biocontrol agents of postharvest fungal
1483 diseases in lemons. **PloS One**, 11(10), e0165590. doi: 10.1371/journal.pone.0165590

1484

1485 Peterson, S. W., Ito, Y., Horn, B. W., Goto, T. (2001). *Aspergillus bombycis*, a new
1486 aflatoxigenic species and genetic variation in its sibling species, *A. nomius*. **Mycologia**, 93(4),
1487 689-703. doi: 10.1080/00275514.2001.12063200

1488

1489 Pfeiffer, I., Golubev, W. I., Farkas, Z., Kucsera, J., Golubev, N. (2004). Mycocin production
1490 in *Cryptococcus aquaticus*. **Antonie Van Leeuwenhoek**, 86, 369-375. doi: 10.1007/s10482-
1491 004-0888-0

1492

1493 Philliskirk, G., Young, T. W. (1975). The occurrence of killer character in yeasts of various
1494 genera. **Antonie van Leeuwenhoek**, 41, 147-151. doi: 10.1007/BF02565046

1495

1496 Pieczynska, M. D., de Visser, J. A. G., Korona, R. (2013). Incidence of symbiotic dsRNA
1497 'killer' viruses in wild and domesticated yeast. **FEMS Yeast Research**, 13(8), 856–859. doi:
1498 10.1111/1567-1364.12086

1499

1500 Pieczynska, M. D., Wloch-Salamon, D., Korona, R., de Visser, J. A. G. (2016). Rapid multiple-
1501 level coevolution in experimental populations of yeast killer and nonkiller strains. **Evolution**,
1502 70(6), 1342-1353. doi: 10.1111/evo.12945

1503

1504 Pintar, J., Starmer, W. T. (2003). The costs and benefits of killer toxin production by the yeast
1505 Pichia kluyveri. **Antonie van Leeuwenhoek**, 83, 89-97. doi: 10.1023/A:0000000089097

1506

1507 Posit team (2023). RStudio: Integrated Development Environment for R. Posit Software, PBC,
1508 Boston, MA. URL http://www.posit.co/.

1509

1510 R Core Team. (2023). R: A Language and Environment for Statistical Computing.; **R
1511 Foundation for Statistical Computing: Vienna, Austria**; Available online: https://www.R-
1512 project.org/

1513

Ramachandran GN, Ramakrishnan C, Sasisekharan V (1963). Stereochemistry of polypeptide chain configurations. **J Mol Biol**. 7(1): 95–99. doi: 10.1016/s0022-2836(63)80023-6.

1516

Rambaut, A., 2016. **Figtree v. 1.4.3**. http://tree.bio.ed.ac.uk/software/figtree/2016.

1518

Ramírez, M., Velázquez, R., Maqueda, M., López-Piñeiro, A., Ribas, J. C. (2015). A new wine *Torulaspora delbrueckii* killer strain with broad antifungal activity and its toxin-encoding double-stranded RNA virus. **Frontiers in microbiology**, 6, 983. doi: 10.3389/fmicb.2015.00983

1523

Robledo-Leal, E., Elisondo-Zatuche, M., Treviño-Rangel, R. J., González, G. M., Hernández-Luna, C. (2016). Isolation of killer yeasts from ants of the genus *Atta* and their effect on the red tomato's fungal pathogen *Geotrichum candidum*. **Mexican Journal of Phytopathology**, 34, 258–269. doi: 10.18781/r.mex.fit.1605-3

1528

Rodrigues, A., Bacci Jr, M., Mueller, U. G., Ortiz, A., Pagnocca, F. C. (2008). Microfungal "weeds" in the leafcutter ant symbiosis. **Microbial Ecology**, 56, 604-614. doi: 10.1007/s00248-008-9380-0

1532

Rodrigues, A., Cable, R. N., Mueller, U. G., Bacci, M., Pagnocca, F. C. (2009). Antagonistic interactions between garden yeasts and microfungal garden pathogens of leaf-cutting ants. **Antonie Van Leeuwenhoek**, 96, 331-342. doi: 10.1007/s10482-009-9350-7

1536

Rodríguez-Cousiño, N., Gómez, P., Esteban, R. (2017). Variation and distribution of LA helper totiviruses in *Saccharomyces* sensu stricto yeasts producing different killer toxins. Toxins, 9(10), 313. doi: doi.org/10.3390/toxins9100313

1540

Rodríguez-Cousiño, N., Maqueda, M., Ambrona, J., Zamora, E., Esteban, R., Ramírez, M. (2011). A new wine *Saccharomyces cerevisiae* killer toxin (Klus), encoded by a double-stranded RNA virus, with broad antifungal activity is evolutionarily related to a chromosomal

host gene. **Applied and environmental microbiology**, 77(5), 1822-1832. doi: 10.1128/AEM.02501-10

Rodriguez Coy, L., Plummer, K. M., Khalifa, M. E., MacDiarmid, R. M. (2022). Mycovirus-encoded suppressors of RNA silencing: Possible allies or enemies in the use of RNAi to control fungal disease in crops. **Frontiers in Fungal Biology**, 3, 965781. doi: 10.3389/ffunb.2022.965781

Ronque, M. U., Feitosa, R. M., Oliveira, P. S. (2019). Natural history and ecology of fungus-farming ants: a field study in Atlantic rainforest. **Insectes Sociaux**, 66(3), 375-387. doi: 10.1007/s00040-019-00695-y

Rowley, P. A. (2017). The frenemies within: viruses, retrotransposons and plasmids that naturally infect *Saccharomyces* yeasts. **Yeast**, 34(7), 279-292. doi: 10.1002/yea.3234

Rupert, C. B., Rusche, L. N. (2022). The Pathogenic Yeast *Candida parapsilosis* Forms Pseudohyphae through Different Signaling Pathways Depending on the Available Carbon Source. **Msphere**, 7(3), e00029-22. doi: 10.1128/msphere.00029-22

Satwika, D., Klassen, R., Meinhardt, F. (2012). Repeated capture of a cytoplasmic linear plasmid by the host nucleus in *Debaryomyces hansenii*. **Yeast**, 29(3-4), 145-154. doi: 10.1002/yea.2893

Schaffrath, R., Meinhardt, F., Klassen, R. (2018). Yeast killer toxins: Fundamentals and applications. **Physiology and Genetics: Selected Basic and Applied Aspects**, 87-118. doi: 10.1007/978-3-319-71740-1_3

Schmitt, M. J., Neuhausen, F. (1994). Killer toxin-secreting double-stranded RNA mycoviruses in the yeasts *Hanseniaspora uvarum* and *Zygosaccharomyces bailii*. **Journal of virology**, 68(3), 1765-1772. doi: 10.1128/jvi.68.3.1765-1772.1994

1575   Schneider, C. A., Rasband, W. S., Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of
1576   image analysis. **Nature methods**, 9(7), 671-675. doi: 10.1038/nmeth.2089

1577

1578   Schultz, T. R., Brady, S. G. (2008). Major evolutionary transitions in ant agriculture.
1579   **Proceedings of the National Academy of Sciences**, 105(14), 5435–5440. doi:
1580   10.1073/pnas.0711024105

1581

1582   Schultz, T. R., Sosa-Calvo, J., Brady, S. G., Lopes, C. T., Mueller, U. G., Bacci Jr, M.,
1583   Vasconcelos, H. L. (2015). The most relictual fungus-farming ant species cultivates the most
1584   recently evolved and highly domesticated fungal symbiont species. **The American Naturalist**,
1585   185(5), 693–703. doi: 10.1086/680501

1586

1587   Segers, G. C., van Wezel, R., Zhang, X., Hong, Y., Nuss, D. L. (2006). Hypovirus papain-like
1588   protease p29 suppresses RNA silencing in the natural fungal host and in a heterologous plant
1589   system. **Eukaryotic Cell**, 5(6), 896-904. doi: 10.1128/EC.00373-05

1590

1591   Segers, G. C., Zhang, X., Deng, F., Sun, Q., Nuss, D. L. (2007). Evidence that RNA silencing
1592   functions as an antiviral defense mechanism in fungi. **Proceedings of the National Academy**
1593   **of Sciences**, 104(31), 12902-12906. doi: 10.1073/pnas.070250010

1594

1595   Seppey, M., Manni, M., Zdobnov, E.M. (2019). BUSCO: Assessing Genome Assembly and
1596   Annotation Completeness. In: Kollmar, M. (eds) Gene Prediction. **Methods in Molecular**
1597   **Biology**, vol 1962. Humana, New York, NY. doi: 10.1007/978-1-4939-9173-0_14

1598

1599   Silva, A., Bacci Jr, M., de Siqueira, C. G., Bueno, O. C., Pagnocca, F. C., Hebling, M. J. A.
1600   (2003). Survival of *Atta sexdens* workers on different food sources. **Journal of Insect**
1601   **Physiology**, 49(4), 307-313. doi: 10.1016/S0022-1910(03)00004-0

1602

1603   Silva, A., Bacci Jr, M., Pagnocca, F. C., Bueno, O. C., Hebling, M. J. A. (2006). Starch
1604   metabolism in *Leucoagaricus gongylophorus*, the symbiotic fungus of leaf-cutting ants.
1605   **Microbiological research**, 161(4), 299-303. doi: 10.1016/j.micres.2005.11.001

1606

1607 Somers, J. M., Bevan, E. A. (1969). The inheritance of the killer character in yeast. **Genetics**
1608 **Research**, 13(1), 71-83. doi: 10.1017/S0016672300002743

1609

1610 Spoel D. V. D., Lindahl E, Hess B, Groenhof G, Mark AE, Berendsen HJC (2005).
1611 GROMACS: Fast, flexible, and free. **J Comput Chem**. 26(16): 1701–1718. doi:
1612 10.1002/jcc.20291.

1613

1614 Stanke, M., Morgenstern, B. (2005). AUGUSTUS: a web server for gene prediction in
1615 eukaryotes that allows user-defined constraints. **Nucleic acids research**, 33(suppl_2), W465-
1616 W467. doi: 10.1093/nar/gki458

1617

1618 Starmer, W. T., Ganter, P. F., Aberdeen, V., Lachance, M. A., Phaff, H. J. (1987). The
1619 ecological role of killer yeasts in natural communities of yeasts. **Canadian journal of**
1620 **microbiology**, 33(9), 783-796. doi: 10.1139/m87-134

1621

1622 Sternberg, L. D. S., Pinzon, M. C., Moreira, M. Z., Moutinho, P., Rojas, E. I., Herre, E. A.
1623 (2007). Plants use macronutrients accumulated in leaf-cutting ant nests. **Proceedings of the**
1624 **Royal Society B: Biological Sciences**, 274(1608), 315-321. doi: 10.1098/rspb.2006.3746

1625

1626 Stopiglia, C. D. O., Heidrich, D., Sorrentino, J. M., Vieira, F. J., Landell, M. F., Valente, P.,
1627 Scroferneker, M. L. (2014). Susceptibility of species within the *Sporothrix schenckii* complex
1628 to a panel of killer yeasts. **Journal of Basic Microbiology**, 54(6), 578-584. doi:
1629 10.1002/jobm.201200516

1630

1631 Studer G, Rempfer C, Waterhouse AM, Gumienny R, Haas J, Schwede T (2019).
1632 QMEANDisCo – Distance Constraints Applied on Model Quality Estimation. **Bioinformatics**.
1633 36(6): 1765–1771. doi: 10.1093/bioinformatics/btz828.

1634

1635 Stumm, C., Hermans, J. M. H., Middelbeek, E. J., Croes, A. F., De Vries, G. J. M. L. (1977).
1636 Killer-sensitive relationships in yeasts from natural habitats. **Antonie van Leeuwenhoek**, 43,
1637 125-128. doi: 10.1007/BF00395667

1638

Suh, S. O., Blackwell, M. (2005). Four new yeasts in the *Candida mesenterica* clade associated
with   basidiocarp-feeding   beetles.   **Mycologia**,   97(1),   167-177.   doi:
10.1080/15572536.2006.11832850

Suh, S. O., McHugh, J. V., Pollock, D. D., Backwell, M. (2005). The beetle gut: a hyperdiverse
source   of   novel   yeasts.   **Mycological   Research**,   109(3),   261–265.   doi:
10.1017/S0953756205002388

Summerell, B. A., Laurence, M. H., Liew, E. C., Leslie, J. F. (2010). Biogeography and
phylogeography of *Fusarium*: a review. **Fungal Diversity**, 44(1), 3-13. doi: 10.1007/s13225-
010-0060-2

Suzuki, C., Nikkuni, S. (1994). The primary and subunit structure of a novel type killer toxin
produced by a halotolerant yeast, *Pichia farinosa*. **Journal of Biological Chemistry**, 269(4),
3041-3046. doi: 10.1016/S0021-9258(17)42044-8

Taylor, D. J., Ballinger, M. J., Bowman, S. M., Bruenn, J. A. (2013). Virus-host co-evolution
under a modified nuclear genetic code. **PeerJ**, 1, e50. doi: 10.7717/peerj.50

Taylor, D. J., Bruenn, J. (2009). The evolution of novel fungal genes from non-retroviral RNA
viruses. **BMC biology**, 7, 1-11. doi: 10.1186/1741-7007-7-88

Thoen, E., et al. (2020). In vitro evidence of root colonization suggests ecological versatility in
the genus *Mycena*. **New Phytologist**, 227(2), 601-612. doi: 10.1111/nph.16545

Travers-Cook, T. J., Jokela, J., Buser, C. C. (2023). The evolutionary ecology of fungal killer
phenotypes.   **Proceedings   of   the   Royal   Society   B**,   290,   20231108.   doi:
10.1098/rspb.2023.1108

Trindade, R. C., Resende, M. A., Silva, C. M., Rosa, C. A. (2002). Yeasts associated with fresh
and frozen pulps of Brasilian tropical fruits. **Systematic and Applied Microbiology**, 25(2),
294-300. doi: 10.1078/0723-2020-00089

1671

1672 Vadasz, A. S., Vadasz, P., Gupthar, A. S., Abashar, M. E. (2003). Theoretical and experimental
1673 evidence of extinction and coexistence of killer and sensitive strains of yeast grown as a mixed
1674 culture in water. **International journal of food microbiology**, 84(2), 157-174. doi:
1675 10.1016/S0168-1605(02)00417-8

1676

1677 Vaughan-Martini, A. (1995). *Saccharomyces barnetti* and *Saccharomyces spencerorum*: two
1678 new species of *Saccharomyces* sensu lato (van der Walt). **Antonie van Leeuwenhoek**, 68, 111-
1679 118. doi: 10.1007/BF00873098

1680

1681 Vijayraghavan, S., Kozmin, S. G., Strope, P. K., Skelly, D. A., Magwene, P. M., Dietrich, F.
1682 S., McCusker, J. H. (2023). RNA viruses, M satellites, chromosomal killer genes, and
1683 killer/nonkiller phenotypes in the 100-genomes *S. cerevisiae* strains. **G3: Genes, Genomes,**
1684 **Genetics**, 13(10), jkad167. doi: 10.1093/g3journal/jkad167

1685

1686 Villan Larios, et al. (2023). Exploring the Mycovirus Universe: Identification, Diversity, and
1687 Biotechnological Applications. **Journal of Fungi**, 9(3), 361. doi: 10.3390/jof9030361

1688

1689 Walker, G. M., Mcleod, A. H., Hodgson, V. J. (1995). Interactions between killer yeasts and
1690 pathogenic fungi. **FEMS microbiology letters**, 127(3), 213-222. doi: 10.1111/j.1574-
1691 6968.1995.tb07476.x

1692

1693 Warnes, G., Bolker, B., Bonebakker, L., Gentleman, R., Huber, W., Liaw, A., Lumley, T.,
1694 Maechler, M., Magnusson, A., Moeller, S., Schwartz, M., Venables, B., Galili, T. (2024).
1695 *gplots*: Various R Programming Tools for Plotting Data. **R package version 3.2.0**. doi:
1696 10.32614/CRAN.package.gplots

1697

1698 Weber, N. A. (1972). The fungus-culturing behavior of ants. **American Zoologist**, 12(3), 577–
1699 587. doi: 10.1093/icb/12.3.577

1700

1701 Wickham H (2016). ggplot2: Elegant Graphics for Data Analysis. Springer-Verlag New York.
1702 ISBN 978-3-319-24277-4, https://ggplot2.tidyverse.org.

1703

Wloch-Salamon, D. M., Gerla, D., Hoekstra, R. F., de Visser, J. A. G. (2008). Effect of dispersal and nutrient availability on the competitive ability of toxin-producing yeast. **Proceedings of the Royal Society B: Biological Sciences**, 275(1634), 535-541. doi: 10.1098/rspb.2007.1461

1708

Wójcik, M., Kordowska-Wiater, M. (2015). The occurrence of killer activity in yeasts isolated from natural habitats. **Acta Biochimica Polonica**, 62(4). doi: 10.18388/abp.2015_1141

1711

Woods, D. R., Bevan, E. A. (1968). Studies on the nature of the killer factor produced by *Saccharomyces cerevisiae*. **Microbiology**, 51(1), 115–126. doi: 10.1099/00221287-51-1-115

1714

Worsham, P. L., Bolen, P. L. (1990). Killer toxin production in *Pichia acaciae* is associated with linear DNA plasmids. **Current genetics**, 18, 77-80. doi: 10.1007/BF00321119

1717

Zorg, J., Kilian, S., Radler, F. (1988). Killer toxin producing strains of the yeasts *Hanseniaspora uvarum* and *Pichia kluyveri*. **Archives of microbiology**, 149, 261-267. doi: 10.1007/BF00422015

A

B

C

Yamadazyma paraaseri LESF 1412 ▲
98
Yamadazyma sp. LESF 1490 ▲
100
Candida sp. LESF 1433 ▲ ☠
97
Candida vrieseae
Candida sp. LESF 1493 ◆
Candida margitis LESF 1378 ●
Candida orthopsilosis LESF 1402 ●
Candida parapsilosis LESF 1347 ★
98
100
Lodderomyces elongisporus LESF 1360 ●
99
Candida albicans LESF 1442 ●
95
Candida sp. LESF 1461 ▲ ☠
Candida sinolaborantium LESF 1467 ▲
100
Scheffersomyces sp. LESF 1457 ▲
81
Meyerozyma guilliermondii LESF 1444 ●
Meyerozyma carpophila LESF 1389 ●
100
Meyerozyma caribbica LESF 1452 ●
Meyerozyma smithsonii
Priceomyces sp. LESF 1391 ●
Schwanniomyces vanrijiae LESF 1478 ▲■◆ ☠
Debaryomyces hansenii LESF 1405 ●
94
Metschnikowia koreensis LESF 1422 ■
99
Metschnikowia sp. LESF 1343 ■
79
Metschnikowia sp. LESF 1383 ▲
71
Metschnikowia sp. LESF 1355 ○
98
99
Candida blattae LESF 1417 ▲◆ ☠
Candida sp. LESF 1395 ●
Candida sp. LESF 1413 ▲
81
Kodamaea ohmeri LESF 1474 ● ☠
Diutina catenulata LESF 1446 ●
Saccharomycetales sp. LESF 1445 ●
Hyphopichia sp. LESF 1344 ▲
Hyphopichia burtonii LESF 1503 ■
Kurtzmaniella sp. LESF 1491 ▲
Kurtzmaniella quercitrusa LESF 1421 ▲◆
Kurtzmaniella sp. LESF 1473 ▲
99
Candida railenensis LESF 1437 ●★
Babjeviella inositovora
70
Limtongozyma sp. LESF 1404 ▲
Limtongozyma cylindracea LESF 1500 ▲
72
Candida spp.
Pichia kudriavzevii LESF 1411 ●
Cyberlindnera sp. LESF 1438 ▲
100
Wickerhamomyces ciferrii LESF 1397 ■ ☠
Wickerhamomyces sp. LESF 1460 ▲
97
Hanseniaspora opuntiae LESF 1470 ●
92
Hanseniaspora thailandica LESF 1297 ●
Grigorovia sp. LESF 1345 ○
100
Jamesozyma spencerorum LESF 1468 ▲ ☠
Kluyveromyces hubeiensis
Starmerella sp. LESF 1454 ●
100
Starmerella bombicola LESF 1472 ●
100
Starmerella sp. LESF 1293 ●
Starmerella etchellsii LESF 1300 ●
Starmerella meliponinorum LESF 1496 ●
72
Wickerhamiella sp. LESF 1495 ★
74
97
100
Wickerhamiella sp. LESF 1506 ●
100
Wickerhamiella occidentalis LESF 1296 ●▲
Yarrowia lipolytica LESF 1385 ●
Sugiyamaella smithiae LESF 1336 ★
Sugiyamaella carassensis LESF 1463 ○▲
100
Sugiyamaella bonitensis LESF 1465 ▲
Sugiyamaella ligni LESF 1459 ▲
89
Sugiyamaella boreocaroliniensis LESF 1366 ★
100
Sugiyamaella floridensis
100
Blastobotrys proliferans LESF 1481 ●
Schizosaccharomyces pombe

Serinales

Pichiales

Phaffomycetales

Saccharomycodales

Saccharomycetales

Dipodascales

0.2

● *Acromyrmex coronatus*   ★ *Mycocepurus goeldii*
■ *Mycetomoellerius tucumanus*   ◆ *Apterostigma goniodes*
▲ *Mycetophylax* aff. *auritus*   ☠ Killer yeast

A Toxin Susceptible Yeasts

Killer Yeasts

W. ciferrii
S. vanrijiae
Wickerhamomyces sp.
Candida sp.
K. ohmeri
K28
S. vanrijiae
D. catenulata
K21
K45
K62
K2
K74
C. blattae
K1
K1L
Klus
C. sinolaborantium

Ogataea sp.
N. castellii
Cyberlindnera sp.
S. bombicola
S. meliponinorum
S. etchellsii
S. cerevisiae
S. mikatae
S. paradoxus
P. kudriavzevii
S. kudriavzevii
M. caribbica
H. thailandica
S. vanrijiae
M. caribbica
Y. paraaseri
N. delphensis
N. castellii
M. koreensis
K. quercitrusa
Cyberlindnera sp.
S. eubayanus
K. africana
D. catenulata
Y. lipolytica
S. carassensis
S. ligni
S. bonitensis
K. ohmeri
Y. riverae
K. spencerorum

B

Yeast isolates

75
50
25
0

Killer
Non-killer

● *A. coronatus*
■ *M. tucumanus*
▲ *M. auritus*
★ *M. goeldii*
◆ *A. goniodes*

Ant species

C

Same source
Different source

Killed
Not killed

Lawn yeast
Killer yeast
Inhibition zone
Methylene blue halo

Clear zone

+halo   -halo

No clear zone

Halo only   -halo

● *Acromyrmex coronatus*
■ *Mycetomoellerius tucumanus*
▲ *Mycetophylax* aff. *auritus*
★ *Mycocepurus goeldii*
◆ *Apterostigma goniodes*

A

S. cerevisiae (Klus)

− CHX   + CHX

Kbp
5 — ◄ Totivirus
4 —
3 —
2 — ◄ Satellite dsRNA (Mlus)
1 —

− +
CHX

B

Elongated growth ratio

**

+   −
Halo

C

CFU×10^2 mL^{-1}

ns

+   −
Halo

D

100 ┌ 83 — *Mycena* spp.
85 │   └ *Roridomyces roridus* (KAJ7614860)
     └ *Penicillium verhagenii* (XP_057019693)
86 ┌ 100 — *Colletotrichum* spp.
84 │   └ *Lasiodiplodia mahajangana* (KAJ8128189)
99 │ ┌ *Lobaria immixta* (MCJ1271286)
   │ └ 98 *Pseudocyphellaria aurata* (MCJ1467775)
88 ┌ 100 — *Trichoderma* spp.
     └ *Aspergillus* spp.
100 ┌ 100 *Kazachstania africana* (XP_003958467)
    │ 82 └ *Kazachstania africana* (XP_003957740)
    │   └ *Suhomyces tanzawaensis* (XP_020062392)
    │ 100 ┌ *Candida anglica* (CAK7918227)
    │   └ *Candida anglica* (CAK7894508)
    │ 100 ┌ *Scheffersomyces stipitis* (XP_001388021)
    │   └ *Scheffersomyces stipitis* (XP_001383947)
    │   **Candida sinolaborantium** Ksino (LESF 1467)
    └ *Kazachstania africana* (XP_003957274)
      Fulvia fulva (XP_047769465)
99 ┌ *Botrytis/Botryotinia* spp.
100 │ └ *Sclerotinia sclerotiorum* (XP_001588904)
95 │   *Cercospora kikuchii* (XP_044657970)
96 ┌ *Fusarium* spp.
   │ *Ramularia collo-cygni* (XP_023627751)
98 ┌ 100 ┌ 100 *Fusarium* spp.
   │   └ *Ilyonectria* spp.
93 │ 100 └ *Thelonectria olida* (KAH6887094)
99 └ *Fusarium* spp.
   Naumovozyma dairenensis (XP 003671032)
Saccharomyces cerevisiae Mlus (ADG64740)

0.5

A

B   C

A

GAL1 ──► Ksino

S. cerevisiae + Ksino

C. sino   L186S   WT   n/a

GAL

DEX

C

Area (cm²)

pH: 3.5 4.0 4.5 5.0 5.5 6.0 6.5

a a   b b b
                  0 0 0

D

Temperature °C: 17 20 25 30 37

b b   a
a   0