Evaluating Plant-Microbe Associations in Response to Environmental Stressors to Enhance

Salt Marsh Restoration

RUNNING TITLE: Salt marsh plant-microbiome restoration

Kai A. Davis[1], Mary-Margaret S. McKinney[2], Rachel K. Gittman[1,3], Ariane L. Peralta[1,*]

[1]Department of Biology, [2]Integrated Coastal Sciences, East Carolina University, Greenville, North

Carolina, USA, [3]Coastal Studies Institute, Wanchese, North Carolina, USA

*Corresponding Author: phone: 252-328-2712, email: peraltaa@ecu.edu

ABSTRACT

Microorganisms can enhance nutrient acquisition or suppress diseases from pathogens, while plants can provide carbon resources and oxygen to root-associated microbes. However, human activities have altered nutrient cycles and disrupted such mutualisms. Therefore, we need to understand how to promote positive plant-microbe associations to aid in restoring coastal wetland ecosystems where human stressors and climate change (e.g., hurricanes, sea-level rise) challenge restoration outcomes. This study examined how salinity stressors influence plant-microbe relationships, where we hypothesize that the presence of microbes will buffer stressor effects. We used a whole sediment inocula approach and exposed marsh cordgrass (*Sporobolus alterniflorus*) plugs to a replicated factorial experiment with three levels of microbiome addition (microbial inocula, autoclaved microbial inocula, no microbe control) and two levels of salinity (<0.5 psu, 20 psu), replicated ten times. We added microbial inocula from the marsh site with autoclaved soilless media and exposed half the plugs to saltwater (20 psu) and half to freshwater (<0.5 psu). Results revealed that marsh microbial inocula additions during early plant development may ameliorate salinity stressors and could be critical for future restoration efforts. The addition of marsh microbial inocula provided a rescue effect from salinity stress, observed in plant height over time. Aboveground and belowground biomass, bacterial diversity H', and bacterial evenness J' were similar across microbial and water treatments; however, water and microbial treatments significantly influenced the bacterial community composition. This work provides evidence that microbiome stewardship buffers against environmental stressors and could promote plant establishment for coastal wetland restoration.

Keywords: soil microbial communities, salinity stress, microbiome stewardship, coastal protection

INTRODUCTION

Salt marshes, coastal wetlands, and other coastal ecosystems aid in ameliorating erosion

and flooding, protect people and property from storm surges, and provide habitat for fisheries and

many other species (Arkema et al., 2013; Buchanan et al., 2022). Despite their high ecological and

economic importance, these coastal habitats are under threat due to anthropogenic activities (e.g.,

land development, dredging) and climate change (e.g., accelerated sea level rise, increased

frequency of storms) (Valiela et al., 2018; Leonardi et al., 2018; Rudershausen et al., 2021;

Buchanan et al., 2022). Unfortunately, current restoration efforts are failing to keep pace with

current and future losses (Gittman et al. 2019), highlighting the necessity for improved salt marsh

restoration techniques (Gittman et al. 2024).

The planting of marsh smooth cordgrass *Sporobolus alterniflorus* (also known as *Spartina*

alterniflora) is a common approach to salt marsh restoration efforts, as the plant is a foundational

species that can withstand a wide range of salinities, inundation periods, and anoxic stress (Gedan

& Bertness, 2010). These foundation or habitat-forming species alter their environment, making it

possible for other organisms to establish and endure within an ecosystem (Bruno, 2000). However,

when commercially raised nursery plants are transferred to field restoration sites, they are exposed

to abiotic stressors (e.g., salinity and wave energy) and biotic factors (e.g., species interactions)

that can induce stress and hinder restoration (Billah et al., 2022). The effect of exposing *S.*

alterniflorus plants to abiotic stressors prior to establishment at a restoration site is unknown. In

addition, the understanding of plant-microbe associations between marsh soil microbiomes and *S.*

alterniflorus and its influence on plant growth and survival is limited. Therefore, an examination of

the effect of salinity stress and microbial inoculation on marsh plant productivity would further the

understanding of ecological mechanisms and aid in restoring coastal wetland ecosystems.

In a broader context, microorganisms can enhance the nutrient acquisition of plants,

support growth, and suppress diseases from pathogens (Mariotte et al., 2018; Farrer et al., 2022). In

turn, plants can provide carbon resources and oxygen to root-associated microorganisms (Mariotte

et al., 2018). Human activities and climate change can and have altered these plant-microbe

relationships in ways that promoted or suppressed plant growth. For example, anthropogenic-

caused eutrophication and pollution have disrupted nutrient cycles and have led to losses in

microbial biodiversity (Cavicchioli et al., 2019), while sea level rise and extreme climatic events

damage plants and disrupt mutualisms (Zabin et al., 2022). Conversely, the practice of microbiome

stewardship — the intentional understanding and manipulation of microbial communities — can

lead to increased microbiome diversity, evenness, and homeostasis (Peixoto et al., 2022). Growing

evidence has also indicated that microbial communities can increase plant survivorship and

growth in high-stress environments, aiding ecological restoration efforts (McHugh & Dighton, 2004;

Mariotte et al., 2018).

With the urgent need for enhanced restoration techniques, we aim to understand how to

promote positive plant-microbe associations to aid in restoring coastal wetland ecosystems where

human stressors and climate change challenge restoration outcomes. In this investigation, we

evaluated the interactive effects of salinity stress and marsh sediment microbial inoculation on *S.*

alterniflorus productivity. Conducting a replicated factorial with two levels of salinity and three

levels of microbial inocula treatment, we measured the plant height and biomass (aboveground

and belowground) of *S. alterniflorus* plugs after a 2-month duration. Additionally, we extracted

genomic DNA from the sediment samples to characterize the root-associated microbiomes of the

laboratory-treated plugs. We hypothesized that *S. alterniflorus* plugs inoculated with restoration-

area microbial communities would have higher survivorship and growth rates than control plugs

when salt-stressed.

METHODS

Microbiome x Salinity Stress Experimental Design

To test our hypothesis that *S. alterniflorus* plugs inoculated with restoration-area microbial

communities would have higher survivorship and growth rates than control plugs when salt-

stressed, we conducted a factorial experiment with three levels of microbiome addition (No

Addition, Autoclaved Inocula, Added Inocula) and two levels of salinity (<0.5 psu, 20 psu),

replicated ten times (n=60) (Figure 2). Marsh sediment was collected from Hammocks Beach State

Park (Swansboro, North Carolina), while the marsh plugs, *Sporobolus alterniflorus*, were purchased

from a commercial plant nursery (Garner's Landscaping, Inc., Newport, North Carolina). To prepare

the microbial treatments, we combined microbial inocula with Fafard 3B potting mix, where 5% of

the total volume was marsh sediment and 95% was autoclaved potting mix (following the methods

of Lau & Lennon, 2011). While autoclaving can influence the chemistry of soil organic matter and

the accessibility of plant nutrients, it creates a benchmark for sterilization and microbial

community establishment. Any variations observed between microbial treatments can be

attributed to the introduction of inocula containing either live or deceased soil microorganisms

(Lau & Lennon, 2011).

The combination of untreated marsh sediment and autoclaved potting mix represented the

'Added Inocula' treatment. For the 'Autoclaved Inocula' treatment, marsh sediment was autoclaved

before being combined, with the purpose of examining whether there were abiotic effects on plant

productivity. For the 'No Addition' treatment, 100% of the volume of the microbial treatment was

comprised of autoclaved potting mix. The plugs were transferred to 4.5" square pots with their

respective treatments on 08 June and allowed to acclimate to the soil conditions for four weeks

under laboratory conditions (room temperature at about 25 °C next to a window for sunlight) . We

measured the maximum plant height and number of live stems biweekly, with changes in average

stem height functioning as a proxy for growth. The first height was recorded on 23 June, which was

the day when the plants were moved from the laboratory to the outside environment. Following

acclimation, weekly salt stress treatment began on 09 July and ran for four weeks.

For the salinity treatment, we exposed each potted plug to 15 minutes of 20 psu saltwater or

freshwater. Saltwater was made with Instant Ocean Sea Salt, with a general ratio of ~300 g Instant

Ocean to 14 L of tap water. This established a concentration of ~20 psu, which represented salinity

levels at Hammocks Beach State Park, where salinity ranges from 20 to 35 psu (personal

communication, park manager). Although the duration of salt exposure in our experimental setup

may be shorter compared to natural field conditions experienced by plants, it was anticipated that

the deposited salt would gradually accumulate in the soil due to evaporation, mimicking real-world

scenarios. This deliberate manipulation allowed us to specifically examine the effects of salt stress

on plant physiology, distinct from the stress induced by prolonged inundation, which typically leads

to anoxia. Thus, our experimental design ensured a focused investigation into salt stress dynamics

without confounding factors associated with prolonged inundation.

Plant and Soil Sample Processing

At the end of the experiment, after eight weeks, we harvested aboveground biomass and

roots, dried the plant biomass samples at 60 °C for 48 hours, and weighed aboveground and root

biomass. For soil bacterial characterization and analysis, we homogenized the remaining soil,

transferred soil into sterile WhirlPak bags, and stored samples at -20 °C until DNA extraction and

16S rRNA amplicon sequencing (described below).

Field Site Description

The field study focused on *S. alterniflorus* growth and survival across an elevation gradient. The

field site chosen for planting was a small section of sandy, eroded shoreline behind a recently

installed offshore sill at Hammocks Beach State Park on Queens Creek in Swansboro, North

Carolina. The section of beach contained roughly equal areas above and below the mean high

water (MHW) line (Figure 1).

Field Plant and Soil Sampling

Commercially grown *S. alterniflorus* plugs were obtained on 1 June 2023 from a local

nursery that routinely grows *S. alterniflorus* plugs from locally collected seeds for salt marsh

restoration projects. The number of stems in each cell in the nursery trays was counted and the

length of the longest leaf in each cell was measured to 0.1 centimeter (cm). All plants were then

trimmed to a total height of 20 cm. Nine rows of plants were installed (n=270): five above MHW

(n=142) and four below (n=128), with approximately 7 cm between plugs and 25 cm between rows

(Figure 1). The remaining plugs were relocated to the laboratory for salinity and microbial

treatments.

The location and elevation of each row were measured at the time of planting using a

Trimble Virtual Reference Station (VRS; Trimble, Sunnyvale, California, USA), Real-Time Kinematic

(RTK), and Global Positioning System (GPS). Elevations ranged from an average of 0.45 m along the

highest row to 0.09 m relative to North American Vertical Datum of 1988 (NAVD88) along the lowest

row. Row 4 was approximately along the MHW line at an elevation of 0.32 m NAVD88. Surviving

planted plugs in each row were counted every two to three weeks throughout the 2023 growing

season, and the height of the longest leaf on each plant was recorded to the nearest 0.1 cm.

Sediment samples were taken, and one plant per row was destructively harvested for biomass

measurements two, six, and 16 weeks after planting.

We sampled field soil on three dates:12 June, 13 July, and 5 October 2023. We sampled

along the transect from row 1 to row 9. At each row, we randomly sampled a plant (above and

belowground biomass) and associated soil using a 5 cm diameter PVC corer to 12 cm depth. We

stored samples on ice and then at 4 °C overnight prior to sample processing for downstream

microbiome analyses. To separate bulk from root-associated soils, we cut aboveground plant

biomass, shook the sediment from the roots into a plastic bag, and then transferred roots and

remaining soils into a sterile tube and mixed with sterile 1X phosphate buffer solution (1X PBS).

After shaking for 5 minutes, we used sterile tweezers to remove roots and transferred cleaned roots

into a paper bag to dry at 60 °C for 48 hours. We centrifuged the remaining root-associated soils,

carefully poured off PBS, and stored these rhizosphere sediments at -20 °C until DNA extractions.

After removing the plant from the field sediment sample, we homogenized the sediment sample,

transferred a subsample to a sterile WhirlPak bag, and stored the bulk soil sample at -20 °C until

DNA extractions.

DNA Extraction and Bacterial Amplicon Sequencing

To characterize the sediment microbiomes of the laboratory-treated plugs as well as any

community composition differences between rhizosphere and bulk field soil samples, we extracted

genomic DNA from sediments using the Qiagen DNeasy PowerSoil Kit, employing a combination of

chemical and mechanical methods to lyse the sample following the manufacturer's protocol. The

isolated DNA was quantified using the Nanodrop spectrophotometer and diluted to a standardized

concentration of 10 ng μl^{-1} to ensure uniformity across all samples. This process facilitated

equimolar concentrations of DNA for downstream applications. Triplicate Polymerase Chain

Reactions (PCR) were subsequently conducted on all extracted samples, serving as an efficient

downstream technique for the amplified DNA. To characterize bacterial communities, we used

barcoded primers (515FB/806R) initially developed by the Earth Microbiome Project (Caporaso et

al. 2012) to target the V4 region of the bacterial 16S subunit of the ribosomal RNA gene (Caporaso

et al., 2012; Apprill et al., 2015; Parada et al., 2016). For each sample, three 50 µL PCR libraries

were prepared by combining 35.75 µL molecular grade water, 5 µL Amplitaq Gold 360 10x buffer, 5

µL MgCl$_2$ (25 mM), 1 µL dNTPs (40mM total, 10mM each), 0.25 µL Amplitaq Gold 360 polymerase, 1

µL 515 forward barcoded primer (10 µM), 1 µL 806 reverse primer (10 µM), and 1 µL DNA template

(10 ng µL^{-1}). Thermocycler conditions for PCR reactions were as follows: initial denaturation (94 °C,

3 minutes); 30 cycles of 94 °C for 45 seconds, 50 °C for 30 seconds, 72 °C for 90 seconds; final

elongation (72 °C, 10 minutes). The triplicate 50 µL PCR libraries were combined and then cleaned

using the Axygen® AxyPrep Magnetic (MAG) Bead Purification Kit (Axygen, Union City, California,

USA). Cleaned PCR products were quantified using QuantIT dsDNA BR assay (Thermo Scientific,

Waltham, Massachusetts, USA) and diluted to a concentration of 10 ng µL^{-1} before pooling libraries

in equimolar concentration of 5 ng µL^{-1}. We sequenced the pooled libraries using the Illumina

MiSeq platform using paired-end (2×250 bp) reads (Illumina Reagent Kit v2, 500 reaction kit) at the

Duke University Center for Genomic and Computational Biology Core Sequencing Facility.

Sequences were processed using the mothur (v1.42.0) (Schloss et al. 2009) MiSeq pipeline

(Kozich et al., 2013). We assembled contigs from the paired-end reads, quality trimmed using a

moving average quality score (minimum quality score 35), aligned sequences to the SILVA rRNA

database (v132) (Quast et al. 2013), and removed chimeric sequences using the VSEARCH

algorithm (Rognes et al. 2016). We created operational taxonomic units (OTUs) by first splitting

sequences based on taxonomic class and then binning into OTUs based on 97% sequence

similarity. Taxonomic identity was assigned using the SILVA rRNA database (v132) (Quast et al.

2013). We used an OTU-based 3% distance threshold instead of using amplicon sequence variant

(ASV) of a single base difference, especially given our high diversity microbial communities based

on short-read (250 bp) amplicon sequences, and that similar broadscale microbiome patterns are

observed using OTU- or ASV-based approaches (Glassman and Martiny 2018). In addition, the OTU-

based approach limits overestimating the number of ASVs due to PCR bias and sequencing errors

(Schloss 2021).

Statistical Analysis

We evaluated the interactive effects of inoculation and salinity on *S. alterniflorus*

aboveground biomass, belowground biomass, root-to-shoot ratio, and maximum plant height using

a series of generalized linear models with inoculation and salinity treatment as fixed effects. All

statistical analyses, calculations, and graphical manipulations were run in the R environment (R

version 4.4.1, 2024; R Studio version 2024.09.0+375, 2024) using the vegan, ade4, picante

packages, and custom functions (Dray et al., 2023; Oksanen et al., 2022).

Prior to microbial statistical analyses, we removed low abundance OTUs that were <10

sequences across the data set. We also omitted two samples where the total sequences were less

than 10,000 reads, which represented issues with the library preparation. Samples were organized

by salinity treatment and microbial treatments. Several soil bacterial diversity metrics were then

computed, including Shannon Diversity (H') and Pielou's Evenness (J') with corresponding ANOVA

tests. These metrics provide insights into community composition by assessing both species

richness and species evenness. Species richness refers to the number of different species present

in a community, while species evenness examines the distribution of individuals among those

species (Morris et al., 2014). Shannon Diversity (H') measures the uncertainty in identifying

individual organisms within a species. In highly diverse and evenly distributed systems, there is

more significant uncertainty regarding the identity of unknown individuals, whereas in less diverse

systems, it is easier to predict the identity of unknown individuals, leading to lower uncertainty

(Morris et al., 2014). Pielou's evenness index (J'), also known as the Shannon evenness index,

complements measures of species richness by providing insights into the distribution of individuals

among different species. The index ranges from 0 to 1, with 0 indicating complete unevenness (i.e.,

one species dominates the community) and 1 indicating perfect evenness (i.e., all species are

equally abundant) (Chao & Ricotta, 2019). Pielou's Index standardizes the Shannon H' to account

for differences in species richness, allowing for comparisons of evenness across communities with

varying species numbers. For water and microbial response variables, we performed Tukey's

Honest Significant Difference (HSD) post-hoc analysis of the ANOVA models using the emmeans

package in R. We compared estimated marginal means (EMMs) to conduct pairwise comparisons

between treatment groups (Lenth 2022).

We examined the microbial community variation explained by inoculation and salinity

treatment using a permutational multivariate analysis of variance (PERMANOVA). Principal

Coordinates Analysis (PCoA) graphs were generated to visualize the multivariate relationships

between samples based on their microbial community compositions, allowing for exploration of

how the varying experimental factors influenced the overall microbial diversity and structure.

To analyze prominent indicator species within the microbial communities, we used the

package 'indicspecies'. Indicator species — selected based on their ability to reflect environmental

conditions — indicate changes in the environment and predict diversity patterns (De Cáceres et al.,

2010). The indicspecies package facilitates this analysis by examining the relationship between

species occurrence or abundance and site classification into groups representing habitat types or

disturbance states (De Cáceres et al., 2010).

RESULTS

Evidence for Marsh Microbiomes Buffering Plants from Salinity Stress

Analysis of variance revealed that microbial inocula manipulation (ANOVA, $F_{2,192}$=4.681,

P=0.010) and date (i.e., time) (ANOVA, $F_{3,192}$=4.468, P=0.004) significantly influenced changes in

plant stem height, while water treatments were insignificant (ANOVA, $F_{1,192}$=2.481, P=0.117) (Table

1). When comparing plants treated with salt water, the change in stem heights over time of plants

with autoclaved restoration-area marsh inocula (Autoclaved Inocula) was less than plants with

added microbial inocula (Added Inocula) (Figure 3, Table 2) (EMMs, estimate=−72.76 ± 24.9,

t(192)=−2.922, P=0.044). Similarly, 'Added Inocula' and freshwater treatment yielded more

productive growth than 'Autoclaved Inocula' and saltwater treatment (EMMs, estimate= 77.79 ±

24.9, t(192)=3.124, P=0.0248) (Figure 3, Table 2). Results also revealed that for 'Autoclaved Inocula'

and 'No Addition' treatments, freshwater consistently showed increased plant growth compared to

saltwater. However, in the 'Added Inocula' treatment, this trend was reversed, with the saltwater

treatment resulting in near-equal or greater median plant height over time compared to its

freshwater counterpart. This may hint at salt marsh microbial addition providing a rescue effect

from salinity stress.

We observed a similar (but weaker) trend with aboveground plant biomass measurements,

where 'Added Inocula' treatment may have buffered salinity stress and resulted in greater

aboveground biomass when compared to freshwater treatment (Figure 4a, Table 3). While not

statistically significant (ANOVA, $F_{2,21}$ = 0.656, P = 0.526), we did observe that the aboveground

biomass tended to be greater under 'Added Inocula' in saltwater conditions (Figure 4a). When

looking at belowground biomass (Figure 4b, Table 4), we observed similar root biomass across the

microbial and water treatments (ANOVA, $F_{2,31}$ = 1.496, P = 0.240), which could represent that the

allocation of the plant's carbon was more directed at aboveground biomass rather than

belowground.

286 *Salinity and Microbiome Addition Influenced Bacterial Composition More than Diversity*

287　　　　The experimental treatments of salinity and marsh microbial inocula additions resulted in

288　　an average bacterial Shannon Diversity H' of 6.448 ± 0.083 (Figure 5a) and Pielou's Evenness J' of

289　　0.850 ± 0.0057 (Figure 5b). Soil bacterial diversity metrics were similar across water (ANOVA, $F_{1,48}$=

290　　0.040, $P = 0.842$) and microbial treatments (ANOVA, $F_{2,48}$= 1.789, $P = 0.178$) (Table 6). Likewise,

291　　evenness metrics were similar across treatments of water (ANOVA, $F_{1,48}$= 0.041, $P = 0.840$) and

292　　microbial addition (ANOVA, $F_{2,48}$= 1.861, $P = 0.167$) (Table 7).

293　　　　　Conversely, microbial and water treatments significantly modified bacterial community

294　　composition (Figure 6, Table 5). The main effects of water (PERMANOVA, R^2=0.1002, P<0.001) and

295　　microbial (PERMANOVA, R^2=0.0863, P<0.001) treatments significantly influenced bacterial

296　　community composition (Table 5). We observed distinct bacterial community composition

297　　associated with water treatment, while there was more overlap in bacterial communities across

298　　microbial treatments for salt compared to freshwater treatments (Figure 6).

299

300　　**Context-Dependent Responses of Bacterial Indicator Taxa Across Treatments**

301　　　　　To further examine bacterial associations with microbial and salinity treatments, we used

302　　indicator species analysis to identify a subset of bacterial taxa representing each experimental

303　　root-associated microbiome in salt and freshwater treatments (Table S1). There were 48 OTUs

304　　significantly associated with freshwater conditions across all inocula treatments from phyla

305　　*Proteobacteria*, *Acidobacteria, Actinobacteria*, *Verrucomicrobia*, *Planctomycetes*, and

306　　*Gemmatimonadetes*; and 37 OTUs were associated with all saltwater treatments from phyla

307　　*Proteobacteria*, *Bacteroidetes*, *Acidobacteria*, and *Verrucomicrobia*. In freshwater treatments,

308　　prominent OTUs represented orders *Myxococcales*, *Burkholderiales*, and *Gp1*, the family

309　　*Desulfuromonadaceae*, and the genus *Dongia*. In contrast, in the saltwater treatments, OTUs

represented taxa from the genus *Opitutus*, family *Rhodospirillaceae*, and order *Myxococcales*

(Table S1).

For the microbial treatments, there were 30 OTUs were significantly associated with the 'No

Addition' control treatment from phyla *Proteobacteria, Planctomycetes*, *Bacteroidetes*,

Acidobacteria, *Verrucomicrobia*, *Actinobacteria*, *Gemmatimonadetes*; 8 OTUs were associated

with 'Autoclaved Inocula' treatment from phyla *Proteobacteria*, *Verrucomicrobia*, *Actinobacteria*;

and 22 OTUs were associated with the Added Inocula' treatment from phyla *Proteobacteria*,

Verrucomicrobia, *Acidobacteria, Actinobacteria*, and *WPS-1* (Table S2). The top OTUs for the 'No

Addition' treatments represented taxa from the families *Rhizomicrobium*, *Plantomycetaceae*,

Sinobacteraceae, and *Chitnophagaceae*; for 'Autoclaved Inocula' treatment indicator species

represented families *Myxococcales* and *Rhodospirillales* and orders *Myxococcales* and

Rhodospirillales; and for the 'Added Inocula' treatment, top indicator species represented genera

Geobacter, *Prosthecobacter* and family *Gp1* (Table S2).

Finally, when characterizing differences in field soil samples, indicator species analysis

revealed 22 OTUs significantly associated with bulk plant soil from phyla *Proteobacteria* and

Bacteroidetes; 26 OTUs were associated with rhizosphere soil from phyla *Proteobacteria*,

Chloroflexi, *Firmicutes*, and two unidentified phyla (Table S3). Noteworthy OTUs from bulk soil

were represented by taxa from the order *Chromatiales*, families *Desulfobulbaceae*,

Rhodobacteraceae, and *Desulfobacteraceae,* and the genus *Robertkochia*. Rhizosphere-

associated OTUs were represented by the families *Methylococcaceae* and *Sinobacteraceae*

(*Solimonadaceae*) and the order *Bacillales*.

Plant-Associated Marsh Microbiomes and Elevation Influenced Community Composition

In support of the short-term abiotic and biotic effects on bacterial communities and plant

growth, the characterization of sediment microbiomes at a newly planted marsh restoration site

revealed that plant associations (bulk vs. rhizosphere) and shoreline location (below and above

mean high waterline) influenced bacterial community composition over time (Figure 7).

DISCUSSION

While understanding the underpinnings of microbial ecology is still in its relative infancy

compared to more contemporary scientific disciplines, the roles of belowground microbial

communities in coastal wetland restoration have been of great interest (Birnbaum & Trevathan-

Tackett, 2023). Many empirical studies have demonstrated the reliance of plants on soil

microbiomes in tolerating environmental stresses, including salinity, inundation, drought, and

limited nutrient availability (reviewed in Farrer et al., 2022; Rodriguez et al., 2008). Furthermore, the

adaptive and evolutionary responses of plants to changing environmental conditions have been

shown to rely heavily on the context of microbial community assemblage, with plants exhibiting

increased fitness when associated with stress-adapted microbiomes (Lau & Lennon, 2012).

Presently, gaps in knowledge hinder the success of complete ecosystem restoration projects

(Duarte et al., 2015); therefore, the foundational knowledge gained from biological conservation

and species restoration studies is of utmost relevance and importance. In our investigation, the

interactive effects of microbial inocula additions and salinity treatment on *S. alterniflorus* height,

biomass, and root-associated microbiomes produced various results. We hypothesized that

inoculating *S. alterniflorus* plugs with salt marsh microbial communities would buffer salt stress

and promote plant growth and survivorship; a subset of our results supports this hypothesis.

Upon analyzing plant height data, we concluded that the addition of marsh microbes

'Added Inocula' at early plant development stages may ameliorate salinity stress (as demonstrated

by the reversal trend in median plant heights, Figure 3). Conversely, autoclaved microbial media

and salinity stress treatment were shown to decrease the rate of growth in plant stem height (Table

2). While intuitive, this provides evidence that the sterilization of belowground community

influences and the addition of nonessential or less-utilized abiotic materials can hinder plant

productivity. These findings are in accordance with a similar study, where greenhouse-raised *S.*

alterniflora that were exposed to saltwater (10 ppt salinity) and microbial consortium (a selection of

plant growth-promoting rhizobacteria) experienced the highest rates of growth when compared to

other treatments (Bledsoe & Boopathy, 2016).

Aboveground biomass results further supported evidence that microbial inocula additions

during early plant development may be critical for establishing stress-tolerant plugs. While not

statistically significant (Tables 3 and 4), we observed similar patterns for the belowground biomass.

The allocation of carbon to aboveground biomass more than belowground is of interest as it may

hint at efficient primary productivity and carbon sequestration but less carbon storage and root

establishment. As salt marsh roots indirectly modulate soil dynamics and facilitate resilience to

abiotic stress (Feagin et al., 2009), additional studies are necessary to determine the consequence

of gradual stressors (e.g., sea-level rise, tidal forces) on the success of plant establishment during

marsh restoration.

When analyzing patterns in bacterial diversity, average bacterial diversity H' and evenness J'

were similar across microbial and water treatments. The 'Autoclaved Inocula' treatment tended to

decrease the ranges of H' and J', while greater ranges existed in the 'Added Inocula' treatment.

Additionally, marsh microbial inocula under salinity stress tended to exhibit greater medians in H'

and J' than that of freshwater, in support of the reversal trend. Throughout contemporary literature,

it is well-accepted that increases in microbial biodiversity and evenness are associated with

increased resistance to plant pathogens, improved soil health and ecosystem functions, increased

plant growth and survivorship, and many other positive benefits (see Banerjee & van der Heijden,

2023; Birnbaum & Trevathan-Tackett, 2023; Farrer et al., 2022; Graham & Knelman, 2023; Lau &

Lennon, 2012).

Despite similar bacterial diversity, microbial and water treatments significantly influenced

community composition, with distinct bacterial communities forming in the salt and freshwater

groups (Table 1, Figure 6). This may indicate the strength of the selective pressure of environmental

conditions in forming community structure and should be considered when applying microbial

inocula for restoration. Further, our results suggest that the elevation gradient and associated

abiotic factors found *in situ* may play roles in varying the bacterial community composition along

the shoreline (Figure 7). The separation along the primary coordinate axis indicates that shoreline

elevation and related environmental factors significantly influence bacterial community structure.

Likewise, distinct clustering patterns between bulk soil and rhizosphere samples (Figure 7) suggest

that the root environment exerts selective pressures, fostering unique bacterial assemblages

compared to the surrounding bulk soil (REFs).

As microbial sampling techniques and high-throughput sequencing continue to improve,

increasing evidence has highlighted the utility of microbial communities as indicator species (Liang

et al., 2023; Ma et al., 2022; Urakawa & Bernhard, 2017). Microorganisms function as rapid and

sensitive bioindicators and exhibit patterns of spatial distribution (e.g., distinct rhizosphere and

bulk soil communities) (Urakawa & Bernhard, 2017; Wang et al., 2016); our findings add to this

growing body of literature. While indicator species analysis revealed dominant microbial

assemblages from phyla *Proteobacteria*, *Acidobacteria*, *Actinobacteria*, *Bacteroidetes*, and

Verrucomicrobia – phyla known to be relatively abundant and ubiquitous in soil (Janssen, 2006) –

there were clear distinctions in OTUs with little overlap between treatments. One such overlapping

bioindicator was between saltwater treatments and rhizosphere soil samples, with and OUT from

the nitrogen-fixing family *Desulfuromonadaceae* (Greene, 2014). However, in general, microbial

bioindicators were unique among water and microbial addition treatments, as well as bulk vs.

rhizosphere samples, further providing evidence that plant-microbe interactions and environmental

conditions uniquely structure soil communities. It is important to note that due to bacteria's high

biodiversity, many remain unculturable and unidentified (Stewart, 2012), which represents 23

bacterial taxa that could not be resolved past the phylum or kingdom level.

The patterns in bacterial community composition revealed support for strong

environmental filtering due to salinity exposure, even under short (15-minute) weekly pulses. The

bacterial community composition was similar among microbial treatments but revealed distinct

bacterial assemblages when under the influence of salinity treatments. The pre-treatment of salt

conditions and sediment microbes could help grasses buffer from abiotic and biotic field stressors

after marsh grass plugs are transplanted and exposed to stressful environmental conditions. This

work provides evidence that microbial stewardship is essential for buffering against ecological

stressors and could promote plant establishment for wetland restoration. Additionally, this work

supports the theory of "habitat-adapted symbioses," highlighting the facultative mutualisms

between marsh plants and their symbiotic microorganisms in tolerating high-stress environments

(Rodriguez et al., 2008). With many stressors projected to intensify in the near future (e.g., extreme

climatic events, sea level rise, compounding stress from invasive species, anthropogenic change,

etc.), marshes will need to adapt to survive, and their associated microorganisms will undergo even

more rapid evolutionary changes (Zabin et al., 2022; Billah et al., 2022; Lau & Lennon, 2012; Mueller

et al., 2020). By incorporating microbial inocula into salt marsh restoration practices, success

could be better leveraged.

Within a broader scope, coastal wetlands and salt marshes are among the most productive

ecosystems on Earth and provide numerous ecological and economic services. Salt marshes

429 support fisheries and habitats for biodiversity, filter contaminants, sequester carbon, buffer storm

430 surges and protect shorelines from erosion, provide tourism, and facilitate many other functions

431 (Arkema et al., 2013; Buchanan et al., 2022). Therefore, preserving and restoring such ecological

432 habitats are vital to humans, wildlife, and the environment, emphasizing the concept of One Health

433 (Peixoto et al., 2022). By evaluating the interactions between *Sporobolus alterniflorus*, marsh

434 sediment microbiomes, and environmental stressors, this study provides insight into ecological

435 dynamics and may aid in salt marsh restoration efforts considering climate and anthropogenic

436 change.

ACKNOWLEDGMENTS

We acknowledge the staff at Hammocks Beach State Park and Garner's Landscaping for their assistance in conducting the field experiment. We thank Gittman lab members Jennifer Fickler and James Kelley for assistance in setting up and maintaining the field research and Peralta lab members Toby Gallegos, Daphka Joseph, Scott Siebor, Dejah Smith, Colin Finlay for lab assistance. We also thank Aeran Coughlin and Jean-Philippe Gibert for microbiome sequencing support. This work was supported by the National Science Foundation (#1845845, #2009185, # 2302609 to A.L.P.), the U.S. Coastal Research Program and the National Oceanographic and Atmospheric Administration National Sea Grant (#2023-0534-02/22-NC-USCRP-3 to R.K.G.) and North Carolina Sea Grant (#PAM-P24-002985-SA-02 to R.K.G. and A.L.P.) and East Carolina University (Undergraduate Research and Creative Activity Award to K.A.D.).

REFERENCES

Apprill, A., McNally, S., Parsons, R., & Weber, L. (2015). Minor revision to V4 region SSU rRNA 806R gene primer greatly increases detection of SAR11 bacterioplankton. *Aquatic Microbial Ecology*, *75*. https://doi.org/10.3354/ame01753

Arkema, K. K., Guannel, G., Verutes, G., Wood, S. A., Guerry, A., Ruckelshaus, M., Kareiva, P., Lacayo, M., & Silver, J. M. (2013). Coastal habitats shield people and property from sea-level rise and storms. *Nature Climate Change*, *3*(10), Article 10. https://doi.org/10.1038/nclimate1944

Banerjee, S., & van der Heijden, M. G. A. (2023). Soil microbiomes and one health. *Nature Reviews Microbiology*, *21*(1), 6–20. https://doi.org/10.1038/s41579-022-00779-w

Billah, M. M., Bhuiyan, M. K. A., Islam, M. A., Das, J., & Hoque, A. R. (2022). Salt marsh restoration: An overview of techniques and success indicators. *Environmental Science and Pollution Research*, *29*(11), 15347–15363. https://doi.org/10.1007/s11356-021-18305-5

Birnbaum, C., & Trevathan-Tackett, S. M. (2023). Aiding coastal wetland restoration via the belowground soil microbiome: An overview. *Restoration Ecology*, *31*(7), e13824. https://doi.org/10.1111/rec.13824

Bledsoe, R., & Boopathy, R. (2016). Bioaugmentation of microbes to restore coastal wetland plants to protect land from coastal erosion. *International Biodeterioration & Biodegradation*, *113*, 155–160. https://doi.org/10.1016/j.ibiod.2016.02.010

Bruno, J. F. (2000). Facilitation of Cobble Beach Plant Communities Through Habitat Modification by Spartina Alterniflora. *Ecology*, *81*(5), 1179–1192. https://doi.org/10.1890/0012-9658(2000)081[1179:FOCBPC]2.0.CO;2

Buchanan, M. K., Kulp, S., & Strauss, B. (2022). Resilience of U.S. coastal wetlands to accelerating

sea level rise. *Environmental Research Communications*, *4*(6), 061001.

https://doi.org/10.1088/2515-7620/ac6eef

Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens, S. M.,

Betley, J., Fraser, L., Bauer, M., Gormley, N., Gilbert, J. A., Smith, G., & Knight, R. (2012).

Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq

platforms. *The ISME Journal*, *6*(8), 1621–1624. https://doi.org/10.1038/ismej.2012.8

Cavicchioli, R., Ripple, W. J., Timmis, K. N., Azam, F., Bakken, L. R., Baylis, M., Behrenfeld, M. J.,

Boetius, A., Boyd, P. W., Classen, A. T., Crowther, T. W., Danovaro, R., Foreman, C. M.,

Huisman, J., Hutchins, D. A., Jansson, J. K., Karl, D. M., Koskella, B., Mark Welch, D. B., …

Webster, N. S. (2019). Scientists' warning to humanity: Microorganisms and climate change.

Nature Reviews Microbiology, *17*(9), Article 9. https://doi.org/10.1038/s41579-019-0222-5

Chao, A., & Ricotta, C. (2019). Quantifying evenness and linking it to diversity, beta diversity, and

similarity. *Ecology*, *100*(12), e02852. https://doi.org/10.1002/ecy.2852

De Cáceres, M., Legendre, P., & Moretti, M. (2010). Improving indicator species analysis by

combining groups of sites. *Oikos*, *119*(10), 1674–1684. https://doi.org/10.1111/j.1600-

0706.2010.18334.x

Dray, S., Dufour, A.-B., Thioulouse, and J., Jombart, with contributions from T., Pavoine, S., Lobry, J.

R., Ollier, S., Borcard, D., Legendre, P., & Chessel, S. B. and A. S. B. on earlier work by D.

(2023). *ade4: Analysis of Ecological Data: Exploratory and Euclidean Methods in*

Environmental Sciences (1.7-22) [Computer software]. https://cran.r-

project.org/web/packages/ade4/index.html

Duarte, C. M., Borja, A., Carstensen, J., Elliott, M., Krause-Jensen, D., & Marbà, N. (2015).

Paradigms in the Recovery of Estuarine and Coastal Ecosystems. *Estuaries and Coasts*,

38(4), 1202–1212. https://doi.org/10.1007/s12237-013-9750-9

Feagin, R. A., Lozada-Bernard, S. M., Ravens, T. M., Möller, I., Yeager, K. M., & Baird, A. H. (2009).

Does vegetation prevent wave erosion of salt marsh edges? *Proceedings of the National*

Academy of Sciences, *106*(25), 10109–10113. https://doi.org/10.1073/pnas.0901297106

Farrer, E. C., Van Bael, S. A., Clay, K., & Smith, M. K. H. (2022). Plant-Microbial Symbioses in Coastal

Systems: Their Ecological Importance and Role in Coastal Restoration. *Estuaries and*

Coasts, *45*(7), 1805–1822. https://doi.org/10.1007/s12237-022-01052-2

Gedan, K. B., & Bertness, M. D. (2010). How will warming affect the salt marsh foundation species

Spartina patens and its ecological role? *Oecologia*, *164*(2), 479–487.

https://doi.org/10.1007/s00442-010-1661-x

Gittman, R. K., Baillie, C. J., Arkema, K. K., Bennett, R. O., Benoit, J., Blitch, S., Brun, J., Chatwin, A.,

Colden, A., Dausman, A., DeAngelis, B., Herold, N., Henkel, J., Houge, R., Howard, R.,

Hughes, A. R., Scyphers, S. B., Shostik, T., Sutton-Grier, A., & Grabowski, J. H. (2019).

Voluntary Restoration: Mitigation's Silent Partner in the Quest to Reverse Coastal Wetland

Loss in the USA. *Frontiers in Marine Science*, *6*. https://doi.org/10.3389/fmars.2019.00511

Gittman, R. K., Baillie, C. J., Cros, A., Grabowski, J. H., McKinney, M.-M., Saccomanno, V. R., Smith,

Carter. S., & DeAngelis, B. (2024). Assessing how restoration can facilitate 30×30 goals for

climate-resilient coastal ecosystems in the United States. *Conservation Biology*, *n/a*(n/a),

e14429. https://doi.org/10.1111/cobi.14429

Glassman, S. I., & Martiny, J. B. H. (2018). Broadscale Ecological Patterns Are Robust to Use of

Exact Sequence Variants versus Operational Taxonomic Units. *mSphere*, *3*(4),

10.1128/msphere.00148-18. https://doi.org/10.1128/msphere.00148-18

Graham, E. B., & Knelman, J. E. (2023). Implications of Soil Microbial Community Assembly for

Ecosystem Restoration: Patterns, Process, and Potential. *Microbial Ecology*, *85*(3), 809–819.

https://doi.org/10.1007/s00248-022-02155-w

Janssen, P. H. (2006). Identifying the Dominant Soil Bacterial Taxa in Libraries of 16S rRNA and 16S

rRNA Genes. *Applied and Environmental Microbiology*, *72*(3), 1719–1728.

https://doi.org/10.1128/AEM.72.3.1719-1728.2006

Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., & Schloss, P. D. (2013). Development of

a Dual-Index Sequencing Strategy and Curation Pipeline for Analyzing Amplicon Sequence

Data on the MiSeq Illumina Sequencing Platform. *Applied and Environmental Microbiology*,

79(17), 5112–5120. https://doi.org/10.1128/AEM.01043-13

Lau, J. A., & Lennon, J. T. (2011). Evolutionary ecology of plant–microbe interactions: Soil microbial

structure alters selection on plant traits. *New Phytologist*, *192*(1), 215–224.

https://doi.org/10.1111/j.1469-8137.2011.03790.x

Lau, J. A., & Lennon, J. T. (2012). Rapid responses of soil microorganisms improve plant fitness in

novel environments. *Proceedings of the National Academy of Sciences*, *109*(35), 14058–

14062. https://doi.org/10.1073/pnas.1202319109

Lenth, R. (2022). emmeans: Estimated marginal means, aka least-squares means. R package

version 1.7. 2, https://github.com/rvlenth/emmeans

Leonardi, N., Carnacina, I., Donatelli, C., Ganju, N. K., Plater, A. J., Schuerch, M., & Temmerman, S.

(2018). Dynamic interactions between coastal storms and salt marshes: A review.

Geomorphology, *301*, 92–107. https://doi.org/10.1016/j.geomorph.2017.11.001

Liang, S., Li, H., Wu, H., Yan, B., & Song, A. (2023). Microorganisms in coastal wetland sediments: A

review on microbial community structure, functional gene, and environmental potential.

Frontiers in Microbiology, *14*. https://doi.org/10.3389/fmicb.2023.1163896

Ma, F., Wang, C., Zhang, Y., Chen, J., Xie, R., & Sun, Z. (2022). Development of Microbial Indicators

in Ecological Systems. *International Journal of Environmental Research and Public Health*,

19(21), Article 21. https://doi.org/10.3390/ijerph192113888

Mariotte, P., Mehrabi, Z., Bezemer, T. M., De Deyn, G. B., Kulmatiski, A., Drigo, B., Veen, G. F. (Ciska),

van der Heijden, M. G. A., & Kardol, P. (2018). Plant–Soil Feedback: Bridging Natural and

Agricultural Sciences. *Trends in Ecology & Evolution*, *33*(2), 129–142.

https://doi.org/10.1016/j.tree.2017.11.005

McHugh, J. M., & Dighton, J. (2004). Influence of Mycorrhizal Inoculation, Inundation Period,

Salinity, and Phosphorus Availability on the Growth of Two Salt Marsh Grasses, Spartina

alterniflora Lois. And Spartina cynosuroides (L.) Roth., in Nursery Systems. *Restoration*

Ecology, *12*(4), 533–545. https://doi.org/10.1111/j.1061-2971.2004.03109.x

Morris, E. K., Caruso, T., Buscot, F., Fischer, M., Hancock, C., Maier, T. S., Meiners, T., Müller, C.,

Obermaier, E., Prati, D., Socher, S. A., Sonnemann, I., Wäschke, N., Wubet, T., Wurst, S., &

Rillig, M. C. (2014). Choosing and using diversity indices: Insights for ecological applications

from the German Biodiversity Exploratories. *Ecology and Evolution*, *4*(18), 3514–3524.

https://doi.org/10.1002/ece3.1155

Mueller, E. A., Wisnoski, N. I., Peralta, A. L., & Lennon, J. T. (2020). Microbial rescue effects: How

microbiomes can save hosts from extinction. *Functional Ecology*, *34*(10), 2055–2064.

https://doi.org/10.1111/1365-2435.13493

Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B.,

Solymos, P., Stevens, M. H. H., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker, B.,

Borcard, D., Carvalho, G., Chirico, M., Caceres, M. D., Durand, S., … Weedon, J. (2022).

vegan: Community Ecology Package (2.6-4) [Computer software]. https://cran.r-

project.org/web/packages/vegan/index.html

Parada, A. E., Needham, D. M., & Fuhrman, J. A. (2016). Every base matters: Assessing small

subunit rRNA primers for marine microbiomes with mock communities, time series and

global field samples. *Environmental Microbiology*, *18*(5), 1403–1414.

https://doi.org/10.1111/1462-2920.13023

Peixoto, R. S., Voolstra, C. R., Sweet, M., Duarte, C. M., Carvalho, S., Villela, H., Lunshof, J. E., Gram,

L., Woodhams, D. C., Walter, J., Roik, A., Hentschel, U., Thurber, R. V., Daisley, B., Ushijima,

B., Daffonchio, D., Costa, R., Keller-Costa, T., Bowman, J. S., … Berg, G. (2022). Harnessing

the microbiome to prevent global biodiversity loss. *Nature Microbiology*.

https://doi.org/10.1038/s41564-022-01173-1

Rodriguez, R. J., Henson, J., Van Volkenburgh, E., Hoy, M., Wright, L., Beckwith, F., Kim, Y.-O., &

Redman, R. S. (2008). Stress tolerance in plants via habitat-adapted symbiosis. *The ISME*

Journal, *2*(4), 404–416. https://doi.org/10.1038/ismej.2007.106

Rognes, T., Flouri, T., Nichols, B., Quince, C., & Mahé, F. (2016). VSEARCH: A versatile open source

tool for metagenomics. *PeerJ*, *4*, e2584. https://doi.org/10.7717/peerj.2584

Rudershausen, P. J., Lombardo, S. M., & Buckel, J. A. (2021). Linking Historical Changes in Salt-

Marsh Coverage to Lost Production of a Nektonic Bioindicator. *Marine and Coastal*

Fisheries, *13*(2), 131–139. https://doi.org/10.1002/mcf2.10147

Schloss, P. D. (2021). Amplicon Sequence Variants Artificially Split Bacterial Genomes into

Separate Clusters. *mSphere*, *6*(4), e00191-21. https://doi.org/10.1128/mSphere.00191-21

Stewart, E. J. (2012). Growing Unculturable Bacteria. *Journal of Bacteriology*, 194(16), 4151–4160.

https://doi.org/10.1128/jb.00345-12

Urakawa, H., & Bernhard, A. E. (2017). Wetland management using microbial indicators. *Ecological*

Engineering, *108*, 456–476. https://doi.org/10.1016/j.ecoleng.2017.07.022

Valiela, I., Lloret, J., Bowyer, T., Miner, S., Remsen, D., Elmstrom, E., Cogswell, C., & Robert Thieler,

E. (2018). Transient coastal landscapes: Rising sea level threatens salt marshes. *Science of*

The Total Environment, *640–641*, 1148–1156.

https://doi.org/10.1016/j.scitotenv.2018.05.235

Wang, M., Yang, P., & Falcão Salles, J. (2016). Distribution of Root-Associated Bacterial

Communities Along a Salt-Marsh Primary Succession. *Frontiers in Plant Science,* 6.

https://doi.org/10.3389/fpls.2015.01188

Zabin, C. J., Jurgens, L. J., Bible, J. M., Patten, M. V., Chang, A. L., Grosholz, E. D., & Boyer, K. E.

(2022). Increasing the resilience of ecological restoration to extreme climatic events.

Frontiers in Ecology and the Environment, *20*(5), 310–318. https://doi.org/10.1002/fee.2471

STATEMENTS AND DECLARATIONS

The authors declare no competing interests.

 TABLES

638 **Table 1.** Summary ANOVA comparing changes in stem height (in mm) due to main effects (water

639 treatment, microbial inocula treatment, date) and the interaction water × microbial treatments ×

640 date. Bold text indicates significant differences ($P \leq 0.05$).

Effect	DF	Sum of Sqs	Mean Sq	F-value	P-value
Microbe TRT	2	100859	50429	4.6807	**0.0104**
Water TRT	1	26730	26730	2.4810	0.1169
Date	3	144419	48140	4.4682	**0.0047**
Water TRT: Microbe TRT	2	18190	9095	0.8442	0.4315
Microbe TRT: Date	6	15666	2611	0.2423	0.9619
Water TRT: Date	3	3457	1152	0.1069	0.9560
Microbe TRT: Water TRT: Date	6	31374	5229	0.4853	0.8188
Residuals	192	2068598	10774		

643   **Table 2.** Summary of estimated marginal means (EMMs) contrasting isolated experimental

644   manipulation effects on plant stem height. Results are averaged over time. P value adjustment:

645   tukey method for comparing a family of 6 estimates. Bold text indicates significant differences (P ≤

646   0.05).

Contrast	*estimate*	SE	DF	*T-ratio*	*P-value*
CONTROL FRESH - Microbe_neg FRESH	-3.31	24.0	192	-0.138	1.0000
CONTROL FRESH - Microbe_pos FRESH	-33.72	24.5	192	-1.378	0.7400
CONTROL FRESH - CONTROL SALT	12.52	24.7	192	0.508	0.9958
CONTROL FRESH - Microbe_neg SALT	44.06	24.9	192	1.770	0.4878
CONTROL FRESH - Microbe_pos SALT	-28.69	24.5	192	-1.173	0.8493
Microbe_neg FRESH - Microbe_pos FRESH	-30.41	24.0	192	-1.267	0.8026
Microbe_neg FRESH - CONTROL SALT	15.83	24.2	192	0.654	0.9865
Microbe_neg FRESH - Microbe_neg SALT	47.37	24.4	192	1.938	0.3823
Microbe_neg FRESH - Microbe_pos SALT	-25.38	24.0	192	-1.058	0.8974
Microbe_pos FRESH - CONTROL SALT	46.24	24.7	192	1.876	0.4204
Microbe_pos FRESH - Microbe_neg SALT	77.79	24.9	192	3.124	**0.0248**
Microbe_pos FRESH - Microbe_pos SALT	5.03	24.5	192	0.206	0.9999
CONTROL SALT - Microbe_neg SALT	31.54	25.1	192	1.257	0.8076
CONTROL SALT - Microbe_pos SALT	-41.22	24.7	192	-1.672	0.5522
Microbe_neg SALT - Microbe_pos SALT	-72.76	24.9	192	-2.922	**0.0444**

651 **Table 3.** Summary ANOVA comparing changes in aboveground biomass due to main effects (water

652 treatment, microbial inocula treatment) and the interaction water × microbial treatments. Bold text

653 indicates significant differences (P ≤ 0.05).

Effect	*DF*	Sum of Sqs	Mean Sq	*F-value*	*P-value*
Microbe TRT	2	15.199	7.5994	2.6148	0.08926
Water TRT	1	2.458	2.4581	0.8458	0.36485
Water TRT: Microbe TRT	2	3.811	1.9054	0.6556	0.52618
Residual	31	90.097	2.9064		

662 **Table 4.** Summary ANOVA comparing changes in belowground biomass due to main effects (water

663 treatment, microbial inocula treatment) and the interaction water × microbial treatments. Bold text

664 indicates significant differences (P ≤ 0.05).

Effect	*DF*	Sum of Sqs	Mean Sq	*F-value*	*P-value*
Microbe TRT	2	2.803	1.40151	1.3003	0.2869
Water TRT	1	0.036	0.03587	0.0333	0.8564
Water TRT: Microbe TRT	2	3.225	1.61240	1.4960	0.2398
Residual	31	33.413	1.07782		

674　　**Table 5.** Summary PERMANOVA comparing microbial community composition due to main effects

675　　(water treatment, microbial inocula treatment) and the interaction water × microbial treatments.

676　　Bold text indicates significant differences (P ≤ 0.05).

Effect	*DF*	Sum of Sqs	R^2	*F-value*	*P-value*
Water TRT	1	0.4953	0.1002	6.2325	**<0.001**
Microbe TRT	2	0.4267	0.0863	2.6845	**<0.001**
Water TRT: Microbe TRT	2	0.2062	0.0417	1.2973	0.1399
Residual	48	3.8147	0.7718		
Total	53	4.9429	1.0000		

687　　**Table 6.** Summary ANOVA comparing changes in Shannon Diversity H' due to main effects (water

688　　treatment, microbial inocula treatment) and the interaction water × microbial treatments. Bold text

689　　indicates significant differences (P ≤ 0.05).

Effect	DF	Sum of Sqs	Mean Sq	*F-value*	*P-value*
Water TRT	1	0.000	0.00033	0.005	0.947
Microbe TRT	2	0.254	0.12716	1.769	0.181
Water TRT: Microbe TRT	2	0.184	0.09182	1.278	0.288
Residual	48	3.450	0.07187		

691
692
693
694

698 **Table 7.** Summary ANOVA comparing changes in Pielou's Evenness J' due to main effects (water

699 treatment, microbial inocula treatment) and the interaction water × microbial treatments. Bold text

700 indicates significant differences ($P \leq 0.05$).

Effect	*DF*	Sum of Sqs	Mean Sq	*F-value*	*P-value*
Water TRT	1	0.000002	0.0000016	0.005	0.946
Microbe TRT	2	0.001336	0.0006682	1.936	0.155
Water TRT: Microbe TRT	2	0.000738	0.0003688	1.068	0.352
Residual	48	0.016569	0.0003452		

FIGURES

Figure 1.

Figure 1. Location of field marsh sediment and plant sampling at Hammocks Beach State Park,

Swansboro, North Carolina, USA. *Sporobolus alterniflorus* transplants were planted along an

elevation gradient. Row 1 represented the highest elevation and row 9 represented the lowest

elevation. At three time points (12 June 2023, 13 July 2023, 5 October 2023), one plant sample and

associated bulk soil were collected and processed to examine bulk and rhizosphere microbiomes.

Marsh sediment from this field site was used as the microbial inocula treatment in the salinity x

microbe experiment.

Figure 2.

Figure 2. Experimental design of salinity/microbe treatments. Factorial experiment with three levels

of microbial addition (No Addition, Autoclaved Inocula, Added Inocula) and two levels of salinity

(<0.5 psu, 20 psu). Treatments were replicated ten times. Microbial inocula were combined with

potting mix and established for four weeks, followed by weekly salt stress treatment, and harvested

after eight weeks.

Figure 3.

Figure 3. Boxplots depicting plant height (in mm) on the y-axis over time on the x-axis in response to

marsh soil microbes and salinity stress treatments. Color represents salinity stress (orange) and

freshwater control (blue). The boxplot is a visual representation of 5 key summary statistics: the

median, the 25% and 75% percentiles, and the whiskers, which represent the feasible range of the

data as determined by 1.5 × the interquartile range. Symbols represent individual data points.

735   **Figure 4.**

736   **(a)**

744   **(b)**

Figure 4. Boxplots depicting aboveground (a) and belowground root (b) biomass (in g) on the y-axis

in response to microbial (No Addition, Autoclaved Inocula, Added Inocula) and salinity (<0.5 psu,

20 psu) treatments. Color represents salinity stress (orange) and freshwater control (blue). The

boxplot is a visual representation of 5 key summary statistics: the median, the 25% and 75%

percentiles, and the whiskers, which represent the feasible range of the data as determined by 1.5 ×

the interquartile range. Symbols represent individual data points.

Figure 5.

(a)

768 **(b)**

Figure 5. Boxplots for bacterial diversity representing Shannon Diversity Index (H') (a) and Pielou's

Evenness (J') (b) associated with microbial (No Addition, Autoclaved Inocula, Added Inocula) and

salinity (<0.5 psu, 20 psu) treatments. Colors indicate salinity treatment (fresh in blue, salt in

orange). Symbols represent individual data points.

Figure 6.

(a)

792 **(b)**

Figure 6. Ordination based on a principal coordinates analysis depicting bacterial community

composition according to salinity and microbial treatments representing average centroid across

treatments (a) and individual samples (b). Symbols are colored according to salinity treatment

(orange = saltwater, blue = freshwater) and shaped by microbial treatment (circles = Autoclaved

Inocula, triangles = Added Inocula, squares = No Addition).

Figure 7.

Figure 7. Ordination based on a principal coordinates analysis depicting bacterial community

composition according to salinity gradient from landward (row 1) to seaward (row 9) and sample

source (bulk soil = circles, root-associated rhizosphere = triangles). Symbols are colored according

to row planted. Row 4 was located at approximately the mean high-water line.