Evaluating Plant-Microbe Associations in Response to Environmental Stressors to Enhance

Salt Marsh Restoration

RUNNING TITLE: Salt marsh plant-microbiome restoration

Kai A. Davis[1], Mary-Margaret McKinney[2], Rachel K. Gittman[1,3], Ariane L. Peralta[1,*]

[1]Department of Biology, [2]Integrated Coastal Sciences, East Carolina University, Greenville, North

Carolina, USA, 3 Coastal Studies Institute, Wanchese, North Carolina, USA

*Corresponding Author: phone: 252-328-2712, email: peraltaa@ecu.edu

15   Author contributions: All authors contributed to the study conception, design, material preparation,

16   and data collection. Kai A. Davis and Ariane L. Peralta performed data analyses. The first draft of the

17   manuscript was written by Kai A. Davis, and all authors commented on previous versions of the

18   manuscript. All authors read and approved the final manuscript. Kai A. Davis, Rachel K. Gittman,

19   and Ariane L. Peralta acquired financial support for the project that resulted in this manuscript.

ABSTRACT

Despite substantial contributions of ecosystem services, mutualistic plant-microbe associations

in wetlands are severely threatened by human activities. Therefore, promoting positive plant-

microbe associations underpins coastal wetland restoration, where human stressors and climate

change challenge successful outcomes. This study examined how salinity stressors influence

plant-microbe relationships, where we hypothesized that the presence of microbes would provide

a rescue effect by buffering abiotic stressors and yielding higher plant biomass. We used a whole

sediment inocula approach and exposed marsh cordgrass (*Sporobolus alterniflorus*) plugs to a

factorial experiment with three levels of microbiome addition (microbial inocula, autoclaved

microbial inocula, no microbe control) and two levels of salinity (<0.5 psu, 20 psu), replicated ten

times. We added marsh-site microbial inocula with autoclaved peat-based greenhouse soil and

exposed half the plugs to saltwater (20 psu) and half to freshwater (<0.5 psu). Results revealed that

marsh microbial inocula additions during early plant development may ameliorate salinity

stressors. Plants treated with microbial inocula and salinity stress exhibited greater aboveground

biomass ($P < 0.09$) than those under freshwater conditions. With live microbial inocula, the median

of aboveground biomass and change in plant height were higher in saltwater compared to

freshwater conditions. Change in plant height, belowground biomass, root-to-shoot ratio, soil

bacterial diversity H', and evenness J' were similar across microbial and salinity treatments.

However, salinity ($P < 0.001$) and microbial treatments ($P < 0.001$) significantly influenced the

bacterial community composition, with distinct assemblages under salinity conditions. This work

supports microbiome stewardship enhancing plant resiliency, improving future wetland restoration

outcomes.

Keywords: soil microbial communities, salinity stress, microbiome stewardship, coastal protection

INTRODUCTION

48
Salt marshes, coastal wetlands, and other coastal ecosystems aid in ameliorating erosion

and flooding, protect people and property from storm surges, and provide habitat for fisheries and

many other species (Arkema et al., 2013; Buchanan et al., 2022). Despite their high ecological and

economic importance, these coastal habitats are under threat due to anthropogenic activities (e.g.,

land development, dredging) and climate change (e.g., accelerated sea level rise, increased mean

global surface temperature, and frequency of heatwaves) (Valiela et al., 2018; Leonardi et al., 2018;

Rudershausen et al., 2021; Buchanan et al., 2022; Trivedi et al., 2022). Unfortunately, current

restoration efforts are failing to keep pace with current and future losses (Gittman et al. 2019),

highlighting the necessity for improved salt marsh restoration techniques (Gittman et al. 2024).

The planting of marsh smooth cordgrass *Sporobolus alterniflorus* (also known as *Spartina*

alterniflora) is a common approach to salt marsh restoration efforts, as the plant is a foundation

species that can withstand a wide range of salinities, inundation periods, and anoxic stress (Gedan

& Bertness, 2010). These foundation or habitat-forming species alter their environment, making it

possible for other organisms to establish and endure within an ecosystem (Bruno, 2000). However,

when commercially raised nursery plants are transplanted to field restoration sites, they are

exposed to novel abiotic stressors (e.g., salinity and wave energy) and biotic factors (e.g., species

interactions) that can induce stress and hinder restoration (Billah et al., 2022). Published studies

evaluating the effects of exposing *S. alterniflorus* plants to abiotic stressors prior to establishment

at a restoration site are lacking (but *see* Sloey et al. 2025). In addition, the understanding of plant-

microbe associations between marsh soil microbiomes and *S. alterniflorus* and its influence on

plant growth and survival is limited. Therefore, an examination of the effect of salinity stress and

microbial inoculation on marsh plant productivity would further the understanding of ecological

mechanisms and aid in restoring coastal wetland ecosystems.

In a broader context, microorganisms can enhance the nutrient acquisition of plants,

support growth, and suppress diseases from pathogens (Mariotte et al., 2018; Farrer et al., 2022;

Pantigoso et al., 2022; Chepsergon & Moleleki, 2023; Singh et al., 2023). In turn, plants can provide

carbon resources and oxygen to root-associated microorganisms (Mariotte et al., 2018;

Chepsergon & Moleleki, 2023). Human activities and climate change can and have altered these

plant-microbe relationships in ways that promoted or suppressed plant growth (Cavicchioli et al.,

2019; Rojas-Sánchez et al., 2025; Singh et al., 2023; Dini-Andreote, 2025). For example,

anthropogenically caused eutrophication and pollution have disrupted carbon, nitrogen, and other

nutrient cycles, leading to losses in microbial biodiversity and reduced fungal colonization of plants

and microbial respiration (Wei et al., 2013; Cavicchioli et al., 2019; Lin et al., 2020). Global change,

such as rising global surface temperatures, heatwaves, droughts, and sea level rise, can directly

and indirectly modulate plant physiology, soil composition and chemistry, and microbial

community assemblages, often leading to states of stress and dysbiosis (Trivedi et al., 2022; Dini-

Andreote, 2025; Zabin et al., 2022). Conversely, the practice of microbiome stewardship, defined as

the responsible and conscious management of microbial communities for beneficial conservation

and restoration outcomes, can minimize or reverse biodiversity loss, as well as preserve key

ecosystem functions and mechanisms (Peixoto et al., 2022; Choudoir et al., 2025). Growing

evidence has also indicated that microbial communities can enhance plant survivorship and

growth in high-stress environments, potentially aiding ecological restoration efforts (McHugh &

Dighton, 2004; Mariotte et al., 2018; Li et al., 2021; Rojas-Sánchez et al., 2025; Dini-Andreote,

2025).

With the urgent need for enhanced restoration techniques, we aim to understand how to

promote positive plant-microbe associations to aid in restoring coastal wetland ecosystems where

human stressors and climate change challenge restoration outcomes. In this investigation, we

evaluated the interactive effects of salinity stress and marsh sediment microbial inoculation on *S.*

alterniflorus productivity. Conducting a replicated factorial experiment with two levels of salinity

and three levels of microbial inocula treatment, we measured the change in plant height and

biomass (aboveground and belowground) of *S. alterniflorus* plugs after a 2-month duration.

Additionally, we extracted genomic DNA from the sediment samples to characterize the soil

microbiome of experimentally manipulated plants. As resident marsh microorganisms are well-

adapted to a hypersaline environment, we hypothesized that *S. alterniflorus* plugs inoculated with

established salt marsh sediment microbial communities would exhibit higher biomass than control

plugs when subjected to salt stress.

METHODS

Microbiome x Salinity Stress Experimental Design

To test our hypothesis that *S. alterniflorus* plugs inoculated with restoration-area microbial

communities would have higher growth rates than control plugs under salinity stress, we

conducted a factorial experiment with three levels of microbiome addition (*No Addition*,

Autoclaved Inocula, *Added Inocula*) and two levels of salinity (<0.5 psu, 20 psu), replicated ten

times (*n*=60) (Figure 1). We collected marsh sediment from Hammocks Beach State Park

(Swansboro, North Carolina), while the marsh plugs, *Sporobolus alterniflorus*, were purchased

from a commercial plant nursery (Garner's Landscaping, Inc., Newport, North Carolina) that

routinely grows *S. alterniflorus* plugs from locally collected seeds for salt marsh restoration

projects. The commercial marsh plugs were procured in 50-cell propagation trays, with one juvenile

plant per 71 cm^3 cell and filled with standard greenhouse mix (Sungro Fafard peat-based potting

mix).

To prepare the microbial treatments, we combined microbial inocula with Fafard 3B potting

mix, with 5% of the total volume being marsh sediment and 95% being autoclaved potting mix

(following the methods of Lau & Lennon, 2011). Specifically, we combined 10 g of marsh sediment

with 190 g of potting mix in an 11.25 cm square pot prior to transplanting the plant plug (25 g

average weight). While autoclaving can influence the chemistry of soil organic matter and the

accessibility of plant nutrients, it creates a benchmark for sterilization and microbial community

establishment. Any variations observed between microbial treatments can be attributed to the

introduction of inocula containing either live or dead soil microorganisms (Lau & Lennon, 2011).

The combination of untreated marsh sediment and autoclaved potting mix represented the

Added Inocula treatment. For *Autoclaved Inocula*, marsh sediment was autoclaved before being

combined, with the purpose of examining whether there were abiotic effects on plant productivity.

While the microbial load may not have been completely erased, autoclaving ensured a decreased

abundance of resident marsh microorganisms (31 sequence reads for autoclaved field soil

compared to >26,000 sequence reads for all other sample types). For *No Addition*, 100% of the

volume of the microbial treatment was comprised of autoclaved potting mix. Commercially grown

S. alterniflorus plugs were obtained on 01 June 2023 and were transferred to 11.25 cm square pots

with their respective treatments on 08 June to begin a four-week acclimation period to their new soil

conditions. The plants were kept under laboratory conditions (room temperature at about 25 °C)

next to a window for sunlight, and were rotated daily to ensure even light exposure.

On 23 June, the plants were transferred from the laboratory to an outside environment in

Greenville, North Carolina, ~70 miles northwest of where the plants were purchased. We placed

plants on outdoor tables, in trays, where they received the same sunlight and precipitation

conditions across all experimental units. We also began conducting biweekly measurements at this

time, recording the maximum plant stem height (in mm) following standard methods (Currin et al.,

2008; Gittman et al., 2018; Mendelssohn & Seneca, 1980; Morris & Haskin, 1990) and the number

of live stems, with changes in stem height (final – initial) functioning as a proxy for growth. The four-

week acclimation period concluded on 09 July, at which point the weekly salinity stress treatment

was initiated and continued for four weeks.

For the salinity treatment, we bottom-watered each potted plug for 15 minutes with either

~20 psu saltwater or ~0.5 psu freshwater. Saltwater was made with Instant Ocean Sea Salt, with a

general ratio of ~300 g Instant Ocean to 14 L of tap water. This established a concentration of ~20

psu, which represented salinity levels at Hammocks Beach State Park, where salinity ranges from

20 to 35 psu (personal communication, park manager). Although the duration of salt exposure in

our experimental setup may be shorter compared to natural field conditions experienced by plants,

it was anticipated that the deposited salt would gradually accumulate in the soil due to

evaporation, mimicking real-world scenarios. This deliberate manipulation allowed us to

specifically examine the effects of salt stress on plant survival and growth, distinct from the stress

induced by prolonged inundation, which typically leads to anoxia. Thus, our experimental design

ensured a focused investigation into salt stress dynamics without confounding stressors

associated with prolonged inundation (Wang et al. 2019). Further, we opted for a short-duration,

saltwater bottom-watering because it could be easily replicated by nurseries. Top-watering and

subsequent freshwater flushing could be prohibitively labor- and equipment-intensive (saltwater

would have to be flushed out of watering systems and equipment) as well as difficult to apply to

multiple trays of plants simultaneously.

Plant and Soil Sample Processing

At the end of the experiment, after eight weeks total, we harvested aboveground biomass

and roots, dried the plant biomass samples at 60 °C for 48 hours, and weighed aboveground and

root biomass. For soil bacterial characterization and analysis, we homogenized the remaining soil,

transferred soil into sterile WhirlPak bags, and stored samples at -20 °C until DNA extraction and

16S rRNA amplicon sequencing (described below).

DNA Extraction and Bacterial Amplicon Sequencing

To characterize the sediment microbiomes of the laboratory-treated plugs as well as any

community composition differences, we extracted genomic DNA from sediments using the Qiagen

DNeasy PowerSoil Kit, employing a combination of chemical and mechanical methods to lyse the

sample following the manufacturer's protocol. The isolated DNA was quantified using the

Nanodrop spectrophotometer and diluted to a standardized concentration of 10 ng μl^{-1} to ensure

uniformity across all samples. This process facilitated equimolar concentrations of DNA for

downstream applications. Triplicate Polymerase Chain Reactions (PCR) were subsequently

conducted on all extracted samples, serving as an efficient downstream technique for the

amplified DNA. To characterize bacterial communities, we used barcoded primers (515FB/806R)

initially developed by the Earth Microbiome Project (Caporaso et al. 2012) to target the V4 region of

the bacterial 16S subunit of the ribosomal RNA gene (Caporaso et al., 2012; Apprill et al., 2015;

Parada et al., 2016). For each sample, three 50 µL PCR libraries were prepared by combining 35.75

µL molecular grade water, 5 µL Amplitaq Gold 360 10x buffer, 5 µL $MgCl_2$ (25 mM), 1 µL dNTPs

(40mM total, 10mM each), 0.25 µL Amplitaq Gold 360 polymerase, 1 µL 515 forward barcoded

primer (10 µM), 1 µL 806 reverse primer (10 µM), and 1 µL DNA template (10 ng μL^{-1}). Thermocycler

conditions for PCR reactions were as follows: initial denaturation (94 °C, 3 minutes); 30 cycles of 94

°C for 45 seconds, 50 °C for 30 seconds, 72 °C for 90 seconds; final elongation (72 °C, 10 minutes).

The triplicate 50 µL PCR libraries were combined and then cleaned using the Axygen® AxyPrep

Magnetic (MAG) Bead Purification Kit (Axygen, Union City, California, USA). Cleaned PCR products

were quantified using QuantIT dsDNA BR assay (Thermo Scientific, Waltham, Massachusetts, USA)

and diluted to a concentration of 10 ng μL^{-1} before pooling libraries in equimolar concentration of 5

ng μL^{-1}. We sequenced the pooled libraries using the Illumina MiSeq platform using paired-end

(2×250 bp) reads (Illumina Reagent Kit v2, 500 reaction kit) at the Duke University Center for

Genomic and Computational Biology Core Sequencing Facility.

Sequences were processed using the mothur (v1.48.3) (Schloss et al. 2009) pipeline (Kozich

et al., 2013). We assembled contigs from the paired-end reads, quality trimmed using a moving

average quality score (minimum quality score 35), aligned sequences to the SILVA rRNA database

(v138.2) (Quast et al. 2013), and removed chimeric sequences using the VSEARCH algorithm

(Rognes et al. 2016). We created operational taxonomic units (OTUs) by first splitting sequences

based on taxonomic class and then binning into OTUs based on 97% sequence similarity.

Taxonomic identity was assigned using the SILVA rRNA database (v138.2) (Quast et al. 2013). We

used an OTU-based 3% distance threshold instead of using amplicon sequence variant (ASV) of a

single base difference, especially given our high diversity microbial communities based on short-

read (250 bp) amplicon sequences, and that similar broadscale microbiome patterns are observed

using OTU- or ASV-based approaches (Glassman and Martiny 2018). In addition, the OTU-based

approach limits overestimation of the number of ASVs due to PCR bias and sequencing errors

(Schloss 2021).

Statistical Analyses

All statistical analyses, calculations, and graphical manipulations were run in the R

environment (Rv4.5.1; R Studio 2025.05.1+513) using the vegan, ade4, picante packages, and

custom functions (Dray et al., 2023; Oksanen et al., 2022). We evaluated the main and interactive

effects of inoculation and salinity on *S. alterniflorus* aboveground biomass, belowground biomass,

root-to-shoot ratio, and change in maximum plant stem height. We tested parametric assumptions

of normality by visualizing residuals using Q-Q plots and conducting Shapiro-Wilk tests. Since the

data violated the normal distribution, we conducted Kruskal-Wallis tests and *post-hoc* Dunn's Test

(Holm *P*-value adjustment, if significance was found) to assess group differences due to

treatments.

Prior to microbial statistical analyses, we removed low-abundance OTUs that were

represented by <10 sequences across the data set. We also omitted two samples where the total

sequences were less than 10,000 reads, which represented issues with the library preparation. We

calculated the relative abundance by dividing each OTU by the total number of sequence reads for

each sample prior to conducting community composition analyses. Samples were organized by

salinity treatment and microbial treatments. Soil bacterial diversity metrics were then computed,

including Shannon Diversity (H') and Pielou's Evenness (J') with corresponding ANOVA tests,

providing insights into community composition by assessing both species richness and species

evenness.

We evaluated the extent to which inoculation and salinity treatment explained variation in

bacterial community composition using permutational multivariate analysis of variance

(PERMANOVA) based on Bray-Curtis dissimilarity. We also conducted a multivariate homogeneity

of group dispersions using the betadisper function in the vegan package (Table S5). We found that

group dispersions according to main effects and the interaction of inoculation and salinity were

similar. Therefore, significant PERMANOVA results are attributed to treatment differences and not

differences in group dispersions. Next, Principal Coordinates Analysis (PCoA), based on Bray-Curtis

dissimilarities, was used to visualize relationships among samples according to bacterial

community composition.

To identify prominent indicator species within the bacterial communities, we used the

package 'indicspecies'. Indicator species—selected based on their ability to reflect environmental

conditions—indicate changes in the environment and predict diversity patterns (De Cáceres et al.,

2010). The indicspecies package facilitates this analysis by examining the relationship between

species occurrence or abundance and site classification into groups representing habitat types or

disturbance states (De Cáceres et al., 2010).

RESULTS

Marsh Microbial Addition and Salinity Stress Influences Plants to Varying Degrees

The effects of salinity (< 0.5 psu and 20 psu) and microbial treatments (*No Addition,*

Autoclaved Inocula, Added Inocula) produced mixed results for *S. alterniflorus* growth and

biomass. A notable pattern emerged where median biomass and change in plant height were higher

in freshwater and saltwater for both the *No Addition* and *Autoclaved Inocula* treatments; however,

this pattern was reversed for the *Added Inocula* treatment, which showed higher medians under

saltwater conditions (though not always statistically significant) (Figure 2A, 2B, 2C).

The change in plant height (final height – initial height) over the course of the experiment

was similar across salinity and microbial treatments (Kruskal-Wallis, X^2(5) = 3.54, P = 0.617)

(Figure 2A). In contrast, aboveground biomass was significantly influenced by the interaction of

microbial and salinity treatments (Kruskal-Wallis, X^2(5) = 11.24, P = 0.047). Pairwise comparisons

revealed a difference in biomass between the *No Addition + Freshwater* and *Autoclaved Inocula +*

Saltwater groups (Table 1, Figure 2B). The *S. alterniflorus* plants in the freshwater control had more

aboveground biomass than plants in the saltwater autoclaved inocula treatment (Z = 2.92, P =

0.003, adj P = 0.052) (Table 1). A similar pattern was observed between *Autoclaved Inocula +*

Saltwater and *Added Inocula + Saltwater* treatments, where plants with salt marsh microorganisms

may have ameliorated salinity stress compared to autoclaved inocula (Z = -2.73, P = 0.006; adj P

=0.090) (Table 1). No other pairwise comparisons were statistically significant for aboveground

biomass analysis. Similarly, microbial and salinity manipulations affected belowground biomass

evenly (Kruskal-Wallis, X^2 (5) = 4.61, P = 0.465) (Figure 2C). Finally, the root-to-shoot ratios were

similar across treatments (Kruskal-Wallis; X^2 (5) = 2.65, P = 0.754) (Figure 2D).

Salinity and Microbial Addition Influenced Bacterial Composition More Than Diversity

269   Salinity and microbial treatments did not influence bacterial diversity. The experimental

treatments of salinity and marsh microbial inocula additions resulted in an average bacterial

Shannon Diversity H' of 6.456 $\pm$ 0.264 (Figure 3A) and Pielou's Evenness J' of 0.850 $\pm$ 0.018 (Figure

3B). Soil bacterial diversity metrics were similar across salinity (ANOVA, $F_{1,48}$ = 0.035, P = 0.853) and

microbial treatments (ANOVA, $F_{2,48}$ = 1.53, P = 0.227) (Table S3). Likewise, evenness metrics were

similar across treatments of salinity (ANOVA, $F_{1,48}$ = 0.048, P = 0.827) and microbial inocula addition

(ANOVA, $F_{2,48}$ = 1.72, P = 0.190) (Table S4).

276   Conversely, microbial and salinity treatments significantly modified bacterial community

277   composition (Table 2). The PERMANOVA, based on a Bray-Curtis dissimilarity matrix, revealed that

278   the main effects of salinity (R^2 = 0.1007, F = 6.245, P < 0.001) and microbial inocula (R^2 = 0.0853, F =

279   2.644, P < 0.001) treatments significantly influenced bacterial community composition (Table 2).

280   There was no significant interaction effect of salinity and microbial treatment on bacterial

281   community structure (R^2 = 0.0396, F = 1.228, P = 0.182).

282   To visualize the effect of treatments on bacterial community composition, we used

Principal Coordinates Analysis (PCoA) based on a Bray-Curtis dissimilarity matrix (Figure 3C, 3D).

We observed distinct bacterial community composition according to saltwater or freshwater

exposure along the second principal coordinate axis (PCoA axis 2, accounting for 11.7% of

variance). A more nuanced separation by microbial inocula was observed along the first principal

coordinate axis (PCoA axis 1, accounting for 21.9% of variance). Within both salinity groups,

communities showed a consistent gradient along PCoA axis 1. *Autoclaved Inocula* treatment

resulted in the most distinct communities, particularly under freshwater conditions, where it was

most separated from the other groups.

Context-Dependent Responses of Bacterial Indicator Taxa Across Treatments

To further examine bacterial associations with microbial and salinity treatments, we

employed indicator species analysis to identify a subset of bacterial taxa that represent the

interaction and main effects of microbial and salinity treatments (Table S1). There were 46 OTUs

significantly associated with freshwater conditions across all inocula treatments from phyla

Proteobacteria, *Acidobacteria, Actinobacteria*, *Verrucomicrobia*, *Planctomycetes*, and

Gemmatimonadetes; and 37 OTUs were associated with all saltwater treatments from phyla

Proteobacteria, *Bacteroidetes*, *Acidobacteria*, *Verrucomicrobia,* and *Spirochaetes*. In freshwater

treatments, prominent OTUs represented orders *Myxococcales*, *Caulobacterales*, *Rhizobiales,* and

Rhodospirillales, and the genus *Opitutus*. In contrast, in saltwater treatments, top OTUs

represented taxa from the orders *Myxococcales, Burkholderiales,* and *Sphingobacteriales*, family

Desulfuromonadaceae, and genus *Dongia* (Table S1).

For the microbial treatments, 28 OTUs were significantly associated with the *No Addition*

control treatment from phyla *Proteobacteria, Planctomycetes*, *Bacteroidetes*, *Acidobacteria*,

Verrucomicrobia, *and Gemmatimonadetes*; 8 OTUs were associated with *Autoclaved Inocula* from

phyla *Proteobacteria* and *Verrucomicrobia*; and 24 OTUs were associated with *Added Inocula* from

phyla *Proteobacteria*, *Verrucomicrobia*, *Acidobacteria,* and *Actinobacteria* (Table S2). The top

OTUs for the *No Addition* treatments represented taxa from the order *Myxococcales*, families

Plantomycetaceae, *Chitnophagaceae,* and genera *Rhizomicrobium* and *Opitutus*. For *Autoclaved*

Inocula, indicator species represented bacterial taxa in the orders *Myxococcales* and

Rhodospirillales, taxa in the family *Comamonadaceae,* and taxa in the genus *Hyphomicrobium*.

Finally, top indicator species for *Added Inocula* represented taxa in the genera *Geobacter*,

Prosthecobacter, Phenylobacterium, and *Opitutus,* and taxa in the class *Gp1* (Acidobacteria)

(Table S2).

DISCUSSION

While understanding the underpinnings of microbial ecology is still in its relative infancy

compared to more contemporary scientific disciplines, the roles of belowground microbial

communities in coastal wetland restoration have been of great interest (Birnbaum & Trevathan-

Tackett, 2023). Many empirical studies have demonstrated the reliance of plants on soil

microbiomes in tolerating environmental stresses, including salinity, inundation, drought, and

limited nutrient availability (reviewed in Farrer et al., 2022; Rodriguez et al., 2008). Furthermore, the

adaptive and evolutionary responses of plants to changing environmental conditions have been

shown to rely heavily on the context of microbial community assemblage, with plants exhibiting

increased fitness when associated with stress-adapted microbiomes (Lau & Lennon, 2012).

Presently, gaps in knowledge hinder the success of complete ecosystem restoration projects

(Duarte et al., 2015); therefore, the foundational knowledge gained from biological conservation

and species restoration studies is of utmost relevance and importance. In our investigation, the

interactive effects of microbial inocula additions and salinity treatment on *S. alterniflorus* height,

biomass, and bulk soil microbiomes produced various results. We hypothesized that inoculating *S.*

alterniflorus plugs with salt marsh microbial communities would buffer salt stress and promote

plant growth; a subset of our results supports this hypothesis.

The effect of salinity on plant growth depended on the microbial treatment. The addition of

live marsh microbes resulted in higher median plant biomass and changes in plant height under

saltwater compared to freshwater, while the opposite pattern was observed for plants receiving

autoclaved microbes or the no addition control. Despite these observable trends, statistical

analysis concluded that there was no significant effect of microbial and salinity treatments on the

change in plant height and belowground biomass. This null result, particularly for bioinoculation

experiments, is not uncommon across both laboratory and field settings and can be attributed to

several factors (reviewed in Compant et al., 2019). Possible explanations include, but are not

limited to, insufficient establishment of microbial communities due to the short 8-week duration of

our design; absent vital satellite taxa (i.e., rare taxa) in the given dosage of microbial inocula; and/or

the multifactorial nature of salinity stress limiting the rescue effect of inocula (e.g., alleviating

osmotic stress, but not ion toxicity), resulting in a plant growth response too subtle to detect

statistically without more precise physiological metrics and/or larger sample size. Recent evidence

has also indicated that endophytic bacteria (those within the plant tissue) provide greater benefits

to host health when under stress than rhizome-associated bacteria (Malik & Arora, 2022; Peng et

al., 2023). In all, the high variability observed in plant growth responses within treatment groups

likely overwhelmed the signal of any potential treatment effect, reducing the statistical power of our

experiment.

The findings from this study obscure a potentially important biological pattern within our

figures. Specifically, *Added Inocula* was the only microbial treatment group where saltwater

displayed a greater median of change in height than those in freshwater (Figure 2A). While this

pairwise comparison (*Added Inocula + Salt* versus *Added Inocula + Fresh*) was not significant

(Dunn's Test; P = 0.07 before adjustment), the reversal of the predicted negative effects of salinity

stress on plant growth is noteworthy and may hint at amelioration. To further investigate whether

marsh microbial inoculation ameliorates abiotic stress, we recommend future experiments with

greater statistical power (increased sample sizes) to address variability in responses, as well as

more enhanced physiological stress markers (e.g., proline accumulation, leaf ion concentration

responses to stress) (Hayat et al., 2012; Azizi et al., 2017).

Aboveground biomass results provided more nuance and clarity on the interaction of salinity

and microbial additions to plant growth. Compared to freshwater control, *Autoclaved Inocula*

under salinity stress experienced a greatly inhibited biomass (Table 1, Figure 2B). These results

provide evidence that the sterilization of belowground microbial communities and the addition of

chemically inert soil components (i.e., sand) can hinder plant productivity, possibly inducing

nutrient limitations for the host plants and nutrient cycling microbes. Comparatively, plants under

salinity stress that were inoculated with marsh microorganisms exhibited greater aboveground

biomass, further supporting a growing body of research that microbial inocula additions during

early plant development may be critical for establishing resilience to environmental stressors

(Trivedi et al., 2021; Peng et al., 2023; Shade, 2023) For example, greenhouse-raised *S. alterniflorus*

that were exposed to saltwater (10 ppt salinity) and microbial consortium (a selection of plant

growth-promoting rhizobacteria) experienced the highest rates of growth when compared to other

treatments (Bledsoe & Boopathy, 2016).

Belowground biomass and root-to-shoot ratios were similar across treatment types (Figure 2C,

2D), providing further insight into the plants' stress response strategy and possible limitations of

our methodology. The short duration of our experiment may have been insufficient to elicit

measurable changes in biomass allocation. Longer-term morphological changes, like root system

restructuring, are often not as prioritized as immediate physiological adjustments (i.e., shoot

growth) (Poorter et al., 2012). Additionally, the physical constraints of the pot sizes may have

limited root expansion and shoot growth, possibly masking treatment effects. Despite this, some

studies have suggested that optimal growth levels can be reached over a broad range of carbon

allocation patterns (Sugiura & Tateno, 2011). As *S. alterniflorus* plants under *Added Inocula + Salt*

exhibited the lowest root:shoot median compared to other treatments (Figure 2D), a compelling

hypothesis for our results is that the beneficial microorganisms (i.e., plant-growth-promoting

rhizobacteria) did not change the plant's allocation of carbon but rather improved the efficiency of

its use (Yang et al., 2009). Future work should investigate how pre-exposing *S. alterniflorus* to

salinity stress and marsh microbiomes influences long-term productivity after transplantation to

field conditions. In a natural, unrestricted setting, the evaluation of carbon allocation between

roots and shoots would be more ecologically relevant than in the restrictive environment of a pot.

Salinity stress and microbial inocula additions reshaped bacterial community composition

without altering overall diversity or evenness. When analyzing patterns in bacterial diversity, average

alpha diversity metrics Shannon H' and Pielou's evenness J' were similar across microbial and

salinity treatments (Table S3 & S4). *Autoclaved Inocula* treatment tended to decrease the ranges of

Shannon diversity (H') and evenness (J'), while greater ranges existed in the *Added Inocula* (Figure

3A, 3B). Additionally, marsh microbial inocula under salinity stress tended to exhibit greater

medians in Shannon diversity (H') and evenness (J') than those of freshwater, in support of the

reversal trend. Past studies have shown that increases in microbial biodiversity and evenness are

associated with increased resistance to plant pathogens, improved soil health and ecosystem

functions, increased plant growth and survivorship, and many other positive benefits (see Banerjee

& van der Heijden, 2023; Birnbaum & Trevathan-Tackett, 2023; Farrer et al., 2022; Graham &

Knelman, 2023; Lau & Lennon, 2012). Despite similar bacterial diversity, microbial and salinity

treatments significantly influenced community composition, with distinct bacterial communities

forming in the salt and freshwater groups (Table 2, Figure 3C, 3D). This may indicate the strength of

the selective filtering of environmental conditions in microbial community assembly, rather than

acting as forces that uniformly suppressed or enhanced microbial richness and evenness. As the

composition of microorganisms mattered more than their sheer diversity in the context of specific

environmental filters, this should be highly considered when applying microbial inocula for

restoration.

410   As microbial sampling techniques and high-throughput sequencing continue to improve,

increasing evidence has highlighted the utility of microbial communities as indicator species (Liang

et al., 2023; Ma et al., 2022; Urakawa & Bernhard, 2017). Microorganisms can function as rapid and

sensitive bioindicators and exhibit patterns of spatial distribution (e.g., distinct rhizosphere and

bulk soil communities) (Urakawa & Bernhard, 2017; Wang et al., 2016); our findings contribute to

this growing body of literature. While indicator species analysis revealed dominant microbial

assemblages from phyla *Proteobacteria*, *Acidobacteria*, *Actinobacteria*, *Bacteroidetes*, and

Verrucomicrobia—phyla known to be relatively abundant and ubiquitous in soil (Janssen, 2006)—

there were clear distinctions in OTUs with little overlap between treatments (Table S1 & S2). Seven

OTUs from the family *Opitutaceae* (phylum *Verrucomicrobia*) overlapped in *Added Inocula*,

saltwater, and freshwater treatments, which have been found in a wide variety of environments but

have been associated with nutrient cycling in low-oxygen areas (Jayakumar et al., 2017). At the

phylum/class level, many indicator OTUs from *Added Inocula* and saltwater were classified into

Verrucomicrobia, Acidobacteria GP1, and *Alphaproteobacteria*, all of which are acknowledged to

contain well-known halotolerant and/or oligotrophic bacteria (Kielak et al., 2016; Zhang et al.,

2024). Out of ten OTUs classified into *Bacteroidetes*, 70% were found exclusively in saltwater

treatments, supporting evidence of halo-tolerance (Canfora et al., 2014), while interestingly, the

remaining three were found in the *No Addition* group. In general, however, microbial bioindicators

were unique among salinity and microbial addition treatments, further providing evidence that

plant-microbe interactions and environmental conditions uniquely structure soil communities (Li et

al., 2021; Zhang et al., 2024). It is important to note that the majority of environmental microbes

remain unculturable and unidentified (Stewart, 2012), which represent 20 bacterial taxa that could

not be resolved past the phylum or kingdom levels. Moreover, microorganisms in nature often live in

complex, synergistic communities, with the vast majority of functions and interactions remaining

out of current technological and scientific grasp (Widder et al., 2016).

The patterns in bacterial community composition revealed support for strong

environmental filtering due to salinity exposure, even under short (15-minute) weekly pulses.

Bacterial community composition was relatively similar among microbial treatments but revealed

distinct bacterial assemblages when under the influence of salinity treatments. As such, the pre-

treatment of saline conditions and resident microbial inocula may provide rescue effects from

abiotic and biotic field stressors after marsh grass plugs are transplanted and exposed to ex situ

environmental conditions. This work provides evidence that microbial stewardship is essential for

buffering against ecological stressors and could promote successful plant establishment for

wetland restoration. Additionally, our work supports the theory of "habitat-adapted symbioses,"

highlighting the facultative mutualisms between marsh plants and their symbiotic microorganisms

in tolerating high-stress environments (Rodriguez et al., 2008). With many stressors projected to

intensify in the near future (e.g., extreme climatic events, sea level rise, droughts and heatwaves,

compounding stress from invasive species, anthropogenic change, etc.), marshes will need to

adapt to survive, and their associated microorganisms will undergo even more rapid evolutionary

changes (Zabin et al., 2022; Billah et al., 2022; Lau & Lennon, 2012; Mueller et al., 2020). By

incorporating microbial inocula into salt marsh restoration practices, success could be better

leveraged.

Within a broader scope, coastal wetlands and salt marshes are among the most productive

ecosystems on Earth and provide numerous ecological and economic services. Salt marshes

support fisheries and habitats for biodiversity, filter contaminants, sequester carbon, buffer storm

surges and protect shorelines from erosion, provide tourism, and facilitate many other functions

(Arkema et al., 2013; Buchanan et al., 2022). The preservation and restoration of these habitats are

thus vital, aligning with the principles of the One Health framework, which emphasizes how the

health of humans, wildlife, and the environment is inextricably linked (Trinh et al., 2018; Peixoto et

al., 2022; Banerjee & van der Heijden, 2023). By evaluating the interactions between *Sporobolus*

alterniflorus, marsh sediment microbiomes, and environmental stressors, this study provides

insight into ecological dynamics and may aid in salt marsh restoration efforts.

Acknowledgments

We acknowledge the staff at Hammocks Beach State Park and Garner's Landscaping for their

assistance in conducting the field experiment. We thank Gittman lab members Jennifer Fickler and

James Kelley for assistance in setting up and maintaining the field research, and Peralta lab

members Toby Gallegos, Daphka Joseph, Scott Siebor, Dejah Smith, and Colin Finlay for lab

assistance. We also thank Aeran Coughlin and Jean-Philippe Gibert for microbiome sequencing

support. This work was supported by the National Science Foundation (#1845845, #2009185, #

2302609 to A.L.P.), the U.S. Coastal Research Program and the National Oceanographic and

Atmospheric Administration National Sea Grant (#2023-0534-02/22-NC-USCRP-3 to R.K.G.) and

North Carolina Sea Grant (#PAM-P24-002985-SA-02 to R.K.G. and A.L.P.) and East Carolina

University (Undergraduate Research and Creative Activity Award to K.A.D.).

REFERENCES

Apprill, A., McNally, S., Parsons, R., & Weber, L. (2015). Minor revision to V4 region SSU rRNA 806R

gene primer greatly increases detection of SAR11 bacterioplankton. *Aquatic Microbial*

Ecology, *75*. https://doi.org/10.3354/ame01753

Arkema, K. K., Guannel, G., Verutes, G., Wood, S. A., Guerry, A., Ruckelshaus, M., Kareiva, P.,

Lacayo, M., & Silver, J. M. (2013). Coastal habitats shield people and property from sea-level

rise and storms. *Nature Climate Change*, *3*(10), Article 10.

https://doi.org/10.1038/nclimate1944

Azizi, S., Tabari, M., & Striker, G. G. (2017). Growth, physiology, and leaf ion concentration

responses to long-term flooding with fresh or saline water of *Populus euphratica*. *South*

African Journal of Botany, 108, 229–236. https://doi.org/10.1016/j.sajb.2016.11.004

Banerjee, S., & van der Heijden, M. G. A. (2023). Soil microbiomes and one health. *Nature Reviews*

Microbiology, *21*(1), 6–20. https://doi.org/10.1038/s41579-022-00779-w

Billah, M. M., Bhuiyan, M. K. A., Islam, M. A., Das, J., & Hoque, A. R. (2022). Salt marsh restoration:

An overview of techniques and success indicators. *Environmental Science and Pollution*

Research, *29*(11), 15347–15363. https://doi.org/10.1007/s11356-021-18305-5

Birnbaum, C., & Trevathan-Tackett, S. M. (2023). Aiding coastal wetland restoration via the

belowground soil microbiome: An overview. *Restoration Ecology*, *31*(7), e13824.

https://doi.org/10.1111/rec.13824

Bledsoe, R., & Boopathy, R. (2016). Bioaugmentation of microbes to restore coastal wetland plants

to protect land from coastal erosion. *International Biodeterioration & Biodegradation*, *113*,

155–160. https://doi.org/10.1016/j.ibiod.2016.02.010

Bruno, J. F. (2000). Facilitation of Cobble Beach Plant Communities Through Habitat Modification by

Spartina Alterniflora. *Ecology*, *81*(5), 1179–1192. https://doi.org/10.1890/0012-

9658(2000)081[1179:FOCBPC]2.0.CO;2

Buchanan, M. K., Kulp, S., & Strauss, B. (2022). Resilience of U.S. coastal wetlands to accelerating

sea level rise. *Environmental Research Communications*, *4*(6), 061001.

https://doi.org/10.1088/2515-7620/ac6eef

Canfora, L., Bacci, G., Pinzari, F., Papa, G. L., Dazzi, C., & Benedetti, A. (2014). Salinity and Bacterial

Diversity: To What Extent Does the Concentration of Salt Affect the Bacterial Community in

a Saline Soil? *PLOS ONE*, *9*(9), e106662. https://doi.org/10.1371/journal.pone.0106662

Caporaso, J. G., Lauber, C. L., Walters, W. A., Berg-Lyons, D., Huntley, J., Fierer, N., Owens, S. M.,

Betley, J., Fraser, L., Bauer, M., Gormley, N., Gilbert, J. A., Smith, G., & Knight, R. (2012).

Ultra-high-throughput microbial community analysis on the Illumina HiSeq and MiSeq

platforms. *The ISME Journal*, *6*(8), 1621–1624. https://doi.org/10.1038/ismej.2012.8

Cavicchioli, R., Ripple, W. J., Timmis, K. N., Azam, F., Bakken, L. R., Baylis, M., Behrenfeld, M. J.,

Boetius, A., Boyd, P. W., Classen, A. T., Crowther, T. W., Danovaro, R., Foreman, C. M.,

Huisman, J., Hutchins, D. A., Jansson, J. K., Karl, D. M., Koskella, B., Mark Welch, D. B., …

Webster, N. S. (2019). Scientists' warning to humanity: Microorganisms and climate change.

Nature Reviews Microbiology, *17*(9), Article 9. https://doi.org/10.1038/s41579-019-0222-5

Chepsergon, J., & Moleleki, L. N. (2023). Rhizosphere bacterial interactions and impact on plant

health. *Current Opinion in Microbiology*, *73*, 102297.

https://doi.org/10.1016/j.mib.2023.102297

Choudoir, M. J., Ishaq, S. L., Beiko, R. G., Silva, D. S., Allen-Vercoe, E., & O'Doherty, K. C. (2025). The

case for microbiome stewardship: What it is and how to get there. *mSystems, 10*(5),

e00062-25. https://doi.org/10.1128/msystems.00062-25

Currin, C. A., Delano, P. C., & Valdes-Weaver, L. M. (2008). Utilization of a citizen monitoring

protocol to assess the structure and function of natural and stabilized fringing salt marshes

in North Carolina. *Wetlands Ecology and Management*, *16*(2), 97–118.

https://doi.org/10.1007/s11273-007-9059-1

De Cáceres, M., Legendre, P., & Moretti, M. (2010). Improving indicator species analysis by

combining groups of sites. *Oikos*, *119*(10), 1674–1684. https://doi.org/10.1111/j.1600-

0706.2010.18334.x

Dini-Andreote, F. (2025). Plant-microbe interactions amid global change. *Cell Host & Microbe*,

33(8), 1217–1221. https://doi.org/10.1016/j.chom.2025.05.028

Dray, S., Dufour, A.-B., Thioulouse, and J., Jombart, with contributions from T., Pavoine, S., Lobry, J.

R., Ollier, S., Borcard, D., Legendre, P., & Chessel, S. B. and A. S. B. on earlier work by D.

(2023). *ade4: Analysis of Ecological Data: Exploratory and Euclidean Methods in*

Environmental Sciences (1.7-22) [Computer software]. https://cran.r-

project.org/web/packages/ade4/index.html

Duarte, C. M., Borja, A., Carstensen, J., Elliott, M., Krause-Jensen, D., & Marbà, N. (2015).

Paradigms in the Recovery of Estuarine and Coastal Ecosystems. *Estuaries and Coasts*,

38(4), 1202–1212. https://doi.org/10.1007/s12237-013-9750-9

Farrer, E. C., Van Bael, S. A., Clay, K., & Smith, M. K. H. (2022). Plant-Microbial Symbioses in Coastal

Systems: Their Ecological Importance and Role in Coastal Restoration. *Estuaries and*

Coasts, *45*(7), 1805–1822. https://doi.org/10.1007/s12237-022-01052-2

Gedan, K. B., & Bertness, M. D. (2010). How will warming affect the salt marsh foundation species

Spartina patens and its ecological role? *Oecologia*, *164*(2), 479–487.

https://doi.org/10.1007/s00442-010-1661-x

Gittman, R. K., Fodrie, F. J., Baillie, C. J., Brodeur, M. C., Currin, C. A., Keller, D. A., Kenworthy, M. D.,

Morton, J. P., Ridge, J. T., & Zhang, Y. S. (2018). Living on the Edge: Increasing Patch Size

Enhances the Resilience and Community Development of a Restored Salt Marsh. *Estuaries*

and Coasts, *41*(3), 884–895. https://doi.org/10.1007/s12237-017-0302-6

Gittman, R. K., Baillie, C. J., Arkema, K. K., Bennett, R. O., Benoit, J., Blitch, S., Brun, J., Chatwin, A.,

Colden, A., Dausman, A., DeAngelis, B., Herold, N., Henkel, J., Houge, R., Howard, R.,

Hughes, A. R., Scyphers, S. B., Shostik, T., Sutton-Grier, A., & Grabowski, J. H. (2019).

Voluntary Restoration: Mitigation's Silent Partner in the Quest to Reverse Coastal Wetland

Loss in the USA. *Frontiers in Marine Science*, *6*. https://doi.org/10.3389/fmars.2019.00511

Gittman, R. K., Baillie, C. J., Cros, A., Grabowski, J. H., McKinney, M.-M., Saccomanno, V. R., Smith,

Carter. S., & DeAngelis, B. (2024). Assessing how restoration can facilitate 30×30 goals for

climate-resilient coastal ecosystems in the United States. *Conservation Biology*, *n/a*(n/a),

e14429. https://doi.org/10.1111/cobi.14429

Glassman, S. I., & Martiny, J. B. H. (2018). Broadscale Ecological Patterns Are Robust to Use of

Exact Sequence Variants versus Operational Taxonomic Units. *mSphere*, *3*(4),

10.1128/msphere.00148-18. https://doi.org/10.1128/msphere.00148-18

Graham, E. B., & Knelman, J. E. (2023). Implications of Soil Microbial Community Assembly for

Ecosystem Restoration: Patterns, Process, and Potential. *Microbial Ecology*, *85*(3), 809–819.

https://doi.org/10.1007/s00248-022-02155-w

Hayat, S., Hayat, Q., Alyemeni, M. N., Wani, A. S., Pichtel, J., & Ahmad, A. (2012). Role of proline

under changing environments. *Plant Signaling & Behavior*, *7*(11), 1456–1466.

https://doi.org/10.4161/psb.21949

Janssen, P. H. (2006). Identifying the Dominant Soil Bacterial Taxa in Libraries of 16S rRNA and 16S

rRNA Genes. *Applied and Environmental Microbiology*, *72*(3), 1719–1728.

https://doi.org/10.1128/AEM.72.3.1719-1728.2006

Jayakumar, A., Chang, B. X., Widner, B., Bernhardt, P., Mulholland, M. R., & Ward, B. B. (2017).

Biological nitrogen fixation in the oxygen-minimum region of the eastern tropical North

Pacific ocean. *The ISME Journal*, *11*(10), 2356–2367. https://doi.org/10.1038/ismej.2017.97

Kielak, A. M., Barreto, C. C., Kowalchuk, G. A., van Veen, J. A., & Kuramae, E. E. (2016). The Ecology

of Acidobacteria: Moving beyond Genes and Genomes. *Frontiers in Microbiology*, *7*, 744.

https://doi.org/10.3389/fmicb.2016.00744

Kozich, J. J., Westcott, S. L., Baxter, N. T., Highlander, S. K., & Schloss, P. D. (2013). Development of

a Dual-Index Sequencing Strategy and Curation Pipeline for Analyzing Amplicon Sequence

Data on the MiSeq Illumina Sequencing Platform. *Applied and Environmental Microbiology*,

79(17), 5112–5120. https://doi.org/10.1128/AEM.01043-13

Lau, J. A., & Lennon, J. T. (2011). Evolutionary ecology of plant–microbe interactions: Soil microbial

structure alters selection on plant traits. *New Phytologist*, *192*(1), 215–224.

https://doi.org/10.1111/j.1469-8137.2011.03790.x

Lau, J. A., & Lennon, J. T. (2012). Rapid responses of soil microorganisms improve plant fitness in

novel environments. *Proceedings of the National Academy of Sciences*, *109*(35), 14058–

14062. https://doi.org/10.1073/pnas.1202319109

Leonardi, N., Carnacina, I., Donatelli, C., Ganju, N. K., Plater, A. J., Schuerch, M., & Temmerman, S.

(2018). Dynamic interactions between coastal storms and salt marshes: A review.

Geomorphology, *301*, 92–107. https://doi.org/10.1016/j.geomorph.2017.11.001

Li, H., La, S., Zhang, X., Gao, L., & Tian, Y. (2021). Salt-induced recruitment of specific root-

associated bacterial consortium capable of enhancing plant adaptability to salt stress. *The*

ISME Journal, *15*(10), 2865–2882. https://doi.org/10.1038/s41396-021-00974-2

Liang, S., Li, H., Wu, H., Yan, B., & Song, A. (2023). Microorganisms in coastal wetland sediments: A

review on microbial community structure, functional gene, and environmental potential.

Frontiers in Microbiology, *14*. https://doi.org/10.3389/fmicb.2023.1163896

Lin, C., Wang, Y., Liu, M., Li, Q., Xiao, W., & Song, X. (2020). Effects of nitrogen deposition and

phosphorus addition on arbuscular mycorrhizal fungi of Chinese fir (Cunninghamia

lanceolata). *Scientific Reports, 10*(1), 12260. https://doi.org/10.1038/s41598-020-69213-6

Liu, F., Hewezi, T., Lebeis, S. L., Pantalone, V., Grewal, P. S., & Staton, M. E. (2019). Soil indigenous

microbiome and plant genotypes cooperatively modify soybean rhizosphere microbiome

assembly. *BMC Microbiology*, *19*(1), 201. https://doi.org/10.1186/s12866-019-1572-x

Ma, F., Wang, C., Zhang, Y., Chen, J., Xie, R., & Sun, Z. (2022). Development of Microbial Indicators

in Ecological Systems. *International Journal of Environmental Research and Public Health*,

19(21), Article 21. https://doi.org/10.3390/ijerph192113888

Malik, G., & Arora, R. (2022). Endophytic Bacteria: Mitigating Abiotic Stress from Inside. In A. K.

Singh, V. Tripathi, A. K. Shukla, & P. Kumar (Eds.), *Bacterial Endophytes for Sustainable*

Agriculture and Environmental Management (pp. 23–43). Springer.

https://doi.org/10.1007/978-981-16-4497-9_2

Mariotte, P., Mehrabi, Z., Bezemer, T. M., De Deyn, G. B., Kulmatiski, A., Drigo, B., Veen, G. F. (Ciska),

van der Heijden, M. G. A., & Kardol, P. (2018). Plant–Soil Feedback: Bridging Natural and

Agricultural Sciences. *Trends in Ecology & Evolution*, *33*(2), 129–142.

https://doi.org/10.1016/j.tree.2017.11.005

McHugh, J. M., & Dighton, J. (2004). Influence of Mycorrhizal Inoculation, Inundation Period,

Salinity, and Phosphorus Availability on the Growth of Two Salt Marsh Grasses, *Spartina*

alterniflora Lois. And *Spartina cynosuroides* (L.) Roth., in Nursery Systems. *Restoration*

Ecology, *12*(4), 533–545. https://doi.org/10.1111/j.1061-2971.2004.03109.x

Mendelssohn, I. A., & Seneca, E. D. (1980). The influence of soil drainage on the growth of salt

marsh cordgrass Spartina alterniflora in North Carolina. *Estuarine and Coastal Marine*

Science, *11*(1), 27–40. https://doi.org/10.1016/S0302-3524(80)80027-2

Morris, J. T., & Haskin, B. (1990). A 5-yr Record of Aerial Primary Production and Stand

Characteristics of Spartina Alterniflora. *Ecology*, *71*(6), 2209–2217.

https://doi.org/10.2307/1938633

Mueller, E. A., Wisnoski, N. I., Peralta, A. L., & Lennon, J. T. (2020). Microbial rescue effects: How

microbiomes can save hosts from extinction. *Functional Ecology*, *34*(10), 2055–2064.

https://doi.org/10.1111/1365-2435.13493

Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R., O'Hara, R. B.,

Solymos, P., Stevens, M. H. H., Szoecs, E., Wagner, H., Barbour, M., Bedward, M., Bolker, B.,

Borcard, D., Carvalho, G., Chirico, M., Caceres, M. D., Durand, S., … Weedon, J. (2022).

vegan: Community Ecology Package (2.6-4) [Computer software]. https://cran.r-

project.org/web/packages/vegan/index.html

Pantigoso, H. A., Newberger, D., & Vivanco, J. M. (2022). The rhizosphere microbiome: Plant–

microbial interactions for resource acquisition. *Journal of Applied Microbiology*, *133*(5),

2864–2876. https://doi.org/10.1111/jam.15686

Parada, A. E., Needham, D. M., & Fuhrman, J. A. (2016). Every base matters: Assessing small

subunit rRNA primers for marine microbiomes with mock communities, time series and

646  global field samples. *Environmental Microbiology*, *18*(5), 1403–1414.

647  https://doi.org/10.1111/1462-2920.13023

648 Peixoto, R. S., Voolstra, C. R., Sweet, M., Duarte, C. M., Carvalho, S., Villela, H., Lunshof, J. E., Gram,

649  L., Woodhams, D. C., Walter, J., Roik, A., Hentschel, U., Thurber, R. V., Daisley, B., Ushijima,

650  B., Daffonchio, D., Costa, R., Keller-Costa, T., Bowman, J. S., … Berg, G. (2022). Harnessing

651  the microbiome to prevent global biodiversity loss. *Nature Microbiology*.

652  https://doi.org/10.1038/s41564-022-01173-1

653 Peng, M., Jiang, Z., Zhou, F., & Wang, Z. (2023). From salty to thriving: Plant growth promoting

654  bacteria as nature's allies in overcoming salinity stress in plants. *Frontiers in Microbiology,*

655  *14.* https://doi.org/10.3389/fmicb.2023.1169809

656 Poorter, H., Niklas, K. J., Reich, P. B., Oleksyn, J., Poot, P., & Mommer, L. (2012). Biomass allocation

657  to leaves, stems and roots: Meta-analyses of interspecific variation and environmental

658  control. *New Phytologist*, 193(1), 30–50. https://doi.org/10.1111/j.1469-8137.2011.03952.x

659 Rodriguez, R. J., Henson, J., Van Volkenburgh, E., Hoy, M., Wright, L., Beckwith, F., Kim, Y.-O., &

660  Redman, R. S. (2008). Stress tolerance in plants via habitat-adapted symbiosis. *The ISME*

661  *Journal*, *2*(4), 404–416. https://doi.org/10.1038/ismej.2007.106

662 Rognes, T., Flouri, T., Nichols, B., Quince, C., & Mahé, F. (2016). VSEARCH: A versatile open source

663  tool for metagenomics. *PeerJ*, *4*, e2584. https://doi.org/10.7717/peerj.2584

664 Rojas-Sánchez, B., de los Santos-Villalobos, S., Alarcón, J. J. V., Chávez-Bárcenas, A. T., Orozco-

665  Mosqueda, Ma. del C., & Santoyo, G. (2025). Optimizing milpa agrosystems with beneficial

666  microbes and their ecological interactions: A review. *Discover Applied Sciences*, *7*(2), 104.

667  https://doi.org/10.1007/s42452-025-06503-6

Rudershausen, P. J., Lombardo, S. M., & Buckel, J. A. (2021). Linking Historical Changes in Salt-

Marsh Coverage to Lost Production of a Nektonic Bioindicator. *Marine and Coastal*

Fisheries, *13*(2), 131–139. https://doi.org/10.1002/mcf2.10147

Schloss, P. D. (2021). Amplicon Sequence Variants Artificially Split Bacterial Genomes into

Separate Clusters. *mSphere*, *6*(4), e00191-21. https://doi.org/10.1128/mSphere.00191-21

Shade, A. (2023). Microbiome rescue: Directing resilience of environmental microbial

communities. *Current Opinion in Microbiology*, 72, 102263.

https://doi.org/10.1016/j.mib.2022.102263

Singh, B. K., Delgado-Baquerizo, M., Egidi, E., Guirado, E., Leach, J. E., Liu, H., & Trivedi, P. (2023).

Climate change impacts on plant pathogens, food security and paths forward. *Nature*

Reviews Microbiology, *21*(10), Article 10. https://doi.org/10.1038/s41579-023-00900-7

Sloey, T. M., Gieser, K. D., Pounders, K., Báez, S. E., & Whytlaw, J. L. (2025). Efficacy of stress

acclimation techniques in coastal wetland plants (Spartina alterniflora and S. patens):

Implications for restoration and nursery production. *Wetlands Ecology and Management*,

33(5), 64. https://doi.org/10.1007/s11273-025-10079-8

Stewart, E. J. (2012). Growing Unculturable Bacteria. *Journal of Bacteriology*, 194(16), 4151–4160.

https://doi.org/10.1128/jb.00345-12

Sugiura, D., & Tateno, M. (2011). Optimal Leaf-to-Root Ratio and Leaf Nitrogen Content Determined

by Light and Nitrogen Availabilities. *PLOS ONE*, 6(7), e22236.

https://doi.org/10.1371/journal.pone.0022236

Trinh, P., Zaneveld, J. R., Safranek, S., & Rabinowitz, P. M. (2018). One Health Relationships Between

Human, Animal, and Environmental Microbiomes: A Mini-Review. *Frontiers in Public Health,*

6. https://www.frontiersin.org/articles/10.3389/fpubh.2018.00235

Trivedi, P., Mattupalli, C., Eversole, K., & Leach, J. E. (2021). Enabling sustainable agriculture

through understanding and enhancement of microbiomes. New Phytologist, 230(6), 2129–

2147. https://doi.org/10.1111/nph.17319

Trivedi, P., Batista, B. D., Bazany, K. E., & Singh, B. K. (2022). Plant–microbiome interactions under a

changing world: Responses, consequences and perspectives. *New Phytologist, 234*(6),

1951–1959. https://doi.org/10.1111/nph.18016

Urakawa, H., & Bernhard, A. E. (2017). Wetland management using microbial indicators. *Ecological*

Engineering, *108*, 456–476. https://doi.org/10.1016/j.ecoleng.2017.07.022

Valiela, I., Lloret, J., Bowyer, T., Miner, S., Remsen, D., Elmstrom, E., Cogswell, C., & Robert Thieler,

E. (2018). Transient coastal landscapes: Rising sea level threatens salt marshes. *Science of*

The Total Environment, *640–641*, 1148–1156.

https://doi.org/10.1016/j.scitotenv.2018.05.235

Wang, F., Kroeger, K. D., Gonneea, M. E., Pohlman, J. W., & Tang, J. (2019). Water salinity and

inundation control soil carbon decomposition during salt marsh restoration: An incubation

experiment. *Ecology and Evolution*, *9*(4), 1911–1921. https://doi.org/10.1002/ece3.4884

Wang, M., Yang, P., & Falcão Salles, J. (2016). Distribution of Root-Associated Bacterial

Communities Along a Salt-Marsh Primary Succession. *Frontiers in Plant Science,* 6.

https://doi.org/10.3389/fpls.2015.01188

Wei, C., Yu, Q., Bai, E., Lü, X., Li, Q., Xia, J., Kardol, P., Liang, W., Wang, Z., & Han, X. (2013). Nitrogen

deposition weakens plant–microbe interactions in grassland ecosystems. *Global Change*

Biology, 19(12), 3688–3697. https://doi.org/10.1111/gcb.12348

Widder, S., Allen, R. J., Pfeiffer, T., Curtis, T. P., Wiuf, C., Sloan, W. T., Cordero, O. X., Brown, S. P.,

Momeni, B., Shou, W., Kettle, H., Flint, H. J., Haas, A. F., Laroche, B., Kreft, J.-U., Rainey, P.

B., Freilich, S., Schuster, S., Milferstedt, K., ... Soyer, O. S. (2016). Challenges in microbial

ecology: Building predictive understanding of community function and dynamics. *The ISME*

Journal, *10*(11), 2557–2568. https://doi.org/10.1038/ismej.2016.45

Yang, J., Kloepper, J. W., & Ryu, C.-M. (2009). Rhizosphere bacteria help plants tolerate abiotic

stress. *Trends in Plant Science*, 14(1), 1–4. https://doi.org/10.1016/j.tplants.2008.10.004

Zabin, C. J., Jurgens, L. J., Bible, J. M., Patten, M. V., Chang, A. L., Grosholz, E. D., & Boyer, K. E.

(2022). Increasing the resilience of ecological restoration to extreme climatic events.

Frontiers in Ecology and the Environment, *20*(5), 310–318. https://doi.org/10.1002/fee.2471

Zhang, G., Bai, J., Zhai, Y., Jia, J., Zhao, Q., Wang, W., & Hu, X. (2024). Microbial diversity and

functions in saline soils: A review from a biogeochemical perspective. *Journal of Advanced*

Research, *59*, 129–140. https://doi.org/10.1016/j.jare.2023.06.015

STATEMENTS AND DECLARATIONS

The authors declare no competing interests.

FIGURES

Figure 1.

Figure 1. Experimental design of salinity and marsh sediment microbial inocula treatments.

Factorial experiment includes three levels of microbial addition (*No Addition*, *Autoclaved Inocula*,

Added Inocula) and two levels of salinity (<0.5 psu, 20 psu). Treatments were replicated ten times.

Microbial inocula were combined with potting mix and established for four weeks, followed by

weekly salt stress treatment, and harvested after eight weeks.

Figure 2.

Figure 2. Boxplots depicting the change in plant height (in mm) **(A)**, aboveground biomass **(B)**,

belowground root biomass **(C)**, and root-to-shoot ratio **(D)** on the y-axes, in response to marsh soil

microbes (*No Addition, Autoclaved Inocula, Added Inocula*) and salinity stress (<0.5 psu FRESH, 20

psu SALT) treatments on the x-axes. Color represents salinity stress (orange) and freshwater control

(blue), while symbols represent individual data points. The boxplot is a visual representation of 5

key summary statistics: the median, the 25% and 75% percentiles, and the whiskers, which

represent the feasible range of the data as determined by 1.5 × the interquartile range. Change in

plant height, belowground biomass, and root:shoot ratio results were not significant (NS). For

aboveground biomass **(B)**, different letters above boxplots indicate significant difference at

adjusted *P* < 0.10 based on Dunn's test pairwise comparisons (see also Table 1).

Figure 3.

Figure 3. Boxplots for bacterial diversity representing Shannon Diversity Index (H') **(A)** and Pielou's

Evenness (J') **(B)** associated with microbial and salinity treatments. The average Shannon Diversity

H' equaled 6.456 ± 0.264, and Pielou's Evenness J' equaled 0.850 ± 0.018, with no significant

differences found across treatments. Also included are ordination plots based on a principal

coordinates analysis depicting bacterial community composition according to salinity and

microbial treatments, representing the average centroid across treatments **(C)** and individual

samples **(D).** Main effects of salinity (PERMANOVA: R^2 = 0.1007, P < 0.001) and microbial

(PERMANOVA: R^2 = 0.0853, P < 0.001) treatments significantly influenced bacterial community

composition. No significant effect of salinity × microbial treatment on community structure

(PERMANOVA: R^2 = 0.0396, P = 0.182) (See Table 2). Symbols are colored according to salinity

treatment (orange = saltwater (SALT), blue = freshwater (FRESH)) and shaped by microbial

treatment (circles = *Autoclaved Inocula*, triangles = *Added Inocula*, squares = *No Addition*).

803 **Table 1.** Non-parametric *post-hoc* Dunn's test for pairwise comparisons of aboveground biomass

804 across microbial and salinity treatments. *P*-value adjustment: Holm method. *Note*: Kruskal-Wallis

805 test statistic: X^2 (5) = 11.24, *P* = 0.047.

Comparison	*Z Statistic*	*P-value*	*Adjusted P-value*
No Addition Fresh – No Addition Salt	1.18	0.240	1.00
No Addition Fresh – Autoclaved Inocula Fresh	1.68	0.092	1.00
No Addition Salt – Autoclaved Inocula Fresh	0.573	0.566	1.00
No Addition Fresh – Autoclaved Inocula Salt	2.92	0.003	**0.052**
No Addition Salt – Autoclaved Inocula Salt	1.92	0.055	0.709
Autoclaved Inocula Fresh – Autoclaved Inocula Salt	1.33	0.182	1.00
No Addition Fresh – Added Inocula Fresh	1.49	0.135	1.00
No Addition Salt – Added Inocula Fresh	0.372	0.710	1.00
Autoclaved Inocula Fresh – Added Inocula Fresh	-0.196	0.844	0.844
Autoclaved Inocula Salt– Added Inocula Fresh	-1.52	0.128	1.00
No Addition Fresh – Added Inocula Salt	0.372	0.710	1.00
No Addition Salt – Added Inocula Salt	-0.859	0.390	1.00
Autoclaved Inocula Fresh – Added Inocula Salt	-1.41	0.159	1.00
Autoclaved Inocula Salt – Added Inocula Salt	-2.73	0.006	**0.090**
Added Inocula Fresh – Added Inocula Salt	-1.20	0.228	1.00

807 **Table 2.** Summary PERMANOVA results comparing microbial community composition (based on

808 the Bray-Curtis dissimilarity matrix) across main effects (salinity treatment, microbial inocula

809 treatment) and the interaction of salinity × microbial treatments. Bold text indicates significant

810 differences (*P* ≤ 0.05).

Effect	*DF*	Sum of Sqs	R^2	*F-value*	*P-value*
Salinity TRT	1	0.4931	0.1007	6.245	**<0.001**
Microbe TRT	2	0.4176	0.0853	2.644	**<0.001**
Salinity TRT × Microbe TRT	2	0.1940	0.0396	1.228	0.182
Residual	48	3.7900	0.7743		
Total	53	4.8946	1.0000		