An increase in reactive oxygen species underlies neonatal cerebellum repair

Anna Pakula[1,#], Salsabiel El Nagar[1,#], N. Sumru Bayin[1,2,3,#], Jens Bager Christensen[2,3], Daniel N. Stephen[1], Adam James Reid[2], Richard Koche[4], Alexandra L. Joyner[1,5,*]

[1]Developmental Biology Program, Sloan Kettering Institute, New York, NY, USA

[2]Gurdon Institute, Cambridge University, Cambridge, UK

[3]Department of Physiology, Development and Neuroscience, Cambridge University, Cambridge, UK

[4]Center for Epigenetics Research, Memorial Sloan Kettering Cancer Center, New York, NY

[5]Biochemistry, Cell and Molecular Biology Program and Neuroscience Program, Weill Cornell Graduate School of Medical Sciences, New York, NY, USA

[#]These authors contributed equally to the work and share first authorship

* Correspondence:
joynera@mskcc.org

Keywords: Nestin-expressing progenitors, NEPs, granule cell progenitors, ROS, regeneration

Abstract

The neonatal mouse cerebellum shows remarkable regenerative potential upon injury at birth, wherein a subset of Nestin-expressing progenitors (NEPs) undergoes adaptive reprogramming to replenish granule cell progenitors that die. Here, we investigate how the microenvironment of the injured cerebellum changes upon injury and contributes to the regenerative potential of normally gliogenic-NEPs and their adaptive reprogramming. Single cell transcriptomic and bulk chromatin accessibility analyses of the NEPs from injured neonatal cerebella compared to controls show a temporary increase in cellular processes involved in responding to reactive oxygen species (ROS), a known damage-associated molecular pattern. Analysis of ROS levels in cerebellar tissue confirm a transient increased one day after injury at postanal day 1, overlapping with the peak cell death in the cerebellum. In a transgenic mouse line that ubiquitously overexpresses human mitochondrial catalase (mCAT), ROS is reduced 1 day after injury to the granule cell progenitors, and we demonstrate that several steps in the regenerative process of NEPs are curtailed leading to reduced cerebellar growth. We also provide evidence that microglia are involved in one step of adaptive reprogramming by regulating NEP replenishment of the granule cell precursors. Collectively, our results highlight that changes in the tissue microenvironment regulate multiple steps in adaptive reprogramming of NEPs upon death of cerebellar granule cell progenitors at birth, highlighting the instructive roles of microenvironmental signals during regeneration of the neonatal brain.

Introduction

The microenvironment surrounding a brain injury and the cellular responses elicited in the remaining cells are key determinants of how efficiently a repair process will unfold. An important factor underlying the effectiveness of regenerative responses to an injury is the plasticity of the stem/progenitor cells in a tissue (Burda and Sofroniew, 2014). The degree to which the microenvironment and specific cell types within it provide pro- or anti-regenerative factors is highly context dependent. The neonatal mouse cerebellum has a remarkable capacity to regenerate cells ablated around birth (Wojcinski et al., 2017, Bayin N. S., 2021, Bayin et al., 2018, Altman and Anderson, 1971). Thus, the cerebellum provides an ideal system to study the roles that signals in the microenvironment play in key steps of the repair process in the brain.

The cerebellum is a folded hindbrain structure that is critical for motor coordination. It also participates in higher order social and cognitive behaviors through its circuit connections with all other brain regions (Badura et al., 2018, Buckner, 2013, Burda and Sofroniew, 2014, Salman and Tsai, 2016, Strick et al., 2009, Tomlinson et al., 2013). Compared to the rest of the brain, the cerebellum has protracted development, as its major growth occurs during the first two weeks after birth in mice and at least six months surrounding birth in humans (Altman and Bayer, 1997, Rakic and Sidman, 1970, Dobbing and Sands, 1973). This timing of the major growth of the cerebellum makes it susceptible to injury around birth. Indeed, cerebellar hypoplasia is the second leading risk factor for autism spectrum disorders and cerebellar injury around birth can have devastating outcomes and significant effects on subsequent quality of life (Tsai et al., 2018, Stoodley et al., 2017, Wang et al., 2014). Therefore, it is critical to better understand the regenerative processes that allow repair of the cerebellum.

All the cell types in the cerebellum are derived from two progenitor zones, the embryonic rhombic lip and the ventricular zone that give rise to the excitatory neurons, or the inhibitory neurons and glia, respectively (Leto et al., 2015, Joyner and Bayin, 2022). During postnatal growth, the rhombic lip-derived granule cell precursors (GCPs) cover the surface of the cerebellum in a structure named the external granule layer (EGL) and continue to proliferate in a sonic hedgehog (SHH) dependent manner for two weeks after birth in mice (Wechsler-Reya and Scott, 1999, McMahon et al., 2003, Corrales et al., 2006). Following their exit from the cell cycle, the granule cells (GC) migrate inwards to form the internal granule layer (IGL). Other SHH-dependent progenitor populations of the neonatal cerebellum are either gliogenic Nestin-Expressing Progenitors (NEPs)

that express SOX2 and generate astroglia (astrocytes and Bergman glia) or neurogenic-NEPs

that generate late born interneurons (Bayin N. S., 2021, Cerrato et al., 2018, Parmigiani et al.,

2015). Gliogenic-NEPs reside either in the Bergmann glia layer (BgL) intermixed with Purkinje

cells and generate Bergmann glia (Bg) and astrocytes, or in the white matter in the center of the

lobules (folds) and generate astrocytes. Neurogenic-NEPs are restricted to the white matter and

produce interneurons that migrate outwards to the outermost molecular layer (Bayin N. S., 2021,

Brown et al., 2020, De Luca et al., 2015). Surprisingly, when the GCPs are killed upon injury soon

after birth, the gliogenic-NEPs in the BgL (BgL-NEPs) undergo adaptive reprogramming to

generate GCPs and replenish the EGL via a transitory cellular state that involves upregulation of

the neurogenic gene *Ascl1* to promote a glial-to-neural fate switch (Wojcinski et al., 2017, Bayin

N. S., 2021). Adaptive reprogramming involves multiple sequential stages starting with increased

proliferation of BgL-NEPs, then a fate switch to neuronal progenitors, migration to the site of injury

(EGL) and acquisition of a GCP identity. The full repertoire of injury-induced signals that initiate

and govern adaptive reprogramming remains to be discovered.

In the adult brain, numerous cell types communicate and provide a concerted response to injury,

including astrocytes, microglia (macrophages of the brain) and stem cells of the neurogenic

niches (Frik et al., 2018). The timelines of the cellular responses of each cell type to injury - cell

death, activation of microglia, reactive gliosis, proliferation, scar formation and cellular remodeling

- have been delineated for specific adult brain injuries, particularly in the cerebral cortex. For

example, upon traumatic brain injury cells release damage-associated molecular patterns

(DAMPs), which act as an inflammatory stimulus and activate microglia that can lead to gliosis,

eventually causing neurotoxicity and scarring (Donat et al., 2017). However, the cellular

composition and microenvironment of the early postnatal brain are very different from the adult.

In the neonatal cerebellum, microglia are immature (Li et al., 2019) and are still being generated,

and NEPs and GCPs are actively proliferating and producing astroglia and neurons. Therefore,

the existing knowledge on how adult brain cells react to injury might not apply to the neonatal

cerebellum. For example, in the spinal cord, while neonatal microglia and astrocytes facilitate

scarless repair, the same cells in the adult promote scarring upon spinal cord injury in mice (Li et

al., 2020). It is thus important to study the microenvironment of the neonatal brain during repair

to determine what factors promote or inhibit regeneration.

Dying cells release many factors, including reactive oxygen species (ROS) that activate signaling

cascades in neighboring cells. However, little is known about how these signals regulate brain

repair, especially during development. The level of ROS during homeostasis is regulated by
metabolic processes, and typically is increased following injury (Niethammer, 2016). Furthermore,
ROS can directly react with proteins that regulate proliferation, viability, quiescence or
differentiation and metabolism (Bigarella et al., 2014, Tan and Suda, 2018). Thus, ROS are
considered key signaling molecules that participate in the crosstalk between progenitor cell fate
decisions and metabolic switches in a context- and cell type-dependent manner (Bigarella et al.,
2014). One significant mechanism by which ROS signaling is implicated during inflammatory
responses following an injury is through the activation of microglia, which in turn can lead to more
ROS production (Smith et al., 2022). This process is critical as it can potentially promote repair.
However, the role of ROS signaling during adaptive reprogramming of NEPs following neonatal
cerebellar injury is not known.
127
Here we first delineate the sequential changes in the microenvironment upon injury (focused
irradiation) to the mouse hindbrain at postnatal day 1 (P1). We then demonstrate a requirement
for a transient increase in ROS levels at ~24 hours (hr) post injury for cerebellar regeneration.
Single cell RNA-sequencing (scRNA-seq) and bulk ATAC-seq analyses revealed increased ROS
signaling compared to controls that peaks 24 hr after injury in NEPs, particularly those in the BgL,
demonstrating that ROS is an acute signal associated with the NEP response to GCP death. A
functional role of ROS signaling was established using a transgene (*mCAT*) that expresses the
human mitochondrial Catalase which can reduce ROS levels broadly. Several key steps in
adaptive reprogramming were abrogated in *mCAT*/+ mice leading to reduced replenishment of
the EGL and a smaller adult cerebellum. Finally, we show that the density of microglia is reduced
at P5 in irradiated *mCAT* mice compared to controls and provide evidence that microglia play a
role in the step of replenishing the EGL with BgL-derived GCPs during adaptive reprogramming.
140

Materials and Methods

Animals

143 All the mouse experiments were performed according to protocols approved by the Institutional
144 Animal Care and Use Committee of Memorial Sloan Kettering Cancer Center (MSKCC) (protocol
145 no. 07-01-001). Animals were housed on a 12-hour light/dark cycle and given access to food and
146 water ad libitum.
147
Two mouse lines were used in this study: *Nes-Cfp* (JAX #034387) (Encinas et al., 2006) and
mCAT (JAX #016197) (Schriner et al., 2005). Animals were maintained on an outbred Swiss

Webster background. Both sexes were used for analyses except for the genomics experiments
(scRNA-seq and ATAC-seq) where males were used.

EdU administration

5-ethynyl-2'-deoxyuridine (EdU) stock was dissolved in sterile phosphate-buffered saline (PBS)
at 10 mg/mL and a dose of 5 μg/g was intraperitoneally injected into animals 1 hour prior to
euthanasia.

PLX5622 administration

PLX5622 powder was provided by Plexxikon under a Materials Transfer Agreement. PLX5622
powder was first diluted in DMSO at 20mM and then diluted 1X in PBS just before intraperitoneal
injection into newborn pups. Injections were given every day from P1 to P8 at a dose of 10µg/g
of body weight. Control pups were injected with PBS-DMSO vehicle control.

Irradiation

P1 mice were anesthetized by hypothermia and given a single dose of ~5Gy γ-irradiation in an X-
RAD 225Cx (Precision X-ray) Microirradiator in the MSKCC Small-Animal Imaging Core facility.
A 5-mm diameter collimator was used to target the hindbrain from the left side of the animal.

Tissue preparation and histology
For immunocytochemistry, animals younger than P8 were sacrificed and then brains were
dissected, fixed in 4% paraformaldehyde for 24 hr at 4°C, cryoprotected in 30% sucrose in
phosphate-buffered saline (PBS) until they sank and then frozen in Cryo-OCT (Tissue-Tek). Older
animals were anesthetized and then perfused with cold PBS followed by 4% paraformaldehyde
prior to brain dissection. Frozen brains were cryosectioned sagittally at 14 μm and slides stored
at -20°C. Midline cerebellar sections were used for quantification in all downstream analyses.

For immunofluorescence staining, slides were allowed to warm to room temperature (RT) and
washed 3 times in PBS. Then, tissues were blocked for one hour with blocking buffer (5% BSA
(w/v) in 1XPBS with 0.1% Triton-X) at RT. Primary antibodies diluted in the blocking buffer were
placed on slides for overnight incubation at 4°C. Slides were then washed in PBS with 0.1%
Triton-X and incubated with fluorophore-conjugated secondary antibodies diluted in the blocking
buffer for 1 hr at RT. Following washes in PBS with 0.1% Triton-X after the secondary antibody
incubation, nuclei were counterstained with Hoechst (1:3000) and the slides were mounted using

mounting media (Electron Microscopy Sciences). Primary antibodies used are described in Table

S1 and secondary antibodies were Alexa Fluor-conjugated secondary antibodies (1:1000).

EdU was detected using a Click-it EdU (Invitrogen, C10340) assay with Sulfo-Cyanine5 azide

(Lumiprobe Corporation, A3330).

For TUNEL staining, after primary antibody incubation and washes, sections were permeabilized

in PBS with 0.5% TritonX-100 for 10 minutes (min) and then pre-incubated in TdT buffer (30mM

Tris HCl, 140 mm sodium cacodylate and 1mM $CoCl_2$) for 15 min at RT. Slides were then

incubated for 1hr at 37°C in TUNEL reaction solution containing Terminal Transferase and

Digoxigenin-11dUTP (Roche). After the TUNEL reaction and washes, slides were incubated with

a secondary antibody solution which included a sheep anti-dixogenin-rhodamine (Roche) for the

visualization of TUNEL reaction.

Image acquisition and analysis

Images were collected with a DM6000 Leica microscope, a NanoZoomer Digital Pathology

microscope (Hamamatsu Photonics), or an LSM880 confocal microscope (Zeiss). Images were

processed using NDP.view2 software, ImageJ software (NIH, Bethesda, MA, USA) and

Photoshop (Adobe).

Cell dissociation for FACS and flow cytometry

Cerebella were dissected into ice-cold 1x Hank's Buffered Salt Solution (Gibco) and dissociated

using Accutase (Innovative Cell Technologies) at 37°C for 10-15 min. After dissociation, Accutase

was diluted with 3x excess volume of neural stem cell media (Neurobasal medium, supplemented

with N2, B27 (without vitamin A)), and nonessential amino acids (Life Technologies, Gibco) and

cells were filtered using a 40 μm mesh cell strainer. After 5 min of centrifugation in a chilled

centrifuge at 500g, the pellet was resuspended in neural stem cell media and strained using

strainer cap tubes (Falcon) for downstream experimentation. All centrifugation was performed at

4°C and cells were kept on ice when possible.

Flow cytometry analysis for ROS and mitochondria mass

For MitoSOX and MitoTracker analyses, cells were incubated for 30 min at 4°C with 5 μM of

MitoSOX and 100 μM of MitoTracker (Thermo Fisher Scientific) to assess mitochondrial

superoxide production and mitochondrial mass, respectively. Data were collected using a LSR

Fortessa flow cytometer (BD) and analyzed using FlowJo software. The gating for the MitoSOX
or MitoTracker high populations (top 90%) were performed as previously described (Clutton et
al., 2019)
221

Multiplexed sc-RNAseq
Sample preparation
2-4 *Nes-Cfp/+* cerebella/replicates from male control non-irradiated pups or pups that were
irradiated at P1 were collected at P1 (control only), P2, P3 and P5 and dissociated as described
above. All conditions were performed in 2 replicates for nonIR and IR conditions, except for P5.
Following dissociation, CFP+ cells were immediately sorted on a BD FACS Aria sorter (BD
Biosciences) using a 100 μm nozzle. 50,000 Nes-CFP+ cells from each sample were processed
for scRNA-seq. Cells were sorted into neural stem cell media.
230

Multiplexing, droplet preparation, sequencing and data processing
The scRNA-Seq of FACS-sorted Nes-CFP+ cell suspensions was performed on a Chromium
instrument (10X genomics) following the user guide manual for 3′ v3.1. In brief, 50,000 FACS-
sorted cells from each condition were multiplexed using CellPlex reagents (10x Genomics) as
described by the manufacturer's protocol. P3 nonIR replicate #2 did not yield sufficient cells after
multiplexing. The viability of cells was above 95%, as confirmed with 0.2% (w/v) Trypan Blue
staining and barcoded cells were pooled into a single sample in PBS containing 1% bovine serum
albumin (BSA) to a final concentration of 700–1,300 cells per μl. 2-3,000 cells were targeted for
each sample. Samples were multiplexed together on one lane of a 10X Chromium following
the 10x Genomics 3' CellPlex Multiplexing protocol and a total of ~30,000 cells were targeted for
droplet formation. Cells were captured in droplets. After the reverse transcription and cell
barcoding in droplets, emulsions were broken, and cDNA was purified using Dynabeads MyOne
SILANE followed by PCR amplification per the manufacturer's instruction. Final libraries were
sequenced on an Illumina NovaSeq S4 platform (R1 – 28 cycles, i7 – 8 cycles, R2 – 90 cycles)
by the MSKCC core facility.
246

Data Analysis
Single-cell RNA-seq fastq files were demultiplexed using sharp (https://github.com/hisplan/sharp)
and initially mapped to the mouse mm10 reference genome using Cell Ranger v6.0.1 (Zheng et
al., 2017). The Cell Ranger BAM files for individual samples were then converted back to fastq
files using *bamtofastq* (Cell Ranger v7.0.1), with --reads-per-fastq=1000000000. Reads were then

mapped to the GRCm39 mouse reference with Gencode annotations (vM30) using STARsolo

v2.7.10a (--soloFeatures Gene Velocyto, --soloType CB_UMI_Simple, --soloCBwhitelist 3M-

february-2018.txt, --soloUMIlen 12, --soloCellFilter EmptyDrops_CR, --soloMultiMappers EM)

(Kaminow et al., 2021, Frankish et al., 2021).

The STARsolo output was used to create Seurat (v4.3) objects for each sample with spliced and

unspliced read count matrices (Hao et al., 2021). The objects were integrated, by running

NormalizeData and FindVariableFeatures (selection.method = "vst", nfeatures = 3000) on each

one, then SelectIntegrationFeatures and FindIntegrationAnchors on the list of objects and finally

IntegrateData with the anchors. Counts were then normalized with SCTransform. Based on

manual analysis, cells were filtered out where number of detected genes was ≤ 1,500, number of

detected transcripts was ≥ 40,000 and mitochondrial gene percentage ≥ 5%. To determine cell

cycle phases, the Kowalczyk et al. (Kowalczyk et al., 2015) cell cycle markers were used,

assuming the gene names, capitalized to the title case, to be orthologous between mouse and

human with the CellCycleScoring function. SCTransform was used to regress out the "Cell Cycle

difference" score (S score – G2M score). Dimension reduction was performed using RunPCA and

RunUMAP (dims = 1:20, n.neighbors = 30). For clustering, FindNeighbors was run using the first

20 principal components, then FindClusters with the Leiden algorithm with the default settings

(Traag et al., 2019). Clusters were annotated using known markers: GCPs

(*Atoh1+/Barhl1+/Tubb3+*), BgL-NEPs (*Hopx+*), ependymal cells (*Foxj1+*), immature interneurons

(*Pax2+*), Neurogenic NEPs (*Ascl1+*), astrocytes (*Slc6a11+*), oligodendrocytes (*Olig1+*),

meninges (*Col3a1+/Vtn+/Dcn+*), microglia (*Ly86+/Fcer1g+*).

The differential gene expression analyses were performed individually on *Hopx+* gliogenic-NEPs,

Ascl1+ neurogenic-NEPs, and GCPs following the same computational workflow. First, clusters

containing *Hopx*-NEPs (clusters 2, 3, 6, 10), *Ascl1*-NEPs (clusters 5, 8, 11), or GCPs (clusters 1,

4, 7, 12, 14) were subsetted from the original data set based on the expression of *Hopx*, *Ascl1*,

and *Atoh1*, *Barhl1* and *Rbfox*, respectively. Second, the subsetted cells were divided according

to whether the cells were from P2 or P3+P5 and split based on their biological replicate. The split

data sets were normalized using NormalizeData with default parameters, and the 3,000 top

variable features were computed using FindVariableFeatures with default settings. Re-integration

of the data sets was subsequently performed using SelectIntegrationFeatures,

FindIntegrationAnchors, and IntegrateData all with default parameters as previously described,

except for IntegrateData in the *Ascl1+* neurogenic-NEP analyses where k.weight = 50 was used.

Following re-integration, SCTransform with default parameters was used to normalize
mitochondrial read percentage, cell cycle difference score, number of features, and number of
counts. Dimension reduction was thereafter performed using RunPCA with default parameters
and RunUMAP with default parameters except dims = 1:40, repulsion.strength = 0.1, and min.dist
= 0.5. Clustering was subsequently performed using FindNeighbors with default settings except
dim = 30 and FindClusters with default settings for resolutions between 0.1 and 3 using the
original Louvain algorithm. A final resolution which gave a high silhouette score with a relatively
low negative silhouette proportion was selected for individual data sets. To allow downstream
DESeq2 analyses, count matrices were constructed by aggregating counts from cells from the
same treatment condition and biological replicate using AggregateExpression. The aggregated
count matrices were converted into a DESeq2 dataset object using DESeqDataSetFromMatrix,
grouping the samples by treatment conditions. Genes with fewer than 10 counts were filtered out.
DESeq2 (v1.36.0) was used to perform the differential expression analyses using a negative
binomial distribution and default settings (Love et al., 2014). The results were visualized using
EnhancedVolcano with a fold change cut-off of ±0.5 and an adjusted p-value threshold of 0.05.
301
302 The GO term analyses were performed on differentially expressed genes from the DESeq2 results
303 filtered with a log2 fold change threshold of ±0.5 and an adjusted p-value threshold of 0.05 for
304 each comparison. The probability weight function was computed using nullp with default
305 parameters and the mm8 mouse genome. The category enrichment testing was performed using
306 goseq with default parameters from the goseq package (v1.48.0).
307
Bulk ATAC-seq
Sample preparation
FACS-sorted Nes-CFP+ cells (30,000-50,000 per replicate) were isolated from control or
irradiated (at P1) P2 cerebella. 2-3 cerebella were pooled for each sample. Cells were
immediately processed for nuclei preparation and transposition using the OMNI-ATAC protocol
(Corces et al., 2017). Sequencing was performed at the MSKCC genomics core facility using the
Illumina NovaSeq S4 platform.
315
Data Processing and Analysis
Raw sequencing reads were 3' trimmed and filtered for quality and adapter content using version
0.4.5 of TrimGalore (https://www.bioinformatics.babraham.ac.uk/projects/trim_galore), with a
quality setting of 15, and running version 1.15 of cutadapt and version 0.11.5 of FastQC. Version

2.3.4.1 of bowtie2 (http://bowtie-bio.sourceforge.net/bowtie2/index.shtml) was used to align reads

to mouse assembly mm10 and alignments were deduplicated using MarkDuplicates in version

2.16.0 of Picard Tools. Enriched regions were discovered using MACS2

(https://github.com/taoliu/MACS) with a p-value setting of 0.001, filtered for blacklisted regions

(http://mitra.stanford.edu/kundaje/akundaje/release/blacklists/   mm10-

mouse/mm10.blacklist.bed.gz), and a peak atlas was created using +/- 250 bp around peak

summits. The BEDTools suite (http://bedtools.readthedocs.io) was used to create normalized

bigwig files. Version 1.6.1 of featureCounts (http://subread.sourceforge.net) was used to build a

raw counts matrix and DESeq2 was used to calculate differential enrichment for all pairwise

contrasts. Peak-gene associations were created by assigning all intragenic peaks to that gene,

while intergenic peaks were assigned using linear genomic distance to transcription start site.

Network analysis was performed using the assigned genes to differential peaks by running

enrichplot::cnetplot in R with default parameters. Composite and tornado plots were created using

deepTools v3.3.0 by running computeMatrix and plotHeatmap on normalized bigwigs with

average signal sampled in 25 bp windows and flanking region defined by the surrounding 2 kb.

Motif signatures were obtained using Homer v4.5 (http://homer.ucsd.edu) on differentially

enriched peak *regions.*

Data and code availability

The scRNA-seq data was submitted to ArrayExpress (Accession E-MTAB-13353). Bulk ATAC-

seq data has been submitted to GEO (GSE269342). The code used to carry out the scRNA-seq

analysis can be found on GitHub repository: https://github.com/BayinLab/Pakula_et_al_23

Quantification and statistical analysis

For detecting TUNEL, IBA1, GFAP, CFP and SOX2, images were captured using a DM600 Leica

fluorescent microscope and subsequently quantified on ImageJ Software (NIH). Measurements

were conducted on lobules 3, 4 and 5 of midsagittal sections unless indicated in Figure legends.

Positive cells were counted and densities were calculated based on BgL length, EGL area, WM

area or on the whole cerebellum without the EGL (outside EGL) as indicated in the figures. For

the cerebellar section area, images were acquired using a Nanozoomer2.0 HT slide scanner

(Hamamatsu). Midsagittal sections were selected and exported for manual analysis using ImageJ

software. For EGL thickness, images of lobules 3, 4 and 5 from midsagittal sections were obtained

using a DM600 Leica fluorescent microscope and analyzed on ImageJ. EGL thickness was

calculated as the EGL area divided by the EGL perimeter.

Prism (GraphPad) was used for all statistical analyses. Statistical tests performed in this study

were Welch's two-tailed t-test and Two-way analysis of variance (ANOVA) followed by post hoc

analysis with Tukey's multiple comparison tests. A p-value ≤ 0.05 was considered as significant.

Results are presented as the mean ± SEM. At least three biological samples and 2 to 3 sections

per sample were analyzed for each experiment to ensure reproducibility and the sample sizes are

reported in the Results section for significant data. For qualitative analysis, midsagittal sections

from at least 4 samples were observed.

362
363   **Results**
364

scRNA-seq of NEPs during adaptive reprogramming reveals increased cellular stress, ROS

signaling and DNA damage

To investigate the molecular changes that NEP subpopulations undergo upon injury to the EGL,

in particular an increase in the signaling pathways associated with injury induced cellular stress

and ROS, we performed multiplexed scRNA-seq of CFP+ cells isolated by fluorescence-activated

cell sorting (FACS) of cerebella from *Nes-Cfp/+* transgenic neonates either irradiated at P1 (IR;

P2, P3, P5) or non-irradiated (nonIR; P1, P2, P3, P5) (Figure 1A, Supplementary Figure 1A, B).

Following the filtering out of poor quality cells and integration of replicates and the two conditions,

the clustering of 11,878 cells (6,978 nonIR and 4,900 IR) was performed (Hao et al., 2021). The

analysis revealed the expected three distinct groups of cells: gliogenic-NEPs and astrocytes

(*Hopx*+ clusters 2, 3, 6, 10), neurogenic-NEPs (*Ascl1*+ clusters 5, 8, 11) and GCP (*Atoh1*+

clusters 1, 4, 7, 12, 14) that were present at each stage and in both conditions (Figure 1B-E,

Supplementary Figure 1C, Table S2). These groups of cells were further subdivided into

molecularly distinct clusters based on their cell cycle profiles or developmental stages (Table S2).

In addition, oligodendrocyte progenitors (cluster 15), microglia (clusters 17, 21), ependymal cells

(clusters 13, 18, 19) and meninges (cluster 16) were detected (Figure 1B, D, Table S2). Cluster

20 represented low-quality cells and was omitted from downstream analyses. As expected, the

GCP clusters were enriched in the cells from irradiated mice and at P5 (Supplementary Figure

1C).

Our further analyses focused on changes in the signaling pathways associated with injury induced

cellular stress and ROS. We previously showed that a subset of the *Hopx*+ gliogenic-NEPs that

are present in the BgL are the ones that undergo adaptive reprograming following GCP death

(Wojcinski et al., 2017, Bayin N. S., 2021). We therefore assessed the immediate and later effects

of GCP death on *Hopx+* gliogenic-NEPs by performing differential expression analyses between

nonIR and IR gliogenic-NEPs (*Hopx+,* clusters 2, 3, 6, 10) at P2, or at P3 and P5 (P3+5). 24 hr

after injury at P1, 132 genes in gliogenic-NEP clusters were significantly upregulated in IR

compared to 34 genes that were upregulated in nonIR P2 cells (adjusted p-value≤0.05, Figure

1F, Table S3). The significantly increased genes included *Cdkn1a, Phlda3, Ass1* and *Bax,* all of

which have been implicated as increased in response to DNA damage and in ROS signaling, as

well as in anti-apoptotic functions (Figure 1G) (Masgras et al., 2012, Bensellam et al., 2019, Qiu

et al., 2014, Jiang et al., 2008). Indeed, some of the top gene ontology (GO) terms associated

with the genes upregulated in gliogenic-NEPs with injury were related to response to irradiation,

DNA damage, the P53 pathway and ROS metabolic processes, whereas the top GO terms

associated with the genes upregulated in the nonIR cells at P2 were related to metabolic

processes (p-value≤0.05, Figure 1H, Table S4). Interestingly, the transcriptional changes

observed at P2 were less pronounced at later time points, where although some of the top

differentially expressed genes at P2 were still significantly upregulated at P3+5 stages combined

(e.g. *Cdkn1a, Phlda3, Ass1, Bax*), the increase in expression levels of these genes upon injury

and/or the number of cells expressing them gradually declined after P2 in IR gliogenic-NEPs when

compared to their nonIR counterparts (Figure 1G). Genes upregulated in IR P3+5 gliogenic NEPs

were associated with similar GO terms to those at P2, such as response to irradiation and the

P53 pathway, however, "ROS metabolic processes" was no longer a significantly enriched GO

term (p-value≤0.05, Supplementary Figure 2A, F, Table S3-4).

To assess whether the injury induced transcriptional signatures are specific to gliogenic-NEPs,

we performed the same differential expression analysis on nonIR and IR neurogenic-NEPs

(*Ascl1+,* clusters 5, 8, 11) and GCPs (*Atoh1+,* clusters 1,4,7,12,14) at P2, or at P3+P5 to identify

the immediate and later changes upon injury at birth. Similar to the gliogenic-NEPs, P2 IR

neurogenic-NEPs showed significant upregulation of genes associated with GO terms related to

stress response and apoptosis following injury, although the ROS related GO term was not

prominent (Supplementary Figure 2B, G, Table S3-4). Interestingly, although IR neurogenic-

NEPs at P3+5 had only 27 genes that were significantly upregulated following injury (adjusted p-

value≤0.05, Table S3, Supplementary Figure 2C), the genes included ones associated with

neural stem cells and BgL-NEPs (e.g. *Id1, Apoe, Ednrb*). The latter genes could represent the

transitory *Ascl1+* BgL-NEP population that induces *Ascl1* expression during adaptive

reprogramming and would be included in the *Ascl1+* neurogenic clusters or reflect the delayed

neurogenesis of neurogenic-NEPs previously demonstrated (Bayin N. S., 2021). Consistent with

the latter, the nonIR P3+5 neurogenic-NEPs showed an increase in mature neuron markers

compared to IR cells (adjusted p-value<0.05, Supplementary Figure 2C, H, Tables S3-4).

In contrast to the gliogenic- and neurogenic-NEP subtypes, P2 IR GCPs showed upregulation of

genes enriched in GO terms such as neural differentiation, axonogenesis and nonIR P2 GCPs

showed increased expression of genes involved in cell cycle and mitosis (Supplementary Figure

2D, I). This result could reflect the death of highly proliferative GCPs after irradiation and sparing

of only postmitotic granule cells upon irradiation at P1. P3+5 IR GCPs showed increased

expression of genes associated with BgL-NEPs (*Id1* and *Gdf10*, adjusted p-value ≤ 0.05) and

genes associated with GO terms such as cell cycle and cell fate commitment whereas P3+5 nonIR

GCPs showed enrichment for GO terms related to cell migration and neurogenesis (adjusted p-

value ≤0.05, Supplementary Figure 2D, E, I, J, Tables S3-4). This result could reflect the delayed

neurogenesis of GCPs following injury. Interestingly, we did not observe significant enrichment

for GO terms associated with cellular stress response in the GCPs that survived the irradiation

compared to controls (Table S4). Collectively, these results indicate that the gliogenic- and

neurogenic-NEP subtypes transiently upregulate stress response genes upon GCP death, but

only gliogenic-NEPs have a strong upregulation of ROS signaling.

Injury induces changes in NEP chromatin landscape at P2

We next tested whether GCP death at birth induces changes to the chromatin landscape of NEPs

that reflect the altered gene expression observed with scRNA-seq, by performing bulk ATAC-seq

on FACS-isolated CFP+ cells from P2 control and injured *Nes-Cfp/+* pup cerebella (Figure 1A).

P2 was chosen as it is the stage when GCPs contribute the least to the total Nes-CFP+ FACS

population and to identify the immediate effects of the injury on NEP chromatin. Analysis of

differentially open chromatin showed that injury induces major changes to the chromatin

landscape of the NEPs (1168 differentially open regions, adjusted p-value<0.05, Figure 1I. Table

S5). Of interest, *Cdkn1a* and *Phlda3*, two genes stimulated by ROS and injury (Bensellam et al.,

2019, Masgras et al., 2012) and that were upregulated in gliogenic-NEPs after irradiation (Figure

1F-G, Supplementary Figure 2B) exhibited new accessible regions around their gene bodies

compared to the nonIR P2 Nes-CFP+ cells (Figure 1J). Known motif analysis in the regions with

increased accessibility upon injury showed enrichment for binding motifs of the JUN/AP1

transcriptional complex (p-value = 10^{-14}, % of target sequences with motif = 15%, Table S6) which

is known to act in response to cellular stress and be activated by ROS. In addition, the DNA

binding site for FOXO3, a transcription factor that regulates ROS levels, had increased chromatin

accessibility (p-value = 0.001, % of target sequences with motif = 22.41%) (Table S6) (Filosto et

al., 2003, Auten and Davis, 2009, Hagenbuchner et al., 2012). Finally, gene network analysis of

ATAC-seq data revealed an active transcriptional network involved in regulating cell death and

apoptosis (Figure 1K). Furthermore, some of the genes involved in this response, such as *Ppara*,

Egln3, *Foxo3*, *Jun* and *Nos1ap,* have been implicated as upregulated with increased ROS levels

or involved in ROS signaling (Devchand et al., 2004, Kaelin, 2005, Hagenbuchner et al., 2012,

Filosto et al., 2003). In summary, our ATAC-seq data analysis along with the scRNA-seq provide

strong evidence that irradiation causes increased ROS signaling predominantly in the BgL-NEPs

upon GCP death by inducing transcriptional and epigenetic changes within 1 day after injury (P2).

In addition, genes related to cell survival and death, cellular stress and DNA damage are

upregulated in NEPs shortly upon injury, possibly as a means to overcome the cellular effects of

injury and induce adaptive reprogramming of NEPs to replenish the lost cells.

Transient increase in cerebellar ROS during apoptosis of granule cell precursors

To validate that the transient increase in expression of genes associated with cellular stress and

ROS signaling in NEPs is due to an increase in ROS upon cerebellar injury, we quantified ROS

levels in whole cerebellum samples using a mitochondrial superoxide indicator (MitoSOX) via flow

cytometry. A significant increase in ROS (cells present in the top 90% MitoSOX+ intensity) was

observed specifically at P2 (p=0.0005, n≥10) and not at P3 or P5 in IR pups compared to nonIR

(Figure 2A, B and Supplementary Figure 3A, B). Furthermore, quantification of mitochondrial

mass with MitoTracker revealed a reduction only at P2 (p=0.0005, n≥10) in IR pups (Figure 2C

and Supplementary Figure 3C-E). Additionally, quantification of TUNEL staining in midline sagittal

sections of the cerebella showed a large increase in cell death in the EGL of injured cerebella at

P2 (p=0.0040, n=4), but not at P3 compared to the controls (Figure 2D, E; see also Figure 3K, L).

Outside the EGL there was a small increase in cell death after injury that was only significant at

P3 (p=0.0379, n≥3) (Figure 2F, G). Thus, a transient increase in ROS in the cerebellum one day

after irradiation at P1 correlates with the timing of the major death of GCPs.

Altered glial microenvironment following death of granule cell precursors

Given the potential importance of glial cells to regenerative cellular responses after injury, we next

asked whether the glial microenvironment of the cerebellum changes during early postnatal

development in response to irradiation at P1. We first analyzed the astrocyte marker GFAP, since
it is generally upregulated in soon after brain injury (Burda and Sofroniew, 2014). Most astrocytes,
including the specialized Bg, express GFAP in the adult cerebellum, but the cells are generated
by gliogenic-NEPs during the first two weeks after birth. We therefore determined the normal
location and timing of initiation of GFAP expression in these cells during postnatal cerebellum
development. GFAP was first detected in rare astrocytes at P2 located in the white matter (WM)
below the lobules, with strong expression in all WM astrocytes at P5 and later stages (P8 and
P12) (Figure 3A-D, observed in n=4 mice/stage). In contrast, GFAP expression in astrocytes in
the developing IGL and in the glial processes of Bg that project through the molecular layer (ML)
and EGL was not detectable until P5 and was much stronger at P8 and P12 (Figure 3A-D).
Interestingly, after injury at birth, GFAP expression was prematurely upregulated in the WM
astrocytes below the lobules at P2 and in the remaining astrocytes at P5, including in the fibers
of Bg that extend through the ML and EGL, compared to nonIR cerebella (Figure 3A-H, observed
in n=4 mice/stage). GFAP expression was similar in all glia in both conditions at P8 and P12.
These results reveal that astrocytes and Bg react to EGL injury caused by irradiation at P1 by
initiating GFAP expression earlier than normal in the deep WM, and then the IGL and in Bg.
508
The macrophages of the brain, microglia, are generated in the early embryo but their main
increase in cell number occurs during neonatal development in mice (Hammond et al., 2018). We
found that most microglia were located in the WM of the cerebellum, and that the density of IBA1+
microglia increased in the WM between P2 (235.1 ± 18.3 cells/mm^2) and P3 (344.3 ± 53.1
cells/mm^2) and then was maintained at P5 (350.4 ± 40.2 cells/mm^2) (Fig. 3I-K, Supplementary
Figure 3F). Since the area of the cerebellum is increasing between P2 and P5, active microglia
production and/or infiltration must continue to occur between P2 and P5. As expected, after
irradiation the density of microglia in the injured EGL was significantly increased (~3 fold) at P2
but not at P3 or P5 compared to controls (p=0.0331, n=4) (Fig. 3L-N, Supplementary Figure 3G).
No difference in the microglial density in the WM was detected between conditions at both time
points (Fig. 3I-L; Supplementary Figure 3F). Thus, as expected microglia density was increased
in the EGL one day after injury when the maximum GCP cell death occurs.
521

522   **Decreasing ROS impairs cerebellar repair**
523
Given the transient increase in ROS signaling in gliogenic-NEPs during peak GCP death and
recruitment of microglia to the EGL, and the later astroglial response, we tested whether an
increase in ROS is necessary for adaptive reprogramming and cerebellar repair following EGL

injury at P1. To reduce the level of ROS we utilized an *mCAT* transgenic mouse line that
expresses the human mitochondrial catalase ubiquitously from a *CMV* promoter (Schriner et al.,
2005). The transgene is expected to reduce mitochondrial ROS levels in all cells by catalyzing
the breakdown of hydrogen peroxide into water and oxygen, hence protecting the cells from
oxidative damage. We confirmed that mCAT protein is expressed throughout the cerebellum
using immunohistochemical (IHC) staining of cerebellar sections (Figure 4A, B). MitoSOX flow
cytometry revealed a significant decrease in ROS in the cerebella of *mCAT/+* mice 1 day after
irradiation (P2) (p=0.0163, n≥8) and not at P3 or P5 compared to control IR mice and no baseline
decrease in ROS at any stage in nonIR mice (Figure 4C and Supplementary Figure 4A, B).
Mitochondrial mass, as measured by MitoTracker flow cytometry, was reduced at P2 in *mCAT/+*
IR compared to nonIR *mCAT/+* pups with no significant difference observed between *mCAT/+*
and control IR mice (Fig. 4D, Supplementary Figure 4C, D). The level of cell death in the EGL
and density of IBA1+ microglia in the EGL and elsewhere in the cerebellum were similar between
mCAT/+ and littermate control IR mice at P2 and P3 (Fig. 4E, F, Supplementary Figure 4G, J).
There was a slight increase in cell death outside the EGL at P2 in IR *mCAT/+* compared to nonIR
mCAT/+ mice but not compared to IR controls (Supplementary Figure 4F, E). Thus, the *mCAT*
transgene counteracts the transient increase in cerebellar ROS following EGL injury but does
have a major effect on GCP death or the infiltration of microglia to the injured EGL (Supplementary
Figure 4 F-J).
546
We next determined the regenerative efficiency of *mCAT/+* mice by analyzing the area of sections
of cerebella from nonIR and IR P30 *mCAT/+* mice compared to littermate controls. Strikingly, the
cross-sectional area of the medial cerebellum (vermis) of IR *mCAT/+* adult mice was significantly
reduced compared to IR controls (p=0.0040, n≥6) (Fig. 4G-K). Analysis of cerebellar area across
ages (P3, 5, 8 and 12) revealed that the vermis sectional area of the IR *mCAT/+* cerebella was
only significantly reduced at P30 compared to IR controls, however at P12 it was reduced in
mCAT/+ IR cerebella compared to *mCAT/+* nonIR mice, whereas it was not significantly different
between control IR and control nonIR mice (Fig. 4L, Supplementary Figure 4 K-N). These results
indicate that cerebellar growth begins to be reduced at P12 in *mCAT/+* mice following injury. Thus,
a reduction in ROS at the time of cell death in the EGL leads to a diminution of cerebellar recovery.
557
558   **Reduced regeneration in *mCAT* mice is associated with reduced adaptive reprogramming**
559   **at P5.**
560

Given that a decrease in ROS following EGL injury reduces regeneration of the neonatal
cerebellum, we determined whether specific stages of the adaptive reprogramming process are
altered in *mCAT*/+ mice compared to controls. First, we analyzed the replenishment of the EGL
by BgL-NEPs in vermis lobules 3-5. Interestingly, we found that although the thickness of the EGL
in IR mice of both genotypes was similarly reduced compared to nonIR mice at P5 (p=0.0041
control and p=0.0005 *mCAT*/+; n≥4) by P8 the control EGL was a similar thickness to the control
nonIR whereas the thickness of the *mCAT*/+ IR EGL was significantly reduced compared to
mCAT/+ nonIR mice (p=0.035, n=4)(Figure 5A, B). A key regenerative process that contributes
to the expansion of the EGL following injury is the migration of BgL-NEPs to the EGL. Strikingly,
the density of CFP+ cells in the EGL (mainly BgL-derived NEPs) was significantly decreased in
mCAT/+ IR mice compared to control IR mice at P5 (p=0.0002, n=4) but not P3 (Figure 5C-G,
Supplementary Figure 5A). Furthermore, the density of NEPs (CFP+ or SOX2+ cells) in the BgL
was significantly decreased at P5 in *mCAT*/+ IR cerebella compared to controls (p=0.0010, n≥5)
but not at other stages (Figure 5H, Supplementary Figure 5B-D). These results indicate that BgL-
NEPs have a blunted response to EGL injury, and therefore do not fully expand and contribute to
the replenishment of GCPs in the EGL after irradiation.

577
Microglia contribute to one aspect of adaptive reprogramming.
Given that several steps in adaptive reprogramming were decreased specifically at P5 in *mCAT*/+
IR cerebella, we asked whether microglia/macrophages could be involved in any of the processes.
We first determine the density of IBA1+ cells in the EGL and WM of vermis lobules 3-5 at P5 in
nonIR and IR mice of both genotypes. As expected, the density of microglia in the EGL was very
low in the nonIR control and *mCAT/*+ cerebella (Figure 6A-E). Interestingly, whereas control IR
mice had a similar number of IBA1+ cells in the WM as nonIR mice of both genotypes at P5, the
mCAT/+ IR mice had a lower density of microglia/macrophages in the WM compared to control
IR mice at P5 (p=0.0012, n≥3) (Figure 6A-D, F). This result raised the question of whether
macrophages/microglia play a role in adaptive reprogramming. We, therefore, tested whether
reducing the density of IBA1+ microglia/macrophages after birth would alter adaptive
reprogramming at P5 or cerebellar regeneration at later stages. Since macrophages and
cerebellar microglia are dependent on colony stimulating factor 1 (CSF1) for their survival, we
administered PLX5622, a small molecule inhibitor of CSF receptor 1 (CSFR1), to pups every day
from P0-5 (PLX treatment) (Kana et al., 2019, Tan et al., 2021). As expected, IBA1+ cells were
significantly decreased in the cerebellum of PLX-treated mice at P5 compared to their controls,
both nonIR and IR (p=0.0015 and p=0.0059, n=3 and n=5, respectively) (Figure 6G,

Supplementary Figure 6B-E). The thickness of the EGL was not significantly altered at P5 in PLX-
treated IR mice compared to IR controls (Figure 6H). Interestingly, similar to *mCAT/+* mice, the
density of Nes-CFP+ cells in the EGL was significantly decreased in PLX-treated IR mice
compared to IR controls at P5 (p=0.0008, n=5) (Figure 6I-M). In contrast, the density of SOX2+
cells in the BgL, corresponding to the gliogenic BgL-NEPs, were unchanged in the PLX-treated
and control mice, whether irradiated or not, suggesting that the decrease in expansion of BgL-
NEPs caused by ROS is not mediated by microglia (Supplementary Figure 6F). When mice were
treated with PLX from P0-8 and allowed to age to P30, we found the cerebellar vermis section
area was not decreased in PLX-treated IR mice compared to IR controls (Figure 6N). Thus,
reducing the density of IBA1+ microglia/macrophages in neonatal mice reduces the recruitment
of Nes-CFP+ cells to the EGL at P5, but does not have a long-term significant impact on
regeneration of the cerebellum.
607

Discussion
We demonstrate that a transient increase in ROS signaling after cerebellar injury to the EGL is
critical for adaptive reprogramming and full recovery of cerebellar growth. ROS likely acts as an
alarm signal shortly after injury and is primarily detected by BgL-NEPs. scRNA-seq at P1-5 and
bulk ATAC-seq at P2 of NEPs following targeted irradiation at P1 revealed a rapid increase in
transcriptional and epigenetic changes associated with upregulation of ROS and stress related
pathways in NEPs that peaked at P2. A transient upregulation in ROS at P2 was confirmed using
flow cytometry and found to correlate with the timing of cell death in the EGL one day after
irradiation. By reducing mitochondrial ROS levels across all cell types at P2 using a *mCAT*
transgene, we uncovered that ROS is required for several steps of adaptive reprogramming of
BgL-NEPs. In addition, we found that microglia are reduced in injured *mCAT/+* pups, which is
consistent with prior evidence that ROS can trigger immune cell recruitment in other systems (Kim
et al., 2010, Mehl et al., 2022). Moreover, temporary depletion of microglia caused a reduction in
the number of NEPs that migrate into the EGL at P5 following injury at P1, but no long-term
reduction of cerebellar size. Thus, we identified key transcriptomic and epigenomic changes in
cerebellar NEPs upon GCP ablation at birth and discovered roles for the tissue microenvironment,
especially ROS and a more limited role of microglia during neonatal cerebellum regeneration.
625

scRNA-seq analysis of NEPs from nonIR (P1-3, P5) and nonIR (P2, P3, P5) showed an increase
in genes associated with stress responses after irradiation in both the gliogenic and neurogenic
subpopulations, but not in the GCPs. Furthermore, a predominant increase in ROS signaling was

detected only in the gliogenic-NEPs one day after injury (P2). The injury-induced ROS and stress-
related gene signatures were not observed in the later P3+5 gliogenic- and neurogenic-NEPs,
suggesting that the increase in ROS levels is an early injury induced signal affecting the NEP
transcriptome. Our bulk ATAC-seq data generated at P2 revealed that the open chromatin regions
in the IR NEPs were enriched for transcription factor binding motifs related to ROS signaling and
stress induced transcription factors such as FOXO3 and AP1 (Table S6). Thus, cerebellar injury
during development induces transcriptional and epigenomic signatures in the NEPs and ROS
signaling could be a key driver of NEP adaptive reprogramming. Our results are in line with other
regeneration systems where a temporary increase in ROS is observed upon cell death or injury
and is considered to be a DAMP (Niethammer, 2016).

Interestingly, although both gliogenic- and neurogenic-NEPs showed induction of cellular stress
related genes upon injury, upregulation of ROS signaling and related genes was only observed
in the *Hopx*-expressing gliogenic NEPs that undergo adaptive reprogramming. Whether the
upregulation of ROS signaling primarily in the BgL-NEPs but not in the *Ascl1+* neurogenic-NEPs
is due to BgL-NEPs being in proximity to the dying GCPs after injury or their direct contact due to
their radial projections remains to be determined. The ability of BgL-NEPs to respond to GCP
death via upregulating ROS signaling and impaired regeneration upon reduction of ROS levels,
shows that ROS signaling is involved in triggering adaptive reprogramming upon injury.

The cellular composition of the neonatal cerebellum is dramatically different from the adult. During
the early postnatal period, we found that astrocytes in the WM below the lobules are the first to
initiate GFAP expression at P2 and that by P5 all astrocytes express a high level of GFAP. In
contrast, Bg express a low level of GFAP at P5 and reach a high level by P8. Interestingly, we
found that injury leads to an increase in the level of GFAP expression in each type of astroglia
when they first initiate expression, deep WM astrocytes at P2 and Bg (and all astrocytes) at P5.
Once adaptive reprogramming is nearing completion (P8), the GFAP levels remain similar in all
astroglia in nonIR and IR cerebella. These results suggest that the neonatal astrocytes respond
to injury differently than in the adult where GFAP upregulation is observed immediately after injury
(Burda and Sofroniew, 2014), since Bg and astrocytes in the lobules have a delayed response to
injury with GFAP not being upregulated for several days. Furthermore, at birth the microglia have
not fully expanded in number in the cerebellum and previous scRNA-seq showed that the
neonatal and adult microglia are transcriptionally distinct (Hammond et al., 2019). Perhaps
neonatal microglia are anti-inflammatory and pro-regenerative upon injury in neonates, in contrast

to adult microglia where upon traumatic brain injury they can inhibit regeneration (Donat et al.,

2017). Collectively, the differences in glial responses to injury in the neonatal cerebellum and

adult brain likely contributes to the permissiveness of the neonatal cerebellum to regeneration.

Details of the molecular changes that the neonatal cerebellar microglia/macrophages and

astrocytes undergo upon GCP injury and their molecular crosstalk with NEPs remain to be

determined.

We demonstrated the significance of ROS activation in the NEP reprogramming process using

an *mCAT* transgenic mouse line in which human Catalase (CAT) is expressed in mitochondria, a

protein that can lead to a reduction in hydrogen peroxide (H_2O_2) and thus lower ROS. However,

in *mCAT* neonatal cerebella we found that the percentage of MitoSOX high cells are comparable

to control mice under nonIR conditions. Importantly, however, one day after irradiation at P1 when

the percentage of MitoSOX high cells increases in control IR mice, the *mCAT* transgene reduces

the percentage of MitoSOX high cells, such that it remains at the baseline nonIR *mCAT* level.

Therefore, human CAT expression in mitochondria in this model inhibits the injury-induced

increase in ROS levels without affecting the homeostatic production of superoxide. Of possible

relevance, in this mouse model the observed change in ROS levels is likely global, impacting all

cell types. The specific impact of increased mCAT in BgL-NEPs or microglia on their recruitment

and function after an injury remains to be determined with new cell type-specific tools.

Our experiments depleting microglia using PLX5622 indicate that microglia/macrophages are

involved in the regeneration of the EGL following irradiation. Previous studies have demonstrated

a dual role of microglia in promoting and inhibiting regenerative processes within the nervous

system (Lee et al., 2021, Wang et al., 2020). Our data support the idea that microglia are involved

in the adaptive reprogramming of NEPs to GCPs by promoting their replenishment of GCPs in

the EGL. While the direct mechanisms remain to be discovered, this finding highlights that the

neonatal microglia support regeneration upon injury to the newborn cerebellum. PLX-treatment

for 8 days after irradiation did not reduce the later growth of the injured cerebellum. One possibility

is that after the cessation of PLX administration, regeneration proceeds normally. Alternatively, it

raises the possibility that an additional regenerative process is required in parallel to microglial

signaling.

Collectively, we have delineated the spatiotemporal cellular changes in the cerebellar glial

microenvironment upon ablation of GCPs at birth and highlight ROS signaling as a key stimulator

of adaptive reprogramming of NEPs. How DAMP-glia-progenitor crosstalk is orchestrated remains

to be untangled. Understanding how microenvironmental responses shape repair processes is a

crucial first step towards developing strategies to promote regeneration.

Acknowledgements

We thank past and present members of the Joyner laboratory for discussions and technical help.

We would like to thank Dr Ronan Chaligne and his team for their support in the multiplexed

scRNA-seq experiments. We are grateful to the MSKCC Animal Imaging Core, Flow cytometry

Core, Center for Comprehensive Medicine and Pathology, Integrated Genomics Operation,

Single-cell Analytics and Innovation Laboratory and Epigenetics Computational Laboratory teams

for technical services and support. An XRad 225Cx Microirradiator was purchased by support

from a Shared Resources Grant from the MSKCC Geoffrey Beene Cancer Research Center. We

gratefully acknowledge the support of the Gurdon Institute Scientific Computing Facility.

This work was supported by grants from the NIH to ALJ (R01NS092096) and NSB NIH NINDS

K99 NS112605-01. Addition funding was provided to ALJ from an NCI Cancer Center Support

Grant (CCSG, P30 CA08748) and the Cycle for Survival, to SE from a Francois Wallace Monahan

Fellowship; to NSB from a Wellcome Career Development Award (227294/Z/23/Z), Royal Society

(RGS\R1\231143) and Cambridge Stem Cell Institute Seed Funding, and to JBC from a University

of Cambridge School of Biological Sciences DTP PhD Studentship and Peter and Emma

Thomsen's Scholarship (1051). Gurdon Institute is supported by Wellcome Core Grant (203144)

and CRUK Grant (C6946/A24843).

References:

References:

ALTMAN, J. & ANDERSON, W. J. 1971. Irradiation of the cerebellum in infant rats with low-level
x-ray: histological and cytological effects during infancy and adulthood. *Exp Neurol,* 30**,**
492-509.
ALTMAN, J. & BAYER, S. A. 1997. *Development of the cerebellar system in relation to its*
evolution, structure, and functions, Boca Raton, CRC Press.
AUTEN, R. L. & DAVIS, J. M. 2009. Oxygen Toxicity and Reactive Oxygen Species: The Devil
Is in the Details. *Pediatric Research,* 66**,** 121-127.
BADURA, A., VERPEUT, J. L., METZGER, J. W., PEREIRA, T. D., PISANO, T. J., DEVERETT,
B., BAKSHINSKAYA, D. E. & WANG, S. S. 2018. Normal cognitive and social
development require posterior cerebellar activity. *Elife,* 7.
BAYIN N. S., M. D., STEPHEN N. D., LAO Z., SIMS P. A., JOYNER A. L. 2021. Injury induced
ASCL1 expression orchestrates a transitory cell state required for repair of the neonatal
cerebellum. *Sci Adv,* 7**,** eabj1598.
BAYIN, N. S., WOJCINSKI, A., MOURTON, A., SAITO, H., SUZUKI, N. & JOYNER, A. L. 2018.
Age-dependent dormant resident progenitors are stimulated by injury to regenerate
Purkinje neurons. *Elife,* 7.
BENSELLAM, M., CHAN, J. Y., LEE, K., JOGLEKAR, M. V., HARDIKAR, A. A., LOUDOVARIS,
T., THOMAS, H. E., JONAS, J. C. & LAYBUTT, D. R. 2019. Phlda3 regulates beta cell
survival during stress. *Sci Rep,* 9**,** 12827.
BIGARELLA, C. L., LIANG, R. & GHAFFARI, S. 2014. Stem cells and the impact of ROS
signaling. *Development,* 141**,** 4206-18.
BROWN, A. M., WHITE, J. J., VAN DER HEIJDEN, M. E., ZHOU, J., LIN, T. & SILLITOE, R. V.
2020. Purkinje cell misfiring generates high-amplitude action tremors that are corrected
by cerebellar deep brain stimulation. *Elife,* 9.
BUCKNER, R. L. 2013. The cerebellum and cognitive function: 25 years of insight from anatomy
and neuroimaging. *Neuron,* 80**,** 807-15.
BURDA, J. E. & SOFRONIEW, M. V. 2014. Reactive gliosis and the multicellular response to
CNS damage and disease. *Neuron,* 81**,** 229-48.
CERRATO, V., PARMIGIANI, E., FIGUERES-ONATE, M., BETIZEAU, M., APRATO, J.,
NANAVATY, I., BERCHIALLA, P., LUZZATI, F., DE'SPERATI, C., LOPEZ-
MASCARAQUE, L. & BUFFO, A. 2018. Multiple origins and modularity in the
spatiotemporal emergence of cerebellar astrocyte heterogeneity. *PLoS Biol,* 16**,**
e2005513.
CLUTTON, G., MOLLAN, K., HUDGENS, M. & GOONETILLEKE, N. 2019. A Reproducible,
Objective Method Using MitoTracker® Fluorescent Dyes to Assess Mitochondrial Mass
in T Cells by Flow Cytometry. *Cytometry A,* 95**,** 450-456.
CORCES, M. R., TREVINO, A. E., HAMILTON, E. G., GREENSIDE, P. G., SINNOTT-
ARMSTRONG, N. A., VESUNA, S., SATPATHY, A. T., RUBIN, A. J., MONTINE, K. S.,
WU, B., KATHIRIA, A., CHO, S. W., MUMBACH, M. R., CARTER, A. C., KASOWSKI,
M., ORLOFF, L. A., RISCA, V. I., KUNDAJE, A., KHAVARI, P. A., MONTINE, T. J.,
GREENLEAF, W. J. & CHANG, H. Y. 2017. An improved ATAC-seq protocol reduces
background and enables interrogation of frozen tissues. *Nature Methods,* 14**,** 959-962.
CORRALES, J. D., BLAESS, S., MAHONEY, E. M. & JOYNER, A. L. 2006. The level of sonic
hedgehog signaling regulates the complexity of cerebellar foliation. *Development,* 133**,**
1811-21.
DE LUCA, A., PARMIGIANI, E., TOSATTO, G., MARTIRE, S., HOSHINO, M., BUFFO, A.,
LETO, K. & ROSSI, F. 2015. Exogenous Sonic hedgehog modulates the pool of
GABAergic interneurons during cerebellar development. *Cerebellum,* 14**,** 72-85.

DEVCHAND, P. R., ZIOUZENKOVA, O. & PLUTZKY, J. 2004. Oxidative Stress and
Peroxisome Proliferator–Activated Receptors. *Circulation Research,* 95**,** 1137-1139.
DOBBING, J. & SANDS, J. 1973. Quantitative growth and development of human brain. *Arch*
Dis Child, 48**,** 757-67.
DONAT, C. K., SCOTT, G., GENTLEMAN, S. M. & SASTRE, M. 2017. Microglial Activation in
Traumatic Brain Injury. *Front Aging Neurosci,* 9**,** 208.
ENCINAS, J. M., VAAHTOKARI, A. & ENIKOLOPOV, G. 2006. Fluoxetine targets early
progenitor cells in the adult brain. *Proc Natl Acad Sci U S A,* 103**,** 8233-8.
FILOSTO, M., TONIN, P., VATTEMI, G., SAVIO, C., RIZZUTO, N. & TOMELLERI, G. 2003.
Transcription factors c-Jun/activator protein-1 and nuclear factor-kappa B in oxidative
stress response in mitochondrial diseases. *Neuropathol Appl Neurobiol,* 29**,** 52-9.
FRANKISH, A., DIEKHANS, M., JUNGREIS, I., LAGARDE, J., LOVELAND, J. E., MUDGE, J.
M., SISU, C., WRIGHT, J. C., ARMSTRONG, J., BARNES, I., BERRY, A., BIGNELL, A.,
BOIX, C., CARBONELL SALA, S., CUNNINGHAM, F., DI DOMENICO, T.,
DONALDSON, S., FIDDES, I. T., GARCIA GIRON, C., GONZALEZ, J. M., GREGO, T.,
HARDY, M., HOURLIER, T., HOWE, K. L., HUNT, T., IZUOGU, O. G., JOHNSON, R.,
MARTIN, F. J., MARTINEZ, L., MOHANAN, S., MUIR, P., NAVARRO, F. C. P.,
PARKER, A., PEI, B., POZO, F., RIERA, F. C., RUFFIER, M., SCHMITT, B. M.,
STAPLETON, E., SUNER, M. M., SYCHEVA, I., USZCZYNSKA-RATAJCZAK, B.,
WOLF, M. Y., XU, J., YANG, Y. T., YATES, A., ZERBINO, D., ZHANG, Y.,
CHOUDHARY, J. S., GERSTEIN, M., GUIGO, R., HUBBARD, T. J. P., KELLIS, M.,
PATEN, B., TRESS, M. L. & FLICEK, P. 2021. Gencode 2021. *Nucleic Acids Res,* 49**,**
D916-D923.
FRIK, J., MERL-PHAM, J., PLESNILA, N., MATTUGINI, N., KJELL, J., KRASKA, J., GOMEZ,
R. M., HAUCK, S. M., SIRKO, S. & GOTZ, M. 2018. Cross-talk between monocyte
invasion and astrocyte proliferation regulates scarring in brain injury. *EMBO Rep,* 19.
HAGENBUCHNER, J., KUZNETSOV, A., HERMANN, M., HAUSOTT, B., OBEXER, P. &
AUSSERLECHNER, M. J. 2012. FOXO3-induced reactive oxygen species are regulated
by BCL2L11 (Bim) and SESN3. *J Cell Sci,* 125**,** 1191-203.
HAMMOND, T. R., DUFORT, C., DISSING-OLESEN, L., GIERA, S., YOUNG, A., WYSOKER,
A., WALKER, A. J., GERGITS, F., SEGEL, M., NEMESH, J., MARSH, S. E.,
SAUNDERS, A., MACOSKO, E., GINHOUX, F., CHEN, J., FRANKLIN, R. J. M., PIAO,
X., MCCARROLL, S. A. & STEVENS, B. 2019. Single-Cell RNA Sequencing of Microglia
throughout the Mouse Lifespan and in the Injured Brain Reveals Complex Cell-State
Changes. *Immunity,* 50**,** 253-271 e6.
HAMMOND, T. R., ROBINTON, D. & STEVENS, B. 2018. Microglia and the Brain:
Complementary Partners in Development and Disease. *Annu Rev Cell Dev Biol,* 34**,**
523-544.
HAO, Y., HAO, S., ANDERSEN-NISSEN, E., MAUCK, W. M., 3RD, ZHENG, S., BUTLER, A.,
LEE, M. J., WILK, A. J., DARBY, C., ZAGER, M., HOFFMAN, P., STOECKIUS, M.,
PAPALEXI, E., MIMITOU, E. P., JAIN, J., SRIVASTAVA, A., STUART, T., FLEMING, L.
M., YEUNG, B., ROGERS, A. J., MCELRATH, J. M., BLISH, C. A., GOTTARDO, R.,
SMIBERT, P. & SATIJA, R. 2021. Integrated analysis of multimodal single-cell data.
Cell, 184**,** 3573-3587 e29.
JIANG, J., HUANG, Z., ZHAO, Q., FENG, W., BELIKOVA, N. A. & KAGAN, V. E. 2008.
Interplay between bax, reactive oxygen species production, and cardiolipin oxidation
during apoptosis. *Biochem Biophys Res Commun,* 368**,** 145-50.
JOYNER, A. L. & BAYIN, N. S. 2022. Cerebellum lineage allocation, morphogenesis and repair:
impact of interplay amongst cells. *Development,* 149.
KAELIN, W. G. 2005. ROS: Really involved in Oxygen Sensing. *Cell Metabolism,* 1**,** 357-358.

KAMINOW, B., YUNUSOV, D. & DOBIN, A. 2021. STARsolo: accurate, fast and versatile
822   mapping/quantification of single-cell and single-nucleus RNA-seq data. *bioRxiv***,**
823   2021.05.05.442755.
824   KANA, V., DESLAND, F. A., CASANOVA-ACEBES, M., AYATA, P., BADIMON, A., NABEL, E.,
825   YAMAMURO, K., SNEEBOER, M., TAN, I. L., FLANIGAN, M. E., ROSE, S. A., CHANG,
826   C., LEADER, A., LE BOURHIS, H., SWEET, E. S., TUNG, N., WROBLEWSKA, A.,
827   LAVIN, Y., SEE, P., BACCARINI, A., GINHOUX, F., CHITU, V., STANLEY, E. R.,
828   RUSSO, S. J., YUE, Z., BROWN, B. D., JOYNER, A. L., DE WITTE, L. D., MORISHITA,
829   H., SCHAEFER, A. & MERAD, M. 2019. CSF-1 controls cerebellar microglia and is
830   required for motor function and social interaction. *J Exp Med,* 216**,** 2265-2281.
831   KIM, D., YOU, B., JO, E. K., HAN, S. K., SIMON, M. I. & LEE, S. J. 2010. NADPH oxidase 2-
832   derived reactive oxygen species in spinal cord microglia contribute to peripheral nerve
833   injury-induced neuropathic pain. *Proc Natl Acad Sci U S A,* 107**,** 14851-6.
834   KOWALCZYK, M. S., TIROSH, I., HECKL, D., RAO, T. N., DIXIT, A., HAAS, B. J.,
835   SCHNEIDER, R. K., WAGERS, A. J., EBERT, B. L. & REGEV, A. 2015. Single-cell
836   RNA-seq reveals changes in cell cycle and differentiation programs upon aging of
837   hematopoietic stem cells. *Genome Res,* 25**,** 1860-72.
838   LEE, J. W., CHUN, W., LEE, H. J., KIM, S. M., MIN, J. H., KIM, D. Y., KIM, M. O., RYU, H. W. &
839   LEE, S. U. 2021. The Role of Microglia in the Development of Neurodegenerative
840   Diseases. *Biomedicines,* 9.
841   LETO, K., ARANCILLO, M., BECKER, E. B., BUFFO, A., CHIANG, C., DING, B., DOBYNS, W.
842   B., DUSART, I., HALDIPUR, P., HATTEN, M. E., HOSHINO, M., JOYNER, A. L., KANO,
843   M., KILPATRICK, D. L., KOIBUCHI, N., MARINO, S., MARTINEZ, S., MILLEN, K. J.,
844   MILLNER, T. O., MIYATA, T., PARMIGIANI, E., SCHILLING, K., SEKERKOVA, G.,
845   SILLITOE, R. V., SOTELO, C., UESAKA, N., WEFERS, A., WINGATE, R. J. &
846   HAWKES, R. 2015. Consensus Paper: Cerebellar Development. *Cerebellum*.
847   LI, Q., CHENG, Z., ZHOU, L., DARMANIS, S., NEFF, N. F., OKAMOTO, J., GULATI, G.,
848   BENNETT, M. L., SUN, L. O., CLARKE, L. E., MARSCHALLINGER, J., YU, G., QUAKE,
849   S. R., WYSS-CORAY, T. & BARRES, B. A. 2019. Developmental Heterogeneity of
850   Microglia and Brain Myeloid Cells Revealed by Deep Single-Cell RNA Sequencing.
851   *Neuron,* 101**,** 207-223 e10.
852   LI, Y., HE, X., KAWAGUCHI, R., ZHANG, Y., WANG, Q., MONAVARFESHANI, A., YANG, Z.,
853   CHEN, B., SHI, Z., MENG, H., ZHOU, S., ZHU, J., JACOBI, A., SWARUP, V.,
854   POPOVICH, P. G., GESCHWIND, D. H. & HE, Z. 2020. Microglia-organized scar-free
855   spinal cord repair in neonatal mice. *Nature,* 587**,** 613-618.
856   LOVE, M. I., HUBER, W. & ANDERS, S. 2014. Moderated estimation of fold change and
857   dispersion for RNA-seq data with DESeq2. *Genome Biol,* 15**,** 550.
858   MASGRAS, I., CARRERA, S., DE VERDIER, P. J., BRENNAN, P., MAJID, A., MAKHTAR, W.,
859   TULCHINSKY, E., JONES, G. D. D., RONINSON, I. B. & MACIP, S. 2012. Reactive
860   oxygen species and mitochondrial sensitivity to oxidative stress determine induction of
861   cancer cell death by p21. *J Biol Chem,* 287**,** 9845-9854.
862   MCMAHON, A. P., INGHAM, P. W. & TABIN, C. J. 2003. Developmental roles and clinical
863   significance of hedgehog signaling. *Curr Top Dev Biol,* 53**,** 1-114.
864   MEHL, L. C., MANJALLY, A. V., BOUADI, O., GIBSON, E. M. & TAY, T. L. 2022. Microglia in
865   brain development and regeneration. *Development,* 149.
866   NIETHAMMER, P. 2016. The early wound signals. *Curr Opin Genet Dev,* 40**,** 17-22.
867   PARMIGIANI, E., LETO, K., ROLANDO, C., FIGUERES-ONATE, M., LOPEZ-MASCARAQUE,
868   L., BUFFO, A. & ROSSI, F. 2015. Heterogeneity and Bipotency of Astroglial-Like
869   Cerebellar Progenitors along the Interneuron and Glial Lineages. *J Neurosci,* 35**,** 7388-
870   402.

QIU, F., CHEN, Y. R., LIU, X., CHU, C. Y., SHEN, L. J., XU, J., GAUR, S., FORMAN, H. J.,
ZHANG, H., ZHENG, S., YEN, Y., HUANG, J., KUNG, H. J. & ANN, D. K. 2014. Arginine
starvation impairs mitochondrial respiratory function in ASS1-deficient breast cancer
cells. *Sci Signal,* 7**,** ra31.
RAKIC, P. & SIDMAN, R. L. 1970. Histogenesis of cortical layers in human cerebellum,
particularly the lamina dissecans. *J Comp Neurol,* 139**,** 473-500.
SALMAN, M. S. & TSAI, P. 2016. The Role of the Pediatric Cerebellum in Motor Functions,
Cognition, and Behavior: A Clinical Perspective. *Neuroimaging Clin N Am,* 26**,** 317-29.
SCHRINER, S. E., LINFORD, N. J., MARTIN, G. M., TREUTING, P., OGBURN, C. E., EMOND,
M., COSKUN, P. E., LADIGES, W., WOLF, N., VAN REMMEN, H., WALLACE, D. C. &
RABINOVITCH, P. S. 2005. Extension of murine life span by overexpression of catalase
targeted to mitochondria. *Science,* 308, 1909-11.
SMITH, A. N., SHAUGHNESS, M., COLLIER, S., HOPKINS, D. & BYRNES, K. R. 2022.
Therapeutic targeting of microglia mediated oxidative stress after neurotrauma. *Front*
Med (Lausanne), 9**,** 1034692.
STOODLEY, C. J., D'MELLO, A. M., ELLEGOOD, J., JAKKAMSETTI, V., LIU, P., NEBEL, M.
B., GIBSON, J. M., KELLY, E., MENG, F., CANO, C. A., PASCUAL, J. M.,
MOSTOFSKY, S. H., LERCH, J. P. & TSAI, P. T. 2017. Altered cerebellar connectivity in
autism and cerebellar-mediated rescue of autism-related behaviors in mice. *Nat*
Neurosci, 20**,** 1744-1751.
STRICK, P. L., DUM, R. P. & FIEZ, J. A. 2009. Cerebellum and nonmotor function. *Annu Rev*
Neurosci, 32**,** 413-34.
TAN, D. Q. & SUDA, T. 2018. Reactive Oxygen Species and Mitochondrial Homeostasis as
Regulators of Stem Cell Fate and Function. *Antioxid Redox Signal,* 29**,** 149-168.
TAN, I. L., ARIFA, R. D. N., RALLAPALLI, H., KANA, V., LAO, Z., SANGHRAJKA, R. M.,
SUMRU BAYIN, N., TANNE, A., WOJCINSKI, A., KORSHUNOV, A., BHARDWAJ, N.,
MERAD, M., TURNBULL, D. H., LAFAILLE, J. J. & JOYNER, A. L. 2021. CSF1R
inhibition depletes tumor-associated macrophages and attenuates tumor progression in
a mouse sonic Hedgehog-Medulloblastoma model. *Oncogene,* 40**,** 396-407.
TOMLINSON, S. P., DAVIS, N. J. & BRACEWELL, R. M. 2013. Brain stimulation studies of non-
motor cerebellar function: a systematic review. *Neurosci Biobehav Rev,* 37**,** 766-89.
TRAAG, V. A., WALTMAN, L. & VAN ECK, N. J. 2019. From Louvain to Leiden: guaranteeing
well-connected communities. *Sci Rep,* 9**,** 5233.
TSAI, P. T., RUDOLPH, S., GUO, C., ELLEGOOD, J., GIBSON, J. M., SCHAEFFER, S. M.,
MOGAVERO, J., LERCH, J. P., REGEHR, W. & SAHIN, M. 2018. Sensitive Periods for
Cerebellar-Mediated Autistic-like Behaviors. *Cell Rep,* 25**,** 357-367 e4.
WANG, J., HE, X., MENG, H., LI, Y., DMITRIEV, P., TIAN, F., PAGE, J. C., LU, Q. R. & HE, Z.
2020. Robust Myelination of Regenerated Axons Induced by Combined Manipulations of
GPR17 and Microglia. *Neuron,* 108**,** 876-886.e4.
WANG, S. S., KLOTH, A. D. & BADURA, A. 2014. The cerebellum, sensitive periods, and
autism. *Neuron,* 83**,** 518-32.
WECHSLER-REYA, R. J. & SCOTT, M. P. 1999. Control of neuronal precursor proliferation in
the cerebellum by Sonic Hedgehog [see comments]. *Neuron,* 22**,** 103-14.
WOJCINSKI, A., LAWTON, A. K., BAYIN, N. S., LAO, Z., STEPHEN, D. N. & JOYNER, A. L.
2017. Cerebellar granule cell replenishment postinjury by adaptive reprogramming of
Nestin(+) progenitors. *Nat Neurosci,* 20, 1361-1370.
ZHENG, G. X., TERRY, J. M., BELGRADER, P., RYVKIN, P., BENT, Z. W., WILSON, R.,
ZIRALDO, S. B., WHEELER, T. D., MCDERMOTT, G. P., ZHU, J., GREGORY, M. T.,
SHUGA, J., MONTESCLAROS, L., UNDERWOOD, J. G., MASQUELIER, D. A.,
NISHIMURA, S. Y., SCHNALL-LEVIN, M., WYATT, P. W., HINDSON, C. M.,
BHARADWAJ, R., WONG, A., NESS, K. D., BEPPU, L. W., DEEG, H. J., MCFARLAND,

C., LOEB, K. R., VALENTE, W. J., ERICSON, N. G., STEVENS, E. A., RADICH, J. P.,
MIKKELSEN, T. S., HINDSON, B. J. & BIELAS, J. H. 2017. Massively parallel digital
transcriptional profiling of single cells. *Nat Commun,* 8**,** 14049.
925

Figure 1: Injury induces ROS and cell stress signaling reflected by changes in the transcriptome and chromatin landscape of gliogenic-NEPs.

(A) Schematic summarizing the experimental plan.

(B-C) UMAP projections of 11,878 cells (6,978 nonIR and 4,900 IR) showing cluster annotations (B), treatment (black: nonIR, red: IR) and the age of the samples (red: P1, green: P2, blue: P3, purple: P5) (C).

(D) Dot plot showing the expression levels of key marker genes used for cluster annotation (gliogenic-NEPs: *Hopx, Gdf10, Slc1a3*, neurogenic-NEPs: *Ascl1, Ptf1a*, immature neurons: *Pax2*, GCPs: *Atoh1*, postmitotic neurons: *Tubb3*, microglia: *Cx3cr1*, oligodendrocyte progenitors: *Olig2*, oligodendrocytes: *Gpr17*, Ependymal cells: *Foxj1*).

(E) Feature plots showing *Hopx* (gliogenic-NEPs), *Ascl1* (neurogenic-NEPs) and *Atoh1* (GCPs) expression highlighting the three main populations of interest. Clusters containing *Hopx*-NEPs (clusters 2, 3, 6, 10), *Ascl1*-NEPs (clusters 5, 8, 11), or GCPs (clusters 1, 4, 7, 12, 14) were subsetted from the original data set and were divided according to age (P2 or P3+P5) for the downstream differential expression analyses.

(F) Volcano plot showing differentially expressed genes in the P2 gliogenic-NEPs (red: adjusted p-value≤0.05, log$_2$fold-change>|1|).

(G) Violin plots showing some of the top differentially expressed genes in P2 gliogenic-NEPs and how their expression changes over time with respect to their expression in control cells.

(H) Top GO terms associated with differentially expressed genes in P2 gliogenic-NEPs that were either upregulated in nonIR (top panel) or IR (bottom panel) cells (adjusted p-value≤0.05, Table S3).

(I) Heatmap showing differentially open chromatin regions in P2 nonIR and IR NEPs, identified by bulk ATAC-seq (1168 differentially open regions, adjusted p-value<0.05, Table S4).

(J) Tracks highlighting the injury-induced open chromatin regions around *Cdkn1a* and *Phlda3,* the top differentially expressed genes identified in (F).

(K) Gene network analysis of ATAC-seq data revealed an active transcriptional network involved in regulating cell death and apoptosis. Genes colored in red (*Ppara*, *Egln3*, *Foxo3*, *Jun* and *Nos1ap)* have been implicated as upregulated with increased ROS levels or involved in ROS signaling.

Figure 2: Cerebellar injury at P1 results in increased superoxide production, a reduction in mitochondria and increased cell death in the EGL that peaks 24h after injury.

(A) Examples of flow cytometry analysis of mitochondrial ROS at P2 from nonIR and IR cerebella using MitoSOX dye. Gating determined the top 90% MitoSOX signal (MitoSOX high cells).

(B, C) Quantification of MitoSOX high (B) and MitoTtracker high (C) expression in nonIR and IR cerebella at P2.

(D, E) Quantification of TUNEL+ cell density in the EGL at P2 (D) and P3 (E) in lobules 3-5 of nonIR and IR mice.

(F, G) Quantification of TUNEL+ cell density outside the EGL at P2 (F) and P3 (G) in lobules 3-5 of nonIR and IR mice.

EGL, External granular layer; SSC, side scatter; P, postnatal day; nonIR, non-irradiated; IR, irradiated. All statistical significance was determined using an unpaired t-test and data are represented as mean ± SEM.

Figure 3: Cerebellar injury at P1 induces transient microglial recruitment to the EGL and prolonged astroglial microenvironment changes in the cerebellum.

(A-H) Immunohistochemical (IHC) staining of medial sagittal cerebellar sections for GFAP (green) in lobules 4/5 of nonIR and IR cerebellum at the stages indicated. Nuclei were counterstained with Hoechst. (B') and (F') show high-power images of white dashed line boxes in (B) and (F), respectively.

(I, J) Quantification of IBA1+ cell density in the WM at P2 (I) and P3 (J) in lobules 3-5 of nonIR and IR mice.

(K, L) IHC staining of medial sagittal cerebellar sections for IBA1 and TUNEL in lobule 3 of nonIR and IR cerebellum at P2. Nuclei were counterstained with Hoechst. White matter (WM) and external granular layer (EGL) are delineated by white dotted lines and dashed lines, respectively. High-power image in (L) of the area indicated by the white dashed line represents an IBA1+ cell present in the EGL. White arrowheads indicate additional IBA1+ cells in the EGL.

(M, N) Quantification of IBA1+ cell density in the EGL at P2 (M) and P3 (N) in lobules 3-5 of nonIR and IR mice. EGL, External granular layer; WM, White matter; P, postnatal day; nonIR, non-irradiated; IR, irradiated. Scale bar: A and E 100µm, B, C, D, E, F, G and H: 250µm, B' and F': 50µm, I and J: 50µm. All statistical significance was determined using an unpaired t-test and data are represented as mean ± SEM.

Figure 4: Reduction of ROS impairs adaptive reprogramming and cerebellar repair.

(A, B) IHC staining of medial sagittal cerebellar sections for human catalase in control (A) AND *mCAT/+* mice (B) at P7. Nuclei were counterstained with Hoechst (blue). Similar staining was seen in four *mCAT/+* mice.

(C) Quantification of MitoSOX high expression at P2 in control and *mCAT/+* cerebella, with and without irradiation at P1 (Two-way ANOVA, $F_{(1,34)}$=6.768, p=0.0136).

(D) Quantification of MitoTracker high expression at P2 in control and *mCAT/+* cerebella, with and without irradiation at P1 (Two-way ANOVA, $F_{(1,31)}$=25.06, p<0.0001).

(E) Quantification of TUNEL+ cell density in the EGL at P2 in control and *mCAT/+* cerebella, with and without irradiation at P1 (Two-way ANOVA, $F_{(1,14)}$=87.56, p<0.0001).

(F) Quantification of IBA1+ cell density in the EGL at P2 in control and *mCAT/+* cerebella, with and without irradiation at P1 (Two-way ANOVA, F(1,14)=15.58, p=0.0015).

(G-J) Hematoxylin and eosin staining on mid-sagittal sections of P30 control and *mCAT/+* cerebellum with or without irradiation.

(K) Quantification of P30 cerebellar mid-sagittal section area in controls and *mCAT/+* nonIR and IR mice (Two-way ANOVA, $F_{(1,20)}$=11.82, p=0.0026).

(L) Graph showing the average area of mid-sagittal cerebellar sections at P3, P5, P8, P12 and P30 in control and *mCAT/+* non-irradiated and irradiated mice. Detailed statistics are shown in Supplementary Figure 4.

EGL, External granular layer; P, postnatal day; nonIR, non-irradiated; IR, irradiated. Scale bar: A and B: 100μm, F-I: 1mm. Significant *Tukey's post hoc* multiple comparison tests are shown in the figures and data are represented as mean ± SEM.

Figure 5: Reduced ROS impairs expansion of BgL-NEPs and their recruitment to the EGL after injury.

(A, B) Quantification of EGL thickness at P5 (Two-way ANOVA, $F_{(1,21)}$=36.64, p<0.0001) (A) and P8 (Two-way ANOVA, $F_{(1,12)}$=11.34, p=0.0056)(B) in lobules 3-5 of *Nes-Cfp/+* control and *Nes-Cfp/+; mCAT/+* mutant mice with and without irradiation at P1

(C-F) IHC staining of medial sagittal cerebellar sections showing expression of CFP (green) in the lobules 4/5 of *Nes-Cfp/+* control and *Nes-Cfp/+; mCAT/+* mutant mice at P5. Nuclei were counterstained with Hoechst (blue). (D') and (F') show a high-power images of the yellow boxed area in the single channel CFP. EGL is delineated by the dashed white lines.

(G, H) Quantification of CFP+ cell density in the EGL (Two-way ANOVA, $F_{(1,19)}$=5.192, p=0.0359) (G) and BgL (Two-way ANOVA, $F_{(1,17)}$=6.191, p=0.0223) (H) at P5 in *Nes-Cfp/+* control or *Nes-Cfp/+; mCAT/+* mutant non-irradiated and irradiated mice.

EGL, External granular layer; BgL, Bergmann glia Layer; P, postnatal day; nonIR, non-irradiated; IR, irradiated. Scale bar: D-F: 100µm. Significant *Tukey's post hoc* multiple comparison tests are shown in the figures and data are represented as mean ± SEM.

Figure 6: Microglia promote recruitment of NEPs to the EGL during cerebellar adaptive reprogramming after injury.

(A-D) IHC staining of medial sagittal cerebellar sections for IBA1 (red) in control and *mCAT/+* mice at P5. Nuclei were counterstained with Hoechst (blue).

(E, F) Quantification of IBA1+ cell density in the external granular layer (E) and white matter (Two-way ANOVA, $F_{(1,13)}$=24.74, p=0.0003) (F) at P5 on midsagittal sections of lobules 3-5 in the cerebellum of control and *mCAT/+* animals, with or without irradiation.

(G-J) IHC staining of medial sagittal cerebellar sections at P5 for CFP (green) in lobules 4/5 of *Nes-Cfp/+* mice treated with PLX5622 or control DMSO with or without irradiation. Nuclei were counterstained with Hoechst (blue). (H') and (J') show a high-power image of area indicated by yellow boxes. EGL is delineated by the white dashed lines.

(K) Quantification of IBA1+ cell density in the white matter at P5 on mid-sagittal sections in lobules 3-5 of *Nes-Cfp/+* mice treated with PLX5622 or control DMSO, with or without irradiation (Two-way ANOVA, $F_{(1,12)}$=42.40, p<0.001).

(L) Quantification of EGL thickness at P5 in lobules 3-5 of *Nes-Cfp/+* mice treated with PLX5622 or control DMSO with or without irradiation (Two-way ANOVA, $F_{(1,12)}$=109.5, p<0.001).

(M) Quantification of CFP+ cells density in the EGL at P5 on mid-sagittal sections in lobules 3-5 of *Nes-Cfp/+* mice treated with PLX5622 or control DMSO with or without irradiation (Two-way ANOVA, $F_{(1,12)}$=10.62, p=0.0068).

(N) Measurement of cerebellar mid-sagittal section area at P30 in controls or mice treated with PLX, with or without irradiation at P1 (Two-way ANOVA, $F_{(1,12)}$=13.29, p=0.0034).

EGL, External granular layer; WM, White matter; P, postnatal day; nonIR, non-irradiated; IR, irradiated. Scale bar: A-D and G-J: 250 µm. Significant *Tukey's post hoc* multiple comparison tests are shown in the figures and data are represented as mean ± SEM.

A

B

	P1	P2		P3		P5	
	NON-IR	NON-IR	IR	NON-IR	IR	NON-IR	IR
Replicate 1	598	564	1085	1293	896	866	2407
Replicate 2	643	826	880	-	1330	-	-
Total	1241	1390	1965	1293	2226	866	2407

C

Supplementary Figure 1. scRNA-seq quality metrics and number of cells sequenced in each condition and biological replicate.

(A) Violin plots showing the number of features, RNA and percent mitochondrial RNA count across the biological replicates of the scRNA-seq data set after filtering the bad quality cells (cells were filtered out where number of detected genes was ≤ 1500, the number of detected transcripts was ≥ 40,000 and mitochondrial gene percentage ≥ 5%).

(B) Number of cells from each replicate and condition used for downstream analyses after filtering.

(C) UMAPs showing the distribution of cells across different clusters based on the samples.

Supplementary Figure 2: Injury induces distinct transcriptional changes in NEP subtypes and GCPs during adaptive reprograming.
(A-E) Volcano plot showing differentially expressed genes in the P3+5 gliogenic-NEPs (A), P2 and P3+5 neurogenic NEPs (B, C) and P2 or P3+5 GCPs (D, E) (red: adjusted p-value≤0.05, $\log_2$fold-change=1, Table S2).
(F-J) Top GO terms associated with differentially expressed genes in the P3+5 gliogenic-NEPs (F), P2 or P3+5 neurogenic NEPs (G, H) and P2 or P3+5 GCPs (I, J) that were either upregulated in nonIR (top panel) or IR (bottom panel) (p-value≤0.05).

Supplementary Figure 3: Irradiation of cerebella at P1 results in increased superoxide production and cell death and recruitment of microglia to the EGL that peaks at 24h.

(A, B) Quantification of high MitoSOX expression in nonIR and IR cerebella at P3 (A) and P5 (B).

(C) Examples of flow cytometry analysis of mitochondria at P2 from nonIR and IR cerebella using MitoTracker dye. Gating determined the top 90% MitoTracker signal (MitoTracker high cells).

(D, E) Quantification of MitoTracker high expression in nonIR and IR cerebella at P3 (D) and P5 (E).

(F) Quantification of IBA1+ cell density in the WM at P5 in lobules 3-5 of nonIR and IR mice.

(G) Quantification of IBA1+ cell density in the EGL at P5 in lobules 3-5 of nonIR and IR mice.

EGL, External granular layer; WM, White matter; P, postnatal day; nonIR, non-irradiated; IR, irradiated. All statistical significance was determined using an unpaired t-test and data are represented as mean ± SEM.

Supplementary Figure 4: Reduction of ROS impairs adaptive reprogramming and cerebellar repair.
(A, B) Quantification of high MitoSOX expression at P3 (A) and P5 (B) in control and *mCAT/+* cerebella, with and without irradiation at P1.
(C, D) Quantification of MitoTracker high expression at P3 (C) and P5 (D) in control and *mCAT/+* cerebella, with and without irradiation at P1.
(E-G) Quantification of TUNEL+ cell density outside EGL at P2 (Two-way ANOVA, $F_{(1,10)}$=14.20, p=0.0037) (E), at P3 (G), and in the EGL at P3 (F) in lobules 3-5 of nonIR and IR mice.
(H-J) Quantification of IBA1+ cell density in WM at P2 (H), at P3 (J), and in the EGL at P3 (I) in lobules 3-5 of nonIR and IR mice.
(K-N) Quantification of cerebellar midsagittal section area in controls and *mCAT/+* nonIR and IR mice at P3 (K), P5 (Two-way ANOVA, $F_{(1,15)}$=28.52, p<0.001) (L), P8 (Two-way ANOVA, $F_{(1,12)}$=21.21, p=0.0006) (M) and P12 (Two-way ANOVA, $F_{(1,14)}$=9.682, p=0.0077) (N).
EGL, External granular layer; WM, White matter; P, postnatal day; nonIR, non-irradiated; IR, irradiated. Significant *Tukey's post hoc* multiple comparison tests are shown in the figures and data are represented as mean ± SEM.

Supplementary Figure 5: Reduced ROS impairs expansion of BgL-NEPs and their migration to the EGL after injury.
(A) Quantification of CFP+ cell density in the EGL at P3 in *Nes-Cfp/+* control or *Nes-Cfp/+ mCAT* mutant nonIR and IR mice. (Two-way ANOVA, $F_{(1,14)}=33.77$, $p<0.0001$).
(B) Quantification of CFP+ cell normalized on BgL length at P5 in *Nes-Cfp/+* control or *Nes-Cfp/+ mCAT* mutant nonIR and IR mice.
(C, D) Quantification of SOX2+ NEP cell density on BgL length at P8 (C) and P12 (Two-way ANOVA, $F_{(1,12)}=12.50$, $p=0.0033$) (D) in control or *mCAT* mutant nonIR and IR mice.
EGL, External granular layer; BgL, Bergmann glia Layer; P, postnatal day; nonIR, non-irradiated; IR, irradiated. Significant *Tukey's post hoc* multiple comparison tests are shown in the figures and data are represented as mean ± SEM.

Supplementary Figure 6: Microglia promote recruitment of NEPs to the EGL during cerebellar adaptive reprogramming after injury.
(A) Quantification of IBA1+ cell density in the WM at P8 in control or *mCAT* mutant nonIR and IR mice.
(B-E) Immunostaining of medial sagittal cerebellar sections at P5 showing expression of IBA1 (red) in mice treated with PLX5622 or control DMSO, with or without irradiation.
(F) Quantification of SOX2+ NEP cell density in the BgL at P5 in control or PLX treated nonIR and IR mice.
BgL, Bergmann glia Layer; WM, White Matter; P, postnatal day; nonIR, non-irradiated; IR, irradiated. Scale bar: 500 μm. Data are represented as mean ± SEM.

Supplementary Tables

Antigen	Species	Concentration	References	Source
Catalase	Rabbit	1 to 100	01-05-030000	Athens Research & Technology
GFAP	Chicken	1 to 500	ab4674	Abcam
GFP	Rat	1 to 1000	440484	Nacalai Tesque
IBA1	Rabbit	1 to 500	019-19741	Wako Chemicals
SOX2	Goat	1 to 500	AF2018	R&D System

Table S1: List of antibodies and related information.

Table S2. Marker genes expressed by cluster in scRNA-seq dataset (irradiated at P1 (IR; P2, P3, P5) or non-irradiated (nonIR; P1, P2, P3, P5). pct1: % cells in a cluster that express the gene; pct2: % cells that express the gene outside the given cluster.

Table S3. Pseudobulk differential expression analysis between nonIR and IR gliogenic-NEPs (*Hopx+,* clusters 2, 3, 6, 10), neurogenic-NEPs (*Ascl1+,* clusters 5, 8, 11) and GCPs (*Atoh1+,* clusters 1,4,7,12,14) at P2, or at P3 and P5 (P3+5).

Table S4. GO Term analyses of differentially expressed genes (Table S3) of nonIR and IR gliogenic-NEPs (*Hopx+,* clusters 2, 3, 6, 10), neurogenic-NEPs (*Ascl1+,* clusters 5, 8, 11) and GCPs (*Atoh1+,* clusters 1,4,7,12,14) at P2, or at P3 and P5 (P3+5).

Table S5. Differentially open peaks at P2 identified by bulk ATAC-seq from nonIR and IR NEPs.

Table S6. Motif analysis of regions with increased accessibility in IR NEPs compared to the nonIR at P2.