What's in it for the dogs? Assessing the outcomes of a prison-based dog training

program from an animal behavior and welfare perspective

Parizad Baria-Unwalla[1,2]

Gabriela Munhoz Morello[1,2]

Maria Queiroz[3]

I. Anna S. Olsson[1,2]

Ana Catarina Vieira de Castro[1,2*]

1. IBMC - Instituto de Biologia Molecular e Celular, Universidade do Porto, Porto,

Portugal

2. i3S – Instituto de Investigação e Inovação em Saúde, Universidade do Porto,

Porto, Portugal

3. DTC Social, R. de Vila Verde nº 7, Touguinhó, Portugal

*E-mail: ana.castro@ibmc.up.pt

ORCID

Parizad Baria-Unwalla: https://orcid.org/0000-0001-7701-170X

Gabriela Munhoz Morello: https://orcid.org/0000-0002-5439-969X

I. Anna S. Olsson: https://orcid.org/0000-0002-4369-8723

Ana Catarina Vieira de Castro: https://orcid.org/0000-0002-3357-7282

Abstract

Prison-Based Dog Training Programs (PBDTPs) are gaining popularity across the

world for their benefits to inmates in terms of mental and emotional health, and

reduced recidivism. However, the implications for the dogs have never been studied.

The aim of the present study was to assess the outcomes of a PBDTP from an animal

behavior and welfare perspective. Shelter dogs (N=42) were transported to and from

the prisons twice a week where they received training from inmates for a total

duration of 12 weeks. Dogs were tested for potential improvements in sociability and

basic training skills using the Temperament Test (Barnard et al., 2019; Valsecchi et

al., 2011) and the Basic Education Test (adapted from the Canine Good Citizen Test

by AKC). Dog welfare was assessed using behavioral, physiological, and cognitive

measures, namely stress-related behaviors and overall behavior states during training

sessions, salivary cortisol levels, and cognitive bias tests. Results showed that

participating in the PBDTP improved the dogs' sociability towards humans and

increased playful behavior, and improved basic training skills including not jumping

on people, walking on a leash without pulling, responding to commands (sit, down,

and staying in place) and staying calm when separated from the handler. Furthermore,

behaviors indicative of stress were generally rare during training sessions and no

impact of the PBDTP was found on the levels of salivary cortisol nor on the dogs'

affective states (as measured with cognitive bias tests). In conclusion, the present

study suggests that PBDPTs are beneficial for dogs, with animals showing no

indicators of compromised welfare while displaying improved behavior skills, which

will likely facilitate their future rehoming.

Keywords: Dog Training; Prison-Based Dog Training Program; Shelter Dogs; Dog

Adoption; Dog Behavior; Animal Welfare

Introduction

Correctional facilities across the world, including in USA (Cooke & Farrington, 2016), UK (Mercer et al., 2015), Australia (Humby & Barclay, 2018), Netherlands (Karkdijk, 2022), and Japan (Koda et al., 2015) use different forms of prison-based dog programs to improve inmates' social and emotional functioning and to reduce recidivism (Duindam et al., 2020; Grommon et al., 2018; Jasperson, 2013). Dog training programs are one such form, along with other programs, including therapeutic interventions such as Animal Assisted Therapy.

Prison-based dog programs are based on the understanding that human-animal interactions bring psychosocial benefits including reduced stress, increased social support and emotional well-being (Cooke & Farrington, 2016). The beneficial outcomes of Prison-Based Dog Training Programs (PBDTPs) on human participants (primarily inmates, but also staff at the correctional facility and the community at large) include increased offender self-control (Currie, 2018; Davis, 2007; Furst, 2006), increased self-esteem and self-efficacy (Gilger, 2007; Strimple, 2003), increased empathy for dogs and humans (Currie, 2018; Davis, 2007), increased emotional intelligence, and increased employability (Cooke & Farrington, 2016).

The popularity of PBDTPs has been on the rise worldwide in recent years which may be in part due to the positive media attention they receive and the public appeal of connecting shelter dogs and prison inmates, two vulnerable groups that need social contact (Britton & Button, 2005; Nimer & Lundahl, 2007). In the United States alone, over 330 correctional facilities have implemented some form of PBDTPs (Cooke, 2019). Research into PBDTPs has nearly exclusively focused on the impact on human participants (Cooke & Farrington, 2016), whereas there is hardly any data on the effect on dog health and welfare (Glenk & Foltin, 2021).

Dog training programs are potentially beneficial to dogs in several ways.

Shelter dogs have higher chronic cortisol levels than owned dogs (Dudley et al., 2015;

Sandri et al. 2015), which is an indicator of higher stress levels in shelter dogs.

However, interactions with humans at shelters, including walking and play, can

reduce cortisol levels in dogs (Menor-Campos et al., 2011; Shiverdecker et al., 2013),

highlighting the potential benefits to dogs of positive human interactions. Many

PBDTPs bring in dogs from shelters (Cooke & Farrington, 2016), which can

potentially reduce their stress levels. PBDTPs may also create indirect benefits for the

participating dogs. The training that these dogs receive may make them more

attractive to potential adopters, and potentially facilitate their integration in a new

family. Additionally, PBDTPs impart useful skills to dogs who sometimes go on to

become service or working dogs and might have been euthanized otherwise (Deaton,

2005). However, participating in the training program also implies potentially

stressful experiences, including change of environment, transport and interactions

with unfamiliar and initially unexperienced handlers. Failing to consider the welfare

of dogs in PBDTPs may put participants, facilitators, and dogs at risk (Johnson &

Bruneau, 2019).

There has been almost no research into how PBDTPs affect dogs, with a

consequent lack of evidence regarding efficacy and welfare impact of these programs

(Johnson and Bruneau, 2019). Hennessy et al (2006) found that shelter dogs

participating in a 3-week PBDTP exhibited increased training performance and

decreased behavioral arousal in a novel situation from before to after the program as

compared to control dogs that remained at the shelter. However, no differences were

found between groups in physiological measures, with plasma cortisol levels

remaining constant and pituitary hormone (ACHT) increasing from before to after the

program. More recently, Leornardi et al (2024) found that, after participation in a

PBDTP, shelter dogs improved their performance in training tasks, showed increased

levels of relaxed behaviors in their home kennels and received enhanced subjective

ratings of behavior by shelter staff. Taken together, the results of these two studies

suggest that PBDTPs improve dog training performance and do not impair dog

welfare, with Leonardi et al (2024) indeed suggesting a positive effect of participation

in the program on the welfare of the animals.

While the results of these studies are promising, they are limited: Hennessy et

al (2006) only used physiological measures of stress (which provide information on

the arousal but not on the valence of the animals' emotions, e.g., Broom and Johnson,

2019), and the main welfare measure used by Leonardi et al (2024) comprised

subjective scoring by non-blind raters. In addition, it is important to assess how

participation in a PBDTP affects dogs both inside and outside of the training

environment (i.e. the prisons).

Therefore, the aim of the present study was to perform a comprehensive

evaluation of the effects of PBDTPs on dog behavioral prowess and welfare. To that

end, we analyzed several indicators of dog sociability and basic training before and

after participation in a PBDTP and combined physiological, behavioral and cognitive

measures to assess dog welfare before, during and after the program. Given the rising

popularity of PBDTPs, systematic research on the topic is of critical importance.

Animals, Materials and Methods

Ethics Statement

Approvals were obtained from the i3S Committee for Ethics and Responsible

Conduct in Research (N9/CECRI/2022) and from the i3S Animal Welfare and Ethics

Review Body (internal reference 2021-29). All the involvement of the inmates and the

dogs in the PBDTP was under the management and the responsibility of DTC Social

(see below on 'The Prison-Based Dog Training Program'). No identifying

information was collected or recorded about inmates in the program for the purposes

of the present study, and all results are presented without any disclosure of personal

information. During training sessions at the prisons, datasheets were used to code dog

behavior. These beared no information regarding the identity of the inmates training

each dog and no visual recordings were made of sessions in the prisons. All collected

data was coded on MS Excel, and stored in encrypted folders on two different hard

disks, exclusively for research use. Video recordings from the Temperament Tests

(TTs), Basic Education Tests (BETs) and cognitive bias tests (described below) only

featured the authors of the present study and the participating dogs.

No invasive or stressful tests were conducted with the animals: saliva

sampling was non-invasive and implied only brief low-stress handling, behavior

assessments during training sessions were purely observational and the behavior tests

only implied tasks that are part of regular dog-human interactions for family dogs.

The researchers worked under the close supervision of the DTC Social team,

who were responsible for ensuring the necessary health and safety protocols.

The Prison-Based Dog Training Project

Pelos2 is a PBDTP conducted by DTC Social, a Portuguese NGO that works

with dogs, and funded by Portugal Inovação Social, a public initiative to promote

social innovation in Portugal. The program, based on the training of shelter dogs by

prison inmates, has a twofold aim: 1) imparting inmates with occupational skills and

reducing the impact of incarceration on their mental health, and 2) granting dogs with

educational skills and attempt at promoting their successful rehoming.

The research team at i3S (i.e., the authors of this paper) collaborated with

DTC Social to collect research data on dogs participating in the program in three male

prisons in the north of the country: Santa Cruz do Bispo, Vale do Sousa, and Paços de

Ferreira

Participants

Expecting high variability in the sample, even relatively large differences

might yield small effect sizes. Hence, we powered this experiment to detect a Cohen's

f=.25. Power analysis (G*Power) for a repeated-measures design deemed N=34

sufficient (α=.05, β=0.8). Forty-five dogs were recruited for the present study and

equally allocated by DTC Social to three male prisons: Paços de Ferreira (Prison 1),

Vale do Sousa (Prison 2), and Santa Cruz do Bispo (Prison 3). During the course of

the program, however, three dogs dropped out (two due to health issues and one due

to behavior management issues at the prison); hence we ended up with a final sample

of 42 animals.

The participating dogs were selected by DTC Social from the Plataforma de

Acolhimento e Tratamento Animal (PATA), a municipal companion animal shelter in

the Porto region. Selected dogs were between 1-8 years of age, healthy, without

visible signs of aggression or extreme fear (as determined by DTC trainers in a brief

interaction with the animals), and had been at the shelter for at least three weeks

before the PBDTP commenced. While it was attempted to recruit an equal number of

male and female dogs, the predefined eligibility criteria and availability of dogs at

PATA did not allow this, hence we ended up with 12 female and 33 male dogs in the

program. Detailed demographics are depicted in Appendix 1.

For the duration of the program, these dogs were not available for immediate

adoption (i.e., potential adopters were able to place the dogs on hold but could only

take them home once their participation in the program was complete). The

information regarding their participation in the training program was displayed on a

sign at their home enclosures at PATA.

Experimental Design and Data Collection

Training Program

The basic training skills imparted to the dogs included sit, down, stay, come

when called, loose leash walking, and not jumping on people. The socialization and

handling skills included sociability with dogs and humans, handling for hygiene,

grooming and veterinary procedures, resource guarding, confidence, and

independence. Only reward-based methods (positive reinforcement training) were

used. An overview of the training plan followed by the DTC Team is presented in

Appendix 2.

The dogs were picked up from PATA at 9 am and returned at 4 pm, from

Monday to Thursday. Each dog was allocated to one prison, and transported there two

days a week (on alternate days). On these days, they participated in three different

sessions, each with a different inmate. Each session was 45 minutes in duration,

including walking, playtime, and 15 minutes of dedicated training. In between

training sessions, the dogs remained resting in their crates in the van for 15 to 90

minutes (15 minutes between subsequent sessions, up to 90 minutes during lunch

break). The vans were equipped with ventilators and were always parked in shaded

areas, with the doors at the back and side left ajar enough that there was enough

ventilation at all times. On Friday, Saturday, and Sunday, all dogs remained at PATA.

At PATA, the dogs were kept in individual kennels and were not routinely handled

beyond the basic human interactions around feeding and kennel cleaning. They were

fed between 8:00 and 8:30 am every day at the shelter and had water available

throughout the day, including during the training sessions at the prisons.

The program lasted 12 weeks each, for four different batches of dogs. The first

two batches included 6 dogs each, the third batch included 17 dogs and the fourth

batch 16 dogs (after accounting for dropouts).

Testing

A detailed schedule for the testing timeline is provided in Figure 1.

Figure 1. Study timeline. Dog sociability and basic training indicators were assessed with TT - Temperament Test and BET – Basic Education

Test, respectively, and dog welfare with salivary cortisol concentration (B1, B2, C1-C3, T1-T3, F1, F2), stress behaviors (StrB) and cognitive

bias tasks (CB). Dogs were tested with TT, BET and CB before (pre-program, week 0) and after participation in the PBDTP (post-program,

week 13). TTs and BETs were conducted over two days at each stage (half of the subjects on each day). Stress behaviors were assessed during

PBDTP training sessions at the prisons (Pr1- Prison 1; Pr2 – Prison 2; Pr3 – Prison 3) at three time-points: week 2, week 7, and week 11. Saliva

samples T1-T3 (T; training) were collected on these same days at the prisons; and samples C1-C3 (C, control) were collected at PATA shelter at

the same time-points (week 2, week 7, week 11) but on a different day, when the dogs were not transported to the prisons and remained at the

shelter. B1 and B2 are the pre-program baseline samples and F1 and F2 the post-program final samples collected at PATA on the same days the

BETs   were   conducted.

For evaluating dog welfare during the participation in the PBDTP, behavioral and physiological measures were used. The behavior of the dogs during the first training session of the day at three time-points of the PBDTP (weeks 2, 7, and 11) was coded live using previously validated ethograms (Appendices 5 and 6) by ACVC and Parizad Baria-Unwalla (PB). Each researcher coded the behavior of three different dogs, following each dog for five minutes. They used continuous sampling for coding stress behaviors, including lip-licking, yawning, moving away, crouches, body-turns, and body-shakes (as per definitions in Appendix 5), and instantaneous scan sampling every 60 seconds for coding overall behavior states (Tense, Low, Relaxed, Excited, as defined in Appendix 6). The ethograms were based on studies of dog welfare in a training context (e.g., Cooper et al, 2014; Vieira de Castro et al, 2020), and include commonly observed behaviors during training and especially those most prone to show differences in stressful vs non-stressful conditions. The two researchers coding dog behavior during the training sessions trained and calibrated themselves for inter-observer reliability from previously recorded dog training videos from a different project, until a high level of agreement was achieved (>90% for the frequency of stress-related behaviors; >70% for overall behavior states). For stress-related behaviors, the reliability was checked using intra-class correlation and for overall behavior states, it was checked using Cohen's Kappa statistic.

Salivary cortisol concentration was used as a physiological measure of welfare, in conjunction with the behavioral measures (Cooper et al., 2014; Vieira de Castro et al., 2020). Samples of dogs' saliva were collected with cotton buds (Salivette®). One of the researchers used a cotton bud and moved it in gentle circular motions on the inside of the dogs' cheeks, while stroking and praising the dog. Before

the program commenced, two baseline samples (B1 and B2) were collected at the

shelter between 11:00 and 15:00 in two different days (the same days on which the

BETs were performed, and immediately after taking the dogs out of their home pens

for testing, see Figure 1). Once the PBDTP started, samples were collected between

12:30 and 14:30 on training days at the prisons during weeks 2, 7, and 11 (training

samples; T1, T2 and T3). The dogs had undergone training sessions in the morning

and had rested in the crates/vans for at least one hour prior to saliva collection.

Samples were also collected during the same hour range at PATA on Fridays of the

same weeks, when the dogs were at the shelter all day (control samples; C1, C2 and

C3). After the program ended, two other samples (F1 and F2) were collected at the

shelter between 11:00 and 15:00 in two different days (once again, the same days on

which the BETs were performed, see Figure 1). The goal of collecting B samples was

to allow investigating (i) whether C samples could be used as true controls for T

samples (i.e., whether the cortisol levels of the animals when at the shelter would not

change with the participation in the PBDTP) and (ii) together with F samples,

examining whether participating in the PBDTP would affect dogs' cortisol levels at

the shelter outside the program period. All samples were labelled with the dog ID,

date, time, stored in individual plastic tubes on ice, and moved as soon as possible to a

-20 Celsius freezer at i3s. At the end of data collection for each batch, the samples

were sent to an external laboratory (DNAtech) where they were tested for cortisol

concentration with ELISA.

Cognitive bias tests were also carried out before (week 0) and after the

PBDTP (week 13) to evaluate the effects of the program in the affective state of the

dogs. Dogs who were unmanageable on the leash during the Temperament Test were

excluded from selection for this test, as well as dogs who showed no interest in food.

The cognitive bias tests were conducted in an indoor room at PATA. The dogs

learned a discrimination between two food bowl locations (positive, which contained

food, and negative, which was empty) and were then tested with an ambiguous

(empty) food bowl, equally spaced between the two training locations. The time (in

seconds) dogs took to reach the food bowls was scored live using a stopwatch. The

entire protocol is detailed in Appendix 9.

Due to logistical constraints, the TT and BET tests were conducted for all

batches (1-4), whereas smaller subsets of dogs were tested for cortisol levels, stress

behaviors and cognitive bias. A total of 42 dogs underwent TT testing during both the

pre- and post-program assessments. However, two dogs (marked with * in

Appendix 1) refused to walk on a leash during the pre-program BET, and thus could

not be tested at that stage. Consequently, 40 dogs were tested in both the pre- and

post-program BET. Moreover, four dogs (marked with ** in Appendix 1) did not

allow handling for saliva collection and were excluded from cortisol sampling to

avoid causing unnecessary stress. C and T samples were then collected for a total of

34 dogs (from batches 2-4) and samples B and F for a sub-sample of 13 dogs (batch

4). Stress behaviors were coded for 33 animals (from batches 3 and 4) and cognitive

bias testing was conducted for a sub-sample of 16 dogs (from batches 3 and 4). Sub-

samples were balanced for prison and sex.

The sequence in which the TT, BET, and cognitive bias tests were

administered during the pre-program assessment was determined with safety and

standardization in mind. Since the dogs were unfamiliar to the research team,

conducting the TTs first allowed for an evaluation of their behavior around humans

and their response to handling in a relatively safe setting, before progressing to the

more intrusive handling required for BET and cognitive bias tests. Cognitive bias tests

were not performed on the first two batches, which were then tested with the BET

immediately following the TT. To maintain consistency across batches, the cognitive

bias test was conducted as the final assessment in the test battery. For the post-

program assessment, we changed the order of the tests to 1) cognitive bias, 2) BET,

and 3) TT, in order to prioritize those with the least potential to influence the

outcomes of subsequent tests (see Figure 1).

Data analysis

In addition to the live scoring of the dogs' performance in BETs and TTs, the

video recordings of the tests were also scored by four observers (using the same

scoring systems - Appendices 3 and 4). Three observers were blind to the stage (pre-

vs post-program) and to the goals of the study, and the fourth observer was non-blind

(assisted in data collection). Inter-observer reliability (for the five scorers) was

calculated for each exercise using intra-class correlation coefficient (ICC, see

Appendices 7a and 7b). The live and video scorings were combined into a final score

for each BET and TT item. The average scores from the video observers comprised

50% of the total weightage. The remaining 50% was allocated to live scores by MQ

and ACVC, both with extensive experience with dogs (details in Appendices 8a and

8b).

The BET measured variables across seven different categories, with one score

per category – 'Approach' (Item 1 - Accepting a friendly stranger), 'Greet' (Item 2 –

Waiting politely for petting), 'Leash' (Item 3 – Walking on a loose leash), 'Crowd'

(Item 4 – Walking through a crowd), 'Commands' (Item 5 – Sit, down and staying in

place), Recall (Item 6 – Coming when called) and 'Separation' (Item 7 – Supervised

separation). A Total BET score was also computed by summing the scores of the

different categories.

Although the TT was designed to measure variables across seven different

categories (Behavior in Kennel, Human Sociability, Behavior on Leash, Cognitive

Skills, Playfulness, Reactivity and Dog Sociability, see Appendix 4), only three

categories i.e. Human Sociability, Dog Sociability, and Playfulness emerged as robust

measures in previous studies (Valsecchi et al, 2011; Barnard et al, 2019), therefore we

limited the analysis to these same three categories. They were calculated as follows:

Human Sociability (HS) = Item 3 + Item 4 + Item 5 + Item 6 + Item 7 + Item 10a +

Item 10b + Item 10c

Dog Sociability (DS) = Item 16a + 16b

Playfulness (PLAY) = Item 13 + Item 14a + Item 14b

A Total TT score was also computed by summing the scores of all the items.

In the cognitive bias test, the dogs' latencies to reach the bowl in the

ambiguous location was adjusted for the latency to reach the training (positive and

negative) locations (Mendl et al, 2010; see Appendix 9), as follows:

$$adjusted\ latency$$
$$= \frac{(mean\ latency\ to\ ambiguous\ location - mean\ latency\ to\ positive\ location)}{(mean\ latency\ to\ negative\ location - mean\ latency\ to\ positive\ location)}\ x\ 100$$

The frequency of stress-related behaviors and the frequency of scans in the

different overall behavior states were input for analysis as recorded live by the

researchers, and salivary cortisol concentrations (in ng/mL) as reported by the

laboratory that ran the ELISAs (DNAtech).

Statistical analysis

Prior to the main modeling, we tested the adequacy of using cortisol levels

obtained on non-training days at the shelter in the course of the PBDTP (C1 to C3) as

control levels (covariates) for cortisol levels obtained on training days at the prisons

(T1 to T3). To this end, we tested if they differed from baseline cortisol levels at the

shelter before the program started (B1 and B2). We also tested if dogs' cortisol levels

at the shelter changed from before to after the PBDTP, by comparing cortisol levels at

the shelter before the program started (B1 and B2) with cortisol levels at the shelter

after the program ended (F1 and F2). Paired t-tests were performed on SAS

University Edition® (SAS Institute Inc., Cary, NC, USA) to compare baseline cortisol

levels measured during the week prior to the start of the PBDTP (average of B1 and

B2) with C1 to C3, and with post-program cortisol levels measured during the week

after the completion of the PBDTP (average of F1 and F2). Some saliva samples did

not have enough volume to allow laboratory analysis, which led to missing values.

When both B1 and B2 levels were available, average B was obtained and used for

comparison; when one of the B readings was missing, the available B level was

considered for the comparisons. The same approach was taken for F cortisol levels.

As a result, paired t-tests were done in sample subsets of distinct number of dogs (see

Appendix 10).

The response variables training cortisol concentration (T), as well as Total TT

Score, Human Sociability, Dog Sociability, Playfulness, Total BET Score, BET

categories 'Approach', 'Greet', 'Leash', 'Commands' and 'Separation', the number

of occurrences of the stress-related behavior Lip-licking, the number of scans in the

behavior states Tense, Excited, and Relaxed, and the adjusted latency to reach the

ambiguous bowl in the cognitive bias task were modeled as functions of the

independent variables time-point (3-level categorical fixed effect for stress-related

behaviors and behavioral states/ 2-level categorical fixed effect for remaining

response variables), prison (3-level categorical fixed effect), the interaction between

these two variables, dog age (discrete numeric fixed effect), and weight score

(discrete fixed effect)[1]. An adapted bi-directional stepwise selection of variables was

used, starting with the independent variable of interest (time-point), followed by

prison, the interaction between these two variables, dog age (discrete fixed effect),

and weight score. Variables reaching significance (i.e. $p \leq 0.05$) were maintained in the

model. Sex and neuter status were also tested as independent variables, but always

one at a time, due to the unbalanced nature of the dataset. The effect of neuter status

in the response variables were only tested for male dogs, as there was only one intact

female dog in the dataset. Control cortisol concentration (C) was used as a covariate

in the model explaining the variation of training cortisol concentration (T) as a

function of the aforementioned effects. Procedure GLIMMIX (for generalized linear

mixed modelling) was used on SAS University Edition® (SAS Institute Inc., Cary,

NC, USA), considering Dog ID as a random effect.

Multicollinearity was checked by regressing each independent variable on the

others in a numerical format. When model residuals failed to reach normality, a more

appropriate data distribution was considered in the models: log-normal for TT

---

[1] Information on dog weight was estimated from visual inspection and input in ranges of 5 kg by DTC team (see Appendix 1). For purposes of statistical analyses, we created a weight score from one to seven, where one was 5-10 kg, two was 10-15 kg, three was 15-20 kg, four was 20-25 kg, five was 25-30 kg, six was 30-35 kg, and seven was 40-45 kg.

category Human Sociability, behavior state Excited, training cortisol concentration

(T), and Poisson for the stress-related behavior Lip-licking, behavior states Relaxed

and Tense, and adjusted latency to reach the ambiguous bowl. To allow for the log-

normal distribution in the models, 0.1 was added to all data points of the response

variables to eliminate all zeros of the datasets. Adjusted latency was normalized prior

to the analysis, so that the maximum adjusted value was 1 and the minimum was 0, to

reach normality of model residuals.

The remaining stress-related behaviors Body Shake, Body Turn, Crouch,

Move Away, and Yawn, as well as the behavior state Low, had no occurrences in

64%, 77%, 83%, 85%, 88%, and 96% of the observations, respectively, rendering

impossible to explain their occurrences as a function of the variables of interest.

Therefore, the occurrences of these stress-related behaviors were dichotomized into

"1" (having occurred at least once) and "0" (no occurrences), and Fisher's exact test

was used through Procedure FREQ on SAS University Edition® (SAS Institute Inc.,

Cary, NC, USA) to evaluate the contingency between the 2-level categorical response

variable ("0" or "1") and Time-point for each of the participating prisons.

BET categories 'Crowd' and 'Recall' were dichotomized into being low ($\leq$

2.92, $\leq$ 1.8, respectively, thresholds based on data distribution) or high (higher than

thresholds), and Procedure GLIMMIX was used considering a binary data

distribution, with a Logit link function, to model probability of score being high as a

function of the aforementioned independent variables.

For all fitted models, least squares means were compared considering 95%

confidence interval (i.e. significance at $p<0.05$). A Tukey-Kramer adjustment was

applied to all multiple comparisons performed in this study.

Results

Dog sociability and education indicators

1. Basic Education Test (BET)

Figure 2 depicts the average scores for the BET categories pre- and post-

program. We found that post-program scores were significantly higher compared to

pre-program scores for all but one category: 'Approach' [$F_{(1,39)}$=10.53, p=0.002],

'Greet' [$F_{(1,39)}$=12.25, p=0.002], 'Leash' [$F_{(1,39)}$=5.39, p=0.025], 'Crowd'

[$F_{(1,439)}$=4.36, p=0.044], 'Commands' [$F_{(1,39)}$=54.63, p<0.001], and 'Separation'. The

only category for which there was no significant effect of stage was 'Recall'

[$F_{(1,39)}$=0.31, p=0.579]. Total BET score also increased significantly from before to

after the training program [Pre-program (Mean±SEM): 13.03±0.30; Post-program

(Mean±SEM): 15.39±0.40; $F_{(1,39)}$=69.95, p<0.001].

Figure 2. Average scores per category (expressed as a percentage of highest possible

score) of the Basic Education Test, before (black bars) and after (white bars) the

program (N=40). *stands for statistically significant differences (p<0.05).

2. Temperament Test (TT)

Figure 3 depicts the average scores for the categories Human Sociability, Dog

Sociability and Playfulness of the Temperament Test before and after the PBDTP. We

found that the scores for Human Sociability and Playfulness were significantly higher

after the PBDTP compared to before the PBDTP [Human Sociability: $F_{(1,41)}$=8.24,

p=0.006; Playfulness: $F_{(1,41)}$=7.01, p=0.012], but we found no significant effect of

stage (pre- vs post- program) on Dog Sociability [$F_{(1,39)}$=0.13, p=0.724]. Results also

showed that the Total TT score increased significantly from pre-program to post-

program [Pre-program: 48.16±1.46; Post-program: 51.07±1.32; $F_{(1,41)}$=16.45,

p=0.0002]. Moreover, Playfulness was significantly affected by sex, with males

exhibiting higher scores than females [$F_{(2,59)}$=4.22, p=0.046].

Figure 3. Average scores for the categories Human Sociability, Dog Sociability and

Playfulness of the Temperament Test (expressed as a percentage of highest possible

score), before (black bars) and after (white bars) the program (N=42). *stands for

statistically significant differences (p<0.05).

Dog stress and welfare indicators

3. Stress-related Behaviors

Lip-licking behavior was affected by time-point [$F_{(2,59)}=14.91$, $p<0.001$], in

that it was more frequent for the first time-point (Week 2) compared to the second

(Week 7; $t=4.25$, $p<0.001$) and third (Week 11; $t=4.89$, $p<0.001$) time-points, as

shown in Figure 4. No effects of time-point were found for the behaviors Yawn,

Move Away, Body Turn, Crouch, and Body Shake, in any of the participating prisons

(p>0.05). We also found that Lip-licking was affected by prison [$F_{(2,30)}=4.17$,

p=0.025], with dogs lip licking more frequently in Prison 2 compared to Prison 1 ($t=-$

2.62, p=0.035).

Figure 4. Average number of occurrences of stress-related behaviors during training sessions at the prisons across three time-points (Week 2, Week 7 and Week 11; N=33). Vertical bars show the SEM. *stands for statistically significant differences (p<0.05).

4. Overall Behavior States

We found an effect of time-point on the frequency of scans Relaxed

$[F_{(2,59)}=3.60,\ p=0.034]$ and Tense $[F_{(2,59)}=3.74,\ p=0.030]$. Pairwise comparisons

revealed that dogs were found more frequently in a Relaxed state and less frequently

in a Tense state for the second time-point compared to the first time-point (Relaxed:

t=-2.60, p=0.031; Tense: t=2.45,p=0.045, see Figure 5). No effect of time-point was

found for the behavior states Excited $[F_{(2,55)}=0.10,\ p=0.902]$ and Low $[F_{(2,59)}=3.64,$

p=0.032]. We also found an effect of neuter status for the behavior state Tense on

male dogs $[F_{(1,59)}=4.19,\ p=0.045]$. This effect was not tested for females because there

was only one intact female. A paired-samples t-test conducted for male dogs revealed

that intact males were found less frequently in Tense states than neutered males (t=-

2.05, p=0.045). Finally, we found an effect of weight score for the behavior state

Relaxed $[F_{(1,59)}=8.25,\ p=0.006]$, with the frequency of scans increasing as weight

score, thus dog weight, increased, and an effect of age on the behavior state Excited

$[F_{(1,61)}=4.30,\ p=0.042]$, with the frequency of scans increasing as dog age decreased.

Figure 5. Average frequency of scans in the different behavioral states during the

training sessions at the prisons in Week 2, Week 7 and Week 11 (N=33). The cases

where the bars do not reach the maximum (5) correspond to classification of the

behavioral state as 'Unknown' (see Appendix 6). *stands for statistically significant

differences.

5. Salivary Cortisol

No differences were found between pre-program baseline cortisol levels (B)

and control cortisol levels obtained on non-training days at the shelter in the course of

the PBDTP (C1 to C3) [B vs C1: $t(10)=1.85$, $p=0.094$; B vs C2: $t(11)=-0.35$, $p=0.732$;

B vs C3: $t(10)=2.10$, $p=0.062$]. Following these results, we concluded that C levels

could be used as appropriate controls against which to compare the cortisol levels at

the prisons in training days (T). Hence, C levels were included in the main cortisol

model as control values. Thereafter, we found no effects of control cortisol level C

[$F_{(1,33)}=0.20$, $p=0.660$], time-point [$F_{(2,33)}=1.41$, $p=0.258$] or prison [$F_{(2,33)}=0.00$,

p=0.996] in cortisol level T. Figure 3 shows the average cortisol concentrations for

C1-C3 and T1-T3. Moreover, we found no differences between pre-program baseline

cortisol levels (B) and post-program cortisol levels (F) [t(9)=1.33, p=0.217].

Figure 6. Average salivary cortisol concentration (ng/ml) on training days at the

prisons (T) and on non-training days at the shelter (C), across three time-points (Week

2, Week 7 and Week 11, N=34).

6. Cognitive Bias

The (adjusted) latency to reach the food bowl in the ambiguous locations did

not change significantly from pre- to post-program [Pre-program: 46.56±14.96 vs

post-program: 47.03±22.70; $F_{(1,15)}$=0.08, p=0.782].

Discussion

This paper presents a comprehensive study of how dogs are affected by

participating in a PBDTP, combining a battery of sociability and basic training

indicators with behavioral, physiological and cognitive indicators of welfare. Our

results show an overall improvement in the basic training and socialization and

handling skills of the dogs as a consequence of participating in the PBDTP, whereas

there was no indication of lasting stress.

Participation in the 12-week training program resulted in an improvement in

sociability towards humans, playful behavior, and in almost all education categories

that were tested. A significant improvement was seen even for BET categories where

the average performance was around 90% already at the start of the program, i.e.

where the dogs already displayed good basic training skills prior to participating in

the PBDTP.

Changing handlers has been found to reduce performance (Jamieson et al.,

2018), and inexperienced handlers are often not timing reinforcement accurately

(Browne et al., 2011), which can impact learning negatively. However, our study

found that the training was effective even though the dogs were trained by several

handlers who had no previous dog training experience and hence were initially

unskilled. The dogs participating in this program were from a shelter and were

available for adoption after completion of the program. Given that behavioral issues

are the most frequently cited reason for relinquishment or rehoming (Patronek et al.,

2022), there is reason to believe that the improvement in sociability and basic training

may help the dogs be successfully adopted i.e., integrate well and not be returned to

the shelter or cause extreme stress to their adopters.

Among all tested sociability and education categories, dogs did not improve

only on their scores for Dog Sociability (sociability towards other dogs) and Recall

(coming when called). Socializing dogs with conspecifics was not part of the designed

training program and, despite the fact that training sessions could have contributed to

improve dog sociability through desensitization and counterconditioning (sessions

were held in groups of 6 dogs, in a way that dogs were exposed to the presence of

others from a distance that was comfortable to them, and they were receiving food

and other potentially appetitive stimuli such as positive human interactions), it may be

the case that targeted training is needed for this specific behavior to improve (e.g.,

Stewart, 2016). Regarding Recall, this behavior was trained during the program for

distances up to the 2 meters as allowed by the leashes on which the dogs were kept

during the training sessions. The 10-meters distance dogs had to cover during testing

was never directly trained, and distance is advised by dog training professionals as an

important variable to work on when training a dog (e.g., Donaldson, 1996).

The second important factor here is the welfare of the participating dogs.

Participating in the program subjected the dogs to several unfamiliar and potentially

stressful experiences. Transport, at least if long (≥50 minutes) has been shown to

affect dog welfare negatively (de Carvalho et al., 2020). The participating dogs were

transported by car to and from the prisons twice a week, at a driving distance of

approximately 23 to 35km, and they were (to our knowledge) initially not used to

neither crating nor transport. The training itself was exclusively reward-based, thus

avoiding methods known to cause stress (Vieira de Castro et al, 2020), but the fact

that they took place in an initially unfamiliar place is a potential source of stress, as is

the handling by several different and initially completely inexperienced handlers, and

the proximity to other dogs. Despite these potentially stressful factors, there was no

evidence that the dogs were overall negatively affected. Participating in the training

program did not affect levels of salivary cortisol, nor performance in the cognitive

bias test of affective state, and behaviors indicative of stress were generally rare

during training sessions.

At the first time-point (week 2 of training in the prisons), lip licks were more

frequent and dogs were often seen in a tense state and less often in a relaxed state

compared to later time-points. The reduction in stress behaviors between the first

(week 2 of training) and the second (week 7 of training) time-points shows that the

dogs did not take long to habituate, and that the combination of transport and training

did not result in long-lasting stress.

As observed by Cobb et al. (2016), there is a large amount of intra-individual

and inter-individual variability and external variables that could influence salivary

cortisol concentration. In line with this, in the present study, two dogs had much

higher C1 cortisol values than the remaining (11) animals, which resulted in the high

variability observed for this time-point. The reasons underlying the elevated values

for these animals are beyond our knowledge, as no demographic factor including sex,

age, weight, neuter status or time spent in the kennel before study commencement

differentiates these animals from the remaining sample. We have tried to control for

these variations in our experimental design by comparing cortisol concentration only

intra-individual, and by standardizing collection timing and conditions as much as

possible. However, the sample size for this welfare measure turned out to be much

smaller than initially planned. Due to several samples not having enough saliva

volume for laboratory analysis and because we used a repeated-measures design, we

ended up with only 7 to 14 subjects per comparison, despite having collected saliva

samples of 34 animals. This small sample size thus prevents any strong conclusion

from being drawn. However, the analysis of stress-related behaviors during training

sessions, for which the sample size was only one less than the intended, suggest that

participating in the PBDTP was not stressful for the animals. In fact, of the six

behaviors analyzed, five were extremely rare (absent in 64-96% of the observations),

and these same behaviors have been shown to occur frequently in a stressful training

context (Vieira de Castro et al, 2020).

The present findings are in line with Leonardi et al (2024) and Hennessy et al

(2006), in that all the three studies found improvements in dog training performance

after participating in a PBDTP, and in line with Hennessy et al (2006) in that both

studies found no changes in cortisol levels from before to after the program. Leonardi

et al (2024) reported improvements in dog welfare, with dogs showing more relaxed

behaviors in their kennels after the PBDTP. In the present study, we also found dogs

to become more relaxed and less tense across the program. However, our assessments

were performed during training sessions in the prisons and not in the shelter as in

Leonardi et al (2024), and hence are not directly comparable. Moreover, in Leonardi

et al (2024), shelter staff rated dogs as more playful, friendly and relaxed, and less

excitable, frustrated, noisy, reactive or vigilant after participating in the PBDTP.

Despite further suggesting that dog welfare improved with the PBDTP, these results

need to be interpreted with caution because they rely on subjective ratings by shelter

staff who were not blind to the treatment (i.e., participation in the PBDTP).

Importantly, the present study only used objective and quantitative welfare indicators

and evaluated welfare both within and outside the training context, thus providing a

more robust assessment of the welfare of dogs participating in PBDTPs. Taken

together, data available up to date suggest that participation in PBDTPs has, at least,

no negative impact on dog welfare.

The present study provides comprehensive data on dog behavior and welfare

as a result of participating in a PBDTP. The fact that it included a sizeable sample

(42 dogs) and three different prison establishments with different sets of inmates and

supervising dog trainers are further strengths of the study. In the vast majority of

PBDTPs, the dogs live full-time at the prisons, as opposed to living in shelters or

other facilities and being transported to the prisons for the training sessions as in the

case of Pelos2. Besides animal housing, PBDTPs vary in the number of inmates with

whom each dog is paired, number of participants, criteria for participation, length of

participation, or the amount of time per day that the participants spend with the dogs

(Cooke, 2019). It is still unclear how different aspects of PBDTPs may affect their

efficacy and impact on animal welfare, and hence the generalization of the present

results is somewhat limited. Behavior during training sessions could only be analyzed

through direct observations, as video recordings would have raised issues with

protecting the identity of a vulnerable population of human participants. To minimize

the risk of bias and maximize the reliability of the direct observations, we made an

effort to train and calibrate the two observers before they started to collect data. As

regards the tests for dog sociability and basic training, the exercises in the Basic

Education Test were originally designed to be performed by the dog's owner or

regular handler, and the fact the dogs here were tested with research personnel may

have affected their performance.

The improvement in basic training and socialization and handling skills, and

the absence of indicators that the participation was stressful for the dogs suggest that

the program can be deemed a success for the long-term benefits it is likely to bring to

the dogs and their future adopters. In future research, it would be important to

evaluate to what extent the improvement in dog behavior actually translates into an

increase in successful adoptions.

Acknowledgements

We would like to thank the entire DTC team for making this study possible,

especially Luca Conde and Inês Vieira, for being so enthusiastic about this

collaboration since the beginning. We are also deeply grateful to the DTC trainers

involved in the Pelos2 program, particularly Rita Pinto, Marco Saraiva, André

Carneiro and José Xavier, as well as to the DTC psychologist Silvia Sousa, who

facilitated the collection of live behavior data at the prisons, ensuring our research did

not interfere with their training schedules.

Our gratitude goes to the staff at the prison establishments of Paços de Ferreira, Vale

do Sousa, and Santa Cruz do Bispo, as well as the shelter staff at PATA, for their

cooperation and flexibility, which greatly contributed to the successful completion of

this study.

We extend our heartfelt thanks to Sofia Morais and Patricia Fachada for their

invaluable assistance with data collection at the shelter. We would also like to express

our appreciation to the students from the Abel Salazar Biomedical Sciences Institute

at the University of Porto and the University of Edinburgh, whose voluntary efforts in

scoring videos strengthened the robustness of our results.

Finally, we are particularly grateful to Dr. Lydia Hopper, Dr. Sarah-Elizabeth

Byosiere, and the two anonymous reviewers at Animal Behavior and Cognition,

whose constructive feedback greatly improved our study protocol. We also thank Ana

Maria Valentim for her valuable input on our final manuscript.

Author Contributions: Parizad Baria-Unwalla: Conceptualization, Data curation,

Formal analysis, Investigation, Methodology, Validation, Visualization, Writing –

original draft, Writing – review & editing; **Gabriela Munhoz Morello:** Formal

analysis, Writing – review & editing; **Maria Queiroz:** Conceptualization, Data

curation, Methodology; **I. Anna S. Olsson:** Conceptualization, Funding acquisition,

Project administration, Resources, Supervision, Validation, Writing – original draft;

Ana Catarina Vieira de Castro: Conceptualization, Data curation, Formal analysis,

Investigation, Methodology, Project administration, Supervision, Validation,

Visualization, Writing – original draft.

References

AKC, American Kennel Club. (2023, July 25). *Canine Good Citizen Test Items*.

AKC. https://www.akc.org/products-services/training-programs/canine-good-

citizen/canine-good-citizen-test-items/

Barnard, S., Kennedy, D., Watson, R., Valsecchi, P. & Arnott, G. (2019). Revisiting a

previously validated temperament test in shelter dogs, including an

examination of the use of fake model dogs to assess conspecific sociability.

Animals, *9*(10), 835.

Britton, D. M. & Button, A. (2005). Prison pups: Assessing the effects of dog training

programs in correctional facilities. *Journal of Family Social Work*, *9*(4), 79–

95.

Broom D.M., Johnson K.G. (2019). *Stress and Animal Welfare.* 2nd ed. Springer;

Berlin, Germany.

Browne, C. M., Starkey, N. J., Foster, T. M. & McEwan, J. S. (2011). Timing of

reinforcement during dog training. In *2nd Canine Science Forum: Clinical*

Applications and Research, 6(1), 58-59.

Cobb, M. L., Iskandarani, K., Chinchilli, V. M., & Dreschel, N. A. (2016). A

687   systematic review and meta-analysis of salivary cortisol measurement in

688   domestic canines. *Domestic Animal Endocrinology, 57*, 31-42.

689   Cooke, B. J. (2019). Exploring Types of Programs: Dog Rescue, Rehabilitation and

690   Training. In *Prison dog programs: Renewal and rehabilitation in correctional*

691   *facilities* (pp. 37-51). Cham: Springer International Publishing.

692   Cooper, J. J., Cracknell, N., Hardiman, J., Wright, H., & Mills, D. (2014). The

693   welfare consequences and efficacy of training pet dogs with remote electronic

694   training collars in comparison to reward based training. *PloS one*, *9*(9),

695   102722.

696   Currie, N. (2008). *A case study of incarcerated males participating in a canine*

697   *training program* (Unpublished doctoral dissertation). Kansas State

698   University, Manhattan.

699   Davis, K. (2007). *Perspectives of youth in an animal-centered correctional vocational*

700   *program: A qualitative evaluation of Project Pooch* (Unpublished research

701   monograph). Portland State University, OR.

702   Deaton, C. (2005). Humanizing Prisons with Animals: A Closer Look at "Cell Dogs"

703   and Horse Programs in Correctional Institutions. *Journal of Correctional*

704   *Education*, *56,* 46–62.

705   de Carvalho, I. R., Nunes, T., de Sousa, L. & Almeida, V. (2020). The combined use

706   of salivary cortisol concentrations, heart rate, and respiratory rate for the

707   welfare assessment of dogs involved in AAI programs. *Journal of Veterinary*

708   *Behavior*, *36,* 26-33.

709   Dudley, E. S., Schiml, P. A. & Hennessy, M. B. (2015). Effects of repeated petting

710   sessions on leukocyte counts, intestinal parasite prevalence, and plasma

cortisol concentration of dogs housed in a county animal shelter. *Journal of*

the American Veterinary Medical Association, 247(11), 1289-1298.

Duindam, H.M., Asscher, J. J., Hoeve, M., Stams, G. J. J., & Creemers, H.E. (2020).

Are we barking up the right tree? A meta-analysis on the effectiveness of

prison-based dog programs. *Criminal justice and behavior*, *47*(6), 749-767.

Donaldson, J. (1996). *The Culture Clash.* James and Kenneth Publishers.

Furst, G. (2006). Prison-based animal programs. *The Prison Journal*, *86,* 407-430.

Gilger, C. G. (2007). *Cell dogs: No effect of dog training programs on prisoners' self-*

efficacy (Unpublished doctoral dissertation). Emporia State University, KS.

Grommon, E., Carson, D. C., & Kenney, L. (2018). An experimental trial of a dog-

training program in a juvenile detention center. *Journal of Experimental*

Criminology, *41*(1), 62–80.

Hennessy, M.B., Morris, A., Linden, F. (2006). Evaluation of the effects of a

socialization program in a prison on behavior and pituitary adrenal hormone

levels of shelter dogs. *Applied Animal Behavior Science*, *99*, 157–171.

Humby, L. & Barclay, E. (2018). Pawsitive solutions: An overview of prison dog

programs in Australia. *The Prison Journal*, *98*(5), 580-603.

Jamieson, L. T. J., Baxter, G. S., & Murray, P. J. (2018). You are not my handler!

Impact of changing handlers on dogs' behaviors and detection

performance. *Animals*, *8*(10), 176.

Jasperson, R. A. (2013). An animal-assisted therapy intervention with female inmates.

Anthrozoös, *26*(1), 135–145.

Johnson, A. & Bruneau L. (2019). Caring for and About Dogs: Animal Welfare

Considerations. In: Jalongo M. (eds) *Prison Dog Programs.* Springer, Cham.

Karkdijk, E. M., Duindam, H. M., Deković, M., Creemers, H. E., & Asscher, J. J.

(2022). A Friend in Prison: Human-Animal Bond, Stress and Self-Esteem of

Detained Juveniles in Dutch Cell Dogs. *Animals*, *12*(5), 646.

Koda, N., Miyaji, Y., Kuniyoshi, M., Adachi, Y., Watababe, G., Miyaji, C., &

Yamada, K. (2015). Effects of a dog-assisted program in a Japanese prison.

Asian Journal of Criminology, 10(3), 193-208.

Mendl, M., Brooks, J., Basse, C., Burman, O., Paul, E., Blackwell, E., Casey, R.

(2010). Dogs showing separation-related behavior exhibit a 'pessimistic'

cognitive bias'. *Current Biology, 20*, R839—R840.

Leonardi, R.J., Vick, S.-J., Buchanan-Smith, H.M. (2024) Participation in a prison-

based training programme is beneficial for rescue dogs. *Animals, 14*, 1530.

Mercer, J., Gibson, K. & Clayton, D. (2015). The therapeutic potential of a prison-

based animal programme in the UK. *The Journal of Forensic Practice, 17*(1),

43-54.

Nimer, J., & Lundahl, B. (2007). Animal assisted therapy: A meta-analysis.

Anthrozoös, *20*, 225-238.

Patronek, G. J., Bradley, J., & Arps, E. (2022). Saving normal: A new look at

behavioral incompatibilities and dog relinquishment to shelters. *Journal of*

Veterinary Behavior, 49, 36-45.

Sandri, M., Colussi, A., Perrotta, M. G., & Stefanon, B. (2015). Salivary cortisol

concentration in healthy dogs is affected by size, sex, and housing

context. *Journal of Veterinary Behavior, 10*(4), 302-306.

Shiverdecker, M. D., Schiml, P. A., & Hennessy, M. B. (2013). Human interaction
moderates plasma cortisol and behavioral responses of dogs to shelter
housing. *Physiology & behavior*, *109*, 75-79.

Stewart, G. Behavior Adjustment Training 2.0. (2016). *New Practical Techniques for*
Fear, Frustration, and Aggression in Dogs. Dogwise Publishing.

Strimple, E. O. (2003). A history of prison inmate-animal interaction programs.
American Behavioral Scientist, 47, 70-78.

Valsecchi, P., Barnard, S., Stefanini, C., & Normando, S. (2011). Temperament test
for re-homed dogs validated through direct behavioral observation in shelter
and home environment. *Journal of Veterinary Behavior, 6*(3), 161-177.

Vieira de Castro, A. C., Fuchs, D., Morello, G. M., Pastur, S., de Sousa, L., & Olsson,
I. A. S. (2020). Does training method matter? Evidence for the negative
impact of aversive-based methods on companion dog welfare. *PLoS*
One, *15*(12), 0225023.