A glial cell derived pathway directs regenerating optic nerve axons toward

the optic chiasm

Running Title: Glia direct regenerating RGC axons

Beth M. Harvey[1*], Melissa Baxter[1], Alexis M. Garcia[1] and Michael Granato[1*]

Department of Cell and Developmental Biology, Perelman School of Medicine, University of

Pennsylvania, Philadelphia, Pennsylvania 19104

* Co-Corresponding authors

email: bharvey@pennmedicine.upenn.edu

email: granatom@pennmedicine.upenn.edu

Abstract

After optic nerve injury, several retinal ganglion cell (RGC) intrinsic signaling pathways have been shown to enhance RGC survival and RGC axonal growth. In contrast, few extrinsic cues have been identified that guide regenerating RGC axons toward and across the optic chiasm. Here, we use live-cell imaging in larval zebrafish to show that regrowing RGC axons initiate growth toward the midline and extend along a trajectory similar to their original projection. From a candidate genetic screen, we identify the glycosyltransferase Lh3 to be required to direct regrowing RGC axons toward the midline during active regeneration. Moreover, we show that transgenic *lh3* expression in *sox10*+ presumptive *olig2*+ oligodendrocytes located near the optic chiasm restores directed axonal growth in *lh3* mutants. Finally, we find that mutants in *collagen 18a1 (col18a1),* a putative Lh3 substrate, display RGC axonal misguidance phenotypes similar to *lh3* mutants, suggesting that *lh3* may act through *col18a1* during regeneration. Combined, these data identify *lh3* as part of a glial derived molecular pathway critical for guiding *in vivo* regenerating RGC axons toward and across the optic chiasm.

Introduction

As a component of the central nervous system, retinal ganglion cell (RGC) axons within the mammalian optic nerve display a poor regenerative capacity. Multiple factors contribute to the minimal regeneration in mammals, including massive injury-induced RGC death and overall limited axonal regrowth (Villegas-Pérez *et al.*, 1993; Berkelaar *et al.*, 1994; Goldberg *et al.*, 2002). Several studies have identified a number of RGC intrinsic signaling pathways that enhance RGC survival and increase long range axonal growth after injury. For example, deletion of the transcriptional repressor KLF4 in RGCs increases axonal regrowth (Moore *et al.*, 2009). Similarly, activation of inflammatory responses within the retina through injection of Zymosan and cAMP enhances cell survival and axonal regrowth (Leon *et al.*, 2000; Goldberg *et al.*, 2002; Yin *et al.*, 2009). Moreover, deleting intrinsic factors in RGCs such as PTEN and TSC1, which function as negative regulators of the mTOR pathway, improves both RGC cell survival and axonal growth (Park *et al.*, 2008; Smith *et al.*, 2009; Norsworthy *et al.*, 2017). However, enhancing RGC axonal growth by manipulating RGC intrinsic programs frequently results in axonal misguidance as axons project inappropriately from the optic tract before and at the optic chiasm during the initial stages of regeneration (Kurimoto *et al.*, 2010; Luo *et al.*, 2013; Vincent Pernet *et al.*, 2013; V. Pernet *et al.*, 2013; Berry, Ahmed and Logan, 2019). This demonstrates a need to identify molecular and cellular mechanisms that guide RGC axons as they regrow toward and across the optic chiasm during regeneration.

Following a Central Nervous System (CNS) injury in mammals, many non-neuronal cells create an extracellular environment that is largely inhibitory to axonal regrowth. The cellular and molecular barriers that form within a lesion site have long been attributed to an influx of inflammatory cells and the upregulation of inhibitory molecules not conducive for robust regeneration (Yiu and He, 2006; Fitch and Silver, 2008). Additional studies in mice have since revealed roles for microglia and oligodendrocytes in functional recovery and remyelination from

CNS injury (Wang *et al.*, 2020; Mendonça *et al.*, 2021; Brennan *et al.*, 2022). Yet due to the

dominance of factors inhibiting axonal regrowth, the cellular and molecular mechanisms that

guide regrowing RGC axons following axonal injury are largely unknown. Recently, spontaneous

CNS regeneration has been described in two mammalian species, the naked mole rat and the

spiny mouse, demonstrating that pro-regenerative environments that support robust CNS

regeneration are not unique to the CNS of non-mammalian vertebrates (Park *et al.*, 2017;

Nogueira-Rodrigues *et al.*, 2022). Therefore, model organisms that exhibit spontaneous robust

CNS axonal regeneration, without an overwhelmingly inhibitory extracellular environment,

provide unique opportunities to identify mechanisms that promote CNS regeneration.

We recently developed a rapid and robust optic nerve transection assay in larval

zebrafish to take full advantage of its regenerative capabilities and optical transparency to study

functional optic nerve regeneration *in vivo* (Harvey, Baxter and Granato, 2019, 2023). At just 5

days post fertilization (dpf), larval zebrafish already possess a functional visual system driving

several visually guided behaviors, such as prey capture (Brockerhoff *et al.*, 1995; Easter and

Nicola, 1996; Gahtan, Tanger and Baier, 2005). Following complete transections of the RGC

axons in 5 dpf larvae, we observe axonal degeneration, subsequent initiation of regrowth and

reinnervation of the optic tecta by 72 hours post transection (hpt) (Harvey, Baxter and Granato,

2019). We have also shown that RGC axonal regeneration in larval zebrafish occurs

independent of RGC death or neurogenesis, and that visual function is restored by 8 days post

transection (Harvey, Baxter and Granato, 2019). Using this optic nerve transection assay, we

performed a candidate genetic screen and identify Lh3, a collagen-modifying

glycosyltransferase, to be critical for directing axonal regrowth toward the optic chiasm during

optic nerve regeneration. We show that following optic nerve transection, *lh3* expression is

upregulated near the site of axonal injury and that Lh3 functions during the time when active

regeneration occurs, specifically during the early stages of regeneration as RGC axons are

growing toward the midline. We also show that Lh3 expression in oligodendrocytes is required

to direct early regrowing RGC axons across the midline along their original path. Furthermore,

we show that mutants in *collagen 18a1 (col18a1),* a putative Lh3 substrate, display RGC axonal

misguidance phenotypes similar to *lh3* mutants, indicating at a glial-derived molecular pathway

in which Lh3 acts through Collagen18a1 to direct RGC axons. Combined, our results

demonstrate a critical pro-regenerative *in vivo* role for oligodendrocytes during early optic nerve

regeneration to direct axon regrowth toward and across the midline.

Results

Lh3 is required for growth across midline during regeneration

To identify molecular mechanisms that guide regrowing RGC axons across the optic

chiasm during optic nerve regeneration, we performed a candidate genetic screen consisting of

35 genes that are expressed in the visual system or have known functions in axon growth and

guidance. Specifically, we used a sharpened tungsten needle to transect the optic nerves of

genetic mutants and their siblings at 5 dpf and assessed at 72 hpt whether RGC axons had

regrown across the optic chiasm to reinnervate the optic tectum. One of the top hits from this

screen was the gene *lh3*, which encodes a multidomain glycosyltransferase that post-

translationally modifies collagen proteins (Heikkinen *et al.*, 2000; C. Wang *et al.*, 2002). To

bypass the requirement of *lh3* during development (Zeller and Granato, 1999; Taler *et al.*,

2019), we took advantage of a previously validated transgene *Tg(hsp70:lh3-myc)*, which

expresses a Myc-tagged Lh3 under the control of the hsp70 heat-shock-activated promoter

(Isaacman-Beck *et al.*, 2015). At 5 dpf, uninjured RGC axons in both *lh3* siblings and mutants

exit the eye and project across the optic chiasm to terminate in the optic tecta (Fig. 1A-D).

Following optic nerve transection, axons in *lh3* siblings grow across the midline and reinnervate

the optic tecta (Fig 1E,G,I, K-L). In contrast, transected optic nerve axons in *lh3* mutants fail to

cross the midline and instead project along aberrant trajectories, often along anterior and

posterior paths or also dorsally or ventrally (Fig 1F,H,J,K-L, magenta arrowheads). These results

reflect a functional role for Lh3 during optic nerve regeneration in directing RGC axons towards

and across the midline at the optic chiasm.

Lh3 **expression is upregulated in the optic chiasm during regeneration**

We next asked whether Lh3 is required during the process of optic nerve regeneration.

For this, we took advantage of the conditional *Tg(hsp70:lh3-myc)* transgene (Isaacman-Beck *et*

al., 2015). We induced Lh3 expression in 5 dpf larvae 4-5 hours prior to performing the optic

nerve transections, and then assessed axonal regrowth across the optic chiasm and

reinnervation of the optic tecta at 72 hpt (Fig. 2A). RGC axons in *lh3* siblings that did or did not

receive a heat shock regrew across the midline and innervated the optic tecta (Fig. 2B,D,F). In

contrast, *lh3* mutants without a heat shock display aberrant axonal growth, while heat shock

induced Lh3 expression in *lh3* mutants restores optic nerve axonal regrowth across the midline

(Fig. 2C,E-F). These data demonstrate that Lh3 acts during the process of optic nerve

regeneration.

We then sought to determine in which cell-type Lh3 was expressed during optic nerve

regeneration. We used *in situ* hybridization chain reaction to detect *lh3* mRNA expression prior

to and after optic nerve transections. Prior to transections, we failed to detect *lh3* mRNA

expression in retinas nor in the region of the optic chiasm (Fig. 2G-I; n = 17 larvae). Similarly, we

failed to detect *lh3* mRNA expression in RGC neurons in the retina after transection (Fig. 2G-H;

n = 41 larvae). In contrast, we detected upregulated *lh3* mRNA expression around the

transected proximal stump of RGC axons and near the midline during early regeneration

between 2 and 6 hpt (Fig. 2G,I; n = 30/41 larvae), demonstrating that *lh3* mRNA is upregulated

following optic nerve transection in the region of the optic chiasm.

Lh3 directs regrowing RGC axonal growth toward their original trajectory

Following optic nerve transections in wild-type larvae, we have previously observed new

RGC axons emerging from the proximal stump of transected optic nerves by 24 hpt and exhibit

robust growth across the optic chiasm by 48 hpt (Harvey, Baxter and Granato, 2019). To

determine whether Lh3 continuously directs regenerating RGC axons once they emerge from

the optic nerve stump, we used live cell imaging and examined the dynamics of regenerating

RGC axons as they approach and cross the optic chiasm, which has not previously been

reported. For simplicity, we removed one eye from 4 dpf larvae to only visualize the regrowth

from a single retina per larva (Fig 3A). Following transections on 5 dpf *lh3* sibling and mutant

larvae, we observe that the portion of axons distal to the transection site undergoes Wallerian

degeneration, as has been observed for other nerves in zebrafish (Fig. 3B-D,G-I,

Supplementary Movies 1 and 2, brackets) (Waller, 1851; Martin *et al.*, 2010; Rosenberg *et al.*,

2012; Bremer *et al.*, 2019). In *lh3* siblings, some axons emerge from the proximal stump of RGC

axons in a range of directions, often extending and retracting rapidly (Fig. 3B-C, Supplementary

Movie 1). By 16 hpt, we observe that a small group of axon fascicles grows toward the midline

and rapidly extends across the midline (Fig. 3D-E, Supplementary Movie 1, white arrowheads).

Over the next few hours, additional fascicles of axons are observed following the growth of the

axons that have crossed the midline (Fig. 3F, Supplementary Movie 1) revealing *in vivo* axonal

dynamics during the early stages of optic nerve regeneration. Axons in *lh3* mutants emerge at

similar timepoints to *lh3* siblings from the proximal stump (Fig 3G-H, Supplementary Movie 2).

However, unlike the *lh3* siblings, axonal growth does not persist across the midline and axons

often grow anteriorly and posteriorly or dorsally and ventrally around the eye (Fig. 3I-K,

Supplementary Movie 2, magenta arrowheads), strongly suggesting that during the early

process of optic nerve regeneration, Lh3 directs the initial regrowth of RGC axons toward the

midline.

We next sought to identify the mechanisms by which Lh3 directs regrowing RGC axons

toward and across the optic chiasm. We first asked whether Lh3 is critical for the initiation of

axonal growth extending from the proximal stump of transected RGC axons, and consequently

whether optic nerve regrowth was delayed in *lh3* mutants. We found no significant difference in

the timing of initiation of axonal regrowth post transection (Fig 4A-B). To further define the

trajectories that RGC axons take as they grow across the optic chiasm, we measured the angle

between distinct axon fascicles and the original projection path before transection (Fig 4C; see

Methods for more details). In optic nerves in *lh3* siblings that regrew across the midline, we

observe that most fascicles grow very close to their original projection path, and few fascicles

strayed from the pre-injury path (Fig 4D). In contrast, in nerves that failed to cross the midline in

lh3 mutants, we found that the paths of axon projections were more random and were not

directed along their original path (Fig 4D). These data strongly suggest that *lh3* is required to

direct RGC axons along their regenerative path, which is very similar to their pre-transection

path, towards and across the optic chiasm.

Glial expression of Lh3 restores RGC axonal growth across the midline

Given that *lh3* mRNA expression is upregulated in area of the optic chiasm (Fig. 2I) and

that *lh3* is required *in vivo* to direct regenerating RGC axons along their original projection path

(Fig. 2F, 3, 4D), we hypothesized that the relevant cell type critical for *lh3* mediated optic nerve

regeneration would be located in the region of the optic chiasm. Transcriptomics studies in mice

have found *lh3* to be expressed most greatly in microglia (Zhang *et al.*, 2014). Moreover,

macrophages and microglia have been shown to clear debris from injured axons and can

promote axonal growth and remyelination during CNS regeneration (Rosenberg *et al.*, 2012;

Wang *et al.*, 2020; Brennan *et al.*, 2022). Therefore, we first focused on macrophages and

microglia as a potential source of Lh3 activity during optic nerve regeneration. In untransected

larvae, we observe sporadic *mpeg+* cells near the optic chiasm (Fig 5A). At 24 hpt, we observe

a greater concentration of *mpeg+* cells surrounding the proximal stump of RGC axons and the

new axonal growth (Fig. 5A; n = 11 larvae). We next used a transgenic rescue approach to test

whether expression of *lh3* in macrophages/microglia is sufficient to restore axonal regrowth

across the optic chiasm in *lh3* mutants. We found that Lh3 expression from a *Tg(mpeg:lh3-myc)*

transgene in *lh3* mutants resulted in an insignificant difference between *lh3* siblings and

mutants, but also no different from animals without the *Tg(mpeg:lh3-myc)* transgene (Fig 5B).

Since RGC axons regrow very closely to their original path, we next sought to examine

oligodendrocytes associated with the optic nerve (Brösamle and Halpern, 2002). Using both a

Tg(olig2:dsred) transgenic line labeling oligodendrocyte lineage cells (Fig 5C, n= 5 untransected

larvae, n = 21 transected larvae) as well as *in situ* hybridization chain reactions for *mbpa* mRNA

in myelinating oligodendrocytes (Fig 5D, n= 6 untransected larvae, n = 5 transected larvae), we

observed oligodendrocytes in the optic chiasm and associated with RGC axons before and after

transection (Kucenas *et al.*, 2008; Ackerman and Monk, 2016). We then tested whether

expression of *lh3* in oligodendrocytes would rescue axonal regrowth across the optic chiasm in

lh3 mutants. *Sox10*, like *olig2,* is a transcription factor that mediates oligodendrocyte lineage

specificity and is maintained in all oligodendrocyte lineage cells (Bergles and Richardson, 2015;

Ackerman and Monk, 2016). We find that *lh3* expression from a *Tg(sox10:lh3-mkate)* transgene

restores RGC regrowth across the midline in otherwise mutant *lh3* larvae (Fig 5E). Furthermore,

we detect *lh3* mRNA expression that overlaps with transgenic labeling of oligodendrocyte cells,

demonstrating that *lh3* is upregulated in oligodendrocytes surrounding the proximal stump of

transected RGC axons and along axonal projection path across the midline during optic nerve

regeneration (Fig. 5F; n = 15/21). Altogether, these findings demonstrate that *lh3* expression in

oligodendrocytes near the proximal stump of transected RGC axons directs early regrowing

RGC axons toward the midline along their original path to reinnervate the optic tecta.

Col18a1 is also critical for directing regenerating RGC axons

We further sought to identify the relevant Lh3 substrate during optic nerve regeneration.

Given the well documented role of Lh3 in post-translationally modifying collagen proteins

(Heikkinen *et al.*, 2000; C. Wang *et al.*, 2002), we focused on collagen substrates. As part of our

initial candidate genetic screen, we examined *col4a5*, which is required for proper RGC axon

targeting of tectal layers and also functions to scaffold guidance cues during both the

development of the visual system and during motor nerve regeneration (Xiao and Baier, 2007;

Xiao *et al.*, 2011; Isaacman-Beck *et al.*, 2015; Murphy, Isaacman-Beck and Granato, 2022). We

also tested *col7a1,* which has been shown to be expressed in the eye (Wullink *et al.*, 2018). We

ultimately found that RGC axon regeneration toward and across the midline is unaffected in

col4a5 and *col7a1* mutants (Fig. S1). We then examined *col18a1*, a collagen associated with

Knobloch syndrome in humans that is characterized by eye abnormalities (Fukai *et al.*, 2002;

Suzuki *et al.*, 2002). Col18a1 has several putative Lh3 sites within X-Lys-Gly- motifs and has

been shown *in vitro* to bind to Lh3 (Risteli *et al.*, 2004; Salo and Myllyharju, 2021). First, we

used *in situ* hybridization chain reaction to detect *col18a1* mRNA expression and observe an

upregulation of *col18a1* mRNA at 12 hpt in the optic chiasm, similar to the expression pattern of

lh3 following optic nerve transection (Fig. 6A; n = 13/14 larvae). We then tested *col18a1* mutants

in our optic nerve transection assay. Prior to transection, RGC axonal projections were

indistinguishable between *col18a1* siblings and mutants (Fig. 6B-E). In contrast, at 72 hpt we

observed regrowing RGC axons projecting away from the CNS midline, similar to what we

observe in *lh3* mutants where regenerating RGC axons grow along aberrant paths (Fig 6F-K).

This demonstrates that during regeneration *col18a1* is critical for directing regenerating RGC

axons toward the midline and suggests that glial-derived Lh3 acts, as least in part, through

Col18a1 during optic nerve regeneration.

<u>Discussion</u>

Regenerating axons often encounter multiple choice points along their paths that they

must navigate properly to robustly reinnervate their original targets. Previous studies have

shown that PTEN inactivation in RGC neurons strongly promotes RGC regrowth following an

optic nerve injury, but that these axons display axonal misguidance at the optic chiasm, proving

the optic chiasm to be one such choice point (Kurimoto *et al.*, 2010; Luo *et al.*, 2013; Vincent

Pernet *et al.*, 2013; V. Pernet *et al.*, 2013; Berry, Ahmed and Logan, 2019). Here, we identify a

mechanism critical for guiding RGC axons toward and across the optic chiasm during optic

nerve regeneration. We find that *lh3* is required during regeneration to direct regenerating RGC

axons toward and across the midline. We find that following optic nerve transection, expression

of *lh3* and its presumptive substrate *col18a1* are upregulated in cells along the regenerative

path near the optic chiasm. Using live cell imaging, we demonstrate that *lh3* is required to direct

regrowing RGC axons along a regenerative trajectory very similar to their original path.

Together, these results provide compelling evidence for a RGC extrinsic molecular pathway that

guides optic axons during regeneration.

Live cell imaging reveals axonal dynamics of early optic nerve regeneration

To date, most optic nerve regeneration studies in adult rodents and fish monitor RGC

axonal regrowth with static time point analyses (Horder, 1971; Murray, 1976; V. Pernet *et al.*,

2013; Yin and Benowitz, 2018). In contrast, peripheral nervous system regeneration, as well as

spinal cord regeneration have been visualized in live larval zebrafish using *in vivo* imaging

studies (Bhatt *et al.*, 2004; Rosenberg *et al.*, 2012; Ceci *et al.*, 2014; Lewis and Kucenas, 2014;

Isaacman-Beck *et al.*, 2015; Bremer *et al.*, 2019). To visualize RGC axonal dynamics during

optic nerve regeneration, we previously established an optic nerve transection assay in larval

zebrafish (Harvey, Baxter and Granato, 2019, 2023). Here, we perform live cell imaging of RGC

axons initiating regrowth and regenerating across the optic chiasm *in vivo*. These timelapses

reveal two intriguing insights into mechanisms that may mediate robust CNS regeneration in

zebrafish. First, we observe that axons distal to the transection site undergo Wallerian

degeneration (Fig. 3B-D,G-I, Movie S1 and Movie S2, brackets) (Waller, 1851). While a

hallmark of the mammalian CNS is a slower rate of axonal fragmentation following axonal injury,

here we observe complete degeneration of the distal axons by about 15 hpt (Vargas and Barres,

2007). While we did not examine whether persistence of degenerating debris would repel axonal

growth, we do observe new axons reaching the midline around the same time axonal debris is

cleared from that region (Fig. 3D). Axonal debris and associated myelin proteins are known to

inhibit axonal regeneration (Vargas and Barres, 2007). But here, we find that Wallerian

degeneration does not appear to be delayed in *lh3* mutants, and we therefore would not

hypothesize axonal debris to contribute to the misguidance of RGC axons in *lh3* mutants.

Second, our live cell imaging reveals that a small group of axon fascicles exhibit strongly

directed growth toward the midline by 16 hpt and subsequently extends across rapidly (Fig. 3D-

E, Movie S1, white arrowheads). Additional fascicles of axons continue to grow and follow the

growth of the axons that initially crossed the midline (Fig. 3F, Movie S1). These axons grow

along a path that almost completely overlaps with their original path (Fig. 4D), and strongly

suggests that there are factors expressed from cells associated with that projection that direct

regrowing RGC axons during optic nerve regeneration.

A novel role for oligodendrocytes in axon guidance during axon regeneration

To determine the relevant cell type through which *lh3* is functioning during optic nerve

regeneration, we use cell-type specific transgenic rescue. Based on the *lh3* and *col18a1* mRNA

expression in the optic chiasm (Fig. 2G-I, 6A), we hypothesized that the relevant cell type would

be one found in that region during the initial stages of regeneration. We observe macrophages

and microglia dynamically infiltrating the chiasm and surrounding regenerating axonal growth

(Fig. 5A). *Tg(mpeg:lh3-myc)* expression partially rescues aberrant axonal growth in *lh3* mutants,

suggesting that *lh3* expression needs to be more specifically spatially regulated (Fig. 5B). We

therefore examined oligodendrocytes and observe oligodendrocyte lineage cells present in the

optic chiasm and we detect *mbpa* mRNA indicative of myelinating oligodendrocytes along the

degenerating axons (Fig. 5C-D). We find that *lh3* expression from a *Tg(sox10:lh3-mkate)*

transgene restores RGC regrowth across the midline in *lh3* mutants (Fig 5E). However, the

transgenic tools we employed here do not allow is to distinguish whether oligodendrocyte

precursor cells or mature oligodendrocytes were the relevant cell type for *lh3* and *col18a1,* as

sox10 and *olig2* are both transcription factors that are expressed in all oligodendrocyte lineage

cells (Bergles and Richardson, 2015; Ackerman and Monk, 2016). Previous studies implicate

oligodendrocytes to play roles in remyelination (Wang *et al.*, 2020; Mendonça *et al.*, 2021) and

axon repulsion or inhibition via myelin-associated glycoprotein (McKerracher *et al.*, 1994;

Mukhopadhyay *et al.*, 1994), NogoA (Chen *et al.*, 2000; GrandPré *et al.*, 2000), oligodendrocyte-

myelin glycoprotein (K. C. Wang *et al.*, 2002) and semaphorin 4D (Moreau-Fauvarque *et al.*,

2003). From our results, we propose a mechanism in which Lh3, and potentially Col18a1,

expressed in oligodendrocytes mediate axon attraction to direct regenerating RGC axons along

their original path, suggesting a novel role for oligodendrocytes in CNS regeneration.

A molecular mechanism for *lh3* and *col18a1* in specifying a regenerative path

We identify *lh3* to be required to direct regenerating RGC axons toward and across the

optic chiasm (Fig 1). We then determined that *col18a1* was critical for directing RGC axons

during optic nerve regeneration, but not another basement membrane *col4a5* or fibrillar *col7a1*,

which demonstrates a specific role for *col18a1* in optic nerve regeneration (Fig 6, Fig S1). But

the question remains: what are the mechanisms by which Lh3 and Col18a1 target RGC axons

to their original path? We cannot exclude a possible mechanism involving RGC axons directly

interacting with Col18a1 in the extracellular matrix through integrins (Elango *et al.*, 2022).

Transcriptomic analysis of RGCs at different stages of regeneration in adult zebrafish shows

that several integrins are upregulated as axons grow toward and across the midline (Dhara *et*

al., 2019). Therefore, the spatial and temporal specificity of Lh3 and Col18a1 expression could

establish the regenerative path for regrowing RGC axons independent of any other guidance

cues.

But in addition to playing structural roles, there is growing evidence showing a critical role

for extracellular matrix proteins to bind and regulate the spatial distribution of axon guidance

molecules in the extracellular space. For decades, glycosaminoglycans, such as heparan

sulfate proteoglycans, have been known to regulate the extracellular localization of Slit repulsive

guidance cues to affect axon guidance in development (Wang and Denburg, 1992; Johnson *et*

al., 2004; Smart *et al.*, 2011). More specifically, *col4a5* has been shown to be critical during

development of the retinotectal projection for localizing Slit1 expression in the superficial layers

of the optic tectum, which mediates lamina-specific axon targeting (Xiao and Baier, 2007; Xiao

et al., 2011). During motor nerve regeneration, spatially restricted *col4a5* and *lh3* are required in

a subset of Schwann cells to possibly create a Slit1a gradient that directs dorsal nerve axons to

their original projection paths (Isaacman-Beck *et al.*, 2015; Murphy, Isaacman-Beck and

Granato, 2022). In this study, our results describe a similar mechanism where *lh3* is required to

direct regrowing RGC axons toward the midline along their original path, but here, we find in

optic nerve regeneration that Lh3 likely acts through Col18a1, not Col4a5. What are possible

guidance cues that Col18a1 could be spatially regulating? Col18a1 has three isoforms and

within the long isoform is a Frizzled-like domain that has been shown to interact with Wnt

molecules (Quélard *et al.,* 2008). Moreover, Wnt signaling has been implicated in both

promoting as well as repelling axonal growth in spinal cord and optic nerve regeneration (Yin *et*

al., 2008; Hollis and Zou, 2012; Yam and Charron, 2013; Strand *et al.*, 2016; Patel, Park and

Hackam, 2017; Wehner *et al.*, 2017). Therefore, we propose another potential mechanism in

which Lh3 is required for Col18a1 expression that possibly spatially restricts Wnts in the

extracellular space along the original RGC axonal projection to direct regenerating RGC axons

during optic nerve regeneration. Altogether, our studies demonstrate that live cell imaging of

optic nerve regeneration in larval zebrafish allows for unprecedented visualization of axonal

growth dynamics and potential interactions with other cell types in the optic chiasm. Our data

here reveal a glial derived molecular pathway that directs regenerating RGC axons across the

optic chiasm during optic nerve regeneration, furthering our understanding of the cellular and

molecular mechanisms that promote robust functional optic nerve regeneration.

Methods

Fish lines and maintenance

All zebrafish (*Danio rerio*) work was performed in compliance with the University of Pennsylvania Institutional Animal Care and Use Committee regulations and raised as previously described(Mullins *et al.*, 1994). All transgenic lines were maintained in the Tupfel long fin genetic background except the *col18a1* line, which were maintain in the AB background. The following transgenic lines and mutants were used: *lh3*^{TV2O5} (Schneider and Granato, 2006), *col18a1*(Isaacman-Beck *et al.*, 2015), *col4a5*^{s510} (Xiao and Baier, 2007), *col7a1*^{sa3473}, *Tg(hsp:lh30-myc)*(Isaacman-Beck *et al.*, 2015), *Tg(sox10:lh3-mkate)*(Isaacman-Beck *et al.*, 2015), *Tg(isl2b:GFP)*(Pittman, Law and Chien, 2008), *Tg(olig2:dsred)*(Kucenas *et al.*, 2008), *Tg*(*mpeg1:Gal4FF*)^{gl25} (Ellett *et al.*, 2011) and *Tg*(*UAS:nfsB-mCherry*)^{c264} (Davison *et al.*, 2007).

Adult zebrafish and embryos were raised at $29°C$ on a 14-h:10-h light:dark cycle.

To generate *Tg(mpeg1.1:lh3-myc)*, Gateway cloning was employed to combine *lh3-myc* into a pDest vector with the mpeg1.1 promoter and Tol2 transposon sites. Tol2 transgenesis was performed as previously described(Suster *et al.*, 2009) into 1-cell stage embryos from *lh3* heterozygous incrosses. Transgenic lines were identified by genotyping for the transgene with forward primer ACAGTCTCTTGCGTCATCAAAACC and reverse primer TGATCACCAGCAGCTCATTG.

Transection Assay

Optic nerve transection assays were performed as previously described(Harvey, Baxter and Granato, 2019, 2023). Briefly, to inhibit melanocyte pigmentation, larvae were raised in phenylthiourea (PTU, 0.2mM in E3 medium) in the dark at $29°C$ beginning at 1 dpf. For live cell

imaging and distal transections, larvae were anesthetized in PTU E3 plus 0.0053% tricaine then mounted in 2.5% low-melt agarose (SeaPlaque, Lonza) prepared with PTU E3 plus 0.016% tricaine ventral-up on a glass microscopy slide. Then one eye was completely removed using forceps at 4 dpf. Larvae recovered in Ringer's solution and then were kept in PTU E3 at 29°C until transected. At 5 dpf, larvae were anesthetized, mounted in agarose, and then optic nerve transections were performed on an Olympus SZX16 fluorescent microscope or a dissecting microscope with a NightSea Stereo Microscope Fluorescence Adapter (RB-GO only wavelength set, bandpass filter) using a sharpened tungsten needle (Fine Science Tools, Tip Diameter: 0.001mm, Rod Diameter: 0.125mm). For proximal transections, optic nerves were transected at the region of the nerve distal to it exiting the eye, yet proximal to the optic chiasm. For distal transections, optic nerves were transected at the region of the nerve distal to the CNS midline, yet proximal to the optic tectum. Following injuries, larvae were removed from the agarose, allowed to recover in Ringer's solution with 0.2mM PTU, then returned to 0.2mM PTU, E3 medium at 29°C. Larvae were inspected for transection efficiency at 16-18 hpt, except larvae that were fixed at early post transection timepoints, and only larvae with complete optic nerve transections with no visible intact axons remaining from the eye to the tectum were kept until fixation at later designated timepoints.

Live Cell Imaging, Processing and Analysis

Larvae were anesthetized in PTU E3 with 0.0053% tricaine then mounted in 2.5% low-melt agarose (SeaPlaque, Lonza) prepared with PTU E3 plus 0.016% tricaine ventral-up on a glass microscopy slide. Then, one eye as well as a portion of the jaw, including the Meckel's cartilage, palatoquadrate, and the most distal parts of the basihyal and ceratohyal structures, were carefully removed using forceps at 4 dpf. Care was taken so as not to damage the heart. Larvae recovered in Ringer's solution and then were kept in PTU E3 at 29°C until transected. At 5 dpf,

larvae were anesthetized again, mounted in agarose, and then optic nerve transections were

performed. Larvae recovered in Ringer's solution and then were kept in PTU E3 at 29°C until

imaged for live cell imaging. Larvae were mounted ventral side up in 35mm glass-bottomed

Petri dishes in 1.5% low-melt agarose with 0.016% tricaine. Larvae were imaged using a Leica

SP8 confocal microscope with the resonant scanner on, equipped with a petri dish heater and

heated water perfusion so fresh E3 water flowed through the petri dish and the temperature was

kept constant at around 28°C during the imaging session. Larvae were imaged from the ventral

side of the brain, which was not covered by any jaw tissue. Image stacks were acquired every

10-15 minutes with a 20x 1.0 NA lens. All images were adjusted for brightness and contrast, and

color assigned using Fiji.

To determine the time of growth initiation extending from the proximal stump of transected RGC

axons, we recorded in which frame axonal extension longer than 1um occurred that continued to

extend for another subsequent frame.

To measure the angle of the original path before transection to the end of distinct axon fascicles,

we first rotated the images so that parasphenoid cartilage was precisely aligned vertically

against a straight line drawn in Fiji. A circle with a radius of 80 uM centered where RGC axons

exit the eye was drawn within Fiji using a Make Oval Macro in timelapse image sequences.

Axons and fascicles terminating farther than the 80 uM region of interest at 20 hpt were used for

measuring. First the angle of the original path to the parasphenoid was determined using frames

before Wallerian degeneration was complete. Then the angle of the axons and fascicles to the

parasphenoid were determined, and then the difference was calculated and graphed.

Heat Shock-Induced *lh3* Rescue

To induce expression of Lh3 from *Tg(hsp:lh3-myc)* transgene, during development: Embryos

were developmentally staged in the late afternoon. No more than 6 embryos were placed per

tube in PCR strip-tubes. Tubes were incubated in a thermal cycler at 28°C until embryos would

reach 11.66 hpf, then heated to 38°C for 10minutes, then held at 28°C until they could be

removed from the tubes.

To induce expression of Lh3 from Tg(hsp:lh3-myc) transgene, during regeneration: Larvae at 5

dpf homozygous for the Tg(hsp:lh3-myc) transgene were placed only one per tube in PCR strip-

tubes. Tubes were incubated in a thermal cycler at 38°C for 30min then held at 29°C until they

could be removed from the tubes. Larvae were transected 4-5 hours later.

Immunostaining

Larvae were stained using methods modified from those previously described (Randlett *et al.*,

2015). Briefly, larvae were fixed in 4% PFA in PBS overnight at 4°C. Larvae were washed in

PBS + 0.25% Triton (PBT), incubated in 150mM Tris-HCl pH 9.0 for 15 min at 70°C, then

washed in PBT. Larvae were permeabilized in 0.05% Trypsin-EDTA for 5 min on ice , washed in

PBT, blocked in PBT containing 1% bovine serum albumin (BSA), 2% normal goat serum (NGS)

and 1% dimethyl sulfoxide (DMSO), and then incubated in primary and secondary antibodies

overnight at 4°C in PBT containing 1% BSA and 1% DMSO. Stained larvae were stored and

mounted in Vectashield (Vector Laboratories) for imaging using confocal microscopy. For static

timepoint analysis of *lh3* and *col18a1* mutants, primary antibodies used were mouse anti-GFP

(JL-8, 1:200, Takara Bio), and secondary antibodies used were goat anti-mouse Alexa 488

(1:500, Molecular Probes). For static timepoint analysis of macrophages/microglia, primary

antibodies used were mouse anti-GFP (JL-8, 1:200, Takara Bio), rabbit anti-RFP (1:200, Abcam

ab62341) and secondary antibodies used were goat anti-mouse Alexa 488 (1:500, Molecular

Probes) and goat anti-rabbit Alexa 594 (1:500, Molecular Probes). For static timepoint analysis

of oligodendrocytes, primary antibodies used were rabbit anti-GFP (1:500, Molecular Probes),

mouse anti-dsred (1:500, BD Pharmingen) and secondary antibodies used were goat anti-rabbit

488 (1:500, Molecular Probes) and goat anti-mouse Alexa 594 (1:500, Molecular Probes).

Fluorescent *in situ* Hybridization with Hybridization Chain Reaction (HCR)

Lh3 (plod3), col18a1 and mbpa mRNA expression were detected using fluorescent *in situ* hybridization with HCR (Molecular Instruments, Los Angeles, CA, USA)(Choi *et al.*, 2018). HCR probes, buffers, and hairpins were purchased from Molecular Instruments. *Tg(isl2b:GFP)* larvae were fixed at 5 dpf with 4% paraformaldehyde in PBS overnight at 4°C. No antibody staining was required together with in situ HCR. The staining was performed as previously described(Shainer *et al.*, 2023).

Static Confocal Imaging, Processing and Analysis

Larvae were imaged on a Zeiss 880 confocal microscope using a 20× objective. Image stacks of optic chiasms or tecta were compressed into maximum intensity projections. All images were adjusted for brightness and contrast in Adobe Photoshop CS5, and color assigned using Fiji.

Statistical Analysis

Data were imported into GraphPad Prism for analysis. Statistical analysis was performed as indicated throughout the text.

Acknowledgments

We would like to thank Dr. Andrea Stout of the UPenn Cell and Developmental Biology Microscopy Core, as well as Dr. Gordon Ruthel of the Penn Vet Imaging Core for providing technical assistance. We thank all members of the Granato laboratory for comments and discussion.

Funding

This work was supported by grants from the National Institutes of Health to B.M.H. (EY032593) and M.G. (EY024861 and NS097914).

Author Contributions

B.M.H. and M.G. conceptualized the experiments; B.M.H. M.B, and A.M.G. performed and

analyzed experiments; B.M.H. wrote the original draft. All authors contributed to editing and

revision of the paper.

Data Availability

All relevant data are within the article and its supplementary information files.

Competing interests

The authors declare no competing interests.

References

References

Ackerman, S.D. and Monk, K.R. (2016) 'The scales and tales of myelination: using zebrafish
and mouse to study myelinating glia', *Brain Research*, 1641(Pt A), pp. 79–91. Available at:
https://doi.org/10.1016/j.brainres.2015.10.011.

Bergles, D.E. and Richardson, W.D. (2015) 'Oligodendrocyte Development and Plasticity', *Cold*
Spring Harbor Perspectives in Biology, 8(2), p. a020453. Available at:
https://doi.org/10.1101/cshperspect.a020453.

Berkelaar, M. *et al.* (1994) 'Axotomy results in delayed death and apoptosis of retinal ganglion
cells in adult rats', *The Journal of Neuroscience: The Official Journal of the Society for*
Neuroscience, 14(7), pp. 4368–4374.

Berry, M., Ahmed, Z. and Logan, A. (2019) 'Return of function after CNS axon regeneration:
Lessons from injury-responsive intrinsically photosensitive and alpha retinal ganglion cells',
Progress in Retinal and Eye Research, 71, pp. 57–67. Available at:
https://doi.org/10.1016/j.preteyeres.2018.11.006.

Bhatt, D.H. *et al.* (2004) 'Cyclic AMP-induced repair of zebrafish spinal circuits', *Science (New*
York, N.Y.), 305(5681), pp. 254–258. Available at: https://doi.org/10.1126/science.1098439.

Bremer, J. *et al.* (2019) 'The ubiquitin ligase PHR promotes directional regrowth of spinal
zebrafish axons', *Communications Biology*, 2, p. 195. Available at:
https://doi.org/10.1038/s42003-019-0434-2.

Brennan, F.H. *et al.* (2022) 'Microglia coordinate cellular interactions during spinal cord repair in
mice', *Nature Communications*, 13(1), p. 4096. Available at: https://doi.org/10.1038/s41467-022-
31797-0.

Brockerhoff, S.E. *et al.* (1995) 'A behavioral screen for isolating zebrafish mutants with visual
system defects', *Proceedings of the National Academy of Sciences of the United States of*
America, 92(23), pp. 10545–10549.

Brösamle, C. and Halpern, M.E. (2002) 'Characterization of myelination in the developing
zebrafish', *Glia*, 39(1), pp. 47–57. Available at: https://doi.org/10.1002/glia.10088.

Ceci, M.L. *et al.* (2014) 'Axon-Schwann cell interactions during peripheral nerve regeneration in
zebrafish larvae', *Neural Development*, 9, p. 22. Available at: https://doi.org/10.1186/1749-8104-
9-22.

Chen, M.S. *et al.* (2000) 'Nogo-A is a myelin-associated neurite outgrowth inhibitor and an
antigen for monoclonal antibody IN-1', *Nature*, 403(6768), pp. 434–439. Available at:
https://doi.org/10.1038/35000219.

Choi, H.M.T. *et al.* (2018) 'Third-generation in situ hybridization chain reaction: multiplexed,
quantitative, sensitive, versatile, robust', *Development (Cambridge, England)*, 145(12), p.
dev165753. Available at: https://doi.org/10.1242/dev.165753.

Davison, J.M. *et al.* (2007) 'Transactivation from Gal4-VP16 transgenic insertions for tissue-
specific cell labeling and ablation in zebrafish', *Developmental Biology*, 304(2), pp. 811–824.
Available at: https://doi.org/10.1016/j.ydbio.2007.01.033.

Dhara, S.P. *et al.* (2019) 'Cellular reprogramming for successful CNS axon regeneration is
driven by a temporally changing cast of transcription factors', *Scientific Reports*, 9(1), p. 14198.
Available at: https://doi.org/10.1038/s41598-019-50485-6.

Easter, S.S. and Nicola, G.N. (1996) 'The development of vision in the zebrafish (Danio rerio)',
Developmental Biology, 180(2), pp. 646–663.

Elango, J. *et al.* (2022) 'The Molecular Interaction of Collagen with Cell Receptors for Biological
Function', *Polymers*, 14(5), p. 876. Available at: https://doi.org/10.3390/polym14050876.

Ellett, F. *et al.* (2011) 'mpeg1 promoter transgenes direct macrophage-lineage expression in
zebrafish', *Blood*, 117(4), pp. e49-56. Available at: https://doi.org/10.1182/blood-2010-10-
314120.

Fitch, M.T. and Silver, J. (2008) 'CNS injury, glial scars, and inflammation: Inhibitory extracellular
matrices and regeneration failure', *Experimental Neurology*, 209(2), pp. 294–301. Available at:
https://doi.org/10.1016/j.expneurol.2007.05.014.

Fukai, N. *et al.* (2002) 'Lack of collagen XVIII/endostatin results in eye abnormalities', *The
EMBO journal*, 21(7), pp. 1535–1544. Available at: https://doi.org/10.1093/emboj/21.7.1535.

Gahtan, E., Tanger, P. and Baier, H. (2005) 'Visual prey capture in larval zebrafish is controlled
by identified reticulospinal neurons downstream of the tectum', *The Journal of Neuroscience:
The Official Journal of the Society for Neuroscience*, 25(40), pp. 9294–9303. Available at:
https://doi.org/10.1523/JNEUROSCI.2678-05.2005.

Goldberg, J.L. *et al.* (2002) 'Retinal ganglion cells do not extend axons by default: promotion by
neurotrophic signaling and electrical activity', *Neuron*, 33(5), pp. 689–702. Available at:
https://doi.org/10.1016/s0896-6273(02)00602-5.

GrandPré, T. *et al.* (2000) 'Identification of the Nogo inhibitor of axon regeneration as a
Reticulon protein', *Nature*, 403(6768), pp. 439–444. Available at:
https://doi.org/10.1038/35000226.

Harvey, B.M., Baxter, M. and Granato, M. (2019) 'Optic nerve regeneration in larval zebrafish
exhibits spontaneous capacity for retinotopic but not tectum specific axon targeting', *PloS One*,
14(6), p. e0218667. Available at: https://doi.org/10.1371/journal.pone.0218667.

Harvey, B.M., Baxter, M. and Granato, M. (2023) 'Assaying Optic Nerve Regeneration in Larval
Zebrafish', *Methods in Molecular Biology (Clifton, N.J.)*, 2636, pp. 191–203. Available at:
https://doi.org/10.1007/978-1-0716-3012-9_10.

Heikkinen, J. *et al.* (2000) 'Lysyl hydroxylase 3 is a multifunctional protein possessing collagen
glucosyltransferase activity', *The Journal of Biological Chemistry*, 275(46), pp. 36158–36163.
Available at: https://doi.org/10.1074/jbc.M006203200.

Hollis, E.R. and Zou, Y. (2012) 'Expression of the Wnt signaling system in central nervous system axon guidance and regeneration', *Frontiers in Molecular Neuroscience*, 5, p. 5. Available at: https://doi.org/10.3389/fnmol.2012.00005.

Horder, T.J. (1971) 'The course of recovery of the retinotectal projection during regeneration of the fish optic nerve', *The Journal of Physiology*, 217 Suppl, pp. 53P-54P.

Isaacman-Beck, J. *et al.* (2015) 'The lh3 Glycosyltransferase Directs Target-Selective Peripheral Nerve Regeneration', *Neuron*, 88(4), pp. 691–703. Available at: https://doi.org/10.1016/j.neuron.2015.10.004.

Johnson, K.G. *et al.* (2004) 'Axonal heparan sulfate proteoglycans regulate the distribution and efficiency of the repellent slit during midline axon guidance', *Current biology: CB*, 14(6), pp. 499–504. Available at: https://doi.org/10.1016/j.cub.2004.02.005.

Kucenas, S. *et al.* (2008) 'CNS-derived glia ensheath peripheral nerves and mediate motor root development', *Nature Neuroscience*, 11(2), pp. 143–151. Available at: https://doi.org/10.1038/nn2025.

Kurimoto, T. *et al.* (2010) 'Long-distance axon regeneration in the mature optic nerve: contributions of oncomodulin, cAMP, and pten gene deletion', *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 30(46), pp. 15654–15663. Available at: https://doi.org/10.1523/JNEUROSCI.4340-10.2010.

Leon, S. *et al.* (2000) 'Lens injury stimulates axon regeneration in the mature rat optic nerve', *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 20(12), pp. 4615–4626.

Lewis, G.M. and Kucenas, S. (2014) 'Perineurial glia are essential for motor axon regrowth following nerve injury', *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 34(38), pp. 12762–12777. Available at: https://doi.org/10.1523/JNEUROSCI.1906-14.2014.

Luo, X. *et al.* (2013) 'Three-dimensional evaluation of retinal ganglion cell axon regeneration and pathfinding in whole mouse tissue after injury', *Experimental Neurology*, 247, pp. 653–662. Available at: https://doi.org/10.1016/j.expneurol.2013.03.001.

Marquart, G.D. *et al.* (2017) 'High-precision registration between zebrafish brain atlases using symmetric diffeomorphic normalization', *GigaScience*, 6(8), pp. 1–15. Available at: https://doi.org/10.1093/gigascience/gix056.

Martin, S.M. *et al.* (2010) 'Wallerian degeneration of zebrafish trigeminal axons in the skin is required for regeneration and developmental pruning', *Development (Cambridge, England)*, 137(23), pp. 3985–3994. Available at: https://doi.org/10.1242/dev.053611.

McKerracher, L. *et al.* (1994) 'Identification of myelin-associated glycoprotein as a major myelin-derived inhibitor of neurite growth', *Neuron*, 13(4), pp. 805–811. Available at: https://doi.org/10.1016/0896-6273(94)90247-x.

Mendonça, H.R. *et al.* (2021) 'Myelination of regenerating optic nerve axons occurs in conjunction with an increase of the oligodendrocyte precursor cell population in the adult mice',

Brain Research Bulletin, 166, pp. 150–160. Available at: https://doi.org/10.1016/j.brainresbull.2020.11.012.

Moore, D.L. *et al.* (2009) 'KLF family members regulate intrinsic axon regeneration ability', *Science (New York, N.Y.)*, 326(5950), pp. 298–301. Available at: https://doi.org/10.1126/science.1175737.

Moreau-Fauvarque, C. *et al.* (2003) 'The transmembrane semaphorin Sema4D/CD100, an inhibitor of axonal growth, is expressed on oligodendrocytes and upregulated after CNS lesion', *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 23(27), pp. 9229–9239. Available at: https://doi.org/10.1523/JNEUROSCI.23-27-09229.2003.

Mukhopadhyay, G. *et al.* (1994) 'A novel role for myelin-associated glycoprotein as an inhibitor of axonal regeneration', *Neuron*, 13(3), pp. 757–767. Available at: https://doi.org/10.1016/0896-6273(94)90042-6.

Mullins, M.C. *et al.* (1994) 'Large-scale mutagenesis in the zebrafish: in search of genes controlling development in a vertebrate', *Current biology: CB*, 4(3), pp. 189–202.

Murphy, P.L., Isaacman-Beck, J. and Granato, M. (2022) 'Robo2 Drives Target-Selective Peripheral Nerve Regeneration in Response to Glia-Derived Signals', *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 42(5), pp. 762–776. Available at: https://doi.org/10.1523/JNEUROSCI.1528-21.2021.

Murray, M. (1976) 'Regeneration of retinal axons into the goldfish optic tectum', *The Journal of Comparative Neurology*, 168(2), pp. 175–195. Available at: https://doi.org/10.1002/cne.901680202.

Nogueira-Rodrigues, J. *et al.* (2022) 'Rewired glycosylation activity promotes scarless regeneration and functional recovery in spiny mice after complete spinal cord transection', *Developmental Cell*, 57(4), pp. 440-450.e7. Available at: https://doi.org/10.1016/j.devcel.2021.12.008.

Norsworthy, M.W. *et al.* (2017) 'Sox11 Expression Promotes Regeneration of Some Retinal Ganglion Cell Types but Kills Others', *Neuron*, 94(6), pp. 1112-1120.e4. Available at: https://doi.org/10.1016/j.neuron.2017.05.035.

Park, K.K. *et al.* (2008) 'Promoting axon regeneration in the adult CNS by modulation of the PTEN/mTOR pathway', *Science (New York, N.Y.)*, 322(5903), pp. 963–966. Available at: https://doi.org/10.1126/science.1161566.

Park, K.K. *et al.* (2017) 'Retinal ganglion cell survival and axon regeneration after optic nerve injury in naked mole-rats', *The Journal of Comparative Neurology*, 525(2), pp. 380–388. Available at: https://doi.org/10.1002/cne.24070.

Patel, A.K., Park, K.K. and Hackam, A.S. (2017) 'Wnt signaling promotes axonal regeneration following optic nerve injury in the mouse', *Neuroscience*, 343, pp. 372–383. Available at: https://doi.org/10.1016/j.neuroscience.2016.12.020.

Pernet, Vincent *et al.* (2013) 'Long-distance axonal regeneration induced by CNTF gene transfer
is impaired by axonal misguidance in the injured adult optic nerve', *Neurobiology of Disease*, 51,
pp. 202–213. Available at: https://doi.org/10.1016/j.nbd.2012.11.011.

Pernet, V. *et al.* (2013) 'Misguidance and modulation of axonal regeneration by Stat3 and
Rho/ROCK signaling in the transparent optic nerve', *Cell Death & Disease*, 4, p. e734. Available
at: https://doi.org/10.1038/cddis.2013.266.

Pittman, A.J., Law, M.-Y. and Chien, C.-B. (2008) 'Pathfinding in a large vertebrate axon tract:
isotypic interactions guide retinotectal axons at multiple choice points', *Development
(Cambridge, England)*, 135(17), pp. 2865–2871. Available at:
https://doi.org/10.1242/dev.025049.

Quélard, D. *et al.* (2008) 'A cryptic frizzled module in cell surface collagen 18 inhibits Wnt/beta-
catenin signaling', *PloS One*, 3(4), p. e1878. Available at:
https://doi.org/10.1371/journal.pone.0001878.

Randlett, O. *et al.* (2015) 'Whole-brain activity mapping onto a zebrafish brain atlas', *Nature
Methods*, 12(11), pp. 1039–1046. Available at: https://doi.org/10.1038/nmeth.3581.

Risteli, M. *et al.* (2004) 'Characterization of collagenous peptides bound to lysyl hydroxylase
isoforms', *The Journal of Biological Chemistry*, 279(36), pp. 37535–37543. Available at:
https://doi.org/10.1074/jbc.M405638200.

Rosenberg, A.F. *et al.* (2012) 'In vivo nerve-macrophage interactions following peripheral nerve
injury', *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*,
32(11), pp. 3898–3909. Available at: https://doi.org/10.1523/JNEUROSCI.5225-11.2012.

Salo, A.M. and Myllyharju, J. (2021) 'Prolyl and lysyl hydroxylases in collagen synthesis',
Experimental Dermatology, 30(1), pp. 38–49. Available at: https://doi.org/10.1111/exd.14197.

Schneider, V.A. and Granato, M. (2006) 'The myotomal diwanka (lh3) glycosyltransferase and
type XVIII collagen are critical for motor growth cone migration', *Neuron*, 50(5), pp. 683–695.
Available at: https://doi.org/10.1016/j.neuron.2006.04.024.

Shainer, I. *et al.* (2023) 'A single-cell resolution gene expression atlas of the larval zebrafish
brain', *Science Advances*, 9(8), p. eade9909. Available at:
https://doi.org/10.1126/sciadv.ade9909.

Smart, A.D. *et al.* (2011) 'Heparan sulfate proteoglycan specificity during axon pathway
formation in the Drosophila embryo', *Developmental Neurobiology*, 71(7), pp. 608–618.
Available at: https://doi.org/10.1002/dneu.20854.

Smith, P.D. *et al.* (2009) 'SOCS3 deletion promotes optic nerve regeneration in vivo', *Neuron*,
64(5), pp. 617–623. Available at: https://doi.org/10.1016/j.neuron.2009.11.021.

Strand, N.S. *et al.* (2016) 'Wnt/β-catenin signaling promotes regeneration after adult zebrafish
spinal cord injury', *Biochemical and Biophysical Research Communications*, 477(4), pp. 952–
956. Available at: https://doi.org/10.1016/j.bbrc.2016.07.006.

Suster, M.L. *et al.* (2009) 'Transgenesis in zebrafish with the tol2 transposon system', *Methods*
in Molecular Biology (Clifton, N.J.), 561, pp. 41–63. Available at: https://doi.org/10.1007/978-1-
60327-019-9_3.

Suzuki, O.T. *et al.* (2002) 'Molecular analysis of collagen XVIII reveals novel mutations,
presence of a third isoform, and possible genetic heterogeneity in Knobloch syndrome',
American Journal of Human Genetics, 71(6), pp. 1320–1329. Available at:
https://doi.org/10.1086/344695.

Taler, K. *et al.* (2019) 'Lysyl hydroxylase 3 is required for normal lens capsule formation and
maintenance of lens epithelium integrity and fate', *Developmental Biology* [Preprint]. Available
at: https://doi.org/10.1016/j.ydbio.2019.10.020.

Vargas, M.E. and Barres, B.A. (2007) 'Why is Wallerian degeneration in the CNS so slow?',
Annual Review of Neuroscience, 30, pp. 153–179. Available at:
https://doi.org/10.1146/annurev.neuro.30.051606.094354.

Villegas-Pérez, M.P. *et al.* (1993) 'Rapid and protracted phases of retinal ganglion cell loss
follow axotomy in the optic nerve of adult rats', *Journal of Neurobiology*, 24(1), pp. 23–36.
Available at: https://doi.org/10.1002/neu.480240103.

Waller, A. (1851) 'Experiments on the Section of the Glosso-Pharyngeal and Hypoglossal
Nerves of the Frog, and Observations of the Alterations Produced Thereby in the Structure of
Their Primitive Fibres', *Edinburgh Medical and Surgical Journal*, 76(189), pp. 369–376.

Wang, C. *et al.* (2002) 'The third activity for lysyl hydroxylase 3: galactosylation of hydroxylysyl
residues in collagens in vitro', *Matrix Biology: Journal of the International Society for Matrix*
Biology, 21(7), pp. 559–566.

Wang, J. *et al.* (2020) 'Robust Myelination of Regenerated Axons Induced by Combined
Manipulations of GPR17 and Microglia', *Neuron*, 108(5), pp. 876-886.e4. Available at:
https://doi.org/10.1016/j.neuron.2020.09.016.

Wang, K.C. *et al.* (2002) 'Oligodendrocyte-myelin glycoprotein is a Nogo receptor ligand that
inhibits neurite outgrowth', *Nature*, 417(6892), pp. 941–944. Available at:
https://doi.org/10.1038/nature00867.

Wang, L. and Denburg, J.L. (1992) 'A role for proteoglycans in the guidance of a subset of
pioneer axons in cultured embryos of the cockroach', *Neuron*, 8(4), pp. 701–714. Available at:
https://doi.org/10.1016/0896-6273(92)90091-q.

Wehner, D. *et al.* (2017) 'Wnt signaling controls pro-regenerative Collagen XII in functional
spinal cord regeneration in zebrafish', *Nature Communications*, 8(1), p. 126. Available at:
https://doi.org/10.1038/s41467-017-00143-0.

Wullink, B. *et al.* (2018) 'Type VII Collagen in the Human Accommodation System: Expression in
Ciliary Body, Zonules, and Lens Capsule', *Investigative Ophthalmology & Visual Science*, 59(2),
pp. 1075–1083. Available at: https://doi.org/10.1167/iovs.17-23425.

Xiao, T. *et al.* (2011) 'Assembly of lamina-specific neuronal connections by slit bound to type IV
collagen', *Cell*, 146(1), pp. 164–176. Available at: https://doi.org/10.1016/j.cell.2011.06.016.

Xiao, T. and Baier, H. (2007) 'Lamina-specific axonal projections in the zebrafish tectum require
the type IV collagen Dragnet', *Nature Neuroscience*, 10(12), pp. 1529–1537. Available at:
https://doi.org/10.1038/nn2002.

Yam, P.T. and Charron, F. (2013) 'Signaling mechanisms of non-conventional axon guidance
cues: the Shh, BMP and Wnt morphogens', *Current Opinion in Neurobiology*, 23(6), pp. 965–
973. Available at: https://doi.org/10.1016/j.conb.2013.09.002.

Yin, Y. *et al.* (2009) 'Oncomodulin links inflammation to optic nerve regeneration', *Proceedings
of the National Academy of Sciences of the United States of America*, 106(46), pp. 19587–
19592. Available at: https://doi.org/10.1073/pnas.0907085106.

Yin, Y. and Benowitz, L.I. (2018) 'In Vitro and In Vivo Methods for Studying Retinal Ganglion Cell
Survival and Optic Nerve Regeneration', *Methods in Molecular Biology (Clifton, N.J.)*, 1695, pp.
187–205. Available at: https://doi.org/10.1007/978-1-4939-7407-8_16.

Yin, Z.-S. *et al.* (2008) 'Repair effect of Wnt3a protein on the contused adult rat spinal cord',
Neurological Research, 30(5), pp. 480–486. Available at:
https://doi.org/10.1179/174313208X284133.

Yiu, G. and He, Z. (2006) 'Glial inhibition of CNS axon regeneration', *Nature Reviews.
Neuroscience*, 7(8), pp. 617–627. Available at: https://doi.org/10.1038/nrn1956.

Zeller, J. and Granato, M. (1999) 'The zebrafish diwanka gene controls an early step of motor
growth cone migration', *Development (Cambridge, England)*, 126(15), pp. 3461–3472.

Zhang, Y. *et al.* (2014) 'An RNA-sequencing transcriptome and splicing database of glia,
neurons, and vascular cells of the cerebral cortex', *The Journal of Neuroscience: The Official
Journal of the Society for Neuroscience*, 34(36), pp. 11929–11947. Available at:
https://doi.org/10.1523/JNEUROSCI.1860-14.2014.

Figure 1.

Figure 1. Lh3 is required for growth across the midline during regeneration. (A-B) Before injury, RGC axons labeled by *Tg(isl2b:GFP)* in siblings and conditional *lh3* mutants cross the midline and (C-D) innervate contralateral tecta. (E-H) In contrast to *lh3* siblings at 72 hours post transection (hpt), RGC axons in *lh3* mutants turn away from the midline, project aberrantly (magenta arrowheads) and do not reinnervate contralateral optic tecta. (A-H) Dashed lines indicate the outline of the eyes; scale bars = 50 µm (I-J) Schematics of RGC axonal growth at 72 hpt for (I) *lh3* siblings and (J) *lh3* mutants. (K-L) Quantification of the percent of nerves that (K) grew across the midline to reinnervate the optic tectum or (L) did not grow across the midline and instead grew along aberrant trajectories at 72 hpt. Each dot represents a technical replicate. For sibling nerves, n = 37; for mutant nerves, n = 46. Data are represented as mean ± SD; ****P<0.001*, t-test. Representative images of chiasms and tecta for each timepoint are of the same fixed larva, though across timepoints are different larvae.

Figure 2.

Figure 2. *Lh3* expression is upregulated in the optic chiasm during early regeneration. (A)

Timeline of induced *lh3* expression with *Tg(hsp70:lh3-myc)* during regeneration. (B-E) At 72

hours post transection (hpt) in *lh3* siblings with no Heat Shock induction, RGC axons cross the

midline to innervate the optic tecta, while RGC axons in *lh3* mutants turn away from the midline,

project aberrantly (C, magenta arrowheads). In *lh3* siblings and mutants that received a heat

shock, RGC axons project across the midline (E, white arrowheads) and innervate the optic

tecta. Dashed lines indicate the outline of the eyes; scale bar = 50 μm. (F) Quantification of

nerves that regenerated across the midline to the optic tectum at 72 hours post transection (hpt)

in *Tg(isl2b:GFP)* larvae with and without heat-shock. Each dot represents a technical replicate.

For sibling nerves, no heat shock n = 38, with heat shock n = 44; for mutant nerves, no heat

shock n = 8, with heat shock n = 12. Data are represented as mean ± SD; **$P<0.01$*, n.s.*$P>0.05$,

t-test. (G-I) *Lh3* mRNA expression in *Tg(isl2b:GFP)* larvae pre-transection and at 2-6 hpt. (G)

White dashed boxes are areas of the retina shown in (H) and the optic chiasm shown in (I).

Scale bars, (G) 50 μm, (H) 20 μm, (I) 50 μm. Axons in the chiasm region at 2-6 hpt have not yet

fully degenerated following transection. Dashed lines indicate the outline of the eyes in (I).

Representative images across timepoints are of different larvae.

Figure 3.

Figure 3. *Lh3* directs the initial regrowing RGC axons. (A) Schematic of larvae expressing

Tg(isl2b:GFP) to label RGCs that were used for *in vivo* imaging. Larvae have one eye removed

one day prior to timelapse imaging. Red box indicates the region imaged in panels B-K. (B,G)

Transected RGC axons separate from the distal portion of the optic nerve that will eventually

degenerate (indicated by brackets in B-D, G-I). (C) In *lh3* siblings, RGC axons initiate growth out

from the proximal stump. (D) A small group of axons (white arrowhead) project directly toward

the midline along the original path of the projection and then (E-F) subsequently additional

axonal growth follows (n = 7/10 nerves). (H) In the absence of *lh3*, axons grow out from the

proximal stump, but (I-K) persist along aberrant trajectories (magenta arrowheads; n = 6/12

nerves). Dashed lines indicate the outline of the eyes; scale bar = 25 μm.

Figure 4.

Figure 4. ***Lh3* is required to direct regrowing RGC axons along their original path.** (A) Representative frames from live timelapses of larvae expressing *Tg(isl2b:GFP)* to label RGCs showing the initial axonal growth that emerges from the injured proximal stump of axons. hpt = hours post transection. (B) Comparison of the time when axon fascicles first emerge from the proximal stump in *lh3* siblings and mutants. Time was determined by the first frame when axon fascicles were observed extending from the proximal stump that extended for at least one other consecutive frame. Data are represented as mean ± SD; n.s.*P>0.05*, t-test. (C) Representative image of a regenerating nerve at 20 hpt and the mark-up to measure angle of growing fascicles

780   from the injured stump of axons. See Methods for more details on quantifications. (D)

781   Quantification at 20 hpt of the angle of axon fascicles from the end of a projection to the path of

782   the original axonal projection (n = 6 sibling nerves that grew across the midline, n = 5 mutant

783   nerves that grew along aberrant paths).

Figure 5.

Figure 5. Lh3 in *sox10* expressing glia restores RGC axonal growth across the midline.

(A) In wild-type larvae expressing *Tg(isl2b:GFP)* to label RGCs, macrophages and microglia

labeled by *Tg*(*mpeg1:Gal4FF*);*Tg*(*UAS:nfsB-mCherry*) are observed in the area of the optic

chiasm before transection. At 24 hours post transection (hpt), an influx of *mpeg+* cells is seen.

(B) Quantification of nerves that regenerated across the midline to the optic tectum in larvae

with and without *Tg(mpeg:lh3-myc)* transgene. Each dot represents a technical replicate. For

sibling nerves, no *mpeg* transgene n = 47, with transgene n = 57; for mutant nerves, no *mpeg*

transgene n = 44, with transgene n = 22. Data are represented as mean ± SD. (C-D)

Oligodendrocyte lineage cells labeled by *Tg(olig2:dsred)* as well as *mbpa* mRNA expression in

myelinating oligodendrocytes are also seen near the injured proximal stump of RGC axons or

along the degenerating original projection before and after transection. (E) Quantification of

nerves that regenerated across the midline to the optic tectum in larvae with and without

Tg(sox10:lh3-mkate) transgene. Each dot represents a technical replicate. For sibling nerves,

no *sox10* transgene n = 38, with transgene n = 66; for mutant nerves, no *sox10* transgene n =

28, with transgene n = 16. Data are represented as mean ± SD; **$P<0.01$, n.s.$P>0.05$, t-test. (F)

Lh3 mRNA expression is upregulated and overlaps with oligodendrocytes labeled by

Tg(olig2:dsred). Representative images across timepoints are of different larvae. Axons in the

chiasm region at 4 and 6 hpt have not yet fully degenerated following transection, while at 24

hpt new axon growth enters the chiasm. Dashed lines indicate the outline of the eyes; scale

bars = 100um.

Figure 6.

Figure 6. The Lh3 candidate substrate Col18a1 is upregulated after transection and directs RGC axons across the midline during regeneration. (A) *Col18a1* mRNA expression pre-transection and at 12 hours post transection (hpt) in larvae expressing *Tg(isl2b:GFP)* to label RGCs. Axons in the chiasm region at 12 hpt have not yet fully degenerated following transection. **(**B-C) Before injury, RGC axons labeled by *Tg(isl2b:GFP)* in siblings and conditional *col18a1* mutants cross the midline and (D-E) innervate contralateral tecta. (F-I) In contrast to *col18a1* siblings at 72 hours post transection (hpt), RGC axons in *col18a1* mutants turn away from the midline, project aberrantly (magenta arrowheads) and do not reinnervate contralateral

optic tecta. Dashed lines indicate the outline of the eyes; scale bar = 50 μm. (J-K) Quantification

of the percent of nerves that (J) grew across the midline to reinnervate the optic tectum or (K)

did not grow across the midline and instead grew along aberrant trajectories at 72 hpt. Each dot

represents a technical replicate. For sibling nerves, n = 60; for mutant nerves, n = 22. Data are

represented as mean ± SD; ****P<0.001*, t-test. Representative images of chiasms and tecta for

each timepoint are of the same fixed larva, though across timepoints are different larvae.