A glial cell derived pathway directs regenerating optic nerve axons toward

the optic chiasm

Running Title: Glia direct regenerating RGC axons

Beth M. Harvey[1*], Melissa Baxter[1], Alexis M. Garcia[1] and Michael Granato[1]

Department of Cell and Developmental Biology, Perelman School of Medicine, University of

Pennsylvania, Philadelphia, Pennsylvania 19104

* Corresponding author

email: bharvey@pennmedicine.upenn.edu

Key words: lh3, collagen18a1, RGC, optic nerve regeneration, axon guidance, zebrafish

Abstract

After optic nerve injury, several retinal ganglion cell (RGC) intrinsic signaling pathways have been shown to enhance RGC survival and RGC axonal growth. In contrast, few extrinsic cues have been identified that guide regenerating RGC axons toward and across the optic chiasm. Here, we use live-cell imaging in larval zebrafish to show that regrowing RGC axons initiate growth toward the midline and extend along a trajectory similar to their original projection. From a candidate genetic screen, we identify the glycosyltransferase Lh3 (also referred to as Plod3) to be required to direct regrowing RGC axons toward the midline during active regeneration. Moreover, we show that transgenic *lh3* expression in *sox10*+ presumptive *olig2*+ oligodendrocytes located near the optic chiasm restores directed axonal growth in *lh3* mutants. Finally, we find that mutants in *collagen 18a1 (col18a1),* a putative Lh3 substrate, display RGC axonal misguidance phenotypes similar to *lh3* mutants, suggesting that *lh3* may act through *col18a1* during regeneration. Combined, these data identify *lh3* as part of a glial derived molecular pathway critical for guiding *in vivo* regenerating RGC axons toward and across the optic chiasm.

Introduction

As a component of the central nervous system, retinal ganglion cell (RGC) axons within

the mammalian optic nerve display a poor regenerative capacity. Multiple factors contribute to the

minimal regeneration in mammals, including massive injury-induced RGC death and overall

limited axonal regrowth (Villegas-Pérez *et al.*, 1993; Berkelaar *et al.*, 1994; Goldberg *et al.*, 2002).

Several studies have identified a number of RGC intrinsic signaling pathways that enhance RGC

survival and increase long range axonal growth after injury. For example, deletion of the

transcriptional repressor KLF4 in RGCs increases axonal regrowth (Moore *et al.*, 2009). Similarly,

activation of inflammatory responses within the retina through injection of Zymosan and cAMP

enhances cell survival and axonal regrowth (Leon *et al.*, 2000; Goldberg *et al.*, 2002; Yin *et al.*,

2009). Moreover, deleting intrinsic factors in RGCs such as PTEN and TSC1, which function as

negative regulators of the mTOR pathway, improves both RGC cell survival and axonal growth

(Park *et al.*, 2008; Smith *et al.*, 2009; Norsworthy *et al.*, 2017). However, enhancing RGC axonal

growth by manipulating RGC intrinsic programs frequently results in axonal misguidance as axons

project inappropriately from the optic tract before and at the optic chiasm during the initial stages

of regeneration (Kurimoto *et al.*, 2010; Luo *et al.*, 2013; Vincent Pernet *et al.*, 2013; V. Pernet *et*

al., 2013; Berry, Ahmed and Logan, 2019). This demonstrates a need to identify molecular and

cellular mechanisms that guide RGC axons as they regrow toward and across the optic chiasm

during regeneration.

Following a Central Nervous System (CNS) injury in mammals, many non-neuronal cells

create an extracellular environment that is largely inhibitory to axonal regrowth. The cellular and

molecular barriers that form within a lesion site have long been attributed to an influx of

inflammatory cells and the upregulation of inhibitory molecules not conducive for robust

regeneration (Yiu and He, 2006; Fitch and Silver, 2008). Additional studies in mice have since

revealed roles for microglia and oligodendrocytes in functional recovery and remyelination from

CNS injury (Wang *et al.*, 2020; Mendonça *et al.*, 2021; Brennan *et al.*, 2022). Yet due to the

dominance of factors inhibiting axonal regrowth, the cellular and molecular mechanisms that guide

regrowing RGC axons following axonal injury are largely unknown. Recently, spontaneous CNS

regeneration has been described in two mammalian species, the naked mole rat and the spiny

mouse, demonstrating that pro-regenerative environments that support robust CNS regeneration

are not unique to the CNS of non-mammalian vertebrates (Park *et al.*, 2017; Nogueira-Rodrigues

et al., 2022). Therefore, model organisms that exhibit spontaneous robust CNS axonal

regeneration, without an overwhelmingly inhibitory extracellular environment, provide unique

opportunities to identify mechanisms that promote CNS regeneration.

We recently developed a rapid and robust optic nerve transection assay in larval zebrafish

to take full advantage of its regenerative capabilities and optical transparency to study functional

optic nerve regeneration *in vivo* (Harvey, Baxter and Granato, 2019, 2023). At just 5 days post

fertilization (dpf), larval zebrafish already possess a functional visual system driving several

visually guided behaviors, such as prey capture (Brockerhoff *et al.*, 1995; Easter and Nicola, 1996;

Gahtan, Tanger and Baier, 2005). Following complete transections of the RGC axons in 5 dpf

larvae, we observe axonal degeneration, subsequent initiation of regrowth and reinnervation of

the optic tecta by 72 hours post transection (hpt) (Harvey, Baxter and Granato, 2019). We have

also shown that RGC axonal regeneration in larval zebrafish occurs independent of RGC death

or neurogenesis, and that visual function is restored by 8 days post transection (Harvey, Baxter

and Granato, 2019). Using this optic nerve transection assay, we performed a candidate genetic

screen and identify Lh3, a collagen-modifying glycosyltransferase whose gene name is referred

to as *lh3* or also *plod3*, to be critical for directing axonal regrowth toward the optic chiasm during

optic nerve regeneration. We show that following optic nerve transection, *lh3* expression is

upregulated near the site of axonal injury and that Lh3 functions during the time when active

regeneration occurs, specifically during the early stages of regeneration as RGC axons are

growing toward the midline. We also show that Lh3 expression in oligodendrocytes is required to

direct early regrowing RGC axons across the midline along their original path. Furthermore, we

show that mutants in *collagen 18a1 (col18a1),* a putative Lh3 substrate, display RGC axonal

misguidance phenotypes similar to *lh3* mutants, indicating at a glial-derived molecular pathway in

which Lh3 acts through Collagen18a1 to direct RGC axons. Combined, our results demonstrate

a critical pro-regenerative *in vivo* role for oligodendrocytes during early optic nerve regeneration

to direct axon regrowth toward and across the midline.

<u>Results</u>

Lh3 is required for growth across midline during regeneration

To identify molecular mechanisms that guide regrowing RGC axons across the optic

chiasm during optic nerve regeneration, we performed a candidate genetic screen consisting of

35 genes that are expressed in the visual system or have known functions in axon growth and

guidance. Specifically, we used a sharpened tungsten needle to transect the optic nerves of

genetic mutants and their siblings at 5 dpf and assessed at 72 hpt whether RGC axons had

regrown across the optic chiasm to reinnervate the optic tectum. One of the top hits from this

screen was the gene *lh3*, which encodes a multidomain glycosyltransferase that post-

translationally modifies collagen proteins (Heikkinen *et al.*, 2000; C. Wang *et al.*, 2002). RGC axon

projections in *lh3* mutants cross the midline and project to the optic tecta in a manner

indistinguishable to their siblings (Zeller *et al.*, 2002). However, we took advantage of a previously

validated transgene *Tg(hsp70:lh3-myc)*, which expresses a Myc-tagged Lh3 under the control of

the *hsp70* heat-shock-activated promoter (Isaacman-Beck *et al.*, 2015) to rescue various

morphological deficits that would affect the survivability of the larvae for the duration of the

regeneration experiments (Zeller and Granato, 1999; Taler *et al.*, 2019) (Fig. 1A). At 5 dpf,

uninjured RGC axons in both *lh3* siblings and mutants exit the eye and project across the optic

chiasm to terminate in the optic tecta (Fig. 1B-E). Following optic nerve transection, axons in *lh3*

siblings grow across the midline and reinnervate the optic tecta (Fig. 1F,H,J,L). In contrast,

transected optic nerve axons in *lh3* mutants fail to cross the midline and instead project along

aberrant trajectories, often along anterior and posterior paths or also dorsally or ventrally (Fig

1G,I,K,L, magenta arrowheads). These results reflect a functional role for Lh3 during optic nerve

regeneration in directing RGC axons towards and across the midline at the optic chiasm.

Lh3 expression is upregulated in the optic chiasm during regeneration

We next asked whether Lh3 is required during the process of optic nerve regeneration. For

this, we took advantage of the conditional *Tg(hsp70:lh3-myc)* transgene (Isaacman-Beck *et al.*,

2015). We induced Lh3 expression in 5 dpf larvae 4-5 hours prior to performing the optic nerve

transections, and then assessed axonal regrowth across the optic chiasm and reinnervation of

the optic tecta at 72 hpt (Fig. 2A). RGC axons in *lh3* siblings that did or did not receive a heat

shock regrew across the midline and innervated the optic tecta (Fig. 2B,D,F). In contrast, *lh3*

mutants without a heat shock display aberrant axonal growth, while heat shock induced Lh3

expression in *lh3* mutants restores optic nerve axonal regrowth across the midline (Fig. 2C,E-F).

These data demonstrate that Lh3 acts during the process of optic nerve regeneration.

We then sought to determine in which cell-type Lh3 was expressed during optic nerve

regeneration. We used *in situ* hybridization chain reaction to detect *lh3* mRNA expression prior to

and after optic nerve transections. Prior to transections, we failed to detect *lh3* mRNA expression

in retinas nor in the region of the optic chiasm (Fig. 2G-I; n = 17 larvae). Similarly, after transection,

we failed to detect *lh3* mRNA expression in RGC neurons in the retina (Fig. 2G-H; n = 41 larvae).

In contrast, we detected upregulated *lh3* mRNA expression around the transected proximal stump

of RGC axons and near the midline during early regeneration between 2 and 6 hpt (Fig. 2G,I; n =

30/41 larvae), demonstrating that *lh3* mRNA is upregulated following optic nerve transection in

the region of the optic chiasm.

Lh3 directs regrowing RGC axonal growth toward their original trajectory

Following optic nerve transections in wild-type larvae, we have previously observed new RGC axons emerging from the proximal stump of transected optic nerves by 24 hpt and exhibit robust growth across the optic chiasm by 48 hpt (Harvey, Baxter and Granato, 2019). To determine whether Lh3 continuously directs regenerating RGC axons once they emerge from the optic nerve stump, we used live cell imaging and examined the dynamics of regenerating RGC axons as they approach and cross the optic chiasm, which has not previously been reported. For simplicity, we removed one eye from 4 dpf larvae to only visualize the regrowth from a single retina per larva, and then performed transections to timelapse *lh3* siblings and mutants (Fig 3A, S1). Following transections on 5 dpf *lh3* sibling and mutant larvae, we observe that the portion of axons distal to the transection site undergoes Wallerian degeneration, as has been observed for other nerves in zebrafish (Fig. 3B-D,G-I, Supplementary Movies 1 and 2, brackets) (Waller, 1851; Martin *et al.*, 2010; Rosenberg *et al.*, 2012; Bremer *et al.*, 2019). In *lh3* siblings, some axons emerge from the proximal stump of RGC axons in a range of directions, often extending and retracting rapidly (Fig. 3B-C, Supplementary Movie 1). By 16 hpt, we observe that a small group of axon fascicles grows toward the midline and rapidly extends across the midline (Fig. 3D-E, Supplementary Movie 1, white arrowheads). Over the next few hours, additional fascicles of axons are observed following the growth of the axons that have crossed the midline (Fig. 3F, Supplementary Movie 1) revealing *in vivo* axonal dynamics during the early stages of optic nerve regeneration. Axons in *lh3* mutants emerge at similar timepoints to *lh3* siblings from the proximal stump (Fig 3G-H, Supplementary Movie 2). However, unlike the *lh3* siblings, axonal growth does not persist across the midline and axons often grow anteriorly and posteriorly or dorsally and ventrally around the eye (Fig. 3I-K, Supplementary Movie 2, magenta arrowheads), strongly suggesting that during the early process of optic nerve regeneration, Lh3 directs the initial regrowth of RGC axons toward the midline.

We next sought to identify the mechanisms by which Lh3 directs regrowing RGC axons

toward and across the optic chiasm. We first asked whether Lh3 is critical for the initiation of

axonal growth extending from the proximal stump of transected RGC axons, and consequently

whether optic nerve regrowth was delayed in *lh3* mutants. We found no significant difference in

the timing of initiation of axonal regrowth post transection (Fig 4A-B). To further define the

trajectories that RGC axons take as they grow across the optic chiasm, we measured the angle

between distinct axon fascicles and the original projection path before transection (Fig 4C; see

Methods for more details). In optic nerves in *lh3* siblings and mutants that regrew across the

midline, we observe that while some fascicles strayed from the pre-injury path, most fascicles

grow very close to their original projection path (Fig 4D; Supplemental Table 1). Nerves in *lh3*

siblings that did not grow toward the midline grew along a mostly single aberrant trajectory (Fig

4D; Supplemental Table 1). In contrast, in nerves that failed to cross the midline in *lh3* mutants,

we found that the aberrant paths of axon projections displayed a much greater range of angles

(Fig 4D; Supplemental Table 1). These data strongly suggest that *lh3* is required to direct RGC

axons along their regenerative path, which is very similar to their pre-transection path, towards

and across the optic chiasm.

Glial expression of Lh3 restores RGC axonal growth across the midline

Given that *lh3* mRNA expression is upregulated in area of the optic chiasm (Fig. 2I) and

that *lh3* is required *in vivo* to direct regenerating RGC axons along their original projection path

(Fig. 2F, 3, 4D), we hypothesized that the relevant cell type critical for *lh3* mediated optic nerve

regeneration would be located in the region of the optic chiasm. Transcriptomics studies in mice

have found *lh3* to be expressed most greatly in microglia (Zhang *et al.*, 2014). Moreover,

macrophages and microglia have been shown to clear debris from injured axons and can promote

axonal growth and remyelination during CNS regeneration (Rosenberg *et al.*, 2012; Wang *et al.*,

2020; Brennan *et al.*, 2022). Therefore, we first focused on macrophages and microglia as a

potential source of Lh3 activity during optic nerve regeneration. In untransected larvae, we

observe sporadic *mpeg+* cells near the optic chiasm (Fig 5A). At 24 hpt, we observe a greater

concentration of *mpeg+* cells surrounding the proximal stump of RGC axons and the new axonal

growth (Fig. 5A; n = 11 larvae). We next used a transgenic rescue approach to test whether

expression of *lh3* in macrophages/microglia is sufficient to restore axonal regrowth across the

optic chiasm in *lh3* mutants. We found that Lh3 expression from a *Tg(mpeg:lh3-myc)* transgene

fails to rescue the *lh3* mutant phenotype, where nerves in *lh3* mutants mostly display aberrant

growth similar to *lh3* mutants without the *Tg(mpeg:lh3-myc)* transgene (Fig 5D). Since RGC axons

regrow very closely to their original path, we next sought to examine oligodendrocytes associated

with the optic nerve (Brösamle and Halpern, 2002). Using both a *Tg(olig2:dsred)* transgenic line

labeling oligodendrocyte lineage cells (Fig 5B, n= 5 untransected larvae, n = 21 transected larvae)

as well as *in situ* hybridization chain reactions for *mbpa* mRNA in myelinating oligodendrocytes

(Fig 5C, n= 6 untransected larvae, n = 5 transected larvae), we observed oligodendrocytes in the

optic chiasm and associated with RGC axons before and after transection (Kucenas *et al.*, 2008;

Ackerman and Monk, 2016). We then tested whether expression of *lh3* in oligodendrocytes would

rescue axonal regrowth across the optic chiasm in *lh3* mutants. *Sox10*, like *olig2,* is a transcription

factor that mediates oligodendrocyte lineage specificity and is maintained in all oligodendrocyte

lineage cells (Bergles and Richardson, 2015; Ackerman and Monk, 2016). We find that *lh3*

expression from a *Tg(sox10:lh3-mkate)* transgene restores RGC regrowth across the midline in

otherwise mutant *lh3* larvae (Fig 5E). Furthermore, we detect *lh3* mRNA expression that overlaps

with transgenic labeling of oligodendrocyte cells, demonstrating that *lh3* is upregulated in

oligodendrocytes surrounding the proximal stump of transected RGC axons and along an axonal

projection path across the midline during optic nerve regeneration (Fig. 5F-G, Supplemental

Movie 3; n = 15/21). Altogether, these findings demonstrate that *lh3* expression in

oligodendrocytes near the proximal stump of transected RGC axons directs early regrowing RGC

axons toward the midline along their original path to reinnervate the optic tecta.

Col18a1 is also critical for directing regenerating RGC axons

We further sought to identify the relevant Lh3 substrate during optic nerve regeneration.

Given the well documented role of Lh3 in post-translationally modifying collagen proteins

(Heikkinen *et al.*, 2000; C. Wang *et al.*, 2002), we focused on collagen substrates. As part of our

initial candidate genetic screen, we examined *col4a5*, which is required for proper RGC axon

targeting of tectal layers and also functions to scaffold guidance cues during both the development

of the visual system and during motor nerve regeneration (Xiao and Baier, 2007; Xiao *et al.*, 2011;

Isaacman-Beck *et al.*, 2015; Murphy, Isaacman-Beck and Granato, 2022). We also tested *col7a1,*

which has been shown to be expressed in the eye (Wullink *et al.*, 2018). We ultimately found that

RGC axon regeneration toward and across the midline is unaffected in *col4a5* and *col7a1* mutants

(Fig. S2). We then examined *col18a1*, a collagen associated with Knobloch syndrome in humans

that is characterized by eye abnormalities (Fukai *et al.*, 2002; Suzuki *et al.*, 2002). Col18a1 has

several putative Lh3 sites within X-Lys-Gly- motifs and has been shown *in vitro* to bind to Lh3

(Risteli *et al.*, 2004; Salo and Myllyharju, 2021). First, we used *in situ* hybridization chain reaction

to detect *col18a1* mRNA expression and observe an upregulation of *col18a1* mRNA at 12 hpt in

the optic chiasm, similar to the expression pattern of *lh3* following optic nerve transection (Fig.

6A; n = 13/14 larvae). We also detect *col18a1* mRNA expression that overlaps with transgenic

labeling of oligodendrocyte cells, which demonstrates that *col18a1* is also upregulated in

oligodendrocytes surrounding the proximal stump of transected RGC axons and along axonal

projection path across the midline during optic nerve regeneration (Fig. 6B, Supplementary Movie

4). We then tested *col18a1* mutants in our optic nerve transection assay. Prior to transection, RGC

axonal projections were indistinguishable between *col18a1* siblings and mutants (Fig. 6C-F). In

contrast, at 72 hpt we observed regrowing RGC axons projecting away from the CNS midline,

similar to what we observe in *lh3* mutants where regenerating RGC axons grow along aberrant

paths (Fig 6G-K). This demonstrates that during regeneration *col18a1* is critical for directing

regenerating RGC axons toward the midline and suggests that glial-derived Lh3 acts, as least in

part, through Col18a1 during optic nerve regeneration.

Discussion

Regenerating axons often encounter multiple choice points along their paths that they must

navigate properly to robustly reinnervate their original targets. Previous studies have shown that

PTEN inactivation in RGC neurons strongly promotes RGC regrowth following an optic nerve

injury, but that these axons display axonal misguidance at the optic chiasm, proving the optic

chiasm to be one such choice point (Kurimoto *et al.*, 2010; Luo *et al.*, 2013; Vincent Pernet *et al.*,

2013; V. Pernet *et al.*, 2013; Berry, Ahmed and Logan, 2019). Here, we identify a mechanism

critical for guiding RGC axons toward and across the optic chiasm during optic nerve

regeneration. We find that *lh3* is required during regeneration to direct regenerating RGC axons

toward and across the midline. We find that following optic nerve transection, expression of *lh3*

and its presumptive substrate *col18a1* are upregulated in cells along the regenerative path near

the optic chiasm. Using live cell imaging, we demonstrate that *lh3* is required to direct regrowing

RGC axons along a regenerative trajectory very similar to their original path. Together, these

results provide compelling evidence for a RGC extrinsic molecular pathway that guides optic

axons during regeneration.

Live cell imaging reveals axonal dynamics of early optic nerve regeneration

To date, most optic nerve regeneration studies in adult rodents and fish monitor RGC

axonal regrowth with static time point analyses (Horder, 1971; Murray, 1976; V. Pernet *et al.*, 2013;

Yin and Benowitz, 2018). In contrast, peripheral nervous system regeneration, as well as spinal

cord regeneration have been visualized in live larval zebrafish using *in vivo* imaging studies (Bhatt

et al., 2004; Rosenberg *et al.*, 2012; Ceci *et al.*, 2014; Lewis and Kucenas, 2014; Isaacman-Beck

et al., 2015; Bremer *et al.*, 2019). To visualize RGC axonal dynamics during optic nerve

regeneration, we previously established an optic nerve transection assay in larval zebrafish

(Harvey, Baxter and Granato, 2019, 2023). Here, we perform live cell imaging of RGC axons

initiating regrowth and regenerating across the optic chiasm *in vivo*. These timelapses reveal two

intriguing insights into mechanisms that may mediate robust CNS regeneration in zebrafish. First,

we observe that axons distal to the transection site undergo Wallerian degeneration (Fig. 3B-D,G-

I, Movie S1 and Movie S2, brackets) (Waller, 1851). While a hallmark of the mammalian CNS is

a slower rate of axonal fragmentation following axonal injury, here we observe complete

degeneration of the distal axons by about 15 hpt (Vargas and Barres, 2007). While we did not

examine whether persistence of degenerating debris would repel axonal growth, we do observe

new axons reaching the midline around the same time axonal debris is cleared from that region

(Fig. 3D). Axonal debris and associated myelin proteins are known to inhibit axonal regeneration

(Vargas and Barres, 2007). But here, we find that Wallerian degeneration does not appear to be

delayed in *lh3* mutants, and we therefore would not hypothesize axonal debris to contribute to the

misguidance of RGC axons in *lh3* mutants.

Second, our live cell imaging reveals that a small group of axon fascicles exhibit strongly

directed growth toward the midline by 16 hpt and subsequently extends across rapidly (Fig. 3D-

E, Supplementary Movie 1, white arrowheads). Additional fascicles of axons continue to grow and

follow the growth of the axons that initially crossed the midline (Fig. 3F, Supplementary Movie 1).

These axons grow along a path that almost completely overlaps with their original path (Fig. 4D),

and strongly suggests that there are factors expressed from cells associated with that projection

that direct regrowing RGC axons during optic nerve regeneration.

A novel role for oligodendrocytes in axon guidance during axon regeneration

To determine the relevant cell type through which *lh3* is functioning during optic nerve

regeneration, we use cell-type specific transgenic rescue. Based on the *lh3* and *col18a1* mRNA

expression in the optic chiasm (Fig. 2G-I, 6A), we hypothesized that the relevant cell type would

be one found in that region during the initial stages of regeneration. We observe macrophages

and microglia dynamically infiltrating the chiasm and surrounding regenerating axonal growth (Fig.

5A). *Tg(mpeg:lh3-myc)* expression fails to rescue aberrant axonal growth in *lh3* mutants,

suggesting that *lh3* expression needs to be more specifically spatially regulated (Fig. 5D). We

therefore examined oligodendrocytes and observe oligodendrocyte lineage cells present in the

optic chiasm and we detect *mbpa* mRNA indicative of myelinating oligodendrocytes along the

degenerating axons (Fig. 5B-C). We find that *lh3* expression from a *Tg(sox10:lh3-mkate)*

transgene restores RGC regrowth across the midline in *lh3* mutants (Fig 5E). However, the

transgenic tools we employed here do not allow is to distinguish whether oligodendrocyte

precursor cells or mature oligodendrocytes were the relevant cell type for *lh3* and *col18a1,* as

sox10 and *olig2* are both transcription factors that are expressed in all oligodendrocyte lineage

cells (Bergles and Richardson, 2015; Ackerman and Monk, 2016). Previous studies implicate

oligodendrocytes to play roles in remyelination (Wang *et al.*, 2020; Mendonça *et al.*, 2021) and

axon repulsion or inhibition via myelin-associated glycoprotein (McKerracher *et al.*, 1994;

Mukhopadhyay *et al.*, 1994), NogoA (Chen *et al.*, 2000; GrandPré *et al.*, 2000), oligodendrocyte-

myelin glycoprotein (K. C. Wang *et al.*, 2002) and semaphorin 4D (Moreau-Fauvarque *et al.*,

2003). From our results, we propose a mechanism in which Lh3 and Col18a1 expressed in

oligodendrocytes mediate axon attraction to direct regenerating RGC axons along their original

path, suggesting a novel role for oligodendrocytes in CNS regeneration.

A molecular mechanism for *lh3* and *col18a1* in specifying a regenerative path

We identify *lh3* to be required to direct regenerating RGC axons toward and across the

optic chiasm (Fig 1). We then determined that *col18a1* is also critical for directing RGC axons

during optic nerve regeneration, but not another basement membrane *col4a5* or fibrillar *col7a1*,

which demonstrates a specific role for *col18a1* in optic nerve regeneration (Fig 6, Fig S2). But the

question remains: what are the mechanisms by which Lh3 and Col18a1 target RGC axons to their

original path? We cannot exclude a possible mechanism involving RGC axons directly interacting

with Col18a1 in the extracellular matrix through integrins (Elango *et al.*, 2022). Transcriptomic

analysis of RGCs at different stages of regeneration in adult zebrafish shows that several integrins

are upregulated as axons grow toward and across the midline (Dhara *et al.*, 2019). Therefore, the

spatial and temporal specificity of Lh3 and Col18a1 expression could establish the regenerative

path for regrowing RGC axons independent of any other guidance cues.

But in addition to playing structural roles, there is growing evidence showing a critical role

for extracellular matrix proteins to bind and regulate the spatial distribution of axon guidance

molecules in the extracellular space. For decades, glycosaminoglycans, such as heparan sulfate

proteoglycans, have been known to regulate the extracellular localization of Slit repulsive

guidance cues to affect axon guidance in development (Wang and Denburg, 1992; Johnson *et*

al., 2004; Smart *et al.*, 2011). More specifically, *col4a5* has been shown to be critical during

development of the retinotectal projection for localizing Slit1 expression in the superficial layers

of the optic tectum, which mediates lamina-specific axon targeting (Xiao and Baier, 2007; Xiao *et*

al., 2011). During motor nerve regeneration, spatially restricted *col4a5* and *lh3* are required in a

subset of Schwann cells to possibly create a Slit1a gradient that directs dorsal nerve axons to

their original projection paths (Isaacman-Beck *et al.*, 2015; Murphy, Isaacman-Beck and Granato,

2022). In this study, our results describe a similar mechanism where *lh3* is required to direct

regrowing RGC axons toward the midline along their original path, but here, we find in optic nerve

regeneration that Lh3 potentially acts through Col18a1, not Col4a5. What are possible guidance

cues that Col18a1 could be spatially regulating? Col18a1 has three isoforms and within the long

isoform is a Frizzled-like domain that has been shown to interact with Wnt molecules (Quélard *et*

al., 2008). Moreover, Wnt signaling has been implicated in both promoting as well as repelling

axonal growth in spinal cord and optic nerve regeneration (Yin *et al.*, 2008; Hollis and Zou, 2012;

Yam and Charron, 2013; Strand *et al.*, 2016; Patel, Park and Hackam, 2017; Wehner *et al.*, 2017).

Therefore, we propose another potential mechanism in which Lh3 is required for Col18a1

expression that possibly spatially restricts Wnts in the extracellular space along the original RGC

axonal projection to direct regenerating RGC axons during optic nerve regeneration. Altogether,

our studies demonstrate that live cell imaging of optic nerve regeneration in larval zebrafish allows

for unprecedented visualization of axonal growth dynamics and potential interactions with other

cell types in the optic chiasm. Our data here reveal a glial derived molecular pathway that directs

regenerating RGC axons across the optic chiasm during optic nerve regeneration, furthering our

understanding of the cellular and molecular mechanisms that promote robust functional optic

nerve regeneration.

Methods

Fish lines and maintenance

All zebrafish (*Danio rerio*) work was performed in compliance with the University of Pennsylvania Institutional Animal Care and Use Committee regulations and raised as previously described (Mullins *et al.*, 1994). All transgenic lines were maintained in the Tupfel long fin genetic background except the *col18a1* line, which were maintain in the AB background. The following transgenic lines and mutants were used: *lh3*^{TV2O5} (Schneider and Granato, 2006), *col18a1* (Isaacman-Beck *et al.*, 2015), *col4a5*^{s510} (Xiao and Baier, 2007), *col7a1*^{sa3473}, *Tg(hsp:lh30-myc)* (Isaacman-Beck *et al.*, 2015), *Tg(sox10:lh3-mkate)* (Isaacman-Beck *et al.*, 2015), *Tg(isl2b:GFP)* (Pittman, Law and Chien, 2008), *Tg(olig2:dsred)* (Kucenas *et al.*, 2008), *Tg*(*mpeg1:Gal4FF*)^{gl25} (Ellett *et al.*, 2011) and *Tg*(*UAS:nfsB-mCherry*)^{c264} (Davison *et al.*, 2007).

Adult zebrafish and embryos were raised at 28°C on a 14-h:10-h light:dark cycle, or 29°C for larvae in PTU E3, see Transection Assay.

To generate *Tg(mpeg1.1:lh3-myc)*, Gateway cloning was employed to combine *lh3-myc* into a pDest vector with the mpeg1.1 promoter and Tol2 transposon sites. Tol2 transgenesis was performed as previously described(Suster *et al.*, 2009) into 1-cell stage embryos from *lh3* heterozygous incrosses. Transgenic lines were identified by genotyping for the transgene with forward primer ACAGTCTCTTGCGTCATCAAAACC and reverse primer TGATCACCAGCAGCTCATTG.

Transection Assay

Optic nerve transection assays were performed as previously described(Harvey, Baxter and Granato, 2019, 2023). Briefly, to inhibit melanocyte pigmentation, larvae were raised in

phenylthiourea (PTU, 0.2mM in E3 medium) in the dark at 29°C beginning at 1 dpf. For live cell

imaging and distal transections, larvae were anesthetized in PTU E3 plus 0.0053% tricaine then

mounted in 2.5% low-melt agarose (SeaPlaque, Lonza) prepared with PTU E3 plus 0.016%

tricaine ventral-up on a glass microscopy slide. Then one eye was completely removed using

forceps at 4 dpf. Larvae recovered in Ringer's solution and then were kept in PTU E3 at 29°C

until transected. At 5 dpf, larvae were anesthetized, mounted in agarose, and then optic nerve

transections were performed on an Olympus SZX16 fluorescent microscope or a dissecting

microscope with a NightSea Stereo Microscope Fluorescence Adapter (RB-GO only wavelength

set, bandpass filter) using a sharpened tungsten needle (Fine Science Tools, Tip Diameter:

0.001mm, Rod Diameter: 0.125mm). Optic nerves were transected at the region of the nerve

distal to it exiting the eye, yet proximal to the optic chiasm. Following injuries, larvae were removed

from the agarose, allowed to recover in Ringer's solution with 0.2mM PTU, then returned to 0.2mM

PTU, E3 medium at 29°C. Larvae were inspected for transection efficiency at 16-18 hpt, except

larvae that were fixed at early post transection timepoints, and only larvae with complete optic

nerve transections with no visible intact axons remaining from the eye to the tectum were kept

until fixation at later designated timepoints.

Heat Shock-Induced *lh3* Rescue

To induce expression of Lh3 from *Tg(hsp:lh3-myc)* transgene, during development: Embryos were

developmentally staged in the late afternoon. No more than 6 embryos were placed per tube in

PCR strip-tubes. Tubes were incubated in a thermal cycler at 28°C until embryos would reach

11.66 hpf, then heated to 38°C for 10 minutes, then held at 28°C until they could be removed from

the tubes.

To induce expression of Lh3 from Tg(hsp:lh3-myc) transgene, during regeneration: Larvae at 5

dpf homozygous for the Tg(hsp:lh3-myc) transgene were placed only one per tube in PCR strip-

tubes. Tubes were incubated in a thermal cycler at 38°C for 30min then held at 29°C until they

could be removed from the tubes. Larvae were transected 4-5 hours later.

Immunostaining

Larvae were stained using methods modified from those previously described (Randlett *et al.*,

2015). Briefly, larvae were fixed in 4% PFA in PBS overnight at 4°C. Larvae were washed in PBS

+ 0.25% Triton (PBT), incubated in 150mM Tris-HCl pH 9.0 for 15 min at 70°C, then washed in

PBT. Larvae were permeabilized in 0.05% Trypsin-EDTA for 5 min on ice , washed in PBT, blocked

in PBT containing 1% bovine serum albumin (BSA), 2% normal goat serum (NGS) and 1%

dimethyl sulfoxide (DMSO), and then incubated in primary and secondary antibodies overnight at

4°C in PBT containing 1% BSA and 1% DMSO. Stained larvae were stored and mounted in

Vectashield (Vector Laboratories) for imaging using confocal microscopy. For static timepoint

analysis of *lh3* and *col18a1* mutants, primary antibodies used were mouse anti-GFP (JL-8, 1:200,

Takara Bio), and secondary antibodies used were goat anti-mouse Alexa 488 (1:500, Molecular

Probes). For static timepoint analysis of macrophages/microglia, primary antibodies used were

mouse anti-GFP (JL-8, 1:200, Takara Bio), rabbit anti-RFP (1:200, Abcam ab62341) and

secondary antibodies used were goat anti-mouse Alexa 488 (1:500, Molecular Probes) and goat

anti-rabbit Alexa 594 (1:500, Molecular Probes). For static timepoint analysis of oligodendrocytes,

primary antibodies used were rabbit anti-GFP (1:500, Molecular Probes), mouse anti-dsred

(1:500, BD Pharmingen) and secondary antibodies used were goat anti-rabbit 488 (1:500,

Molecular Probes) and goat anti-mouse Alexa 594 (1:500, Molecular Probes).

Fluorescent *in situ* Hybridization with Hybridization Chain Reaction (HCR)

Lh3 (plod3), col18a1 and mbpa mRNA expression were detected using fluorescent *in situ*

hybridization with HCR (Molecular Instruments, Los Angeles, CA, USA)(Choi *et al.*, 2018). HCR

probes, buffers, and hairpins were purchased from Molecular Instruments. *Tg(isl2b:GFP)* larvae

were fixed at 5 dpf with 4% paraformaldehyde in PBS overnight at 4°C. No antibody staining was

required together with in situ HCR. The staining was performed as previously described(Shainer

et al., 2023).

Static Confocal Imaging, Processing and Analysis

419 Larvae were imaged on a Zeiss 880 confocal microscope using a 20× objective. Airyscan super-

420 resolution settings were used to obtain Figures 5G and 6B. Image stacks of optic chiasms or tecta

421 were compressed into maximum intensity projections. All images were adjusted for brightness

422 and contrast in Adobe Photoshop CS5, and color assigned using Fiji.

Live Cell Imaging, Processing and Analysis

Larvae were anesthetized in PTU E3 with 0.0053% tricaine then mounted in 2.5% low-melt

agarose (SeaPlaque, Lonza) prepared with PTU E3 plus 0.016% tricaine ventral-up on a glass

microscopy slide. Then, a portion of the jaw, including the Meckel's cartilage, palatoquadrate, and

the most distal parts of the basihyal and ceratohyal structures, as well as one eye were carefully

removed using forceps at 4 dpf. Care was taken so as not to damage the heart. Larvae recovered

in Ringer's solution and then were kept in PTU E3 at 29°C until transected. At 5 dpf, larvae were

anesthetized again, mounted in agarose, and then optic nerve transections were performed.

Larvae recovered in Ringer's solution and then were kept in PTU E3 at 29°C until imaged for live

cell imaging. For live imaging, larvae were mounted ventral side up in 35mm glass-bottomed Petri

dishes in 1.5% low-melt agarose with 0.016% tricaine. Larvae were imaged using a Leica SP8

confocal microscope with the resonant scanner on, equipped with a petri dish heater and heated

water perfusion so fresh E3 water flowed through the petri dish and the temperature was kept

constant at around 28°C during the imaging session. Larvae were imaged from the ventral side

of the brain, which was not covered by any jaw tissue. Image stacks were acquired every 10-15

minutes with a 20x 1.0 NA lens. All images were adjusted for brightness and contrast, and color

assigned using Fiji.

To determine the time of growth initiation extending from the proximal stump of transected RGC

axons, we recorded in which frame axonal extension longer than 1um occurred that continued to

extend for another subsequent frame.

To measure the angle of the original path before transection to the end of distinct axon fascicles,

we first rotated the images so that parasphenoid cartilage was precisely aligned vertically against

a straight line drawn in Fiji. A circle with a radius of 80 uM centered where RGC axons exit the

eye was drawn within Fiji using a Make Oval Macro in timelapse image sequences. Axons and

fascicles terminating farther than the 80 uM region of interest at 20 hpt were used for measuring.

First the angle of the original path to the parasphenoid was determined using frames before

Wallerian degeneration was complete. Then the angle of the axons and fascicles to the

parasphenoid were determined, and then the difference was calculated and graphed.

Statistical Analysis

Data were imported into GraphPad Prism for analysis. Statistical analysis was performed as

indicated throughout the text.

Acknowledgments

We would like to thank Dr. Andrea Stout of the UPenn Cell and Developmental Biology Microscopy

Core, as well as Dr. Gordon Ruthel of the Penn Vet Imaging Core for providing technical

assistance. We thank all members of the Granato laboratory for comments and discussion.

Funding

This work was supported by grants from the National Institutes of Health to B.M.H. (EY032593)

and M.G. (EY024861 and NS097914).

Author Contributions

B.M.H. and M.G. conceptualized the experiments; B.M.H. M.B, and A.M.G. performed and

analyzed experiments; B.M.H. wrote the original draft. All authors contributed to editing and

revision of the paper.

Data Availability

All relevant data are within the article and its supplementary information files.

Competing interests

The authors declare no competing interests.

References

References

Ackerman, S.D. and Monk, K.R. (2016) "The scales and tales of myelination: using zebrafish
and mouse to study myelinating glia," *Brain Research*, 1641(Pt A), pp. 79–91. Available at:
https://doi.org/10.1016/j.brainres.2015.10.011.

Bergles, D.E. and Richardson, W.D. (2015) "Oligodendrocyte Development and Plasticity," *Cold
Spring Harbor Perspectives in Biology*, 8(2), p. a020453. Available at:
https://doi.org/10.1101/cshperspect.a020453.

Berkelaar, M. *et al.* (1994) "Axotomy results in delayed death and apoptosis of retinal ganglion
cells in adult rats," *The Journal of Neuroscience: The Official Journal of the Society for
Neuroscience*, 14(7), pp. 4368–4374.

Berry, M., Ahmed, Z. and Logan, A. (2019) "Return of function after CNS axon regeneration:
Lessons from injury-responsive intrinsically photosensitive and alpha retinal ganglion cells,"
Progress in Retinal and Eye Research, 71, pp. 57–67. Available at:
https://doi.org/10.1016/j.preteyeres.2018.11.006.

Bhatt, D.H. *et al.* (2004) "Cyclic AMP-induced repair of zebrafish spinal circuits," *Science (New
York, N.Y.)*, 305(5681), pp. 254–258. Available at: https://doi.org/10.1126/science.1098439.

Bremer, J. *et al.* (2019) "The ubiquitin ligase PHR promotes directional regrowth of spinal
zebrafish axons," *Communications Biology*, 2, p. 195. Available at:
https://doi.org/10.1038/s42003-019-0434-2.

Brennan, F.H. *et al.* (2022) "Microglia coordinate cellular interactions during spinal cord repair in
mice," *Nature Communications*, 13(1), p. 4096. Available at: https://doi.org/10.1038/s41467-022-
31797-0.

Brockerhoff, S.E. *et al.* (1995) "A behavioral screen for isolating zebrafish mutants with visual
system defects," *Proceedings of the National Academy of Sciences of the United States of
America*, 92(23), pp. 10545–10549.

Brösamle, C. and Halpern, M.E. (2002) "Characterization of myelination in the developing
zebrafish," *Glia*, 39(1), pp. 47–57. Available at: https://doi.org/10.1002/glia.10088.

Ceci, M.L. *et al.* (2014) "Axon-Schwann cell interactions during peripheral nerve regeneration in
zebrafish larvae," *Neural Development*, 9, p. 22. Available at: https://doi.org/10.1186/1749-8104-
9-22.

Chen, M.S. *et al.* (2000) "Nogo-A is a myelin-associated neurite outgrowth inhibitor and an
antigen for monoclonal antibody IN-1," *Nature*, 403(6768), pp. 434–439. Available at:
https://doi.org/10.1038/35000219.

Choi, H.M.T. *et al.* (2018) "Third-generation in situ hybridization chain reaction: multiplexed,
quantitative, sensitive, versatile, robust," *Development (Cambridge, England)*, 145(12), p.
dev165753. Available at: https://doi.org/10.1242/dev.165753.

Davison, J.M. *et al.* (2007) "Transactivation from Gal4-VP16 transgenic insertions for tissue-
specific cell labeling and ablation in zebrafish," *Developmental Biology*, 304(2), pp. 811–824.
Available at: https://doi.org/10.1016/j.ydbio.2007.01.033.

Dhara, S.P. *et al.* (2019) "Cellular reprogramming for successful CNS axon regeneration is
driven by a temporally changing cast of transcription factors," *Scientific Reports*, 9(1), p. 14198.
Available at: https://doi.org/10.1038/s41598-019-50485-6.

Easter, S.S. and Nicola, G.N. (1996) "The development of vision in the zebrafish (Danio rerio),"
Developmental Biology, 180(2), pp. 646–663.

Elango, J. *et al.* (2022) "The Molecular Interaction of Collagen with Cell Receptors for Biological
Function," *Polymers*, 14(5), p. 876. Available at: https://doi.org/10.3390/polym14050876.

Ellett, F. *et al.* (2011) "mpeg1 promoter transgenes direct macrophage-lineage expression in
zebrafish," *Blood*, 117(4), pp. e49-56. Available at: https://doi.org/10.1182/blood-2010-10-
314120.

Fitch, M.T. and Silver, J. (2008) "CNS injury, glial scars, and inflammation: Inhibitory extracellular
matrices and regeneration failure," *Experimental Neurology*, 209(2), pp. 294–301. Available at:
https://doi.org/10.1016/j.expneurol.2007.05.014.

Fukai, N. *et al.* (2002) "Lack of collagen XVIII/endostatin results in eye abnormalities," *The
EMBO journal*, 21(7), pp. 1535–1544. Available at: https://doi.org/10.1093/emboj/21.7.1535.

Gahtan, E., Tanger, P. and Baier, H. (2005) "Visual prey capture in larval zebrafish is controlled
by identified reticulospinal neurons downstream of the tectum," *The Journal of Neuroscience:
The Official Journal of the Society for Neuroscience*, 25(40), pp. 9294–9303. Available at:
https://doi.org/10.1523/JNEUROSCI.2678-05.2005.

Goldberg, J.L. *et al.* (2002) "Retinal ganglion cells do not extend axons by default: promotion by
neurotrophic signaling and electrical activity," *Neuron*, 33(5), pp. 689–702. Available at:
https://doi.org/10.1016/s0896-6273(02)00602-5.

GrandPré, T. *et al.* (2000) "Identification of the Nogo inhibitor of axon regeneration as a
Reticulon protein," *Nature*, 403(6768), pp. 439–444. Available at:
https://doi.org/10.1038/35000226.

Harvey, B.M., Baxter, M. and Granato, M. (2019) "Optic nerve regeneration in larval zebrafish
exhibits spontaneous capacity for retinotopic but not tectum specific axon targeting," *PloS One*,
14(6), p. e0218667. Available at: https://doi.org/10.1371/journal.pone.0218667.

Harvey, B.M., Baxter, M. and Granato, M. (2023) "Assaying Optic Nerve Regeneration in Larval
Zebrafish," *Methods in Molecular Biology (Clifton, N.J.)*, 2636, pp. 191–203. Available at:
https://doi.org/10.1007/978-1-0716-3012-9_10.

Heikkinen, J. *et al.* (2000) "Lysyl hydroxylase 3 is a multifunctional protein possessing collagen
glucosyltransferase activity," *The Journal of Biological Chemistry*, 275(46), pp. 36158–36163.
Available at: https://doi.org/10.1074/jbc.M006203200.

Hollis, E.R. and Zou, Y. (2012) "Expression of the Wnt signaling system in central nervous system axon guidance and regeneration," *Frontiers in Molecular Neuroscience*, 5, p. 5. Available at: https://doi.org/10.3389/fnmol.2012.00005.

Horder, T.J. (1971) "The course of recovery of the retinotectal projection during regeneration of the fish optic nerve," *The Journal of Physiology*, 217 Suppl, pp. 53P-54P.

Isaacman-Beck, J. *et al.* (2015) "The lh3 Glycosyltransferase Directs Target-Selective Peripheral Nerve Regeneration," *Neuron*, 88(4), pp. 691–703. Available at: https://doi.org/10.1016/j.neuron.2015.10.004.

Johnson, K.G. *et al.* (2004) "Axonal heparan sulfate proteoglycans regulate the distribution and efficiency of the repellent slit during midline axon guidance," *Current biology: CB*, 14(6), pp. 499–504. Available at: https://doi.org/10.1016/j.cub.2004.02.005.

Kucenas, S. *et al.* (2008) "CNS-derived glia ensheath peripheral nerves and mediate motor root development," *Nature Neuroscience*, 11(2), pp. 143–151. Available at: https://doi.org/10.1038/nn2025.

Kurimoto, T. *et al.* (2010) "Long-distance axon regeneration in the mature optic nerve: contributions of oncomodulin, cAMP, and pten gene deletion," *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 30(46), pp. 15654–15663. Available at: https://doi.org/10.1523/JNEUROSCI.4340-10.2010.

Leon, S. *et al.* (2000) "Lens injury stimulates axon regeneration in the mature rat optic nerve," *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 20(12), pp. 4615–4626.

Lewis, G.M. and Kucenas, S. (2014) "Perineurial glia are essential for motor axon regrowth following nerve injury," *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 34(38), pp. 12762–12777. Available at: https://doi.org/10.1523/JNEUROSCI.1906-14.2014.

Luo, X. *et al.* (2013) "Three-dimensional evaluation of retinal ganglion cell axon regeneration and pathfinding in whole mouse tissue after injury," *Experimental Neurology*, 247, pp. 653–662. Available at: https://doi.org/10.1016/j.expneurol.2013.03.001.

Marquart, G.D. *et al.* (2017) "High-precision registration between zebrafish brain atlases using symmetric diffeomorphic normalization," *GigaScience*, 6(8), pp. 1–15. Available at: https://doi.org/10.1093/gigascience/gix056.

Martin, S.M. *et al.* (2010) "Wallerian degeneration of zebrafish trigeminal axons in the skin is required for regeneration and developmental pruning," *Development (Cambridge, England)*, 137(23), pp. 3985–3994. Available at: https://doi.org/10.1242/dev.053611.

McKerracher, L. *et al.* (1994) "Identification of myelin-associated glycoprotein as a major myelin-derived inhibitor of neurite growth," *Neuron*, 13(4), pp. 805–811. Available at: https://doi.org/10.1016/0896-6273(94)90247-x.

Mendonça, H.R. *et al.* (2021) "Myelination of regenerating optic nerve axons occurs in conjunction with an increase of the oligodendrocyte precursor cell population in the adult mice,"

Brain Research Bulletin, 166, pp. 150–160. Available at: https://doi.org/10.1016/j.brainresbull.2020.11.012.

Moore, D.L. *et al.* (2009) "KLF family members regulate intrinsic axon regeneration ability," *Science (New York, N.Y.)*, 326(5950), pp. 298–301. Available at: https://doi.org/10.1126/science.1175737.

Moreau-Fauvarque, C. *et al.* (2003) "The transmembrane semaphorin Sema4D/CD100, an inhibitor of axonal growth, is expressed on oligodendrocytes and upregulated after CNS lesion," *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 23(27), pp. 9229–9239. Available at: https://doi.org/10.1523/JNEUROSCI.23-27-09229.2003.

Mukhopadhyay, G. *et al.* (1994) "A novel role for myelin-associated glycoprotein as an inhibitor of axonal regeneration," *Neuron*, 13(3), pp. 757–767. Available at: https://doi.org/10.1016/0896-6273(94)90042-6.

Mullins, M.C. *et al.* (1994) "Large-scale mutagenesis in the zebrafish: in search of genes controlling development in a vertebrate," *Current biology: CB*, 4(3), pp. 189–202.

Murphy, P.L., Isaacman-Beck, J. and Granato, M. (2022) "Robo2 Drives Target-Selective Peripheral Nerve Regeneration in Response to Glia-Derived Signals," *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 42(5), pp. 762–776. Available at: https://doi.org/10.1523/JNEUROSCI.1528-21.2021.

Murray, M. (1976) "Regeneration of retinal axons into the goldfish optic tectum," *The Journal of Comparative Neurology*, 168(2), pp. 175–195. Available at: https://doi.org/10.1002/cne.901680202.

Nogueira-Rodrigues, J. *et al.* (2022) "Rewired glycosylation activity promotes scarless regeneration and functional recovery in spiny mice after complete spinal cord transection," *Developmental Cell*, 57(4), pp. 440-450.e7. Available at: https://doi.org/10.1016/j.devcel.2021.12.008.

Norsworthy, M.W. *et al.* (2017) "Sox11 Expression Promotes Regeneration of Some Retinal Ganglion Cell Types but Kills Others," *Neuron*, 94(6), pp. 1112-1120.e4. Available at: https://doi.org/10.1016/j.neuron.2017.05.035.

Park, K.K. *et al.* (2008) "Promoting axon regeneration in the adult CNS by modulation of the PTEN/mTOR pathway," *Science (New York, N.Y.)*, 322(5903), pp. 963–966. Available at: https://doi.org/10.1126/science.1161566.

Park, K.K. *et al.* (2017) "Retinal ganglion cell survival and axon regeneration after optic nerve injury in naked mole-rats," *The Journal of Comparative Neurology*, 525(2), pp. 380–388. Available at: https://doi.org/10.1002/cne.24070.

Patel, A.K., Park, K.K. and Hackam, A.S. (2017) "Wnt signaling promotes axonal regeneration following optic nerve injury in the mouse," *Neuroscience*, 343, pp. 372–383. Available at: https://doi.org/10.1016/j.neuroscience.2016.12.020.

Pernet, Vincent *et al.* (2013) "Long-distance axonal regeneration induced by CNTF gene transfer is impaired by axonal misguidance in the injured adult optic nerve," *Neurobiology of Disease*, 51, pp. 202–213. Available at: https://doi.org/10.1016/j.nbd.2012.11.011.

Pernet, V. *et al.* (2013) "Misguidance and modulation of axonal regeneration by Stat3 and Rho/ROCK signaling in the transparent optic nerve," *Cell Death & Disease*, 4, p. e734. Available at: https://doi.org/10.1038/cddis.2013.266.

Pittman, A.J., Law, M.-Y. and Chien, C.-B. (2008) "Pathfinding in a large vertebrate axon tract: isotypic interactions guide retinotectal axons at multiple choice points," *Development (Cambridge, England)*, 135(17), pp. 2865–2871. Available at: https://doi.org/10.1242/dev.025049.

Quélard, D. *et al.* (2008) "A cryptic frizzled module in cell surface collagen 18 inhibits Wnt/beta-catenin signaling," *PloS One*, 3(4), p. e1878. Available at: https://doi.org/10.1371/journal.pone.0001878.

Randlett, O. *et al.* (2015) "Whole-brain activity mapping onto a zebrafish brain atlas," *Nature Methods*, 12(11), pp. 1039–1046. Available at: https://doi.org/10.1038/nmeth.3581.

Risteli, M. *et al.* (2004) "Characterization of collagenous peptides bound to lysyl hydroxylase isoforms," *The Journal of Biological Chemistry*, 279(36), pp. 37535–37543. Available at: https://doi.org/10.1074/jbc.M405638200.

Rosenberg, A.F. *et al.* (2012) "In vivo nerve-macrophage interactions following peripheral nerve injury," *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, 32(11), pp. 3898–3909. Available at: https://doi.org/10.1523/JNEUROSCI.5225-11.2012.

Salo, A.M. and Myllyharju, J. (2021) "Prolyl and lysyl hydroxylases in collagen synthesis," *Experimental Dermatology*, 30(1), pp. 38–49. Available at: https://doi.org/10.1111/exd.14197.

Schneider, V.A. and Granato, M. (2006) "The myotomal diwanka (lh3) glycosyltransferase and type XVIII collagen are critical for motor growth cone migration," *Neuron*, 50(5), pp. 683–695. Available at: https://doi.org/10.1016/j.neuron.2006.04.024.

Shainer, I. *et al.* (2023) "A single-cell resolution gene expression atlas of the larval zebrafish brain," *Science Advances*, 9(8), p. eade9909. Available at: https://doi.org/10.1126/sciadv.ade9909.

Smart, A.D. *et al.* (2011) "Heparan sulfate proteoglycan specificity during axon pathway formation in the Drosophila embryo," *Developmental Neurobiology*, 71(7), pp. 608–618. Available at: https://doi.org/10.1002/dneu.20854.

Smith, P.D. *et al.* (2009) "SOCS3 deletion promotes optic nerve regeneration in vivo," *Neuron*, 64(5), pp. 617–623. Available at: https://doi.org/10.1016/j.neuron.2009.11.021.

Strand, N.S. *et al.* (2016) "Wnt/β-catenin signaling promotes regeneration after adult zebrafish spinal cord injury," *Biochemical and Biophysical Research Communications*, 477(4), pp. 952–956. Available at: https://doi.org/10.1016/j.bbrc.2016.07.006.

Suster, M.L. *et al.* (2009) "Transgenesis in zebrafish with the tol2 transposon system," *Methods in Molecular Biology (Clifton, N.J.)*, 561, pp. 41–63. Available at: https://doi.org/10.1007/978-1-60327-019-9_3.

Suzuki, O.T. *et al.* (2002) "Molecular analysis of collagen XVIII reveals novel mutations, presence of a third isoform, and possible genetic heterogeneity in Knobloch syndrome," *American Journal of Human Genetics*, 71(6), pp. 1320–1329. Available at: https://doi.org/10.1086/344695.

Taler, K. *et al.* (2019) "Lysyl hydroxylase 3 is required for normal lens capsule formation and maintenance of lens epithelium integrity and fate," *Developmental Biology* [Preprint]. Available at: https://doi.org/10.1016/j.ydbio.2019.10.020.

Vargas, M.E. and Barres, B.A. (2007) "Why is Wallerian degeneration in the CNS so slow?," *Annual Review of Neuroscience*, 30, pp. 153–179. Available at: https://doi.org/10.1146/annurev.neuro.30.051606.094354.

Villegas-Pérez, M.P. *et al.* (1993) "Rapid and protracted phases of retinal ganglion cell loss follow axotomy in the optic nerve of adult rats," *Journal of Neurobiology*, 24(1), pp. 23–36. Available at: https://doi.org/10.1002/neu.480240103.

Waller, A. (1851) "Experiments on the Section of the Glosso-Pharyngeal and Hypoglossal Nerves of the Frog, and Observations of the Alterations Produced Thereby in the Structure of Their Primitive Fibres," *Edinburgh Medical and Surgical Journal*, 76(189), pp. 369–376.

Wang, C. *et al.* (2002) "The third activity for lysyl hydroxylase 3: galactosylation of hydroxylysyl residues in collagens in vitro," *Matrix Biology: Journal of the International Society for Matrix Biology*, 21(7), pp. 559–566.

Wang, J. *et al.* (2020) "Robust Myelination of Regenerated Axons Induced by Combined Manipulations of GPR17 and Microglia," *Neuron*, 108(5), pp. 876-886.e4. Available at: https://doi.org/10.1016/j.neuron.2020.09.016.

Wang, K.C. *et al.* (2002) "Oligodendrocyte-myelin glycoprotein is a Nogo receptor ligand that inhibits neurite outgrowth," *Nature*, 417(6892), pp. 941–944. Available at: https://doi.org/10.1038/nature00867.

Wang, L. and Denburg, J.L. (1992) "A role for proteoglycans in the guidance of a subset of pioneer axons in cultured embryos of the cockroach," *Neuron*, 8(4), pp. 701–714. Available at: https://doi.org/10.1016/0896-6273(92)90091-q.

Wehner, D. *et al.* (2017) "Wnt signaling controls pro-regenerative Collagen XII in functional spinal cord regeneration in zebrafish," *Nature Communications*, 8(1), p. 126. Available at: https://doi.org/10.1038/s41467-017-00143-0.

Wullink, B. *et al.* (2018) "Type VII Collagen in the Human Accommodation System: Expression in Ciliary Body, Zonules, and Lens Capsule," *Investigative Ophthalmology & Visual Science*, 59(2), pp. 1075–1083. Available at: https://doi.org/10.1167/iovs.17-23425.

Xiao, T. *et al.* (2011) "Assembly of lamina-specific neuronal connections by slit bound to type IV collagen," *Cell*, 146(1), pp. 164–176. Available at: https://doi.org/10.1016/j.cell.2011.06.016.

Xiao, T. and Baier, H. (2007) "Lamina-specific axonal projections in the zebrafish tectum require
the type IV collagen Dragnet," *Nature Neuroscience*, 10(12), pp. 1529–1537. Available at:
https://doi.org/10.1038/nn2002.

Yam, P.T. and Charron, F. (2013) "Signaling mechanisms of non-conventional axon guidance
cues: the Shh, BMP and Wnt morphogens," *Current Opinion in Neurobiology*, 23(6), pp. 965–
973. Available at: https://doi.org/10.1016/j.conb.2013.09.002.

Yin, Y. *et al.* (2009) "Oncomodulin links inflammation to optic nerve regeneration," *Proceedings*
of the National Academy of Sciences of the United States of America, 106(46), pp. 19587–
19592. Available at: https://doi.org/10.1073/pnas.0907085106.

Yin, Y. and Benowitz, L.I. (2018) "In Vitro and In Vivo Methods for Studying Retinal Ganglion Cell
Survival and Optic Nerve Regeneration," *Methods in Molecular Biology (Clifton, N.J.)*, 1695, pp.
187–205. Available at: https://doi.org/10.1007/978-1-4939-7407-8_16.

Yin, Z.-S. *et al.* (2008) "Repair effect of Wnt3a protein on the contused adult rat spinal cord,"
Neurological Research, 30(5), pp. 480–486. Available at:
https://doi.org/10.1179/174313208X284133.

Yiu, G. and He, Z. (2006) "Glial inhibition of CNS axon regeneration," *Nature Reviews.*
Neuroscience, 7(8), pp. 617–627. Available at: https://doi.org/10.1038/nrn1956.

Zeller, J. *et al.* (2002) "Migration of zebrafish spinal motor nerves into the periphery requires
multiple myotome-derived cues," *Developmental Biology*, 252(2), pp. 241–256.

Zeller, J. and Granato, M. (1999) "The zebrafish diwanka gene controls an early step of motor
growth cone migration," *Development (Cambridge, England)*, 126(15), pp. 3461–3472.

Zhang, Y. *et al.* (2014) "An RNA-sequencing transcriptome and splicing database of glia,
neurons, and vascular cells of the cerebral cortex," *The Journal of Neuroscience: The Official*
Journal of the Society for Neuroscience, 34(36), pp. 11929–11947. Available at:
https://doi.org/10.1523/JNEUROSCI.1860-14.2014.

Figure 1.

Figure 1. Lh3 is required for growth across the midline during regeneration. (A) Timeline of

a transection experiment with *lh3* larvae (hpf, hours post fertilization; dpf, days post fertilization;

hpt, hours post transection). (B-C) Before injury, RGC axons labeled by *Tg(isl2b:GFP)* in siblings

and conditional *lh3* mutants cross the midline and (D-E) innervate contralateral tecta. (F-I) In

contrast to *lh3* siblings at 72 hpt, RGC axons in *lh3* mutants turn away from the midline, project

aberrantly (magenta arrowheads) and do not reinnervate contralateral optic tecta. (B-I) Dashed

lines indicate the outline of the eyes; scale bars = 50 μm (J-K) Schematics of RGC axonal growth

at 72 hpt for (J) *lh3* siblings and (K) *lh3* mutants. (L) Quantification of the percent of nerves at 72

hpt that grew across the midline to reinnervate the optic tectum, did not grow across the midline

and instead grew along aberrant trajectories or showed stalled axonal growth. Numbers indicate

total number of transected nerves. **** *P<0.0001*, Chi squared test. Representative images of

chiasms and tecta for each timepoint are of the same fixed larva, though across timepoints are

different larvae.

Figure 2.

Figure 2. *Lh3* expression is upregulated in the optic chiasm during early regeneration. (A)

Timeline of induced Lh3 expression with *Tg(hsp70:lh3-myc)* during development as well as

regeneration (hpf, hours post fertilization; dpf, days post fertilization; hpt, hours post transection).

(B-C) At 72 hpt in *lh3* siblings with no Heat Shock induction, RGC axons cross the midline to

innervate the optic tecta, while RGC axons in *lh3* mutants turn away from the midline and project

aberrantly (C, magenta arrowheads). (D-E) In *lh3* siblings and mutants that received a heat shock,

RGC axons project across the midline (E, white arrowheads) and innervate the optic tecta.

Dashed lines indicate the outline of the eyes; scale bar = 50 μm. (F) Quantification of the percent

of nerves at 72 hpt that grew across the midline to reinnervate the optic tectum, did not grow

across the midline and instead grew along aberrant trajectories or showed stalled axonal growth

in *Tg(isl2b:GFP)* larvae with and without a regeneration heat-shock. Numbers indicate total

number of transected nerves. *****$P<0.0001$*, ***$P<0.01$*, chi squared test. (G-I) *Lh3* mRNA

expression in *Tg(isl2b:GFP)* larvae pre-transection and at 2-6 hpt. (G) White dashed boxes are

areas of the retina shown in (H) and the optic chiasm shown in (I). Scale bars, (G) 50 μm, (H) 20

μm, (I) 50 μm. Axons in the chiasm region at 2-6 hpt have not yet fully degenerated following

transection. Dashed lines indicate the outline of the eyes in (I). Representative images across

timepoints are of different larvae.

Figure 3.

Figure 3. *Lh3* **directs the initial regrowing RGC axons.** (A) Schematic of larvae expressing

Tg(isl2b:GFP) to label RGCs that were used for *in vivo* imaging. Larvae have one eye removed

one day prior to timelapse imaging. Red box indicates the region imaged in panels B-K. (B,G)

Transected RGC axons separate from the distal portion of the optic nerve that will eventually

degenerate (indicated by brackets in B-D, G-I). (C) In *lh3* siblings, RGC axons initiate growth out

from the proximal stump. (D) A small group of axons (white arrowhead) project directly toward the

770   midline along the original path of the projection and then (E-F) subsequently additional axonal

771   growth follows (n = 7/10 nerves). (H) In *lh3* mutants, axons grow out from the proximal stump, but

772   (I-K) persist along aberrant trajectories (magenta arrowheads; n = 6/12 nerves). See also Figure

773   S1. Dashed lines indicate the outline of the eyes; scale bar = 25 μm.

Figure 4.

Figure 4. *Lh3* **is required to direct regrowing RGC axons along their original path.** (A)

Representative frames from live timelapses of larvae expressing *Tg(isl2b:GFP)* to label RGCs

showing the initial axonal growth that emerges from the injured proximal stump of axons. hpt =

hours post transection. (B) Comparison of the time when axon fascicles first emerge from the

proximal stump in *lh3* siblings and mutants. Time was determined by the first frame when axon

fascicles were observed extending from the proximal stump that extended for at least one other

consecutive frame. Data are represented as mean ± SD; n.s.*P>0.05*, t-test. (C) Representative

image of a regenerating nerve at 20 hpt and the mark-up to measure the angle of growing fascicles

from the injured stump of axons. The magenta line marks the parasphenoid cartilage. See

Methods for more details on quantifications. (D) Quantification at 20 hpt of the absolute angle of

axon fascicles from the end of a projection to the path of the original axonal projection. (Tectal

Innervation Across Midline, sibling n = 6 nerves, mutants n = 3 nerves; Aberrant Axonal Growth,

sibling n = 3 nerves, mutants n = 6 nerves). Individual dots are measured fascicle absolute angles,

lines represent means; one-way ANOVA, n.s.*P>0.05*, * P<0.05. See Supplemental Table 1 for

individual angles.

Figure 5.

Figure 5. Lh3 in *sox10* expressing glia restores RGC axonal growth across the midline. (A)

In wild-type larvae expressing *Tg(isl2b:GFP)* to label RGCs, macrophages and microglia labeled

by *Tg*(*mpeg1:Gal4FF*);*Tg*(*UAS:nfsB-mCherry*) are observed in the area of the optic chiasm before

transection. At 24 hours post transection (hpt), an influx of *mpeg+* cells is seen. (B-C)

Oligodendrocyte lineage cells labeled by *Tg(olig2:dsred)* as well as *mbpa* mRNA expression in

myelinating oligodendrocytes are also seen in the chiasm before and after transection, near the

injured proximal stump of RGC axons or along the degenerating original projection. (D)

Quantification of the percent of nerves at 72 hpt that grew across the midline to reinnervate the

optic tectum, did not grow across the midline and instead grew along aberrant trajectories or

showed stalled axonal growth with and without *Tg(mpeg:lh3-myc)* transgene. Numbers indicate

total number of transected nerves. *****P<0.0001*, n.s. *P>0.05*, chi squared test. (E) Quantification

of regeneration with and without *Tg(sox10:lh3-mkate)* transgene. Numbers indicate total number

of transected nerves. *****P<0.0001*, n.s. *P>0.05*, chi squared test. (F) *Lh3* mRNA expression at

4-6 hpt is upregulated and overlaps with oligodendrocytes labeled by *Tg(olig2:dsred)*. scale bars

= 100um (G) Additional images of individual oligodendrocytes labeled by *Tg(olig2:dsred)* co-

expressing *lh3* mRNA (see also Supplementary Movie 3). scale bars = 10um. Representative

images across timepoints are of different larvae. Axons in the chiasm region at 4-6 hpt have not

yet fully degenerated following transection. Dashed lines indicate the outline of the eyes.

Figure 6.

Figure 6. The Lh3 candidate substrate Col18a1 is upregulated after transection and directs

RGC axons across the midline during regeneration. (A) *Col18a1* mRNA expression pre-

transection and at 12 hours post transection (hpt) in larvae expressing *Tg(isl2b:GFP)* to label

RGCs. Axons in the chiasm region at 12 hpt have not yet fully degenerated following transection.

(B) Additional images of individual oligodendrocytes labeled by *Tg(olig2:dsred) co-expressing lh3*

mRNA (see also Supplementary Movie 4). scale bars = 10um. (C-D) Before injury, RGC axons

labeled by *Tg(isl2b:GFP)* in siblings and *col18a1* mutants cross the midline and (E-F) innervate

contralateral tecta. (G-J) In contrast to *col18a1* siblings at 72 hpt, RGC axons in *col18a1* mutants

turn away from the midline, project aberrantly (magenta arrowheads) and do not reinnervate

contralateral optic tecta. Dashed lines indicate the outline of the eyes; scale bar = 50 μm. (K)

Quantification of the percent of nerves at 72 hpt that grew across the midline to reinnervate the

optic tectum, did not grow across the midline and instead grew along aberrant trajectories or

showed stalled axonal growth. Numbers indicate total number of transected nerves. *****P<0.0001*,

chi squared test. Representative images of chiasms and tecta for each timepoint are of the same

fixed larva, though across timepoints are different larvae.

Supplemental Figure 1.

S1 Figure. Related to Figures 3 and 4. Quantification of the percent of nerves from timelapsed

larvae that grew across the midline to reinnervate the optic tectum, did not grow across the

midline and instead grew along aberrant trajectories, or did not exhibit any axonal growth, ie.

stalled, at 72 hpt. Numbers indicate total number of transected nerves. *P<0.00001*, Fisher's

exact test.

Supplemental Figure 2.

S2 Figure. ***Col4a5*** **and** ***col7a1*** **are not required to direct regrowing RGC axons to the midline**

during regeneration. Quantification of the percent of nerves that grew across the midline to

reinnervate the optic tectum, did not grow across the midline and instead grew along aberrant

trajectories, or did not exhibit any axonal growth, ie. stalled, at 72 hpt. Numbers indicate total

number of transected nerves. n.s.*P>0.05*, chi-square test.

Supplemental Information

Supplementary Movie 1. Related to Figure 2, contains the movie which corresponds to the *lh3*

sibling axonal regeneration time series shown in Figure 2B-F.

Supplementary Movie 2. Related to Figure 2, contains the movie which corresponds to the *lh3*

sibling axonal regeneration time series shown in Figure 2G-K.

Supplementary Movie 3. Related to Figure 5G, representative three-dimensional rotation from a

confocal z-stack demonstrating co-expression of *lh3* mRNA with *Tg(olig2:dsred)*

oligodendrocytes.

Supplementary Movie 4. Related to Figure 6B, representative three-dimensional rotation from a

confocal z-stack demonstrating co-expression of *col18a1* mRNA with *Tg(olig2:dsred)*

oligodendrocytes.

Supplemental Table 1. Related to Figure 4, contains raw data of individual angles of fascicles

from nerves that grew either across the midline or aberrantly from *lh3* sibling and mutant larvae.