Title:

β-adrenergic blockers increase cAMP and stimulate insulin secretion through a

PKA/RYR2/TRPM5 pathway in pancreatic β-cells in vitro

Naoya Murao, Risa Morikawa, Yusuke Seino, Kenju Shimomura, Yuko Maejima, Yuichiro

Yamada, Atsushi Suzuki

Department of Endocrinology, Diabetes and Metabolism, Fujita Health University, School of

Medicine, Toyoake, Japan

Corresponding author

Naoya Murao, M.D., Ph.D.

Departments of Endocrinology, Diabetes and Metabolism, Fujita Health University, School

of Medicine, 1-98 Dengakugakubo, Toyoake, Aichi 4701192 Japan

Phone: +81-562-93-2699

Fax: +81-562-95-1879

E-mail: muraonaoy@gmail.com

Running title

β-blockers increase cAMP in pancreatic β-cells

Authors

Naoya Murao[1,2], Risa Morikawa[1], Yusuke Seino[1,2], Kenju Shimomura[3], Yuko Maejima[3],

Yuichiro Yamada[1,2], Atsushi Suzuki[1]

[1] Department of Endocrinology, Diabetes and Metabolism, Fujita Health University,

School of Medicine, Toyoake, Japan

[2] Yutaka Seino Distinguished Center for Diabetes Research, Kansai Electric Power

Medical Research Institute, Kyoto, Japan

[3] Department of Bioregulation and Pharmacological Medicine, Fukushima Medical

University School of Medicine, Fukushima, Japan

Corresponding author

Naoya Murao, M.D., Ph.D.

Departments of Endocrinology, Diabetes and Metabolism, Fujita Health University, School

of Medicine, 1-98 Dengakugakubo, Toyoake, Aichi 4701192 Japan

Phone: +81-562-93-2699

Fax: +81-562-95-1879

E-mail: muraonaoy@gmail.com

Word counts

Number of text pages:　　　20

Number of tables:　　　　　1

Number of figures: 7

Number of references: 58

Number of words

Abstract   152

Introduction   312

Discussion   813

Abbreviations

AC, adenylyl cyclase

BHQ, 2,5-di-*tert*-butylhydroquinone

cAMP, adenosine 3',5'-Cyclic Monophosphate

CICR, Ca^{2+}-induced Ca^{2+} release

ER, endoplasmic reticulum

FKBP, FK-506-binding protein

GPCR, G-protein coupled receptor

GLP-1, glucagon-like peptide

GSIS, glucose-stimulated insulin secretion

iAUC, incremental area under the curve

IP1, inositol 1-phosphate

IP_3, inositol 1,4,5-trisphosphate

K_{ATP} channel, ATP-sensitive K^+ channel

Propranolol-SIS, propranolol-stimulated insulin secretion

PKA, protein kinase A

PKC, protein kinase C

PLCβ, phospholipase Cβ

SERCA, sarcoplasmic/endoplasmic reticulum Ca^{2+} ATPase

TRPM, Transient receptor potential cation channel subfamily M

VDCC, voltage-dependent Ca^{2+} channel

β-blocker-SIS, β-blocker-stimulated insulin secretion

Abstract

β-adrenergic blockers (β-blockers) are extensively used to inhibit β-adrenoceptor activation

and subsequent cAMP production in many cell types. In this study, we characterized the

effects of β-blockers on mouse pancreatic β-cells. Unexpectedly, high doses (100 μM) of β-

blockers (propranolol and bisoprolol) led to a 5–10 fold increase in cAMP levels, enhanced

intracellular influx, and stimulated a 2–4 fold increase in glucose- and glimepiride-induced

insulin secretion in MIN6-K8 clonal β-cells and isolated mouse pancreatic islets. These

effects were observed despite minimal expression of β-adrenoceptors in these cells.

Mechanistically, cAMP increase led to ryanodine receptor 2 (RYR2) phosphorylation via

protein kinase A (PKA), triggering Ca^{2+}-induced Ca^{2+} release (CICR). CICR then activates

transient receptor potential cation channel subfamily M member 5 (TRPM5), resulting in

increased Ca^{2+} influx via voltage-dependent Ca^{2+} channels. These effects contradict the

conventional understanding of the pharmacology of β-blockers, highlighting the variability in

β-blocker actions depending on the experimental context.

Graphical abstract

At high concentrations (> 10 μM), the β-adrenergic blocker propranolol paradoxically

increased intracellular cAMP levels in pancreatic β-cells. This leads to PKA-induced RYR2

phosphorylation and extracellular Ca^{2+} influx, leading to CICR from the ER. CICR activated

TRPM5, which augmented β-cell electrical activity, extracellular Ca^{2+} influx, and insulin

secretion.

Introduction

β-adrenergic blockers (β-blockers) are widely used for experimental and clinical purposes.

The primary mechanism of β-adrenoceptors involves their coupling with Gαs proteins in

trimeric G protein complexes, which subsequently stimulate adenylyl cyclase (AC) to

generate adenosine 3',5'-Cyclic Monophosphate (cAMP) (Brodde et al., 2006). β-blockers

function by blocking the binding of catecholamines to β1- or β2-adrenoceptors, thus

suppressing cAMP generation (Paul, 2021). Nevertheless, their mechanisms of action in cells

outside the cardiovascular system are not well understood.

Pancreatic β-cells secrete insulin to maintain blood glucose levels within a physiological

range. Glucose is the primary stimulator of insulin secretion; glucose metabolism leads to an

increase in intracellular ATP levels, which inhibits ATP-sensitive K^+ (K_{ATP}) channels,

resulting in membrane depolarization and opening of voltage-dependent Ca^{2+} channels,

increasing intracellular Ca^{2+} ($[Ca^{2+}]_i$) levels and thereby stimulating insulin granule

exocytosis (Henquin, 2000; Henquin, 2009).

Among the various adrenoceptor subtypes, β-cells primarily express α2A-adrenoceptors

(Rorsman and Ashcroft, 2018). It is generally recognized that adrenaline suppresses insulin

release from β-cells (Schuit and Pipeleers, 1986; Henquin, 2021; Sluga et al., 2022) by

stimulating α2-adrenoceptors and the trimeric G protein Gαi/o, which in turn inhibits the

production of adenosine 3',5'-Cyclic Monophosphate (cAMP) by adenylyl cyclase (AC)

(Straub and Sharp, 2012). Conversely, studies have shown that β-adrenoceptor agonists (β-

agonists) increase insulin levels in the blood of mice and humans when administered in vivo.

This effect is counteracted by β2-selective β-blockers (Porte, 1967; Cerasi et al., 1969; Imura

et al., 1971; Loubatières et al., 1971; Ahren and Lundquist, 1981). However, these findings

were not reproduced in vitro.

Here, we describe an unexpected effect of β-blockers on β-cells in vitro: when

administered at high concentrations (> 10 μM), β-blockers significantly increased cAMP

levels, $[Ca^{2+}]_i$, and insulin release in mouse β-cells. Our findings highlight the variable effects

of β-blockers depending on the biological context, emphasizing the need for caution when

using these compounds for experimental purposes.

Materials and Methods

Animals

C57BL/6JJcl (RRID:IMSR_JCL:JCL:MIN-0003) mice were purchased from CLEA Japan

(Tokyo, Japan) and used for experiments at 19–20 weeks of age. NSY.B6-*Tyr+*, *A^y*/Hos mice

were a gift from Hoshino Laboratory Animals (Ibaraki, Japan) and were used for experiments

at 19–20 weeks of age. Mice were maintained under specific-pathogen-free conditions at 23

± 2 °C and 55 ± 10% relative humidity with 12-hour light-dark cycles (8am–8pm), with free

access to water and standard chow CE-2 (CLEA Japan, Tokyo, Japan). The health status of

the mice was checked regularly. All experiments were performed using male mice. All

animal experiments were performed with the approval APU22080 by the Institutional Animal

Care and Use Committee of Fujita Health University, complying with the Guidelines for

Animal Experimentation at Fujita Health University and current Japanese guidelines and

regulations for scientific and ethical animal experimentation (Kagiyama et al., 2006).

Cell lines

MIN6-K8 cells were generated as previously described (Iwasaki et al., 2010) and were kindly

provided by Professor Junichi Miyazaki (Osaka University). *Kcnj11^{-/-}* β-cells (clone name:

*Kcnj11^{-/-}*βCL1) were established by sub-cloning MIN6-K8 cells transfected with Cas9

nickase and guide RNA pairs targeting mouse *Kcnj11*, as described previously (Oduori et al.,

2020). All cells were cultured in Dulbecco's modified Eagle's medium (DMEM) containing

4500 mg/L glucose (Sigma-Aldrich, St. Louis, MO, USA, Cat# D5796) supplemented with

10% fetal bovine serum (FBS) (BioWest, Nuaillé, France, Cat# S1400-500) and 5 ppm 2-

mercaptoethanol. The cells were maintained at 37 °C with 5% CO_2.

Reagents

2.8 mM glucose-supplemented Krebs-Ringer bicarbonate buffer-HEPES (2.8G-KRBH) was

prepared by supplementing KRBH (133.4 mM NaCl, 4.7 mM KCl, 1.2 mM KH_2PO_4, 1.2 mM

$MgSO_4$, 2.5 mM $CaCl_2$, 5 mM $NaHCO_3$, 10 mM HEPES, pH 7.4) with 0.1% bovine-serum

albumin (Sigma-Aldrich, Cat# A6003) and 2.8 mM glucose. Ca^{2+}-free KRBH was

formulated by replacing $CaCl_2$ with an equimolar $MgCl_2$ and adding 0.2 mM EGTA

(NACALAI TESQUE, Kyoto, Japan, Cat# 15214-21). For extracellular Na^+ depletion

experiments, control buffer (150 mM NaCl, 5 mM KCl, 1.2 mM KH_2PO_4, 1 mM $MgCl_2$, 2.2

mM $CaCl_2$, 10 mM HEPES, pH 7.4) and Na^+-free buffer [150 mM N-Methyl-D-glucamine

(Tokyo Chemical Industry, Tokyo, Japan, CAS: 6284-40-8, Cat# M0227), 5 mM KCl, 1.2

mM KH_2PO_4, 1 mM $MgCl_2$, 2.2 mM $CaCl_2$, 10 mM HEPES, pH 7.4] were prepared. 30 mM

Na^+ buffer was formulated by mixing the control buffer and Na^+-free buffer at a ratio of 1:4

and supplementing 0.1% bovine-serum albumin (Sigma-Aldrich, Cat# A6003) and 2.8 mM

glucose.

The following reagents were added to KRBH during the stimulation period: 11.1 mM

glucose, (±)-propranolol hydrochloride (FUJIFILM Wako Pure Chemical, Osaka, Japan,

CAS: 318-98-9, Cat# 168-28071), carteolol hydrochloride (Tokyo Chemical Industry, CAS:

51781-21-6, Cat# C3015), nadolol (Sigma-Aldrich, CAS: 42200-33-9, Cat# N1892), timolol

maleate (Tokyo Chemical Industry, CAS: 26921-17-5, Cat# T2905), bisoprolol hemifumarate

(Tokyo Chemical Industry, CAS: 104344-23-2, Cat# B3994), metoprolol tartrate (Tokyo

Chemical Industry, Tokyo, Japan, CAS: 56392-17-7, Cat# M2555), glimepiride (Tokyo

Chemical Industry, CAS: 93479-97-1, Cat# G0395), and carbamylcholine chloride

(carbachol) (Tokyo Chemical Industry, CAS: 51-83-2, Cat# C0596).

The following reagents were added to KRBH during the pre-incubation and stimulation

periods: nifedipine (FUJIFILM Wako Pure Chemical, Osaka, Japan, Cat# 14505781),

thapsigargin (FUJIFILM Wako Pure Chemical, Osaka, Japan, CAS: 67526-95-8, Cat# 209-

17281), MDL-12330A hydrochloride (Sigma-Aldrich, CAS: 40297-09-4, Cat# M184),

SQ22536 (Selleck Chemicals, Houston, TX, USA, CAS: 17318-31-9, Cat# S8283), H-89

hydrochloride (Cayman Chemical, Ann Arbor, MI, USA, CAS: 130964-39-5, Cat#

10010556), phentolamine mesylate (Tokyo Chemical Industry, CAS: 65-28-1, Cat# P1985),

atipamezole hydrochloride (FUJIFILM Wako Pure Chemical, Osaka, Japan, CAS: 104075-

48-1, Cat# 015-25331), dantrolene sodium salt hydrate (Tokyo Chemical Industry, CAS:

14663-23-1, Cat# D3996), triphenylphosphine oxide (Tokyo Chemical Industry, CAS: 791-

28-6, Cat# T0625), YM-254890 (FUJIFILM Wako Pure Chemical, Osaka, Japan, CAS:

568580-02-9, Cat# 257-00631), t-butylhydroquinone (Tokyo Chemical Industry, CAS: 1948-

33-0, Cat# 027-07212), and xestospongin C (FUJIFILM Wako Pure Chemical, Osaka, Japan,

CAS: 244-00721, Cat# C0598). The reagents used for stimulation were stored as a 1000×

concentrate in dimethyl sulfoxide (DMSO) (FUJIFILM Wako Pure Chemical, Osaka, Japan,

Cat# 041-29351) and diluted with KRBH shortly before the experiment. An equal volume of

DMSO was added to the vehicle control.

Isolation of pancreatic islets from mice

Digesting solution was formulated by supplementing 0.1w/v% Collagenase from Clostridium

histolyticum (Sigma-Aldrich, Cat# C6885) to Hanks' balanced salt solution (136.9 mM NaCl,

5.4 mM KCl, 0.8 mM $MgSO_4$, 0.3 mM Na_2HPO_4, 0.4 mM KH_2PO_4, 4.2 mM $NaHCO_3$, 10

mM HEPES, 1.3 mM $CaCl_2$, 2 mM glucose). The mice were euthanized by isoflurane

exposure. Pancreas was digested by 10-min incubation at 37 °C following intraductal

injection of digesting solution. Islets were hand-picked and separated from exocrine tissues,

transferred to 60-mm non-treated plates (AGC Techno Glass, Shizuoka, Japan, Cat# 1010-

060), and cultured overnight in RPMI-1640 (Sigma-Aldrich, Cat# R8758) supplemented with

10% FBS (BioWest, Nuaillé, France, Cat# S1400-500) and 1% penicillin-streptomycin

solution (FUJIFILM Wako Pure Chemical, Osaka, Japan, Cat# 168-23191) at 37 °C and 5%

CO_2 before the experiments.

Insulin secretion and content

Insulin secretion was measured using the static incubation method, as described previously

(Murao et al., 2022; Murao et al., 2024b). To measure insulin secretion from cell lines, the

cells were seeded in 24-well plates (Corning, Glendale, AZ, USA, Cat# 353047) at a density

of 5×10^5 cells/well and cultured for 48 h. The cells were pre-incubated for 30 min with 300

μL/well of 2.8G-KRBH and subsequently stimulated with 300 μL/well of 2.8G-KRBH

containing the specified stimulations for 30 min at 37 °C. The supernatant was collected for

the measurement of secreted insulin, after which the cells were extracted with 0.1% Triton-X

in 2.8G-KRBH for the measurement of insulin content. Insulin was quantified using a

homogeneous time-resolved fluorescence assay (HTRF) Insulin Ultrasensitive kit (Revvity,

Waltham, MA, USA, Cat# 62IN2PEH) in accordance with the manufacturer's instructions.

To measure insulin secretion from islets, overnight cultured islets were rinsed twice with

2.8G-KRBH, followed by pre-incubation with 2.8G-KRBH for 30 min at 37 °C. Size-

matched islets were hand-picked and dispensed in a 96-well plate (Corning, Cat# 353072) at

5 islets/well. KRBH (100 μL/well) containing the specified stimulations was added and

incubated for 30 min at 37 °C. The supernatant was subjected to insulin quantification as

described above.

Measurement of inositol 1-phosphate (IP1)

Intracellular inositol 1-phosphate content was measured as described previously (Murao et al.,

2024a). Briefly, MIN6-K8 cells were seeded in a 96-well plate (Corning, Cat# 353072) at a

density of 1.0×10^5 cells/well and cultured for 48h. The measurement was performed using

IP1 assay buffer (146 mM NaCl, 4.2 mM KCl, 50 mM LiCl, 1 mM CaCl2, 0.5 mM MgCl2,

10 mM HEPES, 0.1% BSA, pH7.4) supplemented with 2.8 mM glucose (2.8G-IP1 assay

buffer). The cells were pre-incubated for 30 min with 30 μL/well of 2.8G-IP1 assay buffer,

followed by the addition of 10 μL/well of IP1 assay buffer containing the specified

stimulations at 4× concentration, and incubation for another 30 min at 37 °C. The reaction

was terminated by the addition of 10 μL/well lysis buffer. The lysate (30 μL/well) was

transferred to a 384-well plate (Corning, Cat #3826) and subjected to IP1 quantification using

the HTRF IP-One Gq Detection Kit (Revvity, Waltham, MA, USA, Cat# 62IPAPEB)

according to the manufacturer's instructions.

Measurement of cAMP

To prevent cAMP hydrolysis, 2.8G-KRBH was supplemented with 100 μM

isobutylmethylxanthine (IBMX) (FUJIFILM Wako Pure Chemical, CAS: 28822-58-4, Cat#

095-03413) for pre-incubation and stimulation. To measure cAMP levels in MIN6-K8 cells,

the cells were seeded in a 96-well plate (Corning, Cat# 353072) at a density of 1.0×10^5

cells/well and cultured for 48h. The cells were pre-incubated for 30 min with 40 μL/well of

2.8G-KRBH, followed by the addition of 20 μL/well of 2.8G-KRBH containing the specified

stimulations at 4× concentration, and incubation for another 30 min at 37 °C. The reaction

was terminated by the addition of 20 μL/well lysis buffer. The lysate (20 μL/well) was

transferred to a 384-well plate (Corning, Cat #3826) and subjected to cAMP quantification

using the HTRF cAMP Gs Dynamic Detection Kit (Revvity, Waltham, MA, USA, Cat#

62AM4PEB) according to the manufacturer's instructions.

To measure cAMP levels in islets, overnight cultured islets were rinsed twice with 2.8G-

KRBH. Size-matched islets were hand-picked and dispensed in a 96-well plate (Corning,

Cat# 353072) at 5 islets/well, followed by pre-incubation with 30 μL/well of 2.8G-KRBH for

30 min at 37 °C. Subsequently, 10 μL/well of 2.8G-KRBH containing the specified

stimulations at 4× concentration was added to the wells and incubated for another 30 min at

37 °C. The reaction was terminated by the addition of 15 μL/well lysis buffer. The lysate (20

μL/well) was transferred to a 384-well plate and subjected to cAMP quantification, as

described above.

Ca^{2+} imaging

Ca^{2+} imaging was performed as previously described (Murao et al., 2024b). Briefly, MIN6-

K8 cells were seeded in a 35 mm glass-bottom dish (Matsunami Glass, Osaka, Japan, Cat#

D11530H) at a density of 1.28×10^5 cells/dish and cultured for 48 h. Subsequently, the cells

were loaded with 1 μM Fluo-4 AM (Dojindo, Kumamoto, Japan, Cat# F312) in 2.8G-KRBH

for 20 min at 37 °C in room air. The cells were loaded with 1 mL of fresh 2.8G-KRBH, and

basal recordings were performed for 300 s (from time -300 to 0). Following the addition of 1

mL KRBH supplemented with stimulations at a 2× concentration, recordings were resumed

for another 600 s (from time 0 to 600). Time-lapse images were obtained using a Zeiss LSM

980 Airyscan2 inverted confocal laser scanning super-resolution microscope equipped with a

Plan Apo 40×, 1.4 Oil DICII objective lens (Carl Zeiss Microscopy, Jena, Germany). The

cells were excited at 488 nm laser with 0.3% output power, and fluorescence emission was

measured at 508-579 nm. The obtained images were analyzed using the ZEN 3.0 imaging

software (Carl Zeiss Microscopy, Jena, Germany, RRID:SCR_021725). Cells were randomly

chosen for analysis for each stimulation, and the number of cells analyzed is indicated in the

figure legends. The fluorescence intensity of the entire cell body (F) was monitored and

normalized to the average fluorescence intensity between -300 and 0 s (F0).

274 Gene silencing using small interfering RNA (siRNA)

Gene silencing was performed as previously described (Murao et al., 2024b). The following

siRNAs were purchased from Dharmacon (Lafayette, CO, USA): *Adra2a* (Cat# M-062217-

01-0005), *Itpr1* (Cat# M-040933-01-0005), *Trpm4* (Cat# M-056098-01-0005), *Trpm5* (Cat#

M-046645-01-0005), and non-targeting siRNA (Cat# D-001206-14-50). Briefly, siRNAs

were reverse-transfected using DharmaFECT 2 transfection reagent (Dharmacon, Lafayette,

CO, USA, Cat# T-2002-03) according to the manufacturer's instructions. The final

concentrations of siRNA and DharmaFECT 2 were 40 nM and 0.4%, respectively. For

insulin secretion experiments, cells were seeded in 24-well plates (Corning, Cat# 353047) at

5×10^5 cells/500 μL/well. For immunoblotting, cells were seeded in 12-well plates (Corning,

Cat# 353043) at 1×10^6 cells/mL/well.

286 Reverse transcription quantitative polymerase chain reaction (RT-qPCR)

RT-PCR was performed as previously described (Murao et al., 2024b). Briefly, cDNA was

prepared using CellAmp Direct Lysis and RT set (Takara Bio, Shiga, Japan, Cat# 3737S/A)

according to the manufacturer's instructions. Quantitative real-time PCR was performed on a

QuantStudio 7 Flex system (Thermo Fisher Scientific, Waltham, MA, USA,

RRID:SCR_020245) using TaqMan Universal Master Mix II with UNG (Thermo Fisher

Scientific, Waltham, MA, USA, Cat# 4440038) and Taqman probes as follows: *Adra2a* (Cat#

Mm07295458_s1), *Itpr1* (Cat# Mm00439907_m1), *Trpm4* (Cat# Mm00613173_m1), *Trpm5*

(Cat# Mm01129032_m1), and TATA-box binding protein (*Tbp*, Cat# Mm01277042_m1).

Relative gene expression was calculated using the $2^{-\Delta\Delta CT}$ method and normalized to *Tbp*.

297   Immunoblotting

Immunoblotting was performed as previously described (Murao et al., 2024b), with slight

modifications. Briefly, cells were lysed with 50 μL/well RIPA buffer supplemented with a

complete protease inhibitor cocktail (Sigma-Aldrich, Cat# 11697498001) and PhosSTOP

(Roche, Basel, Switzerland, Cat# 4906845001). For the detection of ADRA2A, the lysate

was separated on a 7.5% polyacrylamide-SDS gel and transferred to a PVDF membrane. The

membranes were probed with polyclonal anti-ADRA2A rabbit antibody (Thermo Fisher

Scientific, Waltham, MA, USA, Cat# PA1-048, RRID:AB_2225243) overnight at 4 °C. To

detect RYR2 phosphorylation, the lysate was separated on a 5% polyacrylamide-SDS gel and

transferred to a PVDF membrane. The membranes were probed with either an anti-RYR2

rabbit monoclonal antibody (Abcam, Cambridge, UK, Cat# EPR26288-70, 1:300) or an anti-

phospho RyR2 (Ser2808) rabbit polyclonal antibody (Millipore, Burlington, MA, USA, Cat#

ABS2231, 1:300) overnight at 4 °C. The membrane was then probed with HRP-conjugated

polyclonal swine anti-rabbit antibody (Agilent Technologies, Santa Clara, CA, USA, Cat#

P0399, RRID:AB_2617141, 1:2000) and visualized using ECL Prime (Cytiva,

Buckinghamshire, UK, Cat# RPN2232). To visualize the whole protein, the transferred

membrane was stained with Ponceau-S solution (Beacle, Kyoto, Japan, Cat# BCL-PSS-01).

Images were taken using ImageQuant 800 (Cytiva, Buckinghamshire, UK). The images were

quantified   using   ImageJ   (version   1.53k,   https://imagej.nih.gov/ij/index.html,

RRID:SCR_003070).

Statistical Analysis

Sample sizes were estimated from the expected effect size based on previous experiments.

No randomization or blinding was used. For cell line experiments, including insulin secretion,

IP1 measurement, cAMP measurement, RT-qPCR, and immunoblotting, n represents the

number of biological replicates of cells grown in different wells of the same multiwell plate.

Results were confirmed by 2–3 independent experiments performed on different days using

different cell passages. For islet experiments, n represents the number of wells, each

containing five islets. The number of mice used is indicated in the figure legends. Results

were confirmed by two independent experiments performed on different days using different

mice. For Ca^{2+} imaging, n represents the number of different single cells analyzed. Data are

shown as the mean ± standard deviation (SD) along with the plot of individual data points.

For statistical comparisons between two groups, a two-tailed unpaired Welch's unpaired t-

test was used. For comparisons between more than three groups, Welch's one-way analysis

of variance (ANOVA) was followed by pairwise comparisons corrected using Dunnett's

method. Normality of the distribution was confirmed by the Shapiro-Wilk test. P-values are

indicated in the figures. P-values less than 0.05 were considered statistically significant. The

statistical analyses used are indicated in the figure legends. Statistical analyses were

performed using GraphPad Prism 10 (GraphPad Software, Boston, MA, USA,

RRID:SCR_002798).

Results

β-Blockers amplify insulin secretion from MIN6-K8 β-cell lines

To evaluate the impact of β-blockers on insulin release, we used MIN6-K8, a mouse β-cell

line (Iwasaki et al., 2010). The cells were stimulated with propranolol (non-selective β-

blocker) and bisoprolol (β1-selective blocker). In the presence of stimulatory levels (11.1

mM) of glucose, both propranolol and bisoprolol dose-dependently increased insulin

secretion at concentrations greater than 10 μM and 100 μM, respectively (Figure 1, A–B).

At basal glucose levels (2.8 mM), propranolol had no effect on insulin secretion (Figure

1C) or insulin content (Figure S1A). When co-administered with sulfonylurea (K_{ATP} channel

inhibitor) glimepiride, propranolol amplified insulin secretion even at 2.8 mM glucose

(Figure 1C). To evaluate whether propranolol-stimulated insulin secretion (propranolol-SIS)

results from the promotion of K_{ATP} channel closure, its efficacy was examined in $Kcnj11^{-/-}$ β-

cells. As $Kcnj11$ encodes the pore-forming subunit of the K_{ATP} channel, $Kcnj11^{-/-}$ β-cells lack

K_{ATP} channel activity, and their cell membrane is continuously depolarized regardless of

extracellular glucose levels (Oduori et al., 2020). Propranolol increased insulin secretion in

$Kcnj11^{-/-}$ β-cells even at 2.8 mM glucose (Figure 1D), indicating that the primary mechanism

for propranolol-stimulated insulin secretion is not through facilitating K_{ATP} channel closure.

Rather, our results suggest that membrane depolarization by high glucose, sulfonylureas, or

$Kcnj11$ knockout is permissive for propranolol-SIS. Consequently, in subsequent experiments,

β-blocker-stimulated insulin secretion (β-blocker-SIS) was examined in the presence of high

glucose or glimepiride. We found that a wide range of β-blockers, both non-selective and β1-

selective, enhanced insulin release, with nadolol as the sole exception (Figure S1B).

Therefore, the capacity to promote insulin secretion is likely a common feature among

compounds in the β-blocker class.

β-Blockers increase cAMP levels in MIN6-K8 β-cell lines

To elucidate the mechanism of β-blocker-SIS, we investigated whether β-blockers modulate

canonical GPCR signaling pathways in MIN6-K8 cells.

Propranolol and bisoprolol dose-dependently increased intracellular cAMP levels in

MIN6-K8 cells at concentrations that were effective for insulin secretion (Figure 2, A–B).

Two adenylyl cyclase (AC) inhibitors, MDL-12330A and SQ-22536, suppressed cAMP

accumulation induced by 100 μM propranolol and 1 mM bisoprolol (Figure 2C and S2A),

indicating that β-blocker-induced cAMP accumulation is mediated by adenylyl cyclase

activation. Furthermore, these inhibitors also abolished propranolol-SIS (Figure 2, D–E),

indicating that cAMP production via AC is necessary for propranolol-SIS. Notably,

propranolol-SIS was also abolished by H-89, a protein kinase A (PKA) inhibitor (Figure 2F),

suggesting that the cAMP/PKA pathway is essential for propranolol-SIS.

Subsequently, inositol 1-phosphate (IP1), a metabolite of inositol 1,4,5-trisphosphate (IP_3),

was measured to assess the activity of the Gαq/phospholipase Cβ (PLCβ) signaling pathway.

IP1 levels remained unaltered by propranolol, although substantial accumulation of IP1 was

confirmed using carbachol as a positive control (Figure S2B), indicating that propranolol

does not activate the Gαq/PLCβ pathway. This finding is corroborated by the observation that

YM-254890, a specific inhibitor of Gαq/11/14 (Takasaki et al., 2004; Patt et al., 2021),

suppressed glimepiride-induced insulin secretion (Figure S2C), but minimally affected

propranolol-SIS when expressed as a fold-change (Figure S2D). Collectively, these results

demonstrate that propranolol-SIS involves the cAMP/PKA pathway, but not the Gαq/PLCβ

pathway.

Effect of β-blockers in mouse islets

We investigated whether these findings could be applied to primary β-cells. We observed

enhancement of insulin release by propranolol in C57BL/6J (B6) mouse islets when exposed

to 11.1 mM glucose (Figure 3, A–B) or glimepiride (Figure 3C). These findings are consistent

with the results obtained using MIN6-K8 cells. NSY.B6-*Tyr*+, A^y (NSY.B6-A^y) is a

spontaneous type 2 diabetes mouse strain established by crossing the NSY and B6J- A^y

strains, introducing the obesity-related agouti-yellow (A^y) mutation in the agouti (a) gene into

a diabetes-prone NSY background (Ohno et al., 2022). A^y heterozygous male mice (NSY.B6-

A^y/a) exhibited obesity and hyperglycemia; wild-type male mice (NSY.B6-*a/a*) served as lean

controls (Ohno et al., 2022; Murao et al., 2024b). At 2.8 mM glucose, no propranolol-SIS

was observed in either strain (Figure 3, D–E), confirming the glucose dependency of

propranolol-SIS (Figure 1G). Propranolol substantially increased insulin secretion in diabetic

A^y/a islets (Figure 3E), although it did not reach statistical significance in lean *a/a* islets

(Figure 3D). Notably, propranolol also increased intra-islet cAMP levels in both strains

(Figure 3, F–G). These findings indicate that the observed effects are likely a general

characteristic of β-blockers on β-cells rather than an artifact specific to a cell line.

Propranolol potentiates Ca^{2+} response via PKA/RYR2 pathway

To clarify whether propranolol-SIS involves potentiation of intracellular Ca^{2+} ([Ca^{2+}]$_i$), we

conducted Fluo-4 imaging in MIN6-K8 cells. Propranolol enhanced the glimepiride-induced

increase in $[Ca^{2+}]_i$ (Figure 4, A–C). Glimepiride alone elicited oscillatory $[Ca^{2+}]_i$ waves,

whereas a combination of glimepiride and propranolol induced a rapid increase in $[Ca^{2+}]_i$,

followed by rapid oscillations (Figure 4B).

Notably, propranolol alone elicited a transient $[Ca^{2+}]_i$ increase at 0 min, followed by

sporadic spikes (Figure 4, D–E). This propranolol-stimulated Ca^{2+} response was further

investigated to elucidate the involvement of extracellular Ca^{2+} and intracellular Ca^{2+} stores.

Nifedipine, an inhibitor of L-type voltage-dependent Ca^{2+} channels (VDCC), suppressed the

sporadic $[Ca^{2+}]_i$ spikes but not the $[Ca^{2+}]_i$ increase at 0 min indicated by the black arrow

(Figure 4F), suggesting that the $[Ca^{2+}]_i$ spikes represent Ca^{2+} influx through L-type VDCCs.

In contrast, removal of extracellular Ca^{2+} from the stimulation buffer inhibited both the

$[Ca^{2+}]_i$ increase at 0 min and subsequent $[Ca^{2+}]_i$ spikes (Figure 4G). Depletion of intracellular

Ca^{2+} stores   with   2,5-di-tert-butylhydroquinone   (BHQ),   an   inhibitor   of

sarcoplasmic/endoplasmic reticulum Ca^{2+} ATPase (SERCA), exerted the same effect (Figure

4H). These results suggest that the initial $[Ca^{2+}]_i$ increase involves both $[Ca^{2+}]_i$ mobilization

from the endoplasmic reticulum (ER) and extracellular Ca^{2+} influx and is required for the

later VDCC-mediated $[Ca^{2+}]_i$ spikes. Importantly, nifedipine, removal of extracellular Ca^{2+},

and BHQ also abolished propranolol-SIS (Figure S3, A–C), demonstrating that the

propranolol-stimulated Ca^{2+} response is essential for propranolol-SIS.

The mobilization of Ca^{2+} from the ER is mediated by specific ion channels, such as

ryanodine receptors (RYRs) and IP_3 receptors (IP_3Rs), which release Ca^{2+} into the cytosol in

response to signaling events (Guerrero-Hernández et al., 2020). The involvement of RYRs

was investigated using dantrolene, an RYR inhibitor. Dantrolene negated the capacity of

propranolol to potentiate the glimepiride-induced Ca^{2+} response in MIN6-K8 cells (Figure 5,

A–B). Dantrolene paradoxically increased glimepiride-induced insulin secretion, consistent

with a previous report (Figure S3D) (Pian-Smith et al., 1986); however, it decreased

propranolol-IIS, as expressed by the fold change (Figure 5C), suggesting that the

propranolol-stimulated Ca^{2+} response and insulin secretion are dependent on RYR. It has

been reported that the open probability of RYR2 is increased by intracellular cAMP through

PKA-mediated RYR2 phosphorylation at Ser-2808 (Marx et al., 2000; Wehrens et al., 2006).

Remarkably, phosphorylation of RYR2 at Ser-2808 was substantially induced by propranolol

alone and further enhanced by the addition of glimepiride (Figure 5, D–E). These findings

suggest that the propranolol-stimulated Ca^{2+} response is mediated by PKA-induced RYR2

phosphorylation, corroborating the observation that propranolol-SIS is dependent on PKA

activity (Figure 2F).

The involvement of IP$_3$Rs was investigated using xestospongin C, an IP$_3$R inhibitor (Gafni

et al., 1997), and siRNA-mediated knockdown of IP$_3$ receptor 1 (Itpr1). Xestospongin C

marginally decreased propranolol-SIS (Figure S3, E–F). However, depletion of *Itpr1* by

approximately 50% at the transcript level did not affect propranolol-SIS, as expressed by the

fold-change (Figure S3, G–I). These findings suggest that IP$_3$ receptors play a limited, if any,

role in propranolol-SIS.

Involvement of TRPM5 in propranolol-stimulated Ca^{2+} response

We investigated the mechanism by which PKA-mediated $[Ca^{2+}]_i$ mobilization from the ER

leads to VDCC activation. Interestingly, a reduction in extracellular Na^+ from 150 mM to 30

mM suppressed the propranolol-stimulated Ca^{2+} response and insulin secretion in MIN6-K8

cells (Figure 6, A–B), suggesting the involvement of Na^+-permeable channels. Transient

receptor potential cation channel subfamily M member 4 (TRPM4) and member 5 (TRPM5)

are Ca^{2+}-activated channels that permeate monovalent cations, including Na^+ and K^+. They

are gated by $[Ca^{2+}]_i$, most notably Ca^{2+} from the ER (Zhang et al., 2007; Gonzales et al.,

2010). Notably, TRPM5 has been implicated in the augmentation of glucose-induced

depolarizing currents in β-cells (Colsoul et al., 2010), suggesting its potential role as a

mediator linking $[Ca^{2+}]_i$ mobilization from the ER to depolarization. To elucidate the

involvement of TRPM4 and TRPM5, we conducted siRNA-mediated knockdown, which

depleted their mRNA levels by approximately 50% (Figure S4, A–B). Knockdown of *Trpm4*

and *Trpm5* led to slight and moderate decreases in propranolol-SIS, respectively (Figure 6C,

Figure S4C). Furthermore, triphenylphosphine oxide (TPPO), a TRPM5 inhibitor (Palmer et

al., 2010), inhibited the later Ca^{2+} spikes and insulin secretion (Figure 6, D–E). These

findings suggest that TRPM5 is involved in the later phase of the propranolol-stimulated Ca^{2+}

response, contributing to propranolol-SIS.

α2-adrenoceptors are involved in cAMP elevation by propranolol

Using previously published in-house RNA-seq datasets (Hashim et al., 2018), we

investigated the expression of adrenoceptor subtypes in β-cells. MIN6-K8 cells exclusively

expressed α2A-adrenoceptors, while the other subtypes were undetectable (Table 1). Similar

patterns were observed in B6 mouse islets, which exhibited high expression of α2A-

adrenoceptors and minimal levels of α1, β1, and β2 receptors, potentially originating from

non-β-cells. These findings align with previous research indicating that mouse β-cells

exclusively express α2A-adrenoceptors (Rorsman and Ashcroft, 2018).

Interestingly, Pretreatment with atipamezole (α2-adrenoceptor antagonist) and

phentolamine (α1/2-adrenoceptor antagonist) attenuated propranolol-stimulated cAMP

elevation, as expressed by the fold-change (Figure 7, A–B). These inhibitors also abolished

propranolol-SIS (Figure 7, C–D). Although phentolamine elevated glimepiride-induced

insulin secretion, presumably due to its direct inhibitory effect on K_{ATP} channels (Plant and

Henquin, 1990; Proks and Ashcroft, 1997), it potently suppressed propranolol-SIS (Figure

7D). To further validate the involvement of α2A-adrenoceptors, we performed siRNA-

mediated Adra2a knockdown. Although the knockdown efficiency was only moderate at the

protein level (28%, as shown in Figure S5, A–B), propranolol-SIS was decreased as

expressed by the fold change (Figure 7E, S5C). These results indicate that α2-adrenoceptors

are the only adrenoceptor subtype expressed in mouse β-cells, and they play a permissive role

in cAMP elevation and insulin secretion by β-blockers.

Discussion

This study reports a paradoxical off-target effect of β-blockers on mouse pancreatic β-cells:

β-blockers increased cAMP levels, $[Ca^{2+}]_i$, and insulin secretion despite the absence of β-

adrenoceptors in these cells. We used propranolol to elucidate the signaling pathway of β-

blocker-stimulated insulin secretion, which involves cAMP/PKA/RYR2/TRPM5.

Despite the widespread use of β-blockers in clinical and experimental settings, their

concentrations vary significantly. Clinical pharmacokinetic research indicates that in healthy

subjects, the maximum plasma concentration (Cmax) of propranolol rarely surpasses 1 μM

after a single oral administration (Abdu-Aguye et al., 1986; Taegtmeyer et al., 2014; Kim et

al., 2022). The therapeutic range for propranolol is typically considered to be 192 to 385 nM

(0.05 to 0.10 μg/mL), with toxic effects occurring above 7 μM (2 μg/mL) and fatal outcomes

at levels exceeding 11 μM (3 μg/mL) (Kerns, 2007; Sharifpour et al., 2022). Similarly, the in

vivo Cmax of bisoprolol generally remains below 100 nM (Li et al., 2012; Tjandrawinata et

al., 2012).

In experimental contexts, propranolol exhibits near-complete inhibition of β2-

adrenoceptors at nanomolar levels (IC50 < 1 nM) (Boursier et al., 2020). Nevertheless, some

cell experiments have used propranolol at concentrations exceeding 100 μM. These studies

involved various cell types, including hemangioma cells (Munabi et al., 2015; Kum and Khan,

2014), clear cell renal cell carcinoma cells (Shepard et al., 2018), and PC3 and MCF7 cancer

cell lines (Reyes-Corral et al., 2019). Our findings suggest that caution should be exercised

when using high concentrations of β-blockers, as they may produce off-target effects,

contrary to conventional β-blocker pharmacology. Indeed, Reyes-Corral et al. reported that

50 μM propranolol triggered CICR in PC3 and MCF7 cancer cell lines, although they did not

examine cAMP levels (Reyes-Corral et al., 2019).

Signaling pathways linking cAMP and insulin secretion have been studied extensively

(Seino and Shibasaki, 2005; Tengholm and Gylfe, 2017). However, this is the first report of

the PKA/RYR2/TRPM5 pathway. Notably, propranolol evoked a Ca^{2+} response in MIN6-K8

cells even under low-glucose conditions, in which cells are electrically silent. The Ca^{2+}

response comprised an immediate transient increase at 0 min and subsequent spikes, the latter

indicative of Ca^{2+} influx through L-type VDCCs. We demonstrated that the initial increase in

$[Ca^{2+}]_i$ involves $[Ca^{2+}]_i$ mobilization through RYR2, which is facilitated by PKA-mediated

RYR2 phosphorylation at Ser 2808. Supporting our findings, it has been suggested that RYR2

activation in β-cells requires cAMP-dependent phosphorylation (Islam et al., 1998). However,

that report also indicates that cAMP-dependent phosphorylation alone is insufficient to

mobilize Ca^{2+} from the ER when cytosolic $[Ca^{2+}]$ is low (Islam et al., 1998; Islam, 2002);

instead, it only sensitizes RYR2 to Ca^{2+}-induced Ca^{2+} release (CICR) by releasing the

channel from Mg^{2+}-mediated inhibition (Hain et al., 1994) or FKBP12.6 (FK-506-binding

protein 12.6) binding (Marx et al., 2000). Thus, it is reasonable to postulate that propranolol-

stimulated cAMP triggers both initial Ca^{2+} influx and RYR2 phosphorylation, leading to

CICR, reflected in the transient rise in $[Ca^{2+}]_i$ at 0 min. The mediator of propranolol-

stimulated initial Ca^{2+} influx is currently unclear. However, given that it was inhibited by

depletion of extracellular Ca^{2+} and Na^+, but not by nifedipine, cAMP-activated nifedipine-

insensitive Ca^{2+} channels, such as T-type VDCCs, may be involved (Louiset et al., 2017;

Sharma et al., 2023). A similar mechanism has been proposed for glucagon-like peptide 1

(GLP-1)-stimulated insulin secretion: GLP-1-stimulated cAMP production leading to CICR

through a dual effect on Ca^{2+} influx and RYR2 sensitization (Holz et al., 1999).

TRPM5 and/or TRPM4 have been implicated in background inward currents produced by

GLP-1 (Shigeto et al., 2015) and fructose (Kyriazis et al., 2012). TRPM5 functions

downstream of Gαq- or gustducin-coupled GPCR signaling cascades, wherein PLCβ

activation triggers Ca^{2+} release from the IP_3 receptor and protein kinase C (PKC) activation,

both of which facilitate TRPM5 gating (Shigeto et al., 2015). Our findings suggest that

TRPM5 can also be activated downstream of the cAMP/PKA signaling cascade, highlighting

its versatile role in linking GPCR signaling and membrane depolarization. Small background

inward currents, including those from TRPM5, are incapable of inducing depolarization

unless the majority of β-cell KATP channels are closed (Rorsman and Ashcroft, 2018). This

mechanism may explain why membrane depolarization plays a permissive role in

propranolol-SIS.

The exact mechanism by which β-blockers elevate cAMP levels remains unclear. In our

cellular model, α2-adrenoceptors were the sole adrenoceptor subtype expressed and played a

permissive role in propranolol-induced cAMP elevation and insulin secretion. This finding

indicates that propranolol may potentially interact with α2-adrenoceptors, although such an

effect has not been previously documented. Notably, isoproterenol, a β-adrenoceptor agonist,

was found to inhibit GIIS at high concentrations (10–50 μM), and this effect was prevented

by the preinhibition of α2-adrenoceptors (Zielmann et al., 1985).

In summary, we revealed that β-blockers paradoxically increase cAMP levels in β-cells,

highlighting the variable nature of β-blocker effects, which are potentially influenced by

concentration and cell type. These findings underscore the importance of careful

consideration in experimental design when working with β-blockers.

Acknowledgements

The authors extend their gratitude to President Yutaka Seino of Kansai Electric Power

Hospital for his generous support in this research. The authors thank Yoshikazu Hoshino of

Hoshino Laboratory Animals, Inc. for providing NSY.B6-A^y mice. The authors thank

Shihomi Sakai, Asami Yamaguchi, and Megumi Akiyama for their excellent technical

assistance.

Data Availability

The authors declare that all data supporting the findings of this study are contained within the

paper and Supplemental Data.

Authorship Contributions

Participated in research design: Murao.

Conducted experiments: Murao, Morikawa, Shimomura, and Maejima.

Performed data analysis: Murao.

Wrote or contributed to the writing of the manuscript: Murao, Seino, Yamada, and Suzuki.

Footnotes

This study was supported by JSPS KAKENHI Grant Numbers JP22K20869 and JP23K15401

for N.M. Research grants for N.M. were provided by the Japan Association for Diabetes

Education and Care, Daiwa Securities Foundation, Suzuken Memorial Foundation, Japan

Diabetes Foundation, The Hori Sciences and Arts Foundation, Manpei Suzuki Diabetes

Foundation, and Fujita Academy.

Conflict of Interest Statement

No author has any actual or perceived conflicts of interest with the contents of this article.

References

1. Abdu-Aguye I, Dunlop D, Patel P, Turner P. Propranolol pharmacokinetics during the
menstrual cycle. Postgraduate Medical Journal. 1986;62(734):1093-1095.

2. Ahrén B, Lundquist I. Effects of selective and non-selective β-adrenergic agents on insulin
secretion in vivo. European journal of pharmacology. 1981;71(1):93-104.

3. Boursier ME, Levin S, Zimmerman K, et al. The luminescent HiBiT peptide enables
selective quantitation of G protein–coupled receptor ligand engagement and internalization
in living cells. Journal of Biological Chemistry. 2020;295(15):5124-5135.

4. Brodde O-E, Bruck H, Leineweber K. Cardiac adrenoceptors: physiological and
pathophysiological relevance. Journal of pharmacological sciences. 2006;100(5):323-337.

5. Cerasi E, Effendic S, Luft R. Role of adrenergic receptors in glucose-induced insulin
secretion in man. The Lancet. 1969;294(7615):301-302.

6. Colsoul B, Schraenen A, Lemaire K, et al. Loss of high-frequency glucose-induced Ca2+
oscillations in pancreatic islets correlates with impaired glucose tolerance in Trpm5–/–
mice. Proceedings of the National Academy of Sciences. 2010;107(11):5208-5213.

7. Gafni J, Munsch JA, Lam TH, et al. Xestospongins: potent membrane permeable blockers
of the inositol 1, 4, 5-trisphosphate receptor. Neuron. 1997;19(3):723-733.

8. Gonzales AL, Amberg GC, Earley S. Ca2+ release from the sarcoplasmic reticulum is
required for sustained TRPM4 activity in cerebral artery smooth muscle cells. American
Journal of Physiology-Cell Physiology. 2010;299(2):C279-C288.

9. Guerrero-Hernández A, Sánchez-Vázquez VH, Martínez-Martínez E, et al. Sarco-
endoplasmic reticulum calcium release model based on changes in the luminal calcium
content. Calcium Signaling. 2020:337-370.

10.   Hain J, Onoue H, Mayrleitner M, Fleischer S, Schindler H. Phosphorylation
Modulates the Function of the Calcium Release Channel of Sarcoplasmic Reticulum from
Cardiac Muscle (∗). Journal of Biological Chemistry. 1995;270(5):2074-2081.

11.   Hashim M, Yokoi N, Takahashi H, et al. Inhibition of SNAT5 induces incretin-
responsive state from incretin-unresponsive state in pancreatic β-cells: study of β-cell
spheroid clusters as a model. Diabetes. 2018;67(9):1795-1806.

12.   Henquin J-C. Triggering and amplifying pathways of regulation of insulin secretion
by glucose. Diabetes. 2000;49(11):1751-1760.

13.   Henquin J-C. Regulation of insulin secretion: a matter of phase control and amplitude
modulation. Diabetologia. 2009;52(5):739-751.

14.   Henquin J-C. Non-glucose modulators of insulin secretion in healthy humans:(dis)
similarities between islet and in vivo studies. Metabolism. 2021;122:154821.

15.   Holz GG, Leech CA, Heller RS, Castonguay M, Habener JF. cAMP-dependent
mobilization of intracellular Ca2+ stores by activation of ryanodine receptors in pancreatic
β-cells: a Ca2+ signaling system stimulated by the insulinotropic hormone glucagon-like
peptide-1-(7–37). Journal of Biological Chemistry. 1999;274(20):14147-14156.

16.   Imura H, Kato Y, Ikeda M, Morimoto M, Yawata M. Effect of adrenergic-blocking
or-stimulating agents on plasma growth hormone, immunoreactive insulin, and blood free
fatty acid levels in man. The Journal of Clinical Investigation. 1971;50(5):1069-1079.

17.   Islam MS. The ryanodine receptor calcium channel of β-cells: molecular regulation
633   and physiological significance. Diabetes. 2002;51(5):1299-1309.

634   18.   Islam MS, Leibiger I, Leibiger B, et al. In situ activation of the type 2 ryanodine
635   receptor in pancreatic beta cells requires cAMP-dependent phosphorylation. Proceedings
636   of the National Academy of Sciences. 1998;95(11):6145-6150.

637   19.   Iwasaki M, Minami K, Shibasaki T, Miki T, Miyazaki Ji, Seino S. Establishment of
638   new clonal pancreatic β‐cell lines (MIN6‐K) useful for study of incretin/cyclic
639   adenosine monophosphate signaling. Journal of Diabetes Investigation. 2010;1(4):137-142.

640   20.   Kagiyama N, Ikeda T, Nomura T. Japanese Guidelines and regulations for scientific
641   and ethical animal experimentation. In Vivo Models of Inflammation: Volume I. Springer;
642   2006:187-191.

643   21.   Kerns II W. Management of β-adrenergic blocker and calcium channel antagonist
644   toxicity. Emergency medicine clinics of North America. 2007;25(2):309-331.

645   22.   Kim T-E, Kang J-S, An W, Sohn JH. Systemic exposure to propranolol in patients
646   with chronic liver disease and its correlation with portal blood flow. Frontiers in Medicine.
647   2022;9:973606.

648   23.   Kum JJ, Khan ZA. Propranolol inhibits growth of hemangioma-initiating cells but
649   does not induce apoptosis. Pediatric research. 2014;75(3):381-388.

650   24.   Kyriazis GA, Soundarapandian MM, Tyrberg B. Sweet taste receptor signaling in
651   beta cells mediates fructose-induced potentiation of glucose-stimulated insulin secretion.
652   Proceedings of the National Academy of Sciences. 2012;109(8):E524-E532.

653   25.   Li G-f, Wang K, Chen R, Zhao H-r, Yang J, Zheng Q-s. Simulation of the
654   pharmacokinetics of bisoprolol in healthy adults and patients with impaired renal function
655   using whole-body physiologically based pharmacokinetic modeling. Acta Pharmacologica
656   Sinica. 2012;33(11):1359-1371.

657   26.   Loubatières A, Mariani M, Sorel G, Savi L. The action of beta-adrenergic blocking
658   and stimulating agents on insulin secretion. Characterization of the type of beta receptor.
659   Diabetologia. 1971;7(3):127-132.

660   27.   Louiset E, Duparc C, Lenglet S, Gomez-Sanchez CE, Lefebvre H. Role of
661   cAMP/PKA pathway and T-type calcium channels in the mechanism of action of serotonin
662   in human adrenocortical cells. Molecular and cellular endocrinology. 2017;441:99-107.

663   28.   Marx SO, Reiken S, Hisamatsu Y, et al. PKA phosphorylation dissociates FKBP12. 6
664   from the calcium release channel (ryanodine receptor): defective regulation in failing
665   hearts. Cell. 2000;101(4):365-376.

666   29.   Munabi NC, England RW, Edwards AK, et al. Propranolol targets hemangioma stem
667   cells via cAMP and mitogen-activated protein kinase regulation. Stem cells translational
668   medicine. 2016;5(1):45-55.

669   30.   Murao N, Morikawa R, Seino Y, et al. Pyruvate kinase modulates the link between β-
670   cell fructose metabolism and insulin secretion. bioRxiv. 2024:2024.08. 15.608033.

671   31.   Murao N, Morikawa R, Seino Y, et al. Sildenafil amplifies calcium influx and insulin
672   secretion in pancreatic β cells. Physiological Reports. 2024;12(11):e16091.

673   32.   Murao N, Yokoi N, Takahashi H, Hayami T, Minami Y, Seino S. Increased glycolysis
674   affects β-cell function and identity in aging and diabetes. Molecular metabolism.
675   2022;55:101414.

676   33.   Oduori OS, Murao N, Shimomura K, et al. Gs/Gq signaling switch in β cells defines
677   incretin effectiveness in diabetes. The Journal of clinical investigation. 2020;130(12):6639-
678   6655.

679   34.   Ohno T, Miyasaka Y, Yoshida K, et al. A novel model mouse for type 2 diabetes
680   mellitus with early onset and persistent hyperglycemia. Experimental Animals.
681   2022;71(4):510-518.

682   35.   Palmer RK, Atwal K, Bakaj I, et al. Triphenylphosphine oxide is a potent and
683   selective inhibitor of the transient receptor potential melastatin-5 ion channel. Assay and
684   drug development technologies. 2010;8(6):703-713.

685   36.   Patt J, Alenfelder J, Pfeil EM, et al. An experimental strategy to probe Gq
686   contribution to signal transduction in living cells. Journal of Biological Chemistry.
687   2021;296

688   37.   Paul A. Adrenergic Antagonists. Introduction to Basics of Pharmacology and
689   Toxicology: Volume 2: Essentials of Systemic Pharmacology: From Principles to Practice.
690   2021:55-65.

691   38.   Pian-Smith MC, Wiedenkeller DE, Sharp GW. Paradoxical potentiation of stimulated
692   insulin release by dantrolene in rat pancreatic islets. Pancreas. 1986;1(6):501-508.

693   39.   Plant T, Henquin J-C. Phentolamine and yohimbine inhibit ATP-sensitive K+
694   channels in mouse pancreatic beta-cells. British journal of pharmacology. 1990;101(1):115.

695   40.   Porte Jr D. Beta adrenergic stimulation of insulin release in man. Diabetes.
696   1967;16(3):150-155.

697   41.   Proks P, Ashcroft FM. Phentolamine block of KATP channels is mediated by Kir6. 2.
698   Proceedings of the National Academy of Sciences. 1997;94(21):11716-11720.

699   42.   Reyes-Corral M, Sørensen NM, Thrasivoulou C, Dasgupta P, Ashmore JF, Ahmed A.
700   Differential free intracellular calcium release by class II antiarrhythmics in cancer cell
701   lines. Journal of Pharmacology and Experimental Therapeutics. 2019;369(1):152-162.

702   43.   Rorsman P, Ashcroft FM. Pancreatic β-cell electrical activity and insulin secretion: of
703   mice and men. Physiological reviews. 2018;98(1):117-214.

704   44.   Schuit F, Pipeleers D. Differences in adrenergic recognition by pancreatic A and B
705   cells. Science. 1986;232(4752):875-877.

706   45.   Seino S, Shibasaki T. PKA-dependent and PKA-independent pathways for cAMP-
707   regulated exocytosis. Physiological reviews. 2005;85(4):1303-1342.

708   46.   Sharifpour A, Sadeghi M, Zakariaei Z, Soleymani M. Seizures and irreversible
709   cardiogenic shock following propranolol poisoning: report of 2 cases and literature review.
710   Clinical Medicine Insights: Case Reports. 2022;15:11795476221126981.

711   47.   Sharma A, Rahman G, Gorelik J, Bhargava A. Voltage-gated T-type calcium channel
712   modulation by kinases and phosphatases: the old ones, the new ones, and the missing ones.
713   Cells. 2023;12(3):461.

714   48.   Shepard MJ, Bugarini A, Edwards NA, et al. Repurposing propranolol as an
715   antitumor agent in von Hippel-Lindau disease. Journal of neurosurgery. 2018;131(4):1106-
716   1114.

717   49.   Shigeto M, Ramracheya R, Tarasov AI, et al. GLP-1 stimulates insulin secretion by
718   PKC-dependent TRPM4 and TRPM5 activation. The Journal of clinical investigation.
719   2015;125(12):4714-4728.

720   50.   Sluga N, Križančić Bombek L, Kerčmar J, et al. Physiological levels of adrenaline fail
721   to stop pancreatic beta cell activity at unphysiologically high glucose levels. Frontiers in
722   Endocrinology. 2022;13:1013697.

723   51.   Straub SG, Sharp GW. Evolving insights regarding mechanisms for the inhibition of
724   insulin release by norepinephrine and heterotrimeric G proteins. American Journal of
725   Physiology-Cell Physiology. 2012;302(12):C1687-C1698.

726   52.   Taegtmeyer AB, Haschke M, Tchambaz L, et al. A study of the relationship between
727   serum bile acids and propranolol pharmacokinetics and pharmacodynamics in patients with
728   liver cirrhosis and in healthy controls. PloS one. 2014;9(6):e97885.

729   53.   Takasaki J, Saito T, Taniguchi M, et al. A novel Gαq/11-selective inhibitor. Journal of
730   Biological Chemistry. 2004;279(46):47438-47445.

731   54.   Tengholm A, Gylfe E. cAMP signalling in insulin and glucagon secretion. Diabetes,
732   obesity and metabolism. 2017;19:42-53.

733   55.   Tjandrawinata RR, Setiawati E, Yunaidi DA, Santoso ID, Setiawati A, Susanto LW.
734   Bioequivalence study of two formulations of bisoprolol fumarate film-coated tablets in
735   healthy subjects. Drug design, development and therapy. 2012:311-316.

736   56.   Wehrens XH, Lehnart SE, Reiken S, Vest JA, Wronska A, Marks AR. Ryanodine
737   receptor/calcium release channel PKA phosphorylation: a critical mediator of heart failure
738   progression. Proceedings of the National Academy of Sciences. 2006;103(3):511-518.

739   57.   Zhang Z, Zhao Z, Margolskee R, Liman E. The transduction channel TRPM5 is gated
740   by intracellular calcium in taste cells. Journal of Neuroscience. 2007;27(21):5777-5786.

741   58.   Zielmann S, Schütte G, Lenzen S, Panten U. Effects of isoprenaline and glucagon on
742   insulin secretion from pancreatic islets. Naunyn-Schmiedeberg's archives of pharmacology.
743   1985;329:299-304.

Figure 1.

Figure 1. Effect of β-blockers on glucose- and glimepiride-induced insulin secretion in

MIN6-K8 cells.

A. Dose-dependent effect of propranolol on glucose-induced insulin secretion. $n = 4$.

B. Dose-dependent effect of bisoprolol on glucose-induced insulin secretion. $n = 4$.

C. Requirement of glimepiride for propranolol-stimulated insulin secretion at 2.8 mM glucose. $n = 4$.

D. Propranolol-stimulated insulin secretion in a K_{ATP} channel-deficient β-cell line (*Kcnj11*[-/-]). $n = 4$.

All experiments were conducted using MIN6-K8 cells unless otherwise specified. Data are presented as mean ± standard deviation (SD). Glc, glucose. The reagents were added to

755   achieve the following final concentrations unless otherwise specified: glimepiride (GLM), 1

756   μM; and propranolol (Pro), 100 μM. Statistical comparisons were performed using Welch's

757   one-way ANOVA with Dunnett's post-hoc test.

Figure 2.

Figure 2. Characterization of β-blocker-stimulated cAMP elevation and insulin

secretion in MIN6-K8 cells.

A. Effect of propranolol on intracellular cAMP accumulation. n = 4.

B. Effect of bisoprolol on intracellular cAMP accumulation. n = 4.

C. Effects of AC inhibitors on β-blocker-stimulated cAMP elevation. n = 4. The data are

presented as fold change over 2.8 mM glucose alone. See Figure S1B for the original

values.

D. –E. Effects of AC inhibitors on propranolol-SIS. MDL-12330A and SQ-22536 were

used for D and E, respectively. n = 4.

F. Effect of H-89 on propranolol-SIS. n = 4.

All experiments were conducted using MIN6-K8 cells. Data are presented as mean ± SD. Glc,

glucose. The reagents were added to achieve the following final concentrations unless

otherwise specified: glimepiride (GLM) – 1 μM, propranolol (Pro) – 100 μM, bisoprolol

(Bis) – 100 μM, MDL-12330A – 10 μM, SQ-22536 – 1 mM, and H-89 – 10 μM. Statistical

comparisons were made using Welch's one-way ANOVA with Dunnett's post-hoc test for A

and B, and two-way ANOVA with Šídák's post-hoc test for C – F.

Figure 3. β-Blocker-stimulated cAMP elevation and insulin secretion in isolated mouse islets.

A. –B Dose-dependent effect of propranolol on glucose-induced insulin secretion in B6 mouse islets. One mouse was used for each experiment. n = 4.

C. Effect of propranolol on glimepiride-induced insulin secretion in B6 mouse islets. One mouse was used. n = 7.

D.   –E. Effect of propranolol on insulin secretion in islets isolated from (D) NSY.B6-*a/a*

mice (n = 5) and (E) NSY.B6-*A^y/a* mice (n = 8). Islets were pooled from two mice for

each genotype.

F.   –G. Effect of propranolol on cAMP levels in islets isolated from (F) NSY.B6-*a/a* mice

(n = 4–5) and (G) NSY.B6-Ay/a mice (n = 7) were pooled from two mice. Islets were

pooled from two mice for each genotype.

Glc, glucose. Propranolol (Pro): 100 µM. Glimepiride (GLM): 1 µM. Statistical comparisons

were performed using Welch's one-way ANOVA with Dunnett's post-hoc test for A– E, and

Welch's unpaired two-tailed t-test for F and G.

Figure 4.

Figure 4. Characterization of propranolol-stimulated Ca²⁺ response in MIN6-K8 cells.

Intracellular Ca^{2+} levels were measured using Fluo-4. All experiments were performed using MIN6-K8 cells in the presence of 2.8 mM glucose. The averaged time courses of normalized fluorescence intensity (F/F0) are indicated in A, D, and F–H. Representative traces are indicated in B and E.

A. –C. Effects of propranolol on the glimepiride-induced Ca^{2+} response. GLM: n = 8; GLM + Pro: n = 9. C, The magnitude of Ca^{2+} responses was quantified as the incremental area under the curve (iAUC).

D. –E. Propranolol-stimulated Ca^{2+} response. n = 7.

F. Effect of nifefdipine (100 μM) on the glimepiride-induced Ca^{2+} response. n = 12.

G.   Propranolol-stimulated Ca^{2+} response in extracellular Ca^{2+}-free conditions. n = 17.

EGTA: 100 μM. Carbachol (CCh, 50 μM) was added as a positive control at the end of

the recording. The black arrow indicates the $[Ca^{2+}]_i$ increase at 0 min.

H.   Effect of BHQ (15 μM) on glimepiride-induced Ca^{2+} response. n = 19.   20 mM K^+

(20K) was added as a positive control at the end of the recording.

All experiments were conducted using MIN6-K8 cells in the presence of 2.8 mM glucose.

Data are presented as mean ± SD. Glc, glucose. The reagents were added to achieve the

following final concentrations unless otherwise specified: glimepiride (GLM), 1 μM; and

propranolol (Pro), 100 μM. Statistical comparisons were made using Welch's unpaired two-

tailed t-test for C.

Figure 5.

Figure 5. Involvement of ryanodine receptor in propranolol-stimulated Ca^{2+} response in

MIN6-K8 cells.

A. –B. Effect of dantrolene on the propranolol-stimulated Ca^{2+} response in the presence of

glimepiride. The averaged time courses of the normalized fluorescence intensity of Fluo-

4 (F/F0) are indicated. Dan-: n = 9; Dan+: n = 14. GLM + Pro: n = 9. C, 20 mM K^{+}

(20K) was added as a positive control at the end of the recording. The magnitude of Ca^{2+}

responses was quantified as the incremental area under the curve (iAUC) between 0 and

600 s.

C. Effect of dantrolene on propranolol-SIS. The data are presented as fold change. See

Figure S3D for the original values.

D.   –E. The effects of glimepiride and propranolol on RYR2 phosphorylation at Ser-2808

were assessed by immunoblotting. The intensity of phosphorylated RYR2 (Ser-2808)

was normalized to total RYR2 (Ser-2808) and expressed as fold change over 2.8 mM

glucose in E. n = 3.

All experiments were performed using MIN6-K8 cells in the presence of 2.8 mM glucose.

Data are presented as mean ± SD. The reagents were added to achieve the following final

concentrations, unless otherwise specified: dantrolene (Dan), 100 μM; glimepiride (GLM), 1

μM; and propranolol (Pro), 100 μM. Statistical comparison was made using Welch's

unpaired two-tailed t-test for B and C, and Welch's one-way ANOVA with Dunnett's post-

hoc test for E.

Figure 6.

Figure 6. Involvement of TRPM4 and TRPM5 in propranolol-stimulated Ca²⁺ response in MIN6-K8 cells.

Figure 6. Involvement of TRPM4 and TRPM5 in propranolol-stimulated Ca^{2+} response

in MIN6-K8 cells.

A. Effect of low extracellular sodium concentration (30 mM) on the propranolol-stimulated

Ca^{2+} response. The averaged time course of the normalized fluorescence intensity of

Fluo-4 (F/F0) is indicated. n = 12.

B. Effect of low extracellular sodium concentration (30 mM) on propranolol-SIS. n = 4.

C. Effects of *Trpm4* and *Trpm5* knockdown on propranolol-SIS. n = 4. The data are

presented as fold change. See Figure S4C for the original values. $^{*}p < 0.05$, $^{**}p < 0.01$

versus NT (non-targeting) siRNA.

D. Effect of triphenylphosphine oxide on propranolol-stimulated Ca^{2+} response. The

averaged time course of the normalized fluorescence intensity of Fluo-4 (F/F0) is

indicated. n = 12.

E. Effect of triphenylphosphine oxide on propranolol-SIS. n = 4.

All experiments were performed using MIN6-K8 cells in the presence of 2.8 mM glucose.

Data are presented as mean ± SD. The reagents were added at the following final

concentrations unless otherwise specified: glimepiride (GLM), 1 μM; propranolol (Pro), 100

μM; and triphenylphosphine oxide (TPPO), 100 μM. Statistical comparisons were made

using Welch's one-way ANOVA with Dunnett's post-hoc test for C and two-way ANOVA

with Šídák's post-hoc test for B and E.

Figure 7.

Figure 7. Involvement of α2-adrenoceptors in propranolol-stimulated cAMP elevation

and insulin secretion.

A. –B. Effects of α2-adrenoceptor inhibitors on intracellular cAMP accumulation. n = 6.

The data **are** presented **as (A)** original value and as (B) fold change over 2.8 mM

glucose alone.

C. –D. Effects of α2-adrenoceptor inhibitors, (C) atipamezole and (D) phentolamine, on

propranolol-SIS. n = 4. were used. n = 4.

E. Effect of α2A-adrenoceptor knockdown on propranolol-SIS. n = 4. The data are

presented as fold change. See Figure S5C for the original values.

All experiments were performed using MIN6-K8 cells in the presence of 2.8 mM glucose.

Data are presented as mean ± SD. Glc, glucose. The reagents were added to achieve the

following final concentrations, unless otherwise specified: glimepiride (GLM), 1 μM;

propranolol (Pro), 100 μM; phentolamine, 100 μM; and atipamezole, 250 μM. Statistical

comparisons were made using Welch's one-way ANOVA with Dunnett's post-hoc test for B

([****]$p < 0.0001$ versus DMSO), two-way ANOVA with Šídák's post-hoc test for A and C,

two-way ANOVA with uncorrected Fisher's LSD for D, and Welch's unpaired two-tailed t-

test for E.

		FPKM (n = 3)					
		MIN6-K8			B6 islet		
Adra1a	adrenergic receptor, alpha 1a	0.00	0.00	0.00	1.42	0.00	0.00
Adra1b	adrenergic receptor, alpha 1b	0.00	0.00	0.00	0.71	0.00	0.00
Adra1d	adrenergic receptor, alpha 1d	0.00	0.00	0.00	0.00	0.00	0.00
Adra2a	adrenergic receptor, alpha 2a	455.46	442.45	462.39	354.60	373.48	345.46
Adra2b	adrenergic receptor, alpha 2b	0.00	1.00	0.00	0.00	0.00	0.00
Adra2c	adrenergic receptor, alpha 2c	0.00	0.00	0.00	0.00	0.00	0.00
Adrb1	adrenergic receptor, beta 1	0.00	0.00	0.00	2.83	1.13	0.98
Adrb2	adrenergic receptor, beta 2	0.00	0.00	0.00	0.00	1.13	0.00
Adrb3	adrenergic receptor, beta 3	0.00	0.00	0.00	0.00	0.00	0.00

880 **Table 1. Expression profiles of adrenoceptors in MIN6-K8 cells and B6 mouse islets.**

881 Source data: DDBJ Sequence Read Archive (DRA) accession number DRA006332 (Hashim
882 et al., 2018). n = 3 for each sample type.
883

Figure S1.

Figure S1. Effect of β-blockers on insulin secretion and content in MIN6-K8 cells.

A. Effect of propranolol on intracellular insulin content. n = 4.

B. Effect of various non-selective and β1-selective β-blockers on glimepiride-induced
insulin secretion in MIN6-K8 cells. Car, carteolol; Nad, nadolol; Pro, propranolol; Tim,
timolol; Bis, bisoprolol; Met, metoprolol. All β-blockers were used at a concentration of
100 μM. n = 4.

All experiments were performed using MIN6-K8 cells in the presence of 2.8 mM glucose.
Glimepiride (GLM): 1 μM. Propranolol (Pro): 100 μM. Bisoprolol (Bis): 100 μM. Data are
presented as mean ± SD. Statistical comparisons were made using Welch's one-way ANOVA
with Dunnett's post-hoc test for C, and Welch's unpaired two-tailed t-test for A, and two-way
ANOVA with Šídák's post-hoc test for B.
896

Figure S2.

Figure S3. Involvement of intracellular and extracellular Ca^{2+} in propranolol-stimulated insulin secretion in MIN6-K8 cells.

A. Effect of nifedipine (100 μM) on propranolol-SIS. n = 4.

B. Propranolol-SIS under extracellular Ca^{2+}-free conditions. n = 4. EGTA (100 μM) was added to the Ca^{2+}-free buffer.

C. Effect of BHQ (15 μM) on propranolol-SIS. n = 4.

D. Effect of dantrolene (100 μM) on propranolol-SIS. n = 4.

E. –F. Effect of xestospongin C (2 μM) on propranolol-SIS. n = 4. The data are presented in their original value in E and as fold change over glimepiride in F.

G. The knockdown efficiency of *Itpr1* was assessed using RT-qPCR. The mRNA levels
were normalized to those in siNT (non-targeting siRNA)-treated cells. n = 4.

H. –I. Effect of *Itpr1* knockdown on propranolol-SIS. n = 4. The data are presented in their
original value in H and as fold change over glimepiride in I.

All experiments were performed using MIN6-K8 cells. Glc, glucose. Glimepiride (GLM): 1
μM. Propranolol (Pro): 100 μM. Data are presented as mean ± SD. Statistical comparisons
were made using two-way ANOVA with Šídák's post-hoc test for A–C and Welch's unpaired
two-tailed t-test for D–H.
928

Figure S4.

Figure S4. Effects of *Trpm4* and *Trpm5* knockdown on propranolol-stimulated insulin secretion in MIN6-K8 cells.

A. –B. The knockdown efficiencies of *Trpm4* and *Trpm5* were assessed using RT-qPCR. The mRNA levels were normalized to those in siNT (non-targeting siRNA)-treated cells. n = 4.

C. Effect of *Trpm4* and *Trpm5* knockdown on propranolol-SIS. n = 4.

All experiments were performed using MIN6-K8 cells. Glc, glucose. Glimepiride (GLM): 1 μM. Propranolol (Pro): 100 μM. Data are presented as mean ± SD. Statistical comparisons were made using Welch's unpaired two-tailed t-test.

Figure S5.

Figure S5. Effects of *Adra2a* knockdown on propranolol-stimulated insulin secretion in MIN6-K8 cells.

A. –B. Knockdown efficiency of *Adra2a* was assessed by immunoblotting. (B) The intensity of ADRA2A was normalized to that of total protein visualized by Ponceau-S staining and expressed as fold change over 2.8 mM glucose. n = 6.

C. Effect of *Adra2a* knockdown on propranolol-SIS. n = 4.

All experiments were performed using MIN6-K8 cells. Glc, glucose. Glimepiride (GLM): 1 μM. Propranolol (Pro): 100 μM. Data are presented as the mean ± SD. Statistical comparisons were made using Welch's unpaired two-tailed t-test.