What matters to a mouse? Effects of internal and external context on male vocal response to female squeaks

Lauren R Brunner[1,2*], Laura M Hurley[1,2]

[1]Hurley Laboratory, Department of Biology, Indiana University, Bloomington, Indiana, USA

[2]Center for the Integrative Study of Animal Behavior, Indiana University, Bloomington, Indiana, USA

*Corresponding author

Email: laribrun@iu.edu (LRB)

11 Abstract

House mice adjust their signaling behavior depending on the social context of an interaction,

but which aspects of context elicit the strongest responses from these individuals is often

difficult to determine. To explore how internal and external contextual factors influence how of

house mice produce and respond to social signals, we assessed how dominant and subordinate

male mice differed in their ultrasonic vocalization (USV) production in response to playback of

broadband vocalizations (BBVs, or squeaks) when given limited access to a stimulus female. We

used a repeated measures design in which each male was exposed to two types of trials with

different odor conditions: either just female odors (Fem condition) or female odors in addition

to the odors of potential competitors, other males (Fem+Male condition). The presence of

odors from other males in this assay served as a proxy for an "audience" as the male interacted

with the stimulus female. These conditions were replicated for two distinct cohorts of

individuals: males exposed to the odor of familiar competitors in the Fem+Male condition

(Familiar odor cohort), and males exposed to the odor of unfamiliar competitors in the

Fem+Male condition (Unfamiliar odor cohort). By assessing dominance status of the focal

individual and familiarity of the "audience", we are able to explore how these factors may

affect one another as males respond to BBVs. Dominants and subordinates did not differ in

their baseline vocal production (vocalizations produced prior to squeak playback) or response

to squeaks. However, all groups, regardless of dominance status or odor condition, reduced

their vocal production in response to BBV playback. The presence of unfamiliar male odor

prompted mice to decrease their baseline level of calling and decrease the complexity of their

vocal repertoire compared to trials that only included female odor, and this effect also did not

differ across dominance status. Importantly, the presence of male odor did not affect vocal

behavior when the male odor was familiar to the focal individual. These findings suggest that

mice alter their vocal behavior during courtship interactions in response to cues that indicate

the presence of potential competitors, and this response is modulated by the familiarity of

these competitor cues.

Introduction

The ability to process and respond appropriately to conspecific signals is crucial for animals as

they navigate social interactions [1–3]. How an individual processes and responds to signals

may heavily depend on the context of the situation [4–6], and the role of context in vocal

communication in particular is highly documented across taxa [7–11]. Research on how

contextual influences such as the presence of competitors and potential mates, seasonality,

changes in steroid hormones, and resource abundance affect song production and song

composition in birds has revealed that many aspects of communication depend on context, and

prompted advances and questions on how context can influence the signaling in other species

[12,12–16]. These include house mice, studied not only as a model of rodent communication

signals, but also for their potential applicability to human communication disorders [17,18].

House mice (*Mus musculus*) produce a variety of vocalizations that can vary in usage,

structure, and meaning across different contexts [1,10,11,19,20]. Vocal communication in mice

is prominent during courtship interactions, in which calls are used to communicate receptivity

and sexual motivation [1,10,21–24]. One type of call, ultrasonic vocalizations (USVs), are high-

frequency calls (30-110 kHz) produced by both males and females during social interactions

[1,20,24–26]. These calls are produced in large numbers by males during opposite-sex

interactions, and are often referred to as "courtship calls" because of their prosocial function to

facilitate mating [1,24,27]. Male USV production thus may indicate male courtship effort and

motivation [22]. Mouse vocal repertoires, or the composition of different types of syllables

individuals use, are complex and can vary depending on factors such as age, genetics, and

behavioral situation [10,28–30]. One type of USV in particular, 50 kHz harmonic USVs, tend to

be longer in duration and more complex structurally than other classes of USVs. These calls are

closely associated temporally with incidents of male mounting behavior [10].

Another type of call, broadband vocalizations (BBVs), or squeaks, are lower-frequency

calls with a broader frequency range than USVs, featuring many harmonics in. Both male and

female mice produce BBVs across varying contexts, including during distress, handling, and

conspecific interactions [26,31–33]. When produced by females during the early stages of

courtship interactions, these calls are closely correlated with females performing physical

rejection behaviors like kicking, lunging, and attempted escape [11,34]. Thus, during courtship,

female-produced BBVs may be negatively valenced, and function to express a lack of female

receptivity to sexual advances by males. In accordance with the connection of squeaks to

physical rejection, male mice reduce their USV output in response to BBV playback during

limited interactions with female mice [35]. However, little is known about what sorts of

contextual factors may influence this response to BBVs, and which levels of context may have

the most substantial impacts on mouse behavior as they interact.

The goal of this study was to explore the effects of both internal and external contextual

factors on male vocal production and vocal repertoire in response to female squeaks. By

examining both influences inside and outside of the focal individual, we can gain a better

understanding of how these forces interact to affect behavior as a whole. As an internal

contextual factor, we evaluated dominance status in all of our individuals. Some calls are more

associated with the production of behaviors classified as "more dominant", such as aggression

and courting [20]. Dominant and subordinate mice tend to differ in their vocal responses to

female cues, though the direction of this difference varies across studies. For example, one

study found that dominant male mice call more quickly and more often compared to

subordinate males in response to female stimuli [36], while in contrast, another found that

subordinates called more than dominants when interacting with a female [21]. Thus, the effects

of dominance status on vocal production in male mice appear to be highly-context dependent

and worthy of further investigation.

In addition to the playback of BBVs, the external contextual factor of interest to us was

the presence of non-vocal cues that may alter male response to BBVs: specifically, odor from

other male mice. Audience effects, in which individuals adjust their behavior in the presence of

additional individuals, are well-documented examples of context-dependent communication

[37–39]. Studies in songbirds and domesticated fowl have revealed that the presence of

audiences during courtship signaling can influence not only the acoustic properties of song and

song repertoire, but also the motivation to interact with their partner [40–43]. Previous studies

have established that certain strains of mice are susceptible to audience effects when

responding to female cues [37,44]. For example, Seagraves et al. (2016) found that increases in

vocal production in response to live and anesthetized male audience members may be due to

signalers shifting their vocalizations to be directed to males, potentially in a competitive display.

Thus, the presence of an audience may cause mice to adjust their vocal output and the function

of their calling may change, similar to the strategic changes influenced by audiences across

other contexts [37,43,45–47]. On the other hand, competitive male odor cues can decrease

male vocal behavior and courtship effort when interacting with a female [21,48]. Thus,

audience effects can vary depending on the context of an interaction.

Importantly, defining what constitutes an "audience" and an "audience effect" has been

debated across the literature since the term's conception [38,39]. For example, some argue

that true audience member(s) do not engage in the exchange of signals with those in the

communication interaction [39,49]. Some theorists and studies support that audience effects

may not be limited to the physical presence of an individual acting as an audience, and that

cues that indicate the proximity of a "potential" audience may be sufficient to promote these

changes in signaling behavior [50]. For example, male newts (*Lissotriton boscai*) decrease their

courtship effort when the area in which they are interacting with a female contains chemical

cues from other males, potentially indicating a "chemically-mediated" audience effect [48].

Similarly, subordinate male mice that were evaluated for vocal production during an interaction

with a female called less if they were tested in the same cage in which a dominant male and

female had previously interacted, compared to if they were tested in a cage with no social

odors [21]. Thus, the researchers concluded that odor cues left behind by the urine of dominant

male were sufficient to affect subordinate male vocal behavior during an interaction with a

female, which aligns with the framework of audience effects. Because odor cues can imply

audience presence without the possibility of active signal exchange, investigating audience

effects elicited by these cues can ensure that the audience cannot take part in the signaling

interaction, thus the role of the audience as a "non-targeted receiver" is maintained [39,50].

In order to assess the effects of a simulated audience, in the form of conspecific male

odor cues, and dominance status on male behavioral response to BBVs, we employed a recently

established paradigm [35] that allows for the decoupling of BBVs and their frequently

associated physical rejection behaviors. We predicted that the presence of male odor during

these interactions would prompt males to decrease their vocal output in response to BBV

playback more so than if no male "audience" was present, and that this effect would be

dependent on dominance status. It has been reported that dominant mice are more likely to

adjust their behavior in response to social challenges than subordinate mice [51–53], so we

predicted that dominant males would alter their behavior in response to a simulated social

competitor more than subordinates by showing a larger decrease in USV production during BBV

playback if male odor was present.

We investigated the role of familiarity of the perceived "audience" in this study by

assessing behavior of male mice in two distinct cohorts: one exposed to unfamiliar male

conspecific odor, and the other exposed to odors from a male cage-mate with which they had

an established dominance relationship. Familiarity was of particular interest to us because

identity and status of audience members can alter behavior during signaling [54]. We predicted

that more robust differences between dominants and subordinates would be seen in the

familiar odor cohort because of the established dominance relationship between the signaler

and the audience member.

Including cues that may indicate an audience, allows us to greater understand how

males are responding to vocal courtship rejection signals (BBVs) in a more naturalistic

environment, where they are likely to be confronted with the odors of competing males

[55,56]. By assessing both dominance status and familiarity of the audience to the focal

individual, we can further parse apart what role identity plays in this interaction. Surprisingly,

we found that dominance status did not affect vocal behavior, while the presence of additional

male odor dampened vocal production, but only when the odor came from a male unfamiliar to

the focal individual.

Methods

Animals

This study was carried out in strict accordance with the recommendations in the Guide for the

Care and Use of Laboratory Animals of the National Institutes of Health. The protocol was

approved by the Bloomington Institutional Animal Care and Use Committee at Indiana

University (Protocol number: 21–020). Focal subjects consisted of seven-week-old male CBA/J

mice (N = 23) (The Jackson Laboratory, Bar Harbor, ME). Mice were housed in a 14:10 light:dark

cycle with food and water *ad libitum*. All focal and stimulus mice were housed in same-sex pairs

from at least one week prior to behavioral assays until the end of the experiments in order to

allow for the establishment and maintenance of a social dominance relationship within the

cage. Dominance status in male mice was evaluated using multiple measurements of a

standardized tube test in which cage-mates were simultaneously released at opposite ends of a

clear plastic tube and allowed to interact until only one mouse remained in the tube [35,57].

The mouse that retreated from the tube was classified as subordinate, while the mouse that

remained in the tube longer was classified as dominant. The tube used was narrow enough to

prevent the mice from passing over one another [35,50]. All housing pairs underwent 3-4 tube

tests with no more than one test per day, and with tests occurring no more than 4 days prior to

the trial dates. Almost all mice were consistent in their dominance status across all of their

tests, aside from two pairings that switched dominance status after the first test, but were

consistent for the final three tests. At least two days prior to experimental trials, all mice were

given opposite-sex experience with novel female CBA/J mice (The Jackson Laboratory, Bar

Harbor, ME) through multiple 10-minute interactions across the span of 2-3 days [11]. These

interactions occurred in clean cages, in which each focal male was allowed to interact alone

with a stimulus female. All mice received approximately the same amount of social experience

prior to trials (50-80 minutes total with an average of 67.14 minutes, SD = 14.85). A student's t-

test confirmed that individuals across the two cohorts received statistically similar amounts of

social experience ($t(26)$ = -0.142, p = 0.888). All stimulus animals for experimental trials were

sexually experienced adult female CBA/J house mice novel to the focal mouse.

Behavioral assay and experimental conditions

A split-cage playback paradigm previously established in our lab [35] was employed to evaluate

the effects of vocal playback on male mouse vocal behavior. In this paradigm, the focal mouse

and stimulus female are placed on opposite sides of a 12 inch x 6 inch x 6 inch cage with a

plexiglass barrier in the middle with a small hole at the bottom of the barrier that allows for

limited nose-to-nose social investigation. The presence of a live female, opposed to only female

odor or an anesthetized female, is necessary to promote consistent robust vocalization from

the focal male during this assay [35]. Approximately 30g of dirty bedding containing urine,

feces, and body odor [58,59] from the stimulus female and her cage mate was placed on the

focal male's side of the cage to promote robust vocalizations [35,36], while clean bedding was

placed on the female's side of the cage. Dirty bedding from female cages was obtained after 24-

48 hours after the bedding was placed with the animals. A microphone (CM16/CMPA, Avisoft

Bioacoustics, Glienicke/Nordbahn, Germany) and Canon VIXIA video camera were placed above

the cage on the focal male's side to allow for recording of ultrasonic vocalizations (Fig 1A) and

BBV playback (Fig 1B), and non-vocal behavior, respectively, during the interaction. The entire

cage and recording setup were located in a sound-proof chamber (IAC Acoustics, Naperville, IL)

in a room separate from the area in which the colony was housed. All trials were performed

between 9:00 AM – 4:00 PM during the light portion of the animal's light cycle. Mice were

habituated to the sound chambers 1 day prior to their experimental trials for 60 minutes. On

the day of the trials, the focal male mouse was placed in the experimental cage for 60 seconds,

or until they started producing USVs, and subsequently, the focal female was then placed on

the opposite side of the cage. Mice were in this cage for 15-minute playback trials that

consisted of 5 minutes of silence (a baseline period), followed by 5 minutes of exemplar BBV

playback through an ultrasonic speaker powered by Avisoft next to the cage, followed by

another five minutes of silence (a recovery period). Hence forth, these time periods will be

referred to as test periods. The segment of BBV playback was created using five repeated 1-

minute segments of BBVs collected from a previous direct interaction between a male and a

female during a period of frequent physical rejection and BBV vocalization [35]. Each call during

this period was replaced with a single exemplar BBV with a duration, peak frequency, and

percentage of non-linear segments within one standard deviation of the mean for all BBVs in

the interaction in order to eliminate possible differences in behavior over the course of the

playback due to variation in BBV structure [35]. The BBV segment was played at a loudness

> **Fig 1: Representative spectrograms of (A) Ultrasonic vocalizations (USVs) and (B) Broadband vocalizations (BBVs).** The circled vocalization in (A) is an example of a 50 kHz

calibrated to be approximately matched to the intensity that male mice would experience from

BBVs during natural interactions with females, with a maximum of 104.9 dB [35].

Previous use of this paradigm in our lab has shown that males decrease their USV

production during BBV playback relative to the baseline and increase back to baseline levels

during the recovery. This response to BBV playback when males are physically separated from

the stimulus female indicates that playback of BBVs can alter male courtship vocal behavior

even when the correlated physical rejection behaviors are absent [35]. For the current study, a

modified version of this paradigm was created to allow for two distinct experimental conditions

that altered the available olfactory cues: a female-scent-only condition in which only bedding

from the stimulus female's home cage was placed on the focal animal's side of the cage (Fem

condition) and a female scent + male scent condition in which the focal male was exposed to

the 30g of bedding from the stimulus female as well as 30g of bedding from a cage of male mice

(Fem+Male condition; Fig 2).

Both assay conditions were applied to the two separate experiments with separate

> **Fig 2: Experimental assay and design.** (A) Split-cage paradigm used to evaluate the effects of only BBV playback on male vocal behavior, modified to create the Fem+Male condition for this study. Male and female symbols present on the male mouse's side of the cage represent the presence of dirty bedding from male and female mice. (B) Experimental groups broken up by familiarity cohort, dominance status, and odor condition. A question mark icon is used to illustrate unfamiliarity with the odor stimulus in the trial, and an exclamation point icon is used to illustrate familiarity with the odor stimulus in the trial.

cohorts of animals described in the following sections. These experiments differed based on the

origin of the male bedding used in the Fem+Male condition, described below, while the Fem

condition trials in both experiments were identical in design. All individuals in both

experiments/cohorts underwent both a Fem condition trial and a Fem+Male condition trial to

account for individual differences in vocal behavior. Different trials for the same mice occurred

approximately 24 hours apart and the order of the trials was counterbalanced across all

individuals.

Experimental design

Experiment 1 - Unfamiliar male odor cohort: In the Fem+Male condition, individuals in

the unfamiliar male odor cohort (n=12) were exposed to equal amounts of dirty bedding from

the stimulus female's cage and dirty bedding that came from a cage of males unfamiliar to

them. These "odor donations" were from the other cages of males used in this experiment. No

males from separate cages ever interacted with one another, thus all of the "odor donations"

came from males novel to the focal individuals. Dirty bedding containing urine, feces, and body

odor from unfamiliar male odor donors was collected after the unfamiliar donor male was

allowed to explore the cage for 2 hrs. The Fem+Male odor condition in this experiment was

designed this way to simulate the presence of an unfamiliar male competitor during the limited

interaction with the female.

Experiment 2 - Familiar male odor cohort: In a subsequent experiment, a new cohort

(n=16) was established as the familiar male odor cohort. For these individuals, in the Fem+Male

condition they were exposed to equal amounts of dirty bedding from the stimulus female's

cage and dirty bedding from their own home cage, in order to simulate the presence of their

familiar cage mate: the other half of their dominance pairing. Dirty bedding containing urine,

feces, and body odor from the familiar male odor donors was collected from a cage that held

the focal male and their cage mate for 2 hrs.

Audio analysis

USVs from all of the interactions were counted individually by trained individuals using Avisoft

SasLab Pro software (Avisoft Bioacoustics, Glienicke/Nordbahn, Germany). A total of 74,859

USVs were counted manually during the course of this study. Durations of the USVs were

automatically calculated by Avisoft. In order to assess specific components of the individuals'

vocal repertoires, all USVs were manually categorized as either non-harmonic or 50 kHz

harmonic USVs (Fig 1). 50 kHz harmonics were defined as syllables with at least one harmonic

band at 40-50 kHz, as per Hanson and Hurley's (2012) criteria [10], while all other USVs were

categorized as non-harmonic if they did not possess this specific frequency band (Fig 1). 50 kHz

harmonic USVs were specifically denoted because they are highly correlated to male mouse

mounting behavior and could potentially be used to promote receptivity in females during

courtship interactions [10,35].

Non-vocal behavior analysis

Non-vocal behavior was analyzed using Behavioral Observation Research Interactive Software

(BORIS) [60]. The odor condition and cohort of the focal individual was blinded to the analyst

for each trial. Because male and female investigative behavior may be related in this assay [35],

the duration of time both the focal male and stimulus female spent at the hole in the plexiglass

divider was scored for all individuals. Investigation of the window was defined as the mouse

placing their nose at the entrance of (< 1 cm away from the entrance) or through the opening in

the barrier. Investigation of holes in a barrier that separates a focal and stimulus mouse has

also been historically scored to understand overall investigative behavior in assays like these

[7,35].

Statistical analysis

Statistical analyses were performed in IBM SPSS Statistics 28. A Pearson correlation was used to

evaluate the relationship between the baseline USVs produced in the Fem condition and the

baseline USVs produced in the Fem+Male condition across all individuals to assess consistency

in baseline vocal production. Four Pearson correlations were used to evaluate the relationship

between the investigative behavior of the focal male and the stimulus female in each cohort,

for each odor condition. Due to the difference in the Fem+Male conditions across the two

cohorts, separate general linear models were used for each cohort for analysis of vocal and

non-vocal behavior. For all models that used repeated measures, the mouse's individual ID

served as the repeated measure.

To assess the effects of BBV playback across the 5-minute test periods (baseline,

playback, and recovery) and odor treatment on number of non-harmonic calls, a three-way

repeated-measures ANOVA was used for the unfamiliar odor cohort, and an identical three-way

repeated-measures ANOVA was used for the familiar odor cohort (within-subjects factors: test

period and odor treatment; between-subjects factor: dominance status. Analyses on the effects

of odor condition and dominance status on baseline number of USVs in each interaction and

the duration of baseline USVs produced were conducted using identical two-way mixed

ANOVAs for the unfamiliar odor cohort and familiar odor cohort (within-subjects factor: odor

treatment; between-subjects factor: dominance status). Interactions among all three factors

were included in these models.

In order to investigate how the composition of vocal repertoires responded to our

manipulations, we characterized the proportion of harmonic USVs out of all USVs in each test

period of the interaction. To assess the effects of BBV playback across test periods and odor

treatment on proportion of harmonic calls produced, proportions of harmonic calls across test

periods were calculated as: number of harmonic calls in test period/number all calls in test

period. These proportions were then compared using identical binary logistic regressions for

each cohort in which each vocalization was coded as either harmonic or non-harmonic [61–63]

(within-subjects factors: test period and odor treatment; between-subjects factor: dominance

status). Binary logistic regressions were also used to evaluate the effects of odor condition and

dominance status on the baseline proportion of harmonic USVs for the unfamiliar odor cohort

and familiar odor cohort (within-subjects factor: odor treatment; between-subjects factor:

dominance status).

To assess the effects of BBV playback across test period and odor treatment on non-

vocal behavior, two identical three-way repeated-measures ANOVAs were used, one for each

cohort (within-subjects factors: test period and odor treatment; between-subjects factor:

dominance status). There was an error with one individual's video file for the last ten minutes

of the trial, so this file was excluded from analysis for these measures. Interactions among all

three factors were included in these models.

The following datasets violated the ANOVA assumption of normality and thus were

normalized using square root transformations in order to run analyses of variance: USVs across

test periods in the unfamiliar odor cohort, and female investigative behavior in the unfamiliar

odor cohort. Small violations of the homogeneity of variance requirement were noted, but as

these models were the best fit aside from this fact, and because ANOVAs are robust to small

violations of this assumption when sample sizes are equal, we proceeded with ANOVAs [64].

Benjamini-Hochberg corrections (Q value = 0.05) were conducted in order to account for

multiple comparisons [65], and Bonferroni tests were performed as posthoc analyses.

The possibility of outliers in the datasets was investigated using analyses of the number

of standard deviations that each value deviated from the mean of the dataset, and double-

checked using Tukey's method [66,67]. No outliers were revealed in any of the datasets.

Results

Individual baseline vocal production in both odor conditions is positively correlated

All individuals underwent both a Fem condition trial, in which focal individuals were only exposed to the odor of the female stimulus animal, and a Fem+Male condition trial, in which focal individuals were exposed to both odor from the female stimulus and odor from an additional male conspecific (Fig 2B). A Pearson Correlation revealed that the baseline vocal production across both trials was strongly positively correlated across individuals in both the familiar odor cohort and unfamiliar odor cohort ($r = 0.735$, $p < 0.001$).

Dominance status does not alter behavior across cohorts or odor conditions

Dominance relationships for each cage consisted of a dominant individual and a subordinate individual, and the placement of individuals within these dominance relationships remained stable throughout multiple rounds of tube tests. Out of 14 dominance pairings, only 2 pairings changed status at all throughout the tube tests. Both of these switches occurred during the second test and the pairings were subsequently stable throughout the rest of the three tests. When incorporating dominance status in our statistical models as a between-subjects factor, there were no significant effects of dominance status in any behavioral measures (Tables 1-3).

347 **Table 1. Output of models assessing the effects of BBV playback, Fem or Fem+Male odor**

348 **condition, and dominance status on total USV production and harmonic USV production in**

349 **both cohorts.**

	df	Number of USVs in Unfamiliar odor cohort [a]		Proportion of 50 kHz Harmonics in Unfamiliar odor cohort [b]	
Effect	*df*	*F*	*p*	*F*	*p*
Test Period	2	50.538	**<.001**	4.925	**0.011**
Odor Condition	1	12.026	**0.006**	8.429	**0.005**
Dominance Status	1	0.019	0.892	0.702	0.406
Test Period x Odor Condition	2	0.223	0.802	2.236	0.116
Period x Dominance Status	2	0.614	0.551	0.131	0.877
Odor Condition x Dominance Status	1	0.742	0.409	0.022	0.882
Odor Condition x Test Period x Dominance Status	2	1.203	0.321	1.093	0.342

350 [a] Nonharmonic USVs were assessed with three-way mixed ANOVAs.

351 [b] Proportion of 50 kHz harmonic USVs were assessed using binary logistic regressions.

353 **Table 2. Output of two-way mixed ANOVAs assessing the effects of Fem or Fem+Male odor**

354 **condition and dominance status on baseline USV production, baseline harmonic USV**

355 **production, and duration of baseline USVs in both cohorts.**

	df	Baseline number of USVs in Unfamiliar odor cohort		Baseline 50 kHz Harmonics in Unfamiliar odor cohort		Duration of base USVs in Unfami odor cohort	
Effect	*df*	*F*	*p*	*F*	*p*	*F*	*p*
Odor Condition	1	10.005	**0.01**	10.752	**0.004**	5.285	0.044
Dominance Status	1	0.04	0.846	0.036	0.852	0.051	0.826
Odor Condition x Dominance Status	1	0.919	0.36	2.101	0.163	0.576	0.465

356 * p value is not statistically significant after correcting for multiple comparisons.

358 **Table 3. Output of three-way mixed ANOVAs assessing the effects of BBV playback, Fem or**

359 **Fem+Male odor condition, and dominance status on male and female investigation of the**

360 **divider window in both cohorts.**

		Male investigation in Unfamiliar odor cohort		Female investigation in Unfamiliar odor cohort			M
Effect	*df*	*F*	*p*	*F*	*p*	*df*	*F*
Test Period	2	3.965	0.037*	3.841	0.041*	2	4.
Odor Condition	1	0.958	0.353	5.437	0.045*	1	0.
Dominance Status	1	3.315	0.102	0.549	0.478	1	0.
Test Period x Odor Condition	2	0.421	0.663	0.254	0.779	2	0.
Test Period x Dominance Status	2	0.732	0.495	4.109	0.034*	2	2.
Odor Condition x Dominance Status	1	1.962	0.195	0.059	0.814	1	0.
Odor Condition x Test Period x Dominance Status	2	0.585	0.567	0.493	0.619	2	0.

361 * p value is not statistically significant after correcting for multiple comparisons.

363 Additional male odor during limited social interactions decreases

364 baseline vocal output depending on the familiarity of male odor

365 The presence of unfamiliar male odor during the interaction (Fem+Male condition) caused

366 significantly lower baseline levels of USVs, USVs during the first 5 minutes of the interaction,

367 compared to when only female odor was present (Fem condition) (Two-way mixed ANOVA: $F_{1,10}$

Figure 3: **Properties of vocal repertoires.** (A) Baseline USVs produced in the Fem condition vs the Fem+Male condition. (B) Duration of baseline USVs in seconds. (C) Baseline proportion of harmonic USVs (# baseline harmonics/# baseline USVs). Proportion of baseline harmonic USVs was calculated as baseline harmonic USVs /(baseline harmonic USVs + baseline nonharmonic USVs). Blues represent the unfamiliar odor cohort, reds represent the familiar odor cohort. Bars represent the means of the experimental groups, and means from dominants and subordinates are pooled due to the lack of effect of social status. Error bars are calculated from standard errors. Individual points of data noted with dominants as X's
and subordinates as circles.

= 10.005, p = 0.01). The presence of familiar male odor however produced no significant

difference in baseline USV levels compared to when only female odor was present (Two-way

mixed ANOVA: $F_{1,14}$ = 0.084, p = 0.776) (Fig 3A) (Table 2).

Because USVs can vary widely in duration [1], we measured the duration of all baseline

USVs (both harmonic and nonharmonic). A trend of males producing longer baseline USVs in

the Fem condition compared to the Fem+Male condition in unfamiliar male cohort was noted

(Two-way mixed ANOVA: $F_{1,10}$ = 5.285, p = 0.044), though this result was not significant after

correcting for multiple comparisons (Table 2). No effects of odor condition on duration of

baseline USVs were observed in the familiar male odor cohort (Two-way mixed ANOVA: $F_{1,14}$ =

0.005, p = 0.945) (Fig 3B).

Additional male odor during limited social interactions decreases

baseline usage of 50 kHz harmonic USVs depending on the familiarity

of male odor

50 kHz harmonic USVs are calls highly correlated to mounting behavior in male mice [11]. In this

study, harmonic USVs made up a small proportion of all calls; on average, harmonic USVs made

up 7.31% (SD = 0.0701) of all USVs in the interaction. In order to assess whether the frequency

of harmonic USV usage differed across conditions, for each individual we evaluated the

proportions of harmonic USVs produced in each test period out of the total number of all USVs

in that test period. The presence of unfamiliar male odor during the interaction (Fem+Male

condition) caused mice to use a significantly lower proportion of 50 kHz harmonic USVs during

the baseline period compared to when only female odor was present (Fem condition) (Binary

logistic regression: $F_{1,10}$ = 10.752, p = 0.004), though this effect was not present in the familiar

male odor cohort (Binary logistic regression: $F_{1,14}$ = 0.035 p = 0.853) (Fig 3C) (Table 2).

Male mice decrease USV production in response to BBV playback across all conditions

In both the unfamiliar odor cohort and the familiar odor cohort, male mice reduced their calls

during playback of BBVs, and then increased calling rates after playback ceased in both Fem and

Fem+Male trials (Three-way mixed ANOVAs: Significant main effect of test period $F_{2,20}$ = 50.538,

p < 0.001 and $F_{2,28}$ = 34.635, p < 0.001 for the unfamiliar and familiar cohorts, respectively). This

subsequent increase in calling after the playback period did not fully recover to levels similar to

baseline, however, but rather increased to a level between that observed in the baseline and

the recovery. In the unfamiliar odor cohort, individuals produced more USVs in the Fem

condition compared to the Fem+Male condition (Significant main effect of odor condition: $F_{1,10}$

= 12.026, p = 0.006) (Fig 4A), though this effect was not present in the familiar odor cohort ($F_{1,14}$

= 0.454, p = 0.511) (Fig 4B). No interaction effects between odor and test period were present

in either the unfamiliar odor cohort or the familiar odor cohort ($F_{2,20}$ = 0.223, p = 0.802, and $F_{2,30}$

Figure 4: Number of USVs produced across test periods (A) in the unfamiliar odor cohort, and (B) in the familiar odor cohort. Average USV production of dominants is represented by the solid lines, and average USV production of subordinates is represented by dashed lines. Lighter lines denote average USV production of individuals in the Fem condition, while darker lines denote average USV production of individuals in the Fem+Male condition. Error bars are calculated from standard errors.

406   = 0.734, p = 0.489) (Table 1).

407   Because there was no interaction effect between test periods and any other factor in

408   these models, our data showed that differences across odor conditions and familiarity cohorts

409   can be attributed to differences in baseline levels of USV production, rather than differences in

410   vocal response to BBV playback.

Male mice increase usage of 50 kHz harmonics after BBV playback

ends when exposed to unfamiliar male odor

In the unfamiliar male odor cohort, the proportion of 50 kHz harmonic calls used relative to the

total number of calls produced did not change in response to BBV playback; however, after

playback ended, males produced a greater proportion of harmonic calls in the recovery period

(Binary logistic regression: Significant main effect of test period $F_{2,20}$ = 4.925, p = 0.011).

However, posthoc pairwise comparisons revealed this increase in harmonic usage only occurred

in trials with unfamiliar male odor added (t_{55} = 2.344, p = 0.023) (Fig 5A). Overall, in the

unfamiliar male odor cohort, males produced a greater proportion of harmonics in Fem trials

compared to Fem+Male trials (Binary logistic regression: Significant main effect of odor

condition $F_{1,10}$ = 8.429, p = 0.005). In the familiar male odor cohort, no significant effects of any

Figure 5: Proportion of harmonic USVs across test period out of total USVs (A) in the unfamiliar odor cohort and (B) in the familiar odor cohort. Average harmonic USV production of dominants is represented by the solid lines, and average harmonic USV production of subordinates is represented by dashed lines. Lighter lines denote average harmonic USV production of individuals in the Fem condition, while darker lines denote average harmonic USV production of individuals in the Fem+Male condition. Proportion of harmonic USVs for each test period was calculated as harmonic USVs in test period/(harmonic USVs + nonharmonic USVs in test period). The significance bar in 5A) showing increased usage of proportion of harmonic USVs in the recovery period compared to the baseline and playback period is dark blue, representing that it only applies to the Fem+Male conditions, not the Fem conditions. Error bars are calculated from standard errors.

423   factor were observed (Table 1) (Fig 5B).

Male and female investigative behavior does not significantly differ

across conditions, cohorts, nor test periods

In the unfamiliar odor cohort, male investigation of the divider window appeared to slightly

decrease during BBV playback, and investigative behavior did not recover after BBV playback,

with no effects of dominance status or odor condition (Three-way mixed ANOVA: $F_{2,18}$ = 3.965,

p = 0.037) (Fig 6A). In contrast, in the familiar odor cohort, male investigation of the divider

window appeared to slightly decrease during BBV playback and subsequently recover after

playback, with no effects of dominance status or odor condition (Three-way mixed ANOVA: $F_{2,28}$

= 4.391, p = 0.022) (Fig 6C). However, neither of these results was significant after correcting for

Figure 6: Duration of time the focal male (A and C) or the stimulus female (B and D) spent investigating the divider window across test periods in the unfamiliar odor cohort (A and B) and in the familiar odor cohort (C and D). Average duration of investigation in trials with dominant males is represented by the solid lines, and average duration of investigation in trials with subordinates is represented by dashed lines. Lighter lines denote average investigation during trials in the Fem condition, while darker lines denote average investigation in the Fem+Male condition. Error bars are calculated from standard errors.

multiple comparisons.

Stimulus females paired with males in the unfamiliar odor cohort tended to investigate

the divider window more during the Fem+Male condition trials ($F_{1,9}$ = 5.437, p = 0.045), and this

investigation actually increased during BBV playback, and decreased back to baseline levels

after playback ended ($F_{2,18}$ = 3.841, p = 0.041) (Fig 6B). This effect of playback was dependent

on dominance status in that female investigative behavior was only affected by playback when

440 females were paired with subordinate males, and there was no effect when they were paired

441 with dominant males ($F_{2,18}$ = 4.109, p = 0.034). When stimulus females were paired with males

442 in the familiar odor cohort, they decreased their investigative behavior during BBV playback,

443 and this behavior increased back to baseline levels when playback ended ($F_{2,28}$ = 3.949, p =

444 0.031) (Fig 6D), with no effect of odor condition or dominance status. However, again, none of

445 these results were significant after correcting for multiple comparisons.

447 Additional male odor alters the relationship between male and

448 female investigative behavior

449 In the Fem condition trials, the total duration of male investigative behavior did not correlate to

450 the total duration of female investigation behaviors in either the unfamiliar odor cohort or the

451 familiar odor cohort (Pearson correlation: r = -0.373, p = 0.259; r = 0.235, p = 0.381

452 respectively). In the Fem+Male condition trials, the total duration of male investigative

453 behaviors negatively correlated to the total duration of female investigative behaviors for those

454 in the unfamiliar odor cohort (Pearson correlation r = -0.625, p = 0.04), though this effect was

455 not significant after correcting for multiple comparisons. In contrast, in the Fem+Male odor

456 condition trials conducted with the familiar odor cohort, the total duration of male

457 investigation behaviors positively correlated to the total duration of female investigative

458 behaviors (Pearson correlation r = 0.839 p < 0.001), and this effect remained significant after

459 correcting for multiple comparisons.

Discussion

Behavioral decisions during courtship are highly influenced by the interplay between

environmental cues, internal factors, and signals from communicating conspecifics [4]. Thus, by

evaluating how individuals respond to signals in various contexts, we can begin to decipher how

much weight a given factor may have on such decisions. In this study, when interactions

included odor cues from unfamiliar, but not familiar, male conspecifics, male mice appeared to

exhibit reduced courtship motivation compared to interactions when only female odors were

present. This decrease in courtship motivation was suggested by shifts in multiple aspects of

male courtship vocalizations: decreases in call production, call durations, and variety of calling

repertoires. Male mice thus may be evaluating their environment for potential competitors,

discerning the identity of these competitors, and scaling their courtship activities in response to

these cues. The interplay between signals not only from courtship partners but also from

competitors as well as social recognition of those signals is instrumental in motivating

behavioral decisions during mating interactions.

Unfamiliar male odor during courtship decreases male courtship

motivation

In many taxa, vocal signaling can often reflect motivational state [68–73]. USV production is a

well-established proxy for courtship motivation in male mice, as these calls function to

encourage female sexual receptivity and mating [18,21,22,24,74,75]. Furthermore, out of the

possible measurements for courtship motivation in mice, these signals are particularly

important as they can convey information about male fitness, identity, and reproductive state,

which can in turn influence female mate choice [22,26,76]. Changes in courtship motivation can

be spurred by exposure to social cues [69,70]. It is well-established that auditory

communication, like birdsong, exhibits signal flexibility in response to social context [77]. For

example, male cowbirds will modulate the attractiveness of their songs in response to the

presence of competitors, though how they modulate their songs may depend on factors like

whether or not the competitors can easily access the focal individual and if the individual can

gauge the motivational state of the competitor [78,79].

We found that whether males responded to the male "audience" odor was dependent

on whether or not the focal male had social contact with the odor's source. When presented

with unfamiliar male odor during the interaction, males produced fewer calls during the

baseline period (Fig 3A), with a shorter average length of calls than when male odor was not

present (Fig 3B). Similarly, they also shifted the composition of the different types of syllables

they used, their vocal repertoire [29], to produce a lower proportion of 50 kHz harmonic calls

during the baseline period when unfamiliar male odor is present (Fig 3C). These responses may

indicate a decrease in courtship motivation and effort. Not only is number of calls produced

indicative of motivational state, but types of calls produced can also reveal information about

motivation. Across taxa, increasing vocal production through the expression of complex

syllables and longer calls, like 50 kHz harmonics [10,11], is associated with greater energetic

costs. Longer durations of vocalizations have also been associated with increased courtship

display effort in singing mice (*Scotinomys teguina),* and female rodents tend to prefer males

that produce longer "higher-effort" songs [75,80,81]. Thus, calling less and producing fewer

complex harmonic calls and shorter calls thus could indicate decreased reproductive

motivation.

Olfactory cues are often crucial for defining the context of an interaction, particularly

opposite-sex interactions. Therefore, individuals may adjust their behavior in response to these

cues, potentially to avoid expending unnecessary time and energy engaging with individuals

that are not sexually receptive [82–88]. Odor cues can be used to simulate different audiences

and investigate how different identity characteristics alter behavior during sexual interactions

[21,48,50]. However, a previous study from Seagraves et al. (2016) found that, when presented

alone, male mouse urine and male mouse body odor (in the form of saliva, tears, and fur), did

not affect how males vocally responded to female odor, leading the authors to conclude that

male mouse odors were not sufficient to induce an audience effect [44]. Here, we found that

unfamiliar, but not familiar, male odors from dirty bedding were able to prompt a significant

decrease in vocal behavior when the males were interacting with a live female stimulus mouse.

Our results mirror those of several other studies that found that male odor cues were sufficient

to affect courtship behavior [21,48,89]. Our results and the results from Seagraves et al. (2016)

could be conflicting due to methodological factors. Primarily, our odor stimulus contained both

urine and feces, and potentially body odors from allowing the males to interact freely with the

bedding that was later used as the odor stimulus. Odors produced from these different sources

contain different volatile compounds [58,90], thus having urine, feces, and body odor in the

bedding used in the interaction could have served as a more potent stimulus more likely to

induce an affect than urine or body odor alone [58].

Also in contrast to the present results, Seagraves et al. (2016) found that two males

presented with female odors vocalized more when together than when alone, leading the

authors to conclude that the males were acting as an "audience" for one another that

prompted the increase of vocal output [44]. Another interpretation that the authors suggest for

their results, however, is that the female odor was acting as a potential audience to the male-

male interaction, rather than a live male audience to a male interacting with female odor. This

interpretation aligns more succinctly with stricter definitions of audience effects that specify

that the audience does not directly communicate with the members of the interaction [39].

Seagraves et al. (2016) compare the vocal behavior of one live individual presented with odor

to two live individuals presented with odor; it is logical that the addition of another live

individual that is able to vocally interact with the focal individyal would increase calling

compared to when the focal individual is simply presented with female odor. Thus, it is possible

that rather than a male "audience" prompting the increase in calling, it was the presence of a

live interacting individual being present at all in the interaction that prompted increases in

calling. Here, in both our control condition (Fem) and our experimental condition (Fem+Male),

there was always a live female present to promote vocalizations. The results here conflated

with the Seagraves et al. (2016) study together may suggest that the presence of female odor

during a male-male interaction increases vocal production, while the presence of male odor

during a male-female interaction decreases vocal production, emphasizing the effect of context

of the interaction on vocal communication.

It may be logical for males to show decreased courtship motivation when smelling other

males, through the reduction of and simplification of calling behaviors, because cues that

indicate the presence of potential reproductive competitors can decrease sexual motivation

and induce rapid behavioral plasticity [54,91]. Males may "budget" copulatory efforts

depending on factors like risk of competition [50,92–98]. Across taxa, male precopulatory

courtship behaviors can be reduced as the density of competitors increases [92,96]. This could

be due to adaptive changes in reproductive strategies by the males or sexual interference from

the competitors [92,96].

A second cohort of individuals were exposed to the odor of their cage mate, which with

they had an established dominance relationship, and interestingly, when the odor was familiar

to the focal individual, their vocal behavior did not differ from when no odor was present at all.

Not only was there no difference in dominant and subordinate individuals (to be discussed in

more detail in subsequent sections), but number of calls, duration of calls, and repertoire of

calls were not affected by the familiar competitor odor at all. This important distinction

between the responses of individuals to familiar compared to unfamiliar competitor odors

could have occurred due to various reasons. Mice can identify individuals they are familiar with

through chemical cues in urine [99]. After frequent exposure to odor cues from conspecifics,

mice can habituate to these odors, and thus have a less pronounced response to them

[89,100,101]. The dear enemy hypothesis predicts that familiar competitors should elicit less of

a reaction from territory owners than unfamiliar competitors, due to recognition and

knowledge of the benefits and costs of defending against those particular individuals [102–

104]. Our results are consistent with the dear enemy hypothesis in that the cues from familiar

competitors elicit less of a response (less of a decrease in calls) than cues from unfamiliar

competitors. The familiarity of the male cues present could determine whether or not males

pay more attention to the male cues, or ignore them and shift focus to the female. As the dirty

bedding used to simulate the familiar audience came from the home cage of the focal individual

and their cage mate, it is also possible that the males in this cohort could have been affected by

smelling their own odors as well. A potential way for future studies to explore if, in this context,

the decrease in vocal behavior in response to unfamiliar competitor odor is paired with an

increase in behaviors aimed to signal to the competitor could be to assess competitive scent

marking behavior of the focal males when confronted with familiar vs unfamiliar odor

[89,105,106]. If males confronted with unfamiliar odor are shifting their efforts from courting

the females to focusing on competing with potential competitors, we would expect to see an

increase in marking behavior is correlated with a decrease in vocal behavior.

BBVs spur decreased courtship effort in male mice

All experimental groups in this study decreased calling in response to playback of broadband

vocalizations, BBVs, and no other factors measured affected this response (Fig 4). While the

function of BBVs across contexts has been largely unexplored, evidence suggests that in certain

courtship contexts, BBVs serve to communicate a negative affective state of the signaler

[26,34,107,108]. Surprisingly, we found that when individuals were exposed to the odor of an

unfamiliar competitor, the proportion of 50 kHz harmonic USVs they used increased in the

recovery period after exposure to BBV playback, and it was in this period that they used the

highest proportion of harmonics in their calling, even more so than in the baseline period (Fig

5A). More in depth research will have to explore the intricacies of why the discontinuation of

BBV playback might spur a rapid increase in harmonic usage only when cues from an unfamiliar

potential competitor are present, though this could potentially reflect habituation to the

competitor cues as the trial goes on. This result, though difficult to interpret, does however

support that the cessation of BBV playback changes the trajectory of syllable usage in male

mice when cues indicating the presence of an unfamiliar competitor are present.

BBVs may not always produce a negative reproductive response, and are sometimes

correlated with increases in mounting and calling [11,108,109]. It is possible that the potential

different functions of BBVs manifest when BBVs are paired with other nonvocal behaviors, and

the isolated vocal characteristics of the BBVs may indicate a negative valence, and thus

promote decreased courtship effort in receivers without access to additional cues from the

female.

Our results exemplify that these signals are significant and attention-grabbing to male

mice, seemingly regardless of certain identity and environmental factors. BBVs and their

correlated behavioral responses remain a largely untapped line of research; and very few

studies have explored the effects of BBV playback isolated from the associated physical

rejection behaviors often correlated with BBV production. Greater research devoted to this

area of mouse communication could help elucidate how different internal and external factors

may affect responses to these signals.

Lack of an effect of dominance status

In this study, dominance status did not significantly affect any vocal or nonvocal behavioral

measurements in either the unfamiliar odor cohort or the familiar odor cohort. This was not

what we predicted, as it is generally accepted that dominant male mice produce more USVs

than subordinate individuals in response to a novel female [110,111], though USVs are not a

standard way to measure dominance as they do not directly measure access to mates, but

rather reflect mating effort [22]. Because USV production is highly dependent on other

contextual factors, the relationship between USV production and dominance is likely influenced

by finetuned differences in social history and situational context rather than being simply and

strictly linear [112]. For example, certain studies have found the relationship between

dominance and increased USV production depends on the types of cues present and the

recency of social defeat, and can even reverse [7,19,36]. Interestingly, in this study, the

individual with the highest number of baseline calls in all conditions (both odor cohorts in both

the Fem and Fem+Male conditions) was subordinate (Fig 3A), though there was no overall

statistically significant effect of dominance on any measurements. Despite observations that

this strain of mice may not exhibit overt severe dominance in their hierarchies [113], all housed

pairs in this experiment showed consistent and reliable dyadic dominance relationships through

multiple rounds of the tube test. Thus, depending on the context of an interaction, dominance

may affect behaviors such as offensiveness and aggression during tube tests, while not

manifesting in indirect courtship assays like that used in this study. It is also possible that,

because of the smaller sample size used in this study, decreased statistical power in the

statistical models we used may make it more difficult to ascertain an effect of dominance.

Future studies on fewer experimental groups with larger sample sizes within each group could

resolve this issue.

Dominant and subordinate males also tend to differ in their response to social challenges

[114–116]. Dominant rodents tend to be more susceptible to negative health consequences

and behavioral impairment following social challenges compared to subordinate males

[51,52,117,118], however this is most often exhibited during direct interactions with social

competitors. In this study, dominant males and subordinate males were not differentially

affected by odors of competitor males during courtship, thus odors alone may not be sufficient

to induce this effect of the dominance relationship. Salience of cues, interaction context, social

history, and genetic factors may all contribute to how the effects of dominance manifest during

courtship [110,116,119–121], and more research is needed to tease out which of these factors

are most influential on determining the effects of dominance.

Conclusions

By looking at how these multiple factors intertwine to affect behavioral decisions during

courtship, we can begin to assess what "matters" to a mouse as they navigate mating and when

and where to exert effort during courtship. Future studies should incorporate larger sample

sizes to increase statistical power when considering interactions between the variety of

variables we investigated. This study supports the potential negative valence of a relatively ill-

explored female signal: broadband vocalizations, or squeaks, in certain contexts. We

determined that while males robustly adjust their behavior in response to these calls, the

degree of this behavioral response was not altered by any of the internal or external factors we

manipulated. Furthermore, this study posits that cues indicating the presence, or "audience", of

potential competitors can alter male mouse courtship motivation and behavior, but this effect

is dependent on social history with the audience members. Our results support that certain

identity factors, like dominance status, do not affect the relationship between these cues and

the observed behavioral response.

Acknowledgements

660 The authors would like to thank and acknowledge Dr. Kayleigh E. Hood, Ph.D. for providing

661 mentorship, and guidance regarding the split cage assay, and Morgan Leever, M.P.A., M.S. for

662 consulting regarding statistical analysis.

References

1. Holy TE, Guo Z. Ultrasonic songs of male mice. PLoS Biology. 2005;3(12):1–10.

2. Seyfarth RM, Cheney DL. Signalers and Receivers in Animal Communication. Annual
Review of Psychology. 2003;54:145–73.

3. Wiley RH. Signal Detection and Animal Communication. Advances in the Study of Behavior.
2006;36(06):217–47.

4. Hurley LM, Kalcounis-Rueppell MC. State and Context in Vocal Communication of Rodents.
In: Rodent Bioacoustics. 2018. p. 191–221.

5. Leger DW. Contextual sources of information and responses to animal communication
signals. Psychological Bulletin. 1993;113(2):295–304.

6. Palmer CR, Kristan WB. Contextual modulation of behavioral choice. Current Opinion in
Neurobiology. 2011 Aug 1;21(4):520–6.

7. D'Amato FR. Male-male interference in a reproductive context: Effect of social status in
mice. Ethology Ecology and Evolution. 1991;3(4):295–306.

8. Todt D, Naguib M. Vocal Interactions in Birds: The Use of Song as a Model in
Communication. Advances in the Study of Behavior. 2000;29(C):247–96.

9.   Chabout J, Sarkar A, Dunson DB, Jarvis ED. Male mice song syntax depends on social
contexts and influences female preferences. Frontiers in Behavioral Neuroscience.
2015;9(APR):1–16.

10. Hanson JL, Hurley LM. Female presence and estrous state influence mouse ultrasonic
courtship vocalizations. PLoS ONE. 2012;7(7).

11. Finton CJ, Keesom SM, Hood KE, Hurley LM. What's in a squeak? Female vocal signals
predict the sexual behaviour of male house mice during courtship. Animal Behaviour.
2017;126:163–75.

12. Brumm H, Naguib M. Chapter 1 Environmental Acoustics and the Evolution of Bird Song.
In: Advances in the Study of Behavior [Internet]. Elsevier; 2009 [cited 2024 Sep 30]. p. 1–
33. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0065345409400019

13. Mol C, Chen A, Kager RWJ, Ter Haar SM. Prosody in birdsong: A review and perspective.
Neuroscience & Biobehavioral Reviews. 2017 Oct;81:167–80.

14. Opaev AS. The Communicative Value of Complex Singing in Passerine Birds. Biol Bull Russ
Acad Sci. 2022 Dec;49(10):1750–68.

15. Riters LV, Stevenson SA. Using seasonality and birdsong to understand mechanisms
underlying context-appropriate shifts in social motivation and reward. Hormones and
Behavior. 2022 Jun;142:105156.

16. Ritschard M, Brumm H. Zebra finch song reflects current food availability. Evol Ecol. 2012
Jul;26(4):801–12.

17. Fischer J, Hammerschmidt K. Ultrasonic vocalizations in mouse models for speech and
socio-cognitive disorders: Insights into the evolution of vocal communication. Genes,
Brain and Behavior. 2011;10(1):17–27.

18. Granon S, Faure A, Chauveau F, Cressant A, Ey E. Why Should My Mouse Call Me? Acoustic
Communication in Mouse Models of Social Disorders: Ultrasonic Vocalizations as an
Index of Emotional and Motivational States. Handbook of Behavioral Neuroscience.
2018;25:423–31.

19. Lumley LA, Sipos ML, Charles RC, Charles RF, Meyerhoff JL. Social stress effects on
territorial marking and ultrasonic vocalizations in mice. Physiology and Behavior.
1999;67(5):769–75.

20. Sangiamo DT, Warren MR, Neunuebel JP. Ultrasonic signals associated with different types
of social behavior of mice. Nature Neuroscience. 2020;23(3):411–22.

21. D'amato FR. Courtship ultrasonic vocalizations and social status in mice. Animal Behaviour.
1991;41(5).

22. Egnor SER, Seagraves KM. The contribution of ultrasonic vocalizations to mouse courtship.
Current Opinion in Neurobiology. 2016;38(Figure 1):1–5.

23. Ferhat AT, Torquet N, Le Sourd AM, De Chaumont F, Olivo-Marin JC, Faure P, et al.
Recording mouse ultrasonic vocalizations to evaluate social communication. Journal of
Visualized Experiments. 2016;2016(112):1–12.

24. Portfors CV, Perkel DJ. The role of ultrasonic vocalizations in mouse communication.
Current Opinion in Neurobiology. 2014;28:115–20.

25. Neunuebel JP, Taylor AL, Arthur BJ, Roian Egnor SE. Female mice ultrasonically interact
with males during courtship displays. eLife. 2015;4(MAY):1–24.

26. Sugimoto H, Okabe S, Kato M, Koshida N, Shiroishi T, Mogi K, et al. A role for strain
differences in waveforms of ultrasonic vocalizations during Male-Female interaction.
PLoS ONE. 2011;6(7).

27. White NR, Prasad M, Barfield RJ, Nyby JG. 40- and 70-kHz vocalizations of mice (*Mus
musculus*) during copulation. Physiology and Behavior. 1998;63(4):467–73.

28. Scattoni ML, Gandhy SU, Ricceri L, Crawley JN. Unusual repertoire of vocalizations in the
BTBR T+tf/J mouse model of autism. PLoS ONE. 2008;3(8).

29. Heckman J, McGuinness B, Celikel T, Englitz B. Determinants of the mouse ultrasonic vocal
structure and repertoire. Neuroscience and Biobehavioral Reviews. 2016;65.

30. Grimsley JMS, Monaghan JJM, Wenstrup JJ. Development of social vocalizations in mice.
PLoS ONE. 2011;6(3).

31. Ruat J, Genewsky AJ, Heinz DE, Kaltwasser SF, Canteras NS, Czisch M, et al. Why do mice
squeak? Toward a better understanding of defensive vocalization. iScience.
2022;25(7):104657.

32. Knights CB, Gentry C, Bevan S. Partial medial meniscectomy produces osteoarthritis pain-
related behaviour in female C57BL/6 mice. Pain. 2012;153(2):281–92.

33. Anderson LL, Everett-Morgan D, Petkova SP, Silverman JL, Arnold JC. Ictal vocalizations in
the Scn1a+/− mouse model of Dravet syndrome. Epilepsia Open. 2023;8(3):776–84.

34. Keesom SM, Hurley LM. Socially induced serotonergic fluctuations in the male auditory
midbrain correlate with female behavior during courtship. Journal of Neurophysiology.
2016;115(4):1786–96.

35. Hood KE, Long E, Navarro E, Hurley LM. Playback of broadband vocalizations of female
mice suppresses male ultrasonic calls. PloS one. 2023;18(1):e0273742.

36.   Nyby JG, Dizinno GA, Whitney G. Social status and ultrasonic vocalizations of male mice.
747   Behavioral Biology. 1976;18(2):285–9.

748   37.   Marler P, Dufty A, Pickert R. Vocal communication in the domestic chicken: I. Does a
749   sender communicate information about the quality of a food referent to a receiver?
750   Animal Behaviour. 1986;34(PART 1):188–93.

751   38.   Marler P, Dufty A, Pickert R. Vocal communication in the domestic chicken: II. Is a sender
752   sensitive to the presence and nature of a receiver? Animal Behaviour. 1986 Feb;34:194–
753   8.

754   39.   Matos R J, Schlupp I. Performing in front of an audience: signallers and the social
755   environment. In: Animal Communication Networks. Cambridge, UK: Cambridge
756   University Press; 2005. p. 63–83.

757   40.   Baltz AP, Clark AB. Extra-pair courtship behaviour of male budgerigars and the effect of an
758   audience. Animal Behaviour. 1997 May;53(5):1017–24.

759   41.   Evans CS, Marler P. Food calling and audience effects in male chickens, *Gallus gallus*: their
760   relationships to food availability, courtship and social facilitation. Animal Behaviour. 1994
761   May;47(5):1159–70.

762   42.   Sánchez NV, Mennill DJ. Behavioural consequences of conspecific neighbours: a systematic
763   literature review of the effects of local density on avian vocal communication. J Ornithol.
764   2024 Oct;165(4):847–59.

765   43.   Vignal C, Mathevon N, Mottin S. Audience drives male songbird response to partner's
766   voice. Nature. 2004;430(6998):448–51.

767   44.   Seagraves KM, Arthur BJ, Egnor SER. Evidence for an audience effect in mice: Male social
768   partners alter the male vocal response to female cues. Journal of Experimental Biology.
769   2016;219(10):1437–48.

770   45.   Plath M, Schlupp I. Misleading Mollies. Communicative & Integrative Biology.
771   2008;1(2):199–203.

772   46.   Bierbach D, Girndt A, Hamfler S, Klein M, Mücksch F, Penshorn M, et al. Male fish use prior
773   knowledge about rivals to adjust their mate choice. Biology Letters. 2011;7(3):349–51.

774   47.   Castellano S, Friard O, Pilastro A. The audience effect and the role of deception in the
775   expression of male mating preferences. Animal Behaviour. 2016;115:273–82.

776   48.   Aragón P. Conspecific male chemical cues influence courtship behaviour in the male newt
777   *Lissotriton boscai*. Behav. 2009;146(8):1137–51.

49. Matos R J, McGregor PK. The Effect of the Sex of an Audience on Male-Male Displays of
Siamese Fighting Fish (*Betta splendens*). Behaviour. 2002;139(9):1211–21.

50. Johnston RE. Eavesdropping and scent over-marking. In: Animal Communication Networks.
Cambridge, UK: Cambridge University Press; 2005. p. 344–71.

51. D'Amato FR, Rizzi R, Moles A. A model of social stress in dominant mice: Effects on
sociosexual behaviour. Physiology and Behavior. 2001;73(3):421–6.

52. Larrieu T, Cherix A, Duque A, Rodrigues J, Lei H, Gruetter R, et al. Hierarchical Status
Predicts Behavioral Vulnerability and Nucleus Accumbens Metabolic Profile Following
Chronic Social Defeat Stress. Current Biology. 2017;27(14):2202-2210.e4.

53. Chaouloff F. Serotonin, stress and corticoids. Journal of Psychopharmacology.
2000;14(2):139–51.

54. Fernández-Vargas M. Presence of a potential competitor and its individual identity
modulate ultrasonic vocalizations in male hamsters. Animal Behaviour. 2018;145.

55. Chambers LK, Singleton GR, Krebs CJ. Movements and social organization of wild house
mice (*Mus domesticus*) in the wheatlands of northwestern Victoria, Australia. Journal of
Mammalogy. 2000;81(1).

56. Brown RZ. Social Behavior, Reproduction, and Population Changes in the House Mouse
(*Mus musculus L.*). Ecological Monographs. 1953;23(3).

57. Fan Z, Zhu H, Zhou T, Wang S, Wu Y, Hu H. Using the tube test to measure social hierarchy
in mice. Nature Protocols. 2019;14(3):819–31.

58. Johnston RE. Chemical communication in rodents: From pheremones to individual
recognition. Journal of Mammalogy. 2003 Nov;84(4):1141–62.

59. Frynta D, Baladová M, Eliášová B, Lišková S, Landová E. Why not to avoid the smell of
danger? Unexpected behavior of the Cypriot mouse surviving on the island invaded by
black rats. Current Zoology. 2015 Aug 1;61(4):781–91.

60. Friard O, Gamba M. BORIS: a free, versatile open-source event-logging software for
video/audio coding and live observations. Methods in Ecology and Evolution.
2016;7(11):1325–30.

61. Warton DI, Hui FKC. The arcsine is asinine: The analysis of proportions in ecology. Ecology.
2011;92(1).

62. Feder PI, Aume LL, Triplett CA, Simmons JE, Narotsky MG. Analysis of proportional data in
reproductive and developmental toxicity studies: Comparison of sensitivities of logit

transformation, arcsine square root transformation, and nonparametric analysis. Birth
Defects Research. 2020;112(16):1260–72.

63. Pituch KA, Stevens JP. Binary Logistic Regression. Applied Multivariate Statistics for the
Social Sciences. 2020;454–90.

64. Games PA, Winkler HB, Probert DA. Robust Tests for Homogeneity of Variance.
Educational and Psychological Measurement. 1972;32(4).

65. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B
(Methodological). 1995;57(1).

66. Wöhr M, Roullet FI, Hung AY, Sheng M, Crawley JN. Communication Impairments in Mice
Lacking Shank1: Reduced Levels of Ultrasonic Vocalizations and Scent Marking Behavior.
Crusio WE, editor. PLoS ONE. 2011 Jun 9;6(6):e20631.

67. Rousseeuw PJ, Hubert M. Robust statistics for outlier detection. WIREs Data Min & Knowl.
2011 Jan;1(1):73–9.

68. Hafez B, Hafez ESE. Reproductive Behavior. In: Hafez B, Hafez ESE, editors. Reproduction in
Farm Animals [Internet]. 1st ed. Wiley; 2000 [cited 2024 Oct 3]. p. 291–306. Available
from: https://onlinelibrary.wiley.com/doi/10.1002/9781119265306.ch19

69. Tachibana SI, Touhara K, Ejima A. Modification of Male Courtship Motivation by Olfactory
Habituation via the GABAA Receptor in *Drosophila melanogaster*. Marion-Poll F, editor.
PLoS ONE. 2015 Aug 7;10(8):e0135186.

70. Ishii KK, Touhara K. Neural circuits regulating sexual behaviors via the olfactory system in
mice. Neuroscience Research. 2019 Mar;140:59–76.

71. Iwasaki M, Poulsen TM, Oka K, Hessler NA. Singing-Related Activity in Anterior Forebrain
of Male Zebra Finches Reflects Courtship Motivation for Target Females. Coleman MJ,
editor. PLoS ONE. 2013 Nov 29;8(11):e81725.

72. Okanoya K, Screven LA. Rodent Vocalizations: Adaptations to Physical, Social, and Sexual
Factors. In: Dent ML, Fay RR, Popper AN, editors. Rodent Bioacoustics [Internet]. Cham:
Springer International Publishing; 2018 [cited 2024 Oct 3]. p. 13–41. (Springer Handbook
of Auditory Research; vol. 67). Available from: http://link.springer.com/10.1007/978-3-
319-92495-3_2

73. Ventura-Aquino E, Portillo W, Paredes RG. Sexual Motivation: A Comparative Approach in
Vertebrate Species. Curr Sex Health Rep. 2018 Sep;10(3):114–23.

74.   Nunez AA, Nyby JG, Whitney G. The effects of testosterone, estradiol, and
dihydrotestosterone on male mouse (*Mus musculus*) ultrasonic vocalizations. Hormones
and Behavior. 1978;11(3).

75.   Fernández-Vargas M. Vocal Signals of Sexual Motivation in Male and Female Rodents.
Current Sexual Health Reports. 2018;10(4).

76.   Nunez AA, Tan DT. Courtship ultrasonic vocalizations in male Swiss-Webster mice: Effects
of hormones and sexual experience. Physiology and Behavior. 1984;32(5).

77.   Hausberger M, Henry L, Testé B, Barbu S. Contextual Sensitivity and Bird Song: A Basis for
Social Life. In: Oller DK, Griebel U, editors. Evolution of Communicative Flexibility
[Internet]. The MIT Press; 2008 [cited 2024 Oct 2]. p. 121–38. Available from:
https://direct.mit.edu/books/book/2702/chapter/73061/Contextual-Sensitivity-and-Bird-
Song-A-Basis-for

78.   Gersick AS, White DJ. Male cowbirds vary the attractiveness of courtship songs with
changes in the social context. Behav. 2018;155(1):1–25.

79.   West MJ, King AP, Eastzer DH. Validating the female bioassay of cowbird song: Relating
differences in song potency to mating success. Animal Behaviour. 1981 May;29(2):490–
501.

80.   Burkhard TT, Sachs ER, Phelps SM. Female preferences for high vocal effort in singing
mice. Behaviour. 2023;160(3–4).

81.   Burkhard TT, Westwick RR, Phelps SM. Adiposity signals predict vocal effort in Alston's
singing mice. Proceedings of the Royal Society B: Biological Sciences. 2018;285(1877).

82.   Dore AA, McDowall L, Rouse J, Bretman A, Gage MJG, Chapman T. The role of complex
cues in social and reproductive plasticity. Behavioral Ecology and Sociobiology.
2018;72(8).

83.   Johnston RE. Chemical Signals and Reproductive Behavior. In: Pheromones and
Reproduction in Mammals. 1983.

84.   Guevara-Fiore P, Endler JA. Female receptivity affects subsequent mating effort and mate
choice in male guppies. Animal Behaviour. 2018;140.

85.   Jenikejew J, Wauters J, Dehnhard M, Scheumann M. The female effect - How female
receptivity influences faecal testosterone metabolite levels, socio-positive behaviour and
vocalization in male Southern white rhinoceroses. Conservation Physiology. 2021;9(1).

86.   Barrass R. A Quantitative Study of the Behaviour of the Male *Mormoniella Vitripennis*
(Walker) (Hymenoptera, Pteromalidae) Towards Two Constant Stimulus-Situations.
Behaviour. 1961;18(4).

87.   Girard MB, Elias DO, Kasumovic MM. Female preference for multi-modal courtship:
Multiple signals are important for male mating success in peacock spiders. Proceedings of
the Royal Society B: Biological Sciences. 2015;282(1820).

88.   Linley JR, Hinds MJ. Quantity of the male ejaculate influenced by female unreceptivity in
the fly, *Culicoides melleus*. Journal of Insect Physiology. 1975;21(2).

89.   Johnston RE. Individual Odors and Social Communication. Individual Recognition, Kin
Recognition, and Scent Over-Marking. Advances in the Study of Behavior. 2008;38.

90.   Goodrich BS, Gambale S, Penncuik PR, Redhead TD. Volatile compounds from excreta of
laboratory mice (*Mus musculus*): Preliminary examination of composition and effects on
behavior. J Chem Ecol. 1990 Jul;16(7):2107–20.

91.   Spagopoulou F, Vega-Trejo R, Head ML, Jennions MD. Shifts in reproductive investment in
response to competitors lower male reproductive success. American Naturalist.
2020;196(3).

92.   Bretman A, Gage MJG, Chapman T. Quick-change artists: Male plastic behavioural
responses to rivals. Trends in Ecology and Evolution. 2011;26(9).

93.   Parker GA, Ball MA, Stockley P, Gage MJG. Sperm competition games: A prospective
analysis of risk assessment. Proceedings of the Royal Society B: Biological Sciences.
1997;264(1389):1793–802.

94.   Parker GA, Ball MA, Stockley P, Gage MJG. Sperm competition games: Individual
assessment of sperm competition intensity by group spawners. Proceedings of the Royal
Society B: Biological Sciences. 1996;263(1375).

95.   Parker GA. Conceptual developments in sperm competition: A very brief synopsis:
Developments in sperm competition. Philosophical Transactions of the Royal Society B:
Biological Sciences. 2020;375(1813):120–4.

96.   Bretman A, Westmancoat JD, Gage MJG, Chapman T. Males use multiple, redundant cues
to detect mating rivals. Current Biology. 2011;21(7).

97.   Dore AA, Rostant WG, Bretman A, Chapman T. Plastic male mating behavior evolves in
response to the competitive environment. Evolution. 2021 Jan 1;75(1):101–15.

98.   Dixson AF, Anderson MJ. Sexual behavior, reproductive physiology and sperm competition
in male mammals. Physiology & Behavior. 2004 Nov;83(2):361–71.

99.   Brennan PA. The nose knows who's who: Chemosensory individuality and mate
recognition in mice. Hormones and Behavior. 2004;46(3).

100. DelBarco-Trillo J, Ferkin MH. Male mammals respond to a risk of sperm competition
conveyed by odours of conspecific males. Nature. 2004;431(7007).

101. Rowland WJ. Aggression versus courtship in threespine sticklebacks and the role of
habituation to neighbours. Animal Behaviour. 1988;36(2).

102. Utne-Palm AC, Hart PJB. The effects of familiarity on competitive interactions between
threespined sticklebacks. Oikos. 2000;91(2).

103. Ydenberg RC, Giraldeau LA, Falls JB. Neighbours, strangers, and the asymmetric war of
attrition. Animal Behaviour. 1988;36(2).

104. Fisher RA. Evolution and bird sociality. In: Huxley J, Hardy AC, Ford EB, editors. Evolution as
a Process. London: Alen & Unwin; 1954. p. 71–83.

105. Dobly A. Scent marking by common voles *Microtus arvalis* in the presence of a same-sex
neighbour. Acta Theriologica. 2005;50(3).

106. Carazo P, Font E, Desfilis E. Beyond "nasty neighbours" and "dear enemies"? Individual
recognition by scent marks in a lizard (Podarcis hispanica). Animal Behaviour. 2008;76(6).

107. Keesom SM, Finton CJ, Sell GL, Hurley LM. Early-Life Social Isolation Influences Mouse
Ultrasonic Vocalizations during Male-Male Social Encounters. PLoS ONE. 2017;12(1).

108. Wang H, Liang S, Burgdorf J, Wess J, Yeomans J. Ultrasonic Vocalizations Induced by Sex
and Amphetamine in M2, M4, M5 Muscarinic and D2 Dopamine Receptor Knockout
Mice. PLoS ONE. 2008;3(4):1893.

109. Lupanova AS, Egorova MA. Vocalization of sex partners in the house mouse (*Mus
Musculus*). Journal of Evolutionary Biochemistry and Physiology. 2015;51(4).

110. Fulenwider HD, Caruso MA, Ryabinin AE. Manifestations of domination: Assessments of
social dominance in rodents. Genes, Brain and Behavior. 2022;21(3):1–15.

111. Wolff RJ. Mating behaviour and female choice: their relation to social structure in wild
caught House mice (*Mus musculus*) housed in a semi-natural environment. Journal of
Zoology. 1985;207(1).

112. Wang F, Kessels HW, Hu H. The mouse that roared: Neural mechanisms of social hierarchy.
Trends in Neurosciences. 2014;37(11):674–82.

113. Southwick CH, Clark LH. Aggressive behaviour and exploratory activity in fourteen mouse
strains. American Zoologist. 1966;6:559.

114. Ling TJ, Summers CH, Renner KJ, Watt MJ. Opponent recognition and social status
differentiate rapid neuroendocrine responses to social challenge. Physiology & Behavior.
2010 Apr;99(5):571–8.

115. Friesen CN, Maclaine KD, Hofmann HA. Social status mediates behavioral, endocrine, and
neural responses to an intruder challenge in a social cichlid, Astatotilapia burtoni.
Hormones and Behavior. 2022 Sep;145:105241.

116. Cooper MA, Clinard CT, Morrison KE. Neurobiological mechanisms supporting experience-
dependent resistance to social stress. Neuroscience. 2015 Apr;291:1–14.

117. Sgoifo A, Costoli T, Meerlo P, Buwalda B, Pico'-Alfonso MA, De Boer S, et al. Individual
differences in cardiovascular response to social challenge. Neuroscience and
Biobehavioral Reviews. 2005;29(1 SPEC. ISS.).

118. Larrieu T, Sandi C. Stress-Induced Depression: Is Social Rank a Predictive Risk Factor?
BioEssays. 2018;40(7).

119. Tibbetts EA, Pardo-Sanchez J, Weise C. The establishment and maintenance of dominance
hierarchies. Phil Trans R Soc B. 2022 Feb 28;377(1845):20200450.

120. Dewsbury DA. Dominance Rank, Copulatory Behavior, and Differential Reproduction. The
Quarterly Review of Biology. 1982 Jun;57(2):135–59.

121. Chase ID, Seitz K. Self-structuring properties of dominance hierarchies. A new perspective.
In: Advances in Genetics. 2011.

A. Unfamiliar odor cohort

B. Familiar odor cohort

A. Unfamiliar odor cohort

B. Familiar odor cohort

A. Unfamiliar odor cohort

Duration of male investigation (s) vs Test Period (Baseline, Playback, Recovery)

Legend: Dom Fem, Dom Fem+Male, Sub Fem, Sub Fem+Male

B. Unfamiliar odor cohort

Duration of female investigation (s) vs Test Period (Baseline, Playback, Recovery)

C. Familiar odor cohort

Duration of male investigation (s) vs Test Period (Baseline, Playback, Recovery)

Legend: Dom Fem, Dom Fem+Male, Sub Fem, Sub Fem+Male

D. Familiar odor cohort

Duration of female investigation (s) vs Test Period (Baseline, Playback, Recovery)