What matters to a mouse? Effects of internal and external context on

male vocal response to female squeaks

Lauren R Brunner[1,2]*, Laura M Hurley[1,2]

[1]Hurley Laboratory, Department of Biology, Indiana University, Bloomington, Indiana, USA

[2]Center for the Integrative Study of Animal Behavior, Indiana University, Bloomington, Indiana,

USA

*Corresponding author

Email: laribrun@iu.edu (LRB)

Abstract

Individuals plastically adjust their signaling behavior depending on the social context of an interaction, but which aspects of context elicit the strongest responses from individuals is often difficult to determine. To explore how internal and external contextual factors influence the communication of house mice, we employed a split-cage playback assay to assess how dominant and subordinate male mice differed in their ultrasonic vocalization (USV) production in response to playback of broadband vocalizations (BBVs, or squeaks). We assessed how individuals altered their behavior depending on the odors present during the assay: either just female odors (Fem condition) or female odors in addition to the odors of potential competitors, other males (Fem+Male condition). Using this repeated-measures design, we evaluated two distinct cohorts of individuals: males exposed to the odor of familiar competitors in the Fem+Male condition (Familiar odor cohort), and males exposed to the odor of unfamiliar competitors in the Fem+Male condition (Unfamiliar odor cohort). While dominants and subordinates did not differ in their baseline vocal production or response to squeaks, all groups reduced their vocal production in response to BBV playback, and the addition of unfamiliar male odor prompted mice to decrease their baseline levels of calling and decrease the complexity of their vocal repertoire. Importantly, this effect of additional male odor was not present when individuals were presented with male odor familiar to them. These findings suggest that mice alter their vocal behavior during courtship interactions in response to cues that indicate the presence of potential competitors, and this response is modulated by the familiarity of these competitor cues.

Introduction

The ability to process and respond appropriately to conspecific signals is crucial for animals as they navigate social interactions [1–3]. How an individual processes and responds to signals may heavily depend on the context of the situation [4–6], and the role of context in vocal communication in particular is highly documented across taxa [7–11]. Research on how contextual influences such as the presence of competitors and potential mates, seasonality, changes in steroid hormones, and resource abundance affect song production and song composition in birds has revealed that many aspects of communication depend on context, and prompted advances and questions on how context can influence the signaling in other species [12,12–16]. These include laboratory mice, studied not only as a model of rodent communication signals, but also for their potential applicability to human communication disorders [17,18].

House mice (*Mus musculus*), produce a variety of vocalizations that can vary in usage, structure, and meaning across different contexts [1,10,11,19,20]. Vocal communication in mice is prominent during courtship interactions, in which calls are used to communicate receptivity and sexual motivation [1,10,21–24]. One type of call, ultrasonic vocalizations (USVs), are high-frequency calls (30-110 kHz) produced by both males and females during social interactions [1,20,24–26]. These calls are produced in large numbers by males during opposite-sex interactions, and are often referred to as "courtship calls" because of their prosocial function to facilitate mating [1,24,27] (Fig 1A). Male USV production thus may indicate male courtship effort and motivation [22]. Mouse vocal repertoires, or the composition of different types of syllables individuals use, are complex and can vary depending on factors such as age, genetics, and behavioral situation [10,28–30]. One type of USV in particular, 50 kHz harmonic USVs, tend to be

54 longer in duration and more complex structurally than other classes of USVs. These calls are

55 closely associated temporally with incidents of male mounting behavior [10] (Fig 1A).

> **Fig 1: Representative spectrograms of (A) Ultrasonic vocalizations (USVs) and (B) Broadband vocalizations (BBVs).** The circled vocalization in (A) is an example of a 50 kHz

Another type of call, broadband vocalizations (BBVs), or squeaks, are lower-frequency

calls with a broader frequency range than USVs, featuring many harmonics in (Fig 1B). Both male

and female mice produce BBVs across varying contexts, including during distress, handling, and

conspecific interactions [26,31–33]. When produced by females during the early stages of

courtship interactions, these calls are closely correlated with females performing physical

rejection behaviors like kicking, lunging, and attempted escape [11,34]. Thus, during courtship,

female-produced BBVs may be negatively-valenced, and function to express a lack of female

receptivity to sexual advances by males. In accordance with the connection of squeaks to physical

rejection, male mice reduce their USV output in response to BBV playback during limited

interactions with female mice [35]. However, little is known about what sorts of contextual

factors may influence this response to BBVs, and which levels of context may "matter" the most

to mice as they interact.

The goal of this study was to explore the effects of both internal and external contextual

factors on male vocal production and vocal repertoire in response to female squeaks. As an

internal contextual factor, we evaluated dominance status in all of our individuals. Some calls are

more associated with the production of behaviors classified as "more dominant" such as

aggression, and courting [20]. Dominant and subordinate mice tend to differ in their vocal

responses to female cues, though the direction of this difference varies across studies. For

example, one study found that dominant male mice call more quickly and more often compared

to subordinate males in response to female stimuli [36], while in contrast, another found that

subordinates called the most when interacting with a female [7]. However, this was only true

when subordinates were tested in the experimental cage first; if subordinates were recorded in

the same arena in which a dominant male had previously interacted with a female, their calling

was sharply reduced, indicating that some cues from the dominant males were able to stunt the

courtship behavior of subordinate mice. The same effect did not occur when dominants were

tested after subordinates [7]. Thus, the effects of dominance status on vocal production in male

mice appear to be highly-context dependent, and worthy of further investigation.

In addition to the playback of BBVs, the external contextual factor of interest to us was

the presence of non-vocal cues that may alter male response to BBVs: specifically, odor from

other male mice. Audience effects, in which individuals adjust their behavior in the presence of

additional individuals, are well-documented examples of context-dependent communication

[37]. Studies in songbirds and domesticated fowl have revealed that the presence of audiences

during courtship signaling can influence not only the acoustic properties of song and song

repertoire, but also the motivation to interact with their partner [38–41]. Previous studies have

established that certain strains of mice are susceptible to audience effects when responding to

female cues [37,42]. For example, Seagraves et al. (2016) found that increases in vocal production

in response to live and anesthetized male audience members may be due to signalers shifting

their vocalizations to be directed to males, potentially in a competitive display. Thus, the

presence of an audience may cause mice to adjust their vocal output and the function of their

calling may change, similar to the strategic changes influenced by audiences across other

contexts [37,41,43–45].

In order to assess the effects additional male cues and social status on male behavioral

response to BBVs, we employed a recently established paradigm [35] that allows for the

decoupling of BBVs and their frequently associated physical rejection behaviors. We conducted

two types of trials using this paradigm that varied based on the odors presented to the focal male

during the trial: a trial in which the male was exposed to female odor only (Fem odor condition)

and a trial in which the male was exposed to both female odor and odor from additional males

(Fem+Male odor condition). We predicted that the addition of male odor during these

interactions would increase the degree by which males downregulated their vocal output in

response to BBV playback, and that this effect would be dependent on social status. It has been

reported that dominant mice are more sensitive to social challenges than subordinate mice [46–

48], so we predicted that dominant males would alter their behavior in response to a simulated

social competitor more than subordinates by showing a larger decrease in USV production during

BBV playback if additional male odor was present. For the first part of this study, these assays

were conducted using male odor that was familiar to the stimulus animals.

The second half of this study included an additional cohort of animals who differed from

the first cohort in the familiarity of the mice that provided the additional odor during the trials

to assess the effects of novel vs familiar "audience members". For this familiar odor cohort, the

odor from the familiar male would come from a mouse that the focal male had interacted with

(and with whom they had an established hierarchy). This was of particular interest to us because

identity and status of audience members has been shown to alter behavior during signaling in

response to female cues [49]. We predicted that more robust differences between dominants

and subordinates would be seen in the familiar odor cohort because dominants would be

exposed to the odor of an individual known to be subordinate to them, and subordinates would

be exposed to the odor of an individual known to be dominant over them, rather than the odor

of completely unknown male mice. Surprisingly, we found that dominance status did not affect

vocal behavior, while the presence of additional male odor dampened vocal production, but only

when the odor came from a male unfamiliar to the focal individual.

Methods

Animals

Focal subjects consisted of seven-week-old male CBA/J mice (N = 23) (The Jackson Laboratory,

Bar Harbor, ME). Mice were housed in a 14:10 light:dark cycle with food and water *ad libitum*.

All mice were housed in pairs from at least one week prior to behavioral assays until the end of

the experiments in order to allow for the establishment and maintenance of a social hierarchy

within the cage. Dominance status was evaluated using multiple measurements of a standardized

tube test [35,50]. All housing pairs underwent 3-4 tube tests with no more than one test per day,

and with tests occurring no more than 4 days prior to the trial dates. Almost all mice were

consistent in their dominance status across all of their tests, aside from two pairings that

switched dominance status after the first test, but were consistent for the final three tests. At

least two days prior to experimental trials, all mice were given opposite-sex experience with

novel female CBA/J mice (The Jackson Laboratory, Bar Harbor, ME) through multiple 10 minute

interactions across the span of several days [11]. All mice received approximately the same

amount of social experience prior to trials. All stimulus animals for experimental trials were

sexually experienced adult female CBA/J house mice novel to the focal mouse.

Behavioral assay and experimental conditions

A split-cage playback paradigm previously established in our lab [35] was employed to evaluate

the effects of vocal playback on male mouse vocal behavior. In this paradigm, the focal mouse

and stimulus female are placed on opposite sides of a 12 inch x 6 inch x 6 inch cage with a

plexiglass barrier in the middle with a small hole at the bottom of the barrier that allows for

limited nose-to-nose social investigation. Dirty bedding containing urine and feces from the

stimulus female and her cage mate was placed on the focal male's side of the cage to promote

robust vocalizations [35,36], while clean bedding was placed on the female's side of the cage. A

microphone (CM16/CMPA, Avisoft Bioacoustics, Glienicke/Nordbahn, Germany) and Canon VIXIA

video camera were placed above the cage on the focal male's side to allow for recording of

ultrasonic vocalizations and non-vocal behavior, respectively, during the interaction. The entire

cage and recording setup were located in a sound-proof chamber (IAC Acoustics, Naperville, IL).

Mice were placed in this cage for 15-minute playback trials that consisted of 5 minutes of silence

(a baseline period), followed by 5 minutes of exemplar BBV playback through an ultrasonic

speaker powered by Avisoft next to the cage, followed by another five minutes of silence (a

recovery period). The segment of BBV playback was created using five repeated 1-minute

segments of BBVs collected from a previous direct interaction between a male and a female

during a period of frequent physical rejection and BBV vocalization [35]. Each of the calls during

this period were replaced with a single exemplar BBV with a duration, peak frequency, and

percentage of non-linear segments within one standard deviation of the mean for all BBVs in the

interaction in order to eliminate possible differences in behavior over the course of the playback

due to variation in BBV structure [35]. The BBV segment was played at a loudness calibrated to

be approximately matched to the intensity that male mice would experience from BBVs during

natural interactions with females, with a maximum of 104.9 dB [35].

Previous use of this paradigm in our lab has shown that males decrease their USV

production during BBV playback relative to the baseline and increase back to baseline levels

during the recovery. This response to BBV playback when males are physically separated from

the stimulus female indicates that playback of BBVs can alter male courtship vocal behavior even

when the correlated physical rejection behaviors are absent [35]. For the current study, a

modified version of this paradigm was created to allow for two distinct experimental conditions

that altered the available olfactory cues: a female-scent-only condition in which only bedding

from the stimulus female's home cage was placed on the focal animal's side of the cage (Fem

condition) and a female scent + male scent condition in which the focal male was exposed to

equal parts bedding from the stimulus female as well as bedding from a cage of male mice

(Fem+Male condition; Fig 2).

Fig 2: Experimental assay and design.
(A) Split cage paradigm used to evaluate the effects of only BBV playback on male vocal behavior, modified to create the Fem+Male condition for this study. Male and female symbols present on the male mouse's side of the cage represent the presence of dirty bedding from male and female mice. (B) Experimental groups broken up by familiarity cohort, dominance status, and odor condition.

Both assay conditions were applied to the two separate experiments with separate

cohorts of animals described in the following sections. These experiments differed based on the

origin of the male bedding used in the Fem+Male condition, described below, while the Fem

condition trials in both experiments were identical in design. All individuals in both

experiments/cohorts underwent both a Fem condition trial and a Fem+Male condition trial to

account for individual differences in vocal behavior, and the order of the trials was

counterbalanced across all individuals.

Experimental design

Experiment 1 - Unfamiliar male odor cohort: In the Fem+Male condition, individuals in

the unfamiliar male odor cohort (n=12) were exposed to equal amounts of dirty bedding from

the stimulus female's cage and dirty bedding that came from a cage of males unfamiliar to them.

These "odor donations" were from the other cages of males used in this experiment. No males

from separate cages ever interacted with one another, thus all of the "odor donations" came

from males novel to the focal individuals. The Fem+Male odor condition in this experiment was

designed this way to simulate the presence of an unfamiliar male competitor during the limited

interaction with the female.

Experiment 2 - Familiar male odor cohort: In a subsequent experiment, a new cohort

(n=16) was established as the familiar male odor cohort. For these individuals, in the Fem+Male

condition they were exposed to equal amounts of dirty bedding from the stimulus female's cage

and dirty bedding from their own home cage, in order to simulate the presence of their familiar

cage mate: their social hierarchical partner.

Audio analysis

USVs from all of the interactions were counted individually by trained individuals using Avisoft SasLab Pro software (Avisoft Bioacoustics, Glienicke/Nordbahn, Germany). A total of 74,859 USVs were counted manually during the course of this study. Durations of the USVs were automatically calculated by Avisoft. In order to assess specific components of the individuals' vocal repertoires, all USVs were manually categorized as either non-harmonic or 50 kHz harmonic USVs (Fig 1). 50 kHz harmonic USVs were specifically denoted because they are highly correlated to male mouse mounting behavior and could potentially be used to promote receptivity in females during courtship interactions [10,35].

Non-vocal behavior analysis

Non-vocal behavior was analyzed using Behavioral Observation Research Interactive Software (BORIS) [51]. The odor condition and cohort of the focal individual was blinded to the analyst for each trial. The duration of time both the focal male and stimulus female spent at the hole in the plexiglass divider was scored for all individuals. Investigation of the window was defined as the mouse placing their nose at the entrance of (< 1 cm away from the entrance) or through the opening in the barrier.

Statistical analysis

Statistical analyses were performed in IBM SPSS Statistics 28. A Pearson correlation was used to

evaluate the relationship between the baseline USVs produced in the Fem condition and the

baseline USVs produced in the Fem+Male condition across all individuals to assess consistency in

baseline vocal production. Four Pearson correlations were used to evaluate the relationship

between the investigative behavior of the focal male and the stimulus female in each cohort, for

each odor condition. Due to the difference in the Fem+Male conditions across the two cohorts,

distinct models were used for each cohort for analysis of vocal and non-vocal behavior.

To assess the effects of BBV playback across time bins and odor treatment on number of

non-harmonic calls, a three-way repeated measures ANOVA was used for the unfamiliar odor

cohort, and an identical three-way repeated measures ANOVA was used for the familiar odor

cohort (within-subjects factors: time bin (baseline, playback, or recovery) and odor treatment

(Fem or Fem+Male); between-subjects factor: dominance status (dominant or subordinate)).

Analyses on the effects of odor condition and dominance status on baseline number of USVs in

each interaction and the duration of baseline USVs produced were conducted using identical two-

way mixed ANOVAs for the unfamiliar odor cohort and familiar odor cohort (within subjects

factor: odor treatment (Fem or Fem+Male); between-subjects factor: social status (dominant or

subordinate)).

In order to investigate how the composition of vocal repertoires responded to our

manipulations, we characterized the proportion of harmonic USVs out of all USVs in each time

bin of the interaction. To assess the effects of BBV playback across time bins and odor treatment

on proportion of harmonic calls produced, proportions of harmonic calls across bins were

calculated as: number of harmonic calls in bin/number all calls in bin. These proportions were

then compared using identical binary logistic regressions for each cohort [52–54] (within-subjects

factors: time bin (baseline, playback, or recovery) and odor treatment (Fem or Fem+Male);

between-subjects factor: dominance status (dominant or subordinate)). Binary logistic

regressions were also used to evaluate the effects of odor condition and dominance status on

the baseline proportion of harmonic USVs for the unfamiliar odor cohort and familiar odor cohort

(within subjects factor: odor treatment (Fem or Fem+Male); between-subjects factor: social

status (dominant or subordinate)).

To assess the effects of BBV playback across time bins and odor treatment on non-vocal

behavior, two identical three-way repeated measures ANOVAs were used, one for each cohort

(within-subjects factors: time bin (baseline, playback, or recovery) and odor treatment (Fem or

Fem+Male); between-subjects factor: dominance status (dominant or subordinate)).

The following datasets violated the ANOVA assumption of normality and thus were

normalized using square root transformations in order to run analyses of variance: USVs across

time bins in the unfamiliar odor cohort, and female investigative behavior in the unfamiliar odor

cohort. Small violations of the homogeneity of variance requirement were noted, but as these

models were the best fit aside from this fact, and because ANOVAs are robust to small violations

of this assumption when sample sizes are equal, we proceeded with ANOVAs [55]. Benjamini-

Hochberg corrections (Q value = 0.05) were conducted in order to account for multiple

comparisons [56], and Bonferroni tests were performed as posthoc analyses.

Results

Individual baseline vocal production in both odor conditions is

positively correlated

All individuals underwent both a Fem odor condition trial, in which focal individuals were only

exposed to the odor of the female stimulus animal, and a Fem+Male odor condition trial, in which

focal individuals were exposed to both odor from the female stimulus and odor from an

additional male conspecific (Fig 2B). A Pearson Correlation revealed that the baseline vocal

production across both trials was strongly positively correlated across individuals in both the

familiar odor cohort and unfamiliar odor cohort (r = 0.735, p < 0.001) (Fig 3).

Figure 3: Correlation between vocalizations produced in the Fem condition and the Fem+Male condition for all individuals. Individuals in both cohorts are consistent in vocal output across both odor conditions, as evidenced by a positive Pearson's Correlation (r = 0.735, p < 0.001). Red data points represent individuals in the familiar odor cohort, blue data points represent individuals in the unfamiliar odor cohort. Dominants are represented by X's, and subordinates are represented by circles.

Social status does not alter behavior across cohorts or odor conditions

Social hierarchies for each cage consisted of a dominant individual and a subordinate individual,

and the placement of individuals within these hierarchies remained stable throughout multiple

rounds of tube tests. Out of 14 hierarchy pairings, only 2 pairings changed status at all throughout

the tube tests. Both of these switches occurred during the second test and the pairings were

subsequently stable throughout the rest of the three tests. When incorporating social status in

our statistical models as a between-subjects factor, there were no significant effects of social

status in any behavioral measures (Tables 1-3).

Table 1. Output of models assessing the effects of BBV playback, Fem or Fem+Male odor
condition, and dominance status on total USV production and harmonic USV production in
both cohorts.

Effect	df	Number of USVs in Unfamiliar odor cohort [a]		Proportion of 50 kHz Harmonics in Unfamiliar odor cohort [b]	
		F	p	F	p
Time Bin	2	50.538	**<.001**	4.925	**0.011**
Odor Condition	1	12.026	**0.006**	8.429	**0.005**
Dominance Status	1	0.019	0.892	0.702	0.406
Bin x Odor Condition	2	0.223	0.802	2.236	0.116
Bin x Dominance Status	2	0.614	0.551	0.131	0.877
Odor Condition x Dominance Status	1	0.742	0.409	0.022	0.882
Odor Condition x Bin x Dominance Status	2	1.203	0.321	1.093	0.342

285 [a] Nonharmonic USVs were assessed with three-way mixed ANOVAs.

286 [b] Proportion of 50 kHz harmonic USVs were assessed using binary logistic regressions.

288 **Table 2. Output of two-way mixed ANOVAs assessing the effects of Fem or Fem+Male odor**
289 **condition and dominance status on baseline USV production, baseline harmonic USV**
290 **production, and duration of baseline USVs in both cohorts.**

Effect	df	Baseline number of USVs in Unfamiliar odor cohort		Baseline 50 kHz Harmonics in Unfamiliar odor cohort		Duration of ba USVs in Unfami cohort	
		F	p	F	p	F	p
Odor Condition	1	10.005	**0.01**	10.752	**0.004**	5.285	0.04
Dominance Status	1	0.04	0.846	0.036	0.852	0.051	0.82
Odor Condition x Dominance Status	1	0.919	0.36	2.101	0.163	0.576	0.46

291 * p value is not statistically significant after correcting for multiple comparisons.

293 **Table 3. Output of three-way mixed ANOVAs assessing the effects of BBV playback, Fem or**
294 **Fem+Male odor condition, and dominance status on male and female investigation of the**
295 **divider window in both cohorts.**

Effect	df	Male investigation in Unfamiliar odor cohort		Female investigation in Unfamiliar odor cohort		df	Male Famil
		F	p	F	p		F
Time Bin	2	3.965	0.037*	3.841	0.041*	2	4.391

Odor Condition	1	0.958	0.353	5.437	0.045*	1	0.034
Dominance Status	1	3.315	0.102	0.549	0.478	1	0.665
Bin x Odor Condition	2	0.421	0.663	0.254	0.779	2	0.93
Bin x Dominance Status	2	0.732	0.495	4.109	0.034*	2	2.8
Odor Condition x Dominance Status	1	1.962	0.195	0.059	0.814	1	0.062
Odor Condition x Bin x Dominance Status	2	0.585	0.567	0.493	0.619	2	0.063

296 * p value is not statistically significant after correcting for multiple comparisons.

Additional male odor during limited social interactions decreases baseline vocal output depending on the familiarity of male odor

In order to assess how contextual manipulations affected baseline vocal behavior, vocal behavior

during BBV playback, and vocal behavior after playback ended, we employed a fifteen minute

interaction in which the first five minutes consisted of silence, followed by five minutes of BBV

playback, followed by a final five minutes of silence [35]. The addition of unfamiliar male odor

during the interaction (Fem+Male condition) caused significantly lower baseline levels of USVs

compared to when only female odor was present (Fem condition) (Two-way mixed ANOVA: $F_{1,10}$

= 10.005, p = 0.01). The addition of familiar male odor however produced no significant difference

in baseline USV levels compared to when only female odor was present (Two-way mixed ANOVA:

$F_{1,14}$ = 0.084, p = 0.776) (Fig 4A) (Table 2).

Figure 4: Properties of vocal repertoires. (A) Baseline USVs produced in the Fem condition vs the Fem+Male condition. (B) Average duration of baseline USVs in seconds. (C) Baseline proportion of harmonic USVs (# baseline harmonics/# baseline USVs). Proportion of baseline harmonic USVs was calculated as baseline harmonic USVs /(baseline harmonic USVs + baseline nonharmonic USVs). Blues represent the unfamiliar odor cohort, reds represent the familiar odor cohort. Averages from dominants and subordinates are pooled due to the lack of effect of social status. Individual points of data noted with dominants as X's and subordinates as circles.

Because USVs can vary widely in duration [1], we measured the duration of all baseline

USVs (both harmonic and nonharmonic). Males produced significantly longer baseline USVs in

the Fem condition compared to the Fem+Male condition in unfamiliar male cohort (Two-way

mixed ANOVA: $F_{1,10}$ = 5.285, p = 0.044), though this result was not significant after correcting for

multiple comparisons (Table 2). No effects of odor condition on duration of baseline USVs were

observed in the familiar male odor cohort (Two-way mixed ANOVA: $F_{1,14}$ = 0.005, p = 0.945) (Fig

4B).

Additional male odor during limited social interactions decreases

baseline usage of 50 kHz harmonic USVs depending on the familiarity

of male odor

50 kHz harmonic USVs are calls highly correlated to mounting behavior in male mice [11]. In this

study, harmonic USVs made up a small proportion of all calls; on average, harmonic USVs made

up 7.31% (SD = 0.0701) of all USVs in the interaction. In order to assess whether the frequency

of harmonic USV usage differed across conditions, for each individual we evaluated the

proportions of harmonic USVs produced in each time bin out of the total number of all USVs in

that time bin. The addition of unfamiliar male odor during the interaction (Fem+Male condition)

caused mice to use a significantly lower proportion of 50 kHz harmonic USVs during the baseline

period compared to when only female odor was present (Fem condition) (Binary logistic

regression: $F_{1,10}$ = 10.752, p = 0.004), though this effect was not present in the familiar male odor

cohort (Binary logistic regression: $F_{1,14}$ = 0.035 p = 0.853) (Fig 4C) (Table 2).

Male mice decrease USV production in response to BBV playback across

all conditions

In both the unfamiliar odor cohort and the familiar odor cohort, male mice reduced their calls

during playback of BBVs, and then increased calling rates after playback ceased in both Fem and

Fem+Male trials (Three-way mixed ANOVAs: Significant main effect of bin $F_{2,20}$ = 50.538, p < 0.001

and $F_{2,28}$ = 34.635, p < 0.001 for the unfamiliar and familiar cohorts, respectively). This

subsequent increase in calling after the playback period did not fully recover to levels similar to

baseline, however, but rather increased to a level between that observed in the baseline and the

recovery. In the unfamiliar odor cohort, individuals produced more USVs in the Fem odor

condition compared to the Fem+Male odor condition (Significant main effect of odor condition:

$F_{1,10}$ = 12.026, p = 0.006) (Fig 5A), though this effect was not present in the familiar odor cohort

($F_{1,14}$ = 0.454, p = 0.511) (Fig 5B). No interaction effects between odor and bin were present in

either the unfamiliar odor cohort or the familiar odor cohort ($F_{2,20}$ = 0.223, p = 0.802, and $F_{2,30}$ =

0.734, p = 0.489) (Table 1).

Figure 5: Number of USVs produced across time bins (A) in the unfamiliar odor cohort, and (B) in the familiar odor cohort. Average USV production of dominants is represented by the solid lines, and average USV production of subordinates is represented by dashed lines. Lighter lines denote average USV production of individuals in the Fem condition, while darker lines denote average USV production of individuals in the Fem+Male condition.

345 Because there was no interaction effect between time bin and any other factor in these

346 models, our data showed that differences across odor conditions and familiarity cohorts can be

347 attributed to differences in baseline levels of USV production, rather than differences in vocal

348 response to BBV playback.

Male mice increase usage of 50 kHz harmonics after BBV playback ends

when exposed to unfamiliar male odor

In the unfamiliar male odor cohort, the proportion of 50 kHz harmonic calls used relative to the

total number of calls produced did not change in response to BBV playback, however after

playback ended, males produced a greater proportion of harmonic calls in the recovery period

(Binary logistic regression: Significant main effect of bin $F_{2,20}$ = 4.925, p = 0.011). However,

posthoc pairwise comparisons revealed this increase in harmonic usage only occurred in trials

with unfamiliar male odor added (t_{55} = 2.344, p = 0.023) (Fig 6A). Overall, in the unfamiliar male

odor cohort, males produced a greater proportion of harmonics in Fem trials compared to

Fem+Male trials (Binary logistic regression: Significant main effect of odor condition $F_{1,10}$ = 8.429,

p = 0.005). In the familiar male odor cohort, no significant effects of any factor were observed

(Table 1) (Fig 6B).

Figure 6: Proportion of harmonic USVs across time bins out of total USVs (A) in the unfamiliar odor cohort and (B) in the familiar odor cohort. Average harmonic USV production of dominants is represented by the solid lines, and average harmonic USV production of subordinates is represented by dashed lines. Lighter lines denote average harmonic USV production of individuals in the Fem condition, while darker lines denote average harmonic USV production of individuals in the Fem+Male condition. Proportion of harmonic USVs for each time bin was calculated as harmonic USVs in time bin/(harmonic USVs + nonharmonic USVs in time bin). The significance bar in 5A) showing increased usage of proportion of harmonic USVs in the recovery period compared to the baseline and playback period is dark blue, representing that it only applies to the Fem+Male conditions, not the Fem conditions.

Male and female investigative behavior does not significantly differ

across conditions, cohorts, nor time bins

In the unfamiliar odor cohort, male investigation of the divider window appeared to slightly

decrease during BBV playback, and investigative behavior did not recover after BBV playback,

with no effects of dominance status or odor condition (Three-way mixed ANOVA: $F_{2,18}$ = 3.965, p

= 0.037) (Fig 7A). In contrast, in the familiar odor cohort, male investigation of the divider window

appeared to slightly decrease during BBV playback and subsequently recover after playback, with

no effects of dominance status or odor condition (Three-way mixed ANOVA: $F_{2,28}$ = 4.391, p =

0.022) (Fig 7C). However, neither of these results were significant after correcting for multiple

comparisons.

> **Figure 7: Duration of time the focal male (A and C) or the stimulus female (B and D) spent investigating the divider window across time bins in the unfamiliar odor cohort (A and B) and in the familiar odor cohort (C and D).** Average duration of investigation in trials with dominant males is represented by the solid lines, and average duration investigation in trials with subordinates is represented by dashed lines. Lighter lines denote average investigation during trials in the Fem condition, while darker lines denote average investigation in the Fem+Male condition.

Stimulus females paired with males in the unfamiliar odor cohort tended to investigate

the divider window more during the Fem+Male condition trials ($F_{1,9}$ = 5.437, p = 0.045), and this

investigation actually increased during BBV playback, and decreased back to baseline levels after

playback ended ($F_{2,18}$ = 3.841, p = 0.041) (Fig 7B). This effect of playback was dependent on

dominance status in that female investigative behavior was only affected by playback when

females were paired with subordinate males, and there was no effect when they were paired

with dominant males ($F_{2,18}$ = 4.109, p = 0.034). When stimulus females were paired with males in

the familiar odor cohort, they decreased their investigative behavior during BBV playback, and

this behavior increased back to baseline levels when playback ended ($F_{2,28}$ = 3.949, p = 0.031) (Fig

7D), with no effect of odor condition or dominance status. However, again, none of these results

were significant after correcting for multiple comparisons.

385   Additional male odor alters the relationship between male and female

386   investigative behavior

In the Fem condition trials, the total duration of male investigative behavior did not correlate to

the total duration of female investigation behaviors in either the unfamiliar odor cohort or the

familiar odor cohort (Pearson correlation: r = -0.373, p = 0.259; r = 0.235, p = 0.381 respectively)

(Fig 8A-B). In the Fem+Male condition trials, the total duration of male investigative behaviors

negatively correlated to the total duration of female investigative behaviors for those in the

unfamiliar odor cohort (Pearson correlation r = -0.625, p = 0.04) (Fig 8C), though this effect was

not significant after correcting for multiple comparisons. In contrast, in the Fem+Male odor

condition trials conducted with the familiar odor cohort, the total duration of male investigation

behaviors positively correlated to the total duration of female investigative behaviors (Pearson

correlation r = 0.839 p < 0.001) (Fig 8D), and this effect remained significant after correcting for

multiple comparisons.

Figure 8: Correlation between female investigation and male investigation behavior in the (A) unfamiliar odor cohort: Fem condition, (B) familiar odor cohort: Fem condition, (C) unfamiliar odor cohort: Fem+Male condition, and (D) familiar odor cohort: Fem+Male condition. Dominant individuals denoted as X's and subordinates as circles.

399   Discussion

Behavioral decisions during courtship are highly influenced by the interplay between

environmental cues, internal factors, and signals from communicating conspecifics [4]. Thus, by

evaluating how individuals respond to signals in various contexts, we can begin to decipher how

much weight a given factor may have on such decisions. In this study, when interactions included

odors cues from unfamiliar, but not familiar, male conspecifics, male mice exhibited reduced

courtship motivation compared to interactions when only female odors were present. This

decrease in courtship motivation was evidenced by shifts in multiple aspects of male courtship

vocalizations: decreases in call production, call durations, and variety of calling repertoires. Male

mice thus may be evaluating their environment for potential competitors, discerning the identity

of these competitors, and scaling their courtship activities in response to these cues. The

interplay between signals not only from courtship partners but also from competitors as well as

social recognition of those signals is instrumental in motivating behavioral decisions during

mating interactions.

Unfamiliar male odor during courtship decreases male courtship

motivation

Vocal signaling can often reflect motivational state across taxa [57–62]. USV production is a

well-established proxy for courtship motivation in male mice, as these calls function to

encourage female sexual receptivity and mating [18,21,22,24,63,64]. Furthermore, out of the

possible measurements for courtship motivation in mice, these signals are particularly

important as they can convey information about male fitness, identity, and reproductive state,

which can in turn influence female mate choice [22,26,65].Changes in courtship motivation can

be spurred by exposure to social cues [58,59]. Olfactory cues can perform a variety of functions

during social interactions for both males and females. Because these cues are so significant in

defining the context of an interaction, particularly opposite-sex interactions, individuals may

plastically adjust their behavior in response to cues like these odors, potentially to avoid

expending unnecessary time and energy engaging with individuals that are not sexually

receptive [66–72]. In this study, we manipulated the context of courtship interactions by

including odor cues from additional males not present during the interaction in order to assess

how the focal males would behave compared to when no additional male odor cues are

present. While many courtship assays used in laboratory mice are conducted with a single male

and a single female [73], because of the propensity of mice to live in a wide range of social

densities, it is likely that males in natural settings are often confronted with the odors of other

potentially competing males [74,75]. Thus, investigating how these odor cues that simulate a

potential competitive audience may affect how mice produce and respond to signals gives

important insight to how behavioral decisions are made during courtship, and what information

is most impactful to individuals as they choose how to navigate social interactions. It is well-

established that birdsong exhibits signal flexibility in response to social context [76]. For

example, male cowbirds will modulate the attractiveness of their songs in response to the

presence of competitors, though how they modulate their songs may depend on factors like

whether or not the competitors can easily access the focal individual and if the individual can

gauge the motivational state of the competitor [77,78].

As mice can use chemical components within urine to discriminate between individual

identity, familiarity, and dominance [79–82], odor cues can be used to simulate different

audiences, and investigate how different identity characteristics alter behavior during sexual

interactions. We found that whether males responded to this odor was dependent on whether

or not the focal male had social contact with the odor's source. When confronted with unfamiliar

male odor during the interaction, males produced fewer baseline courtship calls (Fig 4A), the calls

produced during the interaction prior to the playback of BBVs, and the average length of these

calls showed a trend in that they shorter than when they interacted with a female with no

additional male odor present (Fig 4B). These responses may indicate a decrease in courtship

motivation and effort. Not only is number of calls produced indicative of motivational state, but

types of calls produced can also reveal information about motivation. Across taxa, increasing

vocal production through the expression of complex syllables and longer calls is associated with

greater energetic costs [22,42,64,83–85]. Longer duration of vocalizations have also been

associated with increased courtship display effort in singing mice, and female rodents tend to

prefer males that produce longer "higher-effort" songs [64,86,87]. Thus, male mice increasing

the usage of more complex and longer-duration elements in songs, like the 50 kHz harmonics

assessed in this study, may reflect increased mating effort.

When males were confronted with unfamiliar male odor during the courtship interaction,

they also shifted the composition of the different types of syllables they used, their vocal

repertoire [29], to produce a lower proportion of 50 kHz harmonic calls during the baseline period

when unfamiliar male odor is present (Fig 4C). These calls are categorized by exhibiting a

harmonic frequency band at 50 kHz, and are longer and more structurally complex than USVs

that do not contain this band (Fig 1). Importantly, these calls are positively correlated with bouts

of mounting [10,11]. That males specifically decrease the use of these calls in their vocal

repertoire when unfamiliar competitor odors are present reflects the intentional function of

these calls to promote mating, and that these competitive odors are significantly affecting

numerous aspects of the male courtship vocal repertoire. Calling less and producing less complex

and shorter calls thus could indicate decreased reproductive motivation.

It may be logical for individuals to show decreased courtship motivation, through the

reduction of and simplification of calling behaviors, as cues that indicate the presence of potential

reproductive competitors can decrease sexual motivation and induce rapid behavioral plasticity

[49,88]. This is supported by the theory of sperm competition and its subsequent behavioral

consequences, in that males may "budget" copulatory efforts depending on factors like risk of

competition [89–95]. Across taxa, male precopulatory courtship behaviors can be reduced as the

density of competitors increase [89,93]. This could be due to adaptive changes in reproductive

strategies by the males, or sexual interference from the competitors [89,93]. Our results are

consistent with this interpretation. This propensity to quickly adjust one's behavior in response

to the socio-sexual environment is especially relevant to house mice, as in nature they can exist

in a wide range of population densities, and are able to be successful at both very high and low

densities [96–99].

Importantly, the specific presence of male odors during sexual interactions can also

influence female mate preferences [100–102] and impact competition tactics of males across

taxa [103,104]. Thus, it could be important for male mice to recognize when other male odors

are present, and "hedge their bets" when it comes to interacting with a female that has multiple

mate options, as to not invest heavily in courting a female that may prefer another male, or may

have even already mated with another male [89,90,103,105].

A second cohort of individuals were exposed to the odor of their social hierarchy partner

during the limited courtship interaction, and interestingly, when the odor was familiar to the

focal individual, their vocal behavior did not differ from when no odor was present at all. We

predicted that familiar male competitor odor would emphasize the vocal response differences in

dominant and subordinate males, as dominants were confronted with the smell of their

subordinate during the interaction, and subordinates with their dominant. Not only was there no

difference in dominant and subordinate individuals (to be discussed in more detail in subsequent

sections), but number of calls, duration of calls, and repertoire of calls were not affected by the

familiar competitor odor at all. This important distinction between the responses of individuals

to familiar compared to unfamiliar competitor odors could have occurred due to various reasons.

Mice can identify individuals they are familiar with through chemical cues in urine [79]. After

frequent exposure to odor cues from conspecifics, mice can habituate to these odors, and thus

have a less pronounced response to them [103,106,107]. The dear enemy hypothesis predicts

that familiar competitors should elicit less of a reaction from territory owners than unfamiliar

competitors, due to recognition and knowledge of the benefits and costs of defending against

those particular individuals [108–110]. Our results are consistent with the dear enemy hypothesis

in that the cues from familiar competitors elicit less of a response (less of a decrease in calls) than

cues from unfamiliar competitors. The familiarity of the male cues present could determine

whether or not males pay more attention to the male cues, or ignore them and shift focus to the

female. A potential way for future studies to explore if, in this context, the decrease in vocal

behavior in response to unfamiliar competitor odor is paired with an increase in behaviors aimed

to signal to the competitor could be to assess competitive scent marking behavior of the focal

males when confronted with familiar vs unfamiliar odor [106,111,112]. If males confronted with

unfamiliar odor are shifting their efforts from courting the females to focusing on competing with

potential competitors, we would expect to see an increase in marking behavior is correlated with

a decrease in vocal behavior.

The relationship between male and female investigative behavior changes direction depending on familiarity of the competitor odor

Any apparent small changes in male and female investigation of the divider window, through

which limited access of the other individual was possible, were not statistically significant after

correction for multiple comparisons (Fig 7; Table 3). However, unexpectedly, when investigating

how male and female investigative behavior related to one another across the different odor

conditions and familiarity cohorts, we found that when trials included odor from familiar males,

there was a positive correlation between male and female investigative behavior (Fig 8D), and

we found no correlation between male and female investigative behavior in any other conditions

(Fig 8A-C). It is possible that the Fem+Male trials having odor from two different males available

to the females would prompt more investigation from the female, though this was not the case

(Fig 7B, 7D). Overall, our data revealed that the relationship between male and female

investigative behavior in the presence of varying social odor conditions is not straightforward as

one might expect, and worthy of future investigation.

BBVs spur decreased courtship effort in male mice

All experimental groups in this study decreased calling in response to playback of broadband

vocalizations, BBVs, and no other factors measured affected this response (Fig 5). While the

function of BBVs across contexts has been largely unexplored, evidence suggests that in certain

courtship contexts, BBVs serve to communicate a negative affective state of the signaler

[26,34,113,114]. Surprisingly, we found that when individuals were exposed to the odor of an

unfamiliar competitor, the proportion of 50 kHz harmonic USVs they used increased in the

recovery period after exposure to BBV playback, and it was in this period that they used the

highest proportion of harmonics in their calling, even more so than in the baseline period (Fig

6A). More in depth research will have to explore the intricacies of why the discontinuation of BBV

playback might spur a rapid increase in harmonic usage only when cues from an unfamiliar

potential competitor are present, though this could potentially reflect habituation to the

competitor cues as the trial goes on. This result, though difficult to interpret, does however

support that the cessation of BBV playback changes the trajectory of syllable usage in male mice

when cues indicating the presence of an unfamiliar competitor are present.

BBVs may not always produce a negative reproductive response, and are sometimes

correlated with increases in mounting and calling [11,114,115]. It is possible that the potential

different functions of BBVs manifest when BBVs are paired with other nonvocal behaviors, and

the isolated vocal characteristics of the BBVs may indicate a negative valence, and thus promote

decreased courtship effort in receivers without access to additional cues from the female.

Our results exemplify that these signals are significant and attention-grabbing to male

mice, seemingly regardless of certain identity and environmental factors. BBVs and their

correlated behavioral responses remain a largely untapped line of research; and very few studies

have explored the effects of BBV playback isolated from the associated physical rejection

behaviors often correlated with BBV production. Greater research devoted to this area of mouse

communication could help elucidate how different internal and external factors may affect

responses to these signals.

Lack of an effect of dominance status

In this study, dominance status did not significantly affect any vocal or nonvocal behavioral

measurements in either the unfamiliar odor cohort or the familiar odor cohort. This was not what

we predicted, as it is generally accepted that dominant male mice produce more USVs than

subordinate individuals in response to a novel female [116,117], though USVs are not a standard

way to measure dominance as they do not directly measure access to mates, but rather reflect

mating effort [22]. Because USV production is highly dependent on other contextual factors, the

relationship between USV production and dominance is likely influenced by finetuned

differences in social history and situational context rather than being simply and strictly linear

[118]. For example, certain studies have found the relationship between dominance and

increased USV production depends on the types of cues present and the recency of social defeat,

and can even reverse [7,19,36]. Interestingly, in this study, the individual with the highest number

of baseline calls in all conditions (both odor cohorts in both the Fem and Fem+Male conditions)

was subordinate (Fig 4A), though there was no overall statistically significant effect of dominance

on any measurements. Despite observations that this strain of mice may not exhibit overt severe

dominance their hierarchies [119], all housed pairs in this experiment showed consistent and

reliable dyadic hierarchy relationships through multiple rounds of the tube test. Thus, depending

on the context of an interaction, dominance may affect behaviors such as offensiveness and

aggression during tube tests, while not manifesting in indirect courtship assays like that used in

this study.

Dominant and subordinate males also tend to differ in their response to social challenges

[120–122]. Dominant rodents tend to be more susceptible to negative health consequences and

behavioral impairment following social challenges compared to subordinate males

[46,47,123,124], however this is most often exhibited during direct interactions with social

competitors. In this study, dominant males and subordinate males were not differentially

affected by odors of competitor males during courtship, thus odors alone may not be sufficient

to induce this hierarchy effect. Salience of cues, interaction context, social history, and genetic

factors may all contribute to how the effects of dominance manifest during courtship

[116,122,125–127], and more research is needed to tease out which of these factors are most

influential on determining the effects of dominance.

Conclusions

By looking at how these multiple factors intertwine to affect behavioral decisions during

courtship, we can begin to assess what "matters" to a mouse as they navigate mating and when

and where to exert effort during courtship. This study supports the potential negative valence of

a relatively ill-explored female signal: broadband vocalizations, or squeaks, in certain contexts.

We determined that while males robustly adjust their behavior in response to these calls, the

degree of this behavioral response was not altered by any of the internal or external factors we

manipulated. Furthermore, this study posits that cues indicating the presence, or "audience", of

potential competitors can alter male mouse courtship motivation and behavior, but this effect is

dependent on social history with the audience members. Furthermore, our results support that

certain identity factors, like dominance status, do not affect the relationship between these cues

and the observed behavioral response.

Acknowledgements

603 The authors would like to thank and acknowledge Dr. Kayleigh E. Hood, Ph.D. for providing

604 mentorship, and guidance regarding the split cage assay, and Morgan Leever, M.P.A., M.S. for

605 consulting regarding statistical analysis.

References

1. Holy TE, Guo Z. Ultrasonic songs of male mice. PLoS Biology. 2005;3(12):1–10.

2. Seyfarth RM, Cheney DL. Signalers and Receivers in Animal Communication. Annual
Review of Psychology. 2003;54:145–73.

3. Wiley RH. Signal Detection and Animal Communication. Advances in the Study of Behavior.
2006;36(06):217–47.

4. Hurley LM, Kalcounis-Rueppell MC. State and Context in Vocal Communication of Rodents.
In: Rodent Bioacoustics. 2018. p. 191–221.

5. Leger DW. Contextual sources of information and responses to animal communication
signals. Psychological Bulletin. 1993;113(2):295–304.

6. Palmer CR, Kristan WB. Contextual modulation of behavioral choice. Current Opinion in
Neurobiology. 2011 Aug 1;21(4):520–6.

7.   D'Amato FR. Male-male interference in a reproductive context: Effect of social status in
mice. Ethology Ecology and Evolution. 1991;3(4):295–306.

8.   Todt D, Naguib M. Vocal Interactions in Birds: The Use of Song as a Model in
Communication. Advances in the Study of Behavior. 2000;29(C):247–96.

9.   Chabout J, Sarkar A, Dunson DB, Jarvis ED. Male mice song syntax depends on social
contexts and influences female preferences. Frontiers in Behavioral Neuroscience.
2015;9(APR):1–16.

10.   Hanson JL, Hurley LM. Female presence and estrous state influence mouse ultrasonic
courtship vocalizations. PLoS ONE. 2012;7(7).

11.   Finton CJ, Keesom SM, Hood KE, Hurley LM. What's in a squeak? Female vocal signals
predict the sexual behaviour of male house mice during courtship. Animal Behaviour.
2017;126:163–75.

12.   Brumm H, Naguib M. Chapter 1 Environmental Acoustics and the Evolution of Bird Song.
In: Advances in the Study of Behavior [Internet]. Elsevier; 2009 [cited 2024 Sep 30]. p. 1–
33. Available from: https://linkinghub.elsevier.com/retrieve/pii/S0065345409400019

13.   Mol C, Chen A, Kager RWJ, Ter Haar SM. Prosody in birdsong: A review and perspective.
Neuroscience & Biobehavioral Reviews. 2017 Oct;81:167–80.

14.   Opaev AS. The Communicative Value of Complex Singing in Passerine Birds. Biol Bull Russ
Acad Sci. 2022 Dec;49(10):1750–68.

15.   Riters LV, Stevenson SA. Using seasonality and birdsong to understand mechanisms
underlying context-appropriate shifts in social motivation and reward. Hormones and
Behavior. 2022 Jun;142:105156.

16.   Ritschard M, Brumm H. Zebra finch song reflects current food availability. Evol Ecol. 2012
Jul;26(4):801–12.

17.   Fischer J, Hammerschmidt K. Ultrasonic vocalizations in mouse models for speech and
socio-cognitive disorders: Insights into the evolution of vocal communication. Genes,
Brain and Behavior. 2011;10(1):17–27.

18.   Granon S, Faure A, Chauveau F, Cressant A, Ey E. Why Should My Mouse Call Me? Acoustic
Communication in Mouse Models of Social Disorders: Ultrasonic Vocalizations as an
Index of Emotional and Motivational States. Handbook of Behavioral Neuroscience.
2018;25:423–31.

19.   Lumley LA, Sipos ML, Charles RC, Charles RF, Meyerhoff JL. Social stress effects on
territorial marking and ultrasonic vocalizations in mice. Physiology and Behavior.
1999;67(5):769–75.

20.   Sangiamo DT, Warren MR, Neunuebel JP. Ultrasonic signals associated with different types
654   of social behavior of mice. Nature Neuroscience. 2020;23(3):411–22.

655   21.   D'amato FR. Courtship ultrasonic vocalizations and social status in mice. Animal Behaviour.
656   1991;41(5).

657   22.   Egnor SER, Seagraves KM. The contribution of ultrasonic vocalizations to mouse courtship.
658   Current Opinion in Neurobiology. 2016;38(Figure 1):1–5.

659   23.   Ferhat AT, Torquet N, Le Sourd AM, De Chaumont F, Olivo-Marin JC, Faure P, et al.
660   Recording mouse ultrasonic vocalizations to evaluate social communication. Journal of
661   Visualized Experiments. 2016;2016(112):1–12.

662   24.   Portfors CV, Perkel DJ. The role of ultrasonic vocalizations in mouse communication.
663   Current Opinion in Neurobiology. 2014;28:115–20.

664   25.   Neunuebel JP, Taylor AL, Arthur BJ, Roian Egnor SE. Female mice ultrasonically interact
665   with males during courtship displays. eLife. 2015;4(MAY):1–24.

666   26.   Sugimoto H, Okabe S, Kato M, Koshida N, Shiroishi T, Mogi K, et al. A role for strain
667   differences in waveforms of ultrasonic vocalizations during Male-Female interaction.
668   PLoS ONE. 2011;6(7).

669   27.   White NR, Prasad M, Barfield RJ, Nyby JG. 40- and 70-kHz vocalizations of mice (*Mus
670   musculus*) during copulation. Physiology and Behavior. 1998;63(4):467–73.

671   28.   Scattoni ML, Gandhy SU, Ricceri L, Crawley JN. Unusual repertoire of vocalizations in the
672   BTBR T+tf/J mouse model of autism. PLoS ONE. 2008;3(8).

673   29.   Heckman J, McGuinness B, Celikel T, Englitz B. Determinants of the mouse ultrasonic vocal
674   structure and repertoire. Neuroscience and Biobehavioral Reviews. 2016;65.

675   30.   Grimsley JMS, Monaghan JJM, Wenstrup JJ. Development of social vocalizations in mice.
676   PLoS ONE. 2011;6(3).

677   31.   Ruat J, Genewsky AJ, Heinz DE, Kaltwasser SF, Canteras NS, Czisch M, et al. Why do mice
678   squeak? Toward a better understanding of defensive vocalization. iScience.
679   2022;25(7):104657.

680   32.   Knights CB, Gentry C, Bevan S. Partial medial meniscectomy produces osteoarthritis pain-
681   related behaviour in female C57BL/6 mice. Pain. 2012;153(2):281–92.

682   33.   Anderson LL, Everett-Morgan D, Petkova SP, Silverman JL, Arnold JC. Ictal vocalizations in
683   the Scn1a+/− mouse model of Dravet syndrome. Epilepsia Open. 2023;8(3):776–84.

684   34.   Keesom SM, Hurley LM. Socially induced serotonergic fluctuations in the male auditory
685   midbrain correlate with female behavior during courtship. Journal of Neurophysiology.
686   2016;115(4):1786–96.

687   35.   Hood KE, Long E, Navarro E, Hurley LM. Playback of broadband vocalizations of female
688   mice suppresses male ultrasonic calls. PloS one. 2023;18(1):e0273742.

689   36.   Nyby JG, Dizinno GA, Whitney G. Social status and ultrasonic vocalizations of male mice.
690   Behavioral Biology. 1976;18(2):285–9.

691   37.   Marler P, Dufty A, Pickert R. Vocal communication in the domestic chicken: I. Does a
692   sender communicate information about the quality of a food referent to a receiver?
693   Animal Behaviour. 1986;34(PART 1):188–93.

694   38.   Baltz AP, Clark AB. Extra-pair courtship behaviour of male budgerigars and the effect of an
695   audience. Animal Behaviour. 1997 May;53(5):1017–24.

696   39.   Evans CS, Marler P. Food calling and audience effects in male chickens, *Gallus gallus*: their
697   relationships to food availability, courtship and social facilitation. Animal Behaviour. 1994
698   May;47(5):1159–70.

699   40.   Sánchez NV, Mennill DJ. Behavioural consequences of conspecific neighbours: a systematic
700   literature review of the effects of local density on avian vocal communication. J Ornithol.
701   2024 Oct;165(4):847–59.

702   41.   Vignal C, Mathevon N, Mottin S. Audience drives male songbird response to partner's
703   voice. Nature. 2004;430(6998):448–51.

704   42.   Seagraves KM, Arthur BJ, Egnor SER. Evidence for an audience effect in mice: Male social
705   partners alter the male vocal response to female cues. Journal of Experimental Biology.
706   2016;219(10):1437–48.

707   43.   Plath M, Schlupp I. Misleading Mollies. Communicative & Integrative Biology.
708   2008;1(2):199–203.

709   44.   Bierbach D, Girndt A, Hamfler S, Klein M, Mücksch F, Penshorn M, et al. Male fish use prior
710   knowledge about rivals to adjust their mate choice. Biology Letters. 2011;7(3):349–51.

711   45.   Castellano S, Friard O, Pilastro A. The audience effect and the role of deception in the
712   expression of male mating preferences. Animal Behaviour. 2016;115:273–82.

713   46.   D'Amato FR, Rizzi R, Moles A. A model of social stress in dominant mice: Effects on
714   sociosexual behaviour. Physiology and Behavior. 2001;73(3):421–6.

47. Larrieu T, Cherix A, Duque A, Rodrigues J, Lei H, Gruetter R, et al. Hierarchical Status
Predicts Behavioral Vulnerability and Nucleus Accumbens Metabolic Profile Following
Chronic Social Defeat Stress. Current Biology. 2017;27(14):2202-2210.e4.

48. Chaouloff F. Serotonin, stress and corticoids. Journal of Psychopharmacology.
2000;14(2):139–51.

49. Fernández-Vargas M. Presence of a potential competitor and its individual identity
modulate ultrasonic vocalizations in male hamsters. Animal Behaviour. 2018;145.

50. Fan Z, Zhu H, Zhou T, Wang S, Wu Y, Hu H. Using the tube test to measure social hierarchy
in mice. Nature Protocols. 2019;14(3):819–31.

51. Friard O, Gamba M. BORIS: a free, versatile open-source event-logging software for
video/audio coding and live observations. Methods in Ecology and Evolution.
2016;7(11):1325–30.

52. Warton DI, Hui FKC. The arcsine is asinine: The analysis of proportions in ecology. Ecology.
2011;92(1).

53. Feder PI, Aume LL, Triplett CA, Simmons JE, Narotsky MG. Analysis of proportional data in
reproductive and developmental toxicity studies: Comparison of sensitivities of logit
transformation, arcsine square root transformation, and nonparametric analysis. Birth
Defects Research. 2020;112(16):1260–72.

54. Pituch KA, Stevens JP. Binary Logistic Regression. Applied Multivariate Statistics for the
Social Sciences. 2020;454–90.

55. Games PA, Winkler HB, Probert DA. Robust Tests for Homogeneity of Variance.
Educational and Psychological Measurement. 1972;32(4).

56. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. Journal of the Royal Statistical Society: Series B
(Methodological). 1995;57(1).

57. Hafez B, Hafez ESE. Reproductive Behavior. In: Hafez B, Hafez ESE, editors. Reproduction in
Farm Animals [Internet]. 1st ed. Wiley; 2000 [cited 2024 Oct 3]. p. 291–306. Available
from: https://onlinelibrary.wiley.com/doi/10.1002/9781119265306.ch19

58. Tachibana SI, Touhara K, Ejima A. Modification of Male Courtship Motivation by Olfactory
Habituation via the GABAA Receptor in *Drosophila melanogaster*. Marion-Poll F, editor.
PLoS ONE. 2015 Aug 7;10(8):e0135186.

59. Ishii KK, Touhara K. Neural circuits regulating sexual behaviors via the olfactory system in
mice. Neuroscience Research. 2019 Mar;140:59–76.

60. Iwasaki M, Poulsen TM, Oka K, Hessler NA. Singing-Related Activity in Anterior Forebrain
of Male Zebra Finches Reflects Courtship Motivation for Target Females. Coleman MJ,
editor. PLoS ONE. 2013 Nov 29;8(11):e81725.

61. Okanoya K, Screven LA. Rodent Vocalizations: Adaptations to Physical, Social, and Sexual
Factors. In: Dent ML, Fay RR, Popper AN, editors. Rodent Bioacoustics [Internet]. Cham:
Springer International Publishing; 2018 [cited 2024 Oct 3]. p. 13–41. (Springer Handbook
of Auditory Research; vol. 67). Available from: http://link.springer.com/10.1007/978-3-
319-92495-3_2

62. Ventura-Aquino E, Portillo W, Paredes RG. Sexual Motivation: A Comparative Approach in
Vertebrate Species. Curr Sex Health Rep. 2018 Sep;10(3):114–23.

63. Nunez AA, Nyby JG, Whitney G. The effects of testosterone, estradiol, and
dihydrotestosterone on male mouse (*Mus musculus*) ultrasonic vocalizations. Hormones
and Behavior. 1978;11(3).

64. Fernández-Vargas M. Vocal Signals of Sexual Motivation in Male and Female Rodents.
Current Sexual Health Reports. 2018;10(4).

65. Nunez AA, Tan DT. Courtship ultrasonic vocalizations in male Swiss-Webster mice: Effects
of hormones and sexual experience. Physiology and Behavior. 1984;32(5).

66. Dore AA, McDowall L, Rouse J, Bretman A, Gage MJG, Chapman T. The role of complex
cues in social and reproductive plasticity. Behavioral Ecology and Sociobiology.
2018;72(8).

67. Johnston RE. Chemical Signals and Reproductive Behavior. In: Pheromones and
Reproduction in Mammals. 1983.

68. Guevara-Fiore P, Endler JA. Female receptivity affects subsequent mating effort and mate
choice in male guppies. Animal Behaviour. 2018;140.

69. Jenikejew J, Wauters J, Dehnhard M, Scheumann M. The female effect - How female
receptivity influences faecal testosterone metabolite levels, socio-positive behaviour and
vocalization in male Southern white rhinoceroses. Conservation Physiology. 2021;9(1).

70. Barrass R. A Quantitative Study of the Behaviour of the Male *Mormoniella Vitripennis*
(Walker) (Hymenoptera, Pteromalidae) Towards Two Constant Stimulus-Situations.
Behaviour. 1961;18(4).

71. Girard MB, Elias DO, Kasumovic MM. Female preference for multi-modal courtship:
Multiple signals are important for male mating success in peacock spiders. Proceedings of
the Royal Society B: Biological Sciences. 2015;282(1820).

72. Linley JR, Hinds MJ. Quantity of the male ejaculate influenced by female unreceptivity in
the fly, *Culicoides melleus*. Journal of Insect Physiology. 1975;21(2).

73. Kondrakiewicz K, Kostecki M, Szadzińska W, Knapska E. Ecological validity of social
interaction tests in rats and mice. Genes Brain and Behavior. 2019 Jan;18(1):e12525.

74. Chambers LK, Singleton GR, Krebs CJ. Movements and social organization of wild house
mice (*Mus domesticus*) in the wheatlands of northwestern Victoria, Australia. Journal of
Mammalogy. 2000;81(1).

75. Brown RZ. Social Behavior, Reproduction, and Population Changes in the House Mouse
(*Mus musculus L.*). Ecological Monographs. 1953;23(3).

76. Hausberger M, Henry L, Testé B, Barbu S. Contextual Sensitivity and Bird Song: A Basis for
Social Life. In: Oller DK, Griebel U, editors. Evolution of Communicative Flexibility
[Internet]. The MIT Press; 2008 [cited 2024 Oct 2]. p. 121–38. Available from:
https://direct.mit.edu/books/book/2702/chapter/73061/Contextual-Sensitivity-and-Bird-
Song-A-Basis-for

77. Gersick AS, White DJ. Male cowbirds vary the attractiveness of courtship songs with
changes in the social context. Behav. 2018;155(1):1–25.

78. West MJ, King AP, Eastzer DH. Validating the female bioassay of cowbird song: Relating
differences in song potency to mating success. Animal Behaviour. 1981 May;29(2):490–
501.

79. Brennan PA. The nose knows who's who: Chemosensory individuality and mate
recognition in mice. Hormones and Behavior. 2004;46(3).

80. Penn D, Potts W. How do major histocompatibility complex genes influence odor and
mating preferences? Advances in Immunology. 1998;69.

81. Veyrac A, Wang G, Baum MJ, Bakker J. The main and accessory olfactory systems of female
mice are activated differentially by dominant versus subordinate male urinary odors.
Brain Research. 2011;1402.

82. Drickamer LC. Responses to odors of dominant and subordinate house mice (*Mus
domesticus*) in live traps and responses to odors in live traps by dominant and
subordinate males. Journal of Chemical Ecology. 1997;23(11).

83. Riede T. Stereotypic Laryngeal and Respiratory Motor Patterns Generate Different Call
Types in Rat Ultrasound Vocalization. Journal of Experimental Zoology Part A: Ecological
Genetics and Physiology. 2013;319(4).

84. Prestwich KN. The Energetics of Acoustic Signaling in Anurans and Insects. American
Zoologist. 1994;34(6):625–43.

85.   Taigen TL, Wells KD. Energetics of vocalization by an anuran amphibian (Hyla versicolor).
Journal of Comparative Physiology B. 1985;155(2).

86.   Burkhard TT, Sachs ER, Phelps SM. Female preferences for high vocal effort in singing
mice. Behaviour. 2023;160(3–4).

87.   Burkhard TT, Westwick RR, Phelps SM. Adiposity signals predict vocal effort in Alston's
singing mice. Proceedings of the Royal Society B: Biological Sciences. 2018;285(1877).

88.   Spagopoulou F, Vega-Trejo R, Head ML, Jennions MD. Shifts in reproductive investment in
response to competitors lower male reproductive success. American Naturalist.
2020;196(3).

89.   Bretman A, Gage MJG, Chapman T. Quick-change artists: Male plastic behavioural
responses to rivals. Trends in Ecology and Evolution. 2011;26(9).

90.   Parker GA, Ball MA, Stockley P, Gage MJG. Sperm competition games: A prospective
analysis of risk assessment. Proceedings of the Royal Society B: Biological Sciences.
1997;264(1389):1793–802.

91.   Parker GA, Ball MA, Stockley P, Gage MJG. Sperm competition games: Individual
assessment of sperm competition intensity by group spawners. Proceedings of the Royal
Society B: Biological Sciences. 1996;263(1375).

92.   Parker GA. Conceptual developments in sperm competition: A very brief synopsis:
Developments in sperm competition. Philosophical Transactions of the Royal Society B:
Biological Sciences. 2020;375(1813):120–4.

93.   Bretman A, Westmancoat JD, Gage MJG, Chapman T. Males use multiple, redundant cues
to detect mating rivals. Current Biology. 2011;21(7).

94.   Dore AA, Rostant WG, Bretman A, Chapman T. Plastic male mating behavior evolves in
response to the competitive environment. Evolution. 2021 Jan 1;75(1):101–15.

95.   Dixson AF, Anderson MJ. Sexual behavior, reproductive physiology and sperm competition
in male mammals. Physiology & Behavior. 2004 Nov;83(2):361–71.

96.   DeLong KT. Population Ecology of Feral House Mice. Ecology. 1967 Jul 1;48(4):611–34.

97.   Evans FC. A Population Study of House Mice (*Mus musculus*) Following a Period of Local
Abundance. Journal of Mammalogy. 1949 Nov;30(4):351.

98.   Southwick CH. Population characteristics of house mice living in English corn ricks: density
relationships. Proceedings of the Zoological Society of London. 1958 Sep;131(2):163–75.

99. Nathan HW, Clout MN, MacKay JWB, Murphy EC, Russell JC. Experimental island invasion
of house mice. Population Ecology. 2015 Apr;57(2):363–71.

100. Asaba A, Okabe S, Nagasawa M, Kato M, Koshida N, Osakada T, et al. Developmental social
environment imprints female preference for male song in mice. PloS one [Internet]. 2014
Feb 5 [cited 2024 Sep 5];9(2). Available from:
https://pubmed.ncbi.nlm.nih.gov/24505280/

101. Mossman CA, Drickamer LC. Odor Preferences of Female House Mice (*Mus domesticus*) in
Seminatural Enclosures. Journal of Comparative Psychology. 1996;110(2).

102. Ronald KL, Zhang X, Morrison MV, Miller R, Hurley LM. Male mice adjust courtship
behavior in response to female multimodal signals. PLoS ONE. 2020;15(4):1–23.

103. DelBarco-Trillo J, Ferkin MH. Male mammals respond to a risk of sperm competition
conveyed by odours of conspecific males. Nature. 2004;431(7007).

104. Rouse J, Bretman A. Exposure time to rivals and sensory cues affect how quickly males
respond to changes in sperm competition threat. Animal Behaviour. 2016 Dec 1;122:1–8.

105. Parker GA. Sperm Competition and Its Evolutionary Consequences in the Insects. Annals of
the Entomological Society of America. 1970;45:525–67.

106. Johnston RE. Individual Odors and Social Communication. Individual Recognition, Kin
Recognition, and Scent Over-Marking. Advances in the Study of Behavior. 2008;38.

107. Rowland WJ. Aggression versus courtship in threespine sticklebacks and the role of
habituation to neighbours. Animal Behaviour. 1988;36(2).

108. Utne-Palm AC, Hart PJB. The effects of familiarity on competitive interactions between
threespined sticklebacks. Oikos. 2000;91(2).

109. Ydenberg RC, Giraldeau LA, Falls JB. Neighbours, strangers, and the asymmetric war of
attrition. Animal Behaviour. 1988;36(2).

110. Fisher RA. Evolution and bird sociality. In: Huxley J, Hardy AC, Ford EB, editors. Evolution as
a Process. London: Alen & Unwin; 1954. p. 71–83.

111. Dobly A. Scent marking by common voles *Microtus arvalis* in the presence of a same-sex
neighbour. Acta Theriologica. 2005;50(3).

112. Carazo P, Font E, Desfilis E. Beyond "nasty neighbours" and "dear enemies"? Individual
recognition by scent marks in a lizard (Podarcis hispanica). Animal Behaviour. 2008;76(6).

113. Keesom SM, Finton CJ, Sell GL, Hurley LM. Early-Life Social Isolation Influences Mouse
Ultrasonic Vocalizations during Male-Male Social Encounters. PLoS ONE. 2017;12(1).

114. Wang H, Liang S, Burgdorf J, Wess J, Yeomans J. Ultrasonic Vocalizations Induced by Sex
and Amphetamine in M2, M4, M5 Muscarinic and D2 Dopamine Receptor Knockout
Mice. PLoS ONE. 2008;3(4):1893.

115. Lupanova AS, Egorova MA. Vocalization of sex partners in the house mouse (*Mus
Musculus*). Journal of Evolutionary Biochemistry and Physiology. 2015;51(4).

116. Fulenwider HD, Caruso MA, Ryabinin AE. Manifestations of domination: Assessments of
social dominance in rodents. Genes, Brain and Behavior. 2022;21(3):1–15.

117. Wolff RJ. Mating behaviour and female choice: their relation to social structure in wild
caught House mice (*Mus musculus*) housed in a semi-natural environment. Journal of
Zoology. 1985;207(1).

118. Wang F, Kessels HW, Hu H. The mouse that roared: Neural mechanisms of social hierarchy.
Trends in Neurosciences. 2014;37(11):674–82.

119. Southwick CH, Clark LH. Aggressive behaviour and exploratory activity in fourteen mouse
strains. American Zoologist. 1966;6:559.

120. Ling TJ, Summers CH, Renner KJ, Watt MJ. Opponent recognition and social status
differentiate rapid neuroendocrine responses to social challenge. Physiology & Behavior.
2010 Apr;99(5):571–8.

121. Friesen CN, Maclaine KD, Hofmann HA. Social status mediates behavioral, endocrine, and
neural responses to an intruder challenge in a social cichlid, Astatotilapia burtoni.
Hormones and Behavior. 2022 Sep;145:105241.

122. Cooper MA, Clinard CT, Morrison KE. Neurobiological mechanisms supporting experience-
dependent resistance to social stress. Neuroscience. 2015 Apr;291:1–14.

123. Sgoifo A, Costoli T, Meerlo P, Buwalda B, Pico'-Alfonso MA, De Boer S, et al. Individual
differences in cardiovascular response to social challenge. Neuroscience and
Biobehavioral Reviews. 2005;29(1 SPEC. ISS.).

124. Larrieu T, Sandi C. Stress-Induced Depression: Is Social Rank a Predictive Risk Factor?
BioEssays. 2018;40(7).

125. Tibbetts EA, Pardo-Sanchez J, Weise C. The establishment and maintenance of dominance
hierarchies. Phil Trans R Soc B. 2022 Feb 28;377(1845):20200450.

126. Dewsbury DA. Dominance Rank, Copulatory Behavior, and Differential Reproduction. The
Quarterly Review of Biology. 1982 Jun;57(2):135–59.

127. Chase ID, Seitz K. Self-structuring properties of dominance hierarchies. A new perspective.
In: Advances in Genetics. 2011.

A. Unfamiliar odor cohort

Legend:
- Dom Fem
- Dom Fem+Male
- Sub Fem
- Sub Fem+Male

B. Familiar odor cohort

Legend:
- Dom Fem
- Dom Fem+Male
- Sub Fem
- Sub Fem+Male

A. Unfamiliar Odor Cohort

B. Familiar Odor Cohort

A. Unfamiliar odor cohort

B. Unfamiliar odor cohort

C. Familiar odor cohort

D. Familiar odor cohort

A. Unfamiliar odor cohort: Fem trials

B. Familiar odor cohort: Fem trials

C. Unfamiliar odor cohort: Fem+Male trials

D. Familiar odor cohort: Fem+Male trials