Training deep learning models on personalized genomic sequences improves variant effect prediction

Adam Y. He
Cornell University
Ithaca, NY 14850
`ayh8@cornell.edu`

Nathan P. Palamuttam
Cornell University
Ithaca, NY 14850
`npp8@cornell.edu`

Charles G. Danko[*]
Cornell University
Ithaca, NY 14850
`dankoc@gmail.com`

Abstract

Sequence-to-function models have broad applications in interpreting the molecular impact of genetic variation, yet have been criticized for poor performance in this task. Here we show that training models on functional genomic data with matched personal genomes improves their performance at variant effect prediction. Variant effect representations are retained even when transfer learning models to unseen cellular contexts and experimental readouts. Our results have implications for interpreting trait-associated genetic variation.

Deep learning models have rapidly become the state of the art for predicting marks of regulatory function from genome sequence. Despite accurately predicting both the pattern of chromatin marks and gene expression across genomic loci, however, several recent studies have highlighted significant limitations in the ability of some models to predict differences between individuals [1, 2]. Existing models struggle most with gene expression predictions, due in part to challenges integrating the *cis*-regulatory effects of distal enhancers [3, 4]. An additional issue is that most sequence-to-function models are trained using a single haploid reference genome as input [4–10]. While this approach is computationally tractable and enables the accumulation of large datasets where matched genome sequences are not known, training on a single genome prevents models from observing the impact of genetic variation on genome function, potentially resulting in poor performance in variant interpretation tasks. Consequently, the extent to which training models using personalized genomes improves performance has become a subject of intense interest [11–13].

We recently described CLIPNET [14], a deep convolutional neural network trained to use input DNA sequences to predict maps of transcription initiation measured using PRO-cap, an assay that maps the 5'-capped ends of nascent RNAs [15]. We trained CLIPNET using functional data and matched diploid genomes from 58 Yoruban lymphoblastoid cell lines (LCLs) (+9 biological replicates) [16]. In addition to accurately predicting transcription initiation at single-nucleotide resolution from genome sequences, CLIPNET can correctly impute the effects of genetic variants on both transcription initiation quantity (tiQTLs) and directionality (diQTLs) [16].

Here, we asked whether training CLIPNET using matched personal genome sequences improved its ability to predict the impact of genetic variation on transcription initiation. We first trained CLIPNET models on random subsets of the 67 PRO-cap libraries used in our original study. When predicting across loci on holdout chromosomes, predictions of the profile (distribution of PRO-cap reads within a given prediction window) and the total quantity of transcription initiation rapidly improved in accuracy as the number of training individuals increased, with diminishing returns beginning at around 20-30 individuals (Fig. 1A). We observed a similar trend when assessing prediction accuracy at genetic variants associated with proximal *cis* changes in transcription initiation in LCLs. Using an L^2-based variant effect score [4, 14], we found that CLIPNET required training on 20-30 individuals before achieving high accuracy at predicting the effects of tiQTLs and diQTLs (Fig. 1B).

---

[*]Corresponding author

Figure 1: **Training deep learning models on personalized genomic sequences improves variant effect prediction.**
(**A**) Schematic of the personalized genomes training scheme used for CLIPNET and the two ablation strategies (training data subsampling and variant masking/reference training). (**B**) Evaluation of subsampled models on transcription initiation profile (left) and quantity (right) prediction ($n = 4901$ loci). PCC = Pearson's correlation coefficient. (**C**) Evaluation of subsampled models on predicting effects of tiQTLs (left, $n = 2057$) and diQTLs (right, $n = 1207$) (PCC between observed and predicted L^2 variant effect scores). In both **B** and **C**, the performance of the original CLIPNET model is indicated by red dashed lines, while the performance of the reference-trained CLIPNET model is indicated by blue dashed lines. (**D**) Comparison of tiQTL (left two panels) and diQTL (right two panels) effect prediction by reference-trained (left panel in each pair) and personalized (right panel in each pair) CLIPNET models.

While the subsampling test indicated that training on more individuals improves model accuracy, these results could also be explained by CLIPNET benefiting from more training data, irrespective of whether the sequence inputs come from personalized genomes or reference genomes. To directly measure the contribution of training on personalized genomes to model performance, we trained a CLIPNET model on the full 67 library dataset [14], using only reference genome sequence with the genetic variants masked. The reference-trained CLIPNET model performed similarly to the personalized CLIPNET model and outperformed most of the subsampled models at predicting initiation profiles and quantities across loci in the genome (Fig. 1B, Supplementary Fig. S1). However, the reference-trained model substantially underperformed the full personalized CLIPNET model at predicting QTL effects, performing roughly on par with subsampled models trained on only 10 or 20 individuals (Fig. 1C-D). Notably, the reference-trained model predicted 17.0% ($n = 350$) of tested tiQTLs and 20.0% ($n = 241$) of tested diQTLs as having no effect (Fig. 1D). These results show that training on larger numbers of functional genomic datasets improves performance across loci, but performance on variant effect prediction tasks improves further by using dataset-matched personal genomes.

We next asked whether the performance gain from using larger functional datasets and matched personal genomes generalizes across cell types and experimental readouts. While the difference in initiation QTL performance was compelling, the existence of linkage disequilibrium between variants in endogenous genomic sequences complicates causality interpretations. To address this, we used massively parallel reporter assays (MPRAs) to as a ground truth for regulatory SNP impact. Although PRO-cap and MPRAs measure distinct biological processes, transcription initiation and enhancer activity are correlated [17, 18], suggesting that models predicting SNP effects on initiation should also perform reasonably well in predicting enhancer activity.

We used transfer learning to adapt CLIPNET to a K562 PRO-cap dataset [19] and evaluated the accuracy of SNP effect prediction against a massively parallel reporter assay (MPRA) ground truth [20] (Fig. 2A). The transfer-learned CLIPNET K562 model showed strong correlations with experimental PRO-cap data, performing similarly to ProCapNet [7], a model trained natively on K562 PRO-cap data using the reference genome, in predictions across loci (Supplementary Fig. S2). Despite similar performance across loci, CLIPNET significantly outperformed ProCapNet in predicting variant impact measured using the MPRA on SNPs from CLIPNET's holdout chromosomes (Fig. 2B-C). Both PRO-cap-based models significantly outperformed Enformer [6], a large multi-task transformer-based model that has been criticized for poor performance on predicting personal gene expression [1, 2].

To assess whether the variant representations learned by training on personalized genome sequences improved variant effect predictions, we also transferred the reference-trained CLIPNET model to K562. Again, the personalized CLIPNET K562 model outperformed the reference-trained CLIPNET K562 model at variant effect prediction, albeit by a smaller margin than was observed in the LCL initiation QTL prediction task (p = 0.038; bootstrap test; Fig. 2B-D). Next, we compared the change in transcription initiation of each SNP predicted by the model to the fold-change measured using the MPRA. The CLIPNET model trained using personal genomes estimated effect sizes that were closer to the MPRA than ProCapNet or the reference-trained CLIPNET model (Fig. 2D). These results reinforce our initiation QTL analysis and suggest that training on personalized genome sequences allows CLIPNET to observe substantial activity differences between nearly identical input DNA sequences, which reference-trained models fail to observe.

Our work has several valuable applications. First, MPRAs are frequently used to identify and validate candidate causal SNPs from genetic association studies [20–24]. Achieving reasonable *in silico* performance by training on both large functional genomic datasets and matched personal genomes has widespread potential for interpreting the molecular effects of trait-associated SNPs and prioritization for experimental validation. Second, enhancer design is an emerging field [25, 26]. Models such as CLIPNET and ProCapNet, which predict transcription initiation from local sequences, are particularly well suited for designing and interpreting enhancer elements, which often show a stereotypical divergent initiation pattern in mammals [15, 18, 27, 28].

Recent work has shown that while fine-tuning gene expression predictions using genetic variation can improve predictions for individual genes, this performance uplift does not generalize to unobserved genes [12, 13]. Expression models are most sensitive to promoters and capture the effects of distal enhancers much less accurately [3, 29, 30]. These results make intuitive sense: learning accurate long-range models that capture the effect of distal enhancers requires fitting two complex and independent functions. First, the impact of DNA sequence on local *cis*-regulatory activity, and second, the complex logic by which different *cis*-regulatory elements integrate their signals across a locus. Both of these complex functions must be learned from a limited set of only ~20,000 genes. Adding to the challenge, most of the highly expressed genes across different cell types—where mistakes are penalized heavily for models trained across different cell types or tissues—are housekeeping genes, which are less likely to be impacted by enhancers [31]. We argue that this complexity, combined with the limited set of training examples, makes the problem even more difficult.

Our results suggest a path forward for developing accurate variant interpretation models that predict gene expression. We propose that the first step is to accurately learn the effect of variants on local *cis*-regulatory activity. Our findings

Figure 2: **CLIPNET transfer learned to predict initiation in K562 outperforms models trained on only reference genome sequences at predicting MPRA variant effects [20].** (**A**) Schematic illustrating the procedure for transfer learning CLIPNET to predicting initiation in K562 and for benchmarking on MPRA variant effect prediction. (**B**) Precision-recall curves for all tested models on classifying emVars ($n = 2038$) vs expressed non-emVars ($n = 4057$). (**C**) Bootstrapped estimates of change in area under the precision-recall curve of all other models compared to CLIPNET. (**D**) Comparison of the scalar predictions of emVar effects between ProCapNet (left), reference-trained CLIPNET transferred to K562 (center), and CLIPNET transferred to K562 (right).

show that local models are substantially improved by incorporating genetic variation and by using larger datasets with multiple independent measurements of the same cell or tissue type. Genetic variation enhances the model's ability to generalize to unseen variants and even to distinct experimental readouts, such as MPRAs. Accurate local models can then be integrated into models trained to predict gene expression, either through heuristics [30] or through long-range transformer-based models trained to learn regulatory logic. We also suggest that training more complex models may benefit from adding biological constraints, for instance by initializing model parameters using experimental chromatin contact data from assays such as Hi-C or Micro-C. This stepwise approach provides a more feasible path to addressing the inherent complexity of gene regulation models, laying the groundwork for better generalizability and numerous downstream applications interpreting trait-associated genetic variation.

Methods

Subsampled model training

We randomly selected $n = 5, 10, 15, 20, 30$ PRO-cap libraries from the 67 used to train CLIPNET, then trained subsampled CLIPNET models on matched personal genome sequences (1000 Genomes Project, `https://ftp.10 00genomes.ebi.ac.uk/vol1/ftp/data_collections/1000_genomes_project/release/20190312_bial lelic_SNV_and_INDEL/`) and PRO-cap tracks ([16], GSE110638) following the methods described in our previous manuscript (notably, using the same ensembling and holdout scheme) [14]. We avoided sampling multiple isogenic replicates from the same individual in each subsampling experiment. For each n, we performed 5 such runs, resulting in a total of 25 subsampled CLIPNET models.

Variant masking / reference sequence training

We retrained a CLIPNET model using the full set of 67 PRO-cap libraries, but instead of using the corresponding personalized genome sequences, we extracted sequences from the hg38 reference genome.

LCL benchmarks

We benchmarked the subsampled and reference-trained CLIPNET models using the tasks described in our previous manuscript [14].

Specifically, for the cross-loci benchmark, we considered a set of 4901 high confidence PRO-cap peaks on the holdout chromosomes. We extracted the hg38 reference sequence and the average RPM-normalized PRO-cap coverage around each of these peaks [16], then measured the performance of the models by the median Pearson's correlation between predicted and observed PRO-cap coverage tracks (profile performance) and the Pearson's correlation between the predicted and observed $\log_{10}$ total PRO-cap coverage (quantity performance). In our previous paper, we reported performance metrics for each model in the CLIPNET ensemble on its unique holdout set; for the sake of brevity here, we only report the performance of the ensemble on the chromosomes that were entirely withheld during training (chromosomes 9, 13, 20, and 21).

For the tiQTL and diQTL benchmarks, we used the same set of 2,057 tiQTLs and 1,027 diQTLs. We measured the predicted and observed QTL effects by taking the L^2 norm of the difference vector between averaged homozygous reference and averaged homozygous alternative tracks. To avoid data leakage, we used the same QTL score compositing scheme we previously described [14].

K562 transfer learning

We used transfer learning to adapt CLIPNET to a K562 PRO-cap dataset. We downloaded the bigWig files (ENCSR261KBX) from the ENCODE data portal [32], then merged isogenic replicates. To maintain consistency across the PRO-cap datasets used in this study, we re-called PINTS peaks [27] rather than using the ENCODE peaks and applied an RPM transform to the PRO-cap tracks for training. As this dataset was mapped to hg38 and K562 is a cancer cell line with a poorly defined karyotype, hg38 was used as the source of genome sequences for transfer learning.

The K562 models were initialized using the weights from the full personalized LCL model, then trained on the K562 dataset. We used the same hyperparameters as when training the LCL models, but with a single initial warmup epoch at a learning rate of 0.0001. To test the contribution of pre-training on personalized genomic sequences, we also applied the same transfer learning procedure to the reference-trained CLIPNET LCL model.

K562 PRO-cap prediction benchmark

We compared the performance of CLIPNET and ProCapNet at cross-loci prediction in K562. Since the two models used different holdout datasets, we decided to evaluate the prediction accuracies of individual CLIPNET and ProCapNet model folds on their respective holdout chromosomes. We calculated the median Pearson's correlation between the predicted and observed profiles, as well as the Pearson's correlation between the predicted and observed $\log_1 0$ transcription initiation quantities (as described above). The custom K562 peak call set was used for both models. CLIPNET was evaluated on 1000 bp sequence windows and 500 bp tracks, whereas ProCapNet was evaluated on 2114 bp sequence windows and 1000 bp tracks.

K562 MPRA benchmark

Siraj et al. recently characterized the impact of hundreds of thousands of non-coding variants in MPRAs across several human cell lines [20]. We chose to analyze the K562 MPRA dataset, as K562 is the only cell line tested in that study with published PRO-cap data, and which was used to train Enformer [6] and ProCapNet [7] models.

To test the ability of CLIPNET K562 (both personal and reference-pre-trained) to SNP effects in this MPRA, we reconstructed the MPRA sequences by inserting the tested 200 bp oligonucleotides (Supplemental Table 2 of [20]) directly upstream of the minimal TATA promoter of pMPRAv3:minP-GFP, (Addgene #109035) as described in [20]. We then extracted 1 kb of sequence from each reporter containing the oligonucleotide insert and the minimal promoter. We predicted the transcription initiation quantity for each allele and measured the predicted effect of each SNP as the $\log_2$ ratio of the reference and alternate allele predictions.

To generate ProCapNet [7] predictions, we simply padded the sequences constructed for CLIPNET out to the 2114 bp context length used by that model using the reporter backbone. We calculated the average predicted initiation quantity across the 7 ProCapNet K562 model replicates (ENCFF976FHE) for each reporter construct, then calculated the SNP effects as described for CLIPNET.

For the Enformer [6] predictions, we used the endogenous genomic sequences surrounding each SNP, as the long context length used for this model makes it impractical to test on the reporter sequences. We extracted 196,608 bp of sequence centered on each SNP (excluding those that would cause spillovers off the edge of the chromosome), then predicted K562 DNase (tracks 33, 34, 35, 121, 122, 123, and 625) and CAGE (tracks 4828, 5111) tracks in the central 8 bins (representing 1024 bp) using the PyTorch implementation of Enformer (`https://github.com/lucidrains/en former-pytorch/`). After some testing, we determined that K562 DNase (specifically, track 122, corresponding to ENCODE accession ENCFF868NHV) was the best predictor for MPRA SNP effect (as described below). We thus set the Enformer SNP predictions as the $\log_2$ ratio between the reference and alternative allele DNase predictions.

We considered the following two benchmarks for MPRA prediction. First, we measured the ability of each model to distinguish between expression modulating variants (emVar) and active non-emVars in K562 (Supplemental Table 3 of [20]). Specifically, we calculated precision-recall curves (PRC), using the square of the SNP prediction for each model as the predictor. To avoid possible data leakage from variants in the pre-training of the personalized CLIPNET model, we only considered variants on the holdout chromosomes for CLIPNET (chromosomes 9, 13, 20, and 21). We also excluded variants that would cause chromosome edge spillovers for Enformer, resulting in 2038 emVars and 4057 non-emVars. We observed that the personalized CLIPNET K562 model achieved the highest area under the PRC (auPRC). To measure the accuracy of this estimate, we performed $n = 1000$ bootstraps of the holdout SNPs, then calculated the difference in auPRC between all other models and the personalized CLIPNET K562 model.

Since both CLIPNET K562 models and ProCapNet K562 produce predictions of transcription initiation, we also quantitatively assessed the SNP effect predictions made by all three models. We generated scatterplots of the $\log_2$FC predictions made by each model against the observed $\log_2$FC values (specifically, mean RNA counts/mean DNA counts per allele). We also calculated Pearson's correlation coefficients and sign mismatches between the predicted and observed $\log_2$FC values.

Declarations

Data availability

This study makes use of publicly available datasets. URLs and accession codes are provided in the methods section. The original CLIPNET model and the transfer-learned K562 model are available at `https://doi.org/10.5281/ze nodo.10408623` and `https://doi.org/10.5281/zenodo.11196189`, respectively. All other supporting data for

this study, including model weights for the ablation tests, are available at `https://doi.org/10.5281/zenodo.138 23014`.

Code availability

Code for performing and plotting the results of the two ablation tests is available at `https://github.com/Danko-L ab/clipnet_ablation`. Code for transfer learning CLIPNET to predicting initiation in K562 and evaluating on the MPRA dataset are is available at `https://github.com/Danko-Lab/clipnet_k562`. This study also makes use of utility scripts developed for the original CLIPNET paper (`https://github.com/Danko-Lab/clipnet`).

Funding

A.Y.H. was supported by an NIH T32 training grant (5T32HD057854) to Research and Career Training in Vertebrate Developmental Genomics at Cornell University. This work was supported by a grant from the National Human Genome Research Institute (R01HG010346) and by the Laboratory Directed Research and Development program at Sandia National Laboratories. Sandia National Laboratories is a multimission laboratory managed and operated by National Technology and Engineering Solutions of Sandia LLC, a wholly owned subsidiary of Honeywell International Inc. for the U.S. Department of Energy's National Nuclear Security Administration under contract DE-NA0003525. Some of the GPU computing in this project was performed on Bridges-2 at the Pittsburgh Supercomputing Center through allocation BIO210011P from the Advanced Cyberinfrastructure Coordination Ecosystem: Services & Support (ACCESS) program, which is supported by National Science Foundation grants #2138259, #2138286, #2138307, #2137603, and #2138296. The content of this manuscript is solely the responsibility of the authors and does not necessarily represent the official views of Cornell University or any funding agency.

Author contributions

A.Y.H and C.G.D conceived of the project. A.Y.H designed the analyses with input from C.G.D. A.Y.H. and N.P.P. implemented the analyses. A.Y.H. and C.G.D. wrote the manuscript together.

Competing Interests

The authors declare no competing interests.

Acknowledgments

We thank Li Yao and Haiyuan Yu for providing a pre-publication version of their PINTS peak-calling software, Hojoong Kwak for assistance with interpreting his group's QTL analysis, Kelly Cochran for advice on working with ProCapNet, and Anshul Kundaje for suggesting the variant masking ablation test.

References

[1] Huang, C. *et al.* Personal transcriptome variation is poorly explained by current genomic deep learning models. *Nat. Genet.* (2023). URL `http://dx.doi.org/10.1038/s41588-023-01574-w`.

[2] Sasse, A. *et al.* Benchmarking of deep neural networks for predicting personal gene expression from DNA sequence highlights shortcomings. *Nat. Genet.* (2023). URL `http://dx.doi.org/10.1038/s41588-023-01524-6`.

[3] Karollus, A., Mauermeier, T. & Gagneur, J. Current sequence-based models capture gene expression determinants in promoters but mostly ignore distal enhancers. *Genome Biol.* **24**, 56 (2023). URL `http://dx.doi.org/10.1186/s13059-023-02899-9`.

[4] Linder, J., Srivastava, D., Yuan, H., Agarwal, V. & Kelley, D. R. Predicting RNA-seq coverage from DNA sequence as a unifying model of gene regulation. *bioRxiv* 2023.08.30.555582 (2023). URL `https://www.biorxiv.org/content/10.1101/2023.08.30.555582v1`.

[5] Avsec, Z. *et al.* Base-resolution models of transcription-factor binding reveal soft motif syntax. *Nat. Genet.* **53**, 354–366 (2021). URL `http://www.nature.com/articles/s41588-021-00782-6`.

[6] Avsec, Z. *et al.* Effective gene expression prediction from sequence by integrating long-range interactions. *Nat. Methods* **18**, 1196–1203 (2021). URL `https://www.nature.com/articles/s41592-021-01252-x`.

[7] Cochran, K. *et al.* Dissecting the cis-regulatory syntax of transcription initiation with deep learning. *bioRxiv* 2024.05.28.596138 (2024). URL `https://www.biorxiv.org/content/10.1101/2024.05.28.596138v1`.

[8] Dudnyk, K., Cai, D., Shi, C., Xu, J. & Zhou, J. Sequence basis of transcription initiation in the human genome. *Science* **384**, eadj0116 (2024). URL `https://www.science.org/doi/abs/10.1126/science.adj0116`. `https://www.science.org/doi/pdf/10.1126/science.adj0116`.

[9] Zhou, J. *et al.* Deep learning sequence-based ab initio prediction of variant effects on expression and disease risk. *Nat. Genet.* **50**, 1171–1179 (2018). URL `https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6094955/`.

[10] Zhou, J. Sequence-based modeling of three-dimensional genome architecture from kilobase to chromosome scale. *Nat. Genet.* **54**, 725–734 (2022). URL `http://dx.doi.org/10.1038/s41588-022-01065-4`.

[11] Bajwa, A., Rastogi, R., Kathail, P., Shuai, R. W. & Ioannidis, N. Characterizing uncertainty in predictions of genomic sequence-to-activity models. In *Machine Learning in Computational Biology*, 279–297 (PMLR, 2024). URL `https://proceedings.mlr.press/v240/bajwa24a.html`.

[12] Drusinsky, S., Whalen, S. & Pollard, K. S. Deep-learning prediction of gene expression from personal genomes. *bioRxiv* 2024.07.27.605449 (2024). URL `https://www.biorxiv.org/content/10.1101/2024.07.27.605449v1.abstract`.

[13] Rastogi, R., Reddy, A. J., Chung, R. & Ioannidis, N. M. Fine-tuning sequence-to-expression models on personal genome and transcriptome data. *bioRxiv* 2024.09.23.614632 (2024). URL `https://www.biorxiv.org/content/10.1101/2024.09.23.614632v1.abstract`.

[14] He, A. Y. & Danko, C. G. Dissection of core promoter syntax through single nucleotide resolution modeling of transcription initiation. *bioRxiv* 2024.03.13.583868 (2024). URL `https://www.biorxiv.org/content/10.1101/2024.03.13.583868`.

[15] Core, L. J. *et al.* Analysis of nascent RNA identifies a unified architecture of initiation regions at mammalian promoters and enhancers. *Nat. Genet.* **46**, 1311–1320 (2014). URL `http://www.nature.com/articles/ng.3142`.

[16] Kristjánsdóttir, K., Dziubek, A., Kang, H. M. & Kwak, H. Population-scale study of eRNA transcription reveals bipartite functional enhancer architecture. *Nat. Commun.* **11**, 5963 (2020). URL `http://dx.doi.org/10.1038/s41467-020-19829-z`.

[17] Andersson, R. *et al.* An atlas of active enhancers across human cell types and tissues. *Nature* **507**, 455–461 (2014). URL `http://dx.doi.org/10.1038/nature12787`.

[18] Tippens, N. D. *et al.* Transcription imparts architecture, function and logic to enhancer units. *Nat. Genet.* **52**, 1067–1075 (2020). URL `http://dx.doi.org/10.1038/s41588-020-0686-2`.

[19] Lis, J. ENCSR261KBX (2021). URL `https://www.encodeproject.org/experiments/ENCSR261KBX/`.

[20] Siraj, L. *et al.* Functional dissection of complex and molecular trait variants at single nucleotide resolution. *bioRxiv* 2024.05.05.592437 (2024). URL `https://www.biorxiv.org/content/10.1101/2024.05.05.592437v1`.

[21] Abell, N. S. *et al.* Multiple causal variants underlie genetic associations in humans. *Science* **375**, 1247–1254 (2022). URL `http://dx.doi.org/10.1126/science.abj5117`.

[22] Ray, J. P. *et al.* Prioritizing disease and trait causal variants at the TNFAIP3 locus using functional and genomic features. *Nat. Commun.* **11**, 1237 (2020). URL `https://www.nature.com/articles/s41467-020-15022-4`.

[23] Tewhey, R. *et al.* Direct identification of hundreds of expression-modulating variants using a multiplexed reporter assay. *Cell* **165**, 1519–1529 (2016). URL `http://dx.doi.org/10.1016/j.cell.2016.04.027`.

[24] Vockley, C. M. *et al.* Massively parallel quantification of the regulatory effects of noncoding genetic variation in a human cohort. *Genome Res.* **25**, 1206–1214 (2015). URL `http://genome.cshlp.org/content/25/8/1206.abstract`.

[25] de Almeida, B. P. *et al.* Targeted design of synthetic enhancers for selected tissues in the drosophila embryo. *Nature* 1–2 (2023). URL `https://www.nature.com/articles/s41586-023-06905-9`.

[26] Taskiran, I. I. *et al.* Cell type directed design of synthetic enhancers. *Nature* 1–3 (2023). URL `https://www.nature.com/articles/s41586-023-06936-2`.

[27] Yao, L. *et al.* A comparison of experimental assays and analytical methods for genome-wide identification of active enhancers. *Nat. Biotechnol.* **40**, 1056–1065 (2022). URL `http://dx.doi.org/10.1038/s41587-022-01211-7`.

[28] Tome, J. M., Tippens, N. D. & Lis, J. T. Single-molecule nascent RNA sequencing identifies regulatory domain architecture at promoters and enhancers. *Nat. Genet.* **50**, 1533–1541 (2018). URL `http://dx.doi.org/10.1038/s41588-018-0234-5`.

[29] Lin, J., Luo, R. & Pinello, L. EPInformer: a scalable deep learning framework for gene expression prediction by integrating promoter-enhancer sequences with multimodal epigenomic data. *bioRxiv* 2024.08.01.606099 (2024). URL `https://www.biorxiv.org/content/10.1101/2024.08.01.606099v1.abstract`.

[30] Martyn, G. E. *et al.* Rewriting regulatory DNA to dissect and reprogram gene expression. *bioRxiv* 2023.12.20.572268 (2023). URL `https://www.biorxiv.org/content/10.1101/2023.12.20.572268v1`.

[31] Gasperini, M. *et al.* A genome-wide framework for mapping gene regulation via cellular genetic screens. *Cell* **176**, 377–390.e19 (2019). URL `https://pubmed.ncbi.nlm.nih.gov/30612741/`.

[32] Luo, Y. *et al.* New developments on the encyclopedia of DNA elements (ENCODE) data portal. *Nucleic Acids Res.* **48**, D882–D889 (2020). URL `http://dx.doi.org/10.1093/nar/gkz1062`.

Supplementary Figures

Figure S1:   **Comparison of reference-trained and personalized CLIPNET predictions.**   (**A**) Initiation profile prediction. (**B**) Initiation quantity prediction.

Figure S2:   **Comparison of CLIPNET and ProCapNet cross-loci predictions in K562.**   (**A**) Initiation profile prediction. (**B**) Initiation quantity prediction. CLIPNET ($n = 9$) and ProCapNet ($n = 7$) models were evaluated on their respective holdout chromosomes.