# Emergence of Dip2-mediated Specific DAG-based PKC Signalling Axis in Eukaryotes

Sakshi Shambhavi[#1,3], Sudipta Mondal[#1], Arnab Chakraborty[2], Nikita Shukla[1], Bapin K. Panda[1], Santhosh Kumar[1,3], Priyadarshan Kinatukara[1], Biswajit Pal[1], Siddhesh S. Kamat[2] and Rajan Sankaranarayanan[1,3]*

[1]CSIR-Centre for Cellular and Molecular Biology, Hyderabad, India

[2]Department of Biology, Indian Institute of Science Education and Research (IISER), Pune, India

[3]Academy of Scientific and Innovative Research (AcSIR), Ghaziabad, India-201002

[#]These authors have contributed equally

*For correspondence: sankar@ccmb.res.in

**Abstract**

Diacylglycerols (DAGs) are used for metabolic purposes and are tightly regulated secondary lipid messengers in eukaryotes. DAG subspecies with different fatty-acyl chains are proposed to be involved in the activation of distinct PKC isoforms, resulting in diverse physiological outcomes. However, the molecular players and the regulatory origin for fine-tuning the PKC pathway are unknown. Here, we show that Dip2, a conserved DAG regulator across Fungi and Animalia, has emerged as a modulator of PKC signalling in yeast. Dip2 maintains the level of a specific DAG subpopulation, required for the activation of PKC-mediated cell wall integrity pathway. Interestingly, the canonical DAG-metabolism pathways, being promiscuous, are decoupled from PKC signalling. We demonstrate that these DAG subspecies are sourced from a phosphatidylinositol pool generated by the acyl-chain remodelling pathway. Furthermore, we provide insights into the intimate coevolutionary relationship between the regulator (Dip2) and the effector (PKC) of DAG-based signalling. Hence, our study underscores the establishment of Dip2-PKC axis about 1.2 billion years ago in Opisthokonta, which marks the rooting of the first specific DAG-based signalling module of eukaryotes.

**Keywords**: Cell wall integrity/ Coevolution/ Diacylglycerol/ Dip2/ Lipid metabolism/ PKC

**Introduction:**

Diacylglycerol (DAG) is a conserved lipid molecule with well-established roles in membrane lipid biogenesis and storage lipid production in all life forms. DAGs are synthesized via *de novo* pathways of lipid metabolism utilising fatty acid precursors, while it can also be generated as a by-product of multiple salvage pathways in eukaryotes (Carrasco & Merida, 2007; Eichmann & Lass, 2015). DAGs are unique as they lack headgroups unlike the other abundant lipid classes. They exhibit diversity due to variations in acyl-chain length and unsaturation. These acyl-chain variations are proposed to be the basis for the multifaceted physiological roles of DAGs in metabolic and signal transduction pathways (Marignani et al, 1996; Milne et al, 2008; Schuhmacher et al, 2020). The signalling function is executed by various 'DAG effector' proteins such as DAG-dependent protein kinases, Chimaerin, Munc13, RasGRP etc. involved in cell-cycle progression, neurotransmitter release, actin cytoskeleton organization and malignant transformation (Colon-Gonzalez & Kazanietz, 2006; Yang & Kazanietz, 2003). Studies conducted ~four decades ago have indicated the role of acyl-chain compositions of DAGs in Protein Kinase C (PKC) activation (Kishimoto et al, 1980; Marignani et al., 1996; Mori et al, 1982, Kamiya et al., 2016; Masayoshi et al., 1987). However, the cellular pathways for such specific acyl-chain-based regulation of signalling events have not been identified till date.

PKC, a serine/threonine protein kinase, is a classic example of a eukaryotic DAG-based signalling molecule. Mammalian PKC isoforms play a key role in a myriad of biological processes like cell differentiation, cell migration, mitochondrial functioning, cell polarity, etc. (Newton, 2018) and its dysregulation is associated with various pathological conditions such as cancer, diabetes, Alzheimer's, heart disease, obesity and age-related metabolic disorders (Etcheberrigaray et al, 2004; Marrocco et al, 2019; Nishizuka, 1984; Schmitz-Peiffer & Biden, 2008). Fungi, on the other hand, harbour a single PKC gene (*PKC1*) that is implicated in regulating the cell wall biosynthesis pathway, rationalizing its essentiality under normal growth

conditions (Levin, 2005). Genetic or pharmacological impairment of Pkc1 in pathogenic fungi

like *Candida albicans, Cryptococcus neoformans etc.,* confers hypersensitivity to various

antifungal drugs and drastically attenuates its proliferation and virulence in murine model

(Heinisch & Rodicio, 2018; LaFayette et al, 2010) suggesting a crucial role of Pkc1 in

pathogenicity of diverse fungal groups. The array of pathologies and disrupted cellular

processes owing to hampered Pkc1 signalling warrants the necessity of stringent regulation of

Pkc1 activity.

Although DAG-based activation of PKC is well established in metazoans, the role of

specific acyl-chain compositions in fungi is still a mystery. Additionally, the molecular players

regulating these signalling DAGs remain underexplored. Recently, we have identified and

functionally characterized a conserved protein family, Disco-Interacting Protein 2 (Dip2)

(earlier annotated as Cmr2 in yeast; SGD ID: S000005619), harbouring two tandem fatty acyl-

AMP ligase-like domains (FLD1 and FLD2) along with a DMAP-binding domain 1 (DBD1)

at the N-terminus (Mondal et al, 2022). We showed that Dip2 regulates specific DAG subspecies

in terms of its acyl chain length and unsaturation across fungi and animals. Dip2 from

*Saccharomyces cerevisiae* has been shown to be involved in converting C36:0 (18:0/18:0) and

C36:1 (18:0/18:1) DAGs to triacylglycerols (TAGs) with corresponding chain lengths. These

DAG subspecies constitute only ~0.5 % of the total lipid pool but are important for maintaining

the organellar and thereby cellular homeostasis. However, the necessity to introduce such strict

regulation of specific DAG subspecies in Opisthokonta remains elusive.

Here, we show that Dip2 has emerged as a modulator of Pkc1 signalling in the model

yeast *Saccharomyces cerevisiae* by regulating the chain-length specific DAGs. By combining

genetic and lipidomic approaches, we have demonstrated that the absence of Dip2, which leads

to the accumulation of C36:0 and C36:1 DAGs, results in hyperactivation of Pkc1. This in turn

elevates the downstream signalling of cell wall integrity (CWI) pathway leading to cell wall

stress resistance. On the contrary, the bulk DAG pool, which is generally acted upon by

canonical DAG metabolizing enzymes, fails to activate Pkc1. The DAG binding domain of

Pkc1 displays a remarkable specificity to certain DAG subspecies *in vitro,* reflecting the *in vivo*

specificity. To understand this metabolic segregation of DAG pool, we have tracked the source

of the Dip2-metabolised DAG subspecies involved in Pkc1 activation. Interestingly, it was

found to be channelled majorly through the phosphatidylinositol (PI) pool enriched with C18:0

fatty acyl chain generated by acyl-chain remodelling process. Furthermore, an extensive

phylogenetic correlation analysis has revealed that Dip2 and PKC co-emerged and coevolved

during early Opisthokonta evolution to establish the selective DAG-based PKC signalling axis.

**Results:**

**Dip2 is involved in the regulation of Pkc1-mediated cell wall integrity pathway**

110           Initial phenotypic screening revealed that Dip2 is necessary for the survival of yeast in

various stress conditions like ER stress, osmotic stress etc (Mondal et al., 2022). However, *Dip2*

knockout yeast (hereafter referred to as *Δdip2*) outperforms the wildtype (*WT*) strain under cell

wall stress conditions (Mondal et al., 2022). This suggested a possible link between Dip2 and

the cell wall integrity (CWI) pathway of yeast, which is governed by Pkc1 and the downstream

mitogen-activated protein kinase (MAPK) signalling cascade. Given the requirement of DAGs

for Pkc1 activation and the role of Dip2 in regulating DAG subspecies, we hypothesized that

Dip2 maintains optimal levels of DAG subspecies necessary for activation of CWI pathway

**(Fig. 1 A)**. To test this, we compared the fitness of *WT* and *Δdip2* yeast in the presence of cell

wall stress-inducing agents like Congo red (CR) and Calcofluor white (CFW) (Roncero &

Duran, 1985), using serial dilution assay in synthetic complete (SC) media (**Fig. 1 B**) as well as

in rich media **(Fig. 1—fig. supplement 1)**. The two biological replicates*, Δdip2*_Colony1 and

*Δdip2*_Colony2 showed increased resistance to cell wall stress, and the phenotype was restored

through genetic complementation with Dip2 under its native promoter. Additionally, we

employed colony forming unit (CFU) assay for *WT* and *Δdip2* under cell wall stress. Counting

the CFU also showed a similar and significant difference in the growth of *WT* and *Δdip2*

colonies in the presence of CR **(Fig. 1 C)**. These observations suggest the regulatory role of

Dip2 in CWI pathway of yeast. Since cell wall stress is associated with Pkc1-mediated

activation of CWI pathway, we sought to test the status of Pkc1 signalling and its probable link

to Dip2-regulated DAG subspecies. First, we probed the phosphorylation levels of a

downstream MAPK effector of Pkc1 pathway, i.e., Slt2 (Suppressor of the LyTic phenotype)

(Lee et al, 1993; Levin, 2005) to quantitatively assess the extent of Pkc1 activation. In agreement

to the CWI phenotype, a ~3-fold increase in phosphorylation of Slt2 (pSlt2) was observed in

*Δdip2* as compared to *WT*, where *WT* treated with CFW is used as a positive control for pSlt2

increase **(Fig. 1 D)**.

Next, we sought to understand if the hyperactivation of Pkc1 in *Δdip2* is DAG

dependent and involves the catalytic activity of Dip2, i.e., conversion of selective DAGs to

TAGs. To answer this, we generated two catalytic mutants of Dip2 (Dip2$^{D523A}$ and Dip2$^{L687A}$),

that abolish its adenylation activity and have been shown to be unable to decrease the selective

DAG levels on complementing *Δdip2*, as reported in our previous study (Mondal et al., 2022).

The mutants were cloned under its native promoter in pYSS01 plasmid and expressed in *Δdip2*

strain **(Fig. 1—fig. supplement 2A)**. The mutants could not diminish the cell wall stress

resistance in *Δdip2*, unlike wildtype Dip2 **(Fig. 1 E)**. Additionally, complementation with the

catalytically inactive Dip2 could not bring back the pSlt2 level in *Δdip2* **(Fig. 1—fig.**

**supplement 2B)**, suggesting a direct correlation between the enzymatic activity of Dip2 and

Pkc1 hyperactivation. Hence, the catalytic activity of Dip2 *i.e.*, the regulation of DAG

subspecies level is essential in maintaining the active pool of Pkc1 for optimal CWI signalling.

Additionally, to confirm that the cell wall stress resistance observed in *Δdip2* is Pkc1

mediated, we inhibited Pkc1's activity using cercosporamide, a selective Pkc1 inhibitor

(Sussman et al, 2004) and observed decrease in the resistance to cell wall stress (**Fig. 1 F**). The

increase in pSlt2 level in *Δdip2* was also brought back to *WT* level upon treatment with

cercosporamide (**Fig. 1 G**). This implies that the hyperactivation of Pkc1 in *Δdip2* causes the

cell wall stress resistance.

These observations intrigued us to ask how Dip2 spatially regulates Pkc1 as both reside

at different subcellular locations inside the cell. Since Pkc1 is a bud site and plasma membrane

localizing protein (Andrews P. and Stark M., 2000), we wondered whether Dip2 translocates

to plasma membrane under cell wall stress. To examine this, we used *WT* strain with

endogenous Dip2 tagged with GFP and subjected it to cell wall stress. As Dip2 predominantly

localizes to the mitochondria, we co-stained cells with Mitotracker Red and found that there is

no change in the spatial organization of Dip2 (**Fig. 1—fig. supplement 3**). This suggests that

Dip2 and Pkc1 could be cross talking through other ways that are yet to be explored.

**Dip2-regulated DAG subspecies activate Pkc1 signalling via selective interaction**

Since Dip2 is known to regulate the levels of C36:0 and C36:1 DAGs in yeast, we asked

whether these specific DAG subspecies play any role in activation of Pkc1 signalling. Firstly,

we examined how the system responds to cell wall stress in terms of the DAG levels. While

tracking the Pkc1 signalling at temporal scale, we observed profound activation of Pkc1

pathway after 20-30 minutes of CFW treatment (**Fig. 2—fig. supplement 1A**). Hence, we

performed the lipidomic analysis of *WT* yeast pulse-treated with CFW for 30 minutes.

Interestingly, the CFW treatment resulted in several fold increase in selective DAGs which

normally constitute only 4-5% of the total DAG pool. We observed a ~8-fold increase in C36:0

DAGs and ~6-fold increase in C36:1 DAG subspecies levels compared to the untreated *WT*

sample, while the other DAG species levels remained unaffected (**Fig. 2 A**). Thus, the data

clearly indicates that Pkc1 activation in yeast might require C36:0 and C36:1 DAG subspecies.

We have also measured TAG subspecies levels in the presence of cell wall stress by CFW and

observed that there is no depletion in the cognate TAG species suggesting that selective DAG

accumulation upon CW stress is not due to TAG lipolysis **(Fig. 2—fig. supplement 1B)**.

Due to the seemingly high selectivity of DAGs for Pkc1 signalling activation *in vivo*,

we sought to answer whether this specificity for C36:0 and C36:1 DAGs is engraved in the

DAG binding domain of Pkc1. Therefore, to unravel the specificity of Pkc1 to interact with

DAG across subspecies level, we performed an *in vitro* lipid binding assay. For this, we

purified the DAG-binding domain of yeast Pkc1, namely conserved region domain 1

(C1domain), expressed under the galactose-inducible promoter in yeast, with GFP at its C-

terminal **(Fig. 2—fig. supplement 2A).** Then, we performed liposome sedimentation assay

where we prepared liposomes containing phosphatidylcholine (PC) with DAGs of different

acyl chain compositions (DAGs were restricted to commercially available ones that are also

abundantly present in yeast. C36:1 DAG is commercially unavailable). After incubating

different liposomes with the protein, we separated both lipid-bound and unbound fractions and

probed with western blot **(Fig 2 B)**. Surprisingly, we observed binding only in the fraction

containing C36:0 liposomes, while there was no binding seen in any other DAG-containing

liposome **(Fig. 2 C)**. Furthermore, we also observed a concentration-dependent increase in the

binding of C1 domain with C36:0 DAGs **(Fig. 2 D and 2 F)**, confirming their direct physical

interaction. On the other hand, no effect on binding was seen even at the higher concentrations

of the other DAG species containing liposomes **(Fig. 2 E-F)**. This points to a clear specificity

of the C1 domain of yeast Pkc1 for the selective DAG subspecies acted upon by Dip2. A

negative control for GFP was also used to rule out any possibility of non-specific binding with

GFP **(Fig. 2—fig. supplement 2B).**

We further asked whether a similar kind of selectivity of Pkc1 C1 domain for Dip2-regulated

DAGs exists in higher eukaryotes as well. We cloned the C1 domain of one of the novel Pkc1s

of *Drosophila melanogaster* (PKC98E) and rat (PKCδ), as yeast Pkc1 knockout phenotype has

been shown to be rescued by mammalian novel PKCs (PKCδ) (Saiz-Baggetto et al., 2023),

suggesting functional homology between the two. After expressing and purifying the C1

domains **(Fig. 2—fig. supplement 3A)**, we performed the liposome sedimentation assay and

observed that PKC98E C1 domain binds the DAGs which were observed to be majorly

accumulated in Δ*Dmdip2* i.e., C32:0 and C34:1 DAGs (Mondal *et al*., 2022). Interestingly,

C36:0, which was not changed significantly on deleting *DIP2* in *Drosophila*, is also found to

bind the C1 domain of PKC 98E **(Fig. 2—fig. supplement 3B)**. This could possibly be due to

the difference in the *in vivo* and the *in vitro* DAG specificity of Dip2 and Pkc, respectively.

Since the lipidomic profile of the *Drosophila DIP2* deleted strain depicts the DAG changes

from its whole-body tissue, it fails to provide an accurate picture of tissue-specific DAG

specificity of Dip2, which would correspond to PKC.

Similarly, for rat PKCδ C1 domain, the liposome binding assay resulted in its binding

with all the three DAGs tested (C32:0, C34:1, C36:0) **(Fig. 2—fig. supplement 3C)** which

also corroborates our previous finding, where we observed the accumulation of multiple DAGs

on knocking-out *DIP2A* from mouse embryonic cells (Mondal *et al*., 2022). Though this

indicates a lack of DAG-specificity in Dip2 and PKC of higher eukaryotes, the presence of

multiple paralogs of Dip2 and PKC isoforms complicate the identification of clear DAG

selectivity. However, the correlation between the lipidomics data from Dip2A knockout mouse

embryonic cells and the liposome sedimentation assay for PKCδ further strengthens our

hypothesis that Dip2-regulated DAG species are involved in activation of PKCs across life

forms.

**The canonical DAG metabolism axis is not involved in Pkc1 signalling**

Since selective DAG-metabolism by Dip2 is associated with Pkc1 signalling, we asked whether manipulating the levels of bulk DAGs inside cell, mainly produced through canonical DAG metabolism can regulate Pkc1 signalling. DAG, being a central intermediary molecule in lipid metabolism, is acted upon by several conserved enzymes for phospholipid biosynthesis or storage lipid (TAGs) production. Majority of the enzymes participating in these processes are known to act on the bulk pool of DAGs, while the specificity for acyl-chain length remains to be elucidated (Li et al, 2020; Rockenfeller et al, 2018). Therefore, we probed two main TAG-forming enzymes, DAG acyltransferase (Dga1) and DAG transacylase (Lro1) along with the key membrane biogenesis enzyme, DAG kinase (Dgk1) for studying the effect of bulk DAG pool on Pkc1 activation and CWI pathway. Our lipidomic analysis of these gene knockouts confirmed that both Dga1 and Lro1 act on almost all the major DAG species and their absence led to ~2-10-fold accumulation of DAGs, consisting of varying chain lengths (**Fig. 3 A and 3 B**). *Δdgk1*, on the other hand, did not affect the steady-state DAG levels in yeast (**Fig. 3 C**), which is in agreement with earlier studies (Adeyo et al, 2011; Li et al., 2020). Surprisingly, none of the mutants of these bulk DAG-acting enzymes showed cell wall stress resistance in the presence of CFW (**Fig. 3 D**) or CR (**Fig. 3—fig. supplement 1**). Quantifying the pSlt2 level in these knockouts also resulted in no change, when compared to total Slt2 levels (**Fig. 3 E**).

To understand how Dip2 responds to the accumulation of bulk DAGs inside the cells, we probed the localization of Dip2 in the absence of *LRO1* and *DGA1*. Upon co-staining the cells with Mitotracker Red, we found that Dip2 remains in the mitochondria (**Fig. 3—fig. supplement 2**). This suggests that the bulk DAG accumulation does not affect Dip2 subcellular localization. This might also be due to the spatial segregation of the bulk DAG pool in the endoplasmic reticulum from the selective DAG pool, which is likely to be present in mitochondria or mitochondria-vacuole contact sites.

In addition, we also generated double deletion mutant of *DGA1* and *LRO1* and

quantified the DAG levels (**Fig. 3 F**). Spot assay for the double knockout shows that they are

sensitive to cell wall stress in the presence of CFW (**Fig. 3 G**) and in the presence of CR (**Fig.**

**3—fig. supplement 2B**). However, there was ~1.6-fold increase in pSlt2 level compared to

*WT*, which is half of what we observe in *Δdip2* (3-fold) (**Fig. 3 H**). Thus, despite having a ~2

to ~3.5-fold accumulation of total DAG pool in the absence of canonical enzymes, Pkc1

mediated CWI signalling remained unaffected. These observations establish that there is a

functional dichotomy in the cellular DAG pools: a metabolically active DAG pool that only

participates in lipid biogenesis activities and signalling DAG pool that activates Pkc1 signalling

cascade. Therefore, it can be proposed that the Dip2-mediated selective DAG axis has been

evolved to utilize a DAG subpopulation as a secondary messenger to facilitate the regulation

of PKC activation.

**Dip2-regulated DAG subspecies for Pkc1 activation originate from phosphatidylinositol**

**and its acyl-chain remodelling by Psi1**

Since canonical DAG metabolism pathway is decoupled from Pkc1 signalling, we

sought to identify the source of Dip2-regulated DAG subspecies involved in Pkc1 activation.

We probed major metabolic routes that can possibly contribute to specific DAG accumulation

in *Δdip2*. We treated *WT* and *Δdip2* with various inhibitors with the aim to cut off the supply

of DAGs from respective lipids (shown as schematic in (**Fig. 4 A**)) and checked for the

reduction in DAG accumulation. Interestingly, inhibition of $PI(4,5)P_2$ (hereafter referred to as

PIP2) to DAG conversion by U73122, a Phospholipase C (PLC) inhibitor (Banfic et al, 2013;

Jun et al, 2004), reduced the specific DAG levels by 80-90% in a concentration-dependent

manner (**Fig. 4 B**), while the other DAG species remained unaffected (**Fig. 4—fig. supplement**

**1A)**. On the other hand, blocking PA to DAG as well as ceramide to DAG synthesis using

propranolol and aureobasidin-A, respectively (Breslow et al, 2010; Morlock et al, 1991; Sasser et

al, 2012; Starr et al, 2016), did not show change in DAG levels **(Fig. 4—fig. supplement 1B-C)**.

To confirm the same, we also generated double knockout of *DIP2* and the major PA to DAG

producing enzyme genes in the cell, i.e., *PAH1* and checked for DAG accumulation. We found

that unlike what we observed in propranolol-treated *Δdip2* which showed no change in DAG

level, double knockout of *PAH1* and *DIP2* showed ~1.5-to-2-fold increase in all the bulk DAGs

(C32:1, C32:2, C34:1) **(Fig. 4—fig. supplement 2A)**. However, we also observed a significant

decrease of ~42% in C36:0 and ~81% in C36:1 DAGs. We also probed the effect of *PAH1*

deletion on CW stress pathway and Pkc1 activation. We found that *Δpah1* and *Δpah1Δdip2*,

both are defective in growth and sensitive to CW stress **(Fig. 4—fig. supplement 2B)**. This

phenotype does not correlate with pSlt2 level **(Fig. 4—fig. supplement 2C)**, suggesting that

the CW stress sensitivity is not linked to Pkc1 activation and may indicate a pleotropic defect

due to alteration in lipid metabolism. However, given the reduction in the level of selective

DAGs in *Δpah1Δdip2*, we cannot rule out the possibility that Pah1 could also be contributing

to the selective DAG accumulation in *Δdip2*. Given this observation, we also asked whether

the absence of Pah1 could affect subcellular localisation of Dip2. Therefore, we used

microscopy to trace Dip2 in *Δpah1* and found no change in its localization suggesting that Pah1

deletion has no impact on Dip2 localization **(Fig. 4—fig. supplement 2D-E)**.

We also deleted *PLC1 in Δdip2* and observed a ~75% reduction in C36:0, C36:1 DAGs

**(Fig. 4 C)**, with no change in other DAG species**,** suggesting that Plc1-dependent hydrolysis

of C36:0 and C36:1 PIP2 pools serves as the predominant source of Dip2-metabolised selective

DAGs. Whether the same source of DAGs is responsible for Pkc1 activation was investigated

by checking *pSlt2* levels. *Δplc1Δdip2* and U73122-treated *Δdip2* showed depleted levels of

*pSlt2*, when compared to *Δdip2*, while Slt2 phosphorylation remained unaffected in *Δplc1* **(Fig.**

**4 D)**. This suggested that the DAGs regulated by Dip2 are sourced via hydrolysis of corresponding PIP2 and are further involved in activating Pkc1 pathway. However, a further depletion of pSlt2 level in *Δplc1Δdip2*, compared to *Δplc1* was surprising as deletion of *DIP2* should not alter the level of pSlt2 level. This might have resulted from a rewiring of pathways upon deleting both the genes, which possibly led to a pleiotropic effect. These observations were also strengthened by the live-cell imaging of Plc1 that showed that Plc1 partially localizes to the mitochondria **(Fig. 4—fig. supplement 3A-B)** and vacuoles **(Fig. 4—fig. supplement 3C-D)**, similar to Dip2 suggesting that it might be producing the selective DAGs locally for Dip2 to act upon.

Next, we asked how exactly the specific PIP2 pool is maintained inside the cell to source these DAGs. Strikingly, a previous study has reported that the absence of an acyltransferase, phosphatidylinositol stearoyl incorporating 1 (Psi1), leads to severe depletion of C36:0 and C36:1 PIP2 species. Psi1 remodels the neo-synthesized phosphatidylinositol by adding stearic acid (C18:0) at the *sn-1* position of its glycerol backbone (Le Guedard et al, 2009), thus enriching the PI and eventually PIP2 pool with specific acyl chains (Doignon et al, 2016). Hence, we checked the DAG levels in *Δdip2* upon deleting *PSI1* (*Δpsi1Δdip2*) and observed a significant decrease in the accumulated DAGs (**Fig. 4E**). It is worth noting here that *PSI1* deletion in *Δdip2* background did not lead to complete depletion of accumulated DAGs in *Δdip2*, but a considerable quantity of DAGs, i.e., ~36% of C36:0 and 40% of C36:1, respectively are channelled from Psi1. In the single deletion of *PSI1*, unexpectedly, we observed increase in almost all the DAGs quantified **(Fig. 4—fig. supplement 4)**, which could be because of a compensation effect the system might have adapted due to decrease in C36:0, C36:1 phosphoinositide and respective DAGs, as reported earlier (Doignon et al., 2016). To test its role in Pkc1 activation, we measured pSlt2 levels in *Δpsi1Δdip2* and found that pSlt2 levels restored back to *WT* levels **(Fig. 4 F)**. In fact, deletion of *PSI1* itself decreases the pSlt2 level

in *WT* by ~35%, which phenocopies *Δpsi1Δdip2,* thereby strengthening the contribution of Psi1

to Pkc1 activation.

Together, these observations emphasize that the acyl chain remodelling of PI by Psi1

for specifically enriching C36:0 and C36:1 ensures channelling of these chain length

compositions towards respective PIP2. The corresponding PIP2 species are then hydrolysed to

selective DAGs by Plc1, which is responsible for Pkc1 activation. Dip2, which acts on these

DAGs, regulates Pkc1 signalling and keeps a check by converting these DAGs to TAGs (**Fig.**

**4 G**). In addition, these observations also reiterate that the bulk DAG pool cannot activate Pkc1

signalling cascade. Therefore, the Psi1-Plc1-Dip2 axis serves not only as a source of DAGs but

also acts as a critical checkpoint for the Pkc1 signalling cascade.

**Co-emergence and coevolution of Dip2-PKC axis underscores the selective nature**

**of DAG-based signalling**

DAGs are universally recognised by a structurally conserved domain called C1 domain

(conserved domain 1) (Hurley et al, 1997), found in different DAG effector proteins including

PKC. To understand the evolutionary relationship between DAG effectors and selective DAG

regulator, Dip2, we probed their distribution across the tree of life. Previously, we showed that

Dip2 had emerged in early opisthokonts. Interestingly, phylogenetic distribution of all known

DAG effectors (Colon-Gonzalez & Kazanietz, 2006) revealed that the majority of these are

concentrated in metazoans and metazoan sister groups (Choanoflagellates, Filasterea and

Ichthyosporea), while PKC is the only DAG effector conserved in fungal groups and all

opisthokonts **(Fig. 5 A)**.

Fungal Pkc1 (hereafter referred to as prototypical PKC) (Schmitz & Heinisch, 2003)

harbours all the typical regulatory domains present in higher eukaryotes, which includes HR1

(homology region 1) domain, C2 domain (conserved domain 2), C1 domain (conserved domain

1) along with the protein kinase domain (Schmitz & Heinisch, 2003). The phylogenetic

distribution suggest that the prototypical PKC had probably undergone multiple gene

duplication, domain shuffling and gene loss events during early metazoan evolution resulting

in a repertoire of PKC isoforms (**Fig. 5—fig. supplement 1A**). Furthermore, we observed the

presence of primitive PKC, consisting of only C1 and protein kinase domains in Microsporidia,

the sister group of Fungi (Bass et al, 2018; James et al, 2013). Interestingly, only two

*Dictyostelium* species of protists out of ~50 genomes (available in KEGG database) harbour

PKC in its primitive form (Goldberg et al, 2006; Keenan et al, 1997; Wang et al, 1996). Thus, the

early DAG effectors with C1 domain had evolved in one of the two major eukaryotic branches,

known as Unikonta (comprising protist, microsporidia, fungi and metazoa). To our surprise,

the other eukaryotic branch, Bikonta, comprising of plants, algae and SAR (**S**tramenopiles,

**A**lveolates, and **R**hizaria) organisms, is completely devoid of the DAG effectors (**Fig. 5 A**).

Overall, our analysis suggests that PKC is the first DAG effector to be recruited in eukaryotes

and evolved exclusively in opisthokonts.

The emergence of a prototypical Pkc with DAG-binding C1 domain in early

opisthokonts suggests potential requirement of DAG regulators for optimal signalling process.

Previously, we have reported that FAAL-like domains from bacteria gave rise to two tandem

FAAL-like domain containing protein, Dip2 across fungi and metazoans (Mondal et al., 2022;

Patil et al, 2021). Interestingly, the phylogenetic profile analysis of PKC (**Fig. 5—fig.**

**supplement 1B**) and Dip2 showed that both are distributed across Fungi and Animalia, but are

totally absent from Archaea, Bacteria, Plantae and Algae, suggesting that both the proteins

show evolutionary co-emergence and co-occurrence (**Fig. 5 B**). It is important to note here that

Dip2 is absent in a major phylum of fungi called Basidiomycota, suggesting a possible gene-

loss event in this phylogenetic branch. Although Basidiomycota harbours C1 domain

containing Pkc1, whether the Pkc1 is regulated by an unknown functional orthologue of Dip2,

or a distinct Pkc1 signalling has been evolved in this branch of fungi, is yet to be understood.

Taken together, our analysis underscores the fact that the emergence of a fully functional PKC

signalling pathway can be marked by the presence of Dip2 from fungi onwards.

To investigate the evolutionary history of Psi1-Plc1 axis, we examined the phylogenetic

distribution pattern and found that Psi1 was evolved in early eukaryotes and is conserved in

both Unikonta and Bikonta. On the other hand, Plc1, consisting of multiple domains such as

PH domain, EF hand domain, PI-PLC domain and C2 domain, emerged in protists and is

distributed across Unikonta, while absent in Bikonta **(Fig. 5—fig. supplement 1C)**. This

evolutionary history parallels that of the relevant DAG effector, PKC.

We then sought to quantitatively assess the phenomenon of co-evolution between Dip2

and PKC. To investigate this, we furthered our bioinformatic analysis using molecular

phylogenies of Dip2 and PKC from the representative fungi across fungal divisions (Ocana-

Pallares et al, 2022). Correlation between the phylogenetic distances (patristic distance) of Dip2

and PKC were calculated using multiple methods like Bayesian method, neighbour joining,

minimum-evolution, and maximum likelihood (Fourment & Gibbs, 2006). The result showed

significantly high association in the pairwise patristic distances of Dip2 and PKC, compared to

other control gene pairs like Pgk1 and Gapdh **(Fig. 5 C-D)**, suggesting their possible co-

evolution across fungal phylogenetic branches.

To calculate the interspecies correlation of genetic divergence rate, we adopted the

approach described for mirrortree algorithm (Edgar et al, 2012; Ochoa & Pazos, 2014; Pazos &

Valencia, 2001). In this case, genetic distances (distance matrix-based) were used to estimate

the evolutionary association between Dip2 and PKC (**Fig. 5E)**, which scored more than a

correlation coefficient of 0.8, an empirical cut-off value suggested by (Pazos & Valencia, 2001).

We further reconfirmed the above observation using a larger set of all available fungal genomes

(**Fig. 5—fig. supplement 2A)**. This observation is further substantiated by a lesser correlation

obtained on comparing Dip2 or PKC with a highly conserved gene like *GAPDH* (**Fig. 5E)**.

Furthermore, the correlation between *DIP2* or *GAPDH* and a diverse set of conserved genes

with related and unrelated functions were found to be below the cut-off value of correlation

coefficient (<0.8) (**Fig. 5F and fig. supplement 2B**). We also confirmed that the correlation

between Dip2-PKC pair is significantly different from this random control set (**Fig. 5—fig.**

**supplement 2C-D**). Therefore, a significant interspecies correlation between Dip2 and PKC

suggested that both the genes had experienced a similar evolutionary pressure, probably due to

their participation in the same physiological axis. Overall, the evolutionary analysis suggests

that Dip2 and PKC are phylogenetically correlated and have coevolved together for successful

establishment of selective DAG-based PKC signalling in Opisthokonta.

**Discussion:**

The discovery of Dip2 as a conserved and selective DAG regulator presented a cellular

scenario, where there are two 'functional' pools of DAG in cell, *viz*., the bulk 'metabolic' pools

of DAG and the minor pools of 'signalling' DAG. The former is involved in membrane

biogenesis and storage lipid biosynthesis, while the later is Dip2-regulated selective DAG pool,

responsible for activating Pkc1 signalling. Here, we provide evidence for the emergence of

Dip2 and the functional diversification of DAG pool, responsible for activation of Pkc1

signalling cascades in yeast. As DAG metabolizing enzymes are generally promiscuous, their

recruitment for selectively activating and fine-tuning the Pkc1 signalling events is a distant

possibility. The rooting of a lipid-based signalling in eukaryotes during early Opisthokonta

evolution possibly necessitated a more precise regulatory mechanism, making a species-

specific regulator such as Dip2 inevitable for DAG-based PKC signalling **(Fig. 6)**.

The metazoan PKC isoforms such as conventional and novel PKCs are well

characterized to be regulated by signalling molecules such as DAGs, calcium,

phosphatidylserine etc. (Rosse et al, 2010). Different combinations of regulatory domains in

PKC isoforms result in highly diversified and mutually exclusive functions in animals (Reyland,

2009). Interestingly, the single prototypical PKC of yeast with all the regulatory domains is

involved in a myriad of physiological processes, for example, cell wall organization,

progression of cell cycle, lipid metabolism, cell polarity, p-body assembly, septin organization

etc. (Dey et al, 2017). Initially, Pkc1 activation was shown to be unaffected by DAGs (Antonsson

et al, 1994), while a few recent reports have suggested the involvement of DAGs in activating

Pkc1 *in vitro* (Dey et al., 2017). We believe the discrepancy among the existing reports has been

cleared as we have observed a selective DAG-mediated Pkc1 and CWI pathway activation in

yeast. The earlier observed mutations in the putative DAG-binding C1 domain of yeast Pkc1

which results in decreased fitness in cell wall stress conditions (Jacoby et al, 1997), corroborates

our findings on the crucial role of DAG in activation of Pkc1-based CWI pathway. Similarly,

a differential role of other domains like HR1 has been proposed in earlier studies (Schmitz et al,

2002). Whether the selective DAG-based Pkc1 activation also requires HR1 domain to bind

Rho1, remains to be seen.

Recent reports have shown that exogenous addition of DAGs with varied acyl chains

have differential effect on the translocation of mammalian PKC isoforms (Schuhmacher et al.,

2020). While this indicates the role of distinct acyl chain-containing DAGs in differentially

activating PKC isoforms, it does not represent a physiological scenario where a cell employs

this. Here, we have shown selective increase in C36:0 and C36:1 DAGs which results in

activation of Pkc1 upon subjecting the yeast cells to cell wall stress. Additionally, the DAG

binding domain of Pkc1 shows binding exclusively with C36:0 DAG and not with other DAGs

*in vitro*. Hence, we have identified the specific DAG subspecies for regulating Pkc1, thereby,

bridging the gap in the understanding of selective DAG-driven PKC activation, that was

speculated thirty years ago (Marignani et al., 1996). Nonetheless, the rationale for selecting the

two DAG subspecies out of numerous possible combinations of acyl chains for activating Pkc1

remains an enigma and needs to be probed. It also opens up possibilities for specific roles of

selective species of lipids that perhaps explains why there is a preservation of diversity in

lipidome during evolution of different life forms.

The canonical DAG metabolizing enzymes, despite acting on C36:0, C36:1 DAGs to a

certain level, are excluded from their regulatory role in Pkc1 signalling. Although the double

deletion of *LRO1* and *DGA1* elevates Pkc1 activation to an extent, this increase is insufficient

to confer resistance to cell wall stress. Whether it is the distinct source of Dip2 regulated DAGs

from the PI pool or the spatial segregation of the two pools inside the cell, which make it

exclusive to Pkc1 pathway regulation needs to be probed. Furthermore, how Pkc1, which is a

bud-site (plasma membrane) protein in yeast (Denis & Cyert, 2005), is regulated by a selective

DAG regulator Dip2 residing at mitochondria-vacuole contact site (Mondal et al., 2022) requires

further investigation. Characterizing the spatiotemporal nature of specific DAG accumulation

that drives these processes would provide clues to the partitioning of Pkc1 activity and is

currently underway.

Although, PKC is a well-studied signalling proteins and DAG-PKC signalling has been

a textbook model for several decades, the evolutionary and regulatory origin of this pathway

remained unknown. Our current work provides compelling evidence that suggests that PKC

was the first DAG effector protein to be evolved and conserved in opisthokonts, while

completely absent in the other eukaryotic supergroup Bikonta which comprises plant, algae

and SAR organisms. The evolution of such distinct lipid-based signalling in early eukaryotic

ancestors is probably a decisive factor for further branching into two subgroups- Unikonta

(mainly Opisthokonta) and Bikonta. Thus, the work also suggests that the emergence of Dip2

around 1.2 billion years ago in Opisthokonta was a unique metabolic innovation to establish a

precisely regulated PKC signalling pathway (**Fig. 6**).

The role of Dip2 in fungal virulence has been highlighted in several studies as its

deletion in various plant and animal pathogenic fungi like *Magnaporthe oryzae* (causing rice

blast), *Cochliobolus heterostrophus* (causing southern corn blight) and *Coccidioides posadasii*

(causing valley fever) renders them avirulent (Lu et al, 2003; Narra et al, 2016; Wang et al, 2016).

As PKC signalling has also been shown to be crucial for pathogenicity of fungi, the

identification of Dip2 as a unique regulator of PKC would open avenues for therapeutic and

pharmacological interventions. Similarly, the phenotypes for PKC hyperactivation and deletion

of Dip2 observed independently in animals suggest a link between Dip2 and regulation of PKC

isoforms. For instance, the abnormalities observed in upregulation of different PKC isoforms

such as axonal bifurcation and outgrowth (Nitta et al, 2017; Zhang et al, 2019), altered dendritic

spine morphogenesis (Calabrese & Halpain, 2005; Ma et al, 2019) etc. phenocopy the defects

reported in Dip2 knockouts (Ma et al., 2019; Nitta et al., 2017).  Moreover, both Dip2A and PKC

have been implicated independently in autistic behaviour in model organisms (Liu et al, 2018;

Ma et al., 2019; Philippi et al, 2005). Similarly, loss of Dip2C, a candidate breast cancer gene (Li

et al, 2017), leading to increased epithelial-mesenchymal transition (EMT) in cancer cell lines

(Larsson et al, 2017), also resembles PKCε overexpression, which not only induces EMT but is

also a biomarker for aggressive breast cancer (Jain & Basu, 2014). These observations provide

a rationale to test the possibility that multiple paralogs of Dip2 might be regulating various

PKC isoforms by acting on specific pool of DAGs in higher eukaryotes.

## Materials and methods:

### Yeast strains and plasmids

Yeast strains used in this study are all BY4741 (MATa ura3Δ0 his3Δ1 leu2Δ0 met15Δ0) (Brachmann et al, 1998) derived from S288C genetic background as listed in (**Appendix 1 table 1**). Single gene knockout strains are retrieved from the public repository Euroscarf, except for *Δdip2* which is generated by homologous recombination from our previous study (Mondal et al., 2022).

Galactose promoter was replaced by the native Dip2 promoter in the plasmid pYSM10 used in our previous study using Gibson assembly cloning. This was also used as the template for site-directed mutagenesis by performing polymerase chain reaction using mutagenic oligonucleotides. These mutants were confirmed by sequencing.

Double knockout strains of *Δpsi1Δdip2* and *Δplc1Δdip2* were generated by knocking out *DIP2* from *Δpsi1* and *Δplc1* strains from Euroscarf library by homologous recombination. Briefly, *Dip2* (Cmr2; SGD ID: S000005619) gene was replaced with a PCR amplified hygromycin resistance cassette from pFA6A-hphMX6 plasmid. The primers were designed in such a way that the PCR product will have flanking sequences homologous to the 5′ and 3′ end of *DIP2* gene. The PCR product was purified and transformed using direct carrier DNA/PEG method-based transformations (Gietz & Schiestl, 2007). Transformed cells were plated on both hygromycin and G418 selection plate for *DIP2* and *PSI1/PLC1* deletion respectively. Thus, the resistant colonies were further validated for deletion using PCR primers amplifying the hygromycin cassette, designed using a web tool Primers-4-Yeast (Yofe & Schuldiner, 2014) as listed in (**Appendix 1 table 2**). Dip2 was tagged with GFP in the knockout strains of *LRO1* and *PAH1* by transforming amplicons produced by using Dip2 specific C-tag primers, using pFA6A-HisMX6 plasmid. For generating Dip2-GFP in *Δdga1*, we transformed Dip2 tagged with GFP under its native promoter in the double knockout of *DIP2* and *DGA1*. All the GFP

tagging was confirmed by western blot using anti-GFP antibody. For C1a-C1b protein

expression, C1a-C1b (413-534 aa) was amplified from yeast genomic DNA and cloned into

galactose inducible pYSM5 vector used in our previous study (Mondal et al., 2022).

**Media and reagents**

Yeast cells were grown in synthetic complete (SC) media, at 30°C in a shaking

incubator with 200 rotation per minute (RPM). SC media constituents include 1.7 g/l yeast-

nitrogen base with ammonium sulphate (BD Difco), 20 g/l glucose supplemented with

appropriate amino acids (Sigma), adenine (Sigma A5665) and uracil (Sigma U0750). G418

(200 µg/mL) and Hygromycin (0.2mg/mL) were used for knockout strain selection. For some

experiments, YP media (yeast extract – 10 g/l, peptone – 20 g/l) supplemented with 2%

dextrose was used. 2% bacto agar along with the above-mentioned SC media constituents was

used for solid media.

**Spot assay**

An overnight primary culture in SC media was diluted to 0.5 $OD_{600}$ nm and grown till

the secondary culture reaches $OD_{600}$ nm of ~2.0. This culture was diluted to 0.2 $OD_{600}$ nm and

was used as the first dilution, which was further diluted into multiple serial dilution stocks

($10^{-1}$, $10^{-2}$, $10^{-3}$, and $10^{-4}$). 10 µl of each dilution stocks were spotted sequentially on SC agar

control plate and SC agar with either Congo red (100 µg/mL) (Sigma) or Calcofluor white

(Sigma) (50 µg/mL) to induce cell wall stress or in combination with cercosporamide (2

µg/mL).

**Colony forming unit assay**

A primary culture was inoculated overnight and was diluted to 0.2 $OD_{600}$ nm in fresh

SC media, cultured till the early log phase or $OD_{600}$ nm of ~0.8. Cells were serially diluted to

0.002 $OD_{600}$ nm in autoclaved filtered Milli-Q water and 80µL of the cells were plated on SC

media agar as well as SC media agar with CR. After incubation for ~48 hours at 30°C, number

of colonies were counted and CFU was calculated.

**Growth curve analysis**

The overnight grown primary culture of *WT* and *Δdip2* was inoculated in fresh SC

media with ~0.2 $OD_{600}$ nm. Growth curve was performed using this secondary culture grown

with and without CR. The cell density was measured starting from an $OD_{600}$ nm of 0.2 at zero

time point to 28 hours. The readings were recorded at an interval of every 4 hours, until it

reaches the stationary phase.

**Protein expression and purification**

C1-GFP (413-534 amino acids) tagged with GFP and 8xHis at its C-terminus was expressed in

*S. cerevisiae*. Wherein, 8L cell culture grown in YP-Galactose media for induction was

harvested at 4000 RPM and was resuspended in buffer A (PBS pH 8.0, 500 mM NaCl, 10%

Glycerol and 5 mM β-mercaptoethanol), protease inhibitor cocktail (PIC), 1mM

phenylmethysulfonyl fluoride (PMSF) and lysed using 0.5 mm glass beads. For the expression

check, cell lysate was mixed with 6X-SDS loading dye and was run on 12% SDS gel and

checked for in-gel GFP fluorescence. The protein was purified by affinity chromatography with

nickel-nitrilotriacetic acid (Ni-NTA) agarose beads using buffer A. Protein was eluted by single-

step elution using Buffer A containing 250 mM imidazole. Following elution, the protein was buffer

exchanged in Buffer B (PBS, 150 mM NaCl, 10% Glycerol, 2mM DTT), concentrated and stored at -

80°C until further use.

The C198E (a.a. 171-304 of *Drosophila* PKC98E) and C1 delta (a.a. 152-280 of mouse PKC

delta) genes were cloned into pETite vector (Lucigen, USA) with a C-terminal TEV cleavage

site, GFP and 6x His tag. E. coli HI-control BL21(DE3) (Lucigen, USA) cells were used for

expression using Isopropyl β- d-1-thiogalactopyranoside (IPTG) induction-based

overexpression. Cells were grown at 37°C until the O.D. reached 0.6-0.8 and then induced

using 0.5mM IPTG and incubated at 18°C for 16 hours. After induction, the cells were

harvested at 7000rpm for 5 minutes at 4°C. Purification was done according to previously

published protocols (ref.). Briefly, the proteins were purified by immobilised Ni-NTA based

affinity chromatography in buffer containing 50mM Tris-HCl pH 8.0, 300mM NaCl, 0.4%

Triton X-100 and 1mM PMSF. The proteins were eluted using gradient elution between 10 and

250mM imidazole. Eluted proteins were further purified by size-exclusion chromatography

using Superdex 200 in buffer containing 20mM Tris-HCl pH 8.0, 150mM KCl. Fractions were

pooled, concentrated and stored at -80°C until further use.

**Liposome co-sedimentation assay**

Liposomes were prepared according to previously published (Dey et al., 2017; Larsson et al,

2020). Briefly, Lipids (800 nano moles of POPC (Avanti polar lipids) and 200 nano moles of

DAGs (Sigma) for 20 mole percent, were dissolved in chloroform: methanol (2:1 v/v), mixed

in glass tubes and dried to thin film under nitrogen gas stream. Following overnight desiccation,

1mL of Hydration buffer (50mM Tris-HCl pH 7.5, 150mM NaCl, 10mM $MgCl_2$, 1.7mM

$CaCl_2$, 10mM β-mercaptoethanol) was added to make 1mM final lipid concentration. The tubes

were vortexed vigorously and subjected to 10 freeze-thaw cycles using liquid nitrogen for 1min

and at 50°C water bath for 3 min.

C1-GFP (7nM) was incubated with liposomes for 10 minutes in ~~a total volume of~~ 300µl of

hydration buffer. After incubation, the reaction mixture was centrifuged at 100,000xg for 1

hour at 4°C. The supernatant was collected in a separate tube and both supernatant and pellet

were subjected to SDS-PAGE, followed by western blotting using Anti-GFP antibody (Cell

Signalling Technology). An in-house linker of 11 amino acids attached with 8xHis and GFP at

the C-terminal (DSLEFIASKLA-GFP) was purified and used to probe for GFP as a negative

control. Images were captured in BioRad ChemiDoc imaging system and analysed using

ImageJ.

**Western blotting**

Overnight grown yeast cells were inoculated in fresh YPD media (10 mL) at 0.2 $OD_{600}$

nm and grown till log phase (2-2.5 $OD_{600}$ nm). Cell pellets were harvested, washed with

Phosphate buffered saline (PBS) and flash frozen using liquid-nitrogen. The pellets were then

dissolved in 120 µl of Lysis Buffer (50 mM Tris-HCl [pH 7.5], 150 mM NaCl, 5 mM EDTA,

1% Nonidet P-40/Triton X-100, 1 mM Sodium Pyrophosphate, 1 mM Sodium Orthovanadate,

20 mM NaF, 1 mM Phenylmethylsulfonyl Fluoride, 1% Protease Inhibitor Cocktail (Nomura &

Inoue, 2015). The dissolved pellets were subjected to lysis with glass beads on a mechanical

vortex for 10 cycles of 30 s. The obtained homogenate was centrifuged at 12000 rpm for 10

min at 4˚C and then the supernatant was boiled with LaemmLi buffer for 15 min. Protein

extracts were separated on SDS-PAGE and then transferred on to PVDF membrane (Millipore)

using wet transfer at 100 V for 2 hours. Membranes were blocked using 5% BSA in Tris-

buffered saline with 0.1% Tween 20 detergent (TBST) and then incubated with appropriate

primary antibodies (Anti-Phospho-p44/42 MAPK (Erk1/2) (Thr202/Tyr204) Cell Signalling

Technology #9101) to measure pSlt2 and p44/42 MAPK (Erk1/2) Antibody Cell Signalling

Technology #9102)  (Sariki et al., 2019) to measure total Slt2 at 1:5000 dilution or TPI1 at

1:10000 dilution (gift from Dr. Palani Murgan's lab, CSIR-CCMB) followed by washing and

incubation with secondary antibody (Anti-rabbit IgG Cell Signalling Technologies #7074

(1:10000). After final wash with TBST, bands were visualised using SuperSignal™ West Pico

PLUS chemiluminescent horseradish peroxidase (HRP) substrate using BioRad Imaging

System. Images were analysed using ImageJ software. For cell wall stress, cells were incubated

with CR or CFW for 30 min before harvesting. For cercosporamide treatment, cells were

incubated for 30 min at a concentration of 5µg/mL followed by harvesting and sample

preparation.

**Lipidomics sample preparation**

Yeast cells were grown overnight in SC media and were diluted in 20 mL media (x6

replicates, total 120 mL media for single strain) to 0.2 $OD_{600}$ nm, cultured and harvested at ~2-

2.5 $OD_{600}$ nm. Inhibitor treated samples were cultured the same way, except that U73122

(0.5mM and 1mM) (Sigma), propranolol (1mM) (Sigma), aureobasidin A (0.025 µg/mL)

(Takara) were added while inoculating for the secondary culture at 0.2 OD. For cell wall stress,

cells were subjected to CFW (50µg/mL), 30 min before harvesting. Lipid isolation was

performed following modified-Folch method as reported previously (Abhyankar et al, 2018;

Kelkar et al, 2019; Kumar et al, 2019; Mondal et al., 2022; Pathak et al, 2018). Briefly, Cell pellets

were washed with PBS and flash frozen in liquid nitrogen. Cells were resuspended in 500 µl

of chilled PBS and were lysed using sonicator at 60% amplitude for 1 sec ON/ 3 sec OFF for

10 cycles with a probe sonicator on ice bed. 10 µl of lysate from each sample was collected for

protein estimation required for data normalization during lipid analysis. 250 µl of PBS was

added, followed by addition of $CHCl_3$ and methanol mix with respective internal standards to

achieve the ratio of 2:1:1 ($CHCl_3$: Methanol: PBS). Mixture was vortexed thoroughly (2 cycles

of 30 s) and was phase separated by centrifugation at 3000 RPM at RT. Bottom phase was

collected in fresh glass vials followed by addition of 100 μl of formic acid (~10%, vol/vol of

lysate volume) to the remaining top phase. After vortexing again, the same way, same volume

of $CHCl_3$, as mentioned in previous step, was added for re-extraction by phase separation.

Bottom phase was pooled with the already separated lipid in the first round. The total lipids

extracted were dried under stream of nitrogen at room temperature. Protein estimation was

performed using Bradford assay reagent (Sigma).

## Lipidome analysis by mass spectrometry

The lipidomic experiment and analysis of the isolated lipid species were performed

according to previously described protocols (Abhyankar et al., 2018; Kelkar et al., 2019;

Kumar et al., 2019; Mondal et al., 2022; Pathak et al., 2018). Semi-quantitative analysis of

lipids was performed using two mass spectrometers: An Agilent 6545 LC-QTOF (quadrupole-

time-of-flight) and a Sciex X500R QTOF employing high-resolution auto MS-MS methods

and multiple reaction monitoring high resolution (MRM-HR) scanning respectively. An

Electrospray ionization (ESI) source was used in both mass spectrometers. The dried lipid

extracts were re-solubilized in 200 μL of 2:1 CHCl3: MeOH and 10 μL was injected into the

mass spectrometers.

The liquid chromatography separation protocol was the same for both instruments. A

Luna C5 column (Phenomenex, 5 μm, 50 × 4.6 mm) coupled to a C5 guard column

(Phenomenex, 4 × 3 mm) was used for LC separation. The solvents used were buffer A: 95:5

H2O: MeOH + 0.1% Formic acid + 10mM ammonium formate and buffer B: 60:35:5 iPrOH:

MeOH: H2O + 0.1% Formic acid + 10mM ammonium formate. Methods were 30 minutes

long, starting with 0.3 mL/min 100% buffer A for 4 minutes, 0.5 mL/min linear gradient to

100% buffer B over 14 minutes, 0.5 mL/min 100% buffer B for 7 minutes, and equilibration

with 0.5 mL/min 100% buffer A for 5 minutes.

The following settings were used for the ESI-MS positive mode analysis on the Agilent

6545 LC-QTOF: drying gas and sheath gas temperature: 320 °C, drying gas and sheath gas

flow rate: 10L/min, fragmentor voltage: 150V, capillary voltage: 4000V, nebulizer (ion source

gas) pressure: 45 psi and nozzle voltage: 1000V. For analysis, a lipid library of DAGs and

TAGs was employed in the form of a Personal Compound Database Library (PCDL), and the

peaks were validated based on relative retention times and fragments obtained.

For the Sciex X500R QTOF, the following settings were used for the ESI-MS positive

mode analysis: source gas temperature: 400°C, spray voltage: 4500V, source gas 1 pressure:

40 psi, source gas 2 pressure: 50 psi. Peaks were quantified using Sciex OS, where the masses

of DAGs and TAGs were curated from the lipid maps structural database (LMSD).

All lipid species were quantified by normalizing areas under the curve to the protein

concentration of the lysate taken at the beginning.

**Live Cell Microscopy**

Endogenous GFP-tagged strains were grown in SC media at 30°C and were harvested

between 1.5-2 $OD_{600}$ nm (log phase). MitoTracker Red CMXRos (Invitrogen Inc) was added

to a final concentration of 50 nM in 1 mL of liquid cell culture and was incubated further for

30 min with shaking. Cells were then washed 5 times with PBS and was resuspended in 20µl

of PBS. 3µl of these cells were immobilized under agar bed on glass slides and visualized under

a Zeiss Apotome.2, inverted widefield fluorescence microscope equipped with a HAL 100

illuminator, Plan Apochromat ×100 oil objective (NA 1.4), and a AxioCam CCD camera.

Images were captured in DIC (Nomarski optics), mCherry (for MitoTracker Red) and FITC

fluorescence mode. All images were processed using ImageJ2 software.

**Phylogenetic profiling**

Sequences for phylogenetic profiling were collected from a stringent BLAST hit search

(Altschul et al, 1990) using the protein sequence from *Saccharomyces cerevisiae* as the query.

A defined list of organisms was made in order to account for the diversity in the fungal kingdom

(Ascomycota, Mucoromycota, Zoopagomycota, Blastocladiomycota, Chytridiomycota,

Holomycota) for all the proteins used for coevolution analysis. For the distribution of PKC

across tree of life, sequences were collected from the representative organisms from protists to

metazoans, consisting of novel, conventional and atypical PKCs. The domain organization of

the collected sequences was verified in Conserved Domains Database (CDD) (Lu et al, 2020) to

filter partial sequences. Multiple sequence alignment was then performed using MAFFT,

imposing E-INS-i model for multidomain proteins. IQ-TREE (Nguyen et al, 2015) was used to

generate the phylogenetic tree and iTOL (Letunic & Bork, 2021) was used to visualize the tree.

**Coevolution Analysis**

Interspecies correlation serves as a good indicator to predict functional relationship

between proteins. Other than *Dip2* and *PKC1*, a set of conserved housekeeping genes (*GAPDH,*

*PGK1, ACT1, HSP60, CYTC, TPI1, CDC42, RPL5*) and known lipid metabolizing enzymes

(*Dga1, Dgk1, Lro1, Tgl3, Faa1, Lcb1, Erg4*) were taken as control proteins. The obtained

alignment was checked in JalView and arranged alphabetically to maintain homogeneity in

comparison. This output alignment file was used to calculate pairwise genetic distances in

MEGA X using bootstrap (1000 repetitions), employing the JTT model (Jones-Taylor

Thornton model) and uniform mutation rates that represent the number of substitutions per site

between two homologous proteins.

The obtained genetic distances were exported as a distance matrix and Pearson correlation

coefficient was calculated using the following equation-

$$r = \frac{\sum_{i=1}^{n}(R_i-\bar{R})(S_i-\bar{S})}{\sqrt{\sum_{i=1}^{n}(R_i-\bar{R})^2}\sqrt{\sum_{i=1}^{n}(S_i-\bar{S})^2}}$$

The correlation coefficients were converted to normally distributed metric using Fisher's *r* to

*z* transformation,

$$r' = \frac{1}{2} ln \left| \frac{1+r}{1-r} \right|$$

where,

*r* = Pearson correlation coefficient

*r'* = Fisher-transformed correlation coefficient

These transformed coefficients (r′) were further compared to generate z score/z test statistic

using the given equation,

$$z = \frac{r'_1 - r'_2}{\sqrt{\frac{1}{N_1-3} + \frac{1}{N_2-3}}}$$

where,

$r'_1$ is the first Fisher-transformed correlation coefficient

$r'_2$ is the second Fisher-transformed correlation coefficient

$N_1$ and $N_2$ denote the number of common organisms in the first and second correlation

respectively. z-scores were calculated for all pairs where $r'_1$, representing Dip2-PKC correlation

coefficient and $r'_2$ representing correlation coefficient for Dip2 with other protein controls,

giving negative values for higher correlations and positive for lower ones. p-values were

calculated from the obtained z-scores.

Patristic distance is the sum of the lengths of the branches that link two terminal nodes in a

tree, denoting the divergence between the two nodes. Using the same collected sequences,

phylogenetic trees were constructed by standard methods- Bayesian Inference, Maximum

Likelihood (using JTT model), Neighbor Joining (NJ) and Minimum Evolution. Patristic

distances were calculated for each phylogenetic model using PATRISTIC software and

correlation coefficients were generated from the obtained distance matrices. Similar

phylogenetic patterns were observed where the homogeneity in correlation coefficient is

maintained. Basically, a higher correlation score represents a stronger relationship between the

rate at which the two proteins have evolved along the multiple branches of their phylogenetic

trees.

**Statistical analysis**

All the statistical analyses were performed in Microsoft Excel and GraphPad Prism

9.3.1 (471). Statistical analysis of the differences between two groups was performed using a

two-tailed, unpaired and parametric, Student's t-test. Error bars represent standard deviations

(SD), except for lipidomic analyses, where error bars are plotted as standard error of mean

(SEM). Significance of differences are marked based on the p-value obtained. $*p < 0.05$; $**p$

$< 0.01$; $***p < 0.001$; $****p < 0.0001$; ns, not significant. All the experiments are replicated

minimally thrice with technical replicates.

**Acknowledgement:**

We thank Dr. Krishnaveni Mishra (University of Hyderabad) and Dr. Purusharth I.

Rajyaguru (Indian Institute of Science) for sharing yeast knockout strains with us. We thank

Dr. Kaustuv Sanyal and Kuladeep Das (Jawaharlal Nehru Centre for Advanced Scientific

Research) for generating *Δpsi1Δdip2* strain for us. We thank Dr. Sriram Varahan (CSIR-

CCMB) for engaging discussions and sharing resources. We thank Saddam Shekh and Aakash

Chandramouli for technical assistance with the LCMS experiments at IISER Pune. SS thanks

University Grants Commission, India; SM and AC thank Council of Scientific and Industrial

Research (CSIR), India, for the research fellowship. RSN thanks healthcare theme projects-

Fundamental and Innovative CSIR in Science of Tomorrow (FIRST; MLP-0162) and Niche

Creation Project (NCP; MLP-0138) of CSIR, India; J.C. Bose Fellowship of SERB, India; and

Centre of Excellence Project of Department of Biotechnology, India. SSK thanks

Swarnajayanti Fellowship from the Science and Engineering Research Board (SERB),

Department of Science and Technology (DST), Government of India (Grant number

SB/SJF/2021-22/01) and the Department of Science & Technology–Funds for Improvement of

S&T Infrastructure Development (DST-FIST) (grant number SR/FST/LSII-043/2016) to the

Department of Biology, IISER Pune for setting up the Biological Mass Spectrometry Facility.

**References:**

1.  Abhyankar V, Kaduskar B, Kamat SS, Deobagkar D, Ratnaparkhi GS (2018) Drosophila
    DNA/RNA methyltransferase contributes to robust host defense in aging animals by regulating
    sphingolipid metabolism. *J Exp Biol* 221

2.  Adeyo O, Horn PJ, Lee S, Binns DD, Chandrahas A, Chapman KD, Goodman JM (2011) The
    yeast lipin orthologue Pah1p is important for biogenesis of lipid droplets. *J Cell Biol* 192: 1043-
    1055

3.  Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search
    tool. *J Mol Biol* 215: 403-410

4.  Antonsson B, Montessuit S, Friedli L, Payton MA, Paravicini G (1994) Protein kinase C in
    yeast. Characteristics of the Saccharomyces cerevisiae PKC1 gene product. *J Biol Chem* 269:
    16821-16828

5.  Banfic H, Bedalov A, York JD, Visnjic D (2013) Inositol pyrophosphates modulate S phase
    progression after pheromone-induced arrest in Saccharomyces cerevisiae. *J Biol Chem* 288:
    1717-1725

6.  Bass D, Czech L, Williams BAP, Berney C, Dunthorn M, Mahe F, Torruella G, Stentiford GD,

Williams TA (2018) Clarifying the Relationships between Microsporidia and Cryptomycota. *J*

*Eukaryot Microbiol* 65: 773-782

7.  Brachmann CB, Davies A, Cost GJ, Caputo E, Li J, Hieter P, Boeke JD (1998) Designer

deletion strains derived from Saccharomyces cerevisiae S288C: a useful set of strains and

plasmids for PCR-mediated gene disruption and other applications. *Yeast* 14: 115-132

8.  Breslow DK, Collins SR, Bodenmiller B, Aebersold R, Simons K, Shevchenko A, Ejsing CS,

Weissman JS (2010) Orm family proteins mediate sphingolipid homeostasis. *Nature* 463: 1048-

1053

9.  Calabrese B, Halpain S (2005) Essential role for the PKC target MARCKS in maintaining

dendritic spine morphology. *Neuron* 48: 77-90

10. Carrasco S, Merida I (2007) Diacylglycerol, when simplicity becomes complex. *Trends*

*Biochem Sci* 32: 27-36

11. Colon-Gonzalez F, Kazanietz MG (2006) C1 domains exposed: from diacylglycerol binding to

protein-protein interactions. *Biochim Biophys Acta* 1761: 827-837

12. Denis V, Cyert MS (2005) Molecular analysis reveals localization of Saccharomyces cerevisiae

protein kinase C to sites of polarized growth and Pkc1p targeting to the nucleus and mitotic

spindle. *Eukaryot Cell* 4: 36-45

13. Dey P, Su WM, Han GS, Carman GM (2017) Phosphorylation of lipid metabolic enzymes by

yeast protein kinase C requires phosphatidylserine and diacylglycerol. *J Lipid Res* 58: 742-751

14. Doignon F, Laquel P, Testet E, Tuphile K, Fouillen L, Bessoule JJ (2016) Requirement of

Phosphoinositides Containing Stearic Acid To Control Cell Polarity. *Mol Cell Biol* 36: 765-780

15. Edgar RS, Green EW, Zhao Y, van Ooijen G, Olmedo M, Qin X, Xu Y, Pan M, Valekunja UK,

Feeney KA *et al* (2012) Peroxiredoxins are conserved markers of circadian rhythms. *Nature*

485: 459-464

16. Eichmann TO, Lass A (2015) DAG tales: the multiple faces of diacylglycerol--stereochemistry,

metabolism, and signaling. *Cell Mol Life Sci* 72: 3931-3952

17. Etcheberrigaray R, Tan M, Dewachter I, Kuiperi C, Van der Auwera I, Wera S, Qiao L, Bank

813           B, Nelson TJ, Kozikowski AP *et al* (2004) Therapeutic effects of PKC activators in Alzheimer's

814           disease transgenic mice. *Proc Natl Acad Sci U S A* 101: 11141-11146

815    18. Fourment M, Gibbs MJ (2006) PATRISTIC: a program for calculating patristic distances and

816           graphically comparing the components of genetic change. *BMC Evol Biol* 6: 1

817    19. Gietz RD, Schiestl RH (2007) High-efficiency yeast transformation using the LiAc/SS carrier

818           DNA/PEG method. *Nat Protoc* 2: 31-34

819    20. Goldberg JM, Manning G, Liu A, Fey P, Pilcher KE, Xu Y, Smith JL (2006) The dictyostelium

820           kinome--analysis of the protein kinases from a simple model organism. *PLoS Genet* 2: e38

821    21. Heinisch JJ, Rodicio R (2018) Protein kinase C in fungi-more than just cell wall integrity.

822           *FEMS Microbiol Rev* 42

823    22. Hurley JH, Newton AC, Parker PJ, Blumberg PM, Nishizuka Y (1997) Taxonomy and function

824           of C1 protein kinase C homology domains. *Protein Sci* 6: 477-480

825    23. Jacoby JJ, Schmitz HP, Heinisch JJ (1997) Mutants affected in the putative diacylglycerol

826           binding site of yeast protein kinase C. *FEBS Lett* 417: 219-222

827    24. Jain K, Basu A (2014) Protein Kinase C-epsilon Promotes EMT in Breast Cancer. *Breast*

828           *Cancer (Auckl)* 8: 61-67

829    25. James TY, Pelin A, Bonen L, Ahrendt S, Sain D, Corradi N, Stajich JE (2013) Shared signatures

830           of parasitism and phylogenomics unite Cryptomycota and microsporidia. *Curr Biol* 23: 1548-

831           1553

832    26. Jun Y, Fratti RA, Wickner W (2004) Diacylglycerol and its formation by phospholipase C

833           regulate Rab- and SNARE-dependent yeast vacuole fusion. *J Biol Chem* 279: 53186-53195

834    27. Keenan C, Phillips P, Thio M, Pears C (1997) Protein kinase C in dictyostelium discoideum.

835           *Biochem Soc Trans* 25: S592

836    28. Kelkar DS, Ravikumar G, Mehendale N, Singh S, Joshi A, Sharma AK, Mhetre A, Rajendran

837           A, Chakrapani H, Kamat SS (2019) A chemical-genetic screen identifies ABHD12 as an

838           oxidized-phosphatidylserine lipase. *Nat Chem Biol* 15: 169-178

29. Kishimoto A, Takai Y, Mori T, Kikkawa U, Nishizuka Y (1980) Activation of calcium and

phospholipid-dependent protein kinase by diacylglycerol, its possible relation to

phosphatidylinositol turnover. *J Biol Chem* 255: 2273-2276

30. Kumar M, Ojha S, Rai P, Joshi A, Kamat SS, Mallik R (2019) Insulin activates intracellular

transport of lipid droplets to release triglycerides from the liver. *J Cell Biol* 218: 3697-3713

31. LaFayette SL, Collins C, Zaas AK, Schell WA, Betancourt-Quiroz M, Gunatilaka AA, Perfect

JR, Cowen LE (2010) PKC signaling regulates drug resistance of the fungal pathogen Candida

albicans via circuitry comprised of Mkc1, calcineurin, and Hsp90. *PLoS Pathog* 6: e1001069

32. Larsson C, Ali MA, Pandzic T, Lindroth AM, He L, Sjoblom T (2017) Loss of Dip2C in RKO

cells stimulates changes in DNA methylation and epithelial-mesenchymal transition. *BMC*

*Cancer* 17: 487

33. Larsson E, Hubert M, Lundmark R (2020) Analysis of Protein and Lipid Interactions Using

Liposome Co-sedimentation Assays. *Methods Mol Biol* 2169: 119-127

34. Le Guedard M, Bessoule JJ, Boyer V, Ayciriex S, Velours G, Kulik W, Ejsing CS, Shevchenko

A, Coulon D, Lessire R *et al* (2009) PSI1 is responsible for the stearic acid enrichment that is

characteristic of phosphatidylinositol in yeast. *FEBS J* 276: 6412-6424

35. Lee KS, Irie K, Gotoh Y, Watanabe Y, Araki H, Nishida E, Matsumoto K, Levin DE (1993) A

yeast mitogen-activated protein kinase homolog (Mpk1p) mediates signalling by protein kinase

C. *Mol Cell Biol* 13: 3067-3075

36. Levin DE (2005) Cell wall integrity signaling in Saccharomyces cerevisiae. *Microbiol Mol Biol*

*Rev* 69: 262-291

37. Li D, Yang SG, He CW, Zhang ZT, Liang Y, Li H, Zhu J, Su X, Gong Q, Xie Z (2020) Excess

diacylglycerol at the endoplasmic reticulum disrupts endomembrane homeostasis and

autophagy. *BMC Biol* 18: 107

38. Li J, Ping JL, Ma B, Chen YR, Li LQ (2017) Dip2C expression in breast cancer and its clinical

significance. *Pathol Res Pract* 213: 1394-1399

39. Liu T, Shi Y, Chan MTV, Peng G, Zhang Q, Sun X, Zhu Z, Xie Y, Sham KWY, Li J *et al*
(2018) Developmental protein kinase C hyper-activation results in microcephaly and behavioral
abnormalities in zebrafish. *Transl Psychiatry* 8: 232

40. Lu S, Wang J, Chitsaz F, Derbyshire MK, Geer RC, Gonzales NR, Gwadz M, Hurwitz DI,
Marchler GH, Song JS *et al* (2020) CDD/SPARCLE: the conserved domain database in 2020.
*Nucleic Acids Res* 48: D265-D268

41. Lu SW, Kroken S, Lee BN, Robbertse B, Churchill AC, Yoder OC, Turgeon BG (2003) A
novel class of gene controlling virulence in plant pathogenic ascomycete fungi. *Proc Natl Acad
Sci U S A* 100: 5980-5985

42. Ma J, Zhang LQ, He ZX, He XX, Wang YJ, Jian YL, Wang X, Zhang BB, Su C, Lu J *et al*
(2019) Autism candidate gene Dip2A regulates spine morphogenesis via acetylation of
cortactin. *PLoS Biol* 17: e3000461

43. Marignani PA, Epand RM, Sebaldt RJ (1996) Acyl chain dependence of diacylglycerol
activation of protein kinase C activity in vitro. *Biochem Biophys Res Commun* 225: 469-473

44. Marrocco V, Bogomolovas J, Ehler E, Dos Remedios CG, Yu J, Gao C, Lange S (2019) PKC
and PKN in heart disease. *J Mol Cell Cardiol* 128: 212-226

45. Milne SB, Ivanova PT, Armstrong MD, Myers DS, Lubarda J, Shulga YV, Topham MK,
Brown HA, Epand RM (2008) Dramatic differences in the roles in lipid metabolism of two
isoforms of diacylglycerol kinase. *Biochemistry* 47: 9372-9379

46. Mondal S, Kinatukara P, Singh S, Shambhavi S, Patil GS, Dubey N, Singh SH, Pal B, Shekar
PC, Kamat SS *et al* (2022) Dip2 is a unique regulator of diacylglycerol lipid homeostasis in
eukaryotes. *Elife* 11

47. Mori T, Takai Y, Yu B, Takahashi J, Nishizuka Y, Fujikura T (1982) Specificity of the fatty
acyl moieties of diacylglycerol for the activation of calcium-activated, phospholipid-dependent
protein kinase. *J Biochem* 91: 427-431

48. Morlock KR, McLaughlin JJ, Lin YP, Carman GM (1991) Phosphatidate phosphatase from
Saccharomyces cerevisiae. Isolation of 45- and 104-kDa forms of the enzyme that are
differentially regulated by inositol. *J Biol Chem* 266: 3586-3593

49. Narra HP, Shubitz LF, Mandel MA, Trinh HT, Griffin K, Buntzman AS, Frelinger JA, Galgiani
JN, Orbach MJ (2016) A Coccidioides posadasii CPS1 Deletion Mutant Is Avirulent and
Protects Mice from Lethal Infection. *Infect Immun* 84: 3007-3016

50. Newton AC (2018) Protein kinase C: perfectly balanced. *Crit Rev Biochem Mol Biol* 53: 208-
230

51. Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ (2015) IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol* 32: 268-
274

52. Nishizuka Y (1984) The role of protein kinase C in cell surface signal transduction and tumour
promotion. *Nature* 308: 693-698

53. Nitta Y, Yamazaki D, Sugie A, Hiroi M, Tabata T (2017) DISCO Interacting Protein 2 regulates
axonal bifurcation and guidance of Drosophila mushroom body neurons. *Dev Biol* 421: 233-
244

54. Nomura W, Inoue Y (2015) Methylglyoxal activates the target of rapamycin complex 2-protein
kinase C signaling pathway in Saccharomyces cerevisiae. *Mol Cell Biol* 35: 1269-1280

55. Ocana-Pallares E, Williams TA, Lopez-Escardo D, Arroyo AS, Pathmanathan JS, Bapteste E,
Tikhonenkov DV, Keeling PJ, Szollosi GJ, Ruiz-Trillo I (2022) Divergent genomic trajectories
predate the origin of animals and fungi. *Nature* 609: 747-753

56. Ochoa D, Pazos F (2014) Practical aspects of protein co-evolution. *Front Cell Dev Biol* 2: 14

57. Pathak D, Mehendale N, Singh S, Mallik R, Kamat SS (2018) Lipidomics Suggests a New Role
for Ceramide Synthase in Phagocytosis. *ACS Chem Biol* 13: 2280-2287

58. Patil GS, Kinatukara P, Mondal S, Shambhavi S, Patel KD, Pramanik S, Dubey N, Narasimhan
S, Madduri MK, Pal B *et al* (2021) A universal pocket in Fatty acyl-AMP ligases ensures
redirection of fatty acid pool away from Coenzyme A-based activation. *Elife* 10

59. Pazos F, Valencia A (2001) Similarity of phylogenetic trees as indicator of protein-protein
interaction. *Protein Eng* 14: 609-614

60. Philippi A, Roschmann E, Tores F, Lindenbaum P, Benajou A, Germain-Leclerc L, Marcaillou

C, Fontaine K, Vanpeene M, Roy S *et al* (2005) Haplotypes in the gene encoding protein kinase

c-beta (PRKCB1) on chromosome 16 are associated with autism. *Mol Psychiatry* 10: 950-960

61. Reyland ME (2009) Protein kinase C isoforms: Multi-functional regulators of cell life and

death. *Front Biosci (Landmark Ed)* 14: 2386-2399

62. Rockenfeller P, Smolnig M, Diessl J, Bashir M, Schmiedhofer V, Knittelfelder O, Ring J, Franz

J, Foessl I, Khan MJ *et al* (2018) Diacylglycerol triggers Rim101 pathway-dependent necrosis

in yeast: a model for lipotoxicity. *Cell Death Differ* 25: 767-783

63. Roncero C, Duran A (1985) Effect of Calcofluor white and Congo red on fungal cell wall

morphogenesis: in vivo activation of chitin polymerization. *J Bacteriol* 163: 1180-1185

64. Rosse C, Linch M, Kermorgant S, Cameron AJ, Boeckeler K, Parker PJ (2010) PKC and the

control of localized signal dynamics. *Nat Rev Mol Cell Biol* 11: 103-112

65. Sasser T, Qiu QS, Karunakaran S, Padolina M, Reyes A, Flood B, Smith S, Gonzales C, Fratti

RA (2012) Yeast lipin 1 orthologue pah1p regulates vacuole homeostasis and membrane fusion.

*J Biol Chem* 287: 2221-2236

66. Schmitz-Peiffer C, Biden TJ (2008) Protein kinase C function in muscle, liver, and beta-cells

and its therapeutic implications for type 2 diabetes. *Diabetes* 57: 1774-1783

67. Schmitz HP, Heinisch JJ (2003) Evolution, biochemistry and genetics of protein kinase C in

fungi. *Curr Genet* 43: 245-254

68. Schmitz HP, Lorberg A, Heinisch JJ (2002) Regulation of yeast protein kinase C activity by

interaction with the small GTPase Rho1p through its amino-terminal HR1 domain. *Mol*

*Microbiol* 44: 829-840

69. Schuhmacher M, Grasskamp AT, Barahtjan P, Wagner N, Lombardot B, Schuhmacher JS, Sala

P, Lohmann A, Henry I, Shevchenko A *et al* (2020) Live-cell lipid biochemistry reveals a role

of diacylglycerol side-chain composition for cellular lipid dynamics and protein affinities. *Proc*

*Natl Acad Sci U S A* 117: 7729-7738

70. Starr ML, Hurst LR, Fratti RA (2016) Phosphatidic Acid Sequesters Sec18p from cis-SNARE

Complexes to Inhibit Priming. *Traffic* 17: 1091-1109

71. Sussman A, Huss K, Chio LC, Heidler S, Shaw M, Ma D, Zhu G, Campbell RM, Park TS,

Kulanthaivel P *et al* (2004) Discovery of cercosporamide, a known antifungal natural product,

as a selective Pkc1 kinase inhibitor through high-throughput screening. *Eukaryot Cell* 3: 932-

943

72. Wang Y, He D, Chu Y, Zuo YS, Xu XW, Chen XL, Zhao WS, Zhang Y, Yang J, Peng YL

(2016) MoCps1 is important for conidiation, conidial morphology and virulence in

Magnaporthe oryzae. *Curr Genet* 62: 861-871

73. Wang Y, Rubin H, Ravid S (1996) Dictyostelium protein kinase C-delta-like protein is

localized in the cell nucleus. *Biol Cell* 86: 103-109

74. Yang C, Kazanietz MG (2003) Divergence and complexities in DAG signaling: looking beyond

PKC. *Trends Pharmacol Sci* 24: 602-608

75. Yofe I, Schuldiner M (2014) Primers-4-Yeast: a comprehensive web tool for planning primers

for Saccharomyces cerevisiae. *Yeast* 31: 77-80

76. Zhang B, Li Z, Zhang R, Hu Y, Jiang Y, Cao T, Wang J, Gong L, Ji L, Mu H *et al* (2019)

PKCgamma promotes axonal remodeling in the cortico-spinal tract via GSK3beta/beta-catenin

signaling after traumatic brain injury. *Sci Rep* 9: 17078

77. Kamiya, Y., Mizuno, S., Komenoi, S., Sakai, H., & Sakane, F. (2016). Activation of

conventional and novel protein kinase C isozymes by different diacylglycerol molecular

species. *Biochemistry and Biophysics Reports*, *7*, 361–366.

https://doi.org/10.1016/j.bbrep.2016.07.017

78. Letunic, I., & Bork, P. (2021). Interactive Tree Of Life (iTOL) v5: An online tool for

phylogenetic tree display and annotation. *Nucleic Acids Research*, *49*(W1), W293–W296.

https://doi.org/10.1093/nar/gkab301

79. Masayoshi, G., Sekiguchi, K., Nomura, H., Kikkawa, U., & Nishizuka, Y. (1987). Further

studies on the specificity of diacylglycerol for protein kinase C activation. *Biochemical and*

*Biophysical Research Communications*, *144*(2), 598–605. https://doi.org/10.1016/S0006-

291X(87)80008-6

80. Saiz-Baggetto, S., Dolz-Edo, L., Méndez, E., García-Bolufer, P., Marí, M., Bañó, M. C.,

Fariñas, I., Morante-Redolat, J. M., Igual, J. C., & Quilis, I. (2023). A Multimodel Study of the

Role of Novel PKC Isoforms in the DNA Integrity Checkpoint. *International Journal of*

*Molecular Sciences*, *24*(21), 15796. https://doi.org/10.3390/ijms242115796

81. Sariki, S. K., Kumawat, R., Singh, V., & Tomar, R. S. (2019). Flocculation of *Saccharomyces*

*cerevisiae* is dependent on activation of Slt2 and Rlm1 regulated by the cell wall integrity

pathway. *Molecular Microbiology*, *112*(4), 1350–1369. https://doi.org/10.1111/mmi.14375

**Main Figures:**

**Figure 1. Deletion of Dip2 leads to increased cell wall stress resistance.**

(**A**) A model depicting the possible role of Dip2 in regulating DAG species required for PKC activation which governs the CWI pathway in yeast by activating the downstream MAPK cascade. Activation of the MAPK cascade results in increased cell wall synthesis thereby strengthening the cell wall. Dip2 has been depicted in its three-domain architecture, harbouring DMAP-binding domain 1 (DBD1), tandem fatty acyl-AMP ligase-like domains (FLD1 and FLD2).

(**B**) Serial dilution assay for *WT*, two biological replicates of *Δdip2* (*Δdip2*_Colony1 and

*Δdip2*_Colony2; *Δdip2*_Colony1 has been used for further experiments). pDip2*::Δdip2*

represents *Δdip2*_Colony1 complemented with Dip2 expressed under its native promoter.

Control plate contains synthetic complete (SC) media and cell wall (CW) stress plate has either

congo red (CR) (100µg/mL) or calcofluor white (CFW) (50µg/mL). N=3.

(**C**) Colony forming units of *WT* and *Δdip2* grown in SC media control and cell wall stress

induced by CR (100µg/mL). Data are represented as mean ± SD (unpaired, two-tailed Student's

t-test; n = 3; ***$p < 0.001$; ns = not significant). N=3.

(**D**) Representative western blot showing pSlt2 (56 kDa) levels in *WT* and *Δdip2* compared to

the *WT* treated with CFW for 30 min, used as a positive control. Bar graph showing

quantification of fold change in the pSlt2 levels, normalised with total Slt2. Triose phosphate

isomerase (Tpi1) (27 kDa) has been used as a loading control. Data are represented as mean ±

SD (unpaired, two-tailed Student's t-test; $n > 3$; ***$p < 0.001$; ns = not significant.

(**E**) Serial dilution assay image in synthetic complete (SC) media control plate and in the

presence of congo red (100µg/mL) for *WT* and *Δdip2* compared with *Δdip2* complemented

with wildtype and catalytically inactive Dip2 mutants (Dip2$^{D523A}$ and Dip2$^{L687A}$), expressed in

a plasmid pYSM7, under its native promoter. N=3.

(**F**) Serial dilution assay for *WT* and *Δdip2* grown in SD media control and in the presence of

CR (100µg/mL) and cercosporamide (2µg/mL). N=3.

(**G**) Representative western blot for *WT*, *Δdip2* and cercosporamide treated *Δdip2* showing the

levels of pslt2 (56 kDa). Quantification of pSlt2 levels for *Δdip2* treated with cercosporamide

(5µg/mL) compared to *WT* and *Δdip2* control, normalised with total Slt2. Phosphoglycerate

kinase (Pgk1) (45 kDa) has been used as a loading control. Data are represented as mean ± SD

(unpaired, two-tailed Student's t-test; $N$=3; **p < 0.01; ns = not significant).

1027

The online version of this article includes the following source data and figure supplement(s)

for figure 1:

**Figure 1 - Source data 1-1B:** Spot assay for WT, *Δdip2* and pDip2*::Δdip2* under cell wall

stress.

**Figure 1 - Source data 2-1C:** Quantification of CFU for WT and *Δdip2* in the presence of CR.

**Figure 1 - Source data 3-1D:** Western blot and quantification for WT, *Δdip2* and WT treated

with CFW.

**Figure 1 - Source data 4-1E:** Spot assay for the catalytic mutants of Dip2

**Figure 1 - Source data 5-1F:** Spot assay for WT, *Δdip2* in the presence of CR and

cercosporamide.

**Figure 1 - Source data 6-1G:** Western blot and quantification for WT, *Δdip2* and *Δdip2* treated

with cercosporamide.

1040

1041

**Figure 2. Dip2-regulated selective DAG subspecies are associated with the activation of PKC signalling.**

(**A**) Lipidomic analysis of *WT* treated with CFW (50µg/mL) for 30 minutes. DAG intensities normalized to the total protein content in *WT*+CFW have been compared to the intensities from the same species in *WT* control sample. Data are represented as mean ± standard error of mean (SEM) (unpaired, two-tailed Student's t-test; N=6). ****$p < 0.0001$; ***$p < 0.001$ ns = not significant.

(**B**) A flowchart for the liposome sedimentation assay explaining the incubation of liposomes with protein, followed by ultracentrifugation and western blotting for supernatant (unbound protein) and pellet (bound protein) fraction.

(**C**) Representative western blot for liposome sedimentation assay with different DAGs showing supernatant fraction (unbound protein) and pellet fraction (liposome bound protein), probed for C1 domain using anti-GFP antibody (43k Da). N > 3.

(**D**) Western blot probing for C1 domain bound to different concentrations of C36:0 DAG containing liposomes (pellet fraction) and the unbound protein (supernatant fraction) using anti GFP antibody. Structure of respective DAGs is shown on the right side. N > 3.

**(E)** Western blot probing for C1 domain at different concentrations of C34:1 DAG containing liposomes in both pellet and supernatant fractions. Structure of respective DAGs have been shown on the right side. N > 3.

**(F)** Bar graph showing quantification of percentage binding of C1-GFP with increasing concentrations of specific (C36:0) and non-specific (C34:1) DAGs. Data are represented as mean ± SD (unpaired, two-tailed Student's t-test; *n*=3; ***p < 0.001; ns = not significant), N=3.

The online version of this article includes the following source data and figure supplement(s) for figure 2:

**Figure 2 - Source data 1- 2B:** Quantification of DAGs in WT and CFW treated WT.

**Figure 2 - Source data 2- 2C:** Western blot for C1-GFP incubated with different DAGs containing liposomes.

**Figure 2 - Source data 3- 2D:** Western blot for C1-GFP incubated C36:0 DAGs containing liposomes.

**Figure 2 - Source data 4- 2E:** Western blot for C1-GFP incubated C34:1 DAGs containing liposomes.

**Figure 2 - Source data 5- 2F:** Quantification for percent binding of C1 domain with C36:0 and C34:1 DAGs containing liposomes.

1079

**Figure 3. PKC activation is independent of bulk DAG metabolism pathway.**

(**A-C**) Lipidomic analysis of DAGs in *Δdga1*, *Δlro1* and *Δdgk1*, compared to *WT*. Fold change

values are mentioned below and are represented as green colour gradient. Data are represented

as mean ± SEM (unpaired, two-tailed Student's t-test; N = 5) ****p < 0.0001; ***p < 0.001;

**p < 0.01; *p < 0.05; ns = not significant.

(**D**) Serial dilution assay for single knockouts of bulk DAG acting enzymes (Dga1, Lro1 and

Dgk1) in the presence of CFW (50µg/mL), compared to SC media control plate. N=3.

(**E**) Representative western blot and quantification of pSlt2 (56 kDa) levels in deletion strains

of DAG metabolizing enzymes. Fold change has been quantified with respect to the total Slt2.

Triose phosphate isomerase (Tpi1) (27 kDa) has been used as a loading control. Data are

represented as mean ± SD (unpaired, two-tailed Student's t-test); N=4; ****$p < 0.0001$; *$p <$

0.05; ns = not significant.

(**F**) DAG subspecies quantification using lipidomics for the double knockout of *LRO1* and

*DGA1*, compared to *WT* and *Δdip2*. Data are represented as mean ± SEM (unpaired, two-tailed

Student's t-test); n = 6; ****$p < 0.0001$; ***$p < 0.00$; *$p < 0.05$; ns = not significant.

(**G**) Serial dilution assay image for the cell wall stress sensitivity of *Δdga1Δlro1,* compared to

*WT* and *Δdip2*, in the presence of CFW (50µg/mL), compared to SC media control plate. N=3.

SC media control image is reused in Figure 3- figure supplement 2.

(**H**) Representative western blot for pSlt2 (56 kDa) estimation in *Δdga1Δlro1,* compared to

total Slt2 and a loading control Glyceraldehyde-3-phosphate dehydrogenase (Gapdh) (36 kDa).

Fold change has been quantified with respect to the total Slt2. Data are represented as mean ±

SD (unpaired, two-tailed Student's t-test); N=3; **$p < 0.01$; ns = not significant.

1102

The online version of this article includes the following source data and figure supplement(s)

for figure 3:

**Figure 3 - Source data 1- 3A:** Quantification of DAG in *Δdga1*.

**Figure 3 - Source data 2- 3B:** Quantification of DAG in *Δlro1*.

**Figure 3 - Source data 3- 3C:** Quantification of DAG in *Δdgk1*.

**Figure 3 - Source data 4- 3D:** Spot assay for single knock outs of DAG acting enzymes.

**Figure 3 - Source data 5- 3E:** Western blots and quantification for single knockouts of DAG

acting enzymes.

**Figure 3 - Source data 6- 3F:** DAG quantification for double knockouts of *LRO1* and *DGA1*.

**Figure 3 - Source data 7- 3G:** Spot assay for double knockouts of *LRO1* and *DGA1 in* the

presence of CFW.

**Figure 3 - Source data 8- 3H:** Western blots and quantification of pSLt2 for double knockouts

of *LRO1* and *DGA1*.

1116

1117

1118

1119

1120

1121

1122

1123

1124

1125

**Figure 4. Psi1-Plc1-Dip2 triad axis regulates selective DAG subspecies to modulate PKC signalling**.

**(A)** Schematic showing various pathways (green arrows) that feed into the production of DAGs in the presence of various enzymes (shown in blue) and different chemical inhibitors (red) blocking the respective pathways.

(**B**) Quantification of selective DAGs in *WT* and *Δdip2* samples treated with two different

concentrations, i.e., 0.5 and 1 μM of U73122. Data are represented as mean ± SEM (unpaired,

two-tailed Student's t-test; N = 5). ****$p < 0.0001$; ***$p < 0.001$; ns = not significant.

(**C**) Specific DAG quantification using lipidomics for deletion of *Δplc1* and *Δplc1Δdip2*. Data

are represented as mean ± SEM (unpaired, two-tailed Student's t-test; n = 6). ****$p < 0.0001$;

***$p < 0.001$ ns = not significant. N=6.

(**D**) Representative western blot image and quantification of pSlt2 (56 kDa) levels for indicated

samples. Fold change has been quantified with respect to the total Slt2. Gapdh (36 kDa) has

been used as a loading control. Data are represented as mean ± SD (unpaired, two-tailed

Student's t-test; N>4; ****$p < 0.0001$; **$p < 0.01$; ns = not significant.

(**E**) Lipidomic analysis of specific DAGs in double knockout of *PSI1* and *DIP2*, compared with

*WT* and *Δpsi1*. Data are represented as mean ± SEM (unpaired, two-tailed Student's t-test; **$p$

$< 0.01$; *$p < 0.05$; ns = not significant. N = 5).

(**F**) Representative western blot image and quantification of pSlt2 (56 kDa) levels for indicated

samples. Two data points for controls (*WT* and *Δdip2*) are same as that of Figure 4D. Fold

change has been quantified with respect to the total Slt2. Phosphoglycerate kinase (Pgk1) (45

kDa) has been used as a loading control. Data are represented as mean ± SD (unpaired, two-

tailed Student's t-test; N>3; ****$p < 0.0001$; ***$p < 0.001$; ns = not significant.

(**G**) Schematic showing Psi1-Plc1-Dip2 axis to regulate Pkc1 signalling where Psi1 remodels

phosphoinositides (PI) to enrich it with C36:0 and C36:1 containing acyl chain, which is

channelled to the selective DAGs via Plc1 and in turn activate Pkc1.

1152

The online version of this article includes the following source data and figure supplement(s)

for figure 4:

**Figure 4 - Source data 1- 4B:** Quantification of DAG in WT and *Δdip2* in the presence of

Plc1 inhibitor.

**Figure 4 - Source data 2- 4C:** Quantification of DAG in double knockout of *DIP2* and *PLC1*.

**Figure 4 - Source data 3- 4D:** Western blot and quantification of pSlt2 in the double knockout

of *DIP2* and *PLC1* and in the presence of Plc1 inhibitor.

**Figure 4 - Source data 4- 4E:** Quantification of DAG in double knockout of *DIP2* and *PSI1*.

**Figure 4 - Source data 5- 4F:** Western blot and quantification of pSlt2 in the double knockout

of *DIP2* and *PSI1*.

1163

1164

1165

1166

1167

1168

**Figure 5. Dip2 and PKC share a parallel history of co-emergence coevolution across Opisthokonta.**

(**A**) Emergence and distribution of DAG effector proteins across tree of life. Green circles represent presence, while red circles represent absence of effector proteins. Yellow circle represents the primitive form of PKC. List of DAG effectors is taken from a previous study (Colon-Gonzalez & Kazanietz, 2006).

(**B**) Phylogenetic profiling of Dip2 and PKC shows their co-emergence in Opisthokonta. Inset shows presence of Dip2 and PKC in different fungal branches (Ocana-Pallares et al., 2022). Green circle represents presence, red represents absence, while yellow circle represents a primitive form of PKC harbouring only C1 and kinase domain.

(**C**) Interspecies coevolution plot (using patristic distance) between fungal Dip2 and PKC using three different phylogenetic algorithms, with the correlation coefficient value represented by r. Number of organisms and significance levels are indicated.

**(D)** Interspecies correlation coefficient values (r) for Dip2 compared with PKC and control

proteins Gapdh and Pgk using various algorithms for calculating phylogenetic distances. r

value is calculated for Gapdh and Pgk the same way. The table contains the exact correlation

coefficient values for all the above-mentioned protein pairs using different algorithms.

(**E**) Coevolution analysis (using genetic distance) between Dip2 and PKC along with their

individual comparison with control protein Gapdh. Number of organisms and significance

levels are indicated.

(**F**) Correlation coefficient values represented as bar graph for Dip2 compared with a set of

housekeeping genes and lipid metabolizing proteins.

1191

The online version of this article includes the following source data and figure supplement(s)

for figure 4:

**Figure 5 - Source data 1- 5C:** Quantification of interspecies correlation coefficient values for

Dip2 and PKC using different algorithms.

**Figure 5 - Source data 2- 5D:** Quantification of interspecies correlation coefficient values for

Dip2, PKC, Gapdh, and Pgk1 using different algorithms.

**Figure 5 - Source data 3- 5E:** Quantification of interspecies correlation coefficient values

using genetic distances for Dip2, PKC and Gapdh.

**Figure 5 - Source data 4- 5F:** Quantification of interspecies correlation coefficient values for

various housekeeping proteins and Dip2.

1202

1203

1204

1205

**Figure 6. A graphical representation depicting the emergence of Dip2-PKC axis and the**

**evolution of selective DAG-based PKC signalling in Opisthokonta.**

Dip2 is recruited parallel to PKC in the tree of life at the root of Opisthokonta evolution ~1.2

billion years ago. This Dip2-PKC axis remains conserved across Fungi and Metazoan. Two

distinct DAG pools are sourced from *de novo* or salvage pathways and acyl chain remodelling

of PIs, leading to metabolic and signalling DAG pool, respectively. Remodelled

phosphoinositides (PI, PIP, PIP2) by Psi1 are enriched with 18:0 acyl chains and channelled to

corresponding PIP2, which forms respective DAG species (C36:0, C36:1) upon hydrolysis by

Plc1. These selective DAG subspecies act as secondary messenger for PKC signalling and

therefore regulate the CWI pathway in yeast. Dip2 maintains the levels of these selective DAGs

by facilitating their conversion to TAGs, thereby creating a diversification of DAG function

into metabolic and signalling pathways.

1218

 **Supplementary Figures:**

1221 **Figure 1. figure supplement 1**

1222 Serial dilution assay in YPD media for WT, *Δdip2*, pDip2::*Δdip2* represents *Δdip2*
1223 complemented with Dip2 expressed under its native promoter. Control plate contains YPD
1224 media and cell wall (CW) stress plate has either congo red (CR) (100μg/mL) or calcofluor
1225 white (CFW) (50μg/mL). N=3.

1226

1227

1228 **Figure 1. figure supplement 2**

**(A)** Western blot image showing expression of wildtype Dip2-GFP-His-HA (219 kDa) and mutant Dip2 expressed in *Δdip2*. *WT* serves as a negative control and Pgk1(45 kDa) blot serves as the loading control.

**(B)** Representative western blot image showing pSlt2 (56 kDa) levels on complementing *Δdip2* with wildtype and catalytic mutant Dip2 (Dip2[D523A] and Dip2[L687A]). Quantification of pSlt2 levels for *WT* and mutant Dip2 complemented *Δdip2* strain, normalised with total Slt2. Phosphoglycerate kinase (Pgk1) (45 kDa) has been used as a loading control. Data are represented as mean ± SD (unpaired, two-tailed Student's t-test; *N*=3; *p < 0.05; ns = not significant).

**Figure 1. figure supplement 2B - Source data 1:** pSlt2 quantification for *Δdip2* complemented by the catalytic mutants of Dip2.

**Figure. 1. figure supplement 3**

**(A)** Microscopy image showing *WT* cells under normal condition (upper panel) and treated with calcofluor white (50µg/mL) (lower panel). Endogenous Dip2 has been tagged with

GFP, while mitochondria have been visualized by using a mitochondria-targeted mCherry

plasmid transformed in Dip2-GFP cells. The enlarged images of representative bud cells

for control and treated, marked with 1 and 2 respectively are shown in inset. Scale bar =

5µm.

**(B-C)** Line scan of signal intensity along the white line shown in inset images of (A), marked

1 and 2, respectively.

1252

1253 **Figure 1. figure supplement 3B - Source data 1**: Line scan intensity quantification for

1254 GFP and mCherry in Dip2-GFP control

1255 **Figure 1. figure supplement 3C- Source data 2**: Line scan intensity quantification for

1256 GFP and mCherry in Dip2-GFP CFW treated.

1257

1258

1259

**Figure 2. figure supplement 1**

**(A)** Representative western blot showing increasing activation of Pkc1 pathway (the level of

pSlt2) in a time-dependent manner upon CFW treatment (50µg/mL). Tpi1 (27 kDa) has been

used as a loading control.

**(B)** Lipidomic quantification of TAGs for *WT* cells under cell wall stress in the presence of

CFW (50µg/mL) compared to *WT* under normal growth condition.

1266

1267 **Figure 2. figure supplement 1B - Source data 1**: TAG quantification for WT and WT

1268 treated with CFW.

1269

1270

1271    **Figure 2. figure supplement 2**

1272    **(A)** Coomassie stained SDS-PAGE gel image of purified Pkc1-C1 domain, expressed and
1273        purified from yeast.

1274    **(B)** Western blot showing no binding (in the pellet fraction) in case of incubation of
1275        liposomes with only GFP as a negative control.

1276

1277

**Figure 2. figure supplement 3**

**(A)** SDS-PAGE gel stained with Coomassie showing purified C1 domain of *Drosophila*
*melanogaster* (C198E-GFP) and *Rattus novergicus* (C1δ-GFP).
**(B)**  Liposome binding assay for Pkc98E C1 domain from *Drosophila melanogaster* for multiple
DAGs, probed using anti-GFP antibody. Bar graph showing the percentage of DAG binding
of Pkc98E C1 domain for different DAG subspecies.
**(C)** Liposome binding assay for Pkcδ C1 domain from *Rattus novergicus* for multiple DAGs,
probed using anti-GFP antibody. Bar graph showing the percentage of DAG binding of Pkcδ
C1 domain for different DAG subspecies.
1287

1288    **Figure 2. figure supplement 3 - Source data 2**: Quantification for percent binding of
1289    Pkc98E and Pkcδ C1 domain with different DAG containing liposomes.

1290

1291

1292

**Figure 3. figure supplement 1**

(A) Serial dilution assay for single deletions of DAG acting enzyme genes (*DGA1*, *LRO1* and *DGK1*) in the presence of CR (100μg/ml)**.** N=3.

(B) Serial dilution assay for the cell wall stress sensitivity *of Δdga1Δlro1*, compared to *WT* and Δ*dip2*, in the presence of CR (100μg/mL), compared to SC media control plate. N=3. SC media control image is same as that of Fig 3G.

1299

1300

1301

**Figure 3. figure supplement 2**

Live cell microscopy images showing the DIC images (left), Dip2 tagged with GFP (green),

Mitotracker Red (red) and merged images for red and green channels (right) for **(A)** *WT*, **(C)**

*Δlro1* and **(E)** *Δdga1*. Scale bar represents 5 µm.

Line scan of signal intensity along the white lines shown in the respective merged images of

**(B)** *WT*, **(D)** *Δlro1* and **(F)** *Δdga1*. Green and red peaks represent Dip2-GFP and mitotracker

red, respectively.

1309

**Figure 3. figure supplement 2B - Source data 1**: Line scan intensity quantification for

Dip2-GFP and MitoTracker Red in WT control.

**Figure 3. figure supplement 3D- Source data 2**: Line scan intensity quantification for

Dip2-GFP and MitoTracker Red in *Δlro1*.

**Figure 3. figure supplement 3F- Source data 3**: Line scan intensity quantification for

Dip2-GFP and MitoTracker Red in *Δdga1*.

1316

1317
1318

1319

**Figure 4. figure supplement 1**

**(A)** Lipidomic analysis showing the levels of bulk DAGs in WT and *Δdip2* on treating with 0.5 and 1 μM of U73122. Data are represented as mean ± SEM (unpaired, two- tailed Student's t-test). N = 6.

**(B)** Lipidomic analysis for the DAG subspecies in *WT* and *Δdip2* in the presence and absence of Propranolol and **(C)** Aureobasidin. Data are represented as mean ± SEM (unpaired, two-tailed Student's t-test; ns = not significant). N = 6.

**Figure 4. figure supplement 1A- Source data 1**: DAG quantification for WT and *Δdip2* treated with U73122.

**Figure 4. figure supplement 1B- Source data 2**: DAG quantification for WT and *Δdip2* treated with propranolol.

**Figure 4. figure supplement 1C- Source data 3**: DAG quantification for WT and *Δdip2* treated with aureobasidinA.

1335

**Figure 4. figure supplement 2**

**(A)** Bar graph showing quantification of DAG subspecies level in *Δpah1* and *Δpah1Δdip2* DKO. Significance in red shows increase in DAG species while the black arrows show decrease in DAGs. Data are represented as mean ± SEM (unpaired, two-tailed Student's t-test; n = 6). ****p < 0.0001; ***p < 0.001 ns = not significant. N=6.

**(B)** Cell wall stress assay for *Δpah1* and *Δpah1Δdip2*, compared with *Δdip2,* in the presence of congo red (100μg/mL).

(C) Western blot and quantification of pSlt2 (56 kDa) in Δ*pah1* and Δ*pah1Δdip2*, normalized to
total Slt2 (56 kDa). Pgk1 (45 kDa) is used as a loading control. Data are represented as mean
± SD (unpaired, two-tailed Student's t-test; *n*=3; **p < 0.01; ns = not significant).

**(D)**       Live cell microscopy image for Dip2-GFP in *Δpah1* co-stained with Mitotracker Red
(red). Scale bar represents 5 µm.

**(E)**       Line scan of signal intensity along the white line shown in the merged image. Green
and red peaks represent Dip2-GFP and mitotracker red, respectively.

1350

**Figure 4. figure supplement 2A- Source data 4**: DAG quantification for *Δpah1 and*
*Δpah1Δdip2*.

**Figure 4. figure supplement 2C- Source data 5**: pSlt2 quantification for *Δpah1 and*
*Δpah1Δdip2*.

**Figure 4. figure supplement 2E- Source data 6**: Line scan intensity quantification for
Dip2-GFP and MitoTracker Red in *Δpah1*.

1357

1358

**Figure 4. figure supplement 3**

**(A)** Microscopy image panel showing DIC image (left), fluorescence for Plc1 tagged with GFP,
mitotracker-red for staining mitochondria and the merged image for both the fluorophores
(right). Scale bar represents 5 µm.

**(B)** Line scan plotted for the fluorescence intensity of Plc1-GFP along with mitotracker-red

across the line shown in the merged panel.

**(C)** Microscopy image panel for Plc1-GFP stained with FM4-64 for visualizing vacuoles. Th

merged image for red and green channel is sown on the right. Scale bar represents 5 µm.

**(D)** Fluorescence intensity plotted for GFP and FM4-64 along the line showing colocalization

of Plc1 with vacuole.

1369

1370   **Figure 4. figure supplement 3B- Source data 7**: Line scan intensity quantification for

1371 Plc1-GFP and MitoTracker Red.

1372   **Figure 4. figure supplement 3D- Source data 8**: Line scan intensity quantification for

1373 Plc1-GFP and FM4-64.

1374

1375

1376

1377   **Figure 4. figure supplement 4**

1378   DAG subspecies quantification for *Δpsi1* measured via lipidomics. Data are represented as

1379   mean ± SEM (unpaired, two-tailed Student's t-test; N=5). ****p < 0.0001; ***p < 0.001,

1380   **p < 0.01, ns = not significant.

1381   **Figure 4. figure supplement 4- Source data 9**: DAG quantification for *Δpsi1*.

1382

1383

1384**Figure 5. figure supplement 1**

1385**(A)** A schematic showing domain architecture and distribution of PKC isoforms (green circles)

1386 across Unikonta. Further classification of PKC isoforms (conventional, novel, atypical and

1387 Protein Kinase C-Related Kinase (PKN)) and domain annotations are indicated.

1388**(B)** Phylogenetic profiling of PKC showing its distribution among protists (grey), microsporidia

1389 (blue), fungi (yellow) and metazoan (green). Types of PKC isoforms found in metazoan are

1390 mentioned in bracket.

1391**(C)** Distribution of Psi1 and Plc1 and their domains across tree of life. Domain organizations of

1392 both Psi1 and Plc1 are shown on the top. Green and red colours represent the presence and

1393 absence of the proteins in respective supergroups mentioned on the left side.

**Figure 5. figure supplement 2**

(**A**) Interspecies correlation plot for Dip2 & Pkc1 and Gapdh with Pkc1 and Dip2 each from all the fungal sequences present in KEGG Genome Database.

(**B**) Correlation coefficient values represented as a bar graph for Gapdh compared to other protein controls. Pink colour window behind the bar graphs shows the empirical cut-off of 0.8 correlation coefficient.

(**C**) Interspecies correlation plot between Dip2 and various control proteins like Ribosomal 60s subunit protein L5 (Rpl5), Cytochrome C (CytC) and Actin.

(**D**) A table representing the correlation coefficient (r-value), Fisher's z-scores (z-score) and p-value for Dip2 compared with Pkc1 and other protein controls.

**Figure 5 - figure supplement 2A- Source data 1:** Quantification of interspecies correlation

coefficient values for Dip, PKC and Gapdh control for all fungal species.

**Figure 5 - figure supplement 2B- Source data 2:** Quantification of interspecies correlation

coefficient values for various housekeeping proteins with Gapdh.

**Figure 5 - figure supplement 2C- Source data 3:** Quantification of interspecies correlation

coefficient values for Dip compared with Rpl5, CytC and Actin.

**Figure 5 - figure supplement 2D- Source data 4:** Quantification of interspecies correlation

coefficient values, Fisher's z-scores (z-score) and p-value for Dip2 compared with Pkc1 and

other protein control.

1415

1416

1417

1418

1419