Citalopram exhibits immune-dependent anti-tumor effects by modulating C5aR1[+] TAMs and CD8[+] T cells

Fangyuan Dong[1, 3, 4, 5, 11], Shan Zhang[2, 11], Kaiyuan Song[6, 11], Luju Jiang[2, 11], Li-Peng Hu[2], Qing Li[2], Xue-Li Zhang[2], Jun Li[2], Mingxuan Feng[7], Zhi-Wei Cai[8], Hong-Fei Yao[8], Rong-Kun Li[9], Hui Li[2], Jie Chen[1, 3, 4, 5], Xiaona Hu[1, 3, 4, 5], Jiaofeng Wang[1, 3, 4, 5], Chongyi Jiang[8], Helen He Zhu[10], Cun Wang[2], Lin-Tai Da[6], Zhi-Gang Zhang[2, #], Zhijun Bao[1, 3, 4, 5, #], Xu Wang[9, #], Shu-Heng Jiang[2, #]

[1]Department of Gastroenterology, Huadong Hospital, Shanghai Medical College, Fudan University, Shanghai 200040, P.R. China

[2]State Key Laboratory of Systems Medicine for Cancer, Shanghai Cancer Institute, Ren Ji Hospital, School of Medicine, Shanghai Jiao Tong University, Shanghai 200240, P.R. China

[3]Shanghai Key Laboratory of Clinical Geriatric Medicine, Shanghai 200040, P.R. China

[4]National Clinical Research Center for Aging and Medicine, Shanghai 200040, P.R. China

[5]Department of Geriatrics, Huadong Hospital, Shanghai Medical College, Fudan University Shanghai 200040, P.R. China

[6]Key Laboratory of Systems Biomedicine (Ministry of Education), Shanghai Center for Systems Biomedicine, Shanghai Jiao Tong University, Shanghai 200240, China

[7]Department of Liver Surgery, Ren Ji Hospital, School of Medicine, Shanghai Jiao Tong University, Shanghai 200127, P.R. China

[8]Department of General Surgery, Hepato-biliary-pancreatic Center, Huadong Hospital, Fudan University, Shanghai 200040, P.R. China

[9]Institute of Oncology, Affiliated Hospital of Jiangsu University, Zhenjiang 212001, P.R. China

[10]State Key Laboratory of Systems Medicine for Cancer, Renji-Med-X Stem Cell Research Center, Shanghai Cancer Institute & Department of Urology, Ren Ji Hospital, Shanghai Jiao Tong University School of Medicine, Shanghai, 200127, China;

[11]These authors contributed equally to this work

30   #Correspondence

31   shjiang@shsci.org (S.H.J.); zhijunbao@fudan.edu.cn (Z.B.); wangxu@ujs.edu.cn (X.W.);

32   zzhang@shsci.org (Z.G.Z).

Abstract

Administration of selective serotonin reuptake inhibitors (SSRIs) is associated with a reduced cancer risk and shows significant anti-tumor effects across multiple tumor types, suggesting the potential for repurposing SSRIs in cancer therapy. Nonetheless, the specific molecular target and mechanism of action of SSRIs remain to be fully elucidated. Here, we reveal that citalopram exerts an immune-dependent anti-tumor effects in hepatocellular carcinoma (HCC). Interestingly, the anti-HCC effects of citalopram are not reliant on its conventional target, the serotonin transporter. Through various drug repurposing approaches, including global reverse gene expression profiling, drug affinity responsive target stability assay, and molecular docking, the complement component 5a receptor 1 (C5aR1) is identified as a new target of citalopram. C5aR1 is predominantly expressed by tumor-associated macrophages (TAMs), and citalopram treatment enhances local macrophage phagocytosis and elicits $CD8^+$ T anti-tumor immunity. At a systemic level, citalopram treatment results in decreased serum 5-HT levels, which correlates with increased activation of $CD8^+$ T cells. Additionally, citalopram promotes αCD3/αCD28-stimulated $CD8^+$ T cell activation, expansion, and cytotoxicity. Depletion of $CD8^+$ T cells partially hinders the anti-HCC effects of citalopram. Collectively, our study reveals multiple immunomodulatory roles of citalopram in inducing anti-tumor immunity, and provides a basis for considering the repurposing of SSRIs as promising anticancer agents for HCC treatment.

Key words: Drug repurposing, Drug discovery, SLC6A4, SERT, Anti-tumor immunity

Introduction

The development of anti-tumor drugs encounters numerous challenges, such as a high failure rate, substantial costs, limited bioavailability, safety concerns, and lengthy design and testing procedures [1]. As different diseases may share common pathological mechanisms, there is a growing focus on repurposing existing drugs for new applications, referred to as "new uses for old drugs" [2, 3]. Additionally, exploring new disease-relevant targets for established drugs and broadening their indications offers significant translational value [4].

Selective serotonin reuptake inhibitors (SSRIs; i.e., citalopram, escitalopram, fluoxetine, fluvoxamine, paroxetine, and sertraline) are frequently prescribed to treat conditions such as depression, anxiety, insomnia, and chronic pain [5]. SSRIs target the serotonin transporter (SERT, encoded by the *SLC6A4* gene), blocking the reuptake of serotonin (5-hydroxytryptamine, 5-HT) into the presynaptic neuron, ultimately increasing 5-HT levels in the brain [6]. Recent population-based epidemiological studies suggest that SSRIs have the potential to reduce the risk of many cancers, including kidney, breast, colorectal, and liver cancers [7-10]. The anxiety and depression symptoms among cancer patients is increasingly recognized [11]. Considering the established safety profile and known side effects of SSRIs, repurposing these drugs as potential anti-tumor agents, particularly for cancer patients with comorbid depression, could offer dual benefits. This approach may not only help improve neurological function but also potentially impede tumor progression. Hence, repurposing SSRIs in this context could present a cost-effective option. Nevertheless, the mechanisms of actions responsible for the anti-tumor effects of SSRIs are still largely unknown.

In this study, using hepatocellular carcinoma (HCC) as a model, we revealed that citalopram, an SSRI, exhibits anti-tumor properties not through its traditional target SERT, but instead via partially immune-dependent mechanisms. Through a comprehensive approach involving global reverse gene expression profiling, drug affinity responsive target stability (DARTS) [12], and molecular docking analyses, we identified the complement component 5a receptor 1 (C5aR1), predominantly expressed by tumor-associated macrophages (TAMs), as a target of citalopram. C5aR1 play critical roles in fostering an immunosuppressive microenvironment [13, 14] and boosting the metastatic potential of cancer cells [15]. Given its significant impact on cancer progression, C5aR1 has surfaced as a promising therapeutic

target for potential cancer interventions. Here, we further elucidated the mechanism underlying the connection between C5aR1$^+$ TAMs and citalopram-related phenotypes. Additionally, after citalopram treatment, a decline in serum 5-HT levels was observed, which is linked to the activation of CD8$^+$ T cells. More importantly, citalopram directly stimulates CD8$^+$ T cell activation, expansion, and cytotoxicity under *in vitro* conditions. Collectively, the anti-tumor effects of citalopram in HCC involve multiple immune-dependent mechanisms.

Results

The immune-dependent anti-tumor effects of citalopram in HCC

To evaluate the therapeutic effects of citalopram, we engrafted mouse Hepa1-6 and Hep53.4 cells into different strains of recipient mice, including the immune-competent C57BL/6 mice and immune-compromised Rag1$^{-/-}$ mice. Citalopram was given by i.p. administration daily for 30 days after bearing palpable tumors (~50 mm^3). Tumor growth was significantly retarded in both two mouse models. Of note, this impairment was more pronounced in immune-competent C57BL/6 mice, indicating an immune-dependent effect of citalopram (Figures 1A-B). This finding was further validated at molecular levels, as evidenced by reduced staining of the proliferation index Ki67 and increased expression of cleaved caspase 3 (CCS3) (Figure S1).

As SERT is the known target of SSRIs, we tested the therapeutic effect of citalopram in the absence of SERT (Figures 1C-D). In mouse HCC cells, SERT knockdown failed to affect cell proliferation or cell apoptosis *in vitro* and *in vivo* (Figures 1E-F and S2A-B). However, citalopram delayed the growth of SERT-silenced HCC tumors, and its suppressive effect was more proficient in immune-competent hosts compared with Rag1$^{-/-}$ ones (Figures 1E-F and S2C-F). Together, these data suggest that citalopram might target other molecules instead of SERT to exhibit an immune-dependent anti-tumor effects in HCC.

C5aR1 is a direct target of citalopram

Previously, using drug-induced gene signature networks and target prediction method, we identified a repertoire of targets related to citalopram treatment [16]. The top two hits are

glucose transporter 1 (GLUT1) and C5aR1 (Figures 2A-B). Citalopram can reverse the

Warburg effect to inhibit HCC by targeting GLUT1 directly [16]. Considering the significant role

of C5aR1 in immune modulation [17], we investigated whether the immune-dependent anti-

tumor effects of citalopram are mediated through C5aR1. Immunohistochemical analysis

showed that the positive staining of C5aR1 was predominantly found in scattered immune

cells (Figure 2C). Single-cell sequencing data indicated that C5aR1 was expressed at higher

levels in monocytes/macrophages within the tumor microenvironment (TME) compared with

other cell types (Figure 2D). Co-immunofluorescence analysi corroborated that CD163$^+$

macrophages presented significant C5aR1 immunoreactivity (Figure 2E).

The DARTS assay is widely recognized technique used to identify potential protein targets

for small molecules. This approach involves conferring proteolytic protection of target proteins

by their interaction with small molecules [18]. Combined DARTS assay with immunoblotting

analysis, we investigated the stability of C5aR1 against pronase in the presence or absence

of citalopram treatment. Because C5aR1 is expressed by macrophages, THP-1 cells were

employed for the DARTS assay. Citalopram did not affect the basal protein level of C5aR1 in

macrophages (Figure 2F). However, citalopram could enhance the stability of C5aR1 against

pronase treatment. Of note, this effect was specific to C5aR1, as citalopram did not impact

the proteolytic susceptibility of β-actin (Figures 2F). Moreover, the stability of C5aR1 against

pronase was enhanced in a dose-dependent manner following treatment with citalopram

(Figures 2G). Besides citalopram, other 5 SSRIs (escitalopram, fluoxetine, fluvoxamine,

paroxetine, and sertraline) also played protective roles of C5aR1 against pronase (Figure

S3A).

To address citalopram/C5aR1 interaction, *in silico* docking analysis was conducted.

Previous studies revealed two different inhibitor-binding sites in C5aR1, namely, the

orthosteric site on the extracellular side and the allosteric site outside the transmembrane

bundle [19]. Therefore, we docked the structure of citalopram into both orthosteric and allosteric

sites of human C5aR1 (PDB id: 6C1Q), respectively. *In silico* docking of citalopram into the

orthosteric site predicted two binding conformations without overlap (Figures 2H and S3B). In

pose-1, the fluorophenyl and the cyanophtalane group of citalopram stacked with the side

chains of Y258 and R206, and the amino group of citalopram interacted with the negatively

charged residue E199 (Figure 2I, left). In pose-2, two negatively charged residues, D282 and D37, surrounded the amino group (Figure 2I, middle). Considering the accessibility, D282 orienting towards the binding cavity of C5aR1 might play a more pivotal role in the citalopram binding event proposed by pose-2. Additionally, the cyanophtalane group of citalopram interacted with hydrophobic residues L92 and P113 of C5aR1. The fluorine was inserted into a narrow space lying over the side chain of W102 (Figure 2I, right). For the allosteric site, residue W213 played a critical role in inhibitor binding via hydrogen-bond interactions established by the amino group in the indole ring [20]. According to the best pose from cluster 2, the citalopram formed a hydrogen-bond via the oxygen atom in the cyanophtalane group with W213 (Figure 2I, right). Inspired by these molecular docking results, we generated six C5aR1 mutants to figure out the real binding site of citalopram in HEK293T cells (Figure S3C). Indeed, citalopram had no protective effects on C5aR1 stability with the transfection of HA-C5aR1 E199 and D282 mutant against pronase (Figure 2J). Consistently, mutations on E199 and D282 reduce C5a binding affinity to C5aR1 [21]. Furthermore, docking analysis and DARTS assay revealed that other four SSRIs could bind to the D282 site of C5aR1 (Figures S3D-E). These findings demonstrate that SSRIs adopt the orthosteric site to bind to C5aR1 and electrostatic interactions between C5aR1 D282 and the amino groups of SSRIs play an instrumental role in the binding mechanisms.

Citalopram targets C5aR1 in tumor-associated macrophages (TAMs)

C5aR1, the receptor for anaphylatoxin C5a, is profoundly implicated in tumor progression via inducing an immunosuppressive microenvironment in which TAMs are involved [14]. As C5aR1 is localized in TAMs within the TME of HCC, we explored whether citalopram exerted anti-tumor functions by targeting C5aR1 on TAMs. Given that citalopram also targets GLUT1 in HCC [16], we generated a Hepa1-6 clone with complete knockdown of GLUT1 (GLUT1^{KD}) to dissect C5aR1-dependent roles (Figure 3A). Interestingly, citalopram displayed an inhibitory effect on GLUT1^{KD} tumors in immune-component C57BL/6 mice but not Rag1$^{-/-}$ ones (Figure 3B), underscoring the contribution of C5aR1. To delineate the involvement of macrophages, we first depleted mouse macrophages with clodro-Liposomes and found that citalopram was ineffective in controlling GLUT1^{KD} tumors (Figure S4), implying the influences of citalopram on

macrophages. As the second line of evidence, we engrafted GLUT1^{KD} Hepa1-6 cells into the

syngeneic *C5ar1$^{+/-}$* and *C5ar1$^{-/-}$* mice. Compared with *C5ar1$^{+/-}$* recipients, tumor growth was

attenuated in *C5ar1$^{-/-}$* recipients (Figure 3C), whereas C5a deposition remained unchanged

(Figure 3D). Third, we co-injected *C5ar1$^{+/-}$* versus *C5ar1$^{-/-}$* bone marrow-derived macrophages

(BMDMs) with GLUT1^{KD} Hepa1-6 cells into syngeneic recipient *C5ar1$^{-/-}$* mice to analyze the

therapeutic effect of citalopram. Implantation of *C5ar1$^{+/-}$* BMDMs into *C5ar1$^{-/-}$* recipients

restored tumor growth compared with their *C5ar1$^{-/-}$* counterparts and citalopram cannot

restrict tumor growth in r*C5ar1$^{-/-}$*; d*C5ar1$^{-/-}$* mice; in contrast, tumor growth kinetics were

considerably diminished in r*C5ar1$^{-/-}$*; d*C5ar1$^{+/-}$* mice upon citalopram treatment (Figure 3E),

suggesting that C5aR1^{+} TAM rewiring enables the tumoricidal effects of citalopram. Notably,

C5a deposition was not affected by citalopram treatment (Figure 3F).

To answer how C5aR1^{+} macrophages mediate citalopram-induced anti-tumor effects, we

probed macrophage functions. Phagocytosis assay showed that isolated TAMs from *C5ar1$^{-/-}$*

recipients had higher phagocytic capacity of HCC cells than that of *C5ar1$^{+/-}$* recipients (Figure

3G), and citalopram increased the phagocytosis of TAMs from r*C5ar1$^{-/-}$*; d*C5ar1$^{+/-}$* mice

(Figure 3H). Moreover, C5a-mediated macrophage phagocytic impairment was reversed by

C5aR1 knockdown or treatment with citalopram or different SSRIs (Figures S5A-C). C5a was

able to suppress, while citalopram was effective in bolstering phagocytosis in macrophages

expressing C5aR1^{WT} but not C5aR1^{D282A} (Figures S5D-E). Despite several residues of mouse

C5aR1 protein being different from that of human C5aR1, citalopram shared a similar binding

model with them (Figure S6). In orthotopic GLUT1^{KD} Hepa1-6 models, flow cytometry

revealed a significant reduction of TAMs in the citalopram group compared with the control

group (Figure 3I). Moreover, TAM polarization was shifted to a proinflammatory state (Figure

3J). Further phenotyping of tumor immune subsets in immunocompetent C57BL/6 mice

showed that citalopram increased the infiltration of CD8^{+} T and dendritic cells while

decreasing the infiltration of CD4^{+} T and B220^{+} B cells (Figure 3K). No significant changes

were observed in splenic immune subsets upon citalopram treatment (Figure S7).

Next, we sought to understand whether C5aR1^{+} TAMs impact on CD8^{+} T cells.

Intratumoral CD8^{+} T cells from tumor-bearing *C5ar1$^{-/-}$* mice had increased frequency and

expression of GZMB, IFN-γ, and TNF-α compared with that from *C5ar1$^{+/-}$* mice (Figure 3L).

Likewise, the frequency and cytotoxicity of CD8$^+$ T cell from r*C5ar1*$^{-/-}$; d*C5ar1*$^{+/-}$ mice were

reduced compared with r*C5ar1*$^{-/-}$; d*C5ar1*$^{-/-}$ mice, and this effect was partially reversed by

citalopram treatment (Figure 3M). Furthermore, depletion of CD8$^+$ T cells abrogated the effect

of macrophage C5aR1 deficiency on tumor growth (Figure 3N), suggesting that CD8$^+$ T cells

are needed for the pro-tumor phenotype of C5aR1$^+$ TAMs. Clinically, data from TCGA cohort

reinforced the link between C5aR1 expression and an immunosuppressive TME as

demonstrated by the associations with TAMs, exhausted T cells, immune checkpoint

molecules, and effector Tregs (Figure 3O).

214   Citalopram-induced activation of CD8$^+$ T cells involves a serotonergic-dependent

215   mechanism

Considering the fact that citalopram targets GLUT1 and CD8$^+$ T cell function is reliant on

glycolytic metabolism, we investigated the *in vivo* effects of citalopram on T cells. As a result,

citalopram significantly invigorated CD8$^+$ T cell function and glycolytic metabolism in

intrahepatic tumors from WT C57BL/6 mice (Figures 4A-B). To simulate the role of metabolic-

associated fatty liver disease (MAFLD) in the development of cirrhosis and HCC[22, 23], we fed

C57BL/6 mice a choline-deficient, amino acid-defined high-fat diet (CDAHFD) to induce

metabolic dysfunction-associated steatohepatitis (MASH)/fibrosis prior to orthotopic Hepa1-6

cell implantation. Likewise, CD8$^+$ T cell function and glycolytic metabolism were enhanced in

MASH mice (Figures 4C-D). GLUT1-deficient cytotoxic T cells remain phenotypically and

functionally unaffected [24], indicating that other glucose uptake mechanisms exist in cytotoxic

T cells. GLUT1 and GLUT3 are essential for the activation and effector function of T cells [24, 25].

We revealed that intratumoral CD8$^+$ T cells massively upregulated GLUT3 but not GLUT1

(Figure S8A), and this finding was supported by single cell sequencing analysis (Figure S8B),

suggesting the GLUT3-dependent glucose uptake in CD8$^+$ T cells.

Given the wide range of peripheral 5-HT actions, we hypothesized that citalopram

modulates the immune system through the serotonergic mechanisms. Indeed, citalopram

administered for 2 weeks in tumor-bearing mice led to reduced serum 5-HT levels (Figure 4E),

which were accompanied with a reduction of low-grade systemic inflammatory responses in

the MASH mice (Figure 4F). Compared to the control C57BL/6 mice, tumor-bearing *Tph1*$^{-/-}$

mice, which lack peripheral 5-HT, exhibited slowed tumor growth and increased effector function of intratumoral CD8$^+$ T cells (Figures 4G-I). Interestingly, citalopram reduced serum 5-HT to a level as observed in *Tph1*$^{-/-}$ mice, and displayed a superior inhibitory effect on tumor growth and CD8$^+$ T cell function compared with that induced by *Tph1* deficiency (Figures 4G-I). This finding suggests a dual immunostimulatory roles of citalopram on CD8$^+$ T cells, particularly the serotonergic-dependent mechanism. Next, we investigated the direct effects of citalopram on T cell activation, expansion, and cytotoxicity. Citalopram enhanced αCD3/αCD28-stimulated CD8$^+$ T cell proliferation as indicated by CFSE staining (Figure S9A). Moreover, citalopram promoted the TCR-instigated activation of CD8$^+$ T cells and the generation of effector CD8$^+$ T cells as evidenced by the cell surface expression of CD44 and CD62L (Figure S9B). Likewise, the cytotoxic activity of activated CD8$^+$ T was enhanced by citalopram, as reflected by GZMB, IFNγ, and TNF-α staining (Figures S9C-E).

Finally, to examine the immune-dependent anti-tumor effects of citalopram, we depleted either CD8$^+$ or CD4$^+$ T cells from GLUT1^{KD} Hepa1-6 tumor-bearing mice and found that depletion of CD8$^+$ T cells, rather than CD4$^+$ T cells, increased tumor growth in citalopram-treated mice but had marginal effects in control mice (Figure 4J), showing that CD8$^+$ T cells are needed in the effector mechanisms of citalopram.

Discussion

Repurposing approved non-oncology drugs for cancer therapy offers cost-effectiveness, an accelerated development timeline, and a known safety profile as key advantages. Accumulating studies have deciphered the inverse link between SSRIs use and cancer risk. However, there is a scarcity of studies investigating the potential repurposing SSRIs for cancer treatment and their mechanisms of action. Non-oncology drugs may have mechanisms of action that are different from traditional cancer treatments. Previously, we and others have uncovered several new targets of SSRIs, including GLUT1 in HCC, SMPD1 in glioblastoma [26], and VDAC1 in autophagy [27]. Here, we further broaden the knowledge of how SSRIs impact the immune system to exert anti-tumor effects. By targeting C5aR1 on TAMs, citalopram enhances the phagocytic activity of TAMs and boosts local anti-tumor immune responses. Furthermore, treatment with citalopram results in a reduction in circulating 5-HT

levels and this phenomenon is akin to the deficiency of *Tph1* and enhances the activation of CD8[+] T cells. Therefore, these insights shed light on the immunomodulatory role of SSRIs in the context of cancer treatment (Figure 5).

C5aR1 is expressed on TAMs within the TME, which helps to explain why citalopram has better anti-tumor effects in immune-competent mice. In addition to TAMs, C5aR1 is also expressed by neutrophils, mast cells, and myeloid-derived suppressor cells in cancers [14, 28, 29]. Although the functional impacts of citalopram on these C5aR1-expressing immune cells have not been formally determined, macrophage depletion assay and C5aR1[-/-] BMDM co-injection experiment suggested that citalopram exerted anti-tumor functions *in vivo* through C5aR1-expressing TAMs. C5aR1[+] macrophages are profoundly implicated in squamous carcinogenesis and targeting C5aR1 reprograms macrophages to recruit cytotoxic CD8[+] T cells [14]. In alignment with this, citalopram treatment increased CD8[+] T cell filtration and their cytotoxic activities in HCC, and this effect was largely dependent on C5aR1[+] macrophages. One possible explanation is that citalopram targets C5aR1 to enhance macrophage phagocytosis and subsequent antigen presentation and/or cytokine production, which promotes T cell recruitment and activity as well as modulate other aspects of tumor immunity. Moreover, citalopram induces TAM phenotypic polarization towards to a proinflammatory state, which supports anti-tumor immune response within the TME. Therefore, we provide a mechanistic insight regarding how citalopram target C5aR1 activity in TAMs, the central findings are important to our understanding of citalopram-induced anti-tumor activities and may lay a basis for upcoming mechanistic studies aimed to decipher specific tumor-promoting roles of this defined TAM population.

While we highlight the anti-HCC effects of SSRIs through targeting C5aR1, it is possible that SSRIs impact tumor progression by modulating the systemic environment via serotonergic mechanisms, especially concerning the activation of CD8[+] T cells. Apart from its role in regulating hemostasis, 5-HT exerts immunoregulatory functions on circulating immune cells [30]. In this study, it was observed that both citalopram treatment and *Tph1* deficiency enhanced CD8[+] T cell cytotoxic activity. Consistently, peripheral 5-HT has been shown to dampen the effector functions of CD8[+] T cells within tumors [31]. Similarly, treatment with SSRIs such as fluoxetine increased the influx of CD8[+] T cells and reduced the growth of syngeneic

pancreatic and colorectal tumors [31]. In addition to the traditional receptor-dependent signaling

transduction, 5-HT can modulate CD8$^+$ T activity through post-translational modification

known as serotonylation [32]. It remains to be explored whether serotonylation contributes to

the immunomodulatory effects of citalopram, as citalopram directly stimulates CD8+ T cell

proliferation, activation, and cytotoxicity *in vitro*. The observed reduction in serum 5-HT levels

and activation of CD8$^+$ T cells following SSRI treatment further highlight the complex interplay

between neurotransmitters, the immune system, and cancer progression. Immunotherapy,

especially immune checkpoint inhibitors (ICIs), has revolutionized the management of HCC [33].

The immune microenvironments play crucial roles in the response or resistance of HCC to

immunotherapies [34]. However, limited molecular biomarkers are available to guide clinical

decision-making for ICI therapy of HCC. Given SSRIs can increase CD8$^+$ T cell infiltration as

well as cytotoxic activities by targeting C5aR1$^+$ TAMs, one can suppose that C5aR1$^+$ TAMs

within the HCC microenvironment might be used as a molecular marker for SSRIs use in

combination with ICI therapy.

In conclusion, our study suggests the potential repurposing of SSRIs for the treatment of

HCC and highlights C5aR1 as a direct target of citalopram. We elucidate the immune-

dependent mechanisms of action of citalopram, involving the regulation of both TAMs and

CD8$^+$ T cells. Combining SSRIs with ICIs or other first-line therapies could enhance the

treatment efficacy of current standard therapies.

Materials and Methods

Animal models.

In the subcutaneous xenograft model, Rag1$^{-/-}$ C57BL/6N mice (male, 6-8-week-old, Cyagen

Biosciences, C001197), wild type immune-competent C57BL/6 mice (male or female, 6-8-

week-old) or *Tph1*$^{-/-}$ mice (male, 6-8-week-old, Cyagen Biosciences, S-KO-05541), were

maintained on a 12-hour day/night cycle with ad libitum access to food and water. For the

pharmacological inhibition study, 5×10^6 mouse HCC cells (wild type, shControl, sh*Slc6a4* or

sh*Slc2a1* Hepa1-6 and Hep53.4) in 100 µl HBS were injected subcutaneously in the lower

back. When bore visible tumors, mice were given intraperitoneal injections of vehicle or

citalopram (5 mg/kg body weight, Selleck, Shanghai, China) daily (n = 5-7 per group).

In the orthotopic xenograft model, 5×10^6 Hepa1-6 cells transfected with luciferase-

expressing lentiviruses were orthotopically implanted into the liver of WT or MASH C57BL/6

mice. The day following tumor cell implantation, mice were treated with citalopram (5 mg/kg

body weight) or DMSO. In the diet-induced MASH model, mice were fed a choline-deficient,

amino acid-defined high-fat diet (45 kcal% fat) supplemented with 0.1% methionine (CDAHFD

diet, A06071309, from Research Diets) for a duration of 6 weeks.

For the C5aR1 knockdown study, C57BL/6J-*C5ar1*[em1Cya] mice were purchased from

Cyagen Biosciences (S-KO-01274, Suzhou, China). 5×10^6 sh*Slc2a1* Hepa1-6 cells were

injected into the lower back of 6-8-week-old male *C5ar1*[+/-] or *C5ar1*[-/-] C57BL/6 mice. Tumor

volume was measured every week using a digital caliper. Four weeks later, mice were

sacrificed and xenograft tumors were analyzed.

To determine the association between citalopram and C5aR1, 2×10^6 GLUT1[KO] Hepa1-6

cells were mixed 1:1 with BMDMs (*C5ar1*[+/-] or *C5ar1*[-/-]) in growth factor-reduced Matrigel

(#354230, Corning, Shanghai, China) and injected into the lower back of 6-8-week-old male

C5ar1[-/-] C57BL/6 mice. To acquire BMDMs, bone marrow was harvested from the femurs and

tibias of *C5ar1*[+/-] or *C5ar1*[-/-] mice. Then, isolated cells were cultured in RPMI 1640

supplemented with 10% FBS, stimulated with recombinant CSF-1 (20 ng/ml, 416-ML, R&D

systems), and maintained at 37°C in a humidified atmosphere with 5% CO_2. Differentiation of

BMDMs was confirmed by flow cytometric analysis of F4/80 (28463-1-AP, Proteintech) and

CD45 (20103-1-AP, Proteintech) expression. One week after cancer cell implantation, mice

were subjected to treatment with citalopram (5 mg/kg body weight) or DMSO. At indicated

time point, mice were sacrificed and xenograft tumors were analyzed.

In all animal studies, tumor volume was assessed using caliper measurements and

calculated using the formula: tumor = 1/2(length $\times$ width^2), where length represents the

largest diameter and width indicates the smallest diameter. At the experimental endpoint,

mice were humanely euthanized, tumors were excised, and tumor weight was recorded. All

animals received proper care in accordance with the guidelines outlined in the "Guide for the

Care and Use of Laboratory Animals" established by the National Academy of Sciences and

published by the National Institutes of Health (NIH publication 86-23 revised 1985). The

protocols for animal studies were approved by the ethics committee of the Ren Ji Hospital,

School of Medicine, Shanghai Jiao Tong University (approval number 202201427 and RA-2021-096).

Cell culture and reagents.

The following cell lines were included: Hepa1-6, Hep53.4, THP-1, and HEK293T. All cells were cultured in a medium recommended by ATCC, supplemented with 10% fetal bovine serum (FBS; Gibco, USA) and 1% (v/v) streptomycin-penicillin (Sigma-Aldrich, Shanghai, China), and maintained at 37°C in a humidified atmosphere with 5% CO_2. Prior to conducting cell experiments, all the mentioned cell lines underwent testing for short tandem repeat profiling and mycoplasma contamination to ensure their authenticity and purity. Citalopram (S4749), escitalopram (S4064), fluoxetine (S1333), fluvoxamine (S1336), paroxetine (S3005), and sertraline (S4053) were all obtained from Selleck (Shanghai, China).

Plate colony formation assay.

For cell proliferation analysis, a total of 1,000 indicated HCC cells were seeded into 6-well plates with three replicates. After treatment with citalopram for 10-14 days, the plates were washed with phosphate-buffered saline (PBS) twice, fixed with methanol for 10 min, and stained with 0.5% (w/v) crystal violet solution for 10 min. Colonies containing more than 30 cells were counted under the microscope.

Detection of Caspase-3/7 activity.

Cell apoptosis in HCC cells upon citalopram treatment was assessed using the Apo-ONE® Homogeneous Caspase-3/7 Assay (Promega, Madison, WI, USA) as described previously [35]. To normalize the data, the cell number in the same plate was detected using CellTiter-Blue (Promega, G8081). The relative Caspase-3/7 activity was determined by calculating the ratio of the signals obtained from the Apo-ONE assay to those from the CellTiter-Blue assay.

Gene knockdown and knockout.

The specific shRNA oligonucleotides targeting *Slc6a4* and *Slc2a1* were custom-synthesized

by Genepharma (Shanghai, China). Lentivirus packaging was carried out in HEK293T cells

using Lipofectamine 2000 (Invitrogen, Carlsbad, CA), and virus titers were determined

following standard procedures. HCC cells (Hep53.4 and Hepa1-6) were infected with the

virus-containing supernatant in the presence of 6 µg/ml polybrene (Sigma-Aldrich, H9268, St.

Louis, MO) when they reached 70% confluence. Cells expressing the shRNAs were selected

using puromycin (2 µg/ml, Gibco, A1113802).

For CRISPR/Cas9-mediated gene knockout, THP-1 cells were initially transfected with

the lentiCas9-Blast vector to facilitate the expression of Cas9. To generate *C5aR1*$^{-/-}$ THP-1

cells, a lentiviral system utilizing guide RNA (sgRNA) targeting C5AR1 was utilized. The

lentivirus was produced in HEK293T cells by transfecting lentiGuide-Puro along with

packaging plasmids pVSVg and psPAX2. The transfection was carried out using Fugene6

(E2693, Promega) in OptiMEM. After two days, the supernatants were filtered through a 0.45

µm filter, and Cas9-expressing cells were infected with the lentivirus in the presence of 6

µg/ml polybrene. Individual colonies were then seeded into 96-well plates, allowed to reach

confluence, and subsequently subjected to knockout efficiency analysis. The sequences for

shRNA and sgRNA are provided below: sh*Slc6a4*,

CACCGCCTCCTACTATAACACCATCCGAAGATGGTGTTATAGTAGGAGGC; sh*Slc2a1*,

CACCGGGAGAAGAAGGTCACCATCTCGAAAGATGGTGACCTTCTTCTCCC; *C5AR1*

sgRNA, CTTCAGTCAACACGTTCCGG.

Quantitative real-time PCR (qRT-PCR).

RNA was extracted from CD8^{+} T cells using the RNAiso Plus reagent (Takara, Japan) and

subjected to reverse transcription using the PrimeScript RT-PCR kit (Takara, Japan). Then,

qRT-PCR was performed using the 7500 Real-time PCR system (Applied Biosystems, USA).

The housekeeping gene *Gapdh* was used to normalize gene expression. Primer sequences

used in this study were shown below: *Slc2a1* forward, 5'-CAGTTCGGCTATAACACTGGTG-3';

Slc2a1 reverse, 5'-GCCCCCGACAGAGAAGATG-3'; *Slc2a3* forward, 5'-

ATGGGGACAACGAAGGTGAC-3'; *Slc2a3* reverse, 5'-GTCTCAGGTGCATTGATGACTC-3';

Gapdh forward, 5'-AGGTCGGTGTGAACGGATTTG-3'; *Gapdh* reverse, 5'-

TGTAGACCATGTAGTTGAGGTCA-3'.

Immunoblotting.

Cellular lysates were extracted using RIPA lysis buffer (P0013B, Beyotime, Shanghai, China) supplemented with protease and phosphatase inhibitor cocktails (ab201119, Abcam, Shanghai, China). The protein concentration was quantified with a BCA Protein Assay Kit (Pierce Biotechnology, USA). The proteins were then separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and transferred onto polyvinylidene difluoride (PVDF, Millipore) membranes. Subsequently, the membranes were blocked with 5% (m/v) skim milk for 1 h at room temperature before being incubated with specific antibodies overnight at 4°C. On the following day, the membranes were washed three times with PBS and then incubated with HRP-conjugated secondary antibodies for 45 min. Subsequently, an ECL chemiluminescence assay (SB-WB012, Share-bio, Shanghai, China) was performed using a Bio-Spectrum Gel Imaging System (Bio-Rad). The antibodies utilized in this study included: SERT (1:500, ProteinTech, 19559-1-AP), GLUT1 (1:2,000, Cell Signaling Technology, #73015), C5aR1 (1:1,500, ProteinTech, 21316-1-AP), and β-actin (1:2,000, Abcam, ab8226). The information for secondary antibodies is as follows: goat anti-rabbit IgG (H+L) cross-adsorbed secondary antibody (1:5,000, Thermo Fisher Scientific, G-21234) and goat anti-mouse IgG (H+L) cross-adsorbed secondary antibody (1:5,000, Thermo Fisher Scientific, G-21040).

Immunohistochemistry and immunofluorescence analysis.

Paraffin-embedded sections of mouse HCC tissues were subjected to immunohistochemical analysis of Ki67, cleaved caspase 3, and C5aR1 expression. Immunohistochemical analysis was performed as reported previously [35]. The primary antibodies used in this study were shown as follows: Ki67 (1:400, Cell Signaling Technology, #9449), cleaved caspase 3 (1:400, Cell Signaling Technology, #9661), and C5aR1 (1:1,200, ProteinTech, 21316-1-AP). For immunofluorescence analysis, paraffin sections (5 μm) of HCC samples were deparaffinized and rehydrated with graded ethanol, followed by microwave heating-based antigen retrieval. Then tissue sections were blocked with 10% donkey serum (Sigma-Aldrich, D9663, USA) for 1 h at room temperature and incubated with primary antibodies at 4°C overnight. The

following antibodies were used: C5aR1 (1:50, Proteintech, 10375-1-AP) and CD163 (1:200, Abcam, ab182422). The next day, tissue sections were incubated with donkey anti-rabbit Alexa Fluor 594 (1:400, Jackson ImmunoResearch, #711-585-152) and donkey anti-mouse Alexa Fluor 488 (1:400, Jackson ImmunoResearch, #715-545-150) for 30 min at room temperature. Finally, 4',6-diamidino-2-phenylindole (DAPI) staining was performed to visualize the nuclei. Confocal microscope equipped with a digital camera (Nikon, Japan) was used for high-resolution imaging.

Bioinformatics analysis and single cell sequencing data.

Correlation analysis was performed with GEPIA2 (http://gepia2.cancer-pku.cn/#index) [36]. The following gene signatures were included for analysis, TAMs (*CD33*, *MARCO*, *CXCL5*, *SULT1C2*, *MSR1*, *CTSK*, *PTGDS*, *COLEC12*, *GPC4*, *PCOLCE2*, *CHIT1*, *CLEC5A*, *CCL7*, *FN1*, *CD163*, *GM2A*, *BCAT1*, *RAI14*, *COL8A2*, *CHI3L1*, *ATG7*, *CD84*, *MS4A4A*, *EMP1*, *CYBB*, *CD68*, and *CD11b*), exhausted T cells (*HAVCR2*, *TIGIT*, *LAG3*, *PDCD1*, *CXCL13* and *LAYN*), immune checkpoint (*IDO1*, *IDO2*, *PDCD1*, *CD274*, *PDCD1LG2*, *CTLA4*, *LAG3*, *HAVCR2*, *C10orf54*, *BTLA*, *ICOS*, and *TNFRSF9*), and effector Treg cells (*FOXP3*, *CTLA4*, *CCR8*, and *TNFRSF9*). The single-cell transcriptome expression of C5aR1 and GLUT family members in the tumor microenvironment was identified with GSE125449 and GSE140228-10X.

DARTS experiments.

THP-1 or HEK293T cells were lysed using NP40 lysis buffer (Thermo Fisher Scientific, 89842Y). Following centrifugation at 12,000 rpm for 10 min, the supernatant containing the proteins was collected, and the protein concentration was determined using a BCA Protein Assay Kit (Pierce Biotechnology, USA). Samples of equivalent quality were then treated with specific SSRIs and a vehicle at 37°C for 4 h. Subsequently, pronase (10 μg/mL, Roche, 10165921001) or distilled water was added and incubated at room temperature for specified time points (10 and 30 min). The reaction was terminated by adding protease inhibitors, and the resulting products were harvested and subjected to analysis using Western blotting.

 ***In Silico* docking study.**

 To predict the binding mode of citalopram/C5aR1, molecular docking was performed using

 the AutoDock 4.2.6 software package [37]. The receptor structure for C5aR1 (PDB id: 6C1Q) [20]

 was sourced from the RCSB protein data bank (https://www.rcsb.org/). Due to the lack of

 available structural information, the structure of mouse C5aR1 was predicted using a local

 version of the ColabFold (AlphaFold2) software [38, 39]. The original substrates and other

 heteroatoms in these crystal structures were removed for further docking procedure. The

 absolute structure of citalopram (CID: 2771), fluoxetine (CID: 3386), fluvoxamine (CID:

 5324346), paroxetine (CID: 43815), and sertraline (CID: 68617) were sourced from the

 PubChem database (https://pubchem.ncbi.nlm.nih.gov/). In each docking scenario, molecular

 docking was conducted following a standard protocol [40], with a modification in the number of

 genetic algorithm runs to 500 to comprehensively explore the binding modes. The binding

 energy and cluster information were directly extracted from the AutoDock output.

 Biochemical analysis.

 The serum levels of inflammatory cytokines (TNF-α, IL-1β, and IL-6) were analyzed using

 commercial ELISA kits following the manufacturer's instructions (R&D Systems). The cytokine

 concentrations were calculated using a standard curve, and the data were presented in

 picograms per milliliter. Serum 5-HT levels were assessed using the 5-HT ELISA Kit from

 Abcam (ab133053), following the manufacturer's guidelines.

 Isolation of TAMs from xenograft tumor tissues.

 Fresh sterile tumor tissues from subcutaneous xenograft model were washed with PBS two

 times, minced with scalpels, and then digested using a Tumor Dissociation Kit (Miltenyi Biotec,

 cat #130-095-929) at room temperature for 30 min. Digested tumor tissues were filtered

 through Falcon 70Micrometer Cell Strainer (Corning, #352350) and centrifuged at 1,500 rpm

 for 5 min. Red blood cells were removed using Red Blood Cell Lysis Solution (Miltenyi Biotec,

 cat #130-094-183). Then, TAMs were selected by CD11b microbeads (Miltenyi Biotec, cat

 #130-049- 601) according to the manufacturer's instructions.

Isolation of immune cells and flow cytometry.

Mice were euthanized, and fresh tumor tissues and spleen were harvested. Spleens were mechanically disrupted through 70 µm Nylon mesh (Corning, #352350) to release single cells. The cells were washed with PBS, followed by the lysis of red blood cells with Red Blood Cell Lysis Solution (Miltenyi Biotec, cat #130-094-183). Tumor tissues were cut into small pieces with surgical scissors and digested with 2 mg/mL collagenase IV (Sigma) for 20 min and filtered with 70 µm Nylon mesh. The isolated cells were collected with PBS and subjected to centrifuged at 1,500 rpm for 5 min. To enrich immunocytes from tumor tissues, harvested cells were centrifuged through different concentrations of discontinuous gradients of 40/80 Percoll (GE Healthcare, cat #17-0891-01) at 350 × g for 30 min. The middle pellet was washed with PBS, subjected to red blood cell lysis for 3 min.

For staining molecules expressed on cell surface, single cell suspensions from spleens and tumor tissues were labeled with fluorophore-conjugated antibodies: PerCP-Cyanine 5.5 7AAD (Biolegend, Cat #420404, 1:1,000), FITC anti-mouse CD4 (Biolegend, Cat #100406, 1:1,000), PE anti-mouse Ly6G (Biolegend, Cat #127608, 1:200), APC anti-mouse CD11c antibody (Biolegend, Cat #117310, 1:200), PE-cy7 anti-mouse CD8α (Biolegend, Cat #100722, 1:200), Brilliant Violet 421 anti-mouse CD3 (Biolegend, Cat #100227, 1:200), Alex Fluor 647 anti-mouse F4/80 (Invitrogen, Cat #2604384, 1:200), Brilliant Violet 510 anti-mouse CD45 (Biolegend, Cat #103138, 1:200), PE-cy7 anti-mouse NK1.1 (Biolegend, Cat #156513, 1:200), Brilliant Violet 421 anti-mouse B220 (Biolegend, Cat #103240, 1:200), Brilliant Violet 605 anti-mouse CD11b (Biolegend, Cat #101257, 1:200), APC-CY7 Stain 780 (BD Bioscience, Cat #565388, 1:1,000).

For intracellular staining, single cell suspensions from tumor tissues were stimulated with Cell Stimulation Cocktail (Invitrogen) for 4 h, and labeled with PE anti-mouse CD45 antibody (Biolegend, Cat #103106, 1:200), PerCP-Cyanine 5.5 CD3 (Biolegend, Cat #100325, 1:200), PE-cy7 anti-mouse CD8a (Biolegend, Cat# 100722, 1:200), APC-CY7 Stain 780 (BD Bioscience, Cat #565388, 1:1,000). After fixation and permeabilization using the Fix/Perm kit (ThermoFisher, GAS004), cells were labeled with APC anti-human/mouse Granzyme B Recombinant antibody (Biolegend, Catalog no.372203, 1:200), Brilliant Violet 421 anti-mouse IFN-γ (Biolegend, Cat #505830, 1:200), Brilliant Violet 605 anti-mouse TNF-α (Biolegend, Cat

#506329, 1:200). Flow cytometry was performed in BD FACSCanto II.

T cell activation, expansion and cytotoxicity

Single-cell suspensions were freshly dissected from the spleen and lymph nodes, and red

blood cells were removed using Red Blood Cell Lysis Solution (Miltenyi Biotec, cat #130-094-

183). $CD8^+$ T cells were isolated from the resulting cell suspensions by $CD8^+$ magnetic bead

selection (Miltenyi Biotec, cat #130-117-044). For T cell proliferation assay, isolated $CD8^+$ T

cells were resuspended in PBS supplemented with carboxy fluorescein succinimidyl ester

(CFSE, Invitrogen, 1:10,000) for 8 min at room temperature and washed with RPMI 1640,

supplemented with 10% heat-inactivated FBS. Then, $CD8^+$ T cells were activated by plate-

bound $\alpha CD3\epsilon$ (5 µg/mL, Biolegend, Catalog no.100359) and α-CD28 (2 µg/mL, Biolegend,

Catalog no.102121) in complete RPMI 1640 medium at 37°C and 5% CO_2 for 72 h in the

presence or absence of 5 µM citalopram treatment. Finally, cell proliferation was assessed by

flow cytometry. For assessment of cellular activation and cytotoxicity, $CD8^+$ T cells were

stimulated by Leukocyte Activation Cocktail (GolgiPlug™ BD, Catalog no.550583) for 4 h,

resuspended in PBS/2% FBS containing PE anti-mouse CD62L antibody (Biolegend, Catalog

no.104407, 1:200) and Brilliant Violet 510 anti-mouse/human CD44 Antibody (Biolegend,

Catalog no.103044, 1:200), followed by incubation at room temperature for 30 min and

washed in PBS/2% FBS. After fixation and permeabilization using the Cytofix/Cytoperm

solution (BD Biosciences) for 20 min at 4°C, Brilliant Violet 421 anti-mouse IFN-γ antibody

(Biolegend, Catalog no.505830, 1:200), APC anti-human/mouse Granzyme B Recombinant

antibody (Biolegend, Catalog no.372203, 1:200) and Alexa Fluor 488 anti-mouse TNF-α

antibody (Biolegend, Catalog no. 506313, 1:200) were employed for intracellular staining.

Then, cell activation and cytotoxicity were assessed by flow cytometry.

Macrophage phagocytosis assay

Phagocytosis assays of tumor cells were performed as previously described [41]. Briefly, 5 ×

10^4 phorbol myristate acetate (PMA, 100 ng/ml) primed THP-1 cells and harvested TAMs

were seeded overnight in a 24-well tissue culture plate. The next day, HCC-LM3 cells and

Hepa1-6 cells were labelled with pHrodo™ Green (Thermo, Cat #P35373, 1:1,000). After

incubating macrophages in serum-free medium for 2 h, the macrophages were labeled with

CellTracker™ Deep Red Dye (Thermo, Cat #C34565, 6 ng/mL). Then, 5×10^4 predyed tumor

cells were added to the macrophages, in the presence or absence of C5a stimulation and

different SSRIs (5 µM). After incubation at 37 °C for 4 h, macrophages were extensively

washed and imaged with Confocal Laser Scanning Microscope (Leica).

Macrophage depletion assay

The method for eliminating macrophages was reported previously [42]. Briefly, C57BL/6 mice

were intraperitoneally injected with clodronate liposomes (1.4 mg/20 g body weight) to

remove macrophages. Clodronate liposomes (Yeasen, 40337ES10, Shanghai, China) were

given every three days for 12 days before tumor inoculation. After inoculation of sh*Slc2a1*

(GLUT1^{KD}) Hepa1-6 cells, clodronate liposomes were also administered every three days

until the end of the experiment. When the mice were sacrificed, liver and tumor tissues were

extracted, and F4/80 staining (1:250, Cell Signaling Technology, #70076) was performed to

determine the efficiency of macrophage deletion.

CD4$^+$ T cell and CD8$^+$ T cell depletion experiments

For CD4$^+$ T and CD8$^+$ T cell depletion, mice were treated intraperitoneally with 200 µg of

specific monoclonal antibodies (Ultra-LEAF™ Purified anti-mouse CD8a Antibody, Biolegend,

cat #100763; and Ultra-LEAF™ Purified anti-mouse CD4 Antibody, Biolegend, cat #100457)

the day before the implantation of cancer cells and then with 100 µg twice a week for the

entire duration of the experiment.

Statistical analysis.

The sample sizes for in vitro and in vivo studies determined based on similar studies in the

field and through pilot studies. Details regarding sample sizes and biological replicates can be

found in the figure legends. Continuous variables are presented as means ± SD. For

comparisons between two independent groups, a two-tailed Student's t-test was employed.

Statistical analyses involving more than two groups were conducted using either a one-way

analysis of variance (ANOVA) with Tukey's method or two-way ANOVA with Dunnett multiple

comparisons, as specified. GraphPad Prism 9 (GraphPad Software Inc., San Diego, CA) was

used for statistical analyses. Kaplan-Meier survival curves were generated and analyzed

using the log-rank test. A p-value less than 0.05 was considered statistically significant. In all

figures, statistical significance is represented as follows: $*p < 0.05$; $**p < 0.01$; $***p < 0.001$.

ACKNOWLEDGMENTS

The research was supported by grants from National Natural Science Foundation of China

(82001469 and 82071581), Shanghai Pilot Program for Basic Research- Shanghai Jiao Tong

University (21TQ1400225), Natural Science Foundation of Shanghai (22ZR1459500), "Rising

Stars of Medical Talents" Youth Development Program (Youth Medical Talents-Specialist

Program), Clinical Medicine Research Center Construction Project of Huadong Hospital

(LCZX2202), Shanghai Outstanding Young Medical Personnel Training Program & Excellence

Project of Shanghai Municipal Health Commission (20224Z0009), and Key specialized

diseases construction of Huadong Hospital (ZDZB2225).

AUTHOR CONTRIBUTIONS

Conceptualization: S.H.J.; Methodology: F.D., S.H.J., K.S., S.Z., L.J.J. and J.J.; Formal

Analysis: F.D., S.H.J., S.Z. and L.J.J.; Resources: M.X.F., X. W. and S.H.J.; Investigation: all

authors; Supervision: S.H.J., X. W. and Z.G.Z.; Funding Acquisition: S.H.J., Z.G.Z., Z.B. and

F.D.; Writing-Original Draft: F.D., K.S., S.Z., L.J.J. and S.H.J; Writing-Review & Editing: all

authors.

DECLARATION OF INTERESTS

The authors declare no conflicts of interest.

References

1. Pushpakom S, Iorio F, Eyers PA, et al. Drug repurposing: progress, challenges and recommendations. Nature Reviews Drug Discovery 2019;18:41-58.
2. Schipper LJ, Zeverijn LJ, Garnett MJ, et al. Can Drug Repurposing Accelerate Precision

623    Oncology? Cancer Discovery 2022;12:1634-1641.

624 3. Foretz M, Guigas B, Viollet B. Metformin: update on mechanisms of action and repurposing
625    potential. Nature Reviews Endocrinology 2023;19:460-476.

626 4. Xia Y, Sun M, Huang H, et al. Drug repurposing for cancer therapy. Signal Transduction and
627    Targeted Therapy 2024;9.

628 5. Fluyau D, Mitra P, Jain A, et al. Selective serotonin reuptake inhibitors in the treatment of
629    depression, anxiety, and post-traumatic stress disorder in substance use disorders: a Bayesian
630    meta-analysis. European Journal of Clinical Pharmacology 2022;78:931-942.

631 6. Singh I, Seth A, Billesbolle CB, et al. Structure-based discovery of conformationally selective
632    inhibitors of the serotonin transporter. Cell 2023;186:2160-+.

633 7. Lee MJ, Huang CW, Chen YL, et al. Association between selective serotonin reuptake
634    inhibitors and kidney cancer risk: A nationwide population-based cohort study. International
635    Journal of Cancer 2021;148:1331-1337.

636 8. Fischer A, Rennert HS, Rennert G. Selective serotonin reuptake inhibitors associated with
637    increased mortality risk in breast cancer patients in Northern Israel. International Journal of
638    Epidemiology 2022;51:807-816.

639 9. Zhang NQ, Sundquist J, Sundquist K, et al. Combined Use of Aspirin and Selective Serotonin
640    Reuptake Inhibitors Is Associated With Lower Risk of Colorectal Cancer: A Nested Case-
641    Control Study. American Journal of Gastroenterology 2021;116:1313-1321.

642 10. Bhagavathula AS, Woolf B, Rahmani J, et al. Selective serotonin reuptake inhibitor use and
643    the risk of hepatocellular carcinoma: a systematic review and dose-response analysis of cohort
644    studies with one million participants. European Journal of Clinical Pharmacology
645    2022;78:547-555.

646 11. Carlson LE, Ismaila N, Addington EL, et al. Integrative Oncology Care of Symptoms of
647    Anxiety and Depression in Adults With Cancer: Society for Integrative Oncology-ASCO
648    Guideline. Journal of Clinical Oncology 2023;41:4562-+.

649 12. Lomenick B, Hao R, Jonai N, et al. Target identification using drug affinity responsive target
650    stability (DARTS). Proceedings of the National Academy of Sciences of the United States of
651    America 2009;106:21984-21989.

652 13. Beach C, MacLean D, Majorova D, et al. Improving radiotherapy in immunosuppressive
653    microenvironments by targeting complement receptor C5aR1. Journal of Clinical
654    Investigation 2023;133.

655 14. Medler TR, Murugan D, Horton W, et al. Complement C5a Fosters Squamous Carcinogenesis
656    and Limits T Cell Response to Chemotherapy. Cancer Cell 2018;34:561-+.

657 15. Ajona D, Zandueta C, Corrales L, et al. Blockade of the Complement C5a/C5aR1 Axis
658    Impairs Lung Cancer Bone Metastasis by CXCL16-mediated Effects. American Journal of
659    Respiratory and Critical Care Medicine 2018;197:1164-1176.

660 16. Dong F, He K, Zhang S, et al. SSRI antidepressant citalopram reverses the Warburg effect to
661    inhibit hepatocellular carcinoma by directly targeting GLUT1. bioRxiv
662    2024:2024.07.17.603851.

663 17. Pio R, Ajona D, Ortiz-Espinosa S, et al. Complementing the Cancer-Immunity Cycle.
664    Frontiers in Immunology 2019;10.

665 18. Lomenick B, Hao R, Jonai N, et al. Target identification using drug affinity responsive target
666    stability (DARTS). Proc Natl Acad Sci U S A 2009;106:21984-9.

667 19. Robertson N, Rappas M, Dore AS, et al. Structure of the complement C5a receptor bound to
668    the extra-helical antagonist NDT9513727. Nature 2018;553:111-114.

669 20. Liu H, Kim HR, Deepak R, et al. Orthosteric and allosteric action of the C5a receptor
670    antagonists. Nat Struct Mol Biol 2018;25:472-481.

671 21. Wang Y, Liu W, Xu Y, et al. Revealing the signaling of complement receptors C3aR and
672    C5aR1 by anaphylatoxins. Nat Chem Biol 2023.

673 22. Eslam M, Newsome PN, Sarin SK, et al. A new definition for metabolic dysfunction-
674    associated fatty liver disease: An international expert consensus statement. Journal of
675    Hepatology 2020;73:202-209.

676 23. Du DY, Liu C, Qin MY, et al. Metabolic dysregulation and emerging therapeutical targets for
677    hepatocellular carcinoma. Acta Pharmaceutica Sinica B 2022;12:558-580.

678 24. Macintyre AN, Gerriets VA, Nichols AG, et al. The Glucose Transporter Glut1 Is Selectively
679    Essential for CD4 T Cell Activation and Effector Function. Cell Metabolism 2014;20:61-72.

680 25. Hochrein SM, Wu H, Eckstein M, et al. The glucose transporter GLUT3 controls T helper 17
681    cell responses through glycolytic-epigenetic reprogramming. Cell Metabolism 2022;34:516-+.

682 26. Bi JF, Khan A, Tang J, et al. Targeting glioblastoma signaling and metabolism with a re-

683   purposed brain-penetrant drug. Cell Reports 2021;37.

684   27.   Hwang HY, Shim JS, Kim D, et al. Antidepressant drug sertraline modulates AMPK-MTOR
685   signaling-mediated autophagy via targeting mitochondrial VDAC1 protein. Autophagy
686   2021;17:2783-2799.

687   28.   Ou B, Liu Y, Yang X, et al. C5aR1-positive neutrophils promote breast cancer glycolysis
688   through WTAP-dependent m6A methylation of ENO1. Cell Death Dis 2021;12:737.

689   29.   Ding P, Li L, Li L, et al. C5aR1 is a master regulator in Colorectal Tumorigenesis via Immune
690   modulation. Theranostics 2020;10:8619-8632.

691   30.   Costa LHA, Santos BM, Branco LGS. Can selective serotonin reuptake inhibitors have a
692   neuroprotective effect during COVID-19? European Journal of Pharmacology 2020;889.

693   31.   Schneider MA, Heeb L, Beffinger MM, et al. Attenuation of peripheral serotonin inhibits
694   tumor growth and enhances immune checkpoint blockade therapy in murine tumor models.
695   Science Translational Medicine 2021;13.

696   32.   Wang X, Fu SQ, Yuan X, et al. A GAPDH serotonylation system couples CD8+T cell
697   glycolytic metabolism to antitumor immunity. Molecular Cell 2024;84.

698   33.   Zheng Y, Wang S, Cai J, et al. The progress of immune checkpoint therapy in primary liver
699   cancer. Biochim Biophys Acta Rev Cancer 2021;1876:188638.

700   34.   Llovet JM, Castet F, Heikenwalder M, et al. Immunotherapies for hepatocellular carcinoma.
701   Nat Rev Clin Oncol 2022;19:151-172.

702   35.   Jiang SH, Li J, Dong FY, et al. Increased Serotonin Signaling Contributes to the Warburg
703   Effect in Pancreatic Tumor Cells Under Metabolic Stress and Promotes Growth of Pancreatic
704   Tumors in Mice. Gastroenterology 2017;153:277-291 e19.

705   36.   Tang Z, Kang B, Li C, et al. GEPIA2: an enhanced web server for large-scale expression
706   profiling and interactive analysis. Nucleic Acids Res 2019;47:W556-W560.

707   37.   Morris GM, Huey R, Lindstrom W, et al. AutoDock4 and AutoDockTools4: Automated
708   docking with selective receptor flexibility. J Comput Chem 2009;30:2785-91.

709   38.   Mirdita M, Schutze K, Moriwaki Y, et al. ColabFold: making protein folding accessible to all.
710   Nature Methods 2022;19:679-+.

711   39.   Jumper J, Evans R, Pritzel A, et al. Highly accurate protein structure prediction with
712   AlphaFold. Nature 2021;596:583-+.

713   40.   Forli S, Huey R, Pique ME, et al. Computational protein-ligand docking and virtual drug
714   screening with the AutoDock suite. Nat Protoc 2016;11:905-19.

715   41.   Chen J, Zhong MC, Guo HJ, et al. SLAMF7 is critical for phagocytosis of haematopoietic
716   tumour cells via Mac-1 integrin. Nature 2017;544:493-+.

717   42.   Jiang SH, Zhu LL, Zhang M, et al. GABRP regulates chemokine signalling, macrophage
718   recruitment and tumour progression in pancreatic cancer through tuning KCNN4-mediated
719   Ca(2+) signalling in a GABA-independent manner. Gut 2019;68:1994-2006.

Fig. 1. The immune-dependent and SERT-independent anti-tumor effects of citalopram

in HCC. (**A**, **B**) Mouse HCC cells, Hepa1-6 or Hep53.4, were subcutaneously injected into

Rag1$^{-/-}$ C57BL/6 or immunocompetent C57BL/6 mice (n = 5-7 per group). When bore visible

tumors, 5 mg/kg citalopram was treated daily for 15 or 25 days. Tumors were excised after

mice were sacrificed, and the tumor weight was measured. (**C**, **D**) Western blotting showed

the knockdown efficiency of SERT in Hepa1-6 and Hep53.4 cells. (**E**, **F**) shControl and

sh*Slc6a4* Hepa1-6 and Hep53.4 cells were subcutaneously injected into Rag1$^{-/-}$ C57BL/6 or

immunocompetent C57BL/6 mice (n = 5-6 per group). When bore visible tumors, 5 mg/kg

citalopram was treated daily for 15 or 25 days. Tumors were excised after mice were

sacrificed, and the tumor weight was measured. In all panels, *$p < 0.05$, **$p < 0.01$, ***$p <$

0.001. Values as mean ± SD and compared by one-way ANOVA multiple comparisons with

Tukey's method (for bar chart comparison) and two-way ANOVA with Dunnett's multiple

comparisons (for survival curve comparison).

Fig. 2. C5aR1 is a direct target of citalopram.

(**A**) Gene Set Enrichment Analysis (GSEA) identified GLUT1 and C5aR1 as two top hits

related to SSRI-induced gene changes. (**B**) GSEA of HCC RNAseq data (TCGA cohort) with

the SSRI-related gene signature. Sample grouping was made based on the median

expression of C5aR1. (**C**) Representative immunohistochemical images showed the

expression pattern and cellular distribution of C5aR1 in human HCC tissues. Scale bar, 50

µm. (**D**) Single cell RNA sequencing analysis showed the expression pattern of C5aR1 with

the immune microenvironment of HCC. (**E**) Co-immunofluorescence of C5aR1 (green) with

CD163 (red) in HCC samples. Scale bar, 10 µm. (**F**) The DARTS assay and immunoblot

analysis showed C5aR1 protein stability against 5 µg/mL pronase in the presence and

absence of 100 µM citalopram treatment. (**G**) The DARTS assay and immunoblot analysis

showed C5aR1 protein stability against 5 µg/mL pronase in the presence of different

concentration of citalopram treatment. (**H**) The overall conformation of citalopram binding to

C5aR1. (**I**) Representative models of citalopram in pose-1 (left), pose-2 (middle) and allosteric

site (right). Several polar interactions were indicated by black dashed lines. (**J**) HEK293T

cells were transfected with either WT or mutant C5aR1 expression plasmids for 48 h, followed

by DARTS assay with immunoblotting analysis of C5aR1 protein levels. In all panels, *$p <$

0.05, **$p < 0.01$. Values as mean ± SD and compared by the Student's t test (**F**) or one-way

ANOVA multiple comparisons with Tukey's method among groups (**G**, **J**).

Fig. 3. Citalopram targets C5aR1[+] TAMs

(**A**) Western blotting showed the knockdown efficiency of GLUT1 in mouse Hepa1-6 cells. (**B**) GLUT1[KD] Hepa1-6 cells were subcutaneously injected into the Rag1[-/-] or immunocompetent C57BL/6 mice, and mice were treated with 5 mg/kg citalopram when bore visible tumors; three weeks later, tumor burden was examined (n = 6-7 per group). (**C**) The growth kinetics of GLUT1[KD] Hepa1-6 tumors in *C5ar1*[+/-] and *C5ar1*[-/-] C57BL/6 host (n = 7). (**D**) Immunofluorescence analysis of C5a deposition in GLUT1[KD] Hepa1-6 tumors from *C5ar1*[+/-] and *C5ar1*[-/-] C57BL/6 host. Scale bar, 50 μm. (**E, F, H**) GLUT1[KD] Hepa1-6 cells admixed with *C5ar1*[+/-] and *C5ar1*[-/-] BMDMs from donor (d) mice were subcutaneously implanted into

syngeneic recipient (r) mice. The therapeutic effect of citalopram (**E**), C5a deposition (**F**), and

macrophage phagocytosis (**H**) in this model were analyzed. Scale bar, 50 µm. (**G**) The

phagocytic capacity of macrophages isolated from GLUT1^{KD} Hepa1-6 tumors in *C5ar1$^{+/-}$* and

C5ar1$^{-/-}$ C57BL/6 host. (**I-K**) Flow cytometry showed the infiltration of CD45^{+}CD11b^{+}F4/80^{+}

macrophages (**I**), CD206^{+} TAMs and CD11b^{+} TAMs (**J**), tumor-infiltrating lymphocytes (**K**) in

tumor tissues from orthotopic xenograft model, which generated in immunocompetent

C57BL/6 mice with Hepa1-6 cells (n = 5 per group). (**L, M**) Measurement of CD8^{+} T cell

function in tumor tissues from the groups mentioned in **C** and **E**. (**N**) The growth kinetics of

GLUT1^{KD} Hepa1-6 tumors in *C5ar1$^{+/-}$* and *C5ar1$^{-/-}$* C57BL/6 host upon CD8^{+} T cell deletion (n

= 7). (**O**) Correlation analysis of C5aR1 expression and immune checkpoint molecules, gene

signatures of TAMs, exhausted T cells, and effector Tregs in the TCGA cohort (n = 371). In all

panels, *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$; ns, non-significant. Values as mean ± SD and

compared by two-way ANOVA with Dunnett's multiple comparisons (**B**, **C**, **E**, **N**), Student's t

test (**G**, **H**, **I-L**), one-way ANOVA multiple comparisons with Tukey's method (**B**, **M**), and the

Spearman's rank correlation methods (**O**).

Fig. 4. Citalopram activates CD8⁺ T cells.

(**A**, **B**) Measurement of CD8⁺ T cell function and glycolysis in orthotopic tumor tissues from

WT C57BL/6 mice (n = 5 per group). (**C**, **D**) Measurement of CD8⁺ T cell function and

glycolysis in orthotopic tumor tissues from MASH mice (n = 5 per group); Basal ECAR

indicates glycolysis after the addition of glucose, and ΔECAR represents the difference

between oligomycin-induced ECAR and 2-DG-induced ECAR. (**E**) Serum 5-HT levels in

GLUT1^KD Hepa1-6 tumor-bearing mice fed with chow diet or CDAHFD, with the presence or

absence of citalopram treatment (n = 5 per group). (**F**) Serum TNF-α, IL-1β, and IL-6 levels in

GLUT1^KD Hepa1-6 tumor-bearing mice fed with chow diet or CDAHFD, with the presence or

absence of citalopram treatment (n = 5 per group). (**G**) Serum 5-HT levels in WT C57BL/6

and *Tph1*[-/-] mice, with the presence or absence of citalopram treatment (n = 5 per group). (**H**)

Tumor growth of WT and *Tph1*[-/-] mice after subcutaneous injection of Hepa1-6 cells and

treatment with citalopram. (**I**) Measurement of CD8$^+$ T cell function in tumor tissues from the

groups mentioned in **H**. (**J**) The therapeutic effect of citalopram on GLUT1^{KD} Hepa1-6 tumor

was tested in the presence or absence of CD4$^+$ T or CD8$^+$ T cell deletion. In all panels, *$p <$

0.05, **$p <$ 0.01, ***$p <$ 0.001; ns, non-significant. Values as mean ± SD and compared by the

Student's t test (**A**-**F**), one-way ANOVA multiple comparisons with Tukey's method (**I**), and

two-way ANOVA with Dunnett's multiple comparisons (**H**, **J**).

Fig. 5. Mechanism model.

Model depicting the molecular mechanism by which citalopram inhibit the Warburg effect and

elicit an anti-tumor response in HCC. Citalopram not only inhibits glycolytic metabolism of

cancer cells by targeting GLUT1 (Ref 16) but also enhances macrophage-driven anti-tumor

immunity and induces a systemic immunostimulatory effect on CD8$^+$ T cell functions through

serotonergic mechanisms.

Supplementary Materials

Figure S1. Citalopram inhibits HCC cell proliferation and promotes cell apoptosis in immune-competent and immune-deficient mouse models. (**A**, **B**) Immunohistochemical analysis of cleaved caspase-3 (CCS3) and Ki67 in Hepa1-6-bearing subcutaneous xenograft tumors from Rag1$^{-/-}$ or immunocompetent C57BL/6 mice, treated with DMSO or 5 mg/kg citalopram. (**C**, **D**) Immunohistochemical analysis of CCS3 and Ki67 in Hep53.4-bearing subcutaneous xenograft tumors from Rag1$^{-/-}$ or immunocompetent C57BL/6 mice, treated with DMSO or 5 mg/kg citalopram. In all panels, **$p < 0.01$, ***$p < 0.001$. Scale bar, 50 μm. Values as mean ± SD and compared by the Student's t test.

Figure S2. Citalopram suppresses HCC cell proliferation and promote cell apoptosis in

a SERT-independent manner. (**A**) Plate colony formation assay revealed the effect of SERT

knockdown alone or combined with citalopram treatment on the long-term cell proliferation of

Hepa1-6 and Hep53.4 cells (n = 6 per group). (**B**) Caspase-3/7 activity in Hepa1-6 and

Hep53.4 cells upon SERT knockdown or combined treatment with 5 μM citalopram (n = 6 per

group). (**C**, **D**) Immunohistochemical analysis of CCS3 and Ki67 in shControl and sh*Slc6a4*

Hepa1-6-bearing subcutaneous xenograft tumors from Rag1$^{-/-}$ or immunocompetent C57BL/6

mice, treated with DMSO or 5 mg/kg citalopram. Scale bar, 50 µm. (**E**, **F**)

Immunohistochemical analysis of CCS3 and Ki67 in shControl and sh*Slc6a4* Hep53.4-

bearing subcutaneous xenograft tumors from Rag1$^{-/-}$ or immunocompetent C57BL/6 mice,

treated with DMSO or 5 mg/kg citalopram. Scale bar, 50 µm. In all panels, *p < 0.05, **p <

0.01, ***p < 0.001. Scale bar, 50 µm. Values as mean ± SD and compared by one-way

ANOVA multiple comparisons with Tukey's method among groups.

Figure S3. Identification of C5aR1 as a direct target for citalopram. (**A**) The DARTS

assay and immunoblot analysis showed C5aR1 protein stability against 5 µg/mL pronase in

the presence of different concentrations of SSRIs treatment (0, 1, 10, 50, and 100 µM). (**B**)

For C5aR1, the predicted binding energy distribution of the clusters with poses more than 50.

(**C**) Sequencing analysis showed the successful generation of six C5aR1 mutants. (**D**) The

best-scored complex models of C5aR1 with other four different SSRIs. (**E**) HEK293T cells

were transfected with either WT or mutant C5aR1 (D282A) expression plasmids for 48 h,

followed by DARTS assay with immunoblotting analysis of C5aR1 protein levels. In all panels,

*p < 0.05. Values as mean ± SD and compared by one-way ANOVA multiple comparisons

with Tukey's method among groups (**E**).

Figure S4. Macrophage depletion mitigates the anti-tumor effects of citalopram. (**A**)
Schematic depicting macrophage blockade with clodronate liposomes. 12 days before tumor
inoculation, C57BL/6 mice were pretreated with clodronate liposomes or phosphate-buffered
saline (PBS) liposomes. Subsequently, GLUT1^{KD} Hepa1-6 cells were subcutaneously injected
into the C57BL/6 mice. The effect of citalopram (5 mg/kg) on the tumor burden was evaluated
after 18 days of drug treatment. (**B**) Immunofluorescence analysis of F4/80^{+} macrophages in
the liver and tumor tissues of indicated groups. (**C**) In C57BL/6 mice, the effect of citalopram
on the GLUT1^{KD} Hepa1-6 xenograft tumors was measured in the presence of macrophage
deletion (n = 7 per group). (**D**) Immunohistochemical analysis of CCS3 and Ki67 in GLUT1^{KD}
Hepa1-6-bearing subcutaneous xenograft tumors, treated with DMSO or 5 mg/kg citalopram
(n = 7 per group). Scale bar, 50 μm. Values as mean ± SD and compared by the Student's t
test (**C**, **D**) and two-way ANOVA with Dunnett's multiple comparisons (**C**).

Figure S5. Citalopram reverses C5a-mediated macrophage phagocytic impairment via targeting C5aR1. (**A**) Western blotting and immunofluorescence analysis showed SERT protein levels in Cas9-sgControl, -sg*C5ar1* THP-1 subclones. (**B**) Effects of C5aR1 deficiency on the macrophage phagocytosis of HCC-LM3 in the presence or absence of C5a stimulation. (**C**) Effects of different SSRIs on the macrophage phagocytosis of HCC-LM3 in the presence of C5a stimulation. (**D**) Reconstituted expression of WT and D282A mutant C5aR1 in C5aR1^{KO} THP-1 cells. (**E**) The effects of citalopram on macrophage phagocytosis in the absence of C5aR1 with reconstituted expression of C5aR1^{WT} or C5aR1^{D282A}. In all panels, *p < 0.05, **p < 0.01, ***p < 0.001. Values as mean ± SD and compared by one-way ANOVA multiple comparisons with Tukey's method among groups. Data are representative of three independent experiments (**B**, **C**, **E**).

Figure S6. Predicted binding modes of citalopram to human and mouse C5aR1. (**A**)

Conformations of orthosteric binding sites in human (light blue) and mouse (orange) C5aR1.

The conformation of human C5aR1 was obtained from the crystal structure (PDB id: 6c1q).

The structure of mouse C5aR1 was predicted using the ColabFold (AlphaFold2) software. (**B**)

The predicted binding modes of citalopram to human (light blue) and mouse (orange) C5aR1.

The conformations of citalopram were shown in pink (binding mode 1) or deep green (binding

mode 2) sticks. For mouse C5aR1, green sticks indicate residues set to flexible in the

molecular docking process.

Figure S7. Flow cytometry analysis of immune cells in tumor tissues and spleen.

(**A**, **B**) Gating strategies used for flow cytometry analysis of tumor and splenic lymphocytes.

Panel A: Identification of CD4$^+$ T cells, CD8$^+$ T cells, and dendritic cells (DC). Panel B:

Identification of B220$^+$ B cells, tumor-associated macrophages (TAMs), and nature killer (NK)

cells. (**C**) Flow cytometry showed the infiltration of CD45$^+$CD11b$^+$F4/80$^+$ macrophages, CD4$^+$

T cells, CD8$^+$ T cells, B220$^+$ B cells, CD11c$^+$ DC cells, and NK1.1$^+$ NK cells in spleen tissues

from orthotopic xenograft model, which generated in immunocompetent C57BL/6 mice with

Hepa1-6 cells (n = 5 per group). Values as mean ± SD and compared by the Student's t test.

Figure S8. The expression pattern of Glut1 and Glut3 in CD8⁺ T cells. (**A**) Real-time

qPCR analysis of *Glut1* and *Glut3* expression in intratumoral CD8⁺ T cells (n = 3 per group).

(**B**) Single cell analysis of the expression of glucose transporter members in two HCC cohorts.

Values as mean ± SD and compared by the Student's t test (**A**), **$p < 0.01$.

Figure S9. The *in vitro* effects of citalopram treatment on CD8+ T cell function. (**A**)

Splenic CD8+ T cells isolated from C57BL/6 mice were stimulated with plate-bound α-CD3

plus α-CD28 for 72 h, and T-cell proliferation (CFSE staining) upon citalopram treatment was

determined by flow cytometry (n = 3 per group); ACT: activated T cells. (**B**) Splenic CD8+ T

cells from C57BL/6 mice were stimulated with plate-bound α-CD3 and α-CD28 for 72 h, and

CD44 and CD62L levels in activated CD8+ T cells upon citalopram treatment were determined

by flow cytometry (n = 3 per group). (**C-E**) Intracellular GZMB, IFN-γ, and TNF-α were

detected in activated CD8+ T cells upon citalopram treatment (n = 3 per group). In all panels,

**p < 0.01, ***p < 0.001. Values as mean ± SD and compared by the Student's t test.

912