Title: Host transfer triggers widespread plasticity in gene expression, but not DNA

methylation, in a specialist herbivore

Running Title: Transcriptional plasticity enables host use

Author information

W. Nate Collison [1,2], Syed Adeel Zafar [1], Simon C. Groen [1,2], and Paul D. Nabity [1,3]

[1] Department of Botany and Plant Sciences, University of California, Riverside,

Riverside, CA, USA

[2] Department of Nematology, University of California, Riverside, Riverside, CA, USA

[3] School of BioSciences, University of Melbourne, Melbourne, Australia

Corresponding Author: Paul D. Nabity, School of BioSciences, University of Melbourne,

Melbourne, Australia, Paul.Nabity@unimelb.edu.au

Abstract

Insect herbivores employ many adaptive mechanisms to successfully colonize

host plants. Transcriptional plasticity is one mechanism that enables broad host use in

polyphagous insect herbivores; however, whether specialist herbivores use

transcriptional plasticity or undergo epigenome modifications to cope with host shifts

remains largely unexplored. We measured the transcriptome and epigenome of the

specialist aphid herbivore, *Daktulosphaira vitifoliae*, after transfer to new hosts using

RNA-seq and whole-genome bisulfite sequencing. We found the transcriptome of *D.*

vitifoliae is highly plastic and responsive to host genotype – 3,706 genes (28% of

expressed genes) are significantly differentially expressed in at least one contrast

comparing host genotypes. Most host-responsive genes (2,667 genes) were

upregulated when feeding on the non-native host *Vitis vinifera* and enriched for genes

encoding predicted secretory effectors, detoxification enzymes, and cuticle- and chitin-

associated proteins. We found gene co-expression modules that correlated with

variation in fitness suggesting plasticity in transcriptional networks contributes to *D.*

vitifoliae adaptation to host plants. We also identified a previously undescribed family of

proline-rich secretory proteins conserved across the Aphidomorpha infraorder of

insects. Unlike the transcriptome, DNA methylation did not dramatically respond to host

transfer and positively correlated with stably expressed genes; thus, gene body DNA

methylation may help stabilize gene expression when insects move to new host

environments. Together, these results provide new insight into the molecular

mechanisms of host use in insect herbivores.

Key Words: adaptation, aphids, effectors, grapevines, phenotypic plasticity,

methylome

Introduction

Phenotypic plasticity, or the expression of different phenotypes by the same

genotype, can allow organisms to flexibly respond to environmental changes, and has

thus been widely proposed to contribute to the evolution of species (Ancel, 2000;

Pigliucci & Murren, 2003; Ghalambor et al., 2007; Murren et al., 2015). However, the

extent to which plasticity alters adaptive evolution remains an open question (Vinton

et al., 2022). Insect herbivores are particularly good systems for understanding the

mechanisms of plasticity and adaptation, given their ecological diversity (Futuyma &

Agrawal, 2009; Forister et al., 2015), ability to rapidly adapt to new environments

(Garnas, 2018), and dramatic examples of environment-induced phenotypic variation

(Whitman & Agrawal, 2009). Although plant–herbivore interactions, and evolutionary

ecology more broadly, are well established fields within evolutionary biology, the

molecular mechanisms that underlie species interactions are only beginning to be

understood.

Insect herbivores are among the most species-rich lineages of metazoans, and

this diversity is largely driven by adaptations to specific host plants (Simon et al., 2015).

These adaptations function as a mode of ecological specialization that induces trade-

offs in fitness when an herbivorous insect transitions from ancestral to alternative hosts

(García-Robledo & Horvitz, 2012). Support for this hypothesis comes from the fact that

most insect herbivores are monophagous or oligophagous (i.e. specialists), feeding on

one or very few species of closely related hosts (Bernays & Graham, 1988; Futuyma &

Agrawal, 2009; Forister et al., 2015). For polyphagous, generalist herbivores,

compatibility across a breadth of divergent hosts is hypothesized to be maintained by a

high degree of genomic flexibility (Sexton et al., 2017; Birnbaum & Abbot, 2020).

Indeed, the genomes of generalist insect herbivores contain extraordinary expansions

of chemoreception and detoxification genes (W. Xu et al., 2016; Grbić et al., 2011;

Pearce et al., 2017), which enable host finding and feeding on plants. In contrast,

specialist herbivores appear to lack such expansions of chemoreception and

detoxification genes (Peláez et al., 2023). Additionally, there is increasing evidence that

both generalists and specialist herbivores rely on expanded families of secretory

effector proteins to directly manipulate host plant biology during feeding (Zhao et al.,

2015; Mathers et al., 2017; Rispe et al., 2020; Korgaonkar et al., 2021).

75   Transcriptional plasticity of expanded gene families has been linked to fitness

variation in generalist insect herbivores during interactions with divergent host plants

(Mathers et al., 2017; Snoeck et al., 2018), suggesting transcriptional plasticity plays an

adaptive role in these insects. Most studies that investigated transcriptional responses

of insects to different host plants used generalists, as these insects typically represent

major agricultural pests (Grbić et al., 2011; Mathers et al., 2017; Pearce et al., 2017).

Transcriptional plasticity is hypothesized to be reduced in specialist herbivores relative

to generalists (Birnbaum & Abbot, 2020), given host-specific evolution, i.e. reduced diet

breadth. Indeed, the few studies that measured transcriptomic responses to host

transfer in both specialist and generalist insects found overall reduced transcriptional

plasticity in specialists (Govind et al., 2010; Roy et al., 2016; Schweizer et al., 2017).

From an evolutionary perspective, specialization and the reduced capacity for plasticity

are expected to come at a cost, limiting adaptability of species amidst rapid

environmental change (Futuyma & Moreno, 1988; Price et al., 2003; Forister et al.,

2012; Vamosi et al., 2014; Murren et al., 2015; Day et al., 2016).

90   Dynamic regulation of the epigenome is thought to be a central mechanism

underlying rapid, environment-induced transcriptional plasticity (Duncan et al., 2014;

Kilvitis et al., 2017; Duncan et al., 2022). DNA methylation of cytosines, the covalent

addition of a methyl group to these residues, is a well-known epigenetic mark that

maintains functional gene regulation in many eukaryotic organisms (Zemach et al.,

2010; Schmitz et al., 2019). Cytosine methylation, hereafter DNA methylation, has been

well-studied in some insect lineages, primarily social Hymenoptera for its role in

mediating behavioral and phenotypic plasticity (Yan et al., 2014, 2015). Unlike in plants

and mammals, DNA methylation in insects occurs at relatively low levels and is found

primarily in gene body regions of actively expressed genes. Interestingly, some insects

lack discernable levels of DNA methylation in their genomes, suggesting its role in gene

regulation is dispensable (Bewick et al., 2016; Provataris et al., 2018). In insects with

abundant DNA methylation the observed variation has been linked to developmental

and phenotypic plasticity (Glastad et al., 2014, 2019); however, the direct role of DNA

methylation in regulating transcriptional plasticity in insects remains unclear. Early

studies in honey bees (*Apis mellifera*) found differential DNA methylation coincides with

differential gene expression (Kucharski et al., 2008; Li-Byarlay et al., 2013) whereas

recent studies suggest there is no correlation (Harris et al., 2019; Marshall et al., 2019;

Cardoso-Júnior et al., 2021). In other insects such as the green peach aphid (*Myzus*

persicae), differential DNA methylation correlated strongly with differential expression of

sex-linked genes (Mathers et al., 2019), suggesting DNA methylation may regulate

transcriptional plasticity and the emergence of different phenotypes. Plasticity in DNA

methylation may similarly regulate plasticity in herbivore phenotypes in response to

different host plants, although until our study here, no experiments have tested this

hypothesis. More broadly, the role of transcriptional plasticity and DNA methylation in

herbivore acclimatization and adaptation to hosts remains uncharacterized.

The degree to which phenotypic plasticity, and the underlying epigenetic and

transcriptional plasticity, contribute to adaptation and evolution remains contentious

(Pigliucci & Murren, 2003; Price et al., 2003; Ghalambor et al., 2007; Vinton et al.,

2022). Plasticity may slow the rate of genetic evolution by shielding genotypes from the

effects natural selection because one genotype expresses multiple phenotypes across

environments (Ancel, 2000; Price et al., 2003; Paenke et al., 2007). Alternatively,

plasticity that produces phenotypes which are favored by natural selection may later

become genetically assimilated, a case where plasticity results in adaptive evolution

(Price et al., 2003; Paenke et al., 2007; Lande, 2009). Whether plasticity is adaptive

depends on the strength and direction of selection. Plastic responses may move

individuals towards a fitness optimum (adaptative), away from a fitness optimum

(maladaptive), or have a neutral effect (Pigliucci & Murren, 2003; Ghalambor et al.,

2007). A separate but related question is *to what extent does environment-induced*

plasticity lead to phenotypic adaptation via transgenerational inheritance of epigenetic

marks? The answer to this question similarly remains contentious, and likely dependent

on the lineage or trait in question. The transfer of DNA methylation marks across

generations in mammals seems to be rare, because almost all DNA methylation is

erased before being reestablished during embryogenesis (Morgan et al., 2005; H. J.

Lee et al., 2014; S.-M. Lee & Surani, 2024). In insects, conversely, DNA methylation

erasure during embryogenesis does not occur (Harris et al., 2019; X. Xu et al., 2019),

and DNA methylation can be inherited across generations in *Nasonia* wasps (Wang et

al., 2016) and honey bees (Yagound et al., 2020). Thus, whether inheritance of

epigenetic marks can contribute to phenotypic adaptation in insects remains an open

question.

In our study, we examined host-induced variation in fitness, gene expression,

and DNA methylation in the specialist herbivore *Daktulosphaira vitifoliae* (Fitch). This

gall-forming aphid-like insect, commonly known as grape phylloxera, is in the

Phylloxeridae family (Aphidomorpha infraorder) of Hemipteran insects and feeds strictly

on grapevines (species within the *Vitis* genus of flowering plants) (Granett et al., 2001;

Metz et al., 2017). Although native to North America, *D. vitifoliae* has invaded grape

growing regions across the globe, becoming one of the most infamous pests of

grapevines (Rispe et al., 2020). The accidental introduction into France in the late

1800's caused the 'phylloxeric plague', wiping out thousands of vineyards and

threatening global grape production (Rispe et al., 2020). While agriculturally relevant, *D.*

vitifoliae also provides a model study system for ecological genomics and epigenetics.

Short generation times of asexually reproducing individuals allow for large clonal

populations to be reared and removes any confounding effects of genetic background

on transcriptional or epigenetic plasticity, which is a notable advantage in our study. The

obligate nature of *D. vitifoliae* should also strengthen the ability to detect how host

switching shapes trait plasticity, given the outcome is adapt or die. To explore these

traits, we investigate the transcriptional and epigenetic basis of host use in *D. vitifoliae* as it feeds on different host genotypes for a single generation. We test the hypothesis that plastic responses of the transcriptome mediate variation in *D. vitifoliae* fitness on different hosts, such that the direction of plasticity in transcriptional networks is positively correlated (adaptive) or negatively correlated (maladaptive) with fitness variation. We also analyze DNA methylation in the progeny of *D. vitifoliae* adults that fed on different hosts for one generation to characterize the phylloxerid epigenome. We test the hypothesis that environment-induced plasticity leads to predictable changes in the epigenome of *D. vitifoliae* after one generation in different host environments, providing new evidence, or lack thereof, for the role of DNA methylation in phenotypic adaptation in insects.

Materials and Methods

<u>Plant and insect rearing</u>

A clonal line of *D. vitifoliae* was established from a single female leaf gall collected on *V. rupestris* (cv St. George) near Davis, California U.S.A., in 2021. The clonal population was maintained for successive generations on *V. vinifera* cv Cabernet Sauvignon 33 (Cab33) that were grafted (*Vitis* genotype 1616 and 3309 rootstocks) and self-rooted in UC Riverside greenhouses in Riverside, CA (27 °C +/-3 °C) until experimentation during 2022-2023.

We used nine different grapevine genotypes as hosts for *D. vitifoliae* in our experiments, all of which were clonally propagated from cuttings purchased as rooted, overwintered 1-year-old plants from Inland Desert Nursery (Benton City, WA). Genotypes included self-rooted rootstock hybrids of *V. berlandieri* x *V. rupestris* (99R, 1103P), *V. berlandieri* x *V. riparia* (SO4, 5BB), and *V. riparia* x *V. rupestris* (101-14, SW, 3309). All plants were potted in 11.3 liter round pots with 'UC Mix 3' soil (https://agops.ucr.edu/soil-mixing#uc-soil-mix-iii) and grown in a greenhouse set at 24:27 °C +/- 3 °C (night:day) with a 14h photoperiod. After leaf bud swelling (8 days after potting), half of the plants for each genotype were infested. The remaining plants were potted and infested similarly 6 days later. Plantings were replicated through time to optimize sample collection effort given the time required to sort and count insects.

Host transfer and insect performance assay

For experimental infestation of new host plants (host transfer), galled leaves from

the colony on Cab33 (hereafter 'G0', for generation 0) were excised and placed on the

branch nodes of new host plants. Infestation was monitored every 12 hours to note

hatch and dispersal of 1st nymphal instars (hereafter generation 1, 'G1') onto budding

leaves. Galled leaves used for infestation were removed from branch nodes after two

days to prevent further infestation.

To assess insect performance, 'G1' galls were collected 15 days post infestation

(DPI). Galls were collected using a 4 mm diameter hole punch and carefully opened

under a dissecting microscope to count eggs per gall. Each plant was sampled five

different times, at 15, 17, 19, 21, and 23 DPI. Each time, the five largest galls from each

plant were sampled, selecting galls from leaves that had not been previously sampled

unless the largest remaining galls were on a previously sampled leaf. To determine if *D.*

vitifoliae fitness varied significantly on different hosts, pairwise Wilcoxon rank-sum tests

were performed (R v. 4.3.1).

RNA isolation and sequencing

To quantify *D. vitifoliae* transcriptome plasticity during host transfer events,

twenty G1 adults were collected from galls on unique plants and pooled for each

sample. Samples were collected 15-23 DPI and placed in tubes on ice during collection,

then immediately stored at -80 °C until RNA isolation. RNA was isolated from adults

using the RNeasy Mini Kit (Qiagen) following manufacturer's instructions. RNA

intactness and purity were verified with gel electrophoresis and spectrophotometry

(260:280 > 2; 260:230 >1.7) prior to library preparation using NEBNext® Ultra™ II

Directional RNA Library Prep Kit for Illumina® (New England Biolabs). RNA libraries

were sequenced using an Illumina NovaSeq X system at the DNA Technologies and

Expression Analysis Core of the UC Davis Genome Center, Davis, CA, with an average

yield of 28.9 ($\pm$ 3.29 s.d.) million paired-end reads per sample. Sequencing data were

uploaded to the NCBI Sequence Read Archive (SRA) repository under Bioproject

accession PRJNA1211173.

RNA-seq data processing

Fastqc v. 0.11.9 (https://github.com/s-andrews/FastQC) was used to generate

read and base quality statistics for raw and processed reads. Raw RNA-seq reads were

quality trimmed and filtered using BBTools bbduk (v.38.95; https://jgi.doe.gov/data-and-

tools/software-tools/bbtools/) with default settings plus options for kmer length (k=23)

and hamming distance (hist=1). Trimmed and filtered reads were aligned to the

reference genome (Rispe et al., 2020) using STAR v. 2.7.10b (Dobin et al., 2013) with

default settings plus options for maximum intron length (--alignIntronMax 15000) and

removing alignments that contained non-canonical junctions (--outFilterIntronMotifs

RemoveNoncanonical). On average, 85.8% ($\pm$ 1.49% s.d.) of reads uniquely mapped to

the reference genome, yielding 6.80 Gigabases ($\pm$ 0.74 s.d.) of read coverage (Table

S1). Only uniquely mapped reads were used for downstream analysis.

Differential gene expression analysis

Gene count matrices were generated using the --quantMode GeneCounts option

in STAR and analyzed using edgeR (v.4.0.16) (Robinson et al., 2010). Low count genes

were removed from downstream analysis using filterByExpr with default parameters.

Linear model fitting of count data was performed using limma's voom and lmFit

functions (Law et al., 2014). Genes were considered differentially expressed based on

the empirical Bayes method implemented using limma's 'treat' function with a 1.5-fold-

change filter and BH-adjusted p-value < 0.05. Genes that were differentially expressed

between 'G1' Cab33 and 'G0' (initial *D. vitifoliae* colony on Cab33) were removed for

downstream analysis of host-responsive genes, since these were differentially

expressed across generations on the same host genotype. Fisher's exact testing for

gene functional group enrichment was performed with the R stats package (v.4.3.1).

Functional groups were considered significantly enriched based on a Benjamini

Hochberg (BH)-adjusted p-value cutoff of 0.05 and an odds ratio $\geq$2. Functional groups

were defined based on published genome annotations (Rispe et al., 2020).

Gene co-expression network analysis

Genes with 0.5 counts per million (CPM; about 10 raw gene counts) in at least

four samples were included for network construction using WGCNA (Langfelder &

Horvath, 2008). Gene count matrices were normalized using DESeq2's vst (variance

stabilizing transformation) function (v. 1.42.1). A signed gene co-expression network

was constructed based on hierarchical clustering of dissimilarity in gene connectivity (1-

TOM) with the soft power parameter (β) set to 16 to ensure the network had

approximate scale-free topology. Genes were assigned to modules using the

cutreeDynamic function with minimum module size = 20 and module detection

sensitivity (deepSplit) = 2. Heat maps of module eigengenes and normalized gene

expression values (z-scores of vst-transformed count data) were made using pheatmap

v. 1.0.12 (https://github.com/raivokolde/pheatmap; rows were clustered based on

Euclidean distance. Correlations between *D. vitifoliae* fitness and gene co-expression

modules were tested by linear regression of average fitness on each host genotype and

average module eigengene expression using the lm function in the R stats package v.

4.3.1 where mean fitness = $\beta_0 + \beta_1 \times$ mean eigengene + ϵ. Scatter plots of eigengene-

fitness correlations were printed with results from the linear regression test using

ggplot2 v. 3.5.0 (https://ggplot2.tidyverse.org/). Gene co-expression modules were

tested for overrepresentation/enrichment of gene functional groups based on Fisher's

exact tests as described above.

<u>KEGG term enrichment analysis</u>

eggNOG-mapper v. 2 (Cantalapiedra et al., 2021) was used to assign KEGG

terms to the predicted proteome of *D. vitifoliae* based on precomputed orthology

assignments. eggNOG-mapper assigned KEGG terms to 7,532 proteins out of 15,076

queried. To test for KEGG term enrichment for subsets of genes (e.g. differentially

expressed genes vs all KEGG-annotated genes) we performed hypergeometric over-

representation analysis using the 'enrich' function of the R package clusterProfiler v.

4.10.1 (G. Yu et al., 2012). KEGG terms were considered significantly enriched based

on a BH-adjusted p-value cutoff of 0.05 and a gene count ≥4.

<u>Multiple Sequence Alignment and phylogenetic analysis</u>

To determine sequence similarity and phylogenetic relationships of *D. vitifoliae*

ARabinoGalactan-Like (ARGL) proteins and putative homologs (based on BLASTp

searches), we performed Multiple Sequence Alignment (MSA) using the Clustal Omega

web application (https://www.ebi.ac.uk/jdispatcher/msa/clustalo; Sievers & Higgins,

2018). Alignment results were visualized in a sequence logo plot using the WebLogo

web application (https://weblogo.threeplusone.com; Crooks et al., 2004). Phylogenetic

trees were constructed from the Clustal Omega MSA output using the Interactive Tree

of Life (iTOL) v. 6 web application (https://itol.embl.de; Letunic & Bork, 2024).

<u>Whole-genome bisulfite sequencing</u>

For WGBS, *D. vitifoliae* 'G0' adults and 'G0' and 'G1' eggs were collected from

leaf galls at 15-23 DPI and kept at -80 °C until DNA extraction. For 'G1', DNA was

isolated from pooled eggs collected from 'G1' galls, i.e. progeny of the 'generation 1'

adults used for RNA-seq. Galls from one plant were pooled together to make one

biological replicate. DNA was extracted from 200-250 eggs or 15-20 adults per replicate

using a CTAB protocol based on the method described by Mugford et al. (2020). The

quality and quantity of DNA was tested by running it on a 2% agarose gel and

measurement with a Qubit Fluorometer. We prepared 150 bp paired-end WGBS

libraries and used Illumina NovaSeq at the UC Davis Genome Center

(https://dnatech.ucdavis.edu/#C2). Library prep and sequencing for the 'G0' eggs and

adults was performed similarly by Novogene using Illumina NovaSeq

(https://www.novogene.com/us-en/services/research-services/epigenome-

sequencing/whole-genome-bisulfite-sequencing/). On average, we generated 47.2 ($\pm$

1.20 s.d.) million paired-end reads per sample (Table S1). Sequencing data were

uploaded to the NCBI Sequence Read Archive (SRA) repository under Bioproject

accession PRJNA1211173.

<u>DNA methylation analysis</u>

Read and base quality statistics were generated for raw and processed WGBS

reads using fastqc (version 0.11.9) (https://github.com/s-andrews/FastQC). Raw reads

were initially trimmed and filtered using Trim Galore (version 0.6.7)

(https://github.com/FelixKrueger/TrimGalore) with default parameters plus an 80 bp

minimal read length filter. Fastqc results showed the 5' and 3' ends of reads had

artifactual base sequences that were not removed by quality trimming, and libraries had

poor alignment efficiency when mapped to the bisulfite-converted reference genome

(Rispe et al., 2020). Thus, 25 bp at the 5' ends and 20 bp at the 3' ends were trimmed

from all reads using Trim Galore. Reads were mapped to the reference genome using

bismark v. 0.24.1 (Krueger & Andrews, 2011). On average, 59.4 % ($\pm$ 0.71 s.d) of reads

uniquely mapped to the bisulfite-converted reference genome (Table S1). After

removing PCR duplicate reads, the average coverage was 5.21 Gigabases ($\pm$ 1.64 s.d),

or 17.7x ($\pm$ 5.59 s.d) genome coverage (Table S1). Genome-wide CpG, CHG, and CHH

methylation percentages for each sample were extracted from the bismark alignment

report and plotted using ggplot2.

To determine which CpG sites were significantly methylated, a binomial test was

performed using the bisulfite conversion (C to T) error rate, which was 0.3% ($\pm$ 0 s.d,

n=4) based on the CpG methylation percentage of spiked-in lambda bacteriophage

DNA. CpG sites were filtered based on a BH-adjusted p-value cutoff of 0.001, which

removed CpG sites that had less than ~10% methylation (methylated reads divided by

all aligned reads). CpG sites were annotated with gene information (gene ID, functional

group) and genome features (UTRs, introns, exons, intergenic) using the 'intersect'

command of bedtools v. 2.30.0 (Quinlan & Hall, 2010). To assess global differences in

CpG methylation between samples, principal component analysis (PCA) was performed

using prcomp (R stats package v. 4.3.1). For PCA, we included only CpG sites that

were covered by $\geq$4 reads in all samples, were methylated in at least one sample

(binomial test, BH-adjusted p-value cutoff of 0.001), and had non-zero variance. To

determine if any *D. vitifoliae* genes were differentially methylated after feeding on

different hosts, we used a logistic regression with a F-test and applied a q-value cutoff

of 0.05 and a $\geq$10% methylation difference cutoff using the R package methylKit v.

1.28.0 (Akalin et al., 2012). The 'intersect' command of bedtools v. 2.30.0 was used to

determine if any differentially methylated genes were also differentially expressed (i.e.

host-responsive).

Additionally, we wanted to understand how gene expression plasticity correlated

with DNA methylation. We used all host-by-host contrasts to calculate the mean

absolute value of log2-fold-change in gene expression as a measure of transcriptional

plasticity. The 'G0' (starting colony) adult samples were used to calculate DNA

methylation levels for each gene, since these were sampled at the same developmental

stage (adults) as our RNA-seq data. Transcriptional plasticity and DNA methylation

percentage (% reads with methylated cytosines divided by total aligned reads) was

plotted for each gene in a scatterplot and a LOESS (Locally Estimated Scatterplot

Smoothing) line was fitted to the data to visualize the trend. We then tested for

correlation between transcriptional plasticity and DNA methylation with a Spearman

rank correlation test using the R stats package v. 4.3.1. Lastly, genes were binned into

quartiles based on the mean absolute value of log2-fold-change, then pairwise Wilcoxon

rank-sum tests were performed to test for significant differences in DNA methylation

levels between quartiles.

All scripts for data processing and bioinformatic analyses are available online at

https://github.com/wcoll010/Grape-phylloxera-phenotypic-plasticity-during-host-transfer.

Results

<u>Insect fitness differs significantly across host grapevine genotypes</u>

To determine how fitness differs across grapevine hosts, we experimentally

introduced a clonal line of *D. vitifoliae* to a panel of grapevine genotypes, including a

non-native host (Cab33) and interspecific hybrids of native North American grapevine

species (Figure 1). Fitness varied across host grapevine genotypes (Figure 1A) but the

grapevine genotypes SW (*V. riparia* X *V. rupestris*), 5BB (*V. riparia* X *V. berlandieri*),

and 3309 (*V. riparia* X *V. rupestris*) were incompatible, i.e. fully resistant to colonization.

Attempted feeding by *D. vitifoliae* induced small necrotic lesions on leaves of genotypes

SW and 5BB within six days after experimental infestation (Figure 1D and 1E).

Interestingly, genotype 3309 showed symptomless resistance after six days (Figure 1F)

but eventually showed necrotic lesions in leaves at feeding sites three weeks after

infestation (Figure S1). Fitness was lower on both Cab33 and SO4 (*V. berlandieri x V.*

riparia) compared to the other three susceptible hosts (Figure 1G). All other pairwise

comparisons between host genotypes revealed no differences in *D. vitifoliae* fitness.

<u>*D. vitifoliae* gene expression is highly dynamic and responsive to host grapevine</u>

<u>genotype</u>

To understand the genome-wide variation in *D. vitifoliae* gene expression

induced upon transfer to different host grapevine genotypes, we isolated RNA from *D.*

vitifoliae adults after feeding on new hosts for a single generation and performed RNA-

seq. Clustering of samples based on multidimensional scaling (MDS) indicated host

environment induces distinct changes in *D. vitifoliae* gene expression (Figure 1H).

Differential gene expression analysis revealed 3,706 were host-responsive (1.5-fold-

change in any pairwise comparison, BH-adjusted $p < 0.05$; Table 1), which was 28% of

the 13,125 total expressed genes. We then performed a Fisher's exact test to identify

gene functional groups that are enriched (over-represented) among the host-responsive

genes relative to the genome-wide background, which included all 25,808 *D. vitifoliae*

gene models. Genes encoding predicted secretory effectors (adj. $p = 1.7e-127$, odds

ratio = 3.30), developmental regulators (adj. $p = 1.6e-14$, odds ratio = 4.55), cuticle- and

chitin-associated proteins (adj. $p = 1.5e-06$, odds ratio = 3.03), detoxification enzymes

(adj. $p = 1.9e-05$, odds ratio = 2.14), and proteins involved in general metabolism (adj. p

= 8.9e-03, odds ratio = 5.23) were enriched among host-responsive genes (Table 1).

Pair-wise analysis of gene differential expression (DE) revealed that the non-

native host Cab33 induced the strongest change in gene expression compared to other

hosts; 3,432 (92.6%) of the total 3,706 host-responsive genes show differential

expression in contrasts between Cab33 and the other four host genotypes (Figure 2A).

Only 19 genes were differentially expressed in all contrasts with Cab33 (Figure 2B); 6 of

these genes encoded predicted secretory effectors, including 3 RING-type E3 ligases

(DV3020556, DV3022321, and DV3025927). We further investigated the types of genes

that are up- and down-regulated when *D. vitifoliae* feeds on Cab33 using gene

enrichment analysis. Of the host-responsive genes, 2,667 (72%) were upregulated

during interaction with Cab33 relative to all other hosts. Genes encoding predicted

effectors, cuticle- and chitin-associated proteins, and detoxication enzymes were

enriched among genes upregulated on Cab33 (Table S2). Enriched KEGG terms for

upregulated genes included glutamate receptors, fatty acyl-CoA reductases, cathepsin

B cysteine proteases, nicotinic acetylcholine receptors, and organic acid transporters

(Figure 2C). Genes downregulated during interaction with Cab33 are enriched for

functions related to development (Fisher's exact test, BH-adjusted p = 1.59e-09, odds

ratio = 6.36) and KEGG terms for lactose galactohydrolase/β-glucosidases (PFAM

domain GH1) and serine protease inhibitors (Figure 2D). Overall, these results suggest

that the *D. vitifoliae* transcriptome is highly plastic, especially at genes encoding

secretory proteins including putative effectors and genes involved with response to and

protection against environmental stress.

<u>Four gene co-expression modules co-vary with *D. vitifoliae* fitness</u>

To identify groups of genes that likely function together in shared biological

processes, we performed weighted gene co-expression network analysis using 13,917

genes after removing 'zero genes' that showed negligible expression. A signed gene co-

expression network was constructed that contained 88 modules based on hierarchical

clustering of gene connectivity dissimilarity (Figure 3A and 3B). Gene co-expression

modules were named based on colors randomly assigned by the WGCNA program

(Langfelder & Horvath, 2008). To identify groups of co-regulated genes (modules) that

might underlie differences in *D. vitifoliae* fitness across host genotypes, we tested for

correlation between module eigengenes (first principal component of the module and

the most representative gene) and fitness using linear regression tests. Five

eigengenes significantly correlated with *D. vitifoliae* fitness (Figure 3C-F); we excluded

the 'grey' module from further analysis since it is a bin for genes that cannot be

assigned to co-expression modules and are unlikely to be co-regulated.

We performed Fisher's exact tests to identify gene co-expression modules that

were enriched for host-responsive genes. Twenty-one out of the 88 modules were

enriched for host-responsive genes (odd ratio >2, BH-adjusted p <0.05) (Table S3),

which reflects the high number of transcriptionally plastic genes in the dataset. The

'darkorange2' module was both enriched for host-responsive genes and negatively

correlated with *D. vitifoliae* fitness (Figure 3E). Based on KEGG term enrichment

analysis, 'darkorange2' is overrepresented for genes that encode glycoside hydrolase

family 1 enzymes (PFAM GH1), and show distinct patterns of expression across host

genotypes (Figure S3). A small cluster of 24 genes is highly expressed during

interactions with native hosts 101-14 and 1103 and decreases in expression during

interactions with Cab33 and SO4, which are hosts that maintain relatively low *D.*

vitifoliae fitness. The remaining 66 genes in the module are grouped into a larger cluster

that shows the exact opposite trend in expression, largely increasing in expression

during interaction with Cab33 and SO4 relative to 101-14 and 1103. These results show

the 'darkorange2' module is highly responsive to host environment and may contribute

to fitness variation in *D. vitifoliae* during interaction with non-optimal hosts.

<u>One module contains a cluster of conserved and proline-rich secretory proteins</u>

The 'thistle' module was most strongly correlated with *D. vitifoliae* fitness (R^2 =

0.94, p = 0.005; Figure 3C-F) and enriched with genes encoding predicted secretory

effectors (Fisher's exact test, BH-adjusted p = 0.028, odds ratio = 3.32). Other than the

11 predicted effectors, the remaining 28 genes in this module did not belong to any

predicted functional group. Four predicted effectors were homologous to *Acyrthosiphon*

pisum 'lysine-rich arabinogalactan protein 19-like' (accession XP_001948476.1), based

on previously published BLASTp alignments (Rispe et al., 2020). We termed these

genes as encoding '<u>AR</u>abino<u>G</u>alactan-<u>L</u>ike' (ARGL) proteins. *ARGL* genes show highly

similar patterns of transcript expression across treatments, generally increasing in

expression during interaction with high-fitness native hosts 101-14, 1103, and 99R

relative to Cab33 (Figure 4B). We also noticed that the *ARGL* genes are located in a

tandem cluster in the genome on scaffold 40 (Figure 4A). Within this same ~50-kb

spatial cluster are two more predicted effector-encoding genes (DV3003881 and

DV3003883) and three 'null' genes (DV3003873, DV3003874, DV3003879); two of the

three null genes (DV3003873 and DV3003874) are also assigned to the 'thistle' module,

suggesting they are co-regulated with *ARGL* genes. To determine if *ARGL* genes have

highly similar sequences and shared protein motifs, we performed Multiple Sequence

Alignment (MSA) of the seven ARGL proteins using Clustal Omega (Sievers & Higgins,

2018). A sequence logo plot of the ARGL protein MSA showed that the proteins have a

high proportion of proline residues and are highly conserved, especially at proline

residues (Figure S4). We then constructed phylogenetic trees from the MSA of protein

sequences and coding sequences (CDS) for all 2,743 predicted secretory effector

proteins and found *ARGL* coding sequences clustered together into a monophyletic

group (Figure S5), suggesting they may be members of a gene family that evolved

through tandem duplication. Despite being highly conserved at the amino acid level and

clustered together based on coding sequences, ARGL protein sequences did not cluster

together in a monophyletic group (Figure S6). Thus, ARGL proteins may have some

structural and functional divergence from one another.

To identify putative orthologs of *ARGL* genes, we performed a BLASTp search

using the *ARGL* gene DV3003875 queried against the NCBI non-redundant protein

sequences (nr) database (https://blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE=Proteins). We

then performed reciprocal BLASTp alignments of the top 20 BLASTp hits against the *D.*

vitifoliae genome. Seventeen out of the 20 proteins produced reciprocal best hits for the

DV3003875 *ARGL* gene. We then investigated the evolution of the *ARGL* locus by

performing MSA of the seven ARGL proteins with the 17 putative orthologs of ARGL

DV3003875, which spanned adelgids (*Adelges cooleyi*) and eight aphid species (Table

S4). Remarkably, a roughly 200-amino-acid region of all 27 proteins is highly

conserved, especially at proline residues (Figure 4C and Figure S7). We also searched

the proteins for predicted functional domains using InterPro

(https://www.ebi.ac.uk/interpro/search/sequence/). Notably, all 24 proteins contain

predicted secretory signal peptides but generally lack any known functional domains,

with two exceptions. The *Aphis glycines* 'hypothetical protein AGLY_000967' (accession

KAE9545424.1) and the *Metopolophium dirhodum* 'uncharacterized protein

LOC132951494' (accession XP_060879307.1) contain the InterPro-unintegrated motif

(PRINT entry) PR01217 (Proline-rich extensin signature). Extensin proteins are notably

characterized as cell wall-associated proteins in plants (Moussu & Ingram, 2023),

although whether the PR01217 motif confers any similar function to plant extensin

proteins is unclear. Overall, our analysis of the 'thistle' co-expression module revealed

an under-described group of genes encoding predicted secretory proteins, that are

tightly co-regulated and positively correlated with fitness at the transcript level. ARGL

proteins also are proline-rich, highly conserved, and belong to a putative family of

secretory proteins seemingly widespread in the Aphidomorpha infraorder of Hemipteran

insects. Together, these results suggest *ARGL* genes play an important role in host use

in aphidomorph insects.

DNA methylation in *D. vitifoliae* is mostly stable

In addition to our analysis of the *D. vitifoliae* transcriptome, we analyzed genome-

wide patterns of DNA methylation using WGBS. Our goal was to first understand

patterns of DNA methylation in the *D. vitifoliae* epigenome and then to determine

whether host genotype induced changes in DNA methylation after insects fed on

different hosts for a single generation. DNA methylation in *D. vitifoliae* occurs

exclusively in the CpG context, and about 3.5-4% of CpG sites are methylated (Figure

5A), which is similar to what is reported in other Hemipteran insects (Bewick et al.,

2016; Mathers et al., 2019). CHH and CHG methylation rates were around 0.3%, which

is equal to the false positive rate. As in other insects, DNA methylation in *D. vitifoliae*

occurs mostly in exons and other gene body regions (Figure 5B).

To determine variation in DNA methylation across our samples at a global level,

principal component analysis (PCA) was performed using the 201,448 CpG sites that

were methylated in at least one sample and had non-zero variance. Our PCA revealed

no distinct clustering based on treatment group (host grapevine genotype) except for

the adult samples (Figure 5C). We repeated the PCA twice: once after removing the two

potential outlier samples, and then once more after further removing the adult samples.

The results similarly showed no distinct clustering based on host genotype (Figure S8).

We additionally tested whether any genes were differentially methylated between

insects reared on different host grapevine genotypes using methylKit (Akalin et al.,

2012). Eight genes were host-responsive at the DNA methylation level (Table S5), but

none of these genes were differentially expressed at the transcript level. All 3,706 genes

that were transcriptionally plastic (host-responsive) showed no significant change in

DNA methylation in the subsequent generation. These results show that DNA

methylation in *D. vitifoliae* varies across developmental stages (adults compared to

eggs) but is mostly not responsive to host environment. We also found that host-

induced variation in gene expression does not lead to any significant change in gene
body methylation in progeny.

<u>Gene transcriptional plasticity is negatively correlated with DNA methylation</u>

We wanted to further understand how host-induced transcriptional plasticity
correlates with DNA methylation in *D. vitifoliae*. We hypothesized that transcriptionally
plastic (host-responsive) genes lack DNA methylation, which might explain why host-
responsive genes acquired no significant changes in DNA methylation. This hypothesis
is supported by other data showing that DNA methylation in insects occurs primarily in
gene body regions of stably expressed genes (Glastad et al., 2016; Libbrecht et al.,
2016; Mathers et al., 2019). To quantify the degree of transcriptional plasticity for genes,
mean absolute values of log2 fold changes in gene expression across all host-by-host
contrasts were calculated. Gene-level transcriptional plasticity and DNA methylation
level was plotted for genes that were expressed and had CpG methylation data (n =
12,368) (Figure S9). There was a significant negative correlation between
transcriptional plasticity and DNA methylation (Spearman's ρ = -0.54, p < 2.2e-16). To
further assess correlation between gene expression plasticity and DNA methylation,
genes were binned into quartiles based on transcriptional plasticity and pair-wise
Wilcoxon rank-sum tests were performed. This also showed a negative correlation
between transcriptional plasticity and DNA methylation, with transcriptionally plastic
genes having significantly less methylation compared to stably expressed genes (Figure
5D). These data indicate gene body DNA methylation contributes to gene expression
regulation in *D. vitifoliae* and possibly stabilize transcriptional plasticity when *D. vitifoliae*
faces new host environments.

Discussion

The transcriptional and epigenetic mechanisms that enable parasites and
pathogens to colonize of new hosts remain only partially understood. In this study, we
found the transcriptome of the specialist insect *D. vitifoliae* is remarkably plastic, but
more so when colonizing a novel (non-native) host compared to native hosts. The
transcriptionally plastic genes that increased in expression were enriched for predicted

secretory effectors, detoxification enzymes, and cuticle- and chitin-associated proteins –

genes linked to the herbivore lifestyle (Karban, 2020; Peláez et al., 2023). We found

gene co-expression modules correlated with fitness, suggesting plasticity in

transcriptional networks contributes to *D. vitifoliae* adaptation to host plants. In contrast,

DNA methylation did not respond to host environment but negatively correlated with

transcriptional plasticity. These results highlight that rapid transcriptional change may

enable host shifts in a specialist herbivore similar to how polyphagous herbivores adapt

to feed on divergent hosts (Dermauw et al., 2013; Mathers et al., 2017; Snoeck et al.,

2018). These results also link a lack of DNA methylation to increasingly plastic genes

and suggest gene body methylation plays an important role in constitutive gene

expression during colonization of new hosts. Together these results help us understand

how transcriptional and epigenomic variation enable adaptation to novel hosts by

specialist herbivores.

Because *D. vitifoliae* is a specialist herbivore that evolved to feed on a narrow

range of hosts in the *Vitis* genus, we predicted that transcriptional plasticity induced by

transfer to different compatible hosts would be low; this has been shown for other

specialist insect species in similar experiments (Govind et al., 2010; Q. Yu et al., 2016;

Roy et al., 2016; Birnbaum et al., 2017; Schweizer et al., 2017). Contrary to this

prediction, the *D. vitifoliae* transcriptome was highly plastic. Of all expressed genes,

3,706 (28%) were host responsive (Table 1), and the majority of these were upregulated

while feeding on the non-native host European wine grape *V. vinifera* (cv Cab33) (Table

S2). Predicted detoxification genes were enriched among genes upregulated during

interaction with *V. vinifera*, and included numerous cytochrome P450s, glutathione S-

transferases and UDP-glycosyltransferases, and ABC transporters. These patterns

suggest that all three phases of xenobiotic detoxification are reprogrammed when *D.*

vitifoliae feeds on *V. vinifera* (Heidel-Fischer & Vogel, 2015), likely helping to overcome

novel host defenses. Predicted effectors were also enriched among genes upregulated

while feeding on *V. vinifera* (Table S2). This functional group includes over 2,700

secretory proteins with effector attributes, including putative effectors known to interact

with host grapevine proteins such as RING-type E3 ligases (Zhao et al., 2019).

Interestingly, three RING-type E3 ligases were consistently upregulated when *D.*

vitifoliae fed on the non-native host *V. vinifera* compared to the other hosts, supporting

the hypothesis that protein modification is a key pathway to coopt hosts (Zhao et al.,

2015; Huang et al., 2021). The subset of genes identified here serve as targets for

further functional characterization to validate mechanisms of host adaptation in *D.*

vitifoliae and improve management of grape against *D. vitifoliae.* More broadly, these

results show that at least some lineages of specialist herbivores, despite long-term

ecological specialization, rely on substantial transcriptional plasticity for ecological

persistence, contrary to predictions (Sexton et al., 2017; Birnbaum & Abbot, 2020).

Our analysis of KEGG terms similarly revealed that genes putatively involved

with insect adaptations were overrepresented among genes upregulated while feeding

on the non-native host *V. vinifera*. These genes included ionotropic glutamate receptors

which can function as chemosensory receptors (Benton et al., 2009; Croset et al.,

2010), fatty acyl-CoA reductases (FAR) which can modify cuticle composition (Li et al.,

2020; Di et al., 2023), and cathepsin B cysteine proteases (Figure 2C). Notably, aphid

cathepsin B proteases function as secretory effectors during interaction with divergent

host plants (Mathers et al., 2017). Other recent studies show *M. persicae* cathepsin B

genes are expressed in response to abiotic stresses and important for thermotolerance

(Liu et al., 2023; Zhang et al., 2025). Thus, cathepsin B proteins in related aphids

contribute to adaptation to different environments and may also underlie host adaptation

in *D. vitifoliae*. In summary, we found many of the genes that increase in expression

during colonization of a non-native host are putatively involved with key adaptations in

herbivorous insects – detoxification of plant defense metabolites, direct manipulation of

hosts via effector protein secretion, and cuticle modification – highlighting that specialist

lineages dramatically reprogram critical pathways to colonize new environments.

In addition to differential expression analysis, we wanted to understand how

transcriptional plasticity mediates host use by correlating clusters of co-expressed

genes (gene co-expression modules) with variation in *D. vitifoliae* fitness across

different hosts. Four gene co-expression modules were significantly correlated with

variation in fitness, indicating that plasticity in transcriptional networks can contribute to

host plant adaptation (Figure 3). The 'thistle' co-expression module most strongly

correlated with fitness (Figure 3C) and was enriched with genes coding for predicted

secretory effector proteins (Fisher's exact test, BH-adjusted p = 0.028), while the

remaining genes in the module were not assigned to any predicted functional group.

This module also contained several co-regulated genes that are organized as a spatial

cluster in the genome, encode predicted secretory effector proteins (ARGL; Figure 4),

and are shared among aphids indicating their likely presence in the common ancestor of

the Aphidomorpha lineage. *D. vitifoliae* ARGL proteins and putative ARGL orthologs

show a high degree of amino acid conservation and retained secretory function over the

course of more than 160 million years of evolution (Figure 4C and Table S4)(Ren et al.,

2013), highlighting an important role for ARGL proteins in Aphidomorph biology or

ecology. Some of the putative ARGL orthologs in aphids and adelgids (*A. cooleyi*)

contain predicted proline-rich extensin motifs, and other putative ARGL orthologs are

annotated as 'lysine-rich arabinogalactan protein 19-like', 'vegetative cell wall protein

gp1-like', and 'leucine-rich repeat extensin-like protein 1' (Table S4). Although the

function of insect ARGL proteins remains unclear, a common theme among annotations

is the sensing and maintenance of plant cell walls (Herger et al., 2019; Ellis et al., 2010;

Ferris et al., 2001). As a central regulator of cellular structure and immune signaling in

plants, the plant cell wall is a critical interface for herbivore-host interaction (Wolf, 2022;

Erb & Reymond, 2019), and thus a likely target of insect secretory effectors. Whether

ARGL proteins play a role in host cell wall manipulation or host interaction more broadly

requires further functional genetic analysis, although our computational analyses here

suggest they are important for host use across aphidomorph insects.

In addition to our analysis of transcriptomic plasticity induced by host transfer, we

analyzed DNA methylation in *D. vitifoliae* after host transfer. To our knowledge, no other

studies have measured plasticity in both the transcriptome and epigenome of an

herbivore during interaction with novel hosts. Thus, we lack data for how host plants

shape the epigenome of herbivores and on the correlation between gene expression

variation and epigenetic marks. Importantly, our data were collected across two different

generations of *D. vitifoliae*. We collected RNA-seq data from adult *D. vitifoliae* that fed

on new host plants for one generation and DNA methylation in the progeny eggs, which

allowed us to test for correlation between host-induced plasticity in gene expression and

variation in DNA methylation in the subsequent generation. Although the *D. vitifoliae*

transcriptome was highly plastic across host plants, there was little variation in DNA

methylation in the subsequent generation. We found eight genes that were differentially

methylated across host plants (Table S5); thus, we did find evidence for host-specific

epimutations that could affect host adaptation via change gene expression level

(Bonasio et al., 2012; Libbrecht et al., 2016; Mathers et al., 2019), transcript splicing (Li-

Byarlay et al., 2013; Glastad et al., 2016), or gene mutation rate (Elango et al., 2009).

However, none of these eight genes were differentially expressed across hosts or linked

to fitness variation and samples did not cluster based on host genotype (Figure 5C and

Figure S8). These patterns show that, unlike the transcriptome, DNA methylation is

mostly non-responsive to host plant genotype, and thus unlikely to mediate rapid

phenotypic adaptation. We found a significant negative correlation between

transcriptional plasticity and DNA methylation, however, with transcriptionally plastic

genes being generally depleted in DNA methylation relative to stably expressed genes

(Figure 5D and Figure S9). This is consistent with previous findings in other insects

(Bonasio et al., 2012; Libbrecht et al., 2016; Mathers et al., 2019). Given the adaptive

significance of transcriptional plasticity during host use and host shifts, DNA methylation

may play an important role by limiting plasticity of stably expressed genes for which

environment-induced plasticity is maladaptive.

Taken together, our results suggest that specialist herbivores, despite long-term

ecological specialization, can rapidly reprogram their transcriptomes to invade new

hosts, and that gene body methylation contributes to the regulation of host-induced

transcriptional plasticity. In this example, the switch to the novel host *V. vinifera* comes

at a fitness cost but *D. vitifoliae* still persists, highlighting a mechanism for adaptation to

increased host breadth that may occur with invasions and range expansions. More

broadly, these data provide new evidence for the epigenetic basis of host-induced

transcriptional plasticity and can serve as the basis for future functional genetic work to

validate genes involved with adaptation to host plants in aphidomorph insects.

Acknowledgments

We thank Conner Lay for help with collection of phenotypic data, Hung D. Lai for

greenhouse maintenance, and Holly Clark, Clay Clark, and Wei Zhang at the UC

Riverside genomics core for RNA library preparation. This project was supported by the

following grants: USDA-NIFA Hatch grant CA-R-BPS-5199-H and CDFA grant CDFA-

IAB #21-0426-000-SA awarded to PDN; The National Institute of General Medical

Sciences of the National Institutes of Health grant no. R35GM151194 awarded to SCG.

Author contributions

W.N.C., S.A.Z. and P.D.N. conceived the study. W.N.C. and S.A.Z. performed sample

collections and insect performance assays. S.A.Z. performed the DNA and RNA

extractions. W.N.C. performed data processing and analysis and wrote the manuscript

with advice from S.C.G and P.D.N. All authors read, edited, and approved the

manuscript.

References

Akalin, A., Kormaksson, M., Li, S., Garrett-Bakelman, F. E., Figueroa, M. E., Melnick, A., & Mason, C. E. (2012). methylKit: A comprehensive R package for the analysis of genome-wide DNA methylation profiles. *Genome Biology*, *13*(10), R87. https://doi.org/10.1186/gb-2012-13-10-r87

Ancel, L. W. (2000). Undermining the Baldwin Expediting Effect: Does Phenotypic Plasticity Accelerate Evolution? *Theoretical Population Biology*, *58*(4), 307–319. https://doi.org/10.1006/tpbi.2000.1484

Benton, R., Vannice, K. S., Gomez-Diaz, C., & Vosshall, L. B. (2009). Variant Ionotropic Glutamate Receptors as Chemosensory Receptors in Drosophila. *Cell*, *136*(1), 149–162. https://doi.org/10.1016/j.cell.2008.12.001

Bernays, E., & Graham, M. (1988). On the Evolution of Host Specificity in Phytophagous Arthropods. *Ecology*, *69*(4), 886–892. https://doi.org/10.2307/1941237

Bewick, A. J., Vogel, K. J., Moore, A. J., & Schmitz, R. J. (2016). Evolution of DNA Methylation across Insects. *Molecular Biology and Evolution*, msw264. https://doi.org/10.1093/molbev/msw264

Birnbaum, S. S. L., & Abbot, P. (2020). Gene Expression and Diet Breadth in Plant-Feeding Insects: Summarizing Trends. *Trends in Ecology & Evolution*, *35*(3), 259–277. https://doi.org/10.1016/j.tree.2019.10.014

Birnbaum, S. S. L., Rinker, D. C., Gerardo, N. M., & Abbot, P. (2017). Transcriptional profile and differential fitness in a specialist milkweed insect across host plants varying in toxicity. *Molecular Ecology*, *26*(23), 6742–6761. https://doi.org/10.1111/mec.14401

Bonasio, R., Li, Q., Lian, J., Mutti, N. S., Jin, L., Zhao, H., Zhang, P., Wen, P., Xiang, H., Ding, Y., Jin, Z., Shen, S. S., Wang, Z., Wang, W., Wang, J., Berger, S. L., Liebig, J., Zhang, G., & Reinberg, D. (2012). Genome-wide and Caste-Specific DNA Methylomes of the Ants Camponotus floridanus and Harpegnathos saltator. *Current Biology*, *22*(19), 1755–1764. https://doi.org/10.1016/j.cub.2012.07.042

Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P., & Huerta-Cepas, J. (2021). eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain Prediction at the Metagenomic Scale. *Molecular Biology and Evolution*, *38*(12), 5825–5829. https://doi.org/10.1093/molbev/msab293

Cardoso-Júnior, C. A. M., Yagound, B., Ronai, I., Remnant, E. J., Hartfelder, K., & Oldroyd, B. P. (2021). DNA methylation is not a driver of gene expression reprogramming in young honey bee workers. *Molecular Ecology*, *30*(19), 4804–4818. https://doi.org/10.1111/mec.16098

Crooks, G. E., Hon, G., Chandonia, J.-M., & Brenner, S. E. (2004). WebLogo: A Sequence Logo Generator: Figure 1. *Genome Research*, *14*(6), 1188–1190. https://doi.org/10.1101/gr.849004

Croset, V., Rytz, R., Cummins, S. F., Budd, A., Brawand, D., Kaessmann, H., Gibson, T. J., & Benton, R. (2010). Ancient Protostome Origin of Chemosensory Ionotropic Glutamate Receptors and the Evolution of Insect Taste and Olfaction. *PLoS Genetics*, *6*(8), e1001064. https://doi.org/10.1371/journal.pgen.1001064

Day, E. H., Hua, X., & Bromham, L. (2016). Is specialization an evolutionary dead end? Testing for differences in speciation, extinction and trait transition rates across diverse phylogenies of specialists and generalists. *Journal of Evolutionary Biology*, *29*(6), 1257–1267. https://doi.org/10.1111/jeb.12867

Dermauw, W., Wybouw, N., Rombauts, S., Menten, B., Vontas, J., Grbić, M., Clark, R. M., Feyereisen, R., & Van Leeuwen, T. (2013). A link between host plant adaptation and pesticide resistance in the polyphagous spider mite *Tetranychus urticae*. *Proceedings of the National Academy of Sciences*, *110*(2). https://doi.org/10.1073/pnas.1213214110

Di, Z., Tian, J., Ali, M. Y., Zhang, B., Yang, S., Fu, Y., Li, F., & Luo, C. (2023). Diversity, domain duplication, birth-and-death and adaptive evolution of fatty acyl- CoA reductase genes in HEMIPTERA. *Physiological Entomology*, *48*(2–3), 97–106. https://doi.org/10.1111/phen.12413

Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P., Chaisson, M., & Gingeras, T. R. (2013). STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics*, *29*(1), 15–21. https://doi.org/10.1093/bioinformatics/bts635

Duncan, E. J., Cunningham, C. B., & Dearden, P. K. (2022). Phenotypic Plasticity: What Has DNA Methylation Got to Do with It? *Insects*, *13*(2), 110. https://doi.org/10.3390/insects13020110

Duncan, E. J., Gluckman, P. D., & Dearden, P. K. (2014). Epigenetics, plasticity, and evolution: How do we link epigenetic change to phenotype? *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution*, *322*(4), 208–220. https://doi.org/10.1002/jez.b.22571

Elango, N., Hunt, B. G., Goodisman, M. A. D., & Yi, S. V. (2009). DNA methylation is widespread and associated with differential gene expression in castes of the

honeybee, *Apis mellifera*. *Proceedings of the National Academy of Sciences*, *106*(27), 11206–11211. https://doi.org/10.1073/pnas.0900301106

Ellis, M., Egelund, J., Schultz, C. J., & Bacic, A. (2010). Arabinogalactan-Proteins: Key Regulators at the Cell Surface? *Plant Physiology*, *153*(2), 403–419. https://doi.org/10.1104/pp.110.156000

Erb, M., & Reymond, P. (2019). Molecular Interactions Between Plants and Insect Herbivores. *Annual Review of Plant Biology*, *70*(1), 527–557. https://doi.org/10.1146/annurev-arplant-050718-095910

Ferris, P. J., Woessner, J. P., Waffenschmidt, S., Kilz, S., Drees, J., & Goodenough, U. W. (2001). Glycosylated Polyproline II Rods with Kinks as a Structural Motif in Plant Hydroxyproline-Rich Glycoproteins. *Biochemistry*, *40*(9), 2978–2987. https://doi.org/10.1021/bi0023605

Forister, M. L., Dyer, L. A., Singer, M. S., Stireman, J. O., & Lill, J. T. (2012). Revisiting the evolution of ecological specialization, with emphasis on insect–plant interactions. *Ecology*, *93*(5), 981–991. https://doi.org/10.1890/11-0650.1

Forister, M. L., Novotny, V., Panorska, A. K., Baje, L., Basset, Y., Butterill, P. T., Cizek, L., Coley, P. D., Dem, F., Diniz, I. R., Drozd, P., Fox, M., Glassmire, A. E., Hazen, R., Hrcek, J., Jahner, J. P., Kaman, O., Kozubowski, T. J., Kursar, T. A., … Dyer, L. A. (2015). The global distribution of diet breadth in insect herbivores. *Proceedings of the National Academy of Sciences*, *112*(2), 442–447. https://doi.org/10.1073/pnas.1423042112

Futuyma, D. J., & Agrawal, A. A. (2009). Macroevolution and the biological diversity of plants and herbivores. *Proceedings of the National Academy of Sciences*, *106*(43), 18054–18061. https://doi.org/10.1073/pnas.0904106106

Futuyma, D. J., & Moreno, G. (1988). *THE EVOLUTION OF ECOLOGICAL SPECIALIZATION*.

García-Robledo, C., & Horvitz, C. C. (2012). Jack of all trades masters novel host plants: Positive genetic correlations in specialist and generalist insect herbivores expanding their diets to novel hosts: Quantitative genetics of diet expansion. *Journal of Evolutionary Biology*, *25*(1), 38–53. https://doi.org/10.1111/j.1420-9101.2011.02401.x

Garnas, J. R. (2018). Rapid evolution of insects to global environmental change: Conceptual issues and empirical gaps. *Current Opinion in Insect Science*, *29*, 93–101. https://doi.org/10.1016/j.cois.2018.07.013

Ghalambor, C. K., McKAY, J. K., Carroll, S. P., & Reznick, D. N. (2007). Adaptive versus non-adaptive phenotypic plasticity and the potential for contemporary adaptation in new environments. *Functional Ecology*, *21*(3), 394–407. https://doi.org/10.1111/j.1365-2435.2007.01283.x

Glastad, K. M., Gokhale, K., Liebig, J., & Goodisman, M. A. D. (2016). The caste- and sex-specific DNA methylome of the termite Zootermopsis nevadensis. *Scientific Reports*, *6*(1), 37110. https://doi.org/10.1038/srep37110

Glastad, K. M., Hunt, B. G., & Goodisman, M. A. (2014). Evolutionary insights into DNA methylation in insects. *Current Opinion in Insect Science*, *1*, 25–30. https://doi.org/10.1016/j.cois.2014.04.001

Glastad, K. M., Hunt, B. G., & Goodisman, M. A. D. (2019). Epigenetics in Insects: Genome Regulation and the Generation of Phenotypic Diversity. *Annual Review*

of Entomology, *64*(1), 185–203. https://doi.org/10.1146/annurev-ento-011118-111914

Govind, G., Mittapalli, O., Griebel, T., Allmann, S., Böcker, S., & Baldwin, I. T. (2010). Unbiased Transcriptional Comparisons of Generalist and Specialist Herbivores Feeding on Progressively Defenseless Nicotiana attenuata Plants. *PLoS ONE*, *5*(1), e8735. https://doi.org/10.1371/journal.pone.0008735

Granett, J., Walker, M. A., Kocsis, L., & Omer, A. D. (2001). Biology and Management of Grape Phylloxera. *Annual Review of Entomology*, *46*(1), 387–412. https://doi.org/10.1146/annurev.ento.46.1.387

Grbić, M., Van Leeuwen, T., Clark, R. M., Rombauts, S., Rouzé, P., Grbić, V., Osborne, E. J., Dermauw, W., Thi Ngoc, P. C., Ortego, F., Hernández-Crespo, P., Diaz, I., Martinez, M., Navajas, M., Sucena, É., Magalhães, S., Nagy, L., Pace, R. M., Djuranović, S., … Van De Peer, Y. (2011). The genome of Tetranychus urticae reveals herbivorous pest adaptations. *Nature*, *479*(7374), 487–492. https://doi.org/10.1038/nature10640

Harris, K. D., Lloyd, J. P. B., Domb, K., Zilberman, D., & Zemach, A. (2019). DNA methylation is maintained with high fidelity in the honey bee germline and exhibits global non-functional fluctuations during somatic development. *Epigenetics & Chromatin*, *12*(1), 62. https://doi.org/10.1186/s13072-019-0307-4

Heidel-Fischer, H. M., & Vogel, H. (2015). Molecular mechanisms of insect adaptation to plant secondary compounds. *Current Opinion in Insect Science*, *8*, 8–14. https://doi.org/10.1016/j.cois.2015.02.004

Herger, A., Dünser, K., Kleine-Vehn, J., & Ringli, C. (2019). Leucine-Rich Repeat Extensin Proteins and Their Role in Cell Wall Sensing. *Current Biology*, *29*(17), R851–R858. https://doi.org/10.1016/j.cub.2019.07.039

Huang, W., MacLean, A. M., Sugio, A., Maqbool, A., Busscher, M., Cho, S.-T., Kamoun, S., Kuo, C.-H., Immink, R. G. H., & Hogenhout, S. A. (2021). Parasitic modulation of host development by ubiquitin-independent protein degradation. *Cell*, *184*(20), 5201-5214.e12. https://doi.org/10.1016/j.cell.2021.08.029

Karban, R. (2020). The ecology and evolution of induced responses to herbivory and how plants perceive risk. *Ecological Entomology*, *45*(1), 1–9. https://doi.org/10.1111/een.12771

Kilvitis, H. J., Hanson, H., Schrey, A. W., & Martin, L. B. (2017). Epigenetic Potential as a Mechanism of Phenotypic Plasticity in Vertebrate Range Expansions. *Integrative and Comparative Biology*, *57*(2), 385–395. https://doi.org/10.1093/icb/icx082

Korgaonkar, A., Han, C., Lemire, A. L., Siwanowicz, I., Bennouna, D., Kopec, R. E., Andolfatto, P., Shigenobu, S., & Stern, D. L. (2021). A novel family of secreted insect proteins linked to plant gall development. *Current Biology*, *31*(9), 1836-1849.e12. https://doi.org/10.1016/j.cub.2021.01.104

Kucharski, R., Maleszka, J., Foret, S., & Maleszka, R. (2008). Nutritional Control of Reproductive Status in Honeybees via DNA Methylation. *Science*, *319*(5871), 1827–1830. https://doi.org/10.1126/science.1153069

Lande, R. (2009). Adaptation to an extraordinary environment by evolution of phenotypic plasticity and genetic assimilation. *Journal of Evolutionary Biology*, *22*(7), 1435–1446. https://doi.org/10.1111/j.1420-9101.2009.01754.x

Langfelder, P., & Horvath, S. (2008). WGCNA: An R package for weighted correlation
network analysis. *BMC Bioinformatics*, *9*(1), 559. https://doi.org/10.1186/1471-
2105-9-559
Law, C. W., Chen, Y., Shi, W., & Smyth, G. K. (2014). voom: Precision weights unlock
linear model analysis tools for RNA-seq read counts. *Genome Biology*, *15*(2),
R29. https://doi.org/10.1186/gb-2014-15-2-r29
Lee, H. J., Hore, T. A., & Reik, W. (2014). Reprogramming the Methylome: Erasing
Memory and Creating Diversity. *Cell Stem Cell*, *14*(6), 710–719.
https://doi.org/10.1016/j.stem.2014.05.008
Lee, S.-M., & Surani, M. A. (2024). Epigenetic reprogramming in mouse and human
primordial germ cells. *Experimental & Molecular Medicine*, *56*(12), 2578–2587.
https://doi.org/10.1038/s12276-024-01359-z
Letunic, I., & Bork, P. (2024). Interactive Tree of Life (iTOL) v6: Recent updates to the
phylogenetic tree display and annotation tool. *Nucleic Acids Research*, *52*(W1),
W78–W82. https://doi.org/10.1093/nar/gkae268
Li, D., Dai, Y., Chen, X., Wang, X., Li, Z., Moussian, B., & Zhang, C. (2020). Ten fatty
acyl-CoA reductase family genes were essential for the survival of the destructive
rice pest, NILAPARVATA LUGENS. *Pest Management Science*, *76*(7), 2304–2315.
https://doi.org/10.1002/ps.5765
Libbrecht, R., Oxley, P. R., Keller, L., & Kronauer, D. J. C. (2016). Robust DNA
Methylation in the Clonal Raider Ant Brain. *Current Biology*, *26*(3), 391–395.
https://doi.org/10.1016/j.cub.2015.12.040
Li-Byarlay, H., Li, Y., Stroud, H., Feng, S., Newman, T. C., Kaneda, M., Hou, K. K.,
Worley, K. C., Elsik, C. G., Wickline, S. A., Jacobsen, S. E., Ma, J., & Robinson,
G. E. (2013). RNA interference knockdown of *DNA methyl-transferase 3* affects
gene alternative splicing in the honey bee. *Proceedings of the National Academy*
of Sciences, *110*(31), 12750–12755. https://doi.org/10.1073/pnas.1310735110
Liu, J., Liu, Y., Li, Q., & Lu, Y. (2023). Heat shock protein 70 and Cathepsin B genes are
involved in the thermal tolerance of *Aphis gossypii*. *Pest Management Science*,
79(6), 2075–2086. https://doi.org/10.1002/ps.7384
Marshall, H., Lonsdale, Z. N., & Mallon, E. B. (2019). Methylation and gene expression
differences between reproductive and sterile bumblebee workers. *Evolution*
Letters, *3*(5), 485–499. https://doi.org/10.1002/evl3.129
Mathers, T. C., Chen, Y., Kaithakottil, G., Legeai, F., Mugford, S. T., Baa-Puyoulet, P.,
Bretaudeau, A., Clavijo, B., Colella, S., Collin, O., Dalmay, T., Derrien, T., Feng,
H., Gabaldón, T., Jordan, A., Julca, I., Kettles, G. J., Kowitwanich, K., Lavenier,
D., … Hogenhout, S. A. (2017). Rapid transcriptional plasticity of duplicated gene
clusters enables a clonally reproducing aphid to colonise diverse plant species.
Genome Biology, *18*(1), 27. https://doi.org/10.1186/s13059-016-1145-3
Mathers, T. C., Mugford, S. T., Percival-Alwyn, L., Chen, Y., Kaithakottil, G., Swarbreck,
D., Hogenhout, S. A., & Van Oosterhout, C. (2019). Sex-specific changes in the
aphid DNA methylation landscape. *Molecular Ecology*, *28*(18), 4228–4241.
https://doi.org/10.1111/mec.15216
Metz, M. A., Miller, D. R., Dickey, A. M., Bauchan, G. R., Ochoa, R., Skvarla, M. J., &
Miller, G. L. (2017). Rediscovering digitules in Aphidomorpha and the question of

homology among Sternorrhyncha (Insecta, Hemiptera). *ZooKeys*, *683*, 39–50. https://doi.org/10.3897/zookeys.683.10100

Morgan, H. D., Santos, F., Green, K., Dean, W., & Reik, W. (2005). Epigenetic reprogramming in mammals. *Human Molecular Genetics*, *14*(suppl_1), R47–R58. https://doi.org/10.1093/hmg/ddi114

Moussu, S., & Ingram, G. (2023). The EXTENSIN enigma. *The Cell Surface*, *9*, 100094. https://doi.org/10.1016/j.tcsw.2023.100094

Mugford, S., Wouters, R., C Mathers, T., & Hogenhout, S. (2020). *High quality DNA extraction from very small individual insects v1*. https://doi.org/10.17504/protocols.io.bg6wjzfe

Murren, C. J., Auld, J. R., Callahan, H., Ghalambor, C. K., Handelsman, C. A., Heskel, M. A., Kingsolver, J. G., Maclean, H. J., Masel, J., Maughan, H., Pfennig, D. W., Relyea, R. A., Seiter, S., Snell-Rood, E., Steiner, U. K., & Schlichting, C. D. (2015). Constraints on the evolution of phenotypic plasticity: Limits and costs of phenotype and plasticity. *Heredity*, *115*(4), 293–301. https://doi.org/10.1038/hdy.2015.8

Paenke, I., Sendhoff, B., & Kawecki, T. J. (2007). Influence of Plasticity and Learning on Evolution under Directional Selection. *The American Naturalist*, *170*(2), E47–E58. https://doi.org/10.1086/518952

Pearce, S. L., Clarke, D. F., East, P. D., Elfekih, S., Gordon, K. H. J., Jermiin, L. S., McGaughran, A., Oakeshott, J. G., Papanikolaou, A., Perera, O. P., Rane, R. V., Richards, S., Tay, W. T., Walsh, T. K., Anderson, A., Anderson, C. J., Asgari, S., Board, P. G., Bretschneider, A., … Wu, Y. D. (2017). Genomic innovations, transcriptional plasticity and gene loss underlying the evolution and divergence of two highly polyphagous and invasive Helicoverpa pest species. *BMC Biology*, *15*(1), 63. https://doi.org/10.1186/s12915-017-0402-6

Peláez, Julianne N., et al. "Evolution of chemosensory and detoxification gene families across herbivorous Drosophilidae." G3: Genes, Genomes, Genetics 13.8 (2023): jkad133

Pigliucci, M., & Murren, C. J. (2003). PERSPECTIVE: GENETIC ASSIMILATION AND A POSSIBLE EVOLUTIONARY PARADOX: CAN MACROEVOLUTION SOMETIMES BE SO FAST AS TO PASS US BY? *Evolution*, *57*(7), 1455–1464. https://doi.org/10.1111/j.0014-3820.2003.tb00354.x

Price, T. D., Qvarnström, A., & Irwin, D. E. (2003). The role of phenotypic plasticity in driving genetic evolution. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, *270*(1523), 1433–1440. https://doi.org/10.1098/rspb.2003.2372

Provataris, P., Meusemann, K., Niehuis, O., Grath, S., & Misof, B. (2018). Signatures of DNA Methylation across Insects Suggest Reduced DNA Methylation Levels in Holometabola. *Genome Biology and Evolution*, *10*(4), 1185–1197. https://doi.org/10.1093/gbe/evy066

Quinlan, A. R., & Hall, I. M. (2010). BEDTools: A flexible suite of utilities for comparing genomic features. *Bioinformatics*, *26*(6), 841–842. https://doi.org/10.1093/bioinformatics/btq033

Ren, Z., Zhong, Y., Kurosu, U., Aoki, S., Ma, E., Von Dohlen, C. D., & Wen, J. (2013). Historical biogeography of Eastern Asian–Eastern North American disjunct

Melaphidina aphids (Hemiptera: Aphididae: Eriosomatinae) on Rhus hosts (Anacardiaceae). *Molecular Phylogenetics and Evolution*, *69*(3), 1146–1158. https://doi.org/10.1016/j.ympev.2013.08.003

Rispe, C., Legeai, F., Nabity, P. D., Fernández, R., Arora, A. K., Baa-Puyoulet, P., Banfill, C. R., Bao, L., Barberà, M., Bouallègue, M., Bretaudeau, A., Brisson, J. A., Calevro, F., Capy, P., Catrice, O., Chertemps, T., Couture, C., Delière, L., Douglas, A. E., … Tagu, D. (2020a). The genome sequence of the grape phylloxera provides insights into the evolution, adaptation, and invasion routes of an iconic pest. *BMC Biology*, *18*(1), 90. https://doi.org/10.1186/s12915-020-00820-5

Rispe, C., Legeai, F., Nabity, P. D., Fernández, R., Arora, A. K., Baa-Puyoulet, P., Banfill, C. R., Bao, L., Barberà, M., Bouallègue, M., Bretaudeau, A., Brisson, J. A., Calevro, F., Capy, P., Catrice, O., Chertemps, T., Couture, C., Delière, L., Douglas, A. E., … Tagu, D. (2020b). The genome sequence of the grape phylloxera provides insights into the evolution, adaptation, and invasion routes of an iconic pest. *BMC Biology*, *18*(1), 90. https://doi.org/10.1186/s12915-020-00820-5

Robinson, M. D., McCarthy, D. J., & Smyth, G. K. (2010). edgeR: A Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*, *26*(1), 139–140. https://doi.org/10.1093/bioinformatics/btp616

Roy, A., Walker, W. B., Vogel, H., Chattington, S., Larsson, M. C., Anderson, P., Heckel, D. G., & Schlyter, F. (2016). Diet dependent metabolic responses in three generalist insect herbivores Spodoptera spp. *Insect Biochemistry and Molecular Biology*, *71*, 91–105. https://doi.org/10.1016/j.ibmb.2016.02.006

Schmitz, R. J., Lewis, Z. A., & Goll, M. G. (2019). DNA Methylation: Shared and Divergent Features across Eukaryotes. *Trends in Genetics*, *35*(11), 818–827. https://doi.org/10.1016/j.tig.2019.07.007

Schweizer, F., Heidel-Fischer, H., Vogel, H., & Reymond, P. (2017). Arabidopsis glucosinolates trigger a contrasting transcriptomic response in a generalist and a specialist herbivore. *Insect Biochemistry and Molecular Biology*, *85*, 21–31. https://doi.org/10.1016/j.ibmb.2017.04.004

Sexton, J. P., Montiel, J., Shay, J. E., Stephens, M. R., & Slatyer, R. A. (2017). Evolution of Ecological Niche Breadth. *Annual Review of Ecology, Evolution, and Systematics*, *48*(1), 183–206. https://doi.org/10.1146/annurev-ecolsys-110316-023003

Sievers, F., & Higgins, D. G. (2018). Clustal Omega for making accurate alignments of many protein sequences. *Protein Science*, *27*(1), 135–145. https://doi.org/10.1002/pro.3290

Simon, J.-C., d'Alencon, E., Guy, E., Jacquin-Joly, E., Jaquiery, J., Nouhaud, P., Peccoud, J., Sugio, A., & Streiff, R. (2015). Genomics of adaptation to host-plants in herbivorous insects. *Briefings in Functional Genomics*, *14*(6), 413–423. https://doi.org/10.1093/bfgp/elv015

Snoeck, S., Wybouw, N., Van Leeuwen, T., & Dermauw, W. (2018). Transcriptomic Plasticity in the Arthropod Generalist *Tetranychus urticae* Upon Long-Term Acclimation to Different Host Plants. *G3 Genes|Genomes|Genetics*, *8*(12), 3865–3879. https://doi.org/10.1534/g3.118.200585

Vamosi, J. C., Armbruster, W. S., & Renner, S. S. (2014). Evolutionary ecology of
specialization: Insights from phylogenetic analysis. *Proceedings of the Royal*
Society B: Biological Sciences, *281*(1795), 20142004.
https://doi.org/10.1098/rspb.2014.2004
Vinton, A. C., Gascoigne, S. J. L., Sepil, I., & Salguero-Gómez, R. (2022). Plasticity's
role in adaptive evolution depends on environmental change components.
Trends in Ecology & Evolution, *37*(12), 1067–1078.
https://doi.org/10.1016/j.tree.2022.08.008
Wang, X., Werren, J. H., & Clark, A. G. (2016). Allele-Specific Transcriptome and
Methylome Analysis Reveals Stable Inheritance and Cis-Regulation of DNA
Methylation in Nasonia. *PLOS Biology*, *14*(7), e1002500.
https://doi.org/10.1371/journal.pbio.1002500
Whitman, D., & Agrawal, A. (2009). What is Phenotypic Plasticity and Why is it
Important? In D. Whitman & T. Ananthakrishnan (Eds.), *Phenotypic Plasticity of*
Insects. Science Publishers. https://doi.org/10.1201/b10201-2
Wolf, S. (2022). Cell Wall Signaling in Plant Development and Defense. *Annual Review*
of Plant Biology, *73*(1), 323–353. https://doi.org/10.1146/annurev-arplant-
102820-095312
Xu, W., Papanicolaou, A., Zhang, H.-J., & Anderson, A. (2016). Expansion of a bitter
taste receptor family in a polyphagous insect herbivore. *Scientific Reports*, *6*(1),
23666. https://doi.org/10.1038/srep23666
Xu, X., Li, G., Li, C., Zhang, J., Wang, Q., Simmons, D. K., Chen, X., Wijesena, N., Zhu,
W., Wang, Z., Wang, Z., Ju, B., Ci, W., Lu, X., Yu, D., Wang, Q., Aluru, N.,
Oliveri, P., Zhang, Y. E., … Liu, J. (2019). Evolutionary transition between
invertebrates and vertebrates via methylation reprogramming in embryogenesis.
National Science Review, *6*(5), 993–1003. https://doi.org/10.1093/nsr/nwz064
Yagound, B., Remnant, E. J., Buchmann, G., & Oldroyd, B. P. (2020). Intergenerational
transfer of DNA methylation marks in the honey bee. *Proceedings of the National*
Academy of Sciences, *117*(51), 32519–32527.
https://doi.org/10.1073/pnas.2017094117
Yan, H., Bonasio, R., Simola, D. F., Liebig, J., Berger, S. L., & Reinberg, D. (2015).
DNA Methylation in Social Insects: How Epigenetics Can Control Behavior and
Longevity. *Annual Review of Entomology*, *60*(1), 435–452.
https://doi.org/10.1146/annurev-ento-010814-020803
Yan, H., Simola, D. F., Bonasio, R., Liebig, J., Berger, S. L., & Reinberg, D. (2014).
Eusocial insects as emerging models for behavioural epigenetics. *Nature*
Reviews Genetics, *15*(10), 677–688. https://doi.org/10.1038/nrg3787
Yu, G., Wang, L.-G., Han, Y., & He, Q.-Y. (2012). clusterProfiler: An R Package for
Comparing Biological Themes Among Gene Clusters. *OMICS: A Journal of*
Integrative Biology, *16*(5), 284–287. https://doi.org/10.1089/omi.2011.0118
Yu, Q., Fang, S., Zhang, Z., & Jiggins, C. D. (2016). The transcriptome response of
Heliconius melpomene larvae to a novel host plant. *Molecular Ecology*, *25*(19),
4850–4865. https://doi.org/10.1111/mec.13826
Zemach, A., McDaniel, I. E., Silva, P., & Zilberman, D. (2010). Genome-Wide
Evolutionary Analysis of Eukaryotic DNA Methylation. *Science*, *328*(5980), 916–
919. https://doi.org/10.1126/science.1186366

Zhang, J.-S., Meng, J.-Y., Yang, L., & Zhang, C.-Y. (2025). Identification and expression
1040   analysis of cathepsin B genes in *Myzus persicae* (Hemiptera: Aphididae) and
1041   their response to environmental stresses. *Bulletin of Entomological Research*, 1–
1042   11. https://doi.org/10.1017/S0007485325000136
1043   Zhao, C., Escalante, L. N., Chen, H., Benatti, T. R., Qu, J., Chellapilla, S., Waterhouse,
1044   R. M., Wheeler, D., Andersson, M. N., Bao, R., Batterton, M., Behura, S. K.,
1045   Blankenburg, K. P., Caragea, D., Carolan, J. C., Coyle, M., El-Bouhssini, M.,
1046   Francisco, L., Friedrich, M., … Richards, S. (2015). A Massive Expansion of
1047   Effector Genes Underlies Gall-Formation in the Wheat Pest Mayetiola destructor.
1048   *Current Biology*, *25*(5), 613–620. https://doi.org/10.1016/j.cub.2014.12.057
1049   Zhao, C., Rispe, C., & Nabity, P. D. (2019). Secretory RING finger proteins function as
1050   effectors in a grapevine galling insect. *BMC Genomics*, *20*(1), 923.
1051   https://doi.org/10.1186/s12864-019-6313-x
1052
1053

Table 1: Fisher's exact test for functional group enrichment among host-responsive (HR) genes

Annotation Group	BH-adjusted p-value	Odds ratio	# HR genes	Total # genes
General metabolism	8.9E-03	5.23	6	14
Developmental genes	1.6E-14	4.55	53	135
Predicted effectors	1.7E-127	3.30	804	2,743
Cuticle and chitin related proteins	1.5E-06	3.03	35	116
Other	1.4E-33	2.41	319	1,280
Circadian rhythm	not sig.	2.32	4	16
Detoxification of xenobiotics	1.9E-05	2.14	52	222
Epigenetic machinery	not sig.	1.74	1	5
Odorant or gustatory perception	1.9E-05	1.05	14	107
Small RNA machinery	not sig.	0.63	1	12
null	2.5E-74	0.49	2,417	20,998

1054

1055

Figure 1: (a) Overview of the initial experimental setup; genotype names in red were resistant to *D. vitifoliae,* insects
died and were excluded from downstream analysis. (b-c) *D. vitifoliae* adult (black arrows) within its gall at 3 weeks after
infestation of host genotype 99R. (d-f) Young leaves of resistant host plants at 6 days after infestation; white arrows
indicate sites of leaf necrosis induced by *D. vitifoliae* feeding/probing; horizontal black scale bars are 1 mm in length.
(g) Box plot showing interquartile range and median (horizontal line within box) of *D. vitifoliae* fitness (eggs per gall)
across susceptible host genotypes; genotypes with letters 'a' are significantly different from those with letter 'b' based
on pair-wise Wilcoxon tests (Benjamini-Hochberg [BH]-adjusted p-values are all less than 0.0001). (h) Multi-
dimensional scaling plot of log-CPM (gene counts per million gene counts) for all RNA-seq samples.

Figure 2: Transcriptional plasticity induced by host transfer. (a-b) Venn diagrams showing the number of differentially expressed *D. vitifoliae* genes contrasting *V. vinifera* cv Cab33 (a) or SO4 (b) with all other hosts genotypes. (c-d) KEGG terms enriched among genes upregulated (c) and downregulated (d) during interaction with *V. vinifera* cv Cab33 compared to all other hosts; colors represent Benjamini-Hochberg (BH) adjusted p-values.

Figure 3: Gene co-expression network analysis. (a) Gene dendrogram for *D. vitifoliae* signed (unidirectional) gene co-expression network and module assignment (bottom). (b) Heatmap of module eigengenes (first principal compenent of the module, the 'most representative' gene) for all 88 modules (verticle axis) clustered based on Euclidean distance. (c-f) Scatter plots and results of linear regression tests for four co-expression modules that correlate in expression with *D. vitifoliae* fitness.

1087
Figure 4: Spatial organization and gene expression profile of ARGL genes and multiple sequence alignment of ARGL
proteins. (a) Genome browser view showing the ARGL locus in the *D. vitifoliae* genome; ARGL genes assigned to the
thistle gene co-expression module are highlighted with bold red text. (b) Heatmap of normalized gene expression for
genes in the thistle module; black rectangles highlight the ARGL genes. (c) Sequence logo plot based on multiple

sequence alignment (MSA) of the seven predicted effector proteins encoded at the ARGL locus and the top seventeen
orthologs of ARGL protein DV3003875; letters represent amino acid residues and letter size denotes the degree of
conservation (larger letters are more conserved).
1095
1096
1097
1098

1099
Figure 5: DNA methylation in *D. vitifoliae*. (a) Box plot showing the proportion of methylated cytosines (vertical axis) for
each sequence context (CpG, CHG, and CHH). (b) Bar plot showing CpG methylation levels for each genomic region
(untranslated regions (UTRs), intron, exon, and intergenic); the vertical axis is the total number of methylated CpG sites
divided by the total length (in basepairs) of the genomic region. (c) Principal component analysis (PCA) based on

methylation levels at 201,448 CpG sites significantly methylated in at least one sample and covered by 4 or more reads
in all samples. (d) Width-scaled violin plots showing CpG methylation levels (vertical axis) per transcriptional plasticity
quartile; genes that were expressed and had CpG methylation data (n = 12,368) were binned into quartiles based on
mean absolute value of log2-fold-change in expression (a quantification of transcriptional plasticity induced during
transfer to different hosts); letters above the quartiles indicate significant difference based on pair-wise Wilcoxon tests
(all p-values < 2e-16).
1110
1111
1112

1113
1114

1115
1116 **Figure S1**: The underside of a 3309 (*V. riparia* x *V. rupestris*) leaf 3 weeks after infestation with *D. vitifoliae*. Small black necrotic
1117 lesions formed on the leaves where *D. vitifoliae* attempted to feed.

1118
1119 **Figure S2**: KEGG term enrichment analysis for genes in the darkorange2 co-expression module.

1126

1127
1128 **Figure S3**: A heatmap of normalized gene expression for genes in the darkorange2 module. Black rectangles highlight genes with
1129 glycoside hydrolase KEGG annotation.
1130

1131
1132 **Figure S4**: Sequence logo plot based on multiple sequence alignment (MSA)
1133   of the seven predicted effector proteins at the ARGL locus.

ARGL cluster

1134
1135 **Figure S5**: Phylogenetic tree of all 2,743 predicted *D. vitifoliae* effectors based on multiple sequence alignment (MSA) of coding
1136 sequences (CDS). ARGL coding sequences are highlighted in red with black arrows.
1137
1138
1139

Tree scale: 0.1

ARGL proteins

1140

1141 **Figure S6**: Phylogenetic tree of all 2,743 predicted *D. vitifoliae* effector proteins based on multiple sequence alignment (MSA). ARGL

1142 proteins are highlighted in red with black arrows.

1143

1144
1145 **Figure S7**: The most conserved region of ARGL proteins. A multiple sequence alignment (MSA) of ARGL proteins was performed
1146 using all seven *D. vitifoliae* predicted effector proteins at the ARGL locus and the top seventeen orthologs of ARGL protein
1147 DV3003875. Amino acid residues in the 259 to 417 positions are shown, with gene ID (*D. vitifoliae* genes) or accession number on
1148 the left.

1149

1150 **Figure S8**: Effect of host genotype on *D. vitifoliae* CpG methylation. Principal component analysis (PCA) based on methylation data

1151 for 201,448 CpG sites for all samples minus outliers (left) and only egg samples minus outliers (right).

1152
1153 **Figure S9**: Correlation between transcriptional plasticity and gene body CpG methylation. The scatter plot shows transcriptional
1154 plasticity (mean absolute value of log2-fold-change; vertical axis) and percent CpG methylation (horizontal axis) for genes which are
1155 expressed and have CpG methylation data (n=12,368). Results of Spearman's rank correlation test are printed in the top right of the
1156 plot. A Locally Estimated Scatterplot Smoothing (LOESS) line was fitted to the data (blue line).
1157