Title

Transfer to different host grapevines causes rapid and widespread transcriptional

plasticity in the specialist insect *Daktulosphaira vitifoliae*

Author information

W. Nate Collison [1,2], Syed Adeel Zafar [1], Simon C. Groen [1,2], and Paul D. Nabity [1,3]

[1] Department of Botany and Plant Sciences, University of California, Riverside,

Riverside, CA, USA

[2] Department of Nematology, University of California, Riverside, Riverside, CA, USA

[3] School of BioSciences, University of Melbourne, Melbourne, Australia

Corresponding Author: Paul D. Nabity, School of BioSciences, University of Melbourne,

Melbourne, Australia, Paul.Nabity@unimelb.edu.au

Abstract

Insect herbivores employ many adaptive mechanisms to successfully colonize host plants. Transcriptional plasticity is one important mechanism that enables host use in polyphagous insects that maintain a broad host range. However, whether specialist herbivores also rely on transcriptional plasticity to feed successfully on different hosts remains largely unexplored. Here, we measure genome-wide transcriptional plasticity of grape phylloxera, *Daktulosphaira vitifoliae*, a grapevine (*Vitis* spp.) specialist, after transfer onto different host grapevines for a single generation using RNA-seq. We show that the transcriptome of *D. vitifoliae* is highly plastic and responsive to host genotype – 4,820 genes (37% of the 12,876 expressed genes) are significantly differentially expressed in at least one contrast comparing host genotypes. Genes with functions related to host manipulation (predicted secretory effectors), cuticle and chitin, detoxification of xenobiotics, and development were enriched among host-responsive genes, with predicted secretory effectors being most strongly enriched (Fisher's exact test odds ratio = 8.3, BH-adjusted p-value = 0). We further linked gene co-expression modules to *D. vitifoliae* fitness using weighted gene co-expression network analysis. The co-expression module most strongly correlated with fitness was enriched for predicted secretory effectors, including a novel group of proline-rich genes that exist as a tandem gene cluster. The only module negatively correlated with fitness was enriched for genes encoding glycoside hydrolase enzymes that function in carbohydrate metabolism. Overall, our results show that the phylloxerid transcriptome is highly plastic

and rapidly reprogrammed when transferred onto distinct hosts, which may be an

adaptive mechanism underlying host use in this clonally reproducing herbivore.

Key Words

grape phylloxera (*Daktulosphaira vitifoliae*), grapevines (*Vitis* spp.), ecological

specialization, host use, transcriptional plasticity, effectors

Introduction

As sessile organisms, plants face significant pressure to evolve effective

defenses against a breadth of herbivores. Plant defenses may occur at the

morphological level, such as trichomes and thorns, the chemical level, e.g. biosynthesis

of toxic defensive metabolites, or an indirect level, e.g. attraction of natural enemies that

target herbivores. The evolution of plant defense mechanisms imparts selective

pressure on herbivores to reciprocally evolve mechanisms to modulate or evade plant

defense. Two general strategies represent key adaptations to dampen plant defense –

1) tolerance or avoidance of defense chemicals via detoxification or sequestration, and

(2) direct manipulation of host defenses via delivery of effector molecules into or onto

host cells or tissues. In insect herbivores, plant defensive metabolites are often

detoxified in two phases [1]. In Phase I, plant-derived toxins are modified into less

reactive forms via enzymes such as cytochrome P450 monooxygenases. In Phase II,

the toxin derivatives are converted into hydrophilic forms to facilitate sequestration or

excretion via UDP-glycosyltransferases or other conjugating enzymes. The use of

secretory effector proteins that function as host-manipulating 'molecular weapons' has

been well-studied for microbial plant pathogens [2]. More recently, research on molecular

plant-insect interactions has revealed that insects also use secretory proteins as host-

modifying weapons. This is particularly true for herbivores that maintain intimate

associations with host plants, such as sap-feeding and gall-forming insects [3,4]. Although

experiments that functionally validate insect effector proteins are limited, these proteins

appear to play broad roles in modulating host biological processes including E3-

complex-/ubiquitin-mediated protein turnover, calcium signaling, and phytohormone-

mediated signaling [3–5].

Insect herbivores are among the most species-rich lineages of metazoans, and

this diversity is likely driven by insect adaptations to specific host plants [6]. Host-specific

adaptations of insects are a mode of ecological specialization that may inflict trade-offs

in fitness when an herbivorous insect feeds on ancestral or alternative hosts [7]. This

hypothesis is supported by the fact that most insect herbivores are specialists, feeding

on one or very few species of closely related hosts. How generalist, polyphagous

herbivores maintain compatibility with a breadth of divergent hosts remains largely

unknown. One hypothesis is that generalists maintain a high degree of genomic

flexibility, which enables polyphagy [8]. Indeed, the genomes of generalist insect

herbivores contain extraordinary expansions of genes that function in detoxification of

plant defensive metabolites [9,10,]. Additionally, transcriptional plasticity of expanded gene

families, including ones encoding detoxification proteins and secretory effectors, has

been linked to fitness in generalist insect herbivores during interaction with divergent

host plants [11,12]. Following this same logic, specialist herbivores should possess

genomes that are specialized for specific hosts and thus under less pressure to expand

and acquire new functions relative to generalists. Transcriptional plasticity of genes that

function in host plant use may also be limited or constrained in specialist insect

herbivores, although few studies have tested this hypothesis. Here, we investigate the

transcriptional basis of host use in grape phylloxera *Daktulosphaira vitifoliae* as it feeds

on different host grapevines for a single generation. We generated genome-wide gene

expression and fitness data with the aim of (1) understanding how the oligophagous

specialist *D. vitifoliae* successfully colonizes the non-native host *Vitis vinifera* (European

wine grape), (2) determining which genes show transcriptional plasticity upon transfer to

different host grapevines, and (3) correlating transcriptional plasticity with *D. vitifoliae*

fitness to determine whether plasticity in expression of different genes is adaptive,

neutral, or maladaptive.

Results

D. vitifoliae fitness differs significantly across host grapevine genotypes

To determine the fitness of *D. vitifoliae* on different grapevine hosts, we

experimentally introduced a clonal line of *D. vitifoliae* to a panel of grapevine genotypes,

which included the non-native host *V. vinifera* (cv Cab 33) and interspecific hybrids of

native North American grapevine species (Figure 1A). We collected fitness data

repeatedly over an 8-day period starting 15 days after *D. vitifoliae* introduction, when 1st

nymphal instars had matured to adulthood and begun reproducing. We found that *D.*

vitifoliae fitness varied significantly across host grapevine genotypes. Genotypes SW

(*V. riparia* X *V. rupestris*), 5BB (*V. riparia* X *V. berlandieri*), and 3309 (*V. riparia* X *V.*

rupestris) were incompatible with the *D. vitifoliae* genotype used in our study; *D.*

vitifoliae failed to induce galls and subsequently died. Attempted feeding by *D. vitifoliae*

induced small necrotic lesions on leaves of genotypes SW and 5BB within 6 days after

experimental infestation (Figure 1D and 1E). Interestingly, genotype 3309 showed

symptomless resistance after 6 days (Figure 1F), but eventually showed necrotic

lesions in leaves where *D. vitifoliae* had attempted to feed within 3 weeks after

infestation (Supplemental Figure 1). *D. vitifoliae* was compatible with the remaining five

genotypes including the non-native host *V. vinifera* cv Cab 33. To determine whether *D.*

vitifoliae fitness was significantly different between any of the five susceptible host

genotypes, we performed pair-wise Wilcoxon tests. *D. vitifoliae* fitness was significantly

lower on both *V. vinifera* Cab33 and SO4 (*V. berlandieri x V. riparia*) compared to the

other three hosts (Figure 1G). All other pairwise comparisons between host genotypes

revealed no significant differences in *D. vitifoliae* fitness.

D. vitifoliae gene expression is highly dynamic and responsive to host grapevine

genotype

To understand the genome-wide variation in *D. vitifoliae* gene expression

induced upon transfer onto different host plants, we isolated RNA from *D. vitifoliae*

adults after feeding on new hosts for a single generation and performed short-read

RNA-seq. Four replicates for each *D. vitifoliae* treatment group were sequenced,

totaling 24 samples. These included eight replicates for insects on host *V. vinifera* cv

Cab33 across two different *D. vitifoliae* generations. We targeted 30 million 150-bp

reads per sample (29 +/- 3 million) and 85% of reads uniquely mapped to gene models

in the reference genome after basic quality and adapter trimming (Supplemental Table

1). Gene count matrices were transformed and normalized using a standard pipeline in

edgeR [13], then unsupervised clustering of samples was plotted using multidimensional

scaling (MDS) (Figure 1H). Sample clustering indicated host environment induces

distinct changes in *D. vitifoliae* gene expression; however, the SO4 genotype group

showed high variability relative to the other 4 treatment groups.

We then performed differential gene expression analysis to identify genes that

vary in expression across different host genotypes. We performed pairwise contrasts for

each genotype-by-genotype comparison and concatenated the results to compile a list

of genes differentially expressed in one or more contrasts, i.e. host-responsive genes.

Genes were considered differentially expressed based on a 1.5-fold-change difference

in expression and adjusted p-value cutoff of 0.05. This resulted in 4,820 host-responsive

genes (Figure 2A), which is 37% of the 12,876 expressed genes in our dataset. We

then performed a Fisher's exact test to identify gene functional groups that are enriched

among the host-responsive genes relative to the genome-wide background, which

included all 25,808 *D. vitifoliae* gene models. Genes encoding predicted secretory

effectors (adj. p-value = 0, odds ratio = 3.8), cuticle- and chitin-associated proteins (adj.

p-value = 2.2e-47, odds ratio = 17.9), developmental regulators (adj. p-value = 2.2e-12,

odds ratio = 3.8), and genes with predicted functions in detoxification of plant

metabolites and other environmental molecules (adj. p-value = 2.1e-10, odds ratio =

2.7) were enriched among host-responsive genes (Figure 2A).

Pair-wise analysis of gene differential expression (DE) revealed that host *V.*

vinifera cv Cab33 induced the strongest change in gene expression compared to other

hosts; 4,817 out of the total 4,820 host-responsive genes show DE in pairwise contrasts

between *V. vinifera* cv Cab33 and the other four host genotypes (Figure 2B).

Furthermore, 227 genes show DE in all contrasts with *V. vinifera* cv Cab33 (Figure 2B).

Genes encoding cuticle and chitin associated proteins are significantly enriched among

the 227 overlapping DE genes (adj. p-value = 4.1e-38, odds ratio = 50.1) (Supplemental

Figure 2A). Enrichment analysis for Kyoto Encyclopedia of Genes and Genomes

(KEGG) terms showed that genes encoding oxidoreductase enzymes are enriched

among the 227 overlapping DE genes (Supplemental Figure 2B). We also investigated

the types of genes that are up- and down-regulated when *D. vitifoliae* feeds on *V.*

vinifera cv Cab33. Interestingly 3,933 (81.6 %) of the host-responsive genes are

upregulated during interaction with *V. vinifera* cv Cab33. Genes encoding predicted

secretory effectors, cuticle and chitin associated proteins, and detoxication enzymes are

enriched among genes upregulated on *V. vinifera* cv Cab33 (Supplemental Figure 3).

Enriched KEGG terms for upregulated genes include peroxidase and oxidoreductase

enzymes, sugar and organic acid transporters, integral membrane ion channels, and

fatty acid elongation enzymes (Supplemental Figure 4). Genes downregulated during

interaction with *V. vinifera* cv Cab33 are enriched for functions related to development

(Supplemental Figure 3) and carbohydrate metabolism (Supplemental Figure 4).

Overall, these results suggest that *D. vitifoliae* transcript expression is highly plastic,

especially at genes with predicted functions in host manipulation (secretory effectors)

and interaction with the environment.

<u>Four gene co-expression modules co-vary with *D. vitifoliae* fitness</u>

To identify groups of genes that are strongly correlated in expression pattern and

likely function together in shared biological processes, we performed weighted gene co-

expression network analysis using WGCNA [14]. We included 13,917 genes for network

construction after removing 'zero genes' that showed negligible expression (only genes

with ≥0.5 counts per million gene counts (CPM) in at least 4 samples were kept). We

constructed a signed co-expression network with 88 modules based on hierarchical

clustering of gene connectivity dissimilarity (Figure 3A-B). To identify groups of co-

regulated genes (modules) that might underlie differences in *D. vitifoliae* fitness across

host genotypes, we tested for correlation between module eigengenes (first principal

component of the module and the most representative gene) and fitness. Because our

D. vitifoliae RNA-seq samples were collected from a single host genotype we ran linear

regression tests for averaged insect fitness and averaged eigengenes for each host

genotype and co-expression module. Five eigengenes significantly correlated with *D.*

vitifoliae fitness (Figure 3C-F); since the 'grey' module is a bin for genes that cannot be

assigned to a co-expression module and are unlikely to be co-regulated, we included

only the 'thistle', 'darkolivegreen4', 'darkorange2', and 'antiquewhite4' modules for

further analysis. The darkolivegreen4 module was positively correlated with *D. vitifoliae*

fitness (R^2 = 0.91, p-value = 0.012) (Figure 3D). Darkolivegreen4 is moderately

overrepresented for genes related to detoxification of plant metabolites and other

environmental molecules (adjusted p-value = 0.084) (Supplemental Table 3). The

antiquewhite4 module was also positively correlated with *D. vitifoliae* fitness (R^2 = 0.79,

p-value = 0.044) (Figure 3F). A Fisher's exact test for gene annotation group enrichment

and KEGG term enrichment analysis showed no enrichment for the antiquewhite4

module. However, antiquewhite4 showed an interesting pattern of gene expression

across treatments (Supplemental Figure 5). Genes within the module clustered into two

groups with distinct expression patterns based on hierarchical clustering of vst-

normalized expression. A small cluster of nine genes increased in expression during

interaction with *V. vinifera* cv Cab33 relative to native hosts 101-14, 1103, and 99R,

whereas a larger cluster of 57 genes decreased in expression during interaction with *V.*

vinifera cv Cab33 relative to native hosts. Thus, the antiquewhite4 module appears to

be downregulated in *D. vitifoliae* during interaction with *V. vinifera* relative to native

hosts, which may contribute to the successful colonization of this non-native host.

The darkorgange2 module is enriched for genes that are host-responsive and encode

glycoside hydrolase family 1 proteins

We performed Fisher's exact tests to identify gene co-expression modules that

were enriched for host-responsive genes. 21 out of the 88 modules were enriched for

host-responsive genes (odd ratio >2, BH-adjusted p-value <0.05) (Supplemental Table

1). The darkorange2 module was the only module that was both enriched for host-

responsive genes and significantly co-varying with *D. vitifoliae* fitness. Darkorange2 is

also the only module that was negatively correlated with *D. vitifoliae* fitness (Figure 3C-

F). Based on KEGG term enrichment analysis, darkorange2 is over-represented for

genes that encode glycoside hydrolase family 1 enzymes (PFAM GH1), which mediate

carbohydrate metabolism (Supplemental Figure 6). Similar to the antiquewhite4 module,

genes within the darkorange2 module show distinct patterns of expression across host

genotypes (Figure 6). A small cluster of 24 genes is highly expressed during interaction

with native hosts 101-14 and 1103 and decreases in expression during interaction with

V. vinifera cv Cab33 and SO4, which are hosts that maintain relatively low *D. vitifoliae*

fitness. 4 of the 5 genes with glycoside hydrolase KEGG annotation are within this 24-

gene cluster and show similar patterns of expression (Figure 6). The remaining 66

genes in the module are grouped into a larger cluster that shows the exact opposite

trend in expression, largely increasing in expression during interaction with *V. vinifera* cv

Cab33 and SO4 relative to 101-14 and 1103. These results show the darkorange2

module is highly plastic and variation in module expression may explain differences in

D. vitifoliae fitness across distinct host genotypes.

The thistle module is enriched with predicted effectors including a novel group of

proline-rich secretory proteins

The thistle module was most strongly correlated with *D. vitifoliae* fitness (R^2 =

0.94, p-value = 0.005) and) and was positively correlated with fitness (Figure 3C). The

thistle module is also enriched with genes encoding predicted secretory effectors based

on Fisher's exact test (Supplemental Table 3), including a group of four effectors with

homology to the *Acyrthosiphon pisum* lysine-rich arabinogalactan protein 19-like protein

based on BLAST searches

(https://bipaa.genouest.org/sp/daktulosphaira_vitifoliae/download/annotation/v3.2/func_

annot/)[15]; we termed these genes 'ARGL' effectors, for **AR**abino**G**alactan-**L**ike. We also

noticed that the ARGL genes are in a tandem cluster in the genome on scaffold 40

(Supplemental Figure 7). Within this same ~50-kb spatial cluster are two more predicted

effectors and three 'null' genes; two of the three null genes are also assigned to the

thistle module, suggesting they are co-regulated with the ARGL genes. We performed

new BLASTp searches for ARGL effectors and found that they show homology to one

another and to other proteins from *D. vitifoliae*, *Adelges cooleyi*, and multiple aphid

species (Supplemental Table 4). The ARGL effectors and all proteins that are significant

BLAST hits contain no predicted functional domains based on InterProScan searches.

To determine if the ARGL effectors have highly similar sequences and shared

protein motifs, we performed Multiple Sequence Alignment (MSA) of the seven ARGL

proteins using clustalOmega[16]. The sequence logo plot for the ARGL protein MSA

revealed these proteins have a high proportion of proline residues and are highly

conserved, especially at proline residues (Figure 5A). To understand the evolutionary

relationship among ARGL effectors, we performed MSA for all 2,743 predicted effector

proteins and constructed a phylogenetic tree. Despite being highly conserved at the

amino acid level, AGRL effectors did not cluster together in a monophyletic group

(Supplemental Figure 8), suggesting ARGL proteins have some structural divergence.

The ARGL effectors show similar patterns of transcript expression across treatments,

generally increasing in expression during interaction with high-fitness native hosts 101-

14, 1103, and 99R relative to *V. vinifera* cv Cab33. Overall, our analysis of the thistle

co-expression module suggests ARGL genes may have shared function and play a role

in *D. vitifoliae* host use, albeit through unknown mechanisms.

Discussion

We have presented analyses of a large RNA-seq data set collected from *D.*

vitifoliae adults after feeding on different host grapevines (*Vitis* spp.) for a single

generation, with the aim of understanding the transcriptional basis of host use in this

specialist herbivore. We found that 37% of all expressed genes are host-responsive,

with most genes being upregulated during interaction with the non-native host *V.*

vinifera cv Cab33 relative to native hosts. Genes encoding predicted secretory effectors,

cuticle and chitin associated proteins, and detoxification-related proteins were enriched

among host-responsive genes, suggesting that genes that function in host-manipulation,

detoxification of xenobiotics, and protection against the environment are dynamically

regulated for successful colonization of the non-native host *V. vinifera*. Weighted gene

co-expression network analysis revealed modules co-regulated genes that covary with

D. vitifoliae fitness. Most notably, the co-expression module 'thistle' was strongly

positively correlated with *D. vitifoliae* fitness and enriched with genes encoding

predicted secretory effectors. Five of these genes were co-located in a ~25-kb spatial

cluster in the genome and encode highly conserved and proline-rich secretory proteins,

termed ARGL proteins. These proteins have no known functional domain, thus ARGL

gene function remains unclear. Our findings show that the *D. vitifoliae* transcriptome is

highly plastic and is reprogrammed within a single generation on different host

grapevines. Our analysis of genome-wide gene expression and fitness for *D. vitifoliae*

on different hosts also indicates transcriptional plasticity for a subset of genes may be

adaptive and enable successful colonization of the non-native wine grape *V. vinifera*.

Materials and Methods

<u>Propagation of host grapevines</u>

Nine different grapevine genotypes were used as experimental hosts for *D.*

vitifoliae in our experiments, all of which were clonally propagated from cuttings

purchased as rooted, overwintered 1-year-old plants from Inland Desert Nursery

(Benton City, WA). Genotypes included self-rooted rootstock hybrids of *V. berlandieri* x

V. rupestris), (99R, 1103P), *V. berlandieri* x *V. riparia* (SO4, 5BB), and *V. riparia* x *V.*

rupestris (101-14, SW, 3309). All plants were potted in 3-gallon round pots with 'UC Mix

3' soil (https://agops.ucr.edu/soil-mixing#uc-soil-mix-iii) and grown in a greenhouse

(24:27C +/- 3C for day:night with 14h daylight). After leaf bud swelling (8 days after

potting), half of each genotype were infested with the remaining plants potted and

infested similarly 6 days later. Plantings were replicated through time to optimize

sample collection effort given the time required to sort and count insects.

<u>Phylloxera rearing and experimental infestation of host grapevines</u>

A clonal line of leaf-galling *D. vitifoliae* was established from a single female

collected on *V. rupestris* (cv St. George) near Davis, CA, in 2021. The clonal population

was maintained for successive generations on both grafted (*Vitis* genotype 1616 and

3309 rootstocks) and self-rooted *V. vinifera* cv Cabernet Sauvignon 33 (Cab33) in UCR

greenhouses in Riverside, CA (27+/-3C) until experimentation over 2022-2023. For

infestation, galled leaves from the starting *D. vitifoliae* colony (hereafter generation 0, or

'G0') were excised and placed on the branch nodes of new host plants. Infestation was

monitored every 12 hours to note hatch and dispersal of 1^{st} nymphal instars (hereafter

generation 1, 'G1') onto budding leaves. Galled leaves used for infestation were

removed from branch nodes after two days to prevent further infestation.

<u>Insect performance assay and statistical analysis</u>

To assess plant vigor and insect performance, 'G1' galls were collected 15 days

after infestation (DAI). Galls and tissue from ungalled leaves of similar age and node on

un-infested plants were collected using a 4mm diameter hole punch, placed in tubes on

ice for transport back to the lab. Galls were dissected under a dissecting microscope to

count eggs per gall. Dissected gall tissues and 10 leaf discs per plant were also

weighed for fresh weight, dried at 50C until dry (1-2 days), and then assessed for dry

weight. Each plant was sampled five different times, at 15, 17, 19, 21, and 23 DAI. Each

time the five largest galls from each plant were sampled, selecting galls from leaves that

had not been previously sampled unless the largest remaining galls were on a

previously sampled leaf. Egg production was assessed using a mixed effects linear

model with genotype and day as predictors and plant id as a random effect. Post hoc

comparisons among genotypes were performed using pairwise Wilcoxon tests (R v.

4.3.1).

RNA isolation and RNA sequencing

To quantify *D. vitifoliae* transcriptome plasticity during host switching events,

twenty adults were collected from galls from unique plants and pooled per each sample.

Insects were placed in tubes on ice during collection then immediately stored at -80 °C

until RNA isolation. RNA was isolated from adults using the RNeasy Mini Kit (Qiagen)

following manufacturer's instructions. We ran gel electrophoresis to confirm RNA

intactness and only kept samples that had intact 28S and 18S ribosomal RNA bands.

We used a Nanodrop spectrophotometer to measure RNA purity and only used samples

with 260 nm / 280 nm absorbance ratios above 2 and 260 nm / 230 nm absorbance

ratios above 1.7 for library preparation. Libraries were prepared using a

NEBNext® Ultra™ II Directional RNA Library Prep Kit for Illumina® (New England

Biolabs). Paired-end RNA libraries were sequenced using an Illumina NovaSeq X

system at the DNA Technologies and Expression Analysis Core of the UC Davis

Genome Center, Davis, CA.

RNA-seq data preprocessing

Fastqc v. 0.11.9 (https://github.com/s-andrews/FastQC) was used to generate

read and base quality statistics for raw and processed reads. Raw RNA-seq reads were

quality trimmed and filtered using BBTools bbduk (v.38.95; https://jgi.doe.gov/data-and-

tools/software-tools/bbtools/) with default settings plus options for kmer length (k=23)

and hamming distance (hist=1). Trimmed and filtered reads were aligned to the

reference genome [15] using STAR v. 2.7.10b [17] with default settings plus options for

maximum intron length (--alignIntronMax 15000) and removing alignments that

contained non-canonical junctions (--outFilterIntronMotifs RemoveNoncanonical). Only

uniquely mapped reads were used for downstream analysis.

<u>Differential gene expression analysis</u>

Gene count matrices were generated using the --quantMode GeneCounts option

in STAR, which contained gene counts that coincide with counts produced by htseq-

count with default parameters. Gene count matrices were processed and analyzed

using edgeR (v.4.0.16) [13]. Genes with low counts were removed from downstream

analysis using the filterByExpr command with default parameters. Linear model fitting of

count data was performed using limma's voom and lmFit functions [18]. Genes were

considered differentially expressed at adjusted p-value < 0.05 and log fold-change ≥

1.5. Pie charts were made using the R graphics package (v.4.3.1) to visualize functional

annotations of genes that were differentially expressed. Fisher's exact testing for gene

functional group enrichment was performed with the R stats package (v.4.3.1).

Functional groups were considered significantly enriched based on a BH-adjusted p-

value cutoff of 0.05 and an odds ratio ≥2.

Gene co-expression network analysis

Genes with 0.5 CPM (about 10 raw gene counts) in at least four samples were

included for network construction using WGCNA [14]. Gene count matrices were

normalized using DESeq2 vst (variance stabilizing transformation) function (v. 1.42.1). A

signed gene co-expression network was constructed based on hierarchical clustering of

dissimilarity in gene connectivity (1-TOM) with the soft power parameter (β) set to 16 to

ensure the network had approximate scale-free topology. Genes were assigned to

modules using cutreeDynamic function with minimum module size = 20 and module

detection sensitivity (deepSplit) = 2. Heat maps of module eigengenes and normalized

gene expression (z-scores of vst-transformed count data) were made using pheatmap

(v. 1.0.12) (https://github.com/raivokolde/pheatmap); rows were clustered based on

Euclidean distance. Correlation between *D. vitifoliae* fitness and gene co-expression

modules were tested by running a linear regression test for average fitness on each

host genotype and average module eigengene expression using the lm function in the R

stats package v. 4.3.1: fitness = $\beta_0 + \beta_1 \times$ eigengene + ϵ. Scatter plots of eigengene-

fitness correlations were printed with results from the linear regression test using

ggplot2 v, 3.5.0 (https://ggplot2.tidyverse.org/). Gene co-expression modules were

tested for overrepresentation/enrichment of gene functional groups based on Fisher's

exact tests as described above.

KEGG term enrichment analysis

eggNOG-mapper v. 2 [20] was used to assign KEGG terms to the predicted

proteome of *D. vitifoliae* [15] based on precomputed orthology assignments. eggNOG-

mapper assigned KEGG terms to 7,532 proteins out of 15,076 queried. To test for

KEGG term enrichment for subsets of genes, e.g. differentially expressed genes vs all

KEGG-annotated genes, we used the enrich function of the R package clusterProfiler v.

4.10.1 [21]. KEGG terms were considered significantly enriched based a BH-adjusted p-

value cutoff <0.05 and a gene count $\geq$4.

Multiple Sequence Alignment and phylogenetic analysis

To determine sequence similarity and phylogenetic relationships of ARGL

proteins, we performed Multiple Sequence Alignment (MSA) using the clustalOmega

web application (https://www.ebi.ac.uk/jdispatcher/msa/clustalo) [16]. Alignment results

were visualized in a sequence logo plot using the WebLogo web application

(https://weblogo.threeplusone.com) [22]. Phylogenetic trees were constructed from the

clustalOmega MSA output using the Interactive Tree of Life (iTOL) v. 6 web application

(https://itol.embl.de) [23].

Acknowledgments

We thank Conner Lay for help with collection of phenotypic data, Hung D. Lai for

greenhouse maintenance, and Holly Clark, Clay Clark, and Wei Zhang at the UC

Riverside genomics core for RNA library preparation. USDA-NIFA Hatch grant CA-R-

BPS-5199-H and CDFA grant CDFA-IAB #21-0426-000-SA were awarded to PDN to

support this project.

References

1. Heidel-Fischer, H. M. & Vogel, H. Molecular mechanisms of insect adaptation to plant

secondary compounds. *Current Opinion in Insect Science* **8**, 8–14 (2015).

2. Jones, J. D. G. & Dangl, J. L. The plant immune system. *Nature* **444**, 323–329

(2006).

3. Stuart, J. Insect effectors and gene-for-gene interactions with host plants. *Current*

Opinion in Insect Science **9**, 56–61 (2015).

4. Kaloshian, I. & Walling, L. L. Hemipteran and dipteran pests: Effectors and plant host

immune regulators. *J. Integr. Plant Biol.* **58**, 350–361 (2016).

5. Wang, H., Shi, S. & Hua, W. Advances of herbivore-secreted elicitors and effectors in

plant-insect interactions. *Front. Plant Sci.* **14**, 1176048 (2023).

6. Simon, J.-C. *et al.* Genomics of adaptation to host-plants in herbivorous insects.

Briefings in Functional Genomics **14**, 413–423 (2015).

7. García-Robledo, C. & Horvitz, C. C. Jack of all trades masters novel host plants:

positive genetic correlations in specialist and generalist insect herbivores expanding

their diets to novel hosts: Quantitative genetics of diet expansion. *Journal of*

Evolutionary Biology **25**, 38–53 (2012).

8. Birnbaum, S. S. L. & Abbot, P. Gene Expression and Diet Breadth in Plant-Feeding

Insects: Summarizing Trends. *Trends in Ecology & Evolution* **35**, 259–277 (2020).

9. Grbić, M. *et al.* The genome of Tetranychus urticae reveals herbivorous pest

adaptations. *Nature* **479**, 487–492 (2011).

10. Pearce, S. L. *et al.* Genomic innovations, transcriptional plasticity and gene loss

underlying the evolution and divergence of two highly polyphagous and invasive

Helicoverpa pest species. *BMC Biol* **15**, 63 (2017).

11. Snoeck, S., Wybouw, N., Van Leeuwen, T. & Dermauw, W. Transcriptomic

Plasticity in the Arthropod Generalist *Tetranychus urticae* Upon Long-Term

Acclimation to Different Host Plants. *G3 Genes|Genomes|Genetics* **8**, 3865–3879

(2018).

12. Mathers, T. C. *et al.* Rapid transcriptional plasticity of duplicated gene clusters

enables a clonally reproducing aphid to colonise diverse plant species. *Genome Biol*

18, 27 (2017).

13. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR : a Bioconductor

package for differential expression analysis of digital gene expression data.

Bioinformatics **26**, 139–140 (2010).

14. Langfelder, P. & Horvath, S. WGCNA: an R package for weighted correlation

network analysis. *BMC Bioinformatics* **9**, 559 (2008).

15. Rispe, C. *et al.* The genome sequence of the grape phylloxera provides insights

into the evolution, adaptation, and invasion routes of an iconic pest. *BMC Biol* **18**, 90

(2020).

16. Sievers, F. & Higgins, D. G. Clustal Omega for making accurate alignments of

many protein sequences. *Protein Science* **27**, 135–145 (2018).

17.   Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–

21 (2013).

18.   Law, C. W., Chen, Y., Shi, W. & Smyth, G. K. voom: precision weights unlock

linear model analysis tools for RNA-seq read counts. *Genome Biol* **15**, R29 (2014).

19.   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

20.   Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P. & Huerta-Cepas,

J. eggNOG-mapper v2: Functional Annotation, Orthology Assignments, and Domain

Prediction at the Metagenomic Scale. *Molecular Biology and Evolution* **38**, 5825–

5829 (2021).

21.   Yu, G., Wang, L.-G., Han, Y. & He, Q.-Y. clusterProfiler: an R Package for

Comparing Biological Themes Among Gene Clusters. *OMICS: A Journal of*

Integrative Biology **16**, 284–287 (2012).

22.   Crooks, G. E., Hon, G., Chandonia, J.-M. & Brenner, S. E. WebLogo: A

Sequence Logo Generator: Figure 1. *Genome Res.* **14**, 1188–1190 (2004).

23.   Letunic, I. & Bork, P. Interactive Tree of Life (iTOL) v6: recent updates to the

phylogenetic tree display and annotation tool. *Nucleic Acids Research* **52**, W78–W82

(2024).

Figure 1: (a) Overview of the initial experimental setup. (b-c) A phylloxera (*D. vitifoliae*) adult (black arrows) within its gall at 3 weeks after infestation of host genotype 99R. (d-f) Young leaves of resistant host plants at 6 days after infestation; white arrows indicate sites of leaf necrosis induced by *D. vitifoliae* feeding/probing; horizontal black scale bars are 1 mm in length. (g) Box plot of *D. vitifoliae* fitness (eggs per gall) across susceptible host genotypes; genotypes with letters 'a' are significantly different from those with letter 'b' based on pair-wise Wilcoxon tests (Benjamini-Hochberg [BH]-adjusted p-values are all less than 0.0001). (h) Multi-

488   dimensional scaling plot of log-CPM (gene counts per million gene counts) for all

489   RNA-seq samples.

Figure 2: (a) Pie chart of all host-repsonsive *D. vitifoliae* genes and assigned

functional groups; Fisher's exact test results are displayed for functional groups that

are signficantly enriched among host-responsive genes relative to all genes. (b-c)

Venn diagrams of differentially expressed *D. vitifoliae* genes contrasting *V. vinifera* cv

Cab33 (b) or SO4 (c) with all other hosts genotypes.

Figure 3: (a) Gene dendrogram for *D. vitifoliae* signed gene co-expression network

and module assignment. (b) Heatmap of module eigengenes for all 88 modules

clustered based on Euclidean distance. (c-f) Scatter plot and results of linear

regression tests for four co-expression modules that correlate in expression with *D.*

vitifoliae fitness.

Figure 4: Heatmap of normalized gene expression for genes in the darkorange2

module. Black rectangles highlight genes with glycoside hydrolase KEGG annotation.

Figure 5: Sequence logo plot for the seven predicted effector proteins at the ARGL

518   locus.

Figure 6: Heatmap of normalized gene expression for genes in the thistle module.

521 Black rectangles highlight ARGL effectors.