Skin hydration by natural moisturizing factors, a story of H-bond networking

Marving Martin,[*a,b] Benjamin Chantemargue,[b] Patrick Trouillas,[a, c]

a) INSERM U1248 Pharmacology & Transplantation, Univ. Limoges, CBRS, 2 rue du prof. Descottes, F-87000 Limoges, France
b) InSiliBio, 1 avenue d'Ester, Ester Technopôle, F-87000 Limoges, France
c) RCPTM CATRIN Palacky University, Slechtitelu 27, 783 71, Olomouc, Czech Republic.

* marving.martin@unilim.fr

Abstract

Dry skin is a common condition experienced by many. Besides being particularly present during the cold season, various diseases exist all year round, leading to localized xerosis. To prevent it, the skin is provided with natural moisturizing factors (NMFs). They are small amino acids or derivatives found in the outermost layer of the skin, the *stratum corneum* (SC). They are often claimed to be highly efficient humectants, increasing the water content to maintain the fluidity of the skin. However, alternative mechanisms have been proposed, suggesting that NMFs themselves may act as lipid mobility amplifiers.

This work aims at investigating the role of three NMFs, namely urea (URE), glycerol (GLY) and urocanic acid / urocanate (UCA/UCO) in SC *in silico* models, considering two different levels of humidity. Molecular dynamic simulations showed an increase in the diffusion of the different lipid components in the presence of either high water content, NMFs or both. The membrane properties were modified, as seen by an increase in lipid conformational changes as well as a decrease in lipid chain ordering. URE and GLY exhibited a similar impact, while UCA exhibited a higher impact on the liquid-like layer of SC. By studying NMF-water intermolecular interactions, we highlighted the role of NMF as a regulator of membrane perturbations, while insuring membrane fluidity. This role allows NMFs to prevent

destabilization of the skin membrane in the presence of high-water content. This study, performed at an atomistic resolution, highlighted a strong H-bond network between lipids, involving mainly ceramides but also all other ingredients. This network can be modified in the presence of high-water concentration or NMFs, resulting in modifications of membrane properties, rationalizing hydration effects.

Keywords: natural moisturizing factors; stratum corneum; molecular dynamics; membrane properties; H-bond network; skin hydration

Introduction

Every living organism is provided with a biological interface that maintains its integrity and protects it from the surrounding environmental aggressions. It is a key structure for the organism to evolve and adapt to the outside world threats (*e.g.,* strong climate variations, pollution, mechanic impacts or sun light) [1]. For humans, this interface is the skin, a complex multilayered and multiscale structure, composed of the epidermis, the dermis, and the hypodermis [2]. The skin epidermis is the outer layer, in other words, the first protective barrier of the body. A major structuration of this layer is the *stratum corneum* (SC). It is a brick-and-mortar architecture, where the bricks are corneocyte cells and the mortar is a multilamellar lipid stack. The SC is soft, relatively pliable and capable of tolerating physical constraints or varying osmotic conditions, as long as it stays hydrated and moisturized. The corneocytes are flattened and rigid dead keratinocyte cells, containing small hygroscopic molecules named natural moisturizing factors (NMFs) [3]. NMFs bind and retain water molecules from the ambient atmosphere into the corneocytes' lumen, in turn maintaining adequate skin moisturization, even at low humidity level. This allows enzymes to work under correct physiological conditions for biological processes such as desquamation. The mortar is an intercellular lipid matrix, surrounding and maintaining the corneocytes, which are stacked like in a brick wall. This lipid lattice is made of different ceramides (CER), free fatty acids (FFA), cholesterol (CHL), and water. All these ingredients are organized in a lamellar phase providing a tight, fluid, and effective barrier against transepidermal water loss (TEWL). To support the function of the SC, multiple topical creams have been developed using NMF, water, and skin lipids as the main ingredients. These formulations can be used in cosmetics but also to treat various skin pathologies including xerosis or atopic dermatitis [4,5]. However, it is still quite unclear how NMF molecules and skin lipids interact at the molecular level when these topical formulations are applied to the skin. Recent articles have shown the stabilizing

effect of small polar compounds, such as NMF, on the intercellular lipid fluidity under reduced hydration conditions [6–8]. Glycerol (GLY), urea (URE) and urocanic acid (UCA) demonstrated a water-like behavior, by increasing molecular mobility [9]. These findings pave the way toward a better understanding of the role of NMFs and their skin lipid interplay to maintain the SC fluidity. In this publication, we aim at joining the efforts of molecular modeling and the literature knowledge to rationalize the underlying molecular events of NMF and/or water within the SC lipid matrix. To this aim, we used molecular dynamics (MD) as a powerful tool to sample all molecular interactions and motions of a series of three NMFs embedded into an *in silico* model of the SC intercellular lipid matrix. The three NMFs studied here are URE, GLY and UCA. They were inserted in a SC model of reference and at two different hydration levels. Both UCA and its deprotonated urocanate (UCO) forms were considered, as the pKa of this acid-base couple is about 3. Although the pKa value can be modulated at the membrane interface and inside the membrane, at skin pH, the negatively charged UCO form is likely to be present, possibly predominant, in the skin.

Computational details

The all-atom skin model of reference (REF) was inspired by an early published article [10]. The REF model is designed to reproduce cryo-EM (electron microscopy) structures [11] of the intercellular lipid matrix obtained from near-native human skin. It is composed of fully extended (splayed) CER, FFA, CHL, and water molecules with the following distribution 90:88:88:90, respectively. Three types of ceramides were used in this system, with a physiological acyl chain length distribution [12], namely non-hydroxylated sphingosine (NS), non-hydroxylated phytosphingosine (NP), and esterified ω-hydroxylated sphingosine (EOS). The complete composition of the REF system is given in **Table 1**. According to the original publication, a specific 3D spatial arrangement of the lipids is required to accurately mimic the EM structures, the water permeability and the thermotropic behavior. Hence, 75% of CHL is located close to the CER sphingoid chains, and the remaining 25% CHL being located in close contact with the CER fatty acid chains. FFAs are only positioned close to the fatty acid moieties of CER. To construct the REF model, each molecule topology was built from scratch, using the MarvinSketch software[13]. The REF system, containing 90 water molecules, was manually assembled in the VMD software [14] to form the initial structure (**Figure 1.a-b**). The lipids and water were described by the CHARMM36 forcefield[15] and the TIP3P model, respectively. The adjusted polar head dihedral parameters were collected from the publication's supplementary data[10]. Energy minimization was performed using GROMACS 2021[16]. The minimized structure was then heated to 305K (in the NVT ensemble) for 1ns, followed by a semi-isotropic pressurization maintained at 1 bar for 1ns (in the NPT ensemble). The output structure displayed four different layers namely: short-chain, lipid headgroup, long-chain and liquid-like layers (**Figure 1.c**). The short-chain layer is made of 75% CHL and all CER sphingoid moieties. The long-chain layer contains CER fatty acid moieties, FFA and 25% of CHL. The liquid-like layer is a highly disordered region

containing CER EOS ester moieties. The different physical-chemical parameters of the membranes confirmed that the REF system behaves as expected. For example, the calculated overall thickness was found very similar to the value found in the original publication, at around 10 nm [10,11] (see **Figure 1.c**). The density along the z-axis confirmed the positioning of the different lipids and the presence of water molecules, mainly in the lipid headgroup regions and a smaller amount in the liquid-like layer (see **Figure 2.a**).

The production was run for 500 ns, with both electrostatic and Lennard-Jones cutoffs set at 12 Å, and the longer-range interactions were performed using the Particle-Mesh Ewald procedure. The timestep integration was set to 2 fs and the LINCS algorithm handled hydrogen bond restrain. The REF:WAT system, containing a doubled amount of water molecules, *i.e.,* 180, was created from a fully-relaxed REF model. Both REF and REF:WAT systems were used to create eight other systems containing, each, 72 NMF (URE, GLY, UCA or UCO) molecules (see **Figure 1.d**). Therefore, the following eight systems were studied: four REF:NMF systems (REF:GLY, REF:URE, REF:UCA and REF:UCO) and four doubled-hydrated systems, namely REF:GLY:WAT, REF:URE:WAT, REF:UCA:WAT and REF:UCO:WAT. All NMF geometries were optimized using the Gaussian09[17] program at the MP2/6-31G* level of theory, while the bonded and non-bonded parameters were collected from the CGenFF webserver. UCA and UCO were obtained in *trans*[18]. MD productions were run in triplicate, for each of the eight systems. Most of the analyses were carried out using AmberTools22 packages[19]. The electrostatic potential was assessed with the GROMACS analysis tools and the percentage of CER hairpin conformations (see **Figure 1.a**) was calculated with an in-house Python script.

Table 1. Detailed composition of the lipid matrix model of reference *i.e.*, REF system

compounds		Fatty acid chains length	Sphingoid chains length	Ester chains length	Number of Compound
CER	NP	C20:0	C18:0		4
		C22:0			8
		C24:0			16
		C26:0			24
		C28:0			8
		C30:0			2
	NS	C24:0			16
	EOS	C26:0		C18:2 (Δ9,12)	12
FFA		C20:0			4
		C22:0			8
		C24:0			30
		C26:0			24
		C28:0			8
		C30:0			14
CHL					88
WATER					90
TOTAL					**356**

Figure 1 (a.) REF system composition: CER NS and NP in green, CER EOS in purple, CHL in yellow, FFA in orange and water. (b.) Initially constructed REF structures using extended CER conformation. (c.) Equilibrated REF structure revealing the organization of the different layers along the z-axis. (d) NMF chemical structures.

Results

Figure 2 (a.) Trajectory snapshots of the different systems highlighting the position of water (blue surface) and NMF molecules in van der Waals spheres within the SC lipids (CERs in green, CER EOS in purple, FFA in orange and CHL in yellow represented as balls-and-sticks). The z-axis thickness and XY-area value of each simulation box are averaged from trajectory and replica. The arrows indicate the FFA located in the short chain layer. (b.) Trajectory average area compressibility modulus (K_A) of each system, describing their stiffness.

1. **Skin swelling upon hydration, a matter of lipid packing**

When exposed to a humid environment, the SC has been described to absorb and accumulate a significant amount of water, leading to the swelling of the skin[20]. Although this swelling effect is often attributed to the swelling of corneocytes[20], here we show that it is also partly rationalized by a behavior of the SC lipid matrix at the molecular level, by comparing the MD simulations between the REF system and the REF:WAT system. The lipid matrix thickness (D) increased upon water addition by 0.5 nm (see D_{REF} vs $D_{REF:WAT}$ in **Figure 2.a**). This increase mainly comes from a swelling of the liquid-like region where both FFA and water partition in greater amounts, as humidity increases (see the water blue surface in the middle of snapshot for the REF:WAT compared to that of REF, in **Figure 2.a**). Concomitantly to the thickness increase, the XY area (A) decreased by $c.a.$ 0.9 nm^2 (see A_{REF} vs $A_{REF:WAT}$, in **Figure 2.a**), suggesting a change in the skin elasticity, as seen experimentally[21]. This can also be followed by the area compressibility modulus (Ka)[22] presented in **Figure 2.b**. This parameter estimates the force required to deform the membrane system. Comparison between Ka_{REF} and $Ka_{REF:WAT}$ depicted an increased stiffness of the lipid matrix model by 5 mN/m. This means that as the system swells upon hydration, the XY area decreases even faster, which reflects a global improvement of the lateral packing.

2. **Location of NMFs in the SC lipid matrix and molecular impact**

The three NMF molecules (URE, GLY and UCA) are mainly located in the lipid headgroup and the liquid-like regions (**Figure 2.a** and **Figure S1** for a representative snapshot or the density profile, respectively). In these positions, they induce a similar swelling and lipid packing, as that observed when the water concentration was increased (REF:WAT system). Indeed, an increase in the thickness and the area compressibility modulus was observed

(**Figure 2**), when comparing REF to REF:URE (+0.5 nm, +10 mN/m), to REF:GLY (+0.5nm, +8 mN/m) or to REF:UCA (+0.3 nm, +2 mN/m).

UCA and UCO are bulkier than URE and GLY, and their π-conjugated system makes them relatively rigid. Their orientation with respect to the z axis, perpendicular to the membrane surface, is subjected to rapid movements (**Figure S2**). This induced an increase in the XY area (c.a. +0.5nm^2) in the REF:UCA and REF:UCO systems compared to the REF system. Conversely, according to the area compressibility modulus, these two systems have a more packed lipid configuration than REF: +3mN/m and +50 mN/m for REF:UCA and REF:UCO, respectively (**Figure 2.b**). The global trend is that the NMFs behave as water does in terms of membrane properties. This is in line with a previous study showing the water-like effects of NMF on pig skin [9,23,24].

3. NMFs role upon hydration

In the presence of both NMFs and water increase, the swelling and packing effects were not additive. Namely, no further improvement in lipid packing was seen, as area compressibility modulus means and standard deviation were comparable (*e.g.*, 18±2 mN/m for $Ka_{REF:URE}$ vs 22±4mN/m for $Ka_{REF:URE:WAT}$ in **Figure 2.b**). An increase in the swelling of the lipid matrix was observed but not equal to an addition of NMF and water effects. When comparing D_{REF} to $D_{REF:URE:WAT}$, only a 0.6 nm thickness increase was observed (see in **Figure 2.a**), instead of an expected 1.0 nm increase in case of additive NMF and water effects (see 0.5 nm increase from D_{REF} to $D_{REF:URE}$ and 0.5 nm increase between D_{REF} and $D_{REF:WAT}$ in **Figure 2.a**). The same trend can be observed for the other two NMF molecules (GLY and UCA). This suggests an antagonist effect of water and NMF when they are both added to the physiological (REF) system. As water content increases, the literature has shown a high risk of skin lipid matrix disorganization[9]. This was rationalized by the formation of water pools at

the molecular level [23,25]. The early-stage formation of such water pools can be observed here from the MD simulations, at the liquid-like layer of all systems containing an increased water content (see water aggregation in this region for REF:WAT, REF:NMF and REF:NMF:WAT in **Figure 2.a**). Conversely to all other systems, REF:UCO was the only system for which water addition in REF:UCO:WAT increased the XY area, while maintaining its thickness constant (see 24 ± 0.2 nm^2 and 60 ± 2mN/m for REF:UCO vs 24.3 ± 0.3 nm^2 and 31 ± 17mN/m for REF:UCO:WAT in **Figure 2**). This behavior is rationalized in terms of molecular interactions in the Dicussion section.

4. Lipid mobility

As a result of a high hydration level (REF:WAT), a few FFA molecules migrate from the long-chain to the short-chain layers (see arrows in **Figure 2.a**). This migration suggested an increase in lipid mobility upon hydration. **Figure 3** presents three graphics helping to monitor the lipid dynamics and interlayer crossing events. The diffusion coefficient of FFA increased (*i.e.* by $+ 5.10^8$ cm^2/s, see **Figure 3.a**). The same applies to CHL but to a lower extent than FFA (see **Figure S3.a** in supplementary data). This mobility increase can also be indirectly seen by a lower-ordering parameter (**Figure S4**). The increased diffusion rationalizes the FFA infiltration into the short-chain layers, as observed by the presence of FFA in this region in the REF:WAT system (**Figure 3.b**). The CER lipids also indirectly exhibited a greater mobility in the REF:WAT with respect to the REF system, by changing their conformation, namely from splayed to hairpin, with a hairpin conformation increase by more than 10% (**Figure 3.c**). This change is consistent with the changes in CER conformation proposed by Norlén *et. al.*[26] during the formation of the skin barrier in the SC. In the latter publication, the authors explained that lipid dehydration induced molecular stretch-out of the CER from the deeper to the upper layer of the viable epidermis, resulting in a hairpin-to-splayed

conformational change. Here we show the reverse process splayed-to-hairpin upon hydration, highlighting how crucial is the control of skin hydration, a matter of importance that must be seen at the molecular level.

Figure 3 (a.) Diffusion coefficients of free fatty acid (FFA). (b.) Percentage of FFA found in the short chain layers. (c.) Percentage of ceramides (CER) hairpin conformation found among ceramides. GLY and UCA data are also provided in S.I.

Interestingly, when incorporating the NMFs (URE, UCA, and GLY) into the lipid matrix, similar effects were observed, namely more mobile lipids with increased FFA diffusion coefficients (**Figure 3.a**), so short chain layer infiltration by FFA molecules (arrows in **Figure 2.a** and **Figure 3.b**), and presence of CER hairpin conformation (**Figure 3.c**).

Conversely to the other three NMFs, the presence of UCO (REF:UCO system) dramatically

decreased the FFA diffusion coefficients compare to the REF system.

Figure 4 Percentage of H-bonding interactions established between molecules within each system from (a.) to (j.). An H-bond is defined as when the distance between donor and acceptor is less than 3.5 Å and the angle is greater than 120°. The size of each circle is proportional to the amount of H-bonding interaction established by the molecule. The colored percentage value refers to self-interaction. The sum of heteromolecular H-bonds concerning each compound is calculated,

then averaged and reported are in the tables. Percentages over 100% mean more than one H-bond per molecule pair are formed.

Figure 5 (a.) Overall electrostatic potential measured for each system. (b.) Snapshot of UCO (gray) establishing salt bridge (red) and H-bond (blue) with CER molecule (green), CHL (yellow), sodium anion (gray ball) and water.

Discussion: rationalization of underlying interactions

Behind the structural modifications of the lipid matrix of the SC lies a complex H-bond interaction network between all molecules. **Figure 4** summarizes the characterization of this network. In the REF system, CER lipids exhibit the greatest capacity for H-bond interactions (the average heteromolecular H-bonding interaction involving CER being 106%, in **Figure 4.a**). It is worth mentioning that these lipids maintain at least one H-bond with another molecule all along the simulation. This makes a strong interaction maintaining each component at its respective z-position, preventing lamellar disorganization. As such, CER is a keystone molecule of the lipid matrix organization, and therefore of the skin barrier function, as already suggested [27]. The same interaction was suggested in an earlier article using infrared spectroscopy[28]. Later on, in MD simulations of the lipid matrix using CER in hairpin model, others highlighted H-bonds between CER and the amide and carbonyl groups as key in these non-covalent interactions[29]. CHL and FFA are less involved in the H-bond network (the averaged heteromolecular H-bond being 55% and 59%, respectively, in the REF

system, the lowest seen in **Figure 4.a**). As a consequence, the diffusion coefficients of FFA and CHL are relatively high (**Figure S3**).

In the highly hydrated system (REF:WAT), CER lipids are still the most involved in the H-bond structuring network (**Figure 4.b**). However, the number of H-bonds decreased compared to the REF system (*c.a.* -19%, -15%, -6%, -7% for CER, FFA, CHL and WAT, respectively). This leads to a decrease in the intermolecular friction in this system, allowing higher diffusion values than in the REF system. Another consequence is that water molecules diffuse more and populate the liquid-like layer (**Figure S3**). During these movements, water drags FFA molecules, which can also reach the liquid-like layer. This can also be observed by a significant loss of H-bond interaction between CER and FFA (decrease from 106% to 71% in **Figure 4.a** and **b**). During their migration to the liquid-like layer, FFA molecules often change their orientation from parallel to perpendicular with respect to the z axis. They thus create voids that are quickly fulfilled by CER, where splayed-to-hairpin conformational changes can occur (see animation in **supplementary data**). This cascade of events modifies the lipid organization and packing. This is likely impacted by the increase in water-water H-bonding (from 70% to 38% WAT-WAT in **Figure 4.a** and **b**).

In the presence of NMFs, namely REF:URE, REF:GLY, REF:UCA systems, the Result section reported the same lipid packing and mobility effect as in the presence of high-water content (REF:WAT). This is attributed to the presence of an additional partner (the NMF) in the H-bond network, hence weakening H-bond connections between the other partners (see averaged heteromolecular H-bonds in **Figure 4.c, e, g** and **i**). As being the most bonded molecules, CER maintained a strong connection with water and NMF, while decreasing the number of connections with FFA (*e.g.*, -37% CER-FFA, -19% CER-WAT when comparing REF, in **Figure 4.a**, to REF:URE, in **Figure 4.c**). As a result, FFA molecules diffused at a higher rate in the REF:NMF systems, than in the REF system, but slightly lower than in the

REF:WAT systems. This is attributed to the formation of new FFA-NMF H-bonds, as NMFs are small molecules which easily migrate in the different layers together with water, also dragging FFAs. This weakening effect of H-bonding was also observed on a similar membrane model in the presence of a series of excipients[30].

When the hydration level increases in the presence of NMF (WAT:NMF systems), the water-NMF H-bonding connection is the major impacted observable, with +17%, +19%, +23%, from REF:URE to REF:URE:WAT **Figure 4.c** and **d**), from REF:GLY to REF:GLY:WAT (**Figure 4.e** and **f**) and from REF:UCA to REF:UCA:WAT (**Figure 4.g** and **h**), respectively. This modification highlighted a competition between water and NMFs in the formation of H-bonds. The presence of NMFs modulates the impact of water on the lipid matrix organization and its role in the supramolecular H-bond network.

UCO is the only NMF that does not induce packing effect. Conversely, FFA mobility is drastically decreased (**Figure 3.a**). This is attributed to the negative charge state of UCO, which shifted the electrostatic potential from positive to negative, enabling stronger interactions with its surroundings, mainly by creating salt bridges (**Figure 5**). These strong interactions are mainly established with water, almost blocking any diffusion of the two molecules. When the hydration level increases in the presence of UCO (REF:UCO:WAT), the diffusion of FFA is as in the REF system (**Figure 3.a**), as the UCO interaction capacity is saturated with water. Consequently, FFAs also recover their REF-like mobility (**Figure 3.a**), precluding large mobility events such as FFA layer migration and CER conformational changes.

Conclusion

Within a molecular level perspective, this study has confirmed the individual and mutual effects of water and NMFs when inserted in the SC intercellular lipid matrix. Under physiological conditions, CER are keystones of the supramolecular organization of the lipid matrix, by establishing a solid H-bond network with other lipids, contributing to the stability of the lipid matrix. The physical-chemical properties of this lipid matrix were monitored by following the evolutions of this H-bond network. These observations help rationalize the macroscopic impact of hydration and NMF on the skin.

When the amount of water molecules is increased in the lipid matrix, they locate at the lipid polar head region. In turn, they restructure the interaction network by weakening some H-bonds, especially those involving FFAs, thereby increasing their mobility. Consequently, some FFA molecules migrate in between the layers, inducing voids and splayed-to-hairpin conformational switches in CER. This increases the lateral packing of the lipid matrix. The three NMFs, URE, GLY and UCA, exhibit the same effect as that observed with high-water content, as already experimentally suggested [9]. This study also highlights the beneficial interplay between water and NMFs within a large supramolecular H-bond network, as the two compounds work together through preferential interactions to enhance skin hydration.

Acknowledgments

We thank CALI (CAlcul en Limousin), Florent Di Meo and Xavier Montagutelli for computational resources as well as InSiliBio and INSERM for financial support.

References

(1) Dunn, R. R. The Evolution of Human Skin and the Thousands of Species It Sustains, with Ten Hypothesis of Relevance to Doctors. In *Personalized, Evolutionary, and Ecological Dermatology*; Norman, R. A., Ed.; Springer International Publishing: Cham, 2016; pp 57–66. https://doi.org/10.1007/978-3-319-41088-3_5.

(2) Akat, E.; Yenmiş, M.; Pombal, M. A.; Molist, P.; Megías, M.; Arman, S.; Veselỳ, M.; Anderson, R.; Ayaz, D. Comparison of Vertebrate Skin Structure at Class Level: A Review. *Anat. Rec.* **2022**, *305* (12), 3543–3608. https://doi.org/10.1002/ar.24908.

(3) Gunnarsson, M.; Mojumdar, E. H.; Topgaard, D.; Sparr, E. Extraction of Natural Moisturizing Factor from the Stratum Corneum and Its Implication on Skin Molecular Mobility. *J. Colloid Interface Sci.* **2021**, *604*, 480–491. https://doi.org/10.1016/j.jcis.2021.07.012.

(4) Weber, T. M.; Kausch, M.; Rippke, F.; Schoelermann, A. M.; Filbry, A. W. Treatment of Xerosis with a Topical Formulation Containing Glyceryl Glucoside, Natural Moisturizing Factors, and Ceramide. *J. Clin. Aesthetic Dermatol.* **2012**, *5* (8), 29–39.

(5) Fluhr, J. W.; Moore, D. J.; Lane, M. E.; Lachmann, N.; Rawlings, A. V. Epidermal Barrier Function in Dry, Flaky and Sensitive Skin: A Narrative Review. *J. Eur. Acad. Dermatol. Venereol.* **2024**, *38* (5), 812–820. https://doi.org/10.1111/jdv.19745.

(6) Costa-Balogh, F. O.; Wennerström, H.; Wadsö, L.; Sparr, E. How Small Polar Molecules Protect Membrane Systems against Osmotic Stress: The Urea–Water–Phospholipid System. *J. Phys. Chem. B* **2006**, *110* (47), 23845–23852. https://doi.org/10.1021/jp0632440.

(7) Nowacka, A.; Douezan, S.; Wadsö, L.; Topgaard, D.; Sparr, E. Small Polar Molecules like Glycerol and Urea Can Preserve the Fluidity of Lipid Bilayers under Dry Conditions. *Soft Matter* **2012**, *8* (5), 1482–1491. https://doi.org/10.1039/C1SM06273E.

(8) Choe, C.; Lademann, J.; Darvin, M. E. A Depth-Dependent Profile of the Lipid Conformation and Lateral Packing Order of the Stratum Corneum in Vivo Measured Using Raman Microscopy. *The Analyst* **2016**, *141* (6), 1981–1987. https://doi.org/10.1039/C5AN02373D.

(9) Björklund, S.; Andersson, J. M.; Pham, Q. D.; Nowacka, A.; Topgaard, D.; Sparr, E. Stratum Corneum Molecular Mobility in the Presence of Natural Moisturizers. *Soft Matter* **2014**, *10* (25), 4535–4546. https://doi.org/10.1039/C4SM00137K.

(10) Lundborg, M.; Narangifard, A.; Wennberg, C. L.; Lindahl, E.; Daneholt, B.; Norlén, L. Human Skin Barrier Structure and Function Analyzed by Cryo-EM and Molecular Dynamics Simulation. *J. Struct. Biol.* **2018**, *203* (2), 149–161. https://doi.org/10.1016/j.jsb.2018.04.005.

(11) Iwai, I.; Han, H.; Hollander, L. den; Svensson, S.; Öfverstedt, L.-G.; Anwar, J.; Brewer, J.; Bloksgaard, M.; Laloeuf, A.; Nosek, D.; Masich, S.; Bagatolli, L. A.; Skoglund, U.; Norlén, L. The Human Skin Barrier Is Organized as Stacked Bilayers of Fully Extended Ceramides with Cholesterol Molecules Associated with the Ceramide Sphingoid Moiety. *J. Invest. Dermatol.* **2012**, *132* (9), 2215–2225. https://doi.org/10.1038/jid.2012.43.

(12) Norlén, L.; Nicander, I.; Lundsjö, A.; Cronholm, T.; Forslind, B. A New HPLC-Based Method for the Quantitative Analysis of Inner Stratum Corneum Lipids with Special Reference to the Free Fatty Acid Fraction. *Arch. Dermatol. Res.* **1998**, *290* (9), 508–516. https://doi.org/10.1007/s004030050344.

(13) Csizmadia, P. MarvinSketch and MarvinView: Molecule Applets for the World Wide Web. In *Proceedings of The 3rd International Electronic Conference on Synthetic*

Organic Chemistry; MDPI: Sciforum.net, 1999; p 1775. https://doi.org/10.3390/ecsoc-3-01775.

(14)   Humphrey, W.; Dalke, A.; Schulten, K. VMD: Visual Molecular Dynamics. *J. Mol. Graph.* **1996**, *14* (1), 33–38. https://doi.org/10.1016/0263-7855(96)00018-5.

(15)   Yu, Y.; Krämer, A.; Venable, R. M.; Brooks, B. R.; Klauda, J. B.; Pastor, R. W. CHARMM36 Lipid Force Field with Explicit Treatment of Long-Range Dispersion: Parametrization and Validation for Phosphatidylethanolamine, Phosphatidylglycerol, and Ether Lipids. *J. Chem. Theory Comput.* **2021**, *17* (3), 1581–1595. https://doi.org/10.1021/acs.jctc.0c01327.

(16)   Abraham, M. J.; Murtola, T.; Schulz, R.; Páll, S.; Smith, J. C.; Hess, B.; Lindahl, E. GROMACS: High Performance Molecular Simulations through Multi-Level Parallelism from Laptops to Supercomputers. *SoftwareX* **2015**, *1–2*, 19–25. https://doi.org/10.1016/j.softx.2015.06.001.

(17)   M. J. Frisch, et al. Gaussian, Inc., Wallingford CT. Gaussian 09, 2016.

(18)   Hart, P. H.; Norval, M. The Multiple Roles of Urocanic Acid in Health and Disease. *J. Invest. Dermatol.* **2021**, *141* (3), 496–502. https://doi.org/10.1016/j.jid.2020.07.017.

(19)   Case, D. A.; Cheatham, T. E.; Darden, T.; Gohlke, H.; Luo, R.; Merz, K. M.; Onufriev, A.; Simmerling, C.; Wang, B.; Woods, R. J. The Amber Biomolecular Simulation Programs. *J. Comput. Chem.* **2005**, *26* (16), 1668–1688. https://doi.org/10.1002/jcc.20290.

(20)   Mojumdar, E. H.; Pham, Q. D.; Topgaard, D.; Sparr, E. Skin Hydration: Interplay between Molecular Dynamics, Structure and Water Uptake in the Stratum Corneum. *Sci. Rep.* **2017**, *7* (1), 15712. https://doi.org/10.1038/s41598-017-15921-5.

(21)   Rodrigues, L.; Palma, L.; Tavares Marques, L.; Bujan Varela, J. Dietary Water Affects Human Skin Hydration and Biomechanics. *Clin. Cosmet. Investig. Dermatol.* **2015**, 413. https://doi.org/10.2147/CCID.S86822.

(22)   Doktorova, M.; LeVine, M. V.; Khelashvili, G.; Weinstein, H. A New Computational Method for Membrane Compressibility: Bilayer Mechanical Thickness Revisited. *Biophys. J.* **2019**, *116* (3), 487–502. https://doi.org/10.1016/j.bpj.2018.12.016.

(23)   Spears, M. J.; McKillop, K.; Marshall, J. L.; Stone, K. J.; Lilly, N. A.; Warner, R. R.; Boissy, Y. L. Water Disrupts Stratum Corneum Lipid Lamellae: Damage Is Similar to Surfactants1. *J. Invest. Dermatol.* **1999**, *113* (6), 960–966. https://doi.org/10.1046/j.1523-1747.1999.00774.x.

(24)   Malnati, C.; Fehr, D.; Spano, F.; Bonmarin, M. Modeling Stratum Corneum Swelling for the Optimization of Electrode-Based Skin Hydration Sensors. *Sensors* **2021**, *21* (12), 3986. https://doi.org/10.3390/s21123986.

(25)   Warner, R. R.; Stone, K. J.; Boissy, Y. L. Hydration Disrupts Human Stratum Corneum Ultrastructure. *J. Invest. Dermatol.* **2003**, *120* (2), 275–284. https://doi.org/10.1046/j.1523-1747.2003.12046.x.

(26)   Norlén, L.; Lundborg, M.; Wennberg, C.; Narangifard, A.; Daneholt, B. The Skin's Barrier: A Cryo-EM Based Overview of Its Architecture and Stepwise Formation. *J. Invest. Dermatol.* **2022**, *142* (2), 285–292. https://doi.org/10.1016/j.jid.2021.06.037.

(27)   Okuda, M.; Yoshiike, T.; Ogawa, H. Detergent-Induced Epidermal Barrier Dysfunction and Its Prevention. *J. Dermatol. Sci.* **2002**, *30* (3), 173–179. https://doi.org/10.1016/S0923-1811(02)00106-8.

(28)   Rerek, M. E.; Van Wyck, D.; Mendelsohn, R.; Moore, D. J. FTIR Spectroscopic Studies of Lipid Dynamics in Phytosphingosine Ceramide Models of the Stratum Corneum Lipid Matrix. *Chem. Phys. Lipids* **2005**, *134* (1), 51–58. https://doi.org/10.1016/j.chemphyslip.2004.12.002.

(29)   Wang, E.; Klauda, J. B. Molecular Structure of the Long Periodicity Phase in the Stratum Corneum. *J. Am. Chem. Soc.* **2019**, *141* (42), 16930–16943. https://doi.org/10.1021/jacs.9b08995.

(30)   Wennberg, C.; Lundborg, M.; Lindahl, E.; Norlén, L. Understanding Drug Skin Permeation Enhancers Using Molecular Dynamics Simulations. *J. Chem. Inf. Model.* **2023**, *63* (15), 4900–4911. https://doi.org/10.1021/acs.jcim.3c00625.