Load-based divergence in the dynamic allostery of two TCRs recognizing the same pMHC

Ana C. Chang-Gonzalez[1] , **Aoi Akitsu**[2,3,4] , **Robert J. Mallis**[2,3,5] , **Matthew J. Lang**[6,7] , **Ellis L. Reinherz**[2,3,4] , **Wonmuk Hwang***[1,8,9,10]

*For correspondence:
hwm@tamu.edu (WH)

[†]All authors were involved in project conceptualization and writing. A.C.C.G. prepared and performed the simulations. A.C.C.G. and W.H. carried out analysis.

[1]Department of Biomedical Engineering, Texas A&M University, College Station, TX, USA; [2]Laboratory of Immunobiology, Dana-Farber Cancer Institute, Boston, MA, USA; [3]Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA, USA; [4]Department of Medicine, Harvard Medical School, Boston, MA, USA; [5]Department of Dermatology, Harvard Medical School, Boston, MA, USA; [6]Department of Chemical and Biomolecular Engineering, Vanderbilt University, Nashville, TN, USA; [7]Department of Molecular Physiology and Biophysics, Vanderbilt University, Nashville, TN, USA; [8]Department of Materials Science & Engineering, Texas A&M University, College Station, TX, USA; [9]Department of Physics & Astronomy, Texas A&M University, College Station, TX, USA; [10]Center for AI and Natural Sciences, Korea Institute for Advanced Study, Seoul, Republic of Korea

Abstract   Increasing evidence suggests that mechanical load on the $\alpha\beta$ T cell receptor (TCR) is crucial for recognizing the antigenic peptide-loaded major histocompatibility complex (pMHC) molecule. Our recent all-atom molecular dynamics (MD) simulations revealed that the inter-domain motion of the TCR is responsible for the load-induced catch bond behavior of the TCR-pMHC complex and peptide discrimination. To further examine the generality of the mechanism, we perform all-atom MD simulations of the B7 TCR under different conditions for comparison with our previous simulations of the A6 TCR. The two TCRs recognize the same pMHC and have similar interfaces with pMHC in crystal structures. We find that the B7 TCR-pMHC interface stabilizes under ∼15-pN load using a conserved dynamic allostery mechanism that involves the asymmetric motion of the TCR chassis. However, despite forming comparable contacts with pMHC as A6 in the crystal structure, B7 has fewer high-occupancy contacts with pMHC during the simulation. These results suggest that the dynamic allostery common to the TCR$\alpha\beta$ chassis can amplify slight differences in interfacial contacts into distinctive mechanical responses and potentially nuanced biological outcomes.

Introduction

The A6 TCR$\alpha\beta$ and B7 TCR$\alpha\beta$ (herein we call TCR$\alpha\beta$ simply as TCR) are both specific for the same Tax peptide (LLFGYPVYV) of the human T lymphotropic virus 1 (HTLV-1) bound to HLA-A2 (*Garboczi et al., 1996a*,b; *Ding et al., 1998*). A6 and B7 derive from from T cell clones isolated from two patients with HTLV-1-associated myelopathic/tropical spastic paraparesis (*Utz et al., 1996*; *Ding et al., 1998*). They share the same Vβ germline gene (TRBV6-5) and differ only in the Vα germline gene (A6: TRAV12-2; B7:TRAV29DV5), with sequence similarity of 45% for Vα, 96% for Vβ, and 100% for Cα and Cβ (*Ding et al., 1998*). The only structural differences between the two TCRs are from the

residues of the Vα domain and the highly variable complementarity-determining region 3 (CDR3β) loop of the Vβ domain that is crucial for peptide recognition (*Ding et al., 1998*; *Bourcier et al., 2001*; *Rudolph et al., 2006*). In crystal structures, both TCRs bind in a diagonal orientation to the Tax peptide-bound major histocompatibility complex (pMHC), such that both Vα and Vβ contact the MHC α1 and α2 helices (*Garboczi et al., 1996a*). The similar diagonal binding modes are achieved by interactions involving different CDR residues of A6 and B7 contacting largely the same sets of pMHC residues (*Ding et al., 1998*). Only one out of 17 residues contacting pMHC in B7 is also found in the A6 pMHC interaction (*Ding et al., 1998*). T cell response assays demonstrated that single-residue mutations to the Tax peptide elicit different responses in the two TCRs (*Ding et al., 1998*; *Hausmann et al., 1999*). Interfacial interactions may play a role in this TCR-specific response, as residue charges at the surface of the A6 and B7 variable domains show different electrostatic profiles, where the pocket for the Tax peptide Y5 residue is positively charged in A6 but negatively charged in B7 (*Ding et al., 1998*). Overall, the B7 Vα surface has fewer charged residues exposed than A6 Vα. While A6 and B7 recognize Tax-MHC with similar affinities and kinetics, it has been suggested that they achieve binding via different thermodynamic pathways (*Davis-Harrison et al., 2005*).

However, since αβTCR is a mechanosensor (*Reinherz et al., 2023*), the TCR-pMHC bond lifetime under physiological piconewton (pN) level load, rather than equilibrium binding pathway, should be more functionally relevant. In this regard, we have previously used all-atom MD simulations to show that in A6 (*Chang-Gonzalez et al., 2024*) and JM22 (*Hwang et al., 2020*) TCRs, contacts with pMHC are stabilized when an adequate 15–20 pN force is applied. The force-induced stabilization occurs as the asymmetric domain motion of the TCR chassis leads to weakening of the interface with pMHC either in the absence of an adequate level of force or if the sequence of the bound peptide is incompatible with maintaining contacts in the loaded state. The goal of this study is to determine whether the load-dependent control of the binding with pMHC is also present in B7, and identify any differences with A6 that may impact the response of the T-cell while responding to the same pMHC.

We find that the mechanism of dynamic allostery is largely conserved in B7, yet the loaded state does not stabilize contacts with pMHC as robustly as in the A6. Thus, while both A6 and B7 possess comparable equilibrium binding affinity for pMHC, A6 appears to exhibit a stronger catch bond behavior under load. Given the nuanced and dynamic nature of TCRs against the same pMHC, including the effect of ligand abundance (*Akitsu et al., 2024*), difference in mechanical response between A6 and B7 suggests T-cell clones bearing those TCRs may operate differently *in vivo*. Our study also underscores the importance of dynamics when comparing between TCRs that have similar crystal structures and equilibrium binding affinities.

Results

Our simulation systems include the B7 TCR bound to the Tax-pMHC (*Figure 1*A,B) with no, low, and high extensions to apply different loads, and an isolated B7 TCR (*Table 1*). We also used systems without the C-module (Vαβ and Vαβ-pMHC) to study the role of the C-module. We analyzed TCR-pMHC intermolecular and intra-TCR (intramolecular) interactions and domain motion, to test whether the TCR-pMHC interface is stabilized by load, and to find the underlying allostery mechanism that involves the motion of the TCR chassis.

Stabilization of the TCRαβ-pMHC interface with load

Compared to the no- (B7^0) and low-load (B7^{low}) cases, the number of high-occupancy contacts were more numerous for the high-load case (B7^{high}), indicative of a catch bond behavior. Absence of the C-module (Vαβ-pMHC) also promoted more contacts with pMHC, suggesting the allosteric role of the C-module for binding with pMHC (*Figure 1*C). These results agree well with the behaviors seen in A6 and JM22 TCRs in our previous studies (*Chang-Gonzalez et al., 2024*; *Hwang et al., 2020*). However, the number of pMHC contacts with B7 was reduced compared to A6 (*Figure 1*C). This is

Table 1. Simulation systems constructed based on PDB 1BD2 (*Ding et al., 1998*). Extensions of B7^{low} and B7^{high} were selected to yield average low and high loads among simulations scanning different extensions (see Methods). Average load is calculated after 500 ns. The standard deviation (std) in load as measured in 40-ns intervals after 500 ns is shown in parentheses.

Label	Structure	Time (ns)	Extension (Å)	Load (pN)
V$\alpha\beta$	Vα-Vβ only (no pMHC)	1000	–	–
T$\alpha\beta$	TCR$\alpha\beta$ only (no pMHC)	1000	–	–
V$\alpha\beta$-pMHC	V$\alpha\beta$ with pMHC (no C-module)	1000	–	–
B7^0	TCR$\alpha\beta$-pMHC	1050	–	–
B7^{low}	TCR$\alpha\beta$-pMHC	1000	173.7	9.01 (3.96)
B7^{high}	TCR$\alpha\beta$-pMHC	1000	190.0	14.5 (7.20)

despite their comparable number of contacts with the pMHC in crystal structures and comparable equilibrium binding affinity in solution (*Ding et al., 1998*; *Davis-Harrison et al., 2005*). In fact, we had to modify our simulation protocol to avoid premature breakage of contacts between B7 and pMHC when preparing the system for production run under load (see Methods).

Time-dependent behavior of the TCR-pMHC interface further supports the load-mediated enhancement of binding. For this, we calculated instantaneous force and contact occupancy in 40-ns overlapping intervals starting from 200 ns (*Figure 1*D). B7^{high} had overall more inter- and intramolecular contacts (more orange-yellow circles) than B7^{low} (more purple), suggesting that the increased load stiffens the TCR-pMHC complex. B7^0 had fewer TCR-pMHC contacts while intra-TCR contacts increased (horizontal bars in *Figure 1*D), indicating decoupling of the TCR from pMHC in the absence of load.

Figure 1. B7 TCR-pMHC interface. (A) Overview of the base complex used in simulations. Load was applied by holding the C$_\alpha$ atoms of terminal residues (blue spheres at the ends of "added strands") at a given distance from each other. β2m: β2 microglobulin. (B) Magnified view of red box in panel A showing labeled CDR loops and side chains of peptide residues in stick representation. (C) Number of contacts with greater than 50% average occupancy and 80% maximum instantaneous occupancy after the initial 500 ns. Bars: std. Criteria for counting contacts and values for A6 are from *Chang-Gonzalez et al.* (*2024*). (D) Total contact occupancy measured in 40-ns overlapping intervals starting from 200 ns. TCR-pMHC (top; intermolecular) and intra-TCR (bottom; intramolecular) contacts are shown separately. Intra-TCR contacts exclude Cα-Cβ contacts (Methods). Circles with outline: B7^{high}; without outline: B7^{low}. Horizontal bar below each panel: B7^0.

Occupancy heat maps for individual contact residue pairs show reduced or fragmented contacts for B7^0 and B7^{low} (*Figure 2*A,B, more red compared to blue) while B7^{high} and V$\alpha\beta$-pMHC exhibit more persistent contact profiles (*Figure 2*C,D, more blue compared to red). The heat maps

Figure 2. Load dependent behavior of the B7 TCR-pMHC interface. (A–D) Contact occupancy heat maps. (A) B7^0, (B) B7^{low}, (C) B7^{high}, and (D) V$\alpha\beta$-pMHC. Contacts with overall occupancy greater than 30% and instantaneous occupancy greater than 80% are shown. hb: hydrogen bond, np: nonpolar contact. Gray arrow in panel D denotes the approximate time when the initial adjustment of contacts in V$\alpha\beta$-pMHC ends (*Figure 3*B). (E) Location of V-module residues forming contacts with pMHC with greater than 50% average occupancy. The last frame of each simulation is used for visualization. The backbone of the Tax peptide is shown as a purple tube. CDRs are labeled in the first panel. (F) RMSF of peptide backbone C$_\alpha$ atoms after 500 ns. C$_\alpha$ atoms were aligned to those at the beginning of the production run for RMSF calculation.

and contact counts suggest that interfacial contacts were dominated by MHC-Vα (*Figure 3*A). Comparing average counts of high-occupancy pMHC contacts for both A6 and B7 TCRs indicates that Vα formed more contacts with MHC than Vβ, while Vβ formed more contacts with the peptide than it does with MHC (*Figure 3*A). Temporal progression of the number of contacts was measured via the Hamming distance $\mathcal{H}$, the number of the initial high-occupancy contacts lost over time (*Figure 3*B; see Methods). For B7^0, $\mathcal{H}$ rapidly increased and by 200 ns, most of the initial high-occupancy contacts were lost. While the increase in $\mathcal{H}$ for V$\alpha\beta$-pMHC was comparable to that of B7^{low} (*Figure 3*B), the contact occupancy heat maps reveal that V$\alpha\beta$-pMHC maintained contacts after a brief initial adjustment (gray arrow in *Figure 2*D) while contacts were lost in B7^{low} (red *Figure 2*B).

Location of V-module residues forming contacts with pMHC with greater than 50% average occupancy were concentrated along the peptide for B7^{high} and V$\alpha\beta$-pMHC, but scattered in B7^0 and B7^{low} (*Figure 2*E). This trend was also observed in A6 and JM22. Such concentration of high-occupancy contacts may protect them from breakage by water. In A6 and JM22 TCRs, the greater number of contacts with pMHC in the high-load cases and V$\alpha\beta$-pMHC correlated with larger buried surface area (BSA) of the residues forming contacts (*Chang-Gonzalez et al., 2024*; *Hwang et al., 2020*). B7 did not follow this trend, as B7^{high} and V$\alpha\beta$-pMHC had reduced total and per-residue BSA than B7^0 and B7^{low} (*Figure 3*C). This is likely because the fewer high-occupancy contacts in B7 (*Figure 1*C) tend to be more exposed, making the relationship between the BSA and load less direct. Consistent with this, the total BSA of B7 was 67.4 % (B7^0) to 44.2% (B7^{high}) of the corresponding values of A6.

The residues of pMHC forming greater than 50% average occupancy under high load differed between A6 and B7 (*Figure 3*D, top row), as did the location of V-module residues forming respective contacts with the pMHC residues (*Figure 3*D, bottom row), further highlighting their divergence in the interfacial footprint under load. Despite this, the root-mean square fluctuation (RMSF) of C$_\alpha$

Figure 3. Additional characterization of the TCR-pMHC interface. (A) Number of MHC-Vα, MHC-Vβ, peptide-Vα, and peptide-Vβ contacts in A6 and B7 TCRs. Data for A6 are from *Chang-Gonzalez et al.* (*2024*). (B) Hamming distance $\mathcal{H}$. Histograms were calculated using data after the first 500 ns. (C) Total (pink) and per-residue (blue) BSA for interfacial residues after the first 500 ns. (D) Top row: pMHC residues forming contacts with the V-module with average occupancy greater than 50% in the high-load case. MHC residues are shown as sticks and C$_\alpha$ atoms of the peptide residues are shown as spheres. Viewing direction is the same as in *Figure 2*E. Bottom row: V-module residues forming contacts with pMHC residues in top row, shown as spheres. Residues for B7^{high} correspond to those in *Figure 2*E. (E) Distance between the V-module and pMHC. Histograms were calculated using data after the first 500 ns. For V$\alpha\beta$-pMHC and B7^{high}, avg±std after 500 ns are 32.5±0.31 Å and 31.9±0.30 Å, respectively. Dashed line is the value from the crystal structure (31.6 Å).

atoms of the Tax peptide measured after 500 ns was reduced in B7^{high} and V$\alpha\beta$-pMHC compared to B7^0 and B7^{low} (*Figure 2*F), as observed for A6 (*Chang-Gonzalez et al., 2024*), which supports load-induced stabilization of the complex.

We calculated the distance between the V-module and pMHC as another measure of the interfacial stability (*Figure 3*E; Methods). The distance was stably maintained in V$\alpha\beta$-pMHC and B7^{high} whereas it fluctuated more in B7^0 and B7^{low}. Of note, the former two systems maintained the distance greater than that in the crystal structure by 0.3–0.9 Å. Thus, a slight separation engendered by force or in the absence of constraint imposed by the C-module provides room for adjusting residues to form more stable contacts.

CDR3 positions are controlled by load-dependent Vα-Vβ motion

The greater number of Vα-Vβ contacts in B7^0 (*Figure 4*A) is consistent with the increase in total intra-TCR contact occupancy (horizontal bar in *Figure 1*D bottom panel). Without load this does not translate to a stronger TCR-pMHC interface explained above. B7 in general had fewer Vα-Vβ contacts (11.0–16.3) than A6 (15.9–23.1) (*Chang-Gonzalez et al., 2024*). The ~70% reduction in Vα-Vβ contacts for B7 is comparable to the ~50% reduction in contacts with pMHC between the two TCRs (*Figure 1*C).

Vα-Vβ motion was measured via triads (orientational markers) assigned to respective domains and by performing principal component analysis (PCA) (*Figure 4*B; Methods). PC amplitude was the lowest for B7^{high} and V$\alpha\beta$-pMHC (*Figure 4*C), which is consistent with the greater number of Vα-Vβ contacts. Regarding the direction of motion, the mutually orthogonal PC directions can be difficult to interpret (arrows in *Figure 4*B). We instead measured angles between the matching arms of the two triads named $\angle e_i$ ($i = 1, 2, 3$), to examine the Vα-Vβ motion in structurally interpretable directions (*Figure 4*D) (*Chang-Gonzalez et al., 2024*). For example, $\angle e_1$ is the angle between each $\mathbf{e}_1$

Figure 4. B7 TCR chassis motion. (A) Number of Vα-Vβ contacts with greater than 50% average occupancy and 80% maximum instantaneous occupancy after the initial 500 ns. Bars: std. (B) V-module triads $\{e_1, e_2, e_3\}$. Arrows denote directions of the first 3 PC modes for B7^{high} as an example. CDR3s are labeled. (C) Amplitudes for the first 6 PCs. PCA was performed after 500 ns. Transparent bands: std for PCA performed in three overlapping intervals (500–800 ns, 600–900 ns, and 700–1000 ns). (D) Histograms of the V-module triad angles. (E) CDR3 distance vs. triad angles. Transparent bands: std. (F) Number of V-C contacts for each chain measured with the same criteria as in panel A. (G) Average BOC for B7^{low} and B7^{high}. The V-module of B7^{high} is less bent compared to B7^{low}. The arrows for the first 3 V-C PC modes are shown, where PC1 corresponds to the V-C bending motion. (H) Dot products computed between the BOC PC vectors of listed systems. Values closer to 1.0 denote similar V-C BOC direction of motion. (I) V-C PC amplitudes for the first 6 PC modes. Transparent bands: std measured in the same way as in panel C. (J) Differences in amplitudes for the first 3 PCs between matching V- and H-elements of α and β chains. (K) Histograms of hinge angles defined in panel G. (L) CDR3 distance versus hinge angles. Transparent bands: std. (M) V-C hinge angle trajectories over time.

arm from Vα and Vβ, which describes a 'flapping' or 'twisting' motion of the two domains. Since e_2 and e_3 lie approximately parallel to the Vα-Vβ interface, they vary reciprocally, corresponding to a 'scissoring' motion (*Hwang et al., 2020*).

Measuring the distance between CDR3α and CDR3β ("CDR3 distance") revealed that this distance is the shortest for B7^{high} followed by V$\alpha\beta$-pMHC. Comparing CDR3 distance versus triad angles (*Figure 4*E) shows that CDR3 distance varied in opposite directions with $\angle e_2$ and $\angle e_3$, which reflects their reciprocal relation (opposite slopes in *Figure 4*E). In comparison, the CDR3 distance of the B7 crystal structure is 12.0 Å, which is larger than those of B7^{high} and V$\alpha\beta$-pMHC (*Figure 4*E). The slight separation between the V-module and pMHC (*Figure 3*E) in V$\alpha\beta$-pMHC and B7^{high} allows CDR3 loops to come closer together compared to the crystal structure, akin to pinching the central protrusion of the peptide.

Asymmetric V-C bending in the B7 TCR is suppressed with applied load

Next we considered the motion between the V- and C-modules ("V-C motion"). The number of high-occupancy contacts for the Cα-Cβ interface (26.2–27.5) was considerably greater than those

for Vα-Vβ (11.2–16.3), indicating that the C-module acts as a single base for the V-C motion, as noted for A6 and JM22 TCRs (*Hwang et al., 2020*; *Chang-Gonzalez et al., 2024*). Continuing this general feature, there were fewer high-occupancy contacts for the Vα-Cα interface compared to the Vβ-Cβ interface (*Figure 4*F). With two exceptions, residues involved in Vβ-Cβ contacts were the same regardless of force.

The V-C motion was analyzed by using the bead-on-chain (BOC) model that tracks individual domains and the hinge between them (*Figure 4*G). PC motion directions were compared by calculating the dot products between the corresponding PC vectors (*Figure 4*H). PC1 corresponding to the V-C bending in B7^{high} (*Figure 4*G) was similar in other systems, whereas B7^{low} differed the most (darker color for B7^{low} in *Figure 4*H, PC1). Amplitudes of PC1 shows a clear distinction where the unliganded Tαβ and B7^{low} were more mobile than B7^0 and B7^{high} (*Figure 4*I). Comparing PC amplitudes of the elements of the BOC between α and β chains revealed that Vα moves more relative to the C-module than Vβ (*Figure 4*J), similar to A6 (*Chang-Gonzalez et al., 2024*). Amplitude of the hinge motion in the two chains varied, where Hα had greater amplitude in PC1 for B7^{low} compared to B7^{high} (*Figure 4*J, PC1 in bottom row). This suggests a more pronounced asymmetric motion in B7^{low}. The no-load B7^0 Hβ amplitude was larger compared to Hα (*Figure 4*J, bottom, negative value for PC1 of B7^0). For B7^0, the small PC1 amplitude of the overall V-C motion without load (*Figure 4*I) does not suppress the motional asymmetry between α and β chains, while in B7^{high}, the chassis becomes less mobile under load.

Analogous to the triad angle, the V-C angles (*Figure 4*G, ∠TCRα and ∠TCRβ) reveal the motional asymmetry in addition to PCA. As in the case of low-load A6 (*Chang-Gonzalez et al., 2024*), ∠TCRα of B7^{low} shows a wide bimodal distribution (*Figure 4*K). ∠TCRα decreased at around 700 ns as B7^{low} bent more (*Figure 4*M). This would put the V-module in an unfavorable orientation to bind pMHC (*Hwang et al., 2020*). Also, about the dependence of the CDR3 distance on V-C angles, it was the most steady in B7^{high} (*Figure 4*L). This supports the allosteric mechanism by which the asymmetric V-C motion of the whole TCR controls the Vα-Vβ motion and the V-C orientation that in turn affects the stability of the TCR-pMHC interface.

Concluding Discussion

As a protein-protein complex, TCR-pMHC forms a weak interface. A typical heterodimeric protein-protein interface with BSA comparable to that of TCR-pMHC (~1700 Å^2) has sub-μM binding affinity (*Chen et al., 2013*) while the affinity of the TCR-pMHC complex ranges between μM to hundreds of μM (*Rudolph and Wilson, 2002*; *Rudolph et al., 2006*). Given the low equilibrium affinity, recent findings highlight the importance of mechanosensing, where force generated during immune surveillance of αβ T-cells is utilized to discriminate cognate versus non-cognate pMHCs (reviewed in *Zhu et al.* (*2019*); *Liu et al.* (*2021*); *Reinherz et al.* (*2023*)). Our earlier simulation studies of the JM22 and A6 TCRs showed that the dynamical motion of the TCR chassis is responsible for the force-driven stabilization of the contacts with pMHC, *i.e.,* catch bond formation (*Hwang et al., 2020*; *Chang-Gonzalez et al., 2024*). Examining additional TCRs via all-atom MD simulation can inform how the mechanism is adopted. By finding similarities and differences between A6 and B7 TCRs, which recognize the same Tax pMHC, we can elucidate how mechanical force is utilized by each to impact the differential function of T cells.

The present study shows that the dynamic allostery of the TCR chassis observed in A6 (*Chang-Gonzalez et al., 2024*) and JM22 (*Hwang et al., 2020*) is largely conserved in B7. Cα and Cβ domains form extensive contacts to construct a base, and the Vβ-Cβ contacts are more extensive than Vα-Cα contacts. This leads to an asymmetric V-C motion where Vα is more mobile relative to the C-module than Vβ. In turn, the motional asymmetry affects the relative positioning of CDR loops, as measured by the CDR3 distance as well as the orientation of the TCR-pMHC interface relative to the loading direction. Unless an adequate load is applied to the complex to suppress the motion, destabilization of the interface occurs. As noted in A6 and JM22, the C domain renders a disadvantage to binding

except under force. This is indicated by the more interfacial contacts and decreased Vα-Vβ motion of V$\alpha\beta$-pMHC (no C-module) simulations. Therefore, while the Variable domain dictates TCR fit with pMHC, a logical evolutionary pressure would be for the C domains to maximize discriminatory power by adding instability to the TCR chassis.

While dynamic allostery is overall conserved, A6 and B7 differ in the behavior of the TCR-pMHC interface under load. The crystal structures of the two complexes have a comparable number of TCR-pMHC contacts, with a total of 33 for A6 and 29 for B7 in terms of residue pairs, while the number of distinct atom pairs forming contacts are 46 for A6 (*Garboczi et al., 1996a*) and 63 for B7 (*Ding et al., 1998*). During MD simulations, many of these contacts become transient, resulting in fewer high-occupancy contacts. In the thermally fluctuating state, there is a 4-fold increase in the difference in number of contacts between A6 and B7 over that difference for the static crystal structures (*Figure 1*C). Previously, contacts found in crystal structures have been used to explain differences in responses to point mutations to the Tax peptide. For example, the sensitivity of B7 to the mutation of Y5 of Tax was explained based on the Y5-D30α hydrogen bond and stacking of Y5 with Y101β of B7 (Y101β in the present study was numbered Y104β in *Ding et al.* (*1998*) and *Hausmann et al.* (*1999*)). In comparison, the more amenable pocket for Y5 on A6 was suggested to be responsible for the greater tolerance of A6 under mutations of Y5. In MD simulations of B7, the Y5-D30α hydrogen bond is formed only in B7^{high}, after about 300 ns (*Figure 2*C), and the Y5-Y101β nonpolar contact breaks during simulations of B7^0 and B7^{low} (*Figure 2*A,B) while it persists with 69% occupancy in B7^{high} and 52% occupancy in V$\alpha\beta$-pMHC (*Figure 2*C,D). Changes to T cell specificity by point mutations on Y5 are thus unlikely to arise solely from the size or the polarity of its binding pocket as a static structure. Instead, point mutations affect the organization and dynamics of the surrounding contacts in addition to fit (*Chang-Gonzalez et al., 2024*). In a related vein, intermolecular motional network has been suggested to be a controlling factor for allostery in a peptide-SH3 domain complex (*Gomez et al., 2024*), which aligns with the nonlocal and dynamic role of contacts at the TCR-pMHC interface. The inter-domain motion of TCR enables long-range allosteric discrimination of pMHCs, integrating the motion of the two entities such that they influence each other.

The 14.5-pN force on B7^{high} was applied under 190.0-Å extension (*Table 1*), whereas the corresponding high-load simulation for A6 was with 18.2 pN at a slightly lower 187.7-Å extension (*Chang-Gonzalez et al., 2024*). The lower force of the B7 complex despite the larger extension and the overall smaller number of contacts with pMHC in comparison to A6 (*Figure 1*C) leads us to predict that B7 exhibits a weaker catch bond where the peak of the TCR-pMHC bond lifetime is at a lower force. As a related example, PA59 and PA25 TCRs share the same *TRBV* and *TRBJ* genes, and they share 11-aa CDR3β loops that differ only at a single position (Trp vs. Leu) (*Akitsu et al., 2024*). Recognizing the same pMHC (PA$_{224-233}$/D^b), their maximum bond lifetimes and the peak catch bond forces differ considerably, 75 s at 21 pN for PA59, and 13 s at 15 pN for PA25 (*Akitsu et al., 2024*). These differences may facilitate actions of the corresponding T-cells in tissues that present diverse mechanical environments provided by cell movement, adhesion molecules, tight junction, and non-compliant inflammation.

Likewise, T-cells bearing A6 and B7 may also perform differently *in vivo* depending on tissue localization (*Akitsu et al., 2024*) even though they behave similarly *in vitro* in terms of cytotoxicity and secretion of select cytokines (γ-IFN, MIP-1α, and TNFα) (*Ding et al., 1998*; *Hausmann et al., 1999*). Disparate biological outcomes between structurally similar TCRs recognizing the same pMHC are mechanically possible since slight changes in interfacial contacts can result in altered distribution of loads across the TCR chassis so that changes in its dynamics affect interaction with CD3 signaling subunits (*Reinherz et al., 2023*). Structural details of the dynamic amplification and propagation of the recognition signal warrant further investigation.

Computational Methods

Structure preparation

B7 TCR$\alpha\beta$-pMHC was built from PDB 1BD2 (*Ding et al., 1998*) using CHARMM (*Brooks et al., 2009*; *Hwang et al., 2024*). Non-numeral residue IDs in the PDB were renumbered to follow sequential numbering used in the present study. We used MODELLER (*Šali and Blundell, 1993*) to generate coordinates for missing loops in the Cα domain (S133-K136 and S170-D172 in the PDB numbering scheme) followed by a brief energy minimization. We visually verified MODELLER results, comparing generated loops to those of the related A6 TCR (*Garboczi et al., 1996a*; *Ding et al., 1998*). The constant domain of TCRα (Cα) was also missing coordinates for F204-S210 (F198-S204 after renumbering), which were added with the TCRα linker as detailed below. Disulfide bonds were assigned between cysteine residues as defined in the PDB file. Crystal waters within 2.8 Å from the protein were kept for the truncated structures, and all waters were kept for the full structure.

Histidine protonation state was determined to promote hydrogen bond formation with neighboring residues. The histidine N$^\delta$ atom was protonated as follows: MHC residues 3, 93, 114, 145, 151, 188, 260; β2m residues 13, 51; TCRα all histidine residues; and TCRβ residues 29, 47, 154. For the remaining histidine residues, the N$^\epsilon$ atom was protonated.

As done in *Hwang et al.* (*2020*) and *Chang-Gonzalez et al.* (*2024*), we extended the MHC and TCR$\alpha\beta$ termini as handles for applying positional restraints. For MHC, we used UniProt P01892 to add [276]LSSQPTIPI[284]. For TCRα we used GenBank AAA60627.1 to add [205]CDVKLVEKSFETDT[218]. For TCRβ we used GenBank AAC08953.1 to add [245]CGFTSESYQQGVLSA[259]. We placed an interchain disulfide bond between αC205-βC245. Added strands in the initially straight conformations were relaxed to a state similar to that in *Figure 1*A by performing a series of brief energy minimization and MD simulation with the FACTS implicit solvent model (*Haberthür and Caflisch, 2008*).

Truncated structures were built based on the prepared B7 TCR-pMHC complex as:

- V$\alpha\beta$: The last residues were αP110 and βV113.
- T$\alpha\beta$: The last residues were αD206 and βG246 (no C-terminal strands).
- V$\alpha\beta$-pMHC: includes V$\alpha\beta$, pMHC, and β2m. The last residue of MHC was L276.
- B7^0: includes T$\alpha\beta$, pMHC, and β2m. The last residue of MHC was L276.

MD simulation protocol

Solvation, energy minimization, heating, and equilibration of the B7 complexes followed the protocol in *Chang-Gonzalez et al.* (*2024*), except for systems which include the pMHC, where we modified the preparation protocol prior to production runs as detailed below.

Laddered extensions

Applying the same protocol as done for A6 to achieve the laddered extensions in B7 resulted in substantial breakage of the TCR-pMHC contacts within the first 50 ns in several production runs. To mitigate this, we introduced distance restraints to selected atom pairs forming contacts between the TCR and pMHC to prevent them from breaking during preparatory simulations. This ensured that the complex could structurally adapt as we modified the extension distance, yet all laddered extensions maintained a core set of initial TCR-pMHC contacts. Atom pair distance restraints were removed in production runs.

Twelve atom pairs between TCR and pMHC were selected that were within 5 Å of each other in the equilibration restart file of the TCR-pMHC complex with added linkers. A 2-kcal/[mol·Å^2] flat-bottom harmonic restraint potential was applied to keep the atom pair distance within the value at the end of the equilibration run. Then a 2-ns CPT simulation was carried out while also applying a 1-kcal/[mol·Å^2] harmonic potential to the C$_\alpha$ atoms of the C-terminal MHC (I284), TCRα (T218), and TCRβ (A259) residues. Production run followed upon removing the atom pair distance restraints . During the production run, the 1-kcal/[mol·Å^2] harmonic potential to the TCR and MHC end-residues and a 10 Å distance restraint between αT218 and βA259 were applied. Throughout

the production run, we intermittently measured average and rolling force in 40 ns intervals and found these to be around 10 pN, an ideal target for the low-load simulation. This simulation is the 173.7 Å B7^{low} system reported in *Table 1*.

Using the structure at the end of the 2-ns simulation with the TCR-pMHC atom-pair distance restraints, we increased the extension by 8 Å by moving the center of the 1-kcal/[mol·Å^2] harmonic potential on the end-residue C$_\alpha$ atoms by 4 Å at each end. While keeping the TCR-pMHC atom-pair distance restraints applied, we launched another 2-ns simulation under the increased extension. Atom pair distance restraints were then removed and the production run was launched. The extension averaged after 500 ns of this simulation was 181.7 Å. Following the same way, we increased the extension by another 8 Å, which led to the 190.0 Å B7^{high} system reported in *Table 1*. We also decreased the extension by 8 Å from the initial 173.7-Å extension, which led to a 165.7-Å extension.

Among the 4 extensions tested, the 181.7-Å extension was not selected primarily because the average force of the simulation from 500 ns to 850 ns (the total length of the simulation), was 9.26 pN, only barely higher than the reported load for B7^{low}. For the 165.7-Å simulation, the average force from 500 ns to 900 ns (total length of the simulation) was 15.7 pN. We had observed this high force at low extension for A6 TCR (*Chang-Gonzalez et al., 2024*) and attribute this to folding of the flexible added strands leading to contacts between the stands and the TCR constant domains.

V$\alpha\beta$-pMHC and B7^0

We also applied a 2-kcal/[mol·Å^2] flat-bottom harmonic distance restraint during preparatory simulations of V$\alpha\beta$-pMHC and B7^0. We attempted to use the same set of atom pairs as in the laddered extension simulations, but considerable interface breakage occurred, likely due to changes in interfacial contacts after equilibration in these systems. We thus selected different atom pairs for V$\alpha\beta$-pMHC and B7^0. For consistency, we selected 12 atom pairs, the same number as in the laddered extensions, and distributed in the same way between the Tax peptide or MHC residues to Vα or Vβ residues. The distance restraint was applied to the atom pairs for a 2-ns CPT simulation, then released for production runs.

Systems without load

The following additional restraints were used for systems without load.

- V$\alpha\beta$, T$\alpha\beta$: no positional restraints were applied.
- V$\alpha\beta$-pMHC: we applied a weak 0.01-kcal/[mol·Å^2] harmonic positional restraint to the backbone C$_\alpha$ atoms of MHC α3 (P185-L276) to prevent large transverse rotation of the whole molecule in the orthorhombic box.
- B7^0: we applied a 0.2-kcal/[mol·Å^2] harmonic positional restraint to the backbone C$_\alpha$ atoms of MHC α3 with RMSF less than 0.5 Å calculated from the simulations of B7^{low}. These residues were: L201-Y209, T240-Q242, T259-H263.

Production runs

Production runs were performed similar to *Chang-Gonzalez et al.* (*2024*). We used OpenMM (*Eastman et al., 2017*) with the CHARMM param36 all-atom force field (*MacKerell Jr et al., 2004*) and the particle-mesh Ewald method to calculate long-range electrostatic interactions. We used an Ewald error tolerance of 10^{-4} which is $1/5$ of the default value in OpenMM and a 12 Å cutoff distance for nonbonded interactions. The complexes were simulated at 300 K with a 2-fs time step using the Nose-Hoover integrator in OpenMM. Production run lengths are in *Table 1*.

Trajectory analyses

Analysis methods are detailed in *Chang-Gonzalez et al.* (*2024*). Below we mainly explain B7-specific residue selections. Out of ~1-μs production runs, we excluded the initial 500 ns when calculating the average and standard deviation of the number of contacts, BSA, CDR3 distance, PCA, and triad

and V-C angles. With a coordinate saving rate of 20 ps, this leaves at least 25,000 frames for analysis.

V-module to pMHC distance

The distance from TCR V-module to pMHC (*Figure 3*E) was measured between the center of mass of the C_α atoms of the same residues used to build the V-module triads (described below) to the center of mass of five C_α atoms from each of the central 4 strands forming the β-sheet floor located above the $\alpha1$ and $\alpha2$ helices of MHC (20 MHC atoms in total; *Figure 1*A). These were R6–T10, I23–Y27, Q96–G100, and Y113–A117. RMSF of these residues after 500 ns was below 1.4 Å in all B7 systems, so the measured distance is minimally affected by the intra-domain conformational motion.

CDR3 distance

Distance between CDR3α and CDR3β (*Figure 4*E,L) was measured using the midpoint between backbone C_α atoms of two residues at the base of each CDR3, which are E93 and K97 for CDR3α and S94 and E102 for CDR3β.

V-module triads

We assigned triads (*Figure 4*B) based on the backbone C_α atoms of the stably folded β-sheet core of each variable domain (*Hwang et al., 2020*; *Chang-Gonzalez et al., 2024*). Selected residues for triad assignment of the B7 systems were as follows. For Vα, I19-Y24, F32-K37, H71-I76, and Y87-M92. For Vβ, T20-Q25, M32-Q37, D73-L78, and Y89-S94. Prior to triad assignment we aligned all complexes to the first frame of B7^{low} using the selected residues to monitor the relative motion between the two triads without global translation nor rotation.

V-C BOC

We assigned BOCs (*Figure 4*G) as detailed in *Hwang et al.* (*2020*); *Chang-Gonzalez et al.* (*2024*). To place beads for the C-module, we used the following residues. For Cα, A118-R123, V132-D137, Y153-T158, and S171-S176. For Cβ, L143-T148, L157-N162, L190-R195, and F208-Q213. For the hinges, we used: αN114 for Hα, and βD116 and βL117 for Hβ. We aligned all complexes to the backbone C_α atoms of the selected residues of B7^{low} then built BOCs to monitor the motion of the V-module relative to the C-module.

Acknowledgments

This work was funded by US National Institutes of Health Grants P01AI143565 and R01AI136301. Simulations were performed by using computers at the Texas A&M High Performance Research Computing facility.

References

Akitsu A, Kobayashi E, Feng Y, Stephens HM, Brazin KN, Masi DJ, Kirkpatrick EH, Mallis RJ, Duke-Cohan JS, Booker MA, Vincenzo Cinella and WWF, Holliday EL, Lee JJ, Zienkiewicz KJ, Tolstorukov MY, Hwang W, Lang MJ, Reinherz EL. Parsing digital or analog TCR performance through piconewton forces. Sci Adv. 2024; 10(33):eado4313.

Bourcier KD, Lim DG, Ding YH, Smith KJ, Wucherpfennig K, Hafler DA. Conserved CDR3 regions in T-cell receptor (TCR) CD8+ T cells that recognize the Tax11-19/HLA-A* 0201 complex in a subject infected with human T-cell leukemia virus type 1: relationship of T-cell fine specificity and major histocompatibility complex/peptide/TCR crystal structure. J Virol. 2001; 75(20):9836–9843.

Brooks BR, Brooks III CL, Mackerell Jr AeD, Nilsson L, Petrella RJ, Roux B, Won Y, Archontis G, Bartels C, Boresch Sao. CHARMM: the biomolecular simulation program. J Comput Chem. 2009; 30(10):1545–1614.

Chang-Gonzalez AC, Mallis RJ, Lang MJ, Reinherz EL, Hwang W. Asymmetric framework motion of TCR$\alpha\beta$ controls load-dependent peptide discrimination. eLife. 2024; 13:e91881.

Chen J, Sawyer N, Regan L. Protein–protein interactions: General trends in the relationship between binding
affinity and interfacial buried surface area. ProtSci. 2013; 22(4):510–515.

Davis-Harrison RL, Armstrong KM, Baker BM. Two different T cell receptors use different thermodynamic
strategies to recognize the same peptide/MHC ligand. J Mol Biol. 2005; 346(2):533–550.

Ding YH, Smith KJ, Garboczi DN, Utz U, Biddison WE, Wiley DC. Two human T cell receptors bind in a simi-
lar diagonal mode to the HLA-A2/Tax peptide complex using different TCR amino acids. Immunity. 1998;
8(4):403–411.

Eastman P, Swails J, Chodera JD, McGibbon RT, Zhao Y, Beauchamp KA, Wang LP, Simmonett AC, Harrigan
MP, Stern CD, Wiewiora RP, Brooks BR, Pande VS. OpenMM 7: Rapid development of high performance
algorithms for molecular dynamics. PLoS Comput Biol. 2017; 13(7):e1005659.

Garboczi DN, Ghosh P, Utz U, Fan QR, Biddison WE, Wiley DC. Structure of the complex between human T-cell
receptor, viral peptide and HLA-A2. Nature. 1996a; 384(6605):134–141.

Garboczi DN, Utz U, Ghosh P, Seth A, Kim J, VanTienhoven E, Biddison WE, Wiley DC. Assembly, specific binding,
and crystallization of a human TCR-$\alpha\beta$ with an antigenic Tax peptide from human T lymphotropic virus type
1 and the class I MHC molecule HLA-A2. J Immunol. 1996b; 157(12):5403–5410.

Gomez SM, Gonzalez TI, Vasa SK, Linser R. Allostery at a Protein-Protein Interface Harboring an Intermolecular
Motional Network. Angew Chem Intl Ed. 2024; p. e202411472.

Haberthür U, Caflisch A. FACTS: Fast analytical continuum treatment of solvation. J Comput Chem. 2008;
29(5):701–715.

Hausmann S, Biddison WE, Smith KJ, Ding YH, Garboczi DN, Utz U, Wiley DC, Wucherpfennig KW. Peptide
recognition by two HLA-A2/Tax11–19-specific T cell clones in relationship to their MHC/peptide/TCR crystal
structures. J Immunol. 1999; 162(9):5389–5397.

Hwang W, Austin SL, Blondel A, Boittier ED, Boresch S, Buck M, Buckner J, Caflisch A, Chang HT, Cheng X, Choi
YK, J, Chu JW, Crowley M, Cui Q, Damjanovic A, Deng Y, Devereux M, Ding X, Feig M, et al. CHARMM at 45:
Enhancements in accessibility, functionality, and speed. J Phys Chem B. 2024; 128(41):9976–10042.

Hwang W, Mallis RJ, Lang MJ, Reinherz EL. The $\alpha\beta$TCR mechanosensor exploits dynamic ectodomain allostery
to optimize its ligand recognition site. Proc Natl Acad Sci USA. 2020; 117(35):21336–21345.

Liu B, Kolawole EM, Evavold BD. Mechanobiology of T Cell Activation: To Catch a Bond. Annu Rev Cell Dev Biol.
2021; 37:65–87.

MacKerell Jr AD, Feig M, Brooks CL. Improved treatment of the protein backbone in empirical force fields. J
Am Chem Soc. 2004; 126(3):698–699.

Reinherz EL, Hwang W, Lang MJ. Harnessing $\alpha\beta$ T cell receptor mechanobiology to achieve the promise of
immuno-oncology. Proc Natl Acad Sci USA. 2023; 120(27):e2215694120.

Rudolph MG, Stanfield RL, Wilson IA. How TCRs bind MHCs, peptides, and coreceptors. Annu Rev Immunol.
2006; 24(1):419–466.

Rudolph MG, Wilson IA. The specificity of TCR/pMHC interaction. Curr Op Immunol. 2002; 14(1):52–65.

Šali A, Blundell TL. Comparative protein modelling by satisfaction of spatial restraints. J Mol Biol. 1993;
234(3):779–815.

Utz U, Banks D, Jacobson S, Biddison WE. Analysis of the T-cell receptor repertoire of human T-cell leukemia
virus type 1 (HTLV-1) Tax-specific CD8+ cytotoxic T lymphocytes from patients with HTLV-1-associated dis-
ease: Evidence for oligoclonal expansion. J Virol. 1996; 70(2):843–851.

Zhu C, Chen W, Lou J, Rittase W, Li K. Mechanosensing through immunoreceptors. Nat Immunol. 2019;
20(10):1269–1278.