Genomic and structural insights into Jyvaskylavirus, the first giant virus isolated from

Finland

Gabriel Magno de Freitas Almeida[1] [*], Miika Leppänen[2], Bruna Luiza de Azevedo[3], Jonatas Santos Abrahao[3], Julien Andreani[4,5], Davide Zabeo[6†], Janne Ravantti[7], Iker Arriaga[8], Nicola GA Abrescia[8,9] [*], Lotta-Riina Sundberg[2]

[1] The Norwegian College of Fishery Science, Faculty of Biosciences, Fisheries and Economics, UiT - The Arctic University of Norway, Tromsø, Norway

[2] University of Jyväskylä, Department of Biological and Environmental Science and Nanoscience Center, Jyväskylä, Finland

[3] Universidade Federal de Minas Gerais, Institute of Biological Sciences, Department of Microbiology, Belo Horizonte, Minas Gerais, Brazil

[4] Aix Marseille Univ, MEPHI, Marseille, France

[5] IHU-Méditerranée infection, Marseille, France

[6] Diamond Light Source, Harwell Science and Innovation Campus, Didcot, UK.

[7] University of Helsinki, Molecular and Integrative Biosciences Research Programme, Helsinki, Finland

[8] Structure and Cell Biology of Viruses Lab, CIC bioGUNE, Basque Research and Technology Alliance (BRTA), Derio, Spain.

[9] Ikerbasque, Basque Foundation for Science, Bilbao, Spain.

†Current Address: Department of Chemistry and Molecular Biology, University of Gothenburg, Gothenburg 41390, Sweden.

*Corresponding authors: Gabriel Magno de Freitas Almeida (gabriel.d.almeida@uit.no) and Nicola GA Abrescia (nabrescia@cicbiogune.es)

ABSTRACT

Giant viruses of protists are a diverse and likely ubiquitous group of organisms. Here, we describe Jyvaskylavirus, the first giant virus isolated from Finland. This clade B marseillevirus was found in *Acanthamoeba castellanii* from a composting soil sample in Jyväskylä, Central Finland. Its genome shares similarities with other marseilleviruses, including conserved Nucleocytoviricota genes, histone-like genes, and three unique ORFans. Helium ion microscopy and electron microscopy of infected cells unraveled stages of the Jyvaskylavirus lifecycle, such as single virions attached to the cell membrane, likely triggering endocytosis, and virus-rich vesicle formation within the cytoplasm. We also reconstructed the Jyvaskylavirus particle to 6.3 Å resolution using cryo-EM. The ~2,500 Å diameter virion displays structural similarities to other *Marseilleviridae* giant viruses. The capsid comprises of 9,240 copies of the major capsid protein ORF184, which possesses a double jellyroll fold arranged in trimers forming pseudo-hexameric capsomers. Below the capsid shell, the internal membrane vesicle encloses the genome. Through cross-structural and -sequence comparisons with other *Marseilleviridae* using AI-based software in model building and prediction, we elucidated ORF142 as the penton protein, which plugs the twelve vertices of the capsid. Five additional ORFs were identified, with models predicted and fitted into densities that either cap the capsomers externally or stabilize them internally.

The isolation of Jyvaskylavirus, along with other virus-like particles, not only suggests that these viruses may be widespread in the boreal environment but also provide insights extendable to other marseilleviruses, addressing the previously lacking structural data on minor capsid proteins.

INTRODUCTION

Viruses defy paradigms of classical biology and are agents of change and innovation for

biological research. In 2003 researchers were surprised by the description of the *Acanthamoeba*

polyphaga mimivirus (APMV) (**La Scola et al 2003**). Hidden in plain sight for over a century of

microbiology research, APMV and other giant viruses were evading discovery for three main

reasons: their large size traps them in filters commonly used in virological works, their structure

and size makes them visible by Gram-staining, and their protist hosts are not as well studied as

other host species (**Colson et al 2017 ; Queiroz et al 2022**). Common features of their double-

stranded DNA genomes, similarity in a few core genes, relative independence of host

transcription machinery, partial or complete cytoplasmic replication and formation of viral

factories include giant viruses in the nucleo-cytoplasmic large DNA viruses (NCLDVs) group

(**Iyer et al 2001 ; Boyer et al 2010**). Recently the International Committee of Taxonomy of

Viruses (ICTV) classified giant viruses as members of the phylum *Nucleocytoviricota* of the

kingdom *Bamfordvirae* in the realm *Varidnaviria* (**Walker et al, 2019 ; Simmonds et al 2023**).

Isolation efforts and metagenomic data in the last two decades have revealed that giant viruses

are ubiquitous in the environment (**Moniruzzaman et al 2020 ; Schulz et al 2020**). Giant virus

particles or their DNA have been found from the Antarctica to the Siberian permafrost, in deep-

sea sediments and many other sources including urban environments and diverse sample types

(**Andrade et al 2018 ; Legendre et al 2015 ; Backstrom et al 2021 ; Colson et al 2017**). To

support the sharp rise in metagenomic studies, microbial isolation and characterization are

welcomed, and needed for deeper understanding of these entities and the viral world. Isolation of

giant viruses revealed the intriguing mosaicism of Marseillevirus, the complex structure and

mysterious large genome of Pandoravirus, the remarkable structure and translational potential of

Tupanvirus, and many other characteristics of these organisms that have changed our view on the

concept of viruses and their evolution (**Boyer et al 2009** ; **Phillippe et al 2013** ; **Abrahao et al**

2018). It is expected that many new insights will emerge as more isolates are found (**Aherfi et al**

2016 ; Colson et al 2017). Increased effort in isolating giant viruses might also bring new

discoveries and help in understanding their distribution and importance worldwide. Isolation of

viruses allows for in-depth structural studies using their whole virions.

An increased number of icosahedral NCLVD, thanks to the advances in cryo-EM, are now being

targeted for structural analysis despite the challenges that their very large dimensions pose (>

1,500 Å diameter). Cryo-EM structures for African swine fever virus, *Aureococcus*

anophagefferens virus, *Cafeteria roenbergensis* virus, Faustovirus, Marseillevirus, Medusavirus,

Pacmanvirus, Paramecium bursaria Chlorella virus 1, *Phaeocystis pouchetii* virus , and more

recently for Melbournevirus have elucidated their complex architecture, major capsid protein

fold, and assembly organization (**Andreani et al 2017 ; Andres et al 2020 ; Burton-Smith et**

al 2021 ; Gann et al 2020 ; Klose et al 2016 ; Okamoto et al 2018 ; Shao et al 2022 ; Wang et

al 2019 ; Watanabe et al 2022 ; Xiao et al 2017 ; Yan et al 2005). Another example of the

importance of the combination of isolation and structural studies is the description of FLiP, a

missing link between ssDNA and dsDNA viruses in Finland (**Laanto et al 2017)**. However,

FLiP is a bacteriophage and there have been no previous studies on the isolation of giant viruses

in the Finnish boreal ecosystem or have their structural characterization been described so far.

Microbial ecology and virus-host interactions are still poorly studied in many environments,

including boreal ecosystems. Here, we report the occurrence of giant viruses in Central Finland

and the characterization of Jyvaskylavirus, a new virus belonging to the clade B of the

Marseilleviridae family as determined by genome analysis. This virus was isolated from a

composting soil sample in the city of Jyväskylä. We used helium ion microscopy and

transmission electron microscopy to visualize the early attachment events of the virus to its

amoebal host (*Acanthamoeba castellanii*) and the infected cell. Using cryo-EM, we also

determined the three-dimensional (3D) structure of Jyvaskylavirus at a resolution of 6.3 Å.

Cross-structural and sequence comparison allowed us to identify five proteins that compose the

capsid essential for assembly, with their corresponding models reliably placed into density.

Jyvaskylavirus description is the first step in unveiling the diversity of giant viruses from Finland

and from the Nordic Countries, exemplifying that these viruses are also present in the boreal

ecosystems with a still unknown role for microbial ecology.

RESULTS

Giant viruses are present in Finland

During the summer 2019 a preliminary isolation attempt of local amoebas and giant viruses

using samples collected in Central Finland hinted at the presence of these viruses in Finnish

samples (**Supplementary Material and Supplementary Figure 1**). Although viral-like particles

were seen, working with the local protist host presented challenges that initially hindered the

virus characterization. We followed up this preliminary screening with a larger isolation effort

using three reference host strains: *Acanthamoeba castellanii*, *A. polyphaga* and *Vermamoeba*

vermiformis. Ninety-six environmental samples collected in Central Finland were tested using

the three different amoebal hosts. Of these samples, ten had viral presence confirmed by the

appearance of cytophatic effect (CPE) in cultures of *A. castellanii* followed by visualization of

negative stained viral-like particles by electron microscopy (10.41% isolation success in *A.*

castellanii). However, some of the samples belonged to the same sample group and viral

morphology was similar between some of the isolates. Adjusting for sample type and unique

viral morphologies found, we have then tested 53 unique sample types and found three distinct

viral morphologies (5.66% isolation success in *A. castellanii*) (see **Supplementary Table 1**).

Some of the CPE-positive samples contained mixed viral populations. Samples collected from

experimental aquaria had a mixture of three different viral morphotypes with dimensions ranging

from approximately 200 to 300 nm side-to-side, highlighting a previously unexplored diversity

of giant viruses in this setting (**Figure 1A-C**). One sample from a recirculating aquaculture tank

also possessed the diamond-shaped and loose-capsid morphologies seen in the aquaria samples.

The high amount of nutrients in the fish aquaria or tanks might have enriched for protozoa and

thus favoured our giant virus isolation, in a pattern similar to the original methodology for

isolating giant viruses (**Arslan et al 2011**). However, our attempts to isolate these viruses using

limiting dilutions and more passages did not result in reliable purified samples, making it

impossible to proceed with further characterization.

The isolate from sample 85 had a single morphotype of approximately 200 nm in size, displaying

an icosahedral shape with particles that were mostly genome-filled and a few that were empty

(**Figure 1D**). The empty particles possibly displayed an internal membrane beneath the capsid

(**Figure 1D, centre**).

This sample was collected from the municipal Waste Treatment Centre, Jyväskylä, Finland. The

sample was in soil state and originated from a mixture of 70% garden waste, 15% woodchips and

15% pre-treated biowaste. Virus Ac-85 was chosen as a model to represent the first giant virus

isolate from Finland and named Jyvaskylavirus as a homage to the city of its isolation.

Jyvaskylavirus has a fast replication cycle in its *A. castellanii* host, only causes CPE in *A.*

polyphaga or *V. vermiformis* in high concentrations likely due to cytotoxic effect of the virus

preparation, it is sensitive to chloroform treatment (10% for ten minutes) and is stable for long

periods (up to 109 days) even at 37 °C (**Supplementary Material and Supplementary Figure**

2).

Jyvaskylavirus belongs to the *Marseilleviridae* family

Jyvaskylavirus contains a 359,967 base pairs (bp) genome, with a 42.80% GC content and 388

predicted ORFs coding for proteins with sizes that vary between 99 and 1,525 amino acids. The

positive DNA strand codes for 186 ORFs whereas the other 202 are in the negative strand

(**Figure 2A**). During genome annotation, some of the *Nucleocytoviricota* conserved proteins

were detected, including the DNA polymerase family B and the A32-like packaging ATPase.

The most of Jyvaskylavirus genes (about 67%) code for uncharacterized proteins. The second

major function category is the DNA replication, recombination and repair genes, including three

histone-like proteins (ORFs 215, 216, 320), a typical marseillevirus genome feature (**Figure 2B**)

(**Bryson et al 2022**). Three new ORFans were detected (ORFs 264, 265, 289), representing

genes that have no significant similarity with any other sequences from the database used in this

analysis. ORF 264, 265, and 289 codes for putative proteins with 108, 157 and 119 amino acids,

respectively. We searched for translation-related genes, and found three translation factors,

including a translation initiation factor (ORF 154), an elongation factor (ORF 318) and a peptide

chain release factor (ORF 28). No tRNAs or aminoacyl tRNA synthetases genes were found.

Furthermore, most of the BLASTp best hits for Jyvaskylavirus amino acid sequences matched

Lausannevirus or Port-miou virus that are phylogenetically related to marseilleiviruses from

lineage B. This observation can also be reinforced by the phylogeny based on DNA polymerase

family B, which clusters the Jyvaskylavirus within *Marseilleviridae* family, together with other

marseilleviruses from clade B (**Figure 2C, Supplementary Figure 3A**). When analyzing the

genome synteny of different marseilleviruses genomes, it is shown that Jyvaskylavirus presents

similarity blocks comparable to those from clade B marseilleviruses (**Supplementary Figure**

3B).

Jyvaskylavirus attachment to host cells and extracellular virion clusters by scanning

helium ion microscopy

To visualize the interaction of Jyvaskylavirus virions with its *A. castellanii* host cells we used a

scanning Helium Ion Microscope (HIM). This imaging technology has proven successful in the

study of bacteriophages and their host bacteria in the past (**Leppanen et al 2018**). We adapted

the sample preparation methods to image cultures of *A. castellanii* infected with Jyvaskylavirus.

Sample preparation was made by allowing cells colonize silicon chips treated with poly-l-lysine

followed by infection and fixation, avoiding any need for cell scraping and pelleting. We

successfully imaged the amoeba and Jyvaskylavirus virus in their near-original state at fixation

(**Figure 3**).

By HIM imaging *A. castellanii* cells appear as mainly oval-shaped with spined structures

(acanthopodia) protruding from the cell surface; furthermore, a cell with an elongated shape was

captured likely in the process of cell division (**Figure 3A**). No cysts were seen in these samples.

At a closer range, cell surfaces appear rugose, full of crevices, and contains virions attached to it

(**Figure 3B**). One of strategies for marseillevirus entry is the triggering of an endosomal-

stimulated pathway (**Arantes et al 2016**). We captured a cell with virions attached in regions

showing invaginations of the cell membrane, probably indicating the process of endocytosis

(**Figure 3C**). These craters, surrounded by walls with varying degrees of steepness, differ in size

although the estimated diameter to accommodate at least one virion is about 300 nm (**Figure 3C,**

inset). Alongside intact cells with attached virions, we also observed a broken cell nearby, with

most of its cell contents released (**Figure 3D, centre**). Extracellular vesicles larger than 500 nm

in size containing multiple virions inside were visualized close to burst cells (**Figure 3D, left**

inset). These virion clusters are important for marseilleviruses, which are smaller than other

giant viruses, to reach the required threshold for triggering amoebal phagocytosis and thus

initiating their infection cycle through this entry mechanism (**Arantes et al 2016**). Details of the

ruptured cell contents, revealing the presence of large internal spherical vesicles and intracellular

virions, are shown in **Figure 3D** (right inset). Single virions were also observed on the substrate,

suggesting that they could attach to foraging cells.

Jyvaskylavirus forms intracellular vesicles and viral factories inside *A. castellanii* host cells

Intracellular details of the replication cycle of Jyvaskylavirus were imaged by transmission

electron microscopy using ultrathin sectioning of cells undergoing cytopathic effect at 24 hours

post-infection (hpi). At 24 hpi, the cells are overtaken by virus production, with intracellular

vesicles of varying dimensions (*e.g* 1-2 μm), which contain already apparent virions (**Figure**

4A). Large viral factories are observed, with some occupying almost the entire cell area (**Figure**

4B). The nucleus and other cellular components, such as mitochondria, can be clearly

distinguished within the cell sections (**Figure 4A-C**). Several individual virions also populated

the cytoplasm. One imaged vesicle is juxtaposed to the nuclear membrane and contains several

assembled virions with clear icosahedral shape. These virion-rich intracellular vesicles are likely

the source of the extracellular ones imaged by HIM in **Figure 3D (left inset)**. Membrane-related

structures located near or nearly attached to the luminal side of the large vesicle are discernible

(**Figure 4C-D**). These membrane-derived assembled structures might serve during virus

morphogenesis (**Figure 4D**). Inspection of the interior of viral factories unravels distinct stages

of particle formation (**Figure 4E**). A putative assembly path can be extrapolated from the

analysis of images of particles, which transition from a half-assembled icosahedron to a particle

with an open vertex through the orderly accumulation of capsid proteins (**Figure 4F**). This open

vertex is potentially used for genome packaging, and it is subsequently closed by the plugging of

peripentonal and penton proteins.

Jyvaskylavirus three-dimensional architecture

Jyvaskylavirus icosahedral virion, determined to 6.3 Å resolution as judged by the gold-standard

Fourier shell correlation possesses a diameter of ~ 2,516 Å (vertex-to-vertex) and a triangulation

number $T = 309$ ($h = 7$, $k = 13$) (**Figure 5A**, **Supplementary Figure 4 and Supplementary**

Table 2). The protein capsid shell, approximately 120 Å thick, can be geometrically represented

by trisymmetrons and pentasymmetrons as similarly done with other NCLDVs (**Figure 5B**)

(**Sinkovits & Baker 2010**). The trisymmetron and pentasymmetron comprises 136 and 30

pseudo-hexameric capsomers, respectively, along with one penton complex, containing five

copies of the penton protein (**Figure 5B-C**). Jyvaskylavirus icosahedral asymmetric unit (IAU) is

composed of 51 pseudo-hexameric capsomers plus 1/3 of the capsomer sitting on the icosahedral

3-fold axis (**Figure 5C**). Beneath the capsid, the membrane vesicle follows icosahedral

symmetry, although it also displays a high degree of sphericity (**Figure 5A, left**). This internal

membrane vesicle encloses the genome. Some particles, excluded during 2D classification,

showed heterogeneous membrane morphologies indicating their structural fragility. At the five-

fold axis between the capsid and the membrane, there is relatively weak density suggesting the

presence of additional proteins, along with a clear bulging of the membrane vesicle at the five-

fold (radius of curvature of ~275 Å) (**Figure 5A, left**).

Jyvaskylavirus structural proteins composing the capsid

At 6.3 Å resolution alpha-helical secondary structural motifs are identifiable while the separation

of β-strands became clearer beyond 5 Å. Our density corresponding to the capsomer displays

unequivocally pseudo-hexameric morphology with the characteristic footprint of trimers formed

by vertical double jellyroll (DJR) seen in other viruses of the kingdom *Bamfordviriae* (**Andres et**

al 2020 , Ravantti et al 2020 , Simmonds et al 2023). A cross-section of the density shows the

β-barrel walls, further supporting that the major capsid protein (MCP) possesses a vertical DJR

fold (**Figure 5C, inset**). During genome annotation based on sequence homology with other

marseilleviruses, ORF184 was identified as a potential MCP. We submitted ORF184 to

AlphaFold3, which predicted a model with a DJR fold (**Figure 5D**) (**Abramson et al 2024**). The

fitting of this model into density leaves no doubts about ORF184 being the MCP and a total of

154 copies of the MCP compose the IAU (**Figure 5C, E**).

Then, we structurally identified the penton protein of Jyvaskylavirus as ORF142 by using the

latest version of ModelAngelo software with the hidden Markov model (HMM) search procedure

against the publicly available cryo-EM block-reconstructed capsid densities of the

Melbournevirus, a clade A *Marseilleviridae*, at ~3.5 Å resolution (EMD-37188, 37189, 37190)

(for details see Materials and Methods and **Supplementary Figure 5**) (**Jamali et al 2024**). This

identification was based on the hypothesis of structural conservation among conserved capsid

components and partial protein sequence conservation among members of the *Marseilleviridae.*

Model prediction of the ORF142 in Alphafold3 produced a compact eight stranded β-barrel

(BIDG – CHEF) typical of a jellyroll, elaborated by extended loops between strands DC and EF

and a long C terminal tail (**Figure 6A, left**). The jelly-roll core of the penton proteins showed a

reasonable fit into the corresponding Jyvaskylavirus densities at 6.3 Å resolution (**Figures 5E**

and 6A).

Additional density was observed atop the pseudo-hexameric capsomers, capping the central

region formed by the ORF184 jellyroll towers, and likely stabilizing the trimer from the exterior.

Using the same methodological strategy as for the penton protein, we identified ORF121 as this

cap protein, which possesses a β-barrel fold and forms a trimer, with its N-terminal ends

inserting into the crevice formed by the MCP jellyroll towers (**Figure 6B**). Beneath the capsid

shell, weaker density is visible at varying distances from the virus center and at different

locations within the trisymmetron (**Figures 6B and 7A, top left**). Their positioning resembles

that seen in Tokyovirus and Melbournevirus, which have been linked to pentasymmetron protein

components (PCs) and minor capsid proteins (mCPs) with proposed roles in scaffolding,

cementing, and zippering (**Chihara et al 2022 ; Burton-Smith et al 2021**). However, no

information is currently available regarding their corresponding ORFs and/or 3D models.

Using the above-mentioned approach (**Supplementary Figure 5**), we identified four additional

Jyvaskylavirus proteins - ORF36, ORF97, ORF153, and ORF119 – located beneath the

pentasymmetron capsomers. We predicted their 3D models using AlphaFold3 and positioned

them within the corresponding higher-resolution Melbournevirus densities, where the core of the

different molecules generally fit well within the constraints of the map (**Supplementary Figures**

6-8). We then placed all the newly identified and predicted Jyvaskylavirus 3D models into our

cryo-EM density map at 6.3 Å resolution (**Figure 7**). Intriguingly, ORF119 was also identified as

the protein that runs along the edge of the trisymmetron facets, acting as a glue between adjacent

capsomers belonging to two different trisymmetrons (**Figure 7B** and **Supplementary Figure 8**).

The relative orientation of the ORF119 molecule beneath the pentasymmetron and along the

edges of two trisymmetrons is about 90 degrees. However, not all the visible density beneath the

capsid in the deposited Melbournevirus block-based reconstructed maps could be accounted for

by the fitted structures. Interestingly, all Alphafold3 predicted ORFs presented unordered

regions, mostly at the N- and C-terminal ends, with lengths varying depending on the protein and

their location. While we did not attempt further remodelling of these flexible regions into

Melbournevirus densities, a more complete representation of Melbournevirus is achievable. To

this end, we identified the corresponding Jyvaskylavirus ORFs in Melbournevirus through

sequence comparison (**Supplementary Table 3**).

Further, a distinctive feature of the internal membrane vesicle is the bulging of the lipid bilayer

at the five-fold equivalent to that found in Tokyovirus (**Figure 5A, left** and **Figure 6A**)

(**Chihara et al 2022**). The identified PCs crown this region, and it is plausible that, along with

yet unidentified proteins, they act as effectors of the bulging by tethering the membrane. The

radius of the bulging is likely related to the extent of their localization beneath the

pentasymmetron.

DISCUSSION

Jyvaskylavirus is the first characterized giant virus from Finland. So far the only other isolated

giant viruses from the Nordic Countries are the still uncharacterized Lurbovirus from Sweden

and a collection of viruses capable of infecting microalgae isolated in Southern Norway (**Kördel**

et al 2021 ; **Sandaa et al 2001 ; Johannessen et al 2015**). There is genomic evidence of

NCLDV presence in the Greenland ice sheet and of a high diversity of giant viruses, including

the detection of marseilleviruses, in the Loki's Castle deep-sea vent located in the Mid-Atlantic

Ridge between Iceland and Svalbard (**Perini et al 2024** ; **Backstrom et al 2021**). However, no

virus was isolated in these studies. Jyvaskylavirus belongs to clade B of the *Marseilleviridae*

family, so far making it the northernmost known member of the family. The first marseillevirus

was isolated from France in 2007 (**Boyer et al 2009**). Now there are more than 60 isolates

known, obtained from varied sample sources in seven countries over five continents (**Sahmi-**

Bounsiar et al 2021). Our genetic analysis showed that the Loki's Castle marseilleviruses group

together and that they are not close to Jyvaskylavirus, probably forming unique clusters among

themselves (**Supplementary Figure 3**). Jyvaskylavirus is also unique for being the first found

from a composting sample, demonstrating that these giant viruses are found in water and soil

samples worldwide.

For the characterization of Jyvaskylavirus we integrated complementary imaging techniques,

including the use of Helium Ion Microscopy for imaging the virus and its *A. castellanii* host.

This demonstrates the applicability of this technique to giant viruses and sets the methods for

sample preparation, opening new ways for the study of giant viruses and interactions with their

host.

Ultrathin TEM sectioning enabled us to capture snapshots of a putative assembly pathway,

unravelling the progressive formation of the capsid. Single-particle cryo-EM provided critical

insights into the structure of Jyvaskylavirus and its evolutionary relationship with other viruses.

The 3D reconstructed cryo-EM density at 6.3 Å resolution clearly recapitulates the architecture

($T = 309$; vertex-to-vertex diameter 2,500 Å) and capsid protein organization observed in

Melbournevirus and Tokyovirus, other members of the *Marseilleviridae* family (**Burton-Smith**

et al 2021 ; Chihara et al 2022). Although it is not possible to distinguish individual β-strands

at this resolution, with the aid of Alphafold3, we unequivocally identified and placed the

predicted ORF184 MCP vertical double jelly-roll fold model into density. Additionally, we

identified and predicted the fold of the penton protein encoded by ORF142 to adopt a vertical

jellyroll topology. This model nicely fitted the corresponding density, plugging the center of the

pentasymmetrons. There is evident structural conservation of the penton protein fold with that of

other viruses such as penton proteins P31 in bacteriophage PRD1 or VP9 in archaeal virus

HCIV-1 of the *Bamfordviria* and *Helvetiavirae* kingdoms, possessing vertical double and single

jellyroll MCPs, respectively (**Abrescia et al 2004** ; **Santos-Perez et al 2019**). This type of

penton complex acts like a plug-and-play system, able to incorporate various host-recognizing

vertex proteins. These proteins are interchangeable and adapt to environmental evolutionary

pressures (**Gil-Carton et al 2015**). The clear presence of a network of scaffolding, cementing,

and minor proteins beneath, previously observed in ASFV and Tokyovirus, but also in smaller

viruses such as PBCV-1, adenovirus, and PRD1, reiterates the universal requirement for the

assembly of larger virus of an increase number of ancillary proteins tethering the MCPs at the

vertices, within the facets and along the facets **(Abrescia et al 2004 , Burton-Smith et al 2021 ;**

Liu et al 2010 , Wang et al 2019, Chihara et al 2022, Shao et al 2022). In this study, through

cross-structural and sequence comparisons with available Melbournevirus block-based

reconstructed densities and combination of AI-based modelling software (**Zhu et al 2018**

;**Burton-Smith et al 2021** ;**Jamali et al 2024**), we were able to elucidate the identity and the

fold of five additional viral capsid components in Jyvaskylavirus: ORF36, 97, 119, 153 and 121

(and the corresponding ORFs in Melbournevirus). The former four are located beneath the

pentasymmetron while ORF121 is the protein capping on the top the capsomers and it shares its

fold with Marseillevirus Noumeavirus NMV_189 protein (PDB ID 7QRR) (*e.g.*, rmsd 1.5 Å, 146

Cα aligned) and virophage Zamilion vitis protein Zav19 (PDB ID 7QRJ).

Remarkably, two ORF119 molecules that form a homodimer are also used to glue at their bases

capsomers located at the edge of adjacent trisymmetrons. The orientation of the ORF119

molecule below the pentasymmetron differs from that along the edge of the trisymmetron,

indicating that its binding mode depends on the structural environment. The reuse of the same

protein in different regions of the facet for multiple functions highlights ORFs optimization and

genetic parsimony. Particularly intriguing is that while AlphaFold3 predicts the core of most

identified proteins, satisfying the constraints of the map, some regions are poorly ordered. One

clear case is ORF153 which is predicted to possess about 40% of the residues unstructured. We

propose that these regions together with flexible terminal ends are not merely a limitation of

AlphaFold3's predictive capabilities but rather reflect specific functional characteristics of these

proteins, which may fold and adapt through protein partnering.

It has been shown that the conserved flexibility and function of intrinsically disordered proteins

despite their often-fast rate and tolerance for mutations, plays critical physiological roles across

all organisms and viruses (**Brown et al 2011**). Protein P30 of the lipid-containing bacteriophage

PRD1, structurally determined more than 20 years ago, and predicted as a disordered protein,

was the first to be identified possessing a string-like structure running through adjacent virus

facets, interacting with capsomers along the edges and cementing them together (**Abrescia et al.,**

2004). Giant viruses, however, are structurally more complex, with a larger number of minor

capsid proteins holding the virion together beneath the pentasymmetron and trisymmetron. While

our current study advances our understanding of this complexity by identifying various ORFs

and positioning their models in the density map, deciphering their atomic interactions and

specific roles in particle assembly will require higher resolution for the different components.

Jyvaskylavirus isolation is the first step in better understanding the giant virus diversity in the

boreal environment. Our preliminary bioprospection attempt using a locally isolated amoebal

host suggested a high diversity of unique viral like particles, while our screening of fish tank

samples revealed mixed populations of other novel viruses with unique morphologies. The origin

of the viruses found in aquarium water is unknown, and they could be originated from the

ground water used to keep the fish or the fish microbiome itself. Viruses from groundwater have

been described before, making this substrate an interesting source for screening for new viral

groups. In addition, electron microscopy analyses revealed high structural diversity of giant

viruses and virus-like particles in soil, suggesting a gap in knowledge in the diversity and

ecology of environmental virus-host-interactions (**Hylling et al 2020** ; **Fischer et al 2023**).

Despite the technical challenges that prevented us from characterizing all isolated viruses in this

study, our findings underscore the importance that broader sampling and new isolation attempts

are essential to determine the distribution of these viruses. The use of host species isolated

locally might avoid isolation bias towards the reference strains giving a better idea of viral

diversity and allowing the description of unique viral groups.

In summary, marseilleviruses found in diverse locations and environments worldwide show a

strictly conserved architecture, using equivalent pseudo-hexameric building blocks and a

plethora of ancillary proteins. We suggest that this adaptation relies on the number, fold, and

length of disordered regions in the minor capsid proteins, as well as the membrane composition.

These factors not only facilitate correct assembly but would also modulate the virus's structural

stability across various environments.

MATERIALS AND METHODS

Samples used for screening, amoebal host cultivation and viral isolation

To get an overview on presence of giant viruses in Finland and their virus-host interactions,

water and soil samples were collected from Jyvaskyla (Finland) and used fresh for viral isolation.

Previously collected and stored samples were also used, e.g. aquaculture water samples,

recirculating aquaculture filter pellets and frozen water from experimental aquaria (**Runtuvuori-**

Salmela et al 2020 ;**Almeida et al 2019a** ;**Almeida et al 2019b**). Sample details can be found at

the **Supplementary Table 1**. Water samples were collected directly into Eppendorf tubes. Solid

samples like soil and composting material (roughly one fourth of an Eppendorf tube) were

resuspended in 1ml of PAS buffer (120 mg NaCl, 4 mg MgSO$_4$.7H$_2$O, 4 mg CaCl$_2$.2H$_2$O, 142

mg Na$_2$HPO$_4$, 136 mg KH$_2$PO$_4$ in 1 litre of water) and strongly vortexed for resuspension

(**Thomas et al 2006**). Each sample had the following antibiotic mix added to avoid fungal and

bacterial contamination during the isolation process: penicillin (0.14 mg/ml), gentamycin (50

mg/ml), amphotericin B (0.25 µg/ml), ciprofloxacin (0.004 mg/ml), vancomycin (0.004 mg/ml)

and doxycycline (0.02 mg/ml). Indicated concentrations are the final concentration of each

antibiotic in solution.

Isolation was made using *Acanthamoeba castellanii*, *A. polyphaga* and *Vermamoeba vermiformis*

hosts. All three host strains were kindly donated by Dr. Bernard La Scola and grown using PYG

media at room temperature (~25 °C) (**Jensen et al 1970**). For the isolation process, cells were

mixed with 50 µl of the samples in 96 well plates. Cell density was controlled by resuspending

confluent T25 flasks in 1 milliliter of PYG media and diluting it 1/1000 before adding 200 µl to

each well in the plates. The antibiotics mentioned above were also present in PYG media used

for preparing the isolation plates. Cells were monitored daily for five days for the appearance of

cytopathic effect (CPE). In case no effect appeared, the plates were frozen, thawed and 25 µl of

each well used as samples for a second passage. Three passages of all samples were made in all

three hosts. Negative samples were those that no CPE appeared after the third passage.

Growth curve, chloroform sensitivity and stability

Growth curves were made by infecting confluent cells in 96 well plates with a low MOI (1/1000

dilution of the virus stock). For each dilution four wells were prepared. At the indicated time

points the samples were frozen and then, after two freeze-thaw cycles, 50 µl of each well were

transferred to a new 96 well plate containing confluent *A. castellanii* cells for titration.

Chloroform sensitivity was made by exposing a 900 µl viral aliquot to 10% chloroform for ten

minutes, inside an Eppendorf tube. As control, 900 µl of the same viral stock was mixed with

100 µl of PAS and incubated for ten minutes. After incubation both were serially diluted and

titrated (four replicates per dilution). Stability was evaluated by diluting a viral stock serially (-1

to -11, 1ml aliquots) and storing the viral dilutions at room temperature (~25 °C), cold room (~8

°C) and inside an incubator (37 °C). Samples were stored for 109 days and then titrated. The

incubation time used is this long because of the SARS CoV2-related lockdowns that happened

during this experiment. All titers were calculated by tissue culture infectious dose (TCID)50

(**Reed and Muench 1938**).

Sequencing and genomic data

For DNA extraction, a confluent T25 was infected at a low MOI and the supernatant was

harvested after the appearance of full CPE. Two aliquots of 2ml each were subjected to nuclease

treatment and capsid precipitation by $ZnCl_2$ (**Santos 1991**). After proteinase K treatment the

samples were mixed with ethanol:guanidine and DNA extraction was finished using the

GeneJET Genomic DNA Purification Kit (Thermo Fisher). DNA paired-end libraries (2×250-

bp) were constructed with 1 ng of the viral genome as input using the Nextera XT DNA kit

(Illumina, Inc., San Diego, USA) and sequenced on the Illumina MiSeq for 39-hours, the same

strategy employed by **Brahim Belhaouari et al 2022**.

The assembled genome was submitted to open reading frames prediction (ORFs) using

GeneMarkS (**Besemer et al 2001**). Also, a search for tRNA genes was performed using

ARAGORN program (**Laslett & Bjorn 2004**). Only ORFs having more than 150 nucleotides

were considered to analysis. Similar sequences for each predicted protein were searched using

BLASTp (expect threshold: 10-3) against NCBI non-redundant protein sequences (nr) database.

To perform synteny analysis, complete genome sequences from different marseilleviruses were

obtained in GenBank. The sequences underwent manual curation to correct distortions caused by

the circular topology of marseilleviruses genomes. Then, synteny analysis were conducted using

MAUVE program with default parameters (**Darling et al 2004**).

Phylogenetic trees were constructed based on DNA polymerase and major capsid protein (MCP)

amino acid sequences. The sequences used for alignments were obtained using BLASTp (expect

threshold: 10-3) against NCBI non-redundant protein sequences (nr) database. The alignments

were performed using MUSCLE executed through MEGA X program (**Edgar 2004 ; Kumar et

al 2018**). Maximum-likelihood phylogenetic trees were constructed using IQ-tree software

(version 1.6.12) with 1,000 bootstrap replicates as branches support (**Nguyen et al 2014**). Best-

fit substitution models were selected by ModelFinder algorithm implemented in IQtree

(**Kalyaanamoorthy et al 2017**). The phylogenetic trees were visualized using iTOL (**Letunic &

Bork 2021**).

Microscopy

Samples with noticeable CPE during the viral isolation process were prepared for checking the

presence of negatively stained viral-like particles (VLPs). Two microliters of the lysates

(supernatant straight from the isolation plate) were added to a microscopy grid and incubated for

two minutes at room temperature. The excess liquid was removed with a water soaked Whatman

paper. Then five microliters of 2% phosphotungstic acid was added to the grid and its excess was

removed after an incubation of two minutes. The grids were left for drying at room temperature

for five minutes and used for imaging at a Jeol JEM-1400 electron microscope straight away.

Samples destined for thin sectioning were prepared by infecting a T75 flask with a low

multiplicity of infection (MOI). Twenty-four hours later, when the cytopathic effect started to

appear, cells were harvested from the flask and pelleted by centrifugation (10 minutes at 2000 g).

The pellet was resuspended in 2.5 % glutaraldehyde 0.1 M sodium phosphate buffer and kept

under slow rotation for one hour at room temperature. After the incubation the cells were

pelleted again, resuspended in 0.1 M sodium phosphate buffer and sent for blocking and thin

section preparation. The samples were also imaged at a Jeol JEM-1400 electron microscope.

Substrates for helium ion microscopy were prepared by incubating silicon chips in poly-l-lysine

(Sigma Aldrich) for five minutes followed by two wash steps in sterile water and left to dry

overnight. *A. castellanii* cells were seeded in 24 well plates containing the poly-L-lysine coated

substrates and infected at different MOI. After the appearance of cytopathic effect, the samples

were fixed with 2.5 % glutaraldehyde in 0.1 M sodium cacodylate for two hours followed by

three washes wash with 0.1 M sodium cacodylate buffer. After that, the samples were stained by

incubation for one hour with 1 % osmium tetroxide followed by three washes in 0.1 M sodium

cacodylate and a second staining with 0.1 % tannic acid for twenty minutes. Dehydration was

done with sequential exposure to increasing concentrations of ethanol, using the following

percentages: 35, 50, 70, 85, 95 and 99 %. Each exposure was made for thirty minutes, except for

99 %, which had one thirty minutes exposure followed by a second overnight exposure to ensure

proper dehydration. Ethanol washing was followed up by a critical point drying step using

twenty-four cycles in a Leica EM CPD300 equipment. Macroscopic sample structure did not

change during the drying. Imaging was made using the Zeiss Orion Nanofab Helium Ion

Microscope from the University of Jyväskylä Nanoscience Center with acceleration voltage 30

kV and ion current 0.2-0.3 pA. Electron flood gun was used during imaging to mitigate positive

charging by the ions.

Virus production for cryo-electron microscopy and data collection

A large stock of Jyvaskylavirus was prepared and purified for cryo-EM analysis. Twelve

confluent T75 flasks were infected at a low MOI. After the appearance of full CPE the flasks

were frozen-thaw to lyse still intact cells, and then all the flask contents were moved to falcon

tubes. After one additional freeze-thaw cycle, a brief centrifugation (500 g, 5 minutes, 10 °C)

was made to clear the lysate from cell debris. Then the viruses were pelleted (10.000 g, 60

minutes, 10 °C) and resuspended in 700 µl of PAS. The resuspended viruses were loaded into a

10-50 % sucrose gradient and centrifuged (6500 g, 90 minutes, 10 °C). The viral band of the

gradient was collected, mixed with PAS buffer to dilute the sucrose and the viruses were pelleted

again. The final pellet was resuspended in 150 µl of PAS buffer, aliquoted and shipped at 10 °C

to the CIC bioGUNE for cryo-EM analysis. Upon arrival the sample was vitrified either using a

Vitrorobot Mark IV (Thermofisher) or an Automatic Plunge Freezer EM GP2 (Leica). As a

'quality control' step, some of the grids were inspected using the in-house JEM-2200FS/CR

(JEOL, Ltd.) cryo-TEM equipped with a K2 bioquantum camera. The remaining grids were

shipped for high-resolution imaging at eBIC - Diamond Light Source (Didcot, UK) in line with

democratic access to large infrastructure (**Stuart et al 2016**). Four data collections on a Titan

Krios 300 kV with a K3 camera were performed at a nominal magnification of ×64,000,

resulting in a final pixel size of 1.34 Å/pix or 1.35 Å/pix depending on the microscope used

(**Supplementary Table 2**). Briefly, samples were vitrified on Quantifoil Cu R2/2 or R2/1 300

mesh grids and then collected over 40-45 fractions (with a dose per frame of ≈ 1 e-/Å^2), with

defocus ranges from -0.6 to 3 μm. As we had approximately one particle per hole, different

software were used at each data collection to test which strategy would yield the highest number

of particles (EPU, TOMO5 and Serial EM; for details in each data collection see

Supplementary Table 2). After four data collections we obtained 3,742 useful particles from

17,720 movies.

3D reconstruction of Jyvaskylavirus

MotionCorr2 was used to correct the induced beam-shift across the frames within the movies,

while the individual movies' CTF was estimated using the CTFFIND4 software (**Rohou &**

Grigorieff 2015 ; Zheng et al 2017). Particle auto-picking for the first data collection was

performed in crYOLO, training the model with 10 movies (**Wagner et al 2019**); however, the

picking process was supervised. For the remaining acquisitions, particles were manually selected

in RELION 3.1 due to the limited number of movies available (**Scheres 2011**). Virions were

extracted into a 1,000 x 1,000 pixel box, resulting in a final pixel size of 3.087 Å. For each

dataset, several rounds of 2D and 3D classifications were performed before 3D refinement, with

the initial 3D reference being generated *ab-initio* from a limited number of particles from the

first data collection. Subsequently, the particles were re-extracted and re-centered, then

combined with those from the other data collection after undergoing the same pre-processing

workflow. Further classifications led to a homogeneous class comprising 3,742 particles

(**Supplementary Table 2**), which then underwent 3D icosahedral refinement with the original

pixel size (1.34 Å/pix) on a 2,304 pix box. This was achieved by using the Picasso HPC at the

University of Malaga node of the Spanish Supercomputing Network (RES). Computing

resources used for the above refinement included 3 tasks, 32 cpu per task and 1.6 TB of memory

(using --pad 1). Finally, Ewald sphere correction was applied to the final map, resulting in a

resolution of 6.3 Å and a notable improvement in the map interpretability and FSC curve

(**Supplementary Figure 4**).

Structural analysis of Jyvaskylavirus

Using AlphaFold3 we generated an initial atomic model of the MCP corresponding to the

ORF184 identified in this study by sequence comparison with other *Marseillaviridae*

(**Abramson et al 2024**). The best-ranked predicted model had a confidence level of 95 % based

on the predicted local-distance difference test (pLDDT), which ranged from 28.7 to 98.9 %

across the residues. The MCP model was manually fitted into the density corresponding to the

pseudo-hexameric capsomer using COOT graphic software (**Emsley & Cowtan 2004**). The three

molecules were rigid body refined into the 6.3 Å resolution map using PHENIX real-space

refinement (**Afonine et al 2018**) leading to a CC_{mask} of 52 %.

To identify the penton protein located at the vertices as well as some of the additional visible

densities corresponding to the scaffolding and cementing proteins, we used the following

strategy. Owing that Jyvaskylavirus belongs to the clade B of the *Marseilleviridae* (current

study) and that block-based derived cryo-EM maps at the 2-, 3- and 5-folds of Melbournevirus

(clade A) at about 3.5 Å resolution have been recently made available in the Electron

Microscopy Database (EMD-37189, EMD-37190) we used the latest version of ModelAngelo

software (v1.0.12) without any fasta sequence input to build and identify the likely models for

the penton and ancillary proteins in the Melbournevirus densities **(Burton-Smith and Murata**

2023 ;Jamali et al 2024). The sequences of the resulting identified amino acids for the built

fragments were then parsed using the hidden Markov model (HMM) routine in ModelAngelo

against the list of ORFs of Jyvaskylavirus produced in this study. This was performed with the

expectation of high sequence homology between the two viruses, as indicated by our

comparative sequence analysis (**Figure 2**). For the penton protein, ORF142 was identified, and

similarly ORF36, 97, 153 and 121 for the ancillary under the five-fold and cap protein,

respectively. Then, the corresponding full 3D models were predicted using AlphaFold3 and fitted

into the Melbournevirus and Jyvaskylavirus cryoEM density using the fit-into-map routine in

ChimeraX (**Meng et al 2023**). In the case of Melbourne virus, the corresponding ORFs were

determined by using the Jyvaskylavirus ORFs as templates in blastp

(https://blast.ncbi.nlm.nih.gov/Blast.cgi?PAGE=Proteins).

Data Availability

The Jyvaskylavirus genome has been deposited in GenBank under the accession number PQ284187 as part of BankIt submission 2862875.

Raw stacks movies are uploaded to EMPIAR (EMPIAR-XXXX/*in process*), linked to the EMDB accession code 51613 corresponding to the reconstructed cryo-EM map at 6.3 Å resolution. Predicted Jyvaskylavirus PDB models using ModelAngelo and Alphafold have been deposited at Biostudies under the accession number S-BSST1654.

Declaration of interest

The authors declare no conflicts of interest.

Author contributions

G.M.dF.A, L.R.S. and N.G.A.A. conceived the project. G.M.dF.A. performed bioprospection experiments and virus isolation. G.M.dF.A. and ML performed the EM and HIM imaging. J.A., B.L.A. and J.S.A. were responsible for sequencing and bioinformatics analyses. I.A. performed all cryo-EM experiments, including sample vitrification, image processing, cryo-EM density analysis and implementation of ORFs identification and models fitting under the guidance of N.G.A.A. D.Z. collected cryo-EM data with support from I.A. and N.G.A.A. G.M.dF. L.R.S. and N.G.A.A. wrote the manuscript with input from all authors.

Acknowledgements

We would like to thank Mr Petri Papponen (University of Jyvaskyla, Finland) for help with the electron microscopy, Msc. Marjaana Hassani (University of Jyvaskyla, Finland) for donating the composting sample used for Jyvaskylavirus isolation and the staff at the EM Platform at the CIC bioGUNE for infrastructural support and preliminary cryo-grid screening. Professor Bernard La Scola (Aix Marseille University, France) is thanked for donating the amoeba strains used and for

his technical and intellectual input in this project. Yun Song at the UK's national Electron Bio-

imaging Centre (eBIC) at Diamond Light Source (DLS) is acknowledged for her assistance in

setting up the SerialEM script. We are also grateful to DLS for access and support of the cryo-

EM facilities at eBIC [under proposal CM30316], funded by the Wellcome Trust, MRC and

BBRSC and to the Spanish Supercomputing Network (RES, Red Española de

Supercomputación) for Picasso resources provided by the Supercomputing and Bioinnovation

Center (Malaga University) to BCV-2022-3-0015.

This study was funded by the Research Council of Finland (grant #346772), the Centre for New

Antibacterial Strategies (CANS) at the Arctic University of Norway (project ID #2520855), the

Spanish Ministry of Science, Innovation and Universities, and the State Research Agency (AEI)

for project PID2021-126130OB-I00 (NGAA) and the Severo Ochoa Excellence Accreditation to

CIC bioGUNE (CEX2021-001136-S), as well as by the Conselho Nacional de Desenvolvimento

Científico e Tecnológico (CNPq), the Coordenação de Aperfeiçoamento de Pessoal de Nível

Superior (CAPES), the Fundação de Amparo à Pesquisa do Estado de Minas Gerais

(FAPEMIG), and the Pró-Reitorias de Pesquisa e Pós-Graduação da UFMG (PRPG-UFMG).

JSA is a CNPq researcher.

FIGURE LEGENDS

Figure 1: Transmission electron microscopy images of negative stained virus-like particles

isolated from Finnish samples

A) Left, diamond-shaped virions found in a Recirculating Aquaculture System (RAS) tank

sample and in experimental aquarium samples. Right, enlarged view showing the virion interior

with at least three attachment points to the external capsid. B) Left, virions with loosely

structured capsids found in a RAS tank sample and in experimental aquarium samples. Right,

detailed view of one virion of this morphotype. C) Left, round morphotype found in an

experimental aquarium sample; right, enlarged view of a spherical-shaped virion. D) Left,

clusters of full and empty Jyvaskylavirus virions isolated from a composting soil sample. Center,

an empty capsid showing the double-layered architecture of Jyvaskylavirus; C marks the capsid,

and IM marks the putative internal membrane. Right, a fully packaged Jyvaskylavirus virion.

Figure 2: Jyvaskylavirus genomic data

A) Representative map of Jyvaskylavirus genome features. The G-C content, G-C skew, and

ORFs distribution throughout the DNA sequence are coded by different ring colors as indicated

in the color legend above. The outer blue ring represents the forward strand (positive sense)

whereas the inner blue ring represents the reverse strand (negative sense). This illustrative

genome map was constructed using CGview server (**Grant et al 2008**). B) Number of

Jyvaskylavirus proteins according to the function predicted during genome annotation; n= 388.

C) Maximum-likelihood phylogenetic tree based on DNA polymerase family B amino acid

sequences from different nucleocytoviruses. The Jyvaskylavirus sequence is indicated by a blue

circle. The alignment was performed with MUSCLE and the maximum-likelihood tree was

reconstructed using IQtree software using ultrafast bootstrap (1000 replicates). The best-fit

model selected using ModelFinder (implemented in IQtree) was VT+F+R5. Scale bar indicates

the number of substitutions per site.

Figure 3: Helium ion microscopy images of Jyvaskylavirus attachment to *A. castellanii* cells

A) *A. castellanii* cells with spined structures (acanthopodia); elongated cell at the center. B)

Details of a cell containing several viral particles on its surface (white arrowheads mark virions).

C) Icosahedrally shaped virions near cell surface invaginations appearing as craters; inset, details

of a virion likely starting the infection process through endocytosis. D) Centre, one cell

containing viruses on its surface (white arrowhead) near a burst cell displayed with its ruptured

content. Right inset, details of the burst cell content, showing several vesicles (yellow asterisks)

and viruses (white arrowheads). Left inset, clusters of virions inside extracellular vesicles

indicated by white arrows and individual virions by white arrowheads.

Figure 4: Transmission electron microscopy images of thin sections of *A. castellanii* cells

infected by Jyvaskylavirus

A) Infected cell containing viruses spread over its cytoplasm marked by C (green) and with

intracellular vesicles filled with viruses indicated by white arrows. The nucleus, whose boundary

is highlighted in semi-transparent cyan, is indicated by the letter N (cyan). B) One infected cell

with a large viral factory (VF, red) in its cytoplasm. C) View of an intracellular vesicle with

icosahedral genome-filled virions as marked by a white arrowhead; membrane related structures

nearby the vesicle interior are marked by dark-yellow triangles. NM (light cyan) and M (yellow)

mark the nuclear membrane and the mitochondria, respectively. D) Enlarged view of the region

marked by the black rectangle in C) showing possible membrane-related structure juxtaposed to

or detached from the vesicle interior; black triangles possibly indicated a forming membrane

vesicle. E) Details of virions in different stages of maturation inside the viral factory; DNA-full

particles in white arrowheads, empty particles in blue arrowheads. F) Putative stages of virion

assembly (indicated by the black arrows) as derived from the inspection of distinct particles in E

and other cellular sections. The white elliptical line highlights a capsid aperture, while the red

asterisk indicates an assembling capsid; in the remaining virion images, the capsid appears more

assembled.

Figure 5: Jyvaskylavirus cryo-EM reconstruction.

A) Left, a slab (10 central sections) of the isosurface (low-pass filtered to 15 Å) showing the

interior of the virion colour-coded by radius as from key. Right, isosurface of half the virion with

black arrows indicating the triangulation indices h=7, k=13 ($T = 309$). B) Representation of the

virion using trisymmetron geometry, with each trisymmetron color-coded differently, and a

pentasymmetron marked by a pentagonal black line. C) A schematic of a virus facet with a

trisymmetron marked with a white triangle with three icosahedral asymmetric units (IAU), one

of which is marked by a thick black line. The pseudo-hexameric morphology displayed by a

capsomer is represented by a hexagon coloured in light blue, while the true trimeric state of the

MCP is depicted as a yellow triangle. To build the IAU (excluding the penton protein), 51

pseudo-hexameric capsomers and one-third of the capsomer located at the 3-fold symmetry axis

are required, resulting in a total of 154 major capsid proteins (MCP) forming the IAU. The inset

shows a cut-through of the density along the three-fold axis of a capsomer. D) Alphafold

prediction of the MCP ORF184 shows, with very high confidence, that the fold adopted by the

ORF is a vertical double jellyroll. E) Trimeric model of the capsomers rigid-body fitted into the

original cryo-EM density and rendered in ChimeraX; left, viewed along the trimer fold axis and

on the right, viewed orthogonally to it. The three copies of the MCP are represented in cartoon

and coloured in green, light magenta, and light blue, while the corresponding density is shown in

white transparent surface.

Figure 6: Jyvaskylavirus penton and capsomer-cap proteins

A) Cut-through view of the virion cryo-EM density (binned twice from original size) at the 5-

fold axis rendered in white in ChimeraX, with regions corresponding to different proteins

forming the capsid outlined in black. B) Top, a cartoon representation of the predicted model of

the penton protein, with strands labeled BIDG/CHEF and color-coded by confidence level.

Center and bottom, penton complex fitted into the density, shown from the top and as a cut-

through, respectively. C) Top, atomic model of the trimeric cap in pink cartoon, composed of

three copies of ORF121 with a β-barrel fold inserted on top of the pseudo-hexameric capsomer,

shown in white cartoon. Bottom, cut-through view of the cap model fitted into the original

density, Gaussian filtered and rendered as semi-transparent grey in ChimeraX.

Figure 7: Jyvaskylavirus pentasymmetron protein components and trisymmetron facet

glueing protein

A) Top left, view of the cryo-EM density of Jyvaskylavirus (binned four times from the original

size and Gaussian filtered) as seen from within the virion along the five-fold icosahedral

symmetry axis. The white pentameric line marks the pentasymmetron region from below, while

the white circle highlights the density corresponding to some identified proteins. Further

densities corresponding to cementing, zippering and lattice scaffolding proteins are also visible,

with one of them coloured in dodger-blue (see also panel B). The enlarged inset (top right and

below) shows the spatial organization of the four identified ORFs and modelled using

AlphaFold3, represented as cartoon tubes and coloured as per the legend. The penton proteins are

coloured light brown, while peripentonal capsomers and capsomer-cap proteins are shown as

space-filled atoms and coloured as slate-blue and pink, respectively. Left bottom, the different

protein components fitted into the original 6.3 Å resolution Jyvaskylavirus cryo-EM density

(white semi-transparent) binned to 2.68 Å/pix and rendered in ChimeraX; MCPs and capsomer-

cap proteins have been omitted for clarity. B) Left, view of the cryo-EM density of

Jyvaskylavirus, as shown in the top left panel of A), but along the three-fold icosahedral

symmetry axis. A cementing protein, colored red, runs through the capsomers; however, the

corresponding ORF has not been identified. Densities colored dodger-blue, which glue the

capsomers across two trisymmetrons, correspond to ORF119, as shown in the large inset on the

right. In this inset, dimers of ORF119 are depicted as dodger-blue cartoons fitted into the original

cryo-EM density, Gaussian filtered and rendered as semi-transparent grey in ChimeraX. At the

bottom right, a 90-degree view shows the spatial arrangement between two capsomers (navy

blue), with the cap protein ORF121 (pink cartoon) positioned on top, and ORF119 fitted into the

density at the base of the adjacent MCPs.

REFERENCES

Abrahao J, Silva L, Silva LS, et al. Tailed giant Tupanvirus possesses the most complete

translational apparatus of the known virosphere. Nat Commun. 2018;9(1). doi:10.1038/s41467-

018-03168-1

Abramson, J., Adler, J., Dunger, J. et al. Accurate structure prediction of biomolecular

interactions with AlphaFold 3. Nature 630, 493–500 (2024). https://doi.org/10.1038/s41586-024-

07487-w

Abrescia NG, Cockburn JJ, Grimes JM, Sutton GC, Diprose JM, Butcher SJ, Fuller SD, San

Martín C, Burnett RM, Stuart DI, Bamford DH, Bamford JK. Insights into assembly from

structural analysis of bacteriophage PRD1. Nature. 2004 Nov 4;432(7013):68-74. doi:

10.1038/nature03056. PMID: 15525981.

Afonine PV, Poon BK, Read RJ, Sobolev OV, Terwilliger TC, Urzhumtsev A, Adams PD. Real-

space refinement in PHENIX for cryo-EM and crystallography. Acta Crystallogr D Struct Biol.

2018 Jun 1;74(Pt 6):531-544. doi: 10.1107/S2059798318006551. Epub 2018 May 30. PMID:

29872004; PMCID: PMC6096492.

Aherfi, S., Colson, P., La Scola, B., & Raoult, D. (2016). Giant viruses of amoebas: An update.

In Frontiers in Microbiology. https://doi.org/10.3389/fmicb.2016.00349

Almeida, G. M. F., Laanto, E., Ashrafi, R., & Sundberg, L. R. (2019). Bacteriophage adherence

to   mucus   mediates   preventive   protection   against   pathogenic   bacteria.   MBio.

https://doi.org/10.1128/mBio.01984-19

Almeida, G. M. F., Mäkelä, K., Laanto, E., Pulkkinen, J., Vielma, J., & Sundberg, L. R. (2019).

The fate of bacteriophages in recirculating aquaculture systems (RAS)—towards developing

phage therapy for RAS. Antibiotics. https://doi.org/10.3390/antibiotics8040192

Andrade, A. C. D. S. P., Arantes, T. S., Rodrigues, R. A. L., Machado, T. B., Dornas, F. P.,

Landell, M. F., Furst, C., Borges, L. G. A., Dutra, L. A. L., Almeida, G., Trindade, G. D. S.,

Bergier, I., Abrahão, W., Borges, I. A., Cortines, J. R., De Oliveira, D. B., Kroon, E. G., &

Abrahão, J. S. (2018). Ubiquitous giants: A plethora of giant viruses found in Brazil and

Antarctica. Virology Journal. https://doi.org/10.1186/s12985-018-0930-x

Andreani J, Khalil JYB, Sevvana M, Benamar S, Di Pinto F, Bitam I, Colson P, Klose T,

Rossmann MG, Raoult D, La Scola B. Pacmanvirus, a New Giant Icosahedral Virus at the

Crossroads between Asfarviridae and Faustoviruses. J Virol. 2017 Jun 26;91(14):e00212-17. doi:

10.1128/JVI.00212-17. PMID: 28446673; PMCID: PMC5487549.

Andrés G, Charro D, Matamoros T, Dillard RS, Abrescia NGA. The cryo-EM structure of

African swine fever virus unravels a unique architecture comprising two icosahedral protein

capsids and two lipoprotein membranes. J Biol Chem. 2020 Jan 3;295(1):1-12. doi:

10.1074/jbc.AC119.011196. Epub 2019 Oct 24. PMID: 31649031; PMCID: PMC6952596.

Arantes TS, Rodrigues RA, Dos Santos Silva LK, Oliveira GP, de Souza HL, Khalil JY, de

Oliveira DB, Torres AA, da Silva LL, Colson P, Kroon EG, da Fonseca FG, Bonjardim CA, La

Scola B, Abrahão JS. The Large Marseillevirus Explores Different Entry Pathways by Forming

Giant Infectious Vesicles. J Virol. 2016 May 12;90(11):5246-55. doi: 10.1128/JVI.00177-16.

Erratum in: J Virol. 2021 Aug 25;95(18):e0107121. PMID: 26984730; PMCID: PMC4934737.

Arslan, D., Legendre, M., Seltzer, V., Abergel, C., & Claverie, J. M. (2011). Distant Mimivirus

relative with a larger genome highlights the fundamental features of Megaviridae. Proceedings of

the National Academy of Sciences of the United States of America.

https://doi.org/10.1073/pnas.1110889108

Bäckström, D., Yutin, N., Jørgensen, S. L., Dharamshi, J., Homa, F., Zaremba-Niedwiedzka, K.,

Spang, A., Wolf, Y. I., Koonin, E. V., & Ettema, T. J. G. (2019). Virus genomes from deep sea

sediments expand the ocean megavirome and support independent origins of viral gigantism.

MBio. https://doi.org/10.1128/mBio.02497-18

Besemer J, Lomsadze A, Borodovsky M. GeneMarkS: a self-training method for prediction of

gene starts in microbial genomes. Implications for finding sequence motifs in regulatory regions.

Nucleic Acids Research, v 29, p 2607-2618, 2001. https://doi.org/10.1093/nar/29.12.2607

Boyer M, Madoui MA, Gimenez G, La Scola B, Raoult D: Phylogenetic and phyletic studies of

informational genes in genomes highlight existence of a 4th domain of life including giant

viruses. PLoS One 2010, 5(Suppl 12): e15530.

Boyer, M., Yutin, N., Pagnier, I., Barrassi, L., Fournous, G., Espinosa, L., Robert, C., Azza, S.,

Sun, S., Rossmann, M. G., Suzan-Monti, M., La Scola, B., Koonin, E. V., & Raoult, D. (2009).

Giant Marseillevirus highlights the role of amoebae as a melting pot in emergence of chimeric

microorganisms. Proceedings of the National Academy of Sciences of the United States of

America. https://doi.org/10.1073/pnas.0911354106

Brahim Belhaouari D, Hannat S, Rodrigues R, Aherfi S, La Scola B, Andreani J. Genome

Sequences of Two New Pandoravirus Strains Isolated from Brazil and France. Microbiol Resour

Announc. 2022 Jul 21;11(7):e0013122. doi: 10.1128/mra.00131-22. Epub 2022 Jun 22. PMID:

35731201; PMCID: PMC9302070.

Brown CJ, Johnson AK, Dunker AK, Daughdrill GW. Evolution and disorder. Curr Opin Struct

Biol. 2011 Jun;21(3):441-6. doi: 10.1016/j.sbi.2011.02.005. Epub 2011 Apr 7. PMID: 21482101;

PMCID: PMC3112239.

Bryson TD, De Ioannes P, Valencia-Sánchez MI, Henikoff JG, Talbert PB, Lee R, La Scola B,

Armache KJ, Henikoff S. A giant virus genome is densely packaged by stable nucleosomes

within virions. Mol Cell. 2022 Dec 1;82(23):4458-4470.e5. doi: 10.1016/j.molcel.2022.10.020.

Epub 2022 Nov 11. PMID: 36370708.

Burton-Smith RN, Reddy HKN, Svenda M, Abergel C, Okamoto K, Murata K. The 4.4 Å

structure of the giant Melbournevirus virion belonging to the Marseilleviridae family. bioRxiv

2021.07.14.452405; doi: https://doi.org/10.1101/2021.07.14.452405

Burton-Smith RN, Murata K. Post-acquisition super resolution for cryo-electron microscopy.

bioRxiv 2023.06.09.544325; doi: https://doi.org/10.1101/2023.06.09.544325

Chihara A, Burton-Smith RN, Kajimura N, Mitsuoka K, Okamoto K, Song C, Murata K. A novel

capsid protein network allows the characteristic internal membrane structure of Marseilleviridae

giant viruses. Sci Rep. 2022 Dec 11;12(1):21428. doi: 10.1038/s41598-022-24651-2. PMID:

36504202; PMCID: PMC9742146.

Colson P, La Scola B, Levasseur A, Caetano-Anollés G, Raoult D. Mimivirus: Leading the way

in the discovery of giant viruses of amoebae. Nat Rev Microbiol. 2017;15(4):243-254.

doi:10.1038/nrmicro.2016.197

Darling ACE, Mau B, Blattner FR, Perna NT. Mauve: multiple alignment of conserved genomic

sequence with rearrangements. Genome Research 14:1394–1403. 2004

https://doi.org/10.1101/gr.22897 04

Edgar R C. MUSCLE: multiple sequence alignment with high accuracy and high throughput.

Nucleic Acids Research, v. 32, n. 5, p. 1792-1797, 2004. https://doi.org/10.1093/nar/gkh340

Emsley P, Cowtan K. Coot: model-building tools for molecular graphics. Acta Crystallogr D

Biol Crystallogr. 2004 Dec;60(Pt 12 Pt 1):2126-32. doi: 10.1107/S0907444904019158. Epub

2004 Nov 26. PMID: 15572765.

Fischer MG, Mersdorf U, Blanchard JL. Amazing structural diversity of giant virus-like particles

in forest soil. bioRxiv 2023.06.30.546935; doi: https://doi.org/10.1101/2023.06.30.546935

Gann ER, Xian Y, Abraham PE, Hettich RL, Reynolds TB, Xiao C, Wilhelm SW. Structural and

Proteomic Studies of the Aureococcus anophagefferens Virus Demonstrate a Global Distribution

of Virus-Encoded Carbohydrate Processing. Front Microbiol. 2020 Sep 8;11:2047. doi:

10.3389/fmicb.2020.02047. PMID: 33013751; PMCID: PMC7507832.

Gil-Carton D, Jaakkola ST, Charro D, Peralta B, Castaño-Díez D, Oksanen HM, Bamford DH,

Abrescia NGA. Insight into the Assembly of Viruses with Vertical Single β-barrel Major Capsid

Proteins. Structure. 2015 Oct 6;23(10):1866-1877. doi: 10.1016/j.str.2015.07.015. Epub 2015

Aug 27. PMID: 26320579.

Grant JR, Stothard P. The CGView Server: a comparative genomics tool for circular genomes.

Nucleic Acids Research ,36, 181-184. 2008. doi: 10.1093/nar/gkn179

Hylling, O., Carstens, A. B., Kot, W., Hansen, M., Neve, H., Franz, C. M. A. P., Johansen, A., Ellegaard-Jensen, L., & Hansen, L. H. (2020). Two novel bacteriophage genera from a groundwater reservoir highlight subsurface environments as underexplored biotopes in bacteriophage ecology. Scientific Reports. https://doi.org/10.1038/s41598-020-68389-1

Iyer LM, Aravind L, Koonin E V. Common Origin of Four Diverse Families of Large Eukaryotic DNA Viruses. J Virol. 2001;75(23):11720-11734. doi:10.1128/JVI.75.23.11720-11734.2001

Jamali K, Käll L, Zhang R, Brown A, Kimanius D, Scheres SHW. Automated model building and protein identification in cryo-EM maps. Nature. 2024 Apr;628(8007):450-457. doi: 10.1038/s41586-024-07215-4. Epub 2024 Feb 26. PMID: 38408488; PMCID: PMC11006616.

Jensen, T., Barnes, W. G., & Meyers, D. (1970). Axenic cultivation of large populations of Acanthamoeba castellanii (JBM). The Journal of Parasitology. https://doi.org/10.2307/3277503

Johannessen TV, Bratbak G, Larsen A, Ogata H, Egge ES, Edvardsen B, Eikrem W, Sandaa RA. Characterisation of three novel giant viruses reveals huge diversity among viruses infecting Prymnesiales (Haptophyta). Virology. 2015 Feb;476:180-188. doi: 10.1016/j.virol.2014.12.014. Epub 2014 Dec 26. PMID: 25546253.

Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M, Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW, Kavukcuoglu K, Kohli P, Hassabis D. Highly accurate protein structure prediction with AlphaFold. Nature. 2021 Aug;596(7873):583-589. doi: 10.1038/s41586-021-03819-2. Epub 2021 Jul 15. PMID: 34265844; PMCID: PMC8371605.

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder: Fast

model selection for accurate phylogenetic estimates. Nat. Methods, 14:587-589 (2017).

https://doi.org/10.1038/nmeth.4285

Klose T, Reteno DG, Benamar S, Hollerbach A, Colson P, La Scola B, Rossmann MG. Structure

of faustovirus, a large dsDNA virus. Proc Natl Acad Sci U S A. 2016 May 31;113(22):6206-11.

doi: 10.1073/pnas.1523999113. Epub 2016 May 16. PMID: 27185929; PMCID: PMC4896704.

Kördel, M., Svenda, M., Reddy, H. K. N., Fogelqvist, E., Arsana, K. G. Y., Hamawandi, B.,

Toprak, M. S., Hertz, H. M., & Sellberg, J. A. (2021). Quantitative conversion of biomass in

giant DNA virus infection. Scientific Reports. https://doi.org/10.1038/s41598-021-83547-9

Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular Evolutionary Genetics

Analysis across Computing Platforms. Molecular Biology and Evolution, v. 35, n. 6, p. 1547-

1549, 2018.   doi: 10.1093/molbev/msy096

La Scola B, Audic S, Robert C, Jungang L, de Lamballerie X, Drancourt M, Birtles R, Claverie

JM, Raoult D: A giant virus in amoebae. Science 2003, 299: 2033.

Laanto E, Hoikkala V, Ravantti J, Sundberg LR. Long-term genomic coevolution of host-

parasite interaction in the natural environment. Nat Commun. 2017;8(1). doi:10.1038/s41467-

017-00158-7

Laanto E, Mäntynen S, De Colibus L, et al. Virus found in a boreal lake links ssDNA and

dsDNA viruses. Proc Natl Acad Sci. 2017:201703834. doi:10.1073/pnas.1703834114

Laslett D, Bjorn C. ARAGORN, a program to detect tRNA genes and tmRNA genes in

nucleotide sequences. Nucleic Acids Research, v 32, n 1, p 11-16, 2004.

https://doi.org/10.1093/nar/gkh152

Legendre M, Lartigue A, Bertaux L, et al. In-depth study of Mollivirus sibericum, a new 30,000-

y-old giant virus infecting Acanthamoeba. PNAS. 2015;112(38):5327-5335.

doi:10.1073/pnas.1510795112

Leppänen M, Sundberg L-R, Laanto E, de Freitas Almeida GM, Papponen P, Maasilta IJ.

Imaging Bacterial Colonies and Phage-Bacterium Interaction at Sub-Nanometer Resolution

Using Helium-Ion Microscopy. Adv Biosyst. 2017:1700070. doi:10.1002/adbi.201700070

Letunic I, Bork P. Interactive Tree Of Life (iTOL) v5: an online tool for phylogenetic tree

display and annotation. Nucleic Acids Research, v 49, p 293–296, 2021.

https://doi.org/10.1093/nar/gkab301

Liu H, Jin L, Koh SB, Atanasov I, Schein S, Wu L, Zhou ZH. Atomic structure of human

adenovirus by cryo-EM reveals interactions among protein networks. Science. 2010 Aug

27;329(5995):1038-43. doi: 10.1126/science.1187433. PMID: 20798312; PMCID:

PMC3412078.

Meng EC, Goddard TD, Pettersen EF, Couch GS, Pearson ZJ, Morris JH, Ferrin TE. UCSF

ChimeraX: Tools for structure building and analysis. Protein Sci. 2023 Nov;32(11):e4792. doi:

10.1002/pro.4792. PMID: 37774136; PMCID: PMC10588335.

Moniruzzaman, M., Martinez-Gutierrez, C.A., Weinheimer, A.R. et al. Dynamic genome

evolution and complex virocell metabolism of globally-distributed giant viruses. Nat Commun

11, 1710 (2020). https://doi.org/10.1038/s41467-020-15507-2

Nguyen, L T et al. IQ-TREE: a fast and effective stochastic algorithm for estimating maximum-

likelihood phylogenies. Molecular Biology and Evolution, v. 32, n. 1, p. 268-274, 2014.

Okamoto K, Miyazaki N, Reddy HKN, Hantke MF, Maia FRNC, Larsson DSD, Abergel C,

Claverie JM, Hajdu J, Murata K, Svenda M. Cryo-EM structure of a Marseilleviridae virus

particle reveals a large internal microassembly. Virology. 2018 Mar;516:239-245. doi:

10.1016/j.virol.2018.01.021. PMID: 29407382.

Perini, L., Sipes, K., Zervas, A. et al. Giant viral signatures on the Greenland ice sheet.

Microbiome 12, 91 (2024). https://doi.org/10.1186/s40168-024-01796-y

Pettersen EF, Goddard TD, Huang CC, Couch GS, Greenblatt DM, Meng EC, Ferrin TE. UCSF

Chimera--a visualization system for exploratory research and analysis. J Comput Chem. 2004

Oct;25(13):1605-12. doi: 10.1002/jcc.20084. PMID: 15264254.

Philippe, N., Legendre, M., Doutre, G., Couté, Y., Poirot, O., Lescot, M., Arslan, D., Seltzer, V.,

Bertaux, L., Bruley, C., Garin, J., Claverie, J. M., & Abergel, C. (2013). Pandoraviruses:

Amoeba viruses with genomes up to 2.5 Mb reaching that of parasitic eukaryotes. Science.

https://doi.org/10.1126/science.1239181

Queiroz VF, Rodrigues RAL, de Miranda Boratto PV, La Scola B, Andreani J, Abrahão JS.

Amoebae: Hiding in Plain Sight: Unappreciated Hosts for the Very Large Viruses. Annu Rev

Virol. 2022 Sep 29;9(1):79-98. doi: 10.1146/annurev-virology-100520-125832. Epub 2022 Jun

2. PMID: 35655338.

Ravantti JJ, Martinez-Castillo A, Abrescia NGA. Superimposition of Viral Protein Structures: A

Means to Decipher the Phylogenies of Viruses. Viruses. 2020 Oct 9;12(10):1146. doi:

10.3390/v12101146. PMID: 33050291; PMCID: PMC7600307.

Reed LJ, Muench H. A simple method of estimating fifty-percent endpoints. Am J Hyg.

1938;27:493–497.

Rohou A, Grigorieff N. CTFFIND4: Fast and accurate defocus estimation from electron

micrographs. J Struct Biol. 2015 Nov;192(2):216-21. doi: 10.1016/j.jsb.2015.08.008. Epub 2015

Aug 13. PMID: 26278980; PMCID: PMC6760662.

Runtuvuori-Salmela A, HMT Kunttu, E Laanto, GMF Almeida, K Mäkelä, M Middelboe, L-R

Sundberg. Prevalence of genetically similar Flavobacterium columnare phages across

aquaculture environments reveals a strong potential for pathogen control. bioRxiv

2020.09.23.309583; doi: https://doi.org/10.1101/2020.09.23.309583

Santos-Pérez I, Charro D, Gil-Carton D, Azkargorta M, Elortza F, Bamford DH, Oksanen HM,

Abrescia NGA. Structural basis for assembly of vertical single β-barrel viruses. Nat Commun.

2019 Mar 12;10(1):1184. doi: 10.1038/s41467-019-08927-2. PMID: 30862777; PMCID:

PMC6414509.

Sahmi-Bounsiar, D., Rolland, C., Aherfi, S., Boudjemaa, H., Levasseur, A., La Scola, B., &

Colson, P. (2021). Marseilleviruses: An Update in 2021. In Frontiers in Microbiology.

https://doi.org/10.3389/fmicb.2021.648731

Sandaa RA, Heldal M, Castberg T, Thyrhaug R, Bratbak G. Isolation and characterization of two

viruses with large genome size infecting Chrysochromulina ericina (Prymnesiophyceae) and

Pyramimonas orientalis (Prasinophyceae). Virology. 2001 Nov 25;290(2):272-80. doi:

10.1006/viro.2001.1161. PMID: 11883191.

Santos, M. A. (1991). An improved method for the small scale preparation of bacteriophage

DNA based on phage precipitation by zinc chloride. Nucleic Acids Research.

https://doi.org/10.1093/nar/19.19.5442

Scheres SH. A Bayesian view on cryo-EM structure determination. J Mol Biol. 2012 Jan

13;415(2):406-18. doi: 10.1016/j.jmb.2011.11.010. Epub 2011 Nov 12. PMID: 22100448;

PMCID: PMC3314964.

Schulz, F., Roux, S., Paez-Espino, D. et al. Giant virus diversity and host interactions through

global metagenomics. Nature 578, 432–436 (2020). https://doi.org/10.1038/s41586-020-1957-x

Shao Q, Agarkova IV, Noel EA, Dunigan DD, Liu Y, Wang A, Guo M, Xie L, Zhao X,

Rossmann MG, Van Etten JL, Klose T, Fang Q. Near-atomic, non-icosahedrally averaged

structure of giant virus Paramecium bursaria chlorella virus 1. Nat Commun. 2022 Oct

29;13(1):6476. doi: 10.1038/s41467-022-34218-4. PMID: 36309542; PMCID: PMC9617893.

Simmonds P, Adriaenssens EM, Zerbini FM, Abrescia NGA, Aiewsakun P, Alfenas-Zerbini P,

Bao Y, Barylski J, Drosten C, Duffy S, Duprex WP, Dutilh BE, Elena SF, García ML, Junglen S,

Katzourakis A, Koonin EV, Krupovic M, Kuhn JH, Lambert AJ, Lefkowitz EJ, Łobocka M,

Lood C, Mahony J, Meier-Kolthoff JP, Mushegian AR, Oksanen HM, Poranen MM, Reyes-

Muñoz A, Robertson DL, Roux S, Rubino L, Sabanadzovic S, Siddell S, Skern T, Smith DB,

Sullivan MB, Suzuki N, Turner D, Van Doorslaer K, Vandamme AM, Varsani A, Vasilakis N.

Four principles to establish a universal virus taxonomy. PLoS Biol. 2023 Feb

13;21(2):e3001922. doi: 10.1371/journal.pbio.3001922. PMID: 36780432; PMCID:

PMC9925010.

Sinkovits RS, Baker TS. A tale of two symmetrons: rules for construction of icosahedral capsids

from trisymmetrons and pentasymmetrons. J Struct Biol. 2010 Apr;170(1):109-16. doi:

10.1016/j.jsb.2009.12.003. Epub 2009 Dec 4. PMID: 19963064; PMCID: PMC3834589.

Stuart DI, Subramaniam S, Abrescia NG. The democratization of cryo-EM. Nat Methods. 2016

Jul 28;13(8):607-8. doi: 10.1038/nmeth.3946. PMID: 27467721; PMCID: PMC10720754.

Thomas, V., Herrera-Rimann, K., Blanc, D. S., & Greub, G. (2006). Biodiversity of amoebae

and amoeba-resisting bacteria in a hospital water network. Applied and Environmental

Microbiology. https://doi.org/10.1128/AEM.72.4.2428-2438.2006

Wagner T, Merino F, Stabrin M, Moriya T, Antoni C, Apelbaum A, Hagel P, Sitsel O, Raisch T,

Prumbaum D, Quentin D, Roderer D, Tacke S, Siebolds B, Schubert E, Shaikh TR, Lill P,

Gatsogiannis C, Raunser S. SPHIRE-crYOLO is a fast and accurate fully automated particle

picker for cryo-EM. Commun Biol. 2019 Jun 19;2:218. doi: 10.1038/s42003-019-0437-z. PMID:

31240256; PMCID: PMC6584505.

Walker, P. J., Siddell, S. G., Lefkowitz, E. J., Mushegian, A. R., Dempsey, D. M., Dutilh, B. E.,

Harrach, B., Harrison, R. L., Hendrickson, R. C., Junglen, S., Knowles, N. J., Kropinski, A. M.,

Krupovic, M., Kuhn, J. H., Nibert, M., Rubino, L., Sabanadzovic, S., Simmonds, P., Varsani, A.,

… Davison, A. J. (2019). Changes to virus taxonomy and the International Code of Virus

Classification and Nomenclature ratified by the International Committee on Taxonomy of

Viruses (2019). Archives of Virology. https://doi.org/10.1007/s00705-019-04306-w

Wang N, Zhao D, Wang J, Zhang Y, Wang M, Gao Y, Li F, Wang J, Bu Z, Rao Z, Wang X.

Architecture of African swine fever virus and implications for viral assembly. Science. 2019 Nov

1;366(6465):640-644. doi: 10.1126/science.aaz1439. Epub 2019 Oct 17. PMID: 31624094.

Watanabe R, Song C, Kayama Y, Takemura M, Murata K. Particle Morphology of Medusavirus

Inside and Outside the Cells Reveals a New Maturation Process of Giant Viruses. J Virol. 2022

Apr 13;96(7):e0185321. doi: 10.1128/jvi.01853-21. Epub 2022 Mar 17. PMID: 35297671;

PMCID: PMC9006890.

Xiao C, Fischer MG, Bolotaulo DM, Ulloa-Rondeau N, Avila GA, Suttle CA. Cryo-EM

reconstruction of the Cafeteria roenbergensis virus capsid suggests novel assembly pathway for

giant viruses. Sci Rep. 2017 Jul 14;7(1):5484. doi: 10.1038/s41598-017-05824-w. PMID:

28710447; PMCID: PMC5511168.

Yan X, Chipman PR, Castberg T, Bratbak G, Baker TS. The marine algal virus PpV01 has an

icosahedral capsid with T=219 quasisymmetry. J Virol. 2005 Jul;79(14):9236-43. doi:

10.1128/JVI.79.14.9236-9243.2005. PMID: 15994818; PMCID: PMC1168743.

Zheng SQ, Palovcak E, Armache JP, Verba KA, Cheng Y, Agard DA. MotionCor2: anisotropic

correction of beam-induced motion for improved cryo-electron microscopy. Nat Methods. 2017

Apr;14(4):331-332. doi: 10.1038/nmeth.4193. Epub 2017 Feb 27. PMID: 28250466; PMCID:

PMC5494038.

Zhu D, Wang X, Fang Q, Van Etten JL, Rossmann MG, Rao Z, Zhang X. Pushing the resolution

limit by correcting the Ewald sphere effect in single-particle Cryo-EM reconstructions. Nat

Commun. 2018 Apr 19;9(1):1552. doi: 10.1038/s41467-018-04051-9. PMID: 29674632;

PMCID: PMC5908801.

SUPPLEMENTARY MATERIAL LIST

Supplementary text (docx): Preliminary data about isolation of Finnish giant viruses using a

locally isolated amoeba host; Jyvaskylavirus growth, chloroform sensitivity and stability.

Supplementary Figure 1: Details of our preliminary screening for Finnish giant viruses' effort.

Supplementary Figure 2: Jyvaskylavirus growth curve, sensitivity to chloroform and stability.

Supplementary Figure 3: Phylogenetic tree reconstructed using Major Capsid Protein (MCP)

amino acid sequences belonging to viruses from *Marseilleviridae* family.

Supplementary Figure 4: Fourier Shell Correlation curve of the whole Jyvaskylavirus.

Supplementary Figure 5: Schematic workflow for ORF identification and model prediction

Supplementary Figure 6: Fitting of individual into the corresponding densities of the 5-fold

block Melbourne virus map

Supplementary Figure 7: Fitting of all Jyvaskylavirus protein models into the 5-fold block

Melbourne virus map

Supplementary Figure 8: Fitting of Jyvaskylavirus ORF119 protein model into the 2-fold block

Melbourne virus map

Supplementary Table 1 (xls): Details of the samples used and isolation results.

Supplementary Table 2 (docx): Grid vitrification parameters and cryo-EM data collection.

Supplementary Table 3 (docx): Identified Melbournevirus ORFs through Jyvaskylavirus ORFs.

Almeida et al - Figure 1

Almeida et al - Figure 2

Almeida et al - Figure 3

Almeida et al - Figure 4

A

k = 13
h = 7
T = 309

Radius (Å)   850 930 1100 1150 1200 1250 1300 1350 1400

B

C

Trisymmetron = 136 pseudo-hexagonal capsomers

IAU = 154 vertical double jellyroll MCPs (ORF184)
        + 1 single jellyroll (penton)

D

Alphafold2
predicted MCP
(ORF184)

Nt

Ct

98.8 %

28.7 %

E

~180°

~90°

Almeida et al - Figure 5

Almeida et al - Figure 6

A

5-fold

ORF119 ORF036
ORF097 ORF153

90°

B

90°

Almeida et al - Figure 7