Critical functions and key interactions mediated by the RNase E scaffolding domain in

Pseudomonas aeruginosa

Sandra Amandine Marie Geslain, Stéphane Hausmann, Johan Geiser, George Edward Allen, Diego

Gonzalez and Martina Valentini*

Department of Microbiology and Molecular Medicine, CMU, Faculty of Medicine, University of

Geneva, Geneva, Switzerland.

* To whom correspondence should be addressed. e-mail: Martina.Valentini@unige.ch

Abstract

The RNA degradosome is a bacterial multi-protein complex mediating mRNA processing and

degradation. In Pseudomonadota, this complex assembles on the C-terminal domain (CTD) of RNase

E through short linear motifs (SLiMs) that determine its composition and functionality. In the human

pathogen *Pseudomonas aeruginosa*, the RNase E CTD exhibits limited similarity to that of model

organisms, impeding our understanding of RNA metabolic processes in this bacterium. Our study

systematically maps the interactions mediated by the *P. aeruginosa* RNase E CTD and highlights its

critical role in transcript regulation and cellular functions. We identified the SLiMs crucial for

membrane attachment, RNA binding and complex clustering, as well as for direct binding to the core

components PNPase and RhlB. Transcriptome analyses of RNase E CTD mutants revealed altered

expression of genes involved in quorum sensing, type III secretion, and amino acid metabolism.

Additionally, we show that the mutants are impaired in cold adaptation, pH response, and virulence in

an infection model. Overall, this work establishes the essential role of the RNA degradosome in driving

bacterial adaptability and pathogenicity.

Author summary

Bacteria must rapidly adapt to changing environments, whether facing temperature shifts, nutrient

scarcity, or antibiotic exposure. A key mechanism enabling this adaptability is the regulation of mRNA

levels—the molecular blueprints for protein production. This process is governed by the RNA

degradosome, a multi-protein complex that processes and degrades RNA to control gene expression.

Although the RNA degradosome core function is conserved across bacteria, its composition and

organization differ significantly between species, reflecting diverse lifestyles and environmental

challenges each bacterium encounter.

In this study, we investigated the RNA degradosome in *Pseudomonas aeruginosa*, a bacterial pathogen

causing difficult-to-treat infections in humans. We identified its components and mapped regions within

the complex essential for RNA binding, membrane attachment, and spatial organization. Disrupting

these regions compromised *P. aeruginosa* ability to survive in cold conditions, respond to stress, and

establish an infection. Through this work, we uncovered unique features of the *P. aeruginosa* RNA

degradosome that distinguish it from those of other species, emphasizing the RNA degradosome critical

role in bacterial adaptability and highlight it as a promising target for therapies against *P. aeruginosa*

infections.

Introduction

In bacteria, the processing and degradation of mRNAs is primarily orchestrated by the RNA degradosome, a multi-enzyme complex that includes ribonucleases, RNA helicases, and various other proteins [1-5]. The RNA degradosome plays a crucial role in enabling bacteria to rapidly adapt to changing environmental conditions by swiftly modifying mRNA levels and broadly regulating gene expression [6, 7].

Most of our knowledge on the RNA degradosome derives from studies in *Escherichia coli*, where the RNase E endonuclease serves as the central component [8-11]. While the catalytic activity of *E. coli* RNase E resides in its N-terminal domain (NTD, amino acids 1-529), the C-terminal domain (CTD, amino acids 530-1061), or scaffolding domain, is intrinsically disordered and contains short linear motifs (SLiMs) of 10 to 40 amino acids to which the other RNA degradosome components bind [11, 12]. Core *E. coli* RNA degradosome components, namely the polyribonucleotide phosphorylase PNPase [13], the RNA helicase RhlB [14], and the glycolytic enzyme enolase [15], interact with RNase E in semi-stoichiometric ratios via the PBS (PNPase binding site), HBS (helicase binding site) and EBS (enolase binding cite) SLiM, respectively [16, 17]. The *E. coli* RNase E CTD also includes an amphipathic helix, called membrane targeting sequence (MTS), which tethers the entire complex to the inner membrane, and at least two RNA-binding SLiMs, AR1 (named also RBD) and AR2, which mediate contact with translating ribosomes, small non-coding RNAs and mRNAs [12, 18-21]. Additional proteins, such as RraA [22, 23], RraB [24], CsdA [25], Ppk [26], MinD [27, 28], and Hfq [29], can interact with *E. coli* RNase E in sub-stochiometric ratios and/or under non-standard conditions of growth (reviewed in [30]). Of importance, the RNase E NTD is essential for *E. coli* growth and rRNA processing [31], whereas deletion of the CTD is not lethal and alters mRNA levels without affecting rRNAs [19, 32-34]. This highlights the dual role of RNase E, with the NTD being crucial for overall cell maintenance, and the CTD regulating specific RNA substrates through the assembly of the RNA degradosome.

RNase E orthologs are present in approximately 49% of sequenced bacterial genomes, and nearly all Pseudomonadota (formerly called Proteobacteria) encode at least one RNase E ortholog, with the notable exception of most *Campylobacter* and *Helicobacter* species [5, 30]. The RNase E NTD is

remarkably conserved across most species, while the CTD varies considerably in sequence, length, and

SLiMs composition, even between species belonging to the same order [30]. SLiMs variability is

thought to drive the diversity of RNase E interacting partners, and therefore RNA degradosome

composition, among bacterial species. For instance, in *Caulobacter crescentus,* RNase E interacts with

PNPase, RhlB, RNase D, and the Krebs cycle enzyme aconitase [35, 36]. In *Anabaena* sp. strain PCC

7120, RNase E partners with PNPase, enolase, the RNA helicase CrhB, and RNase II [37-39], while in

Pseudomonas syringae LZ4W, only RNase R and the RNA helicase RhlE were found to co-purify with

RNase E in pull-down assays [40]. Similarly, in *Rhodobacter capsulatus*, purification of the RNA

degradosome complex via glycerol-gradient centrifugation of cell extracts revealed the presence of

RNase E, two unnamed RNA helicases, the Rho transcription termination factor, and two unidentified

proteins of 47 and 36 kDa, with only low amounts of PNPase present in the purified fractions [41].

In these examples, the SLiMs mediating protein-protein interactions were not always characterized.

Moreover, even when the interaction of RNase E with a certain protein partner is conserved across

several species, such as the RNase E-PNPase and RNase E-RhlB interactions, the SLiMs involved in

the binding show little sequence identity [30, 36, 37, 42-44]. Therefore, predicting RNA degradosome

composition based solely on the RNase E CTD sequence does not seem possible. Experimental

characterization of the SLIMs across different bacterial species and their impact on the RNA

degradosome composition and function is essential, providing insights into novel regulatory

mechanisms that enable bacteria to thrive under stress and adapt to diverse environments.

Little is known about the RNA degradosome composition of the opportunistic human pathogen

Pseudomonas aeruginosa, which is a leading cause of nosocomial infections worldwide largely due to

its remarkable adaptability to different environments and resistance to antibiotics [45-53]. A recent

study has revealed that *P. aeruginosa* RNase E co-elutes with PNPase, the PA0428 RNA helicase, and

several other proteins in pull-down assays, although their interaction with RNase E has neither been

confirmed nor mapped [54]. Our laboratory has characterized the PA0428 RNA helicase, named RhlE2,

confirming an RNA-dependent interaction with RNase E and identifying the RhlE2 C-terminal region

as necessary and sufficient for this binding [55, 56]. However, the specific RNase E interaction site

remained unmapped.

In this study, we characterize the SLiMs within the CTD of *P. aeruginosa* RNase E, identifying those

involved in RNA binding (AR1, AR4, and REER-repeats) as well as those mediating interactions with

core RNA degradosome components, which we determine to be PNPase and the RNA helicase RhlB

(NDPR and AR1, respectively). We also show that RhlE2 interacts with the RNase E NTD in pull-down

assays. Protein-protein interactions were probed with multiple experimental approaches, including an

analysis of the subcellular localization dynamics of the complex. Additionally, we performed

transcriptomic and phenotypic analyses of RNase E CTD mutant strains, revealing the critical role of

RNA degradosome complex scaffolding in bacterial virulence, adaptation to cold temperatures and to

pH fluctuations.

The ability of *P. aeruginosa* to cause disease is intrinsically linked to a sophisticated gene regulatory

network that facilitates bacterial survival and virulence. Within this network, the role of post-

transcriptional gene regulation remains only partially explored [57, 58]. By uncovering the composition

and functional dynamics of the RNA degradosome in *P. aeruginosa*, our study not only advances the

understanding of the molecular mechanisms driving bacterial environmental adaptation but also reveals

potential novel drug targets, which could be crucial in the fight against antibiotic-resistant infections.

Results

The P. *aeruginosa* RNase E harbours several uncharacterised SLiMs conserved within the

Pseudomonadales

P. aeruginosa RNase E (Pa RNase E) consists of an NTD (amino acids 1-529) that shares high sequence

identity (70%) and similarity (90%) with the NTD of *E. coli* RNase E (Ec RNase E), and a CTD (amino

acids 530-1057) that shows only 25% identity and 37% similarity with the CTD of Ec RNase E (Fig.

S1). Previous work by Aït-Bara et al. (2015) identified seven different putative SLiMs on the Pa RNase

E CTD (MTS, AR4, REE, M29, AR1, M20 and NDPR) [30], which are shown in comparison to the Ec

RNase E in Fig. S2A. Of note, the MTS is the only SLiM of Pa RNase E CTD with a propensity to form

a secondary structure while in Ec RNase E, the helicase, enolase, and PNPase binding sites also exhibit

this feature (Fig. 1 and [12]).

To refine the prediction of Pa RNase E SLiMs, we performed a multiple sequence alignment of 5000

Pseudomonas RNase E orthologs, identifying conserved motifs (Fig. S3). Several highly conserved

CTD regions overlap with the SLiMs previously identified, with the notable presence of additional

conserved residues located immediately adjacent to these SLiMs (Fig. S2A, Fig. S3). Based on these

data, we updated the SLiMs boundaries to include these conserved residues (Fig. 1B, former SLiMs

boundaries in Fig. S2A). Upon manual inspection of the CTD sequence, we also observed a large

duplicated sequence (residues 635-667 and 682-714) within the CTD region containing the REE and

motif 29 SLiMs. Since motif 29 is defined by the conserved motif REERQPR, we merged the two

SLiMs and renamed the entire segment (residues 614-729) as the "REER-repeats" region (Fig. 1B). The

number of REER repeats varies across *Pseudomonas* species: *P. aeruginosa* strains typically harbour

13 RE[ED]R repeats, whereas the *P. fluorescens*, *P. protegens*, *P. stutzeri*, and *P. syringae* environmental

species show a distribution of repeats peaking around 8-10. The exception is *P. putida*, which exhibits

a high frequency of 20-23 RE[ED]R repeats (Fig. S2B). This variability suggests a high frequency of

indels occurring within the REER-repeats region.

Overall, our in-silico analyses of Pa RNase E CTD sequence refined the number and length of the Pa

RNase E SLiMs, highlighting a potential functional significance of the REER-repeats region in

Pseudomonas species, and suggesting differences in RNA degradosome complex dynamics between *P.*

aeruginosa and *E. coli*.

The REER-repeats, AR1 and AR4 SLiMs mediate RNase E CTD RNA binding and foci formation

Previous works in *E. coli* identified arginine-rich SLiMs as RNA-binding motifs [21, 59]. We therefore

hypothesized that the *P. aeruginosa* AR1 and AR4 SLiMs, along with the newly identified REER-repeat

region, might be involved in RNA binding. To test this hypothesis, we conducted electrophoretic

mobility shift assays (EMSA) to examine the binding of purified RNase E CTD proteins to an *in vitro*

synthesized RNA derived from the *malEF* intergenic region (for details, see methods). We tested wild-

type (WT) CTD and mutated variants where arginine residues in AR1, AR4, or the REER-repeats were

replaced with alanine (named AR1mut, AR4mut, and REERmut, respectively; see Table S4 for a

description of the mutations). Qualitative analysis of the EMSA showed that the WT, AR1mut, and

AR4mut variants bound RNA with similar efficiency, as indicated by comparable gel shift patterns (Fig.

S4A, Fig. 2A). However, the REERmut variant exhibited slightly reduced RNA binding, evidenced by

a delayed shift as compared to the former proteins (Fig. S4B and 2A). The combination of AR1 and

AR4 mutations resulted in an impaired RNA binding which was further exacerbated when the REER

mutation was also introduced: no observable shift in the RNA was detected when incubated with the

protein variant carrying mutations in all three SLiMs, even at an RNA-to-protein ratio of 1:80 (Fig. 2A).

Together, these results highlight the functional redundancy of the AR1, AR4, and REER-repeat SLiMs

in RNA binding, as a complete defect in RNA binding is observed only when all these SLiMs are

simultaneously disrupted.

One notable characteristic of the RNA degradosome is its tendency to form highly dynamic subcellular

foci, also referred to as "clusters," "puncta," or "helical structures" [28, 60]. These foci rapidly

disassemble upon rifampicin treatment, suggesting their formation is RNA-dependent [60, 61]. In *E.*

coli, these foci are membrane-anchored via the MTS SLiM, whereas in *C. crescentus* they remain

cytoplasmic, driven by liquid-liquid phase separation [61, 62]. To investigate the subcellular localisation

of Pa RNase E, we constructed *P. aeruginosa* PAO1 strains expressing msfGFP-tagged Pa RNase E, Pa

RNase E$^{1-588}$, or Pa RNase E$^{1-529}$ from the native *rne* locus. The two RNase E truncations were designed

based on our in-silico analysis of Pa RNase E SLiMs, assuming that they would lack the capacity to

assemble the RNA degradosome. The key difference between RNase E$^{1-588}$ and RNase E$^{1-529}$ lies in the

presence of the MTS (Fig. 1B). Visualization using epifluorescence microscopy revealed that RNase E-

msfGFP forms foci concentrated near the cell edge (Fig 2B). Both CTD truncation mutants showed

impairment in foci assembly, but while the RNase E$^{1-588}$-msfGFP protein localises uniformly around the

edge of the bacteria, the RNase E$^{1-529}$-msfGFP mutant protein additionally fails to localise near the

membrane, consistent with the role of the MTS in membrane anchoring (Fig. 2B-C).

Exposure of Pa RNase E-msfGFP expressing cells to high levels of rifampicin (100 µg/ml) led to the

dissipation of these foci, resulting in a smoother distribution of the signal along the cell periphery, which

phenocopies the localization of the RNase E$^{1-588}$-msfGFP strain (Fig. 2D). Quantitative analysis

confirmed a significant reduction in signal intensity variance upon rifampicin treatment as compared to

the control (DMSO-only), indicating foci dissipation, while the average signal intensity remained

constant, ruling out RNase E- msfGFP proteolysis (Fig. 2E-F, Fig. S4F).

To further verify that Pa RNase E-msfGFP foci formation depends on CTD-mediated RNA binding, we

constructed *P. aeruginosa* strains with chromosomally msfGFP-tagged RNase E variants mutated in

AR1, AR4, and REER-repeats SLiMs, individually and in combination, as described previously for the

EMSA. Visualisation of these variants showed a clear correlation between CTD-mediated RNA binding

and foci formation. Specifically, the RNase E AR1+AR4mut-msfGFP showed less distinct foci and

lower signal variance compared to RNase E-msfGFP and single AR1 or AR4 mutants (Fig. 2B-C). The

RNase E REERmut-msfGFP also had diffuse foci and lower variance, while the RNase E

AR1+AR4+REERmut-msfGFP, with abolished RNA binding, displayed the most severe disruption,

with nearly complete loss of foci and a very narrow signal intensity distribution (Fig. 2B-C and Fig.

S4F). Similar patterns were observed with four other CTD truncations containing different

combinations of RNA-binding SLiMs, ruling out the possibility that this foci dissipation is caused by

the changes in the charge of the CTD conferred by the Arg to Ala mutations (Fig. S4E).

Of note, we noticed an increased signal intensity in all the msfGFP-tagged RNase E mutants tested as

compared to the RNase E-msfGFP (Fig. S4C). Since all constructs are expressed at the native locus,

this could likely reflect higher RNase E protein levels, which suggests that CTD-mediated RNA binding

might be necessary for RNase E autoregulation [63]. To confirm this, we performed a Western Blot on

strains expressing several Strep-tagged RNase E mutants and could indeed observe that RNase E protein

levels are increased upon deletion of the CTD or introduction of AR1+AR4+REERmut (Fig. S4D).

Overall, we conclude that in *P. aeruginosa*, the organisation of the RNA degradosome complex into

visualisable foci is dependent on the RNase E CTD-mediated RNA binding and suggest that this

interaction not only organizes the localisation of the RNA degradosome but could also influence the

regulatory activity of RNase E.

Identification of proteins binding to RNase E scaffold domain

In *P. aeruginosa*, the RNA degradosome was proposed to include the RhlE2 RNA helicase (PA0428),

PNPase, and other associated proteins [54, 55]. However, the direct nature of these interactions and the

specific interaction sites have not been demonstrated nor mapped. Additionally, RNase R was also

suggested to be a *P. aeruginosa* RNA degradosome component, as it is in *P. syringae* LZ4W [30, 40].

To test these hypotheses and identify Pa RNase E CTD interacting proteins, we performed a pull-down

assay using *P. aeruginosa* strains expressing C-terminal Strep II-tagged full-length RNase E (RNase E$^{1\text{-}}$

^{1057}-Strep) or an RNase E truncation missing the entire CTD (RNase E$^{1\text{-}529}$-Strep). As an additional

control, the assay was also performed with the *P. aeruginosa* PAO1 wild-type (WT) to discard proteins

interacting nonspecifically with the Strep-Tactin magnetic beads (Fig. 3A-B).

Eluates of the pull-down experiment were first analysed by silver staining (Fig S5A). The RNase E$^{1\text{-}}$

^{1057}-Strep eluate revealed a high molecular weight band corresponding to RNase E, along with additional

bands absent in the untagged and RNase E$^{1\text{-}529}$-Strep samples. These bands likely represent components

of the RNA degradosome or cleaved RNase E$^{1\text{-}1057}$-Strep intermediates, as RNase E is known to be

prone to cleavage [42, 64]. Mass spectrometry (LC-MS/MS) analysis identified a total of 115 proteins

present in both duplicates of the RNase E$^{1\text{-}1057}$-Strep eluates. 13 proteins (including RNase E) were

enriched by at least 5.5-fold as compared with the untagged sample (Fig. 3A). This stringent threshold

was chosen since core components of the RNA degradosome should be abundantly associated to RNase

E. Of these 12 putative partners, 5 (PNPase, GdhB, ArcB, RhlB and RpoB) were completely absent

from both RNase E$^{1\text{-}529}$-Strep duplicates and 2, namely RpoC and RplL, were absent from one replicate

and present in the second replicate with a peptide count of 2 (Fig. 3B). We selected for further study the

first 5 proteins, which were considered the most promising RNA degradosome candidates as their

interaction with Pa RNase E seemed fully dependent on the CTD.

It is worth noting that the RhlE2 RNA helicase was, as expected [54, 55], among the proteins enriched

in the RNase E$^{1\text{-}1057}$-Strep eluate as compared to the untagged control, but surprisingly it was equally

enriched in the RNase E$^{1\text{-}529}$-Strep eluate condition (Fig 3B), indicating that this interaction occurs via

RNase E NTD and not via the CTD.

RNase E direct interaction with PNPase and RhlB occurs via the NDPR and AR1 SLiMs,

respectively

The interaction between RNase E and the 5 candidates was first verified using an adenylate-cyclase-

based bacterial two-hybrid (BTH) assay [65]. Co-expression of T25-PNPase and T25-RhlB with T18-

RNase E resulted in red colonies on MacConkey agar and a significant increase of β-galactosidase

activity compared to empty vector controls (11-fold or 5-fold increase, respectively), confirming their

interaction (Fig. S6A, Fig. 3C). Co-expression of T25-ArcB with T18-RNase E also resulted in a

significant increase in β-galactosidase activity (5-fold increase) compared to the negative control (Fig.

3C, Fig. S6A). To identify the SLiMs mediating the observed interactions, we generated seven CTD

truncations of RNase E, each lacking an additional SLiM as compared with the longer version. Testing

these truncations for their capacity to interact with PNPase revealed that only T18-RNase E$^{1-1025}$ showed

β-galactosidase activity similar to those of T18-RNase E$^{1-1057}$ upon co-expression with T25-PNPase

(Fig. 3D, Fig. S6B). In contrast, all shorter T18-RNase E CTD truncations except T18-RNase E$^{1-529}$,

and also T18-RNase E$^{940-1057}$, exhibited a significant increase of β-galactosidase activity as compared

with the negative controls but lower than T18-RNase E$^{1-1057}$ (Fig. 3D, Fig. S6B). Overall, this suggests

that PNPase interaction with RNase E relies on a complex binding site involving RNase E residues 529-

588 and 940-1057.

Co-expression of the T18-RNase E CTD truncations with T25-RhlB revealed that the protein-protein

interaction is mediated by residues 733-793: no significant interaction could be seen between T25-RhlB

and T18-RNase E$^{1-733}$, although T25-RhlB could interact with T18-RNase E$^{1-793}$ (and every longer CTD

truncation) or with T18-RNase E$^{733-793}$ about as strongly as with T18-RNase E$^{1-1057}$ (Fig. 3E, Fig. S6C).

Of note, neither T18-RNase E$^{733-761}$ nor T18-RNase E$^{762-793}$ were able to restore β-galactosidase activity

above negative control levels when co-expressed with T25-RhlB (Fig. 3E), suggesting that the entire

AR1 SLiM, as defined in Fig. 1B, is necessary for interaction with RhlB.

Surprisingly, all T18-RNase E CTD truncations, except T18-RNase E$^{1-529}$, significantly interact with

T25-ArcB, suggesting a complex binding that is dependent on residues 529-588 (Fig. 3F, Fig. S6D). Of

note, these residues only contain the MTS SLiM, which is presumably involved in membrane binding.

Building on the bacterial two-hybrid assays results, we successively employed affinity chromatography

with purified proteins to investigate direct binding, minimizing the influence of other protein partners

or RNA. Fusion proteins containing a C-terminal 3xFlag tagged RNase E$^{1-529}$, RNase E$^{572-1057}$ (CTD

with no MTS) were purified and immobilized on anti-Flag magnetic beads. After extensive washing,

the beads were mixed with either purified PNPase or RhlB proteins, and the free or bound fractions

were analysed by SDS-PAGE (Fig. 4).

PNPase was retained on beads containing Flag-RNase E$^{572-1057}$, but not on beads alone or containing

Flag-RNase E$^{1-529}$ (Fig. 4A), indicating that the RNase E CTD is sufficient for direct interaction with

PNPase. Point mutations within the NDPR SLiM (Flag-RNase E$^{572-1057}$_NDPRmut) abolished the

binding, confirming that this SLiM is sufficient for direct interaction, as suggested by the BTH assay

(Fig. 4A). Of note, PNPase is not retained on beads containing Flag-RNase E$^{1-588}$, which suggests that

residues 529-588 are not sufficient to mediate a direct protein-protein interaction, with the NDPR SLiM

being the major binding site (Fig. S5B). Similarly, RhlB was retained on Flag-RNase E$^{572-1057}$ but not

on Flag-RNase E$^{1-529}$ and point mutations in the AR1 SLiM (Flag-RNase E$^{572-1057}$_AR1mut) showed that

this region is crucial for RhlB binding (Fig. 4B).

Contrastingly, ArcB was not retained on beads containing either Flag-RNase E$^{1-1057}$ or Flag-RNase E^{1-}

^{588} under the tested conditions, suggesting an indirect type of interaction (Fig. S5C).

To summarize, these experiments clearly establish PNPase and RhlB as core components of *P.*

aeruginosa RNA degradosome and demonstrate that the NDPR and AR1 SLiMs are essential for their

direct interactions with RNase E.

PNPase and RhlB are dependent on their physical interaction with RNase E for colocalising in

RNA degradosome foci.

Given the foci localization of RNase E, we assessed PNPase and RhlB colocalization with RNase E by

using *P. aeruginosa* strains carrying *rne::mCherry* and either *pnp::msfGFP* or *rhl::msfGFP*

chromosomal fusions. As expected, both PNPase-msfGFP and RhlB-msfGFP formed foci near the

membrane and co-localized with RNase E-mCherry in live cells (Fig. 5A). Overlap coefficients between

the two fluorescence channels were consistently over 0.9, while Pearson's coefficients were slightly

lower (0.880 for PNPase and 0.773 for RhlB), likely due to differences in signal intensity. Of note,

strains expressing any of these chromosomal fusion proteins grow identically to the WT strain at 37°C,

suggesting that all the fusion proteins are functional (Figure S7A). We also assessed the subcellular

localisation of ArcB-msfGFP and observed a smooth distribution rather than foci, which suggests that

the interaction between RNase E and ArcB is likely transient, with ArcB not being a core component of

the complex (Fig. S7C).

To further corroborate the need for specific RNase E domains in these interactions, we examined the

localization of PNPase-msfGFP and RhlB-msfGFP in strains expressing RNase E^{1-1057} (WT), RNase E^{1-}

^{529} (*rne529*), RNase E^{AR1mut} (*rne*AR1^{mut}), or RNase $E^{NDPRmut}$ (*rne*NDPR^{mut}). Both PNPase-msfGFP and

RhlB-msfGFP displayed diffuse cytosolic localization in the *rne529* strain compared to the WT strain

(Fig. 5B). In the *rne*NDPR^{mut} strain, PNPase-msfGFP showed a diffuse cytosolic distribution, while

RhlB-msfGFP was diffuse in the *rne*AR1^{mut} strain, highlighting an impaired recruitment into the RNase

E foci. However, the NDPR or AR1 mutations did not prevent the other protein partner from being

recruited into the foci, indicating that both proteins can be independently recruited into RNase E-driven

foci (Fig. 5B). We also confirmed that RNase E NDPRmut-msfGFP forms foci, ruling out the possibility

that the loss of PNPase foci in the *rne*NDPR^{mut} strain is due to impairment of RNase E-driven foci

formation (Figure S7B). These findings confirm the *in vivo* interaction of Pa PNPase and Pa RhlB with

the NDPR and AR1 SLiMs, respectively, and demonstrate that these interactions are crucial for their

near-membrane foci localization.

Phenotypic changes associated with partial or complete deletion of RNase E CTD

To investigate the biological importance of RNase E CTD-mediated complex scaffolding in *P.*

aeruginosa, we performed some phenotypic assays using the *rne529* and *rne588* strains, as well as

strains bearing the AR1 mutation (*rne*AR1^{mut}), NDPR mutation (*rne*NDPR^{mut}), or AR1+AR4+REER

mutation (*rne*AR1+AR4+REER^{mut}), abolishing RhlB binding, PNPase binding, or CTD-mediated RNA

binding plus RhlB binding, respectively.

On solid media at 37°C, the *rne529*, *rne588* and *rne*AR1+AR4+REER^{mut} strains displayed no severe

growth defect but a significantly reduced colony size, suggesting a growth delay (Fig. 6A top panel,

Fig. 6B). Indeed, the *rne529, rne588* and *rne*AR1+AR4+REER^{mut} strains exhibited a slight but

consistent growth delay at 37°C in liquid culture (Fig. 6C). To verify that the growth delay was caused

by truncations/mutations introduced in the *rne* gene, we electroporated the WT or the two most impaired

mutant strains with an inducible vector allowing in-trans expression of RNase E, RNase E-msfGFP, or

RNase E-mCherry. Cultivation of the strains at 37°C in liquid culture revealed that expression of either

native or fluorescently-tagged RNase E could restore identical growth levels compared to the

electroporated WT strain, additionally confirming that the fluorescently-tagged RNase E protein is fully

functional (Fig S8).

Interestingly, when cultured on solid media at 16°C, the *rne588* strain showed impaired growth, while

the growth defects of the *rne529* and *rne*AR1+AR4+REER^mut strains became more pronounced, with

rne529 barely growing (Fig. 6A, upper middle panel). Since the *rne*AR1^mut and the *rne*NDPR^mut strains

exhibited no significant differences compared to the WT, these results indicate that the cold sensitivity

of the *rne588* strain can be attributed to the loss of RNase E CTD RNA binding. Moreover, the severe

growth defect observed in the *rne529* mutant indicates that additional delocalisation of RNase E from

the membrane exacerbates the cold sensitivity caused by deletion of RNase E RNA-binding SLiMs.

Besides low temperature, we also tested sensitivity to low or high pH (Fig. 6A, lower panels): while

growth at pH≃9.5 is rather similar between the different strains, deletion of the CTD (*rne529* and

rne588) impairs growth at pH≃5.5, although the three CTD point mutant strains are mostly unaffected.

This suggests that, although loss of interaction with PNPase or RhlB separately does not seem to impair

growth, the combined loss of RNase E protein-protein and protein-RNA interactions results in a growth

defect under low pH conditions. Finally, we assessed whether the RNase E CTD mutants might be

impaired in virulence using *Galleria mellonella* larvae, a highly predictive infection model for studying

mammalian infection processes [66]. Infection with PAO1 WT, *rne*AR1^mut and *rne*NDPR^mut strains

resulted in nearly 100% larval mortality within 20 hours post-injection (Fig 6D). In contrast, the larvae

infected with the *rne588* mutant exhibited 50% survival at this time point, while those infected with

*rne*AR1+AR4+REER^mut or *rne529* mutants showed 100% survival, with larval death occurring only

after 21 hours post-injection (Fig 6D). These results strongly indicate that the ability of *P. aeruginosa*

to infect a host is compromised when RNase E CTD-mediated RNA binding is disrupted. This

impairment is likely attributable to several contributing elements: reduced bacterial growth within the

host, deficiencies in virulence factor production, and a decreased ability to endure host immune

responses and host-related stresses.

Effect of RNase E CTD on *P. aeruginosa* transcripts steady-state levels

To better understand the phenotypes observed and evaluate the impact of RNase E CTD loss on *P. aeruginosa* transcriptome, we conducted RNA-sequencing analysis on *P. aeruginosa* WT, *rne529,* and *rne588* mutant strains grown at 37°C until the late-exponential growth phase (OD_{600nm} of 1-1.2). Biological triplicates were analysed and show a high degree of relatedness within each strain (Fig. S9A-B). Under these conditions, the loss of the entire RNase E CTD in the *rne529* strain affected the steady-state levels of 450 transcripts, with 183 downregulated and 267 upregulated (adjusted p-value < 0.05, fold change |FC| ≥ 2) (Fig. 7A). In the *rne588* mutant, 417 transcripts were differentially expressed, with 194 downregulated and 223 upregulated (adjusted p-value < 0.05, fold change |FC| ≥ 2) (Fig. 7B). Differential expression analysis on reads mapping to predicted small RNA genes (sRNAs) identified 14 predicted sRNA genes with significantly enriched reads in the *rne529* mutant and 5 in the *rne588* strain (Table S5).

A comparison of the *rne529* and *rne588* strains revealed that 164 mRNAs were consistently regulated in both mutants (97 upregulated and 67 downregulated, see Figure 7C). Among the strongest commonly downregulated genes were: (i) the *pqsABCD*, *pqsE*, and *pqsH* genes, involved in the synthesis of quorum-sensing signal molecules alkylquinolones PQS (2-heptyl-3-hydroxy-4(1H)-quinolone) and HHQ (2-heptyl-4(1H)-quinolone), which play roles in virulence [67], (ii) the *phzA1B1* and *phzA2B2* phenazine biosynthesis genes, which are important for synthesis of the toxic secondary metabolites phenazine and pyocyanin [68], and (iii) the *agtABCD* gene cluster for 4-aminobutyrate (GABA) and 5-aminovalerate (AMV) uptake [69]. Surprisingly, the two strains exhibited more differences in mRNA levels than shared changes. In the *rne588* strain, the mRNA levels corresponding to type 3 secretion system genes, which deliver toxins to host cells [70], were consistently upregulated, while they did not vary significantly in the *rne529* strain as compared to the WT (Fig 7D). On the other hand, ribosomal and many amino acid metabolism genes (tryptophan, arginine, alanine, aspartate and glutamate) were downregulated only in the *rne529* strain, which could correlate with the general growth defect observed (Fig. 7E and Table S5).

Considering the autoregulation of RNase E reported in *E. coli* and *C. crescentus* [63, 71], we quantified

reads mapping to the 5' untranslated region (UTR) of the *rne* gene (transcription start site determined

in [72]) and observed increases of 2 and 0.96 $\log_2$ folds in the *rne529* and *rne588* mutants, respectively,

compared to the WT (Fig. 7F). Similarly, we observed a 1.43 and 0.88 $\log_2$ fold increase in the transcript

corresponding to the NTD of RNase E in the *rne529* and *rne588* mutants, respectively (Fig. 7F), in

agreement with the observed increase in protein levels (Fig. S4C-D).

Finally, to assess RNA degradation efficiency in our strains, we analysed the 5' reads ends of sequenced

RNA fragments. As previously performed in other studies [56, 73], we calculated the ratio between the

number of loci with 5' ends present only in a mutant and not in the WT, over the sum of loci with 5'

read ends seen in both mutant and WT. We performed the analysis in the three genetic backgrounds

(WT, *rne529*, *rne588*) and for each replica independently to ensure consistency (see methods and Fig

S10). Despite the background of random fragmentation intrinsic to the sequencing protocol, a consistent

increase on the number of 5' ends in the *rne529* and *rne588* strains was observed as compared to the

WT (Fig. 7G), suggesting an accumulation of RNA degradation intermediates and an altered RNA

degradation in the RNase E truncated mutants.

Discussion

The composition and the functions of the RNA degradosome vary among bacterial species, likely

reflecting diverse lifestyles and niche adaptations [30]. Studying the RNA degradosome in various

bacterial species is therefore crucial for elucidating how bacteria respond and adapt to different

environmental conditions and sustain various types of stress. In this study, we characterized the RNA

degradosome of *P. aeruginosa*, a versatile opportunistic pathogen and widely used model organism for

bacterial pathogenesis and environmental adaptation.

In the first part of our study, we focused on the putative SLiMs within the CTD of *P. aeruginosa* RNase

E and assessed their contribution to RNA binding and foci assembly. We successively identified RhlB

and PNPase as core RNA degradosome components, as well as the SLiMs mediating these interactions

(Fig. 8). A notable feature of SLiMs is their poor sequence conservation, even when they mediate

interactions with the same protein partner. Our findings confirm this variability for both PNPase and

RhlB interactions.

PNPase, a known partner of RNase E in several species, binds through SLiMs with significantly

different sequences across organisms. For instance, in *Anabaena* PCC7120, the interaction occurs via a

cyanobacterial SLiM having the conserved residues RRRRRRSSA [37], while in *E. coli* or *C.*

crescentus,   the   sequences   are   WQRPTFAFEGKGAAGGHTATHHASA   [30]   and

APPEKPRRGWWRR [43], respectively. Our pull-down assays and *in vitro* binding assays showed that

PNPase interacts with RNase E in *P. aeruginosa* via the NDPR SLiM, which has yet another distinct

sequence (TGRALNDPREKRRLQREAERLAREAAAAA). These sequence variations may reflect

species-specific interactions, as suggested by experimental work showing lack of heterologous

interactions between RNase E and PNPase of *E. coli* and *Pseudoalteromonas haloplanktis* [42]. Despite

sequence variability, the PNPase interacting SLiM is consistently located towards the end of the RNase

E CTD [30, 37, 43], suggesting convergent evolution in the positioning of the PNPase relative to other

RNA degradosome components and to RNase E catalytic domain.

Of note, a recent study on a *P. aeruginosa* mutant with a 50 bp deletion in the 3' end of the *rne* gene

revealed a hypervirulence phenotype, characterized by enhanced cytotoxicity and siderophore

production [74]. Based on *E. coli* RNA degradosome model, this phenotype was associated with a

possible impairment of PNPase binding to Pa RNase E or a different RNA degradosome activity [74].

Our data reveal that PNPase can interact with the RNase E$^{1-1025}$ truncation (BTH assay), ruling out the

former hypothesis. However, we cannot exclude the possibility that this region might be involved in

interaction with a yet unknown partner.

Regarding RhlB, we show that in *P. aeruginosa*, this RNA helicase directly interacts with RNase E via

the AR1 SLiM (RPRRRSRGQRRRSNRRERQR). Mutation of AR1 disrupts not only the RhlB-RNase

E binding *in vitro*, but also RhlB subcellular localisation near the membrane and co-localization with

RNase E *in vivo*. Interestingly, in *E. coli* and other species the AR1 SLiM mediates RNA interaction

[30, 42]. In *P. aeruginosa*, AR1 also contributes to RNA-binding in addition to RhlB binding. Given

that in *E. coli,* a single arginine mutation within the HBS SLiM (R730A) is sufficient to disrupt RhlB

binding [75], it is possible that only one or a few arginine residues of the AR1 SLiM are involved in

interaction with RhlB, while the others mediate RNA binding. Further experiments are needed to

address this hypothesis and to understand the dynamics of the RhlB-RNA-RNase E complex in *P.*

aeruginosa. Moreover, it has been shown that the RNA-dependent ATPase and unwinding activity of

RhlB are stimulated by binding to RNase E [76]. Future studies will investigate whether it is also the

case in *P. aeruginosa*.

In contrast with the *E. coli* RNA degradosome, we did not find any evidence of an interaction between

RNase E and enolase in *P. aeruginosa* when the bacteria grow in rich medium and standard laboratory

conditions. Although our pull-down and bacterial-two hybrid data indicate a probable interaction

between RNase E and ArcB, a catabolic ornithine carbamoyltransferase from the arginine deaminase

pathway [77], no direct protein-protein binding could be observed *in vitro,* nor were the two proteins

colocalising in vivo. We propose that RNase E indirectly interacts with ArcB, probably in a RNA-

dependent manner. Indeed, the *P. aeruginosa arcDABC* polycistronic transcript undergoes

endonucleolytic cleavage at multiple sites for its processing [78], suggesting that an regulatory

interaction between RNase E and ArcB polypeptide could occur during co-translational RNA cleavage

[18].

Additionally, the RNase E-based RNA degradosome of *P. haloplanktis, P. syringae*, and *R. capsulatus*

do not seem to contain either a metabolic enzyme upon standard growing conditions [40-42]. This lack

of RNase E interaction with metabolic enzymes may reflect distinct physiological roles of the RNA

degradosome in different bacterial species, potentially linked to specific metabolic adaptations or

regulatory mechanisms that modulate RNA stability and processing depending on nutrient availability.

However, we cannot exclude that some protein partners were missed as we chose a stringent threshold

for analysis of the pull-down results. Transient or condition-specific accessory protein partners remain

to be identified, and compositional biases such as the conserved AEPV-rich part of the RNase E CTD

are not yet understood, as no ligand or biological function has been assigned to this region even in well-

studied RNase E orthologs.

It is also important to mention that throughout this study, we chose to focus on core RNA degradosome

components that are identified based on their binding to RNase E CTD, but we cannot exclude that

some functional interactions occur with the NTD. For example, RhlB was found to interact with RNase

E NTD in *C. crescentus* [36], while in *Anabaena* the RNA helicase CrhB, RNase II and enolase were

shown to associate with RNase E via its NTD [38, 39]. Similarly, we found that the *P. aeruginosa*

RhlE2 RNA helicase, previously shown to associate with RNase E in an RNA-dependent manner [55],

interacts with the N-terminal domain of Pa RNase E. Although RhlE2 is not considered a core RNA

degradosome component based on our assumption of CTD-mediated interactions, this association

remains functionally significant [55, 56].

In the second part of our study, we evaluated the importance of RNA degradosome assembly at both

transcriptomic and phenotypic levels through an analysis of *P. aeruginosa* strains carrying RNase E

CTD truncations and mutations. Comparing the *rne588* and *rne529* strains allows us to discuss the

importance of RNA degradosome assembly and membrane attachment in *P. aeruginosa*. The two

mutants exhibited varying degrees of sensitivity to cold growth, with *rne529* being the most affected,

indicating that cold growth requires both RNase E membrane attachment and RNA degradosome

assembly. Loss of the CTD without delocalisation from the membrane (*rne588*) does not affect growth

in *P. aeruginosa* under standard laboratory condition (i.e., 37°C and rich medium), but impacts its

capacity to sustain growth in the cold. Transcripts levels were also generally affected in the mutants, as

well as global RNA degradation, as shown in the 5' read ends analysis. Decreased degradation of

cleaved transcripts could lead to an upregulation of their levels. Therefore, direct and indirect effects

remain to be distinguished and those changes remain to be correlated with proteins levels.

Surprisingly, we did not find any phenotypic changes associated to the loss of RhlB or PNPase binding,

questioning the importance of the RNA degradosome protein association for the complex functionality.

Competition assays could be useful in identifying any fitness defects in our mutants, as done in *E. coli*

for RNase E CTD truncations [19, 62], or different phenotypes or stresses could be tested.

On the other hand, our phenotypic analysis highlights the crucial role of CTD RNA binding for the

functionality of the RNA degradosome complex. Mutations in the SLiMs responsible for RNA binding

(AR1, AR4, and the REER-repeats) not only impair *in vivo* RNA degradosome foci assembly and/or

stability but also lead to decreased *P. aeruginosa* virulence and reduced growth at both 37°C and cold

temperatures. The effect of these mutations on foci formation mirrors the effect observed upon cell

exposure to rifampicin, which decreases RNA levels in the cell, and aligns with previous studies

showing that Ec and Cc RNase E orthologs also assemble foci in a CTD-dependent and RNA-dependent

manner [60-62]. Our data thus suggest that impairing foci formation has a major effect on the activity

of the RNA degradosome in *P. aeruginosa*.

Overall, our study identified core RNA degradosome components in *P. aeruginosa* and underscores the

role of this complex in bacterial adaptation to environmental changes and pathogenesis. By revealing

P. aeruginosa-specific interactions and their functional significance, our findings provide valuable

insights into the molecular mechanisms of adaptation in this pathogen and highlight the potential of the

RNA degradosome as a drug target.

Methods

Bacterial strains and growing conditions

A list of bacterial strains used can be found in Table S1. Unless otherwise specified, *P. aeruginosa* PAO1

was grown at 37°C in NYB (25 g of Difco nutrient broth, 5g of Difco yeast agar per liter) with shaking

at 180 rpm or on Nutrient Agar (NA) (40 g Oxoid blood agar base, 5 g Difco yeast agar per liter). When

required, antibiotics were added to these media at the following concentrations: 100 μg/ml ampicillin,

50 μg/mL kanamycin, 20 μg/mL tetracycline and 10 μg/ml gentamicin for *E. coli*; and 50 μg/mL or 100

μg/mL tetracycline and 50 μg/ml gentamicin for *P. aeruginosa*.

Construction of plasmids used in this study

Information about plasmid construction and primers used in this study can be found in supplementary

tables S2 and S3, respectively. Plasmids pUT18C and pKT25 were generally cloned by restriction

enzyme digestion using EcoRI-HF and BamHI-HF (NEB). When a gene to be inserted could be

internally cleaved by EcoRI and/or BamHI, the enzymes MfeI and/or BglII, respectively, were chosen

instead during primer design to allow ligation of the insert to the vector digested with EcoRI/BamHI

thanks to compatible cohesive ends. For cloning of *gdhB* into pKT25, NEB assembly® was used instead,

due to the presence of many restriction enzymes internal cleavage sites. Ligation reactions were carried

our using T4 DNA ligase (Roche or NEB) at 16°C overnight, or at room temperature for 2-3 hours,

following standard protocols. When cloning inserts into the suicide vectors pME3087 or pEXG2, the

NEBuilder HiFi DNA assembly Master Mix (NEB) was used according to the manufacturer's

recommendations. To constructs RNase E CTD mutants, the DNA was synthesized by GeneArt

(ThermoFisher Scientific) and cloned into appropriate vectors for protein purification or strain

construction. A description of the introduced mutations can be found in Table S4. To construct

pME6032-based expression vectors, restriction enzyme digestion was carried out using EcoRI and KpnI

for both vector and inserts.

Ligation mixtures or the assembly reactions were transformed into chemically competent *E. coli* DH5α

cells by heat-shock, plating on appropriate selective LB agar medium. The resulting colonies were

screened by PCR for the presence of the insert in the plasmid of interest. After plasmid extraction

(Miniprep kit, Sigma-Aldrich/Merck), plasmids were verified by Sanger sequencing at Microsynth.

Construction of *P. aeruginosa* mutant strains

To construct chromosomal gene truncations or insert a Strep-tag or fluorescent protein into the

chromosome, in frame with a gene of interest, the suicide vectors pEXG2 [79] or pME3087 [80] were

used. Briefly, approximately 500 bp regions located immediately upstream or downstream of the region

to be edited, and, when applicable, the tag or fluorescent protein of interest, were amplified by PCR and

inserted together into the vector backbone using NEB assembly®. After sequencing verification of the

constructed plasmid quality, the plasmids were introduced into *P. aeruginosa* PAO1 strain by triparental

mating using the *E. coli* strain HB101 (pRK2013). After selection with the appropriate antibiotic

(tetracycline for pME3087 or gentamicin for pEXG2), merodiploids were resolved as described in [81]

for pME3087 or [79] for pEXG2, and strains with the desired insertion/deletion were identified by

colony PCR.

RNase E sequence analysis

Group-specific homologs of RNase E in *P. aeruginosa*, *P. fluorescens*, *P. protegens*, *P. putida* and *P.*

stutzeri were retrieved from NCBI databases using blastp [82] with *P. aeruginosa* PAO1 RNase E as a

query; only sequences with query coverage > 0.5 were conserved (total of 5000 proteins). The number

of RE[E/D]R motifs in each protein sequence was determined using the vcountPattern function from

the "biostrings" R package [83]. The similarity of the protein sequences within a taxonomic group was

determined using the conserv function (method="similarity") from the bio3D R package on a muscle

alignment of the protein sequences[84, 85].

Electrophoretic motility shift assay

The *malEF* transcript was synthesized with a T7 transcription kit (Invitrogen) on linear DNA template

(172 bp) obtained by PCR with primers p253-p254 (see Table S3). The electrophoretic mobility shift

assay (EMSA) was performed following the protocol of [86]. Briefly, the reaction mixtures (15 μl)

contained 35 nM of RNA, 50mM Tris-HCl, 100mM NaCl, 5mM DTT, 1.0 units/μl RNasin (Promega),

and His10Smt3-RNase E proteins as specified, with protein concentration expressed as a molar ratio of

relative to RNA concentration (see Fig. 2A and S4A-B).

The reactions were incubated at 4 °C for 30 min before loading on 2% agarose gel and run at 150 V for

15 min. RNA bands were visualized by gel staining with SybrGold (Invitrogen).

Epifluorescence microscopy imaging

Phase contrast and fluorescence microscopy were performed on a Zeiss Axio Imager 2 equipped with

an objective alpha Plan-Apochromat 100x/1,46 Oil Ph3 M27 and a ZEISS Axiocam 305 mono CCD

camera. Bacteria were grown until OD_{600nm}= 0.2-0.6 (approx. 2.5 hours) in 3 ml of NYB and were then

placed on an agarose pad containing 1% agarose dissolved in 0.5X PBS. Images were processed with

Fiji (NIH, USA) [87].

Image analysis

All image treatments and analysis were performed using Fiji (NIH, USA)[87]. Representative images

shown had contrast adjusted for display by opening the images in Fiji and using the Reset function in

the Adjust Brightness/Contrast menu, after average background intensity subtraction using the Math ->

Subtract function. These average background intensity values were obtained by measuring a square area

devoid of any cells.

For quantification of signal intensity in Fiji, a line was drawn along the longitudinal axis of bacterial

cells and signal was measured using Plot Profile. Results of this function were saved as .csv files. For

each strain, at least 50 bacterial cells were measured from the raw images, using at least two different

images for quantification. Average background values were subsequently subtracted from each results

file using a .csv file containing the Plot Profile measurements of a linear area devoid of cells for each

image used. Boxplots of fluorescence intensity and signal variance or pooled histograms were plotted

using R [88]. The signal variance was obtained by calculating the variance of the $\log_2$ fold-change

between each measured value and the average of all values for a given measured cell. This was done to

normalize the data and obtain comparable variances without introducing a bias due to differences in

overall signal intensity.

Pull-down identification of RNase E partners

Fresh cultures of PAO1 WT, PAO1 RNase E_{1-529}-Strep, or PAO1 RNase E_{1-1057}-Strep were inoculated to

OD=0.05 from overnight cultures in 500 mL Erlenmeyer flasks. Cultures were grown until approx.

OD_{600nm}=0.8-0.9. 50 mL of each culture was harvested by centrifugation at 4500 $x\ g$ for 10 minutes.

Pellets were stored at -80°C until needed.

Each thawed pellet was then resuspended in 3,25 mL of lysis buffer (for 10 mL of buffer, prepared just

before use: 20 mM Tris (pH 10), 250 mM NaCl, 10% v/v glycerol, 1 EDTA-free protease inhibitor

tablet, 50 μg/mL lysozyme). Samples were incubated the samples on ice for about 30 minutes,

transferred into 15-mL Falcon tubes and sonicated using a thin probe for 6x30s, with 15 seconds break

in between. A pulse of 70% and a power output of 50% were used. After sonication, 32.5 μL of Triton

X100 10% were added to each sample (~0.1% final) before incubation on a rotary wheel at 4°C for 8

minutes. The samples were transferred into 2 mL Eppendorf tubes and centrifuged for 1 hour at 10500

$x\ g$ at 4°C. MagStrep® Strep-Tactin®XT beads (IBA, cat# 2-5090) were washed in wash buffer (20 mM

Tris (pH 10), 250 mM NaCl, 10% v/v glycerol) according to the manufacturer's instructions. One mix

of beads was prepared for all the samples with a volume of beads suspension of 45 μL per sample. Final

resuspension was done in 200 μL wash buffer per sample, plus an additional 10 μL. After centrifugation

of the samples, the supernatant of each sample was regrouped into a 5 mL Eppendorf tube, and 200 μL

of washed beads suspension were added per sample. Incubation on a rotary wheel was done for 2h15 at

4°C. The 5 mL tubes were then placed against a magnetic rack and all the supernatant was carefully

removed. A first wash volume of 500 μL was added gently and agitated by horizontal rotations of the

tubes. Tubes were placed back against the magnetic rack and the wash volume was removed. A second

wash step was performed identically with a volume of 225 μL instead of 500 μL. A third wash step of

225 μL was also performed, and the resuspended beads were transferred into a clean 1.5 mL Eppendorf

tube by gentle pipetting with a 1000 μL tip, before removing the wash volume. The beads were covered

with 10 μL of sterile PBS and stored at 4°C before LC-MS/MS analysis at the Proteomics Core Facility

(University of Geneva). For plots in Fig. 3A-B, the log2 fold-change (log2FC) was calculated using the

following formula: log2(average peptide count of protein x identified in the RNase E$^{1\text{-xxx}}$-Strep / average

peptide count of protein x identified in the untagged sample).When an average count of 0 peptide was

identified in the untagged sample, the log2FC value was instead calculated using the following formula:

log2FC(average peptide count of protein x identified in the RNase E$^{1\text{-xxx}}$-Strep/1).

Bacterial adenylate cyclase two-hybrid assay

Cloned plasmids were introduced into chemically competent BTH101 cells by heat shock, and the cells

were plated in LB agar supplemented with ampicillin and kanamycin. Isolated colonies were then

streaked as a line of about 1-2 cm on freshly prepared Mac Conkey agar medium supplemented with

ampicillin, kanamycin, 0.5 mM isopropyl β-d-thiogalactoside (IPTG), and maltose 1%. The

MacConkey Petri dishes were then incubated at 30°C for 24h followed by an incubation at room

temperature (~23°C) for 24h. The bacteria were imaged by scanning the plates, after which

quantification of β-galactosidase activity was done using ONPG as described in [89].

Protein purification

His$_{10}$Smt3-tagged PNPase, His$_{10}$Smt3-tagged RhlB, and His$_{10}$Smt3-tagged ArcB proteins were purified

from soluble bacterial lysates by nickel-agarose affinity chromatography as described previously in the

supplementary information of Hausmann *et al.* 2021 [55]. The imidazole elution profiles were

monitored by SDS-PAGE. The recombinant His$_{10}$Smt3-tagged proteins were recovered predominantly

in the 250 mM imidazole fraction. Protein concentration was determined using the Bio-Rad dye reagent,

with BSA as the standard, and calculated by interpolation against the BSA standard curve. The His_{10}-

Smt3 tag remained uncleaved in the protein-protein interaction assays.

RNase E^{1-1057}, RNase E^{1-588}, RNase E^{1-529}, RNase $E^{572-1057}$, RNase $E^{572-1057\_NDPRmut}$, and RNase E^{572-}

$^{1057\_AR1mut}$ proteins, each carrying a His_{10}Smt3 tag at the N-terminus and a Twin-Strep-tag®-Flag at the

C-terminus, were purified as described above. The 250 mM imidazole elution fractions were used

directly in the Protein-Protein interaction assays. For the EMSA experiment, the RNase E proteins were

further purified as follows: the 250 mM imidazole fraction were applied to a 1 ml of Strep-Tactin

Sepharose (Iba) column that had been equilibrated with buffer C (50 mM Tris pH8, 200 mM NaCl, 10

% glycerol). The column was washed with 5 ml of buffer C and then eluted stepwise with 1 ml aliquots

of buffer C containing, 2.5 mM D-Desthiobiotin respectively. The elution profiles were monitored by

SDS-PAGE.

Protein-protein interaction assays

Anti-Flag M2 magnetic beads (Sigma M8823) were used for the binding assay. Unless otherwise

specified, a volume of 25 µL of beads suspension was used per reaction sample. The beads were washed

once with 1 mL of buffer H (50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 0.01% Triton X-100), before

being resuspended in fresh buffer H and aliquoted into one Eppendorf tube per sample. Purified RNase

E^{1-529}-Flag, RNase E^{1-588}-Flag RNase $E^{572-1057}$-Flag, RNase $E^{572-1057\_NDPRmut}$- Flag, or RNase E^{572-}

$^{1057\_AR1mut}$- Flag fusion proteins (20-25 µg each) were diluted in buffer H to a final volume of 100 µL

and incubated with the magnetic beads for 1 hour at 4 °C on a rotating device. When using purified

RNase E^{1-1057}-Flag fusion protein instead, a volume of 450 µL of purified protein (in a buffer with 50

mM Tris pH=8, 500 mM NaCl, 10% glycerol and 250 mM imidazole) was incubated with 50 µl of Anti-

Flag M2 magnetic beads (Sigma M8823). After the incubation, the beads were collected using a

magnetic separator and washed three times with 0.25 ml of binding buffer H to remove any unbound

protein.

For affinity chromatography, the anti-Flag magnetic beads (Sigma-Aldrich) with bound RNase E-Flag

fusion proteins, or unbound beads, were incubated with either 5 µg of PNPase (Figure 4A), 5 µg of

RhlB (Figure 4B), or 5 µg of ArcB diluted in buffer H (final volume 100 µL, 50 µl of beads suspension

used for probing ArcB binding) for 1 hour at 4 °C on a rotating device. After incubation, the beads were

collected using the magnet, and the supernatant (free fraction F) was withdrawn. The anti-Flag beads

were then resuspended in 100 µl of binding buffer H and subjected to three rounds of washing of 100

µl each. Following the final wash, the proteins bound to the beads were eluted with 100 µl of 2X LDS

sample buffer (Invitrogen) supplemented with 167 mM DTT. Aliquots (15 µl) of the input protein

sample (PNPase or RhlB), the first supernatant fraction (free fraction F), and the bead-bound fraction

were mixed with 5 µl of 4X LDS sample buffer (Invitrogen), heated at 80 °C for 5 minutes, and analysed

by SDS-PAGE. Polypeptides were visualized using Coomassie Blue staining.

Growth assays on solid or liquid media

To assess the growth of the mutant strains in liquid medium, 200 µL of NYB broth was inoculated to

OD_{600nm}=0.05 in a 96-well plate, using an overnight culture washed twice with fresh medium. The wells

on the edge of the plate were not used. The plate was incubated at 37°C using a Biotek Synergy H1

Hybrid Plate reader. OD_{600nm} values were measured every 30 minutes. Before each OD_{600nm}

measurement, orbital shaking with 282 cpm (3 mm) was performed for 1 minute. Data was retrieved

the following day and plotted after substraction of blank OD_{600nm} values using GraphPad Prism.

When growing strains carrying the pME6032 vector for complementation assays, NYB medium was

supplemented with 50 µg/mL of Tetracycline and 70 µM of IPTG for induction. Cells were resuspended

in supplemented NYB after washing and before inoculating the 96-well plate.

To assess the growth of the mutant strains on solid medium, square Petri dishes (12x12cm) were filled

with 50 mL of warm NA. When testing growth at low vs high pH, either 75 µL of 37% fuming HCl or

400 µL of 5N NaOH was added per 50 mL of NA agar to achieve the desired conditions. A fresh culture

(OD_{600nm}= 0.4-0.8) of each strain was adjusted to OD_{600nm}=1 by centrifugation and serially diluted in a

96-well plate using a multipipette. 5 µL of each dilution series was then spotted using a multipipette

onto pre-dried NA plates (1 hour under a sterile flowhood). After the spots had dried, plates were

incubated upside down at 37°C or 16°C.

Galleria killing assays

Galleria mellonella larvae were used to assess in vivo virulence and/or survival of several RNase E mutant strains. Fresh cultures were adjusted to a suspension of $OD_{600nm}= 1$ and serially diluted to 10^{-6} in Gibco Dulbecco's Phosphate Buffered Saline (dPBS) (Thermo Fisher Scientific). 10 µL of the 10^{-6} dilution were spotted onto NYB agar to verify that the dilutions were similar for the different strains. 10 µL spots were then put onto a clean parafilm and injected using Omnican 50 sterile insulin syringe (Braun) into the second to last leg of the larvae, facing towards the head. As a control, sterile dPBS was injected similarly into larvae to assess background levels of death. For every 10 larvae injected, the time of injection was recorded, and they were placed inside a dark incubator at 37°C. The next day, deaths were recorded every hour. A larva was considered dead when it completely stopped responding to stimuli (touched on the legs or the head with a pipette tip). Time of death was calculated as the interval of time between recorded time of injection and death occurence. Survival curves were plotted and analysed with GraphPad Prism (GraphPad Software, Prism10.2).

RNA-seq sample collection

Independent triplicates cultures of PAO1 WT, PAO1 RNase E_{1-588}, (*rne588*) and PAO1 RNase E_{1-529} (*rne529*) strains were grown in 50 mL Erlenmeyer flasks (10 mL NYB broth) until $OD_{600nm}= 1.0-1.2$. 500 µL of each culture were immediately added into four separate 1.5 mL tubes, each containing 1 mL of RNAlater (Sigma-Aldrich, cat). After harvesting of all three strains, the tubes were centrifuged for 30 minutes at 9600 x *g*. Supernatant was discarded and pellets were stored at -80°C until RNA extraction. The RNA extraction was performed using two tubes of pellets for each strain and replicate, with the Monarch Total RNA Miniprep kit. For lysis of the bacterial cells, the two pellets were resuspended in 250 µL of 1X Tris-EDTA (TE) buffer supplemented with 1 mg/mL of lysozyme and regrouped into one tube for each strain, incubated for 5 minutes at RT, and RNA was subsequently extracted following the manufacturer's instructions. After RNA purification, an additional DNA removal treatment was done using TURBO DNase (Thermo Fisher) according to the manufacturer's instructions. RNA was purified again using the RNA purification columns (dark blue) from the Monarch Total RNA Miniprep kit, and following the priming, washing and elution steps described by the

manufacturer. The absence of residual gDNA contamination was verified by performing a PCR for 35

cycles using primer pairs targeting *rpoD* [55]. Biological triplicate samples for each strain were

prepared. Quality check, ribosomal RNA depletion and RNA sequencing were conducted by Novogene.

RNA sequencing analysis

Reads were mapped using Bowtie2 [90] to *Pseudomonas aeruginosa* PAO1 genome assembly

ASM676v1 and counts were aggregated using summarize Overlaps in the package GenomicAlignments

[91] in R using the ASM676v1 gtf version 2.2 (NCBI accession: GCF_000006765.1). Differential

expression was performed using DESeq2 [92] and transcripts were considered differentially expressed

between two conditions if the absolute value of log2FC between conditions was greater than 1 (i.e. 2-

fold up or down) with a adjusted p-value less than 0.05. To find functional enrichment among

differentially expressed genes groups, the Kyoto Encyclopedia of Genes and Genomes (KEGG) was

used for gene set definitions, downloaded using the package KEGGREST [93] in R. Enrichment was

calculated using a hypergeometric test. All correlations presented are Pearson's product moment

correlation coefficient. Pairwise correlations of log2 Reads Per Kilobase per Million mapped reads

(RPKMs) and hierarchical unsupervised clustering analysis of fold changes relative to WT demonstrate

the similarity in the profile of the biological replicates for each strain. Analysis of 5' read ends per loci

was performed as described by de Araújo et al. [94] except that each dataset read number was down-

sampled to 8 million.

Acknowledgments

We thank Sylvain Guex-Crosier for his assistance with RNA extraction and Nathanaël Guggenheim for

protein purification and assistance with cloning experiments. We also acknowledge the Proteomics Core

Facility and the Bioinformatics Support Platform from the University of Geneva for assistance with the

realisation of LC-MS-MS and bioinformatics analyses, respectively. Work in M.V. laboratory is funded

by Swiss National Science Foundation (PCEFP3_203343 to M.V.) and Fondation Pierre Mercier pour

la Science (to M.V.). D.G. work is supported by the Swiss National Science Foundation

(PZ00P3_180142 to D.G., 2020–2023) and the Velux Foundation (grant 1814 to P.J. and D.G., 2023).

References

1.   Hui MP, Foley PL, Belasco JG. Messenger RNA Degradation in Bacterial Cells. Annual Review of Genetics. 2014;48(1):537-59. doi: 10.1146/annurev-genet-120213-092340. PubMed PMID: 25292357.

2.   Celesnik H, Deana A, Belasco JG. Initiation of RNA decay in *Escherichia coli* by 5' pyrophosphate removal. Mol Cell. 2007;27(1):79-90. doi: 10.1016/j.molcel.2007.05.038. PubMed PMID: 17612492; PubMed Central PMCID: PMCPMC2196405.

3.   Deana A, Celesnik H, Belasco JG. The bacterial enzyme RppH triggers messenger RNA degradation by 5' pyrophosphate removal. Nature. 2008;451(7176):355-8. doi: 10.1038/nature06475. PubMed PMID: 18202662.

4.   Luciano DJ, Vasilyev N, Richards J, Serganov A, Belasco JG. Importance of a diphosphorylated intermediate for RppH-dependent RNA degradation. RNA Biol. 2018;15(6):703-6. Epub 20180529. doi: 10.1080/15476286.2018.1460995. PubMed PMID: 29619898; PubMed Central PMCID: PMCPMC6152431.

5.   Tejada-Arranz A, de Crecy-Lagard V, de Reuse H. Bacterial RNA Degradosomes: Molecular Machines under Tight Control. Trends Biochem Sci. 2020;45(1):42-57. Epub 20191101. doi: 10.1016/j.tibs.2019.10.002. PubMed PMID: 31679841; PubMed Central PMCID: PMCPMC6958999.

6.   Dressaire C, Picard F, Redon E, Loubière P, Queinnec I, Girbal L, Cocaign-Bousquet M. Role of mRNA stability during bacterial adaptation. PLoS One. 2013;8(3):e59059. PubMed PMID: 23516597.

7.   Broglia L, Le Rhun A, Charpentier E. Methodologies for bacterial ribonuclease characterization using RNA-seq. FEMS Microbiol Rev. 2023;47(5). PubMed PMID: 37656885.

8.   Ono M, Kuwano M. A conditional lethal mutation in an *Escherichia coli* strain with a longer chemical lifetime of messenger RNA. J Mol Biol. 1979;129(3):343-57. Epub 1979/04/15. doi: 10.1016/0022-2836(79)90500-x. PubMed PMID: 110942.

9.   Apirion D. Isolation, genetic mapping and some characterization of a mutation in *Escherichia coli* that affects the processing of ribonuleic acid. Genetics. 1978;90(4):659-71. PubMed PMID: 369943.

10.   Jain C, Deana A, Belasco JG. Consequences of RNase E scarcity in *Escherichia coli*. Molecular Microbiology. 2002;43(4):1053-64. doi: https://doi.org/10.1046/j.1365-2958.2002.02808.x.

11.   Caruthers JM, Feng Y, McKay DB, Cohen SN. Retention of core catalytic functions by a conserved minimal ribonuclease E peptide that lacks the domain required for tetramer formation. J Biol Chem. 2006;281(37):27046-51. Epub 20060719. doi: 10.1074/jbc.M602467200. PubMed PMID: 16854990.

12. Callaghan AJ, Aurikko JP, Ilag LL, Günter Grossmann J, Chandran V, Kühnel K, et al. Studies

of the RNA degradosome-organizing domain of the *Escherichia coli* ribonuclease RNase E. J Mol Biol.

2004;340(5):965-79. doi: 10.1016/j.jmb.2004.05.046. PubMed PMID: 15236960.

13. Carpousis AJ, Van Houwe G, Ehretsmann C, Krisch HM. Copurification of E. coli RNAase E

and PNPase: evidence for a specific association between two enzymes important in RNA processing

and degradation. Cell. 1994;76(5):889-900. doi: 10.1016/0092-8674(94)90363-8. PubMed PMID:

7510217.

14. Py B, Higgins CF, Krisch HM, Carpousis AJ. A DEAD-box RNA helicase in the *Escherichia*

coli RNA degradosome. Nature. 1996;381(6578):169-72. doi: 10.1038/381169a0. PubMed PMID:

8610017.

15. Chandran V, Luisi BF. Recognition of enolase in the Escherichia coli RNA degradosome. J Mol

Biol. 2006;358(1):8-15. Epub 20060221. doi: 10.1016/j.jmb.2006.02.012. PubMed PMID: 16516921.

16. Worrall JAR, Górna M, Crump NT, Phillips LG, Tuck AC, Price AJ, et al. Reconstitution and

Analysis of the Multienzyme Escherichia coli RNA Degradosome. Journal of Molecular Biology.

2008;382(4):870-83. doi: https://doi.org/10.1016/j.jmb.2008.07.059.

17. Bandyra KJ, Bouvier M, Carpousis AJ, Luisi BF. The social fabric of the RNA degradosome.

Biochim Biophys Acta. 2013;1829(6-7):514-22. Epub 20130228. doi: 10.1016/j.bbagrm.2013.02.011.

PubMed PMID: 23459248; PubMed Central PMCID: PMCPMC3991390.

18. Tsai YC, Du D, Domínguez-Malfavón L, Dimastrogiovanni D, Cross J, Callaghan AJ, et al.

Recognition of the 70S ribosome and polysome by the RNA degradosome in *Escherichia coli*. Nucleic

Acids Res. 2012;40(20):10417-31. Epub 20120825. doi: 10.1093/nar/gks739. PubMed PMID:

22923520; PubMed Central PMCID: PMCPMC3488216.

19. Leroy A, Vanzo NF, Sousa S, Dreyfus M, Carpousis AJ. Function in *Escherichia coli* of the

non-catalytic part of RNase E: role in the degradation of ribosome-free mRNA. Mol Microbiol.

2002;45(5):1231-43. doi: 10.1046/j.1365-2958.2002.03104.x. PubMed PMID: 12207692.

20. Dendooven T, Paris G, Shkumatov AV, Islam MS, Burt A, Kubańska MA, et al. Multi-scale

ensemble properties of the *Escherichia coli* RNA degradosome. Mol Microbiol. 2022;117(1):102-20.

Epub 20210925. doi: 10.1111/mmi.14800. PubMed PMID: 34415624; PubMed Central PMCID:

PMCPMC7613265.

21. Sinha D, De Lay NR. Target recognition by RNase E RNA-binding domain AR2 drives sRNA

decay in the absence of PNPase. Proc Natl Acad Sci U S A. 2022;119(48):e2208022119. Epub

20221121. doi: 10.1073/pnas.2208022119. PubMed PMID: 36409892; PubMed Central PMCID:

PMCPMC9860253.

22. Lee K, Zhan X, Gao J, Qiu J, Feng Y, Meganathan R, et al. RraA. a protein inhibitor of RNase

E activity that globally modulates RNA abundance in *E. coli*. Cell. 2003;114(5):623-34. PubMed PMID:

13678585.

23.   Górna MW, Pietras Z, Tsai YC, Callaghan AJ, Hernández H, Robinson CV, Luisi BF. The
regulatory protein RraA modulates RNA-binding and helicase activities of the E. coli RNA
degradosome. Rna. 2010;16(3):553-62. Epub 20100127. doi: 10.1261/rna.1858010. PubMed PMID:
20106955; PubMed Central PMCID: PMCPMC2822920.

24.   Gao J, Lee K, Zhao M, Qiu J, Zhan X, Saxena A, et al. Differential modulation of E. coli mRNA
abundance by inhibitory proteins that alter the composition of the degradosome. Mol Microbiol.
2006;61(2):394-406. Epub 20060612. doi: 10.1111/j.1365-2958.2006.05246.x. PubMed PMID:
16771842.

25.   Prud'homme-Généreux A, Beran RK, Iost I, Ramey CS, Mackie GA, Simons RW. Physical and
functional interactions among RNase E, polynucleotide phosphorylase and the cold-shock protein,
CsdA: evidence for a 'cold shock degradosome'. Molecular Microbiology. 2004;54(5):1409-21. doi:
https://doi.org/10.1111/j.1365-2958.2004.04360.x.

26.   Blum E, Py B, Carpousis AJ, Higgins CF. Polyphosphate kinase is a component of the
Escherichia   coli   RNA   degradosome.   Molecular   Microbiology.   1997;26(2):387-98.   doi:
https://doi.org/10.1046/j.1365-2958.1997.5901947.x.

27.   Kannaiah S, Goldberger O, Alam N, Barnabas G, Pozniak Y, Nussbaum-Shochat A, et al.
MinD-RNase E interplay controls localization of polar mRNAs in E. coli. Embo j. 2024;43(4):637-62.
Epub 20240119. doi: 10.1038/s44318-023-00026-9. PubMed PMID: 38243117; PubMed Central
PMCID: PMCPMC10897333.

28.   Taghbalout A, Rothfield L. RNaseE and the other constituents of the RNA degradosome are
components of the bacterial cytoskeleton. Proceedings of the National Academy of Sciences.
2007;104(5):1667-72. doi: 10.1073/pnas.0610491104.

29.   Ikeda Y, Yagi M, Morita T, Aiba H. Hfq binding at RhlB-recognition region of RNase E is
crucial for the rapid degradation of target mRNAs mediated by sRNAs in Escherichia coli. Mol
Microbiol. 2011;79(2):419-32. Epub 20101118. doi: 10.1111/j.1365-2958.2010.07454.x. PubMed
PMID: 21219461.

30.   Ait-Bara S, Carpousis AJ, Quentin Y. RNase E in the gamma-Proteobacteria: conservation of
intrinsically disordered noncatalytic region and molecular evolution of microdomains. Mol Genet
Genomics. 2015;290(3):847-62. Epub 20141129. doi: 10.1007/s00438-014-0959-5. PubMed PMID:
25432321; PubMed Central PMCID: PMCPMC4435900.

31.   Goldblum K, Apririon D. Inactivation of the ribonucleic acid-processing enzyme ribonuclease
E blocks cell division. J Bacteriol. 1981;146(1):128-32. doi: 10.1128/jb.146.1.128-132.1981. PubMed
PMID: 6163761; PubMed Central PMCID: PMCPMC217061.

32.   Bernstein JA, Lin PH, Cohen SN, Lin-Chao S. Global analysis of *Escherichia coli* RNA
degradosome function using DNA microarrays. Proc Natl Acad Sci U S A. 2004;101(9):2758-63. Epub
20040223. doi: 10.1073/pnas.0308747101. PubMed PMID: 14981237; PubMed Central PMCID:
PMCPMC365694.

33.   Kido M, Yamanaka K, Mitani T, Niki H, Ogura T, Hiraga S. RNase E polypeptides lacking a

carboxyl-terminal half suppress a mukB mutation in *Escherichia coli*. J Bacteriol. 1996;178(13):3917-

25. doi: 10.1128/jb.178.13.3917-3925.1996. PubMed PMID: 8682798; PubMed Central PMCID:

PMCPMC232654.

34.   Lopez PJ, Marchand I, Joyce SA, Dreyfus M. The C-terminal half of RNase E, which organizes

the *Escherichia coli* degradosome, participates in mRNA degradation but not rRNA processing in vivo.

Mol Microbiol. 1999;33(1):188-99. doi: 10.1046/j.1365-2958.1999.01465.x. PubMed PMID:

10411735.

35.   Hardwick SW, Chan VS, Broadhurst RW, Luisi BF. An RNA degradosome assembly in

Caulobacter crescentus. Nucleic Acids Res. 2011;39(4):1449-59. Epub 20101015. doi:

10.1093/nar/gkq928. PubMed PMID: 20952404; PubMed Central PMCID: PMCPMC3045602.

36.   Voss JE, Luisi BF, Hardwick SW. Molecular recognition of RhlB and RNase D in the

Caulobacter crescentus RNA degradosome. Nucleic Acids Res. 2014;42(21):13294-305. Epub

20141111. doi: 10.1093/nar/gku1134. PubMed PMID: 25389270; PubMed Central PMCID:

PMCPMC4245959.

37.   Zhang JY, Deng XM, Li FP, Wang L, Huang QY, Zhang CC, Chen WL. RNase E forms a

complex with polynucleotide phosphorylase in cyanobacteria via a cyanobacterial-specific nonapeptide

in the noncatalytic region. RNA. 2014;20(4):568-79. Epub 20140221. doi: 10.1261/rna.043513.113.

PubMed PMID: 24563514; PubMed Central PMCID: PMCPMC3964918.

38.   Yan HD, Qin XX, Wang L, Chen WL. Both Enolase and the DEAD-Box RNA Helicase CrhB

Can Form Complexes with RNase E in *Anabaena* sp. Strain PCC 7120. Applied and environmental

microbiology. 2020;86(13). doi: 10.1128/AEM.00425-20. PubMed PMID: WOS:000546650600011.

39.   Zhou C, Zhang JY, Hu XY, Li CC, Wang L, Huang QY, Chen WL. RNase II binds to RNase E

and modulates its endoribonucleolytic activity in the cyanobacterium *Anabaena* PCC 7120. Nuceic

Acid Res. 2020;48(7):3922-34. doi: 10.1093/nar/gkaa092. PubMed PMID: WOS:000525957700040.

40.   Purusharth RI, Klein F, Sulthana S, Jäger S, Jagannadham MV, Evguenieva-Hackenberg E, et

al. Exoribonuclease R Interacts with Endoribonuclease E and an RNA Helicase in the Psychrotrophic

Bacterium Pseudomonas syringae Lz4W*. Journal of Biological Chemistry. 2005;280(15):14572-8.

doi: https://doi.org/10.1074/jbc.M413507200.

41.   Jäger S, Fuhrmann O, Heck C, Hebermehl M, Schiltz E, Rauhut R, Klug G. An mRNA

degrading complex in *Rhodobacter capsulatus*. Nucleic Acids Res. 2001;29(22):4581-8. doi:

10.1093/nar/29.22.4581. PubMed PMID: 11713307; PubMed Central PMCID: PMCPMC92556.

42.   Aït-Bara S, Carpousis Agamemnon J. Characterization of the RNA Degradosome of

Pseudoalteromonas haloplanktis: Conservation of the RNase E-RhlB Interaction in the

Gammaproteobacteria. Journal of Bacteriology. 2010;192(20):5413-23. doi: 10.1128/jb.00592-10.

43.   Hardwick SW, Gubbey T, Hug I, Jenal U, Luisi BF. Crystal structure of *Caulobacter crescentus*
polynucleotide phosphorylase reveals a mechanism of RNA substrate channelling and RNA
degradosome assembly. Open Biology. 2012;2(4):120028. doi: 10.1098/rsob.120028.

44.   Nurmohamed S, Vaidialingam B, Callaghan AJ, Luisi BF. Crystal Structure of Escherichia coli
Polynucleotide Phosphorylase Core Bound to RNase E, RNA and Manganese: Implications for
Catalytic Mechanism and RNA Degradosome Assembly. Journal of Molecular Biology.
2009;389(1):17-33. doi: https://doi.org/10.1016/j.jmb.2009.03.051.

45.   Green SK, Schroth MN, Cho JJ, Kominos SK, Vitanza-jack VB. Agricultural plants and soil as
a reservoir for *Pseudomonas aeruginosa*. Applied microbiology. 1974;28(6):987-91. Epub 1974/12/01.
PubMed PMID: 4217591; PubMed Central PMCID: PMCPMC186868.

46.   Hardalo C, Edberg SC. *Pseudomonas aeruginosa*: assessment of risk from drinking water.
Critical   reviews   in   microbiology.   1997;23(1):47-75.   Epub   1997/01/01.   doi:
10.3109/10408419709115130. PubMed PMID: 9097014.

47.   Pendleton JN, Gorman SP, Gilmore BF. Clinical relevance of the ESKAPE pathogens. Expert
review of anti-infective therapy. 2013;11(3):297-308. Epub 2013/03/06. doi: 10.1586/eri.13.12.
PubMed PMID: 23458769.

48.   https://www.who.int/news/item/27-02-2017-who-publishes-list-of-bacteria-for-which-new-
antibiotics-are-urgently-needed.

49.   Garcia-Clemente M, de la Rosa D, Máiz L, Girón R, Blanco M, Olveira C, et al. Impact of
Pseudomonas aeruginosa Infection on Patients with Chronic Inflammatory Airway Diseases. Journal of
clinical medicine. 2020;9(12). Epub 2020/12/02. doi: 10.3390/jcm9123800. PubMed PMID: 33255354.

50.   Urwin L, Okurowska K, Crowther G, Roy S, Garg P, Karunakaran E, et al. Corneal Infection
Models: Tools to Investigate the Role of Biofilms in Bacterial Keratitis. Cells. 2020;9(11). Epub
2020/11/14. doi: 10.3390/cells9112450. PubMed PMID: 33182687; PubMed Central PMCID:
PMCPMC7696224.

51.   Costerton JW, Stewart PS, Greenberg EP. Bacterial biofilms: a common cause of persistent
infections.   Science   (New   York,   NY).   1999;284(5418):1318-22.   Epub   1999/05/21.   doi:
10.1126/science.284.5418.1318. PubMed PMID: 10334980.

52.   Lu Q, Eggimann P, Luyt CE, Wolff M, Tamm M, François B, et al. Pseudomonas aeruginosa
serotypes in nosocomial pneumonia: prevalence and clinical outcomes. Critical care (London, England).
2014;18(1):R17. Epub 2014/01/17. doi: 10.1186/cc13697. PubMed PMID: 24428878; PubMed Central
PMCID: PMCPMC4057348.

53.   Hancock RE. Resistance mechanisms in *Pseudomonas aeruginosa* and other nonfermentative
gram-negative bacteria. Clin Infect Dis. 1998;27 Suppl 1:S93-9. Epub 1998/08/26. PubMed PMID:
9710677.

54.   Van den Bossche A, Hardwick SW, Ceyssens PJ, Hendrix H, Voet M, Dendooven T, et al.
Structural elucidation of a novel mechanism for the bacteriophage-based inhibition of the RNA

degradosome. Elife. 2016;5. Epub 20160722. doi: 10.7554/eLife.16413. PubMed PMID: 27447594; PubMed Central PMCID: PMCPMC4980113.

55. Hausmann S, Gonzalez D, Geiser J, Valentini M. The DEAD-box RNA helicase RhlE2 is a global regulator of *Pseudomonas aeruginosa* lifestyle and pathogenesis. Nucleic Acids Res. 2021;49(12):6925-40. doi: 10.1093/nar/gkab503. PubMed PMID: 34151378; PubMed Central PMCID: PMCPMC8266600.

56. Hausmann S, Geiser J, Allen GE, Geslain SAM, Valentini M. Intrinsically disordered regions regulate RhlE RNA helicase functions in bacteria. Nucleic Acids Res. 2024. Epub 20240614. doi: 10.1093/nar/gkae511. PubMed PMID: 38874491.

57. Sonnleitner E, Romeo A, Bläsi U. Small regulatory RNAs in *Pseudomonas aeruginosa*. RNA Biol. 2012;9(4):364-71. Epub 2012/02/18. doi: 10.4161/rna.19231. PubMed PMID: 22336763.

58. Valentini M, Gonzalez D, Mavridou DA, Filloux A. Lifestyle transitions and adaptive pathogenesis of *Pseudomonas aeruginosa*. Current opinion in microbiology. 2018;41:15-20. Epub 2017/11/23. doi: 10.1016/j.mib.2017.11.006. PubMed PMID: 29166621.

59. Nishio S-y, Itoh T. Arginine-rich RNA binding domain and protein scaffold domain of RNase E are important for degradation of RNAI but not for that of the Rep mRNA of the ColE2 plasmid. Plasmid. 2009;62(2):83-7. doi: https://doi.org/10.1016/j.plasmid.2009.04.002.

60. Hamouche L, Poljak L, Carpousis AJ. Polyribosome-Dependent Clustering of Membrane-Anchored RNA Degradosomes To Form Sites of mRNA Degradation in *Escherichia coli*. mBio. 2021;12(5):e0193221. Epub 20210907. doi: 10.1128/mBio.01932-21. PubMed PMID: 34488454; PubMed Central PMCID: PMCPMC8546579.

61. Al-Husini N, Tomares DT, Bitar O, Childers WS, Schrader JM. α-Proteobacterial RNA Degradosomes Assemble Liquid-Liquid Phase-Separated RNP Bodies. Mol Cell. 2018;71(6):1027-39.e14. Epub 20180906. doi: 10.1016/j.molcel.2018.08.003. PubMed PMID: 30197298; PubMed Central PMCID: PMCPMC6151146.

62. Khemici V, Poljak L, Luisi BF, Carpousis AJ. The RNase E of *Escherichia coli* is a membrane-binding protein. Mol Microbiol. 2008;70(4):799-813. Epub 20081002. doi: 10.1111/j.1365-2958.2008.06454.x. PubMed PMID: 18976283; PubMed Central PMCID: PMCPMC7610891.

63. Schuck A, Diwa A, Belasco JG. RNase E autoregulates its synthesis in Escherichia coli by binding directly to a stem-loop in the rne 5' untranslated region. Mol Microbiol. 2009;72(2):470-8. Epub 20090306. doi: 10.1111/j.1365-2958.2009.06662.x. PubMed PMID: 19320830; PubMed Central PMCID: PMCPMC2857391.

64. Liu S-J, Lin G-M, Yuan Y-Q, Chen W, Zhang J-Y, Zhang C-C. A conserved protein inhibitor brings under check the activity of RNase E in cyanobacteria. Nucleic Acids Research. 2024;52(1):404-19. doi: 10.1093/nar/gkad1094.

65. Karimova G, Ullmann A, Ladant D. [5] A bacterial two-hybrid system that exploits a cAMP
signaling cascade in Escherichia coli. In: Thorner J, Emr SD, Abelson JN, editors. Methods in
Enzymology. 328: Academic Press; 2000. p. 59-73.

66. Jander G, Rahme Laurence G, Ausubel Frederick M. Positive Correlation between Virulence
ofPseudomonas aeruginosa Mutants in Mice and Insects. Journal of Bacteriology. 2000;182(13):3843-
5. doi: 10.1128/jb.182.13.3843-3845.2000.

67. Lin J, Cheng J, Wang Y, Shen X. The Pseudomonas Quinolone Signal (PQS): Not Just for
Quorum Sensing Anymore. Front Cell Infect Microbiol. 2018;8:230. Epub 20180704. doi:
10.3389/fcimb.2018.00230. PubMed PMID: 30023354; PubMed Central PMCID: PMCPMC6039570.

68. Gonçalves T, Vasconcelos U. Colour Me Blue: The History and the Biotechnological Potential
of Pyocyanin. Molecules. 2021;26(4). Epub 20210210. doi: 10.3390/molecules26040927. PubMed
PMID: 33578646; PubMed Central PMCID: PMCPMC7916356.

69. Chou HT, Li J-Y, Lu C-D. Functional Characterization of the agtABCD and agtSR Operons for
4-Aminobutyrate and 5-Aminovalerate Uptake and Regulation in Pseudomonas aeruginosa PAO1.
Current Microbiology. 2014;68(1):59-63. doi: 10.1007/s00284-013-0446-y.

70. Hauser AR. The type III secretion system of Pseudomonas aeruginosa: infection by injection.
Nat Rev Microbiol. 2009;7(9):654-65. doi: 10.1038/nrmicro2199. PubMed PMID: 19680249; PubMed
Central PMCID: PMCPMC2766515.

71. Nandana V, Al-Husini N, Vaishnav A, Dilrangi KH, Schrader JM. Caulobacter crescentus
RNase E condensation contributes to autoregulation and fitness. Mol Biol Cell. 2024;35(8):ar104. Epub
20240612. doi: 10.1091/mbc.E23-12-0493. PubMed PMID: 38865176; PubMed Central PMCID:
PMCPMC11321048.

72. Wurtzel O, Yoder-Himes DR, Han K, Dandekar AA, Edelheit S, Greenberg EP, et al. The single-
nucleotide resolution transcriptome of Pseudomonas aeruginosa grown in body temperature. PLoS
Pathog. 2012;8(9):e1002945. Epub 20120927. doi: 10.1371/journal.ppat.1002945. PubMed PMID:
23028334; PubMed Central PMCID: PMCPMC3460626.

73. de Araújo Hugo L, Picinato Beatriz A, Lorenzetti Alan PR, Muthunayake Nisansala S,
Rathnayaka-Mudiyanselage IW, dos Santos Naara M, et al. The DEAD-box RNA helicase RhlB is
required for efficient RNA processing at low temperature in Caulobacter. Microbiology Spectrum.
2023;11(6):e01934-23. doi: 10.1128/spectrum.01934-23.

74. Vaillancourt M, Galdino ACM, Limsuwannarot SP, Celedonio D, Dimitrova E, Broerman M,
et al. A compensatory RNase E variation increases Iron Piracy and Virulence in multidrug-resistant
Pseudomonas aeruginosa during Macrophage infection. PLoS Pathog. 2023;19(4):e1010942. Epub
20230407. doi: 10.1371/journal.ppat.1010942. PubMed PMID: 37027441; PubMed Central PMCID:
PMCPMC10115287.

75. Khemici V, Toesca I, Poljak L, Vanzo NF, Carpousis AJ. The RNase E of *Escherichia coli* has at least two binding sites for DEAD-box RNA helicases: functional replacement of RhlB by RhlE. Molecular Microbiology. 2004;54(5):1422-30. doi: https://doi.org/10.1111/j.1365-2958.2004.04361.x.

76. Worrall JA, Howe FS, McKay AR, Robinson CV, Luisi BF. Allosteric activation of the ATPase activity of the *Escherichia coli* RhlB RNA helicase. J Biol Chem. 2008;283(9):5567-76. PubMed PMID: 18165229.

77. Lüthi E, Mercenier A, Haas D. The arcABC Operon Required for Fermentative Growth of Pseudomonas aeruginosa on Arginine: Tn5-751-assisted Cloning and Localization of Structural Genes. Microbiology. 1986;132(10):2667-75. doi: https://doi.org/10.1099/00221287-132-10-2667.

78. Gamper M, Ganter B, Polito MR, Haas D. RNA processing modulates the expression of the *arcDABC* operon in *Pseudomonas aeruginosa*. Journal of Molecular Biology. 1992;226(4):943-57. doi: https://doi.org/10.1016/0022-2836(92)91044-P.

79. Rietsch A, Vallet-Gely I, Dove SL, Mekalanos JJ. ExsE, a secreted regulator of type III secretion genes in Pseudomonas aeruginosa. Proc Natl Acad Sci U S A. 2005;102(22):8006-11. Epub 20050523. doi: 10.1073/pnas.0503005102. PubMed PMID: 15911752; PubMed Central PMCID: PMCPMC1142391.

80. Laville J, Voisard C, Keel C, Maurhofer M, Défago G, Haas D. Global control in Pseudomonas fluorescens mediating antibiotic synthesis and suppression of black root rot of tobacco. Proc Natl Acad Sci U S A. 1992;89(5):1562-6. doi: 10.1073/pnas.89.5.1562. PubMed PMID: 1311842; PubMed Central PMCID: PMCPMC48492.

81. Ye RW, Haas D, Ka JO, Krishnapillai V, Zimmermann A, Baird C, Tiedje JM. Anaerobic activation of the entire denitrification pathway in Pseudomonas aeruginosa requires Anr, an analog of Fnr. J Bacteriol. 1995;177(12):3606-9. doi: 10.1128/jb.177.12.3606-3609.1995. PubMed PMID: 7768875; PubMed Central PMCID: PMCPMC177071.

82. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J Mol Biol. 1990;215(3):403-10. doi: 10.1016/s0022-2836(05)80360-2. PubMed PMID: 2231712.

83. Pagès HA, P.; Gentleman, R.; DebRoy, S. Biostrings: Efficient manipulation of biological strings. 2024.

84. Grant BJ, Rodrigues AP, ElSawy KM, McCammon JA, Caves LS. Bio3d: an R package for the comparative analysis of protein structures. Bioinformatics. 2006;22(21):2695-6. Epub 20060829. doi: 10.1093/bioinformatics/btl461. PubMed PMID: 16940322.

85. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res. 2004;32(5):1792-7. Epub 20040319. doi: 10.1093/nar/gkh340. PubMed PMID: 15034147; PubMed Central PMCID: PMCPMC390337.

86. Ream JA, Lewis LK, Lewis KA. Rapid agarose gel electrophoretic mobility shift assay for quantitating protein: RNA interactions. Anal Biochem. 2016;511:36-41. Epub 20160802. doi: 10.1016/j.ab.2016.07.027. PubMed PMID: 27495142; PubMed Central PMCID: PMCPMC5002362.

87.   Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82. Epub 20120628. doi: 10.1038/nmeth.2019. PubMed PMID: 22743772; PubMed Central PMCID: PMCPMC3855844.

88.   Team RC. R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing. Vienna, Austria2023.

89.   Bordi C, Lamy MC, Ventre I, Termine E, Hachani A, Fillet S, et al. Regulatory RNAs and the HptB/RetS signalling pathways fine-tune Pseudomonas aeruginosa pathogenesis. Mol Microbiol. 2010;76(6):1427-43. Epub 20100401. doi: 10.1111/j.1365-2958.2010.07146.x. PubMed PMID: 20398205; PubMed Central PMCID: PMCPMC2904497.

90.   Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nature Methods. 2012;9(4):357-9. doi: 10.1038/nmeth.1923.

91.   Lawrence M, Huber W, Pagès H, Aboyoun P, Carlson M, Gentleman R, et al. Software for Computing and Annotating Genomic Ranges. PLOS Computational Biology. 2013;9(8):e1003118. doi: 10.1371/journal.pcbi.1003118.

92.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biology. 2014;15(12):550. doi: 10.1186/s13059-014-0550-8.

93.   B. TDM. KEGGREST: Client-side REST access to the Kyoto Encyclopedia of Genes and Genomes (KEGG). doi:10.18129/B9.bioc.KEGGREST. R package version 1420, https://bioconductororg/packages/KEGGREST2023.

94.   de Araújo HL, Picinato BA, Lorenzetti APR, Muthunayake NS, Rathnayaka-Mudiyanselage IW, Dos Santos NM, et al. The DEAD-box RNA helicase RhlB is required for efficient RNA processing at low temperature in Caulobacter. Microbiol Spectr. 2023:e0193423. Epub 2023/10/18. doi: 10.1128/spectrum.01934-23. PubMed PMID: 37850787.

95.   Erdős G, Pajkos M, Dosztányi Z. IUPred3: prediction of protein disorder enhanced with unambiguous experimental annotation and visualization of evolutionary conservation. Nucleic Acids Research. 2021;49(W1):W297-W303. doi: 10.1093/nar/gkab408.

96.   Xue B, Dunbrack RL, Williams RW, Dunker AK, Uversky VN. PONDR-FIT: a meta-predictor of intrinsically disordered amino acids. Biochim Biophys Acta. 2010;1804(4):996-1010. Epub 20100125. doi: 10.1016/j.bbapap.2010.01.011. PubMed PMID: 20100603; PubMed Central PMCID: PMCPMC2882806.

97.   McLaren RS, Newbury SF, Dance GS, Causton HC, Higgins CF. mRNA degradation by processive 3'-5' exoribonucleases in vitro and the implications for prokaryotic mRNA decay in vivo. J Mol Biol. 1991;221(1):81-95. PubMed PMID: 1920421.

98.   Bolte S, Cordelières FP. A guided tour into subcellular colocalization analysis in light microscopy. J Microsc. 2006;224(Pt 3):213-32. doi: 10.1111/j.1365-2818.2006.01706.x. PubMed PMID: 17210054.

1062

 Figure Legends

 Figure 1: Comparison of *E. coli* (Ec) and *P. aeruginosa* (Pa) RNase E architecture. (A) Ec RNase

 E architecture, adapted from [17]. The NTD is shown is grey, and experimentally defined regions

 necessary for interaction with the membrane, with RNA, or with the three core protein partners are

 highlighted in red, dark blue, and yellow/purple/pink, respectively. **(B)** Pa RNase E architecture based

 on in-silico analyses performed in Fig S1-S2 and in [30]. The SLiM nomenclature was taken from [30],

 while the consensus sequence characterising each SLiM was specifically adapted based on sequence

 conservation within the Pseudomonas genus (Fig S1). MTS: membrane-targeting sequence, AR4:

 arginine-rich motif 4, REER-repeats: region with high frequency of occurrence of "RE[E/D]R" motif,

 M29: motif 29, AR1: arginine-rich motif 1, M20: motif 20, NDPR: motif with a highly conserved NDPR

 sequence. All diagrams were drawn to scale. IUPRED3 [95] and PONDR [96] predictions of disorder

 propensity (probability score) are shown for Ec RNase E **(A)** and Pa RNase E **(B)**.

 Figure 2: RNA binding of Pa RNase E CTD via the AR1, AR4, and REER SLiMs. (A) EMSA with

 purified His-Smt3 tagged RNase E CTD$^{572\text{-}1057}$ variants having either the native sequence (WT) or

 systematic alanine mutation of the arginine residues found within the REER-repeats region (REERmut),

 the AR4 SLiM (AR4mut) or the AR1 SLiM (AR1mut). The *malEF* mRNA intergenic region was used

 as a substrate [97]. Ratios of RNA:protein ranging from 1:10 to 1:80 were tested. See Fig S2 for

 additional EMSA with additional RNA:protein ratios. **(B)** Representative microscopy images of a PAO1

 strain expressing the RNase E-msfGFP fusion or mutated or truncated variants as indicated. **(C)**

 Quantification of signal intensity variance in the strains of panel B as indication of foci disappearance

 (see methods). **(D)** Representative microscopy images of a PAO1 strain expressing the RNase E-

 msfGFP fusion after exposure to rifampicin (+ Rif, exposure to a concentration of 100 μg/mL for 30

 minutes) or to DMSO as a negative control (- Rif). Corrected mean fluorescent signal intensity **(E)** and

 signal intensity variance **(F)** is showed in the graphs (see methods). Each quantification analysis

 involved at least 50 individual cells. The scale is 2 μm. Statistical significance was assessed by

1090   performing t-tests using R. p-values for comparisons tested are indicated as follows: *:<0.05; **:<0.01;

1091   ***:<0.001; ****:<0.0001.

1092

Figure 3: **Identification of scaffold domain-dependent RNase E protein partners by pull-down**

assays and mapping of the interactions by bacterial two-hybrid assays (A) Volcano plot

highlighting the relative enrichment of proteins co-eluting with the RNase E^{1-1057}-Strep bait protein

compared to the untagged negative control. The plot was generated from the average peptide count

obtained in duplicate pull-down experiments. An arbitrary threshold of 5.5-fold change was chosen for

selection of candidate (c) proteins. **(B)** Scatterplot highlighting comparative enrichment of proteins co-

eluting with the RNase E^{1-1057}-Strep bait protein (x axis) or with the RNase E^{1-529}-Strep bait protein (y

axis) compared to the untagged negative control. **(C)** Quantification of β-galactosidase activity in *E.*

coli BTH101 cells co-expressing T18-RNase E with T25-PNPase, T25-RhlB, T25-ArcB, T25-GdhB or

T25-RpoB. Interaction between T18-RNase E and a T25-tagged protein partner results in a significant

restoration of β-galactosidase activity compared with empty vector (e.v.) negative controls. Unpaired

Welch's t-tests was performed using GraphPad Prism to assess differences between every tested protein-

protein interaction and the respective negative control. ***: p-value <0.0001; n.s: not significant (99%

confidence level). **(D-F)** Mapping of the interaction between RNase E and PNPase **(D)**, RhlB **(E)**, or

ArcB **(F)** by quantification of β-galactosidase activity in *E. coli* BTH101 cells co-expressing T25-

PNPase **(D)**, T25-RhlB **(E)**, or T25-ArcB **(F)** and either T18-RNase E or one of eight to ten truncated

T18-RNase E fragments. Empty vector (e.v.) negative controls are included into each separate graph

for comparison.

1111

Figure 4: PNPase and RhlB directly interact with RNase E scaffold domain via NDPR and AR1

SLiMs, respectively. Purified Flag-tagged RNase E^{1-529} or RNase $E^{572-1057}$ proteins having either native

or mutated sequence were used as baits, immobilized to anti-flag magnetic beads and incubated with

purified His-Smt3 tagged PNPase **(A)** or RhlB **(B)** proteins to assess direct protein binding. I: input, F:

free fraction, B: bound fraction. NDPRmut, AR1mut: mutations of NDPR and AR1 SLiMs, respectively

(see Table S4). Binding assays were repeated at least 2 times and representative gels are shown.

1118

Figure 5: **Colocalisation of PNPase or RhlB with RNase E foci is dependent on their interaction with NDPR and AR1 RNase E SLiMs, respectively. (A)** Visualisation of PNPase-msfGFP (upper panels) or RhlB-msfGFP (lower panels) fusion and RNase E-mCherry fusion co-expressed in the same strain. Pearson's and M coefficients were calculated in Fiji using the JaCoP plugin [98]. Three independent images were used, each containing at least 30 cells, and an average coefficient was calculated from the three values obtained with each image. **(B)** Visualisation of PNPase-msfGFP (upper panels) and RhlB-msfGFP (lower panels) in strains expressing either the native RNase E protein (full length) or truncated variants, as indicated above each image. Scale is 2 μm.

1127

Figure 6: Phenotypes affected by deletion or mutation of RNase E scaffolding domain. (A) Growth on NA agar plates at 37°C or 16°C, or under low vs high pH conditions of wild-type (WT), *rne588*, *rne529*, *rne*AR1^{mut}, *rne*NDPR^{mut}, and *rne*AR1+AR4+REER^{mut} strains, as indicated. Experiments were performed in biological independent triplicates; a representative plate is shown. **(B)** Measured colony size for the different mutant strains as indicated. Quantification was done using the Measure function in Image J after drawing a circle around the edge of each colony. 11 to 13 colonies were quantified for each strain. A one-way Anova test was performed using GraphPad and comparing the colony size of each mutant strain to that of the WT strain. ***: p-value <0.0001; n.s: not significant (99% confidence interval). **(C)** Growth in NYB broth at 37°C in 96-well plates of wild-type (WT), *rne588*, *rne529*, *rne*AR1^{mut}, *rne*NDPR^{mut}, and *rne*AR1+AR4+REER^{mut} strains, as indicated. Independent biological triplicates, each with four technical replicates, were used to generate the average growth curve shown. **(D)** Killing curves of *G. mellonella* larvae injected with 10 μL of WT or mutant *P. aeruginosa* strains (as in panel A). The control group was injected with 10 μL of sterile PBS instead. Larvae were incubated at 37°C and larval death was monitored over 13-26 hours post-injection. A larva was considered dead when it completely stopped responding to repeated touching stimuli. Two independent replicates were performed using a total of at least 50 larvae for each strain tested. A Kaplan-Meier test for survival analysis was performed using GraphPad Prism and comparing the survival curve of each mutant strain to that of the WT strain. ***: p-value <0.0001; n.s: not significant (99% confidence interval).

1146

Figure 7: RNA-seq analysis of RNase E CTD truncated mutants reveals broad transcriptome

changes. (A-B) Volcano plots highlighting genes differentially regulated in the **(A)** rne^{1-529} (R529) or

(B) rne^{1-588} (R588) mutant strains compared with the WT strain. Genes significantly up- or down-

regulated (adjusted p-value < 0.05, fold change |FC| ≥ 2) are highlighted in red or blue, respectively.

(C) Scatterplot of genes differentially regulated in rne^{1-529} strain versus genes differentially regulated

in rne^{1-588} strain, highlighting commonalities and differences between the two mutant strains. **(D-E)**

Clustered heat maps of type 3 secretion system **(D)** or ribosomal proteins **(E)** transcript levels in the

RNase E$^{1-529}$ and by about 2-fold in the RNase E$^{1-588}$ strains as compared to the WT strain (fold change).

(F) Differential analyses (log 2-fold change) of normalized total reads counts corresponding to the 5'

UTR region of the PA2976 (*rne*) transcript or RNase E NTD coding sequence in the RNase E$^{1-529}$ and

by about 2-fold in the RNase E$^{1-588}$ strains as compared to the WT strain. **(G)** Proportion of loci with 5'

read ends (sequenced fragments) counted only in a strain (numerator) relative to the 5' read ends counted

in that strain and in a WT replica (denominator). A total of 8 million reads was analysed in each strain

to exclude an effect due to the library depth (see methods). Histograms of individual values calculated

per each strain and replicate is shown in Figure S10.

1162

1163 **Figure 8: Model of RNA degradosome in *P. aeruginosa*.** Numbers corresponds to RNase E SLiM

1164 start-end residues involved in membrane, RNA, RhlB or PNPase interaction. RNA is represented as a

1165 dotted line. See discussion for details.

Figure 1

A

B

Figure 2

Figure 4

Figure 5

A

RNase E-mCherry   PNPase-msfGFP   Merged

Pearson's = 0.880
M1 = 0.910
M2 = 0.943

RNase E-mCherry   RhlB-msfGFP   Merged

Pearson's = 0.773
M1 = 0.937
M2 = 0,924

B

PAO1 WT   PAO1 *rne529*   PAO1 *rne*AR1^mut   PAO1 *rne*NDPR^mut

PNPase-msfGFP

RhlB-msfGFP

Figure 6

Figure 8