Critical functions and key interactions mediated by the RNase E scaffolding domain in *Pseudomonas aeruginosa*

Sandra Amandine Marie Geslain, Stéphane Hausmann, Johan Geiser, George Edward Allen, Diego Gonzalez and Martina Valentini*

Department of Microbiology and Molecular Medicine, CMU, Faculty of Medicine, University of Geneva, Geneva, Switzerland.

* To whom correspondence should be addressed. e-mail: Martina.Valentini@unige.ch

Abstract

The RNA degradosome, a multi-protein complex regulating mRNA in bacteria, assembles on the RNase E C-terminal domain (CTD) in Pseudomonadota (Proteobacteria) via short linear motifs (SLiMs). The composition of the *Pseudomonas aeruginosa* RNA degradosome is largely unknown, and its RNase E CTD shows limited similarity to those in models like *Escherichia coli* or *Caulobacter crescentus*. Our study characterizes the function of conserved SLiMs in *P. aeruginosa* RNase E, including a duplicated sequence termed the "REER-repeats" region. This region, along with AR1 and AR4 SLiMs, mediates RNA binding and localizes the degradosome in subcellular foci. Pull-down and bacterial two-hybrid assays identified PNPase and RhlB as interacting proteins, with direct interactions confirmed via protein binding assays. The NDPR and AR1 SLiMs mediate these interactions, respectively. Transcriptome analyses and phenotypic assays of RNase E CTD mutants revealed growth defects in specific conditions and virulence defects, revealing the importance of RNase E CTD RNA binding and RNA degradosome scaffolding for *P. aeruginosa* adaptability.

Author Summary

In this study, we investigate the RNA degradosome in *Pseudomonas aeruginosa*, a bacterium known

for causing multiple types of human infections and for its multi-drug resistance. The RNA degradosome

is a crucial multi-protein complex that regulates RNA levels, helping bacteria to rapidly adapt to various

environments. We identified the key components of this complex in *P. aeruginosa* and mapped their

interactions. Our findings also reveal that interaction between this complex and RNA is important for

bacterial growth, particularly in cold conditions and during infection. We finally compare our findings

with the RNA degradosome in other species, noting significant evolutionary convergence and unique

characteristics in *P. aeruginosa*.

Our research sheds light on the complex mechanisms bacteria use to manage RNA and adapt to their

surroundings, which could eventually inform new strategies to combat bacterial infections and

resistance.

Introduction

In bacteria, the processing and degradation of mRNAs is primarily orchestrated by the RNA degradosome, a multi-enzyme complex that includes ribonucleases, RNA helicases, and various other proteins [1-5]. The RNA degradosome plays a crucial role in enabling bacteria to rapidly adapt to changing environmental conditions by swiftly modifying mRNA levels and broadly regulating gene expression [6, 7].

Most of our knowledge on the RNA degradosome derives from studies in *Escherichia coli*, where the RNase E endonuclease serves as the central component [8-11]. While the catalytic activity of *E. coli* RNase E resides in its N-terminal domain (NTD, amino acids 1-529), the C-terminal domain (CTD, amino acids 530-1061), or scaffolding domain, is intrinsically disordered and contains short linear motifs (SLiMs) of 10 to 40 amino acids to which the other RNA degradosome components bind [11, 12]. Core *E. coli* RNA degradosome components, namely the polyribonucleotide phosphorylase PNPase [13], the RNA helicase RhlB [14], and the glycolytic enzyme enolase [15], interact with RNase E in semi-stoichiometric ratios via the PBS (PNPase binding site), HBS (helicase binding site) and EBS (enolase binding cite) SLiM, respectively [16, 17]. The *E. coli* RNase E CTD also includes an amphipathic helix, called membrane targeting sequence or MTS, which tethers the entire complex to the inner membrane, and two RNA-binding SLiMs, AR1 (named also RBD) and AR2, which mediate contact with translating ribosomes, small non-coding RNAs and mRNAs [12, 18-21]. Additional proteins, such as RraA [22, 23], RraB [24], CsdA [25], Ppk [26], MinD [27, 28], and Hfq [29], can interact with *E. coli* RNase E in sub-stochiometric ratios or under non-standard conditions of growth (reviewed in [30]). Of importance, the RNase E NTD is essential for *E. coli* growth and rRNA processing [31], whereas deletion of the CTD is not lethal and alters mRNA levels without affecting rRNAs [19, 32-34]. This highlights the dual role of RNase E, with the NTD being crucial overall cell maintenance, and the CTD regulating specific RNA substrates through the assembly of the RNA degradosome.

RNase E orthologs are present in approximately 49% of sequenced bacterial genomes, and nearly all Pseudomonadota (formerly called Proteobacteria) encode at least one RNase E ortholog, with the notable exception of most *Campylobacter* and *Helicobacter* species [5, 30]. The RNase E NTD is remarkably conserved across most species, while the CTD varies considerably in sequence, length, and SLiMs composition, even between species belonging to the same order [30]. SLiMs variability is thought to drive the diversity of RNase E interacting partners, and therefore RNA degradosome composition, among bacterial species. For instance, in *Caulobacter crescentus,* RNase E interacts with PNPase, RhlB, RNase D, and the Krebs cycle enzyme aconitase [35, 36]. In *Anabaena* sp. strain PCC 7120, RNase E partners with PNPase, enolase, the RNA helicase CrhB, and RNase II [37-39], while in *Pseudomonas syringae* LZ4W, only RNase R and the RNA helicase RhlE were found to co-purify with RNase E in pull-down assays [40]. Similarly, in *Rhodobacter capsulatus*, purification of the RNA

degradosome complex via glycerol-gradient centrifugation of cell extracts revealed the presence of

RNase E, two unnamed RNA helicases, the Rho transcription termination factor, and two unidentified

proteins of 47 and 36 kDa, with only low amounts of PNPase present in the purified fractions [41].

In these examples, the SLiMs mediating protein-protein interactions were not always characterized.

Besides, even when the interaction of RNase E with a certain protein partner is conserved across several

species, such as the RNase E-PNPase and RNase E-RhlB interactions, the SLiMs involved in the

binding show little sequence identity [30, 36, 37, 42-44]. Therefore, predicting RNA degradosome

composition based solely on the RNase E CTD sequence does not seem possible. Experimental

characterization of the SLIMs across different bacterial species and their impact on the RNA

degradosome composition and function is therefore essential, providing insights into novel regulatory

mechanisms that enable bacteria to thrive under stress and adapt to diverse environments.

Little is known about the RNA degradosome composition of the opportunistic pathogen *Pseudomonas*

aeruginosa, which is a leading cause of nosocomial infections worldwide largely due to its remarkable

adaptability to different environments and resistance to antibiotics [45-53]. A recent study has revealed

that the φKZ bacteriophage protein Dip interacts with *P. aeruginosa* RNase E, leading to its co-elution

with RNase E but also with PNPase, the PA0428 RNA helicase, and several other proteins in pull-down

assays [54]. Therefore, the RNA degradosome of *P. aeruginosa* was assumed to be composed of these

proteins [5], although their interaction with RNase E has neither been confirmed nor mapped. Our

laboratory has characterized the PA0428 RNA helicase, named RhlE2, confirming an RNA-dependent

interaction with RNase E and identifying the RhlE2 C-terminal region as necessary and sufficient for

this binding [55, 56]. However, the specific RNase E interaction site remained unmapped.

In this study, we characterize the SLiMs within the CTD of *P. aeruginosa* RNase E, identifying those

involved in RNA binding (AR1, AR4, and REER-repeats) as well as those mediating interactions with

core RNA degradosome components, which we determine to be PNPase and the RNA helicase RhlB

(NDPR and AR1, respectively). We also show that RhlE2 interacts with the RNase E NTD in pull-down

assays. Protein-protein interactions were probed with multiple experimental approaches, including an

analysis of the subcellular localization dynamics of the complex. Additionally, we performed

transcriptomic and phenotypic analyses of RNase E CTD mutant strains, revealing the critical role of

RNA degradosome complex scaffolding in bacterial virulence, adaptation to cold temperatures and to

pH fluctuations.

The ability of *P. aeruginosa* to cause disease is intrinsically linked to a sophisticated gene regulatory

network that facilitates bacterial survival and virulence. Within this network, the role of post-

transcriptional gene regulation remains only partially explored [57, 58]. By uncovering the composition

and functional dynamics of the RNA degradosome in *P. aeruginosa*, our study not only advances the

understanding of the molecular mechanisms driving bacterial environmental adaptation but also reveals

potential novel drug targets, which could be crucial in the fight against antibiotic-resistant infections.

Results

Pa RNase E harbours several uncharacterised SLiMs conserved within the Pseudomonadales

P. aeruginosa RNase E (Pa RNase E) consists of an NTD (amino acids 1-529) that shares high sequence identity (70%) and similarity (90%) with the NTD of *E. coli* RNase E (Ec RNase E), and a CTD (amino acids 530-1057) that shows only 25% identity and 37% similarity with the CTD of Ec RNase E (Fig. S1). Aït-Bara and colleagues (2015) identified seven putative SLiMs on the Pa RNase E CTD (MTS, AR4, REE, M29, AR1, M20 and NDPR) [59], which are shown in comparison to the Ec RNase E in Fig. S2A. Of note, the MTS is the only SLIM of Pa RNase E CTD with a propensity to form a secondary structure while in Ec RNase E, the helicase, enolase, and PNPase binding sites also exhibited this feature (Fig. 1 and [12]).

To refine the prediction of Pa RNase E SLiMs, we performed a multiple sequence alignment of 5000 Pseudomonas RNase E orthologs and identified conserved motifs (Fig. S3). Several highly conserved regions of the CTD overlap with the SLiMs previously identified, with the notable presence of additional conserved residues located immediately adjacent to these SLiMs (Fig S2/S3). Based on these data, we redefined the boundaries of the SLiMs to integrate these conserved residues (Fig. 1B, see Fig. S2A for former SLiMs). Upon manual inspection of the CTD sequence, we also observed a large duplicated sequence (residues 635-667 and 682-714) within the CTD region containing the REE and motif 29 SLiMs. Since motif 29 is characterised by the conserved motif REERQPR, we have merged the two SLiMs and renamed the entire region (residues 614-729) as the "REER-repeats" region (Fig. 1B). The number of REER repeats varies across Pseudomonas species: *P. aeruginosa* strains typically harbour 13 RE[ED]R repeats, while the environmental *P. fluorescens*, *P. protegens*, *P. stutzeri*, and *P. syringae* groups show a distribution of repeats peaking around 8-10. Exception is *P. putida*, which exhibits a high frequency of 20-23 RE[ED]R repeats (Fig. S2B). This variability suggests a high frequency of indels occurring within the REER-repeats region.

Overall, our in-silico analyses of Pa RNase E CTD sequence redefined the number and length of the putative Pa RNase E SLiMs, highlighting a potential functional significance of the REER-repeats region in Pseudomonas species, and suggesting different RNA degradosome complex dynamics in *P. aeruginosa* as compared to *E. coli*.

The REER-repeats, AR1 and AR4 SLiMs mediate RNase E CTD RNA binding and foci formation

Previous works in *E. coli* identified arginine-rich SLiMs as RNA-binding motifs [21, 60]. We therefore hypothesized that the *P. aeruginosa* AR1 and AR4 SLiMs, along with the newly identified REER-repeat region, might be involved in RNA binding. To test this hypothesis, we conducted electrophoretic mobility shift assays (EMSA) to examine the binding of purified RNase E CTD proteins to an in vitro synthesized RNA derived from the *malEF* intergenic region (for details, see methods). We tested wild-type (WT) CTD and mutated variants where arginine residues in AR1, AR4, or the REER-repeats were

replaced with alanine (named AR1mut, AR4mut, and REERmut, respectively). Qualitative analysis of

the EMSA showed that the WT, AR1mut, and AR4mut variants bound RNA with similar efficiency, as

indicated by comparable gel shift patterns (Fig. S4A). However, the REERmut variant exhibited slightly

reduced RNA binding, evidenced by a delayed shift as compared to the former proteins (Fig. S4B and

2A). The combination of AR1 and AR4 mutations resulted in an impaired RNA binding which was

further exacerbated when the REER mutation was also introduced: no observable shift in the RNA was

detected when incubated with the protein variant carrying mutations in all three SLiMs, even at an

RNA-to-protein ratio of 1:80 (Fig. 2A). Together, these results highlight the functional redundancy of

the AR1, AR4, and REER-repeat SLiMs in RNA binding, as a significant defect in RNA binding is

observed only when all these SLiMs are simultaneously disrupted.

One notable characteristic of the RNA degradosome is its tendency to form highly dynamic subcellular

foci, also referred to as "clusters," "puncta," or "helical structures" [28, 61]. These foci rapidly

disassemble upon rifampicin treatment, suggesting their formation is RNA-dependent [61, 62]. In *E.*

coli, these foci are membrane-anchored via the MTS SLiM, whereas in *C. crescentus* they remain

cytoplasmic, driven by liquid-liquid phase separation [62, 63]. To investigate the subcellular localisation

of Pa RNase E, we constructed *P. aeruginosa* PAO1 strains expressing msfGFP-tagged Pa RNase E, Pa

RNase E$^{1-588}$, or Pa RNase E$^{1-529}$ from the native *rne* locus. The two RNase E truncations were designed

based on our in-silico analysis of Pa RNase E SLiMs, assuming that they would lack the capacity to

assemble the RNA degradosome. The key difference between RNase E$^{1-588}$ and RNase E$^{1-529}$ lies in the

presence of the MTS (Fig. 1B). Visualization using epifluorescence microscopy revealed that RNase E-

msfGFP forms foci concentrated near the cell edge (Fig 2B). Both CTD truncation mutants showed

impairment in foci assembly, but while the RNase E$^{1-588}$-msfGFP protein localises uniformly around the

edge of the bacteria, the RNase E$^{1-529}$-msfGFP mutant protein additionally fails to localise near the

membrane, consistent with the role of the MTS in membrane anchoring (Fig. 2B).

Exposure of Pa RNase E-msfGFP expressing cells to high levels of rifampicin (100 µg/ml) led to the

dissipation of these foci, resulting in a smoother distribution of the signal along the cell periphery, which

phenocopies the localization of the RNase E$^{1-588}$-msfGFP strain (Fig. 2D). Quantitative analysis

confirmed a significant reduction in signal intensity variance upon rifampicin treatment as compared to

the control (DMSO-only), indicating foci dissipation, while the average signal intensity remained

constant, ruling out RNase E- msfGFP proteolysis (Fig. 2E-F, Fig. S4F).

To further verify that Pa RNase E-msfGFP foci formation depends on CTD-mediated RNA binding, we

constructed *P. aeruginosa* strains with chromosomally msfGFP-tagged RNase E variants mutated in

AR1, AR4, and REER-repeats SLiMs, individually and in combination, as done previously for the

EMSA. Visualisation of these variants showed a clear correlation between CTD-mediated RNA binding

and foci formation. Specifically, the RNase E AR1+AR4mut-msfGFP showed less distinct foci and

lower signal variance compared to RNase E-msfGFP and single AR1 or AR4 mutants (Fig. 2B-C). The

RNase E REERmut-msfGFP also had diffuse foci and lower variance, while the RNase E

AR1+AR4+REERmut-msfGFP, with abolished RNA binding, displayed the most severe disruption,

with nearly complete loss of foci and a very narrow signal intensity distribution (Fig. 2B-C and Fig.

S4F). Similar patterns were observed with CTD truncations containing different combinations of RNA-

binding SLiMs, ruling out the possibility that this foci dissipation is caused by the changed in the charge

of the CTD conferred by the mutations (Fig. S4E).

Of note, we noticed an increased signal intensity in all the msfGFP-tagged RNase E mutants tested as

compared to the RNase E-msfGFP (Fig. S4C). Since all constructs are expressed at the native locus,

this could likely reflect higher RNase E protein levels, which suggests that CTD-mediated RNA binding

might be necessary for RNase E autoregulation [64]. To confirm this, we performed a Western Blot on

strains expressing several Strep-tagged RNase E mutants and could indeed observe that RNase E protein

levels are increased upon deletion of the CTD or introduction of AR1+AR4+REERmut (Fig. S4D).

Overall, we conclude that in *P. aeruginosa*, the organisation of the RNA degradosome complex into

visualisable foci is dependent on the RNase E CTD-mediated RNA binding and suggest that this

interaction not only organizes the localisation of the RNA degradosome but could also influence the

regulatory activity of RNase E.

Identification of proteins binding to RNase E scaffold domain

In *P. aeruginosa*, the RNA degradosome was proposed to include the RhlE2 RNA helicase (called

DeaD), PNPase, and other associated proteins [54, 55]. However, the direct nature of these interactions

and the specific interaction sites have not been demonstrated nor mapped. Additionally, RNase R was

also suggested to be a *P. aeruginosa* RNA degradosome component, as it is in *P. syringae* LZ4W [30,

40]. To test these hypotheses and identify Pa RNase E CTD interacting proteins, we performed a pull-

down assay using *P. aeruginosa* strains expressing C-terminal Strep II-tagged full-length RNase E

(RNase E$^{1-1057}$-Strep) or an RNase E truncation missing the entire CTD (RNase E$^{1-529}$-STREP). As an

additional control, the assay was also performed with the *P. aeruginosa* PAO1 wild-type (WT) to discard

proteins interacting nonspecifically with the Strep-Tactin magnetic beads (Fig. 3A-B).

Eluates of the pull-down experiment were first analysed by silver staining (Fig S5). The RNase E$^{1-1057}$-

Strep eluate revealed a high molecular weight band corresponding to RNase E, along with additional

bands absent in the untagged and RNase E$^{1-529}$-Strep samples. These bands likely represent components

of the RNA degradosome or cleaved RNase E$^{1-1057}$-Strep intermediates, as RNase E is known to be

prone to cleavage [42, 65]. Mass spectrometry (LC-MS/MS) analysis identified a total of 115 proteins

present in both duplicates of the RNase E$^{1-1057}$-Strep eluates. 13 proteins (including RNase E) were

enriched by at least 5.5-fold as compared with the untagged sample (Fig 3A). This stringent threshold

was chosen since core components of the RNA degradosome should be abundantly associated to RNase

E. Of these 12 putative partners, 5 (PNPase, GdhB, ArcB, RhlB and RpoB) were completely absent

from both RNase E$^{1-529}$-Strep duplicates and 2, namely RpoC and RplL, were absent from one replicate

and present in the second replicate with a peptide count of 2. We selected for further study the first 5

proteins, which were considered the most promising RNA degradosome candidates as their interaction

with Pa RNase E seemed fully dependent on the CTD.

It is worth noting that the RhlE2 RNA helicase was, as expected [54, 55], among the proteins enriched

in the RNase E$^{1-1057}$-Strep eluate as compared to the untagged control, but surprisingly it was equally

enriched in the RNase E$^{1-529}$-Strep eluate condition (Fig 1C), indicating that this interaction occurs via

RNase E NTD and not via the CTD.

RNase E direct interaction with PNPase and RhlB occur via the NDPR and AR1 SLiMs,

respectively

The interaction between RNase E and the 5 candidates was first verified using an adenylate-cyclase-

based bacterial two-hybrid (BTH) assay [66]. Co-expression of T25-PNPase and T25-RhlB with T18-

RNase E resulted in red colonies on MacConkey agar and a significant increase of β-galactosidase

activity compared to empty vector controls (11-fold or 5-fold increase, respectively), confirming their

interaction (Fig S6A, Fig. 3C). Co-expression of T25-ArcB with T18-RNase E also resulted in a

significant increase in β-galactosidase activity (5-fold) compared to the negative control (Fig. 3C, Fig.

S6A). To identify the SLiMs mediating the observed interactions, we generated seven CTD truncations

of RNase E, each lacking an additional SLiM as compared with the longer version. Testing these

truncations for their capacity to interact with PNPase revealed that only T18-RNase E$^{1-1025}$ showed β-

galactosidase activity similar to those of T18-RNase E$^{1-1057}$ upon co-expression with T25-PNPase (Fig.

3D, Fig. S6B). In contrast, all shorter T18-RNase E CTD truncations except T18-RNase E$^{1-529}$, and also

T18-RNase E$^{940-1057}$, exhibited a significant increase of β-galactosidase activity as compared with the

negative controls but lower than T18-RNase E$^{1-1057}$ (Fig. 3D, Fig. S6B). Overall, this suggests that

PNPase interaction with RNase E relies on two distinct sites of RNase E CTD: residues 529-588 and

940-1057. Co-expression of the T18-RNase E CTD truncations with T25-RhlB revealed that the

protein-protein interaction is mediated by residues 733-793: no significant interaction could be seen

between T25-RhlB and T18-RNase E$^{1-733}$, although T25-RhlB could interact with T18-RNase E$^{1-793}$ (and

every longer CTD truncation) or with T18-RNase E$^{733-793}$ about as strongly as with T18-RNase E$^{1-1057}$

(Fig. 3E, Fig. S6C). Of note, neither T18-RNase E$^{733-761}$ nor T18-RNase E$^{762-793}$ were able to restore β-

galactosidase activity above negative control levels when co-expressed with T25-RhlB (Fig. 3E),

suggesting that the entire AR1 SLiM, as defined in Fig 1B, is necessary for interaction with RhlB.

Surprisingly, all T18-RNase E CTD truncations, except T18-RNase E$^{1-529}$, significantly interact with

T25-ArcB, suggesting a complex binding that is dependent on residues 529-588 (Fig. 3F, Fig. S6D). Of

note, these residues only contain the MTS SLiM, which is presumably involved in membrane binding.

Building on the bacterial two-hybrid assays results, we successively employed affinity chromatography

with purified proteins to investigate direct binding, minimizing the influence of other protein partners

or RNA. Fusion proteins containing a C-terminal 3xFlag tagged RNase E$^{1-529}$, RNase E$^{572-1057}$ (CTD

with no MTS) or RNase E$^{1-1057}$ were purified and immobilized on anti-Flag magnetic beads. After

extensive washing, the beads were mixed with either purified PNPase, RhlB, or ArcB proteins. Analysis

of the bound and free fraction was performed by SDS-page (Fig. 4).

PNPase was retained on beads containing Flag-RNase $E^{572-1057}$, but not on beads alone or containing

Flag-RNase E^{1-529} (Fig. 4A), indicating that the RNase E CTD is sufficient for direct interaction with

PNPase. Point mutations within the NDPR SLiM (Flag-RNase $E^{572-1057}$_NDPRmut) confirmed that this

SLiM is necessary for this interaction, as suggested by the observed interaction between T18-RNase

$E^{940-1057}$ truncation and T25-PNPase in the BTH assay (Fig. 4A).

Similarly, RhlB was retained on Flag-RNase $E^{572-1057}$ but not on Flag-RNase E^{1-529} and point mutations

in the AR1 SLiM (Flag-RNase $E^{572-1057}$–AR1mut) showed that this region is crucial for RhlB binding

(Fig. 4B).

Contrary to PNPase or RhlB, ArcB was not retained on beads containing Flag-RNase E^{1-1057} under the

tested conditions, suggesting an indirect type of interaction (data not shown). To summarize, these

experiments clearly establish PNPase and RhlB as core components of *P. aeruginosa* RNA

degradosome and demonstrate that the AR1 and NDPR SLiMs are essential for their direct interactions

with RNase E.

PNPase and RhlB are dependent on their physical interaction with RNase E for colocalising in

RNA degradosome foci.

Given the foci localization of RNase E, we assessed PNPase and RhlB colocalization with RNase E by

using *P. aeruginosa* strains carrying *rne::mCherry* and either *pnp::msfGFP* or *rhl::msfGFP*

chromosomal fusions. As expected, both PNPase-msfGFP and RhlB-msfGFP formed foci near the

membrane and co-localized with RNase E-mCherry in live cells (Fig. 5A). Overlap coefficients between

the two fluorescence channels were consistently over 0.9, while Pearson's coefficients were slightly

lower (0.880 for PNPase and 0.773 for RhlB), likely due to differences in signal intensity. Of note,

strains expressing any of these chromosomal fusion proteins grow identically to the WT strain at 37°C,

suggesting that all the fusion proteins are functional (Figure S7A). We also assessed the subcellular

localisation of ArcB-msfGFP and observed a smooth distribution rather than foci, which suggests that

the interaction between RNase E and ArcB is likely transient, with ArcB not being a core component of

the complex (data not shown).

To further corroborate the need for specific RNase E domains in these interactions, we examined the

localization of PNPase-msfGFP and RhlB-msfGFP in strains expressing RNase E^{1-1057} (WT), RNase E^{1-}

^{529} (*rne529*), RNase E^{AR1mut} (*rne*AR1^{mut}), or RNase $E^{NDPRmut}$ (*rne*NDPR^{mut}). Both PNPase-msfGFP and

RhlB-msfGFP displayed diffuse cytosolic localization in the *rne529* strain compared to the WT strain

(Fig. 5B). In the *rne*NDPR^{mut} strain, PNPase-msfGFP showed a diffuse cytosolic distribution, while

RhlB-msfGFP was diffuse in the *rne*AR1^{mut} strain. However, NDPR or AR1 mutations did not prevent

RhlB or PNPase from being recruited into RNase E foci, respectively, indicating that both proteins can

be independently recruited into RNase E-driven foci (Fig. 5B). We also confirmed that RNase E

NDPRmut-msfGFP forms foci, ruling out the possibility that the loss of PNPase foci in the *rne*NDPR^mut

strain is due to impairment of RNase E-driven foci formation (Figure S7B). These findings confirm the

in vivo interaction of Pa PNPase and Pa RhlB with the NDPR and AR1 SLiMs, respectively, and

demonstrate that these interactions are crucial for their near-membrane foci localization.

Phenotypic changes associated with partial or complete deletion of RNase E CTD

To investigate the biological importance of RNase E CTD-mediated complex scaffolding in *P.*

aeruginosa, we performed some phenotypic assays using the *rne529* and *rne588* strains, as well as

strains bearing the AR1 mutation (*rne*AR1^mut), NDPR mutation (*rne*NDPR^mut), or AR1+AR4+REER

mutation (*rne*AR1+AR4+REER^mut), abolishing RhlB binding, PNPase binding, or CTD-mediated RNA

binding, respectively.

At 37°C in liquid culture, only the *rne529, rne588* and *rne*AR1+AR4+REER^mut strains exhibited a slight

but consistent growth delay (Fig. 6B). On solid media at 37°C, the *rne529* and *rne*AR1+AR4+REER^mut

strains also displayed reduced colony size, confirming the growth delay (Fig. 6A, top panel). To verify

that the growth delay was caused by truncations/mutations introduced in the *rne* gene, we electroporated

the WT or the two mutant strains with an inducible vector allowing in trans expression of RNase E,

RNase E-msfGFP, or RNase E-mCherry. Cultivation of the strains at 37°C in liquid culture revealed

that expression of either native or fluorescently-tagged RNase E could restore identical growth levels

compared to the electroporated WT strain, additionally confirming that the fluorescently-tagged RNase

E protein is fully functional (Fig S10).

Unexpectedly, when cultured on solid media at 16°C, the *rne588* strain showed impaired growth, while

the growth defects of the *rne529* and *rne*AR1+AR4+REER^mut strains became more pronounced, with

rne529 barely growing (Fig. 6A, upper middle panel). Since the *rne*AR1^mut and the *rne*NDPR^mut strains

exhibited no significant differences compared to the WT, these results indicate that the cold sensitivity

of the *rne588* strain can be attributed to the loss of RNase E CTD RNA binding. Moreover, the severe

growth defect observed in the *rne529* mutant indicates that additional delocalisation of RNase E from

the membrane exacerbates the cold sensitivity caused by deletion of RNase E RNA-binding SLiMs.

Besides low temperature, we also tested sensitivity to low or high pH (Fig. 6A, lower panels): while

growth at pH≃9.5 is similar between the different strains, deletion of the CTD (*rne529* and *rne588*)

impairs growth at pH≃5.5, although the three CTD point mutant strains are mostly unaffected. This

suggests that, although loss of interaction with PNPase or RhlB separately does not seem to impair

growth, the combined loss of RNase E protein-protein and protein-RNA interactions results in a growth

defect under low pH conditions. Finally, we assessed whether the RNase E CTD mutants might be

impaired in virulence using *Galleria mellonella* larvae, a highly predictive infection model for studying

mammalian infection processes [67]. Infection with PAO1 WT, *rne*AR1^mut and *rne*NDPR^mut strains

resulted in 100% larval mortality within 20 hours post-injection (Fig 6C). In contrast, the larvae infected

with the *rne588* mutant exhibited 50% survival at this time point, while those infected with

*rne*AR1+AR4+REER^{mut} or *rne529* mutants showed 100% survival, with larval death occurring only

after 21 hours post-injection (Fig 6C). These results strongly indicate that the ability of *P. aeruginosa*

to infect a host is compromised when RNase E CTD-mediated RNA binding is disrupted. This

impairment is likely attributable to several contributing elements: reduced bacterial growth within the

host, deficiencies in virulence factor production, and a decreased ability to endure host immune

responses and host-related stresses.

Effect of RNase E CTD on *P. aeruginosa* transcripts steady-state levels

To better understand the phenotypes observed and evaluate the impact of RNase E CTD loss on *P.*

aeruginosa transcriptome, we conducted RNA-sequencing analysis on *P. aeruginosa* WT, *rne529* and

rne588 mutant strains grown at 37°C until the late-exponential growth phase (OD_{600nm} of 1-1.2). Under

these conditions, the loss of the entire RNase E CTD in the *rne529* strain affected the steady-state levels

of 450 transcripts, with 183 downregulated and 267 upregulated (P-value < 0.05, fold change $|FC| \geq 2$)

(Fig. 7A). In the *rne588* mutant, 417 transcripts were differentially expressed, with 194 downregulated

and 223 upregulated (P-value < 0.05, fold change $|FC| \geq 2$) (Fig. 7B). Differential expression analysis

on reads mapping to predicted small RNA genes (sRNAs) identified 14 predicted sRNA genes with

significantly enriched reads in the *rne529* mutant and 5 in the *rne588* strain (Table S3 and Figure S).

A comparison of the *rne529* and *rne588* strains revealed that 164 mRNAs were consistently regulated

in both mutants (97 upregulated and 67 downregulated, see Figure 7C). Among the strongest commonly

downregulated genes were: the *pqsABCD* genes, involved in the synthesis of quorum-sensing signal

molecules alkylquinolones PQS (2-heptyl-3-hydroxy-4(1H)-quinolone) and HHQ (2-heptyl-4(1H)-

quinolone) and in virulence (with the entire *pqsE*, *pqsH* and *phnAB* PQS synthesis cluster being

downregulated in *rne529* strain) [68], the *phzA1B1* and *phzA2B2* phenazine biosynthesis genes, which

are important for synthesis of the toxic secondary metabolites phenazine and pyocyanin [69], and the

agtABCD gene cluster for 4-aminobutyrate (GABA) and 5-aminovalerate (AMV) uptake [70].

Surprisingly, the two strains exhibited more differences in mRNA levels than shared changes. In the

rne588 strain, the mRNA levels corresponding to type 3 secretion system genes, which deliver toxins

to host cells [71], were consistently upregulated, while they did not vary significantly in the *rne529*

strain as compared to the WT (Fig 7D). On the other hand, ribosomal and many amino acid metabolism

genes (tryptophan, arginine, alanine, aspartate and glutamate) were downregulated only in the *rne529*

strain, which could correlate with the general growth defect observed (Fig. 7E and Table S4).

Considering the autoregulation of RNase E reported in *E. coli* and *C. crescentus* [64, 72], we quantified

reads mapping to the 5' untranslated region (UTR) of the *rne* gene (TSS determined in [73]) and

observed increases of 2 and 0.96 $\log_2$ folds in the *rne529* and *rne588* mutants, respectively, compared

to the WT (Fig. 7F). Similarly, we observed a 1.43 and 0.88 $\log_2$ fold increase in the transcript

corresponding to the NTD of RNase E in the *rne529* and *rne588* mutants, respectively (Fig. 7F), in

agreement with the observed increase in protein levels (Fig. S4C-D).

Finally, to assess RNA degradation efficiency in our strains, we analysed the 5' reads ends of sequenced

RNA fragments. As previously performed in other studies [56, 74], we calculated the ratio between the

number of loci with 5' ends present only in a mutant and not in the WT, over the sum of loci with 5'

read ends seen in both mutant and WT. We performed the analysis in the three genetic backgrounds

(WT, *rne529*, *rne588*) and for each replica independently to ensure consistency (see methods and Fig

S9). Despite the background of random fragmentation intrinsic to the sequencing protocol, a consistent

increase on the number of 5' ends in the *rne529* and *rne588* strains was observed as compared to the

WT (Fig. 7G), suggesting an accumulation of RNA degradation intermediates and an altered RNA

degradation in the RNase E truncated mutants.

Discussion

The composition and the functions of the RNA degradosome vary among bacterial species, likely

reflecting diverse lifestyles and niche adaptations [59]. Studying the variations of the RNA

degradosome composition in various bacterial species is therefore crucial for elucidating how bacteria

respond and adapt to different environmental conditions and sustain various types of stress. In this study,

we characterized the RNA degradosome of *P. aeruginosa*, a versatile opportunistic pathogen and widely

used model organism for bacterial pathogenesis and environmental adaptation.

In the first part of our study, we focused on the putative SLiMs within the CTD of *P. aeruginosa* RNase

E and assessed their contribution to RNA binding, as well as to binding the other core RNA

degradosome components, which we identified as RhlB and PNPase via pull-down assays (Fig. 8). A

notable feature of SLiMs is their poor sequence conservation, even when they mediate interactions with

the same protein partner. Our findings confirm this variability for both PNPase and RhlB interactions.

PNPase, a known partner of RNase E in several species, binds through SLiMs with significantly

different sequences across organisms. For instance, in *Anabaena* PCC7120, the interaction occurs via a

cyanobacterial SLiM having the conserved residues RRRRRRSSA [37], while in *E. coli* or *C.

crescentus*, the sequences are WQRPTFAFEGKGAAGGHTATHHASA [30] and

APPEKPRRGWWRR [43], respectively. Our pull-down assays and in vitro binding assays showed that

PNPase interacts with RNase E in *P. aeruginosa* via the NDPR SLiM, which has yet another distinct

sequence (TGRALNDPREKRRLQREAERLAREAAAAA). These sequence variations may reflect

species-specific interactions, as suggested by experimental work showing lack of heterologous

interactions between RNase E and PNPase of *E. coli* and *Pseudoalteromonas haloplanktis* [42]. Despite

sequence variability, the PNPase interacting SLiM is consistently located towards the end of the RNase

E CTD [30, 37, 43], suggesting convergent evolution in the positioning of the PNPase relative to other

RNA degradosome components and to RNase E catalytic domain.

Of note, a recent study on a *P. aeruginosa* mutant with a 50 bp deletion in the 3' end of the *rne* gene
revealed a hypervirulence phenotype, characterized by enhanced cytotoxicity and siderophore
production [75]. Based on *E. coli* RNA degradosome model, this phenotype was associated with a
possible impairment of PNPase binding to Pa RNase E or a different RNA degradosome activity [75].
Our data reveal that PNPase can interact with the RNase E$^{1-1025}$ truncation (BTH assay), ruling out the
former hypothesis. However, we cannot exclude the possibility that this region might be involved in
interaction with a yet unknown partner.

Regarding RhlB, we show that in *P. aeruginosa*, this RNA helicase directly interacts with RNase E via
the AR1 SLiM (RPRRRSRGQRRRSNRRERQR). Mutation of AR1 disrupts not only the RhlB-RNase
E binding *in vitro*, but also RhlB subcellular localisation near the membrane and co-localization with
RNase E *in vivo*. Interestingly, in *E. coli* and other species the AR1 SLiM mediates RNA interaction
[30, 42]. In *P. aeruginosa*, AR1 also contributes to RNA-binding in addition to RhlB binding. Given
that in *E. coli,* a single arginine mutation within the HBS SLiM (R730A) is sufficient to disrupt RhlB
binding [76], it is possible that only one or a few arginine residues of the AR1 SLiM are involved in
interaction with RhlB, while the others mediate RNA binding. Further experiments are needed to
address this hypothesis and to understand the dynamics of the RhlB-RNA-RNase E complex in *P.*
aeruginosa. Moreover, it has been shown that the RNA-dependent ATPase and unwinding activity of
RhlB are stimulated by binding to RNase E [77]. Future studies will investigate whether it is also the
case in *P. aeruginosa*.

In contrast with the *E. coli* RNA degradosome, we did not find any evidence of an interaction between
RNase E and enolase or another metabolic enzyme in *P. aeruginosa* when the bacteria grow in rich
medium and standard laboratory conditions. Although our pull-down and bacterial-two hybrid data
indicate a probable interaction between RNase E and ArcB, a catabolic ornithine carbamoyltransferase
from the arginine deaminase pathway [78], no direct protein-protein binding could be observed in vitro,
nor were the two proteins colocalising in vivo. We propose that RNase E indirectly interacts with ArcB,
probably in a RNA-dependent manner. Indeed, the *arcDABC* polycistronic transcript is cleaved at
multiple sites by RNase E for its processing [79], suggesting that an regulatory interaction between
RNase E and ArcB polypeptide could occur during co-translational RNA cleavage [18].

Additionally, the RNase E-based RNA degradosome of *P. haloplanktis, P. syringae*, and *R. capsulatus*
do not seem to contain either a metabolic enzyme upon standard growing conditions [40-42]. This lack
of RNase E interaction with metabolic enzymes may reflect distinct physiological roles of the RNA
degradosome in different bacterial species, potentially linked to specific metabolic adaptations or
regulatory mechanisms that modulate RNA stability and processing depending on nutrient availability.
However, we cannot exclude that some protein partners were missed. Transient or condition-specific
accessory protein partners remain to be identified, and compositional biases such as the conserved
AEPV-rich part of the RNase E CTD are not yet understood, as no ligand or biological function has
been assigned to this region even in well-studied RNase E orthologs.

It is also important to mention that throughout this study, we chose to focus on core RNA degradosome

components that are identified based on their binding to RNase E CTD, but we cannot exclude that

some functional interactions occur with the NTD. For example, RhlB was found to interact with RNase

E NTD in *C. crescentus* [36], while in *Anabaena* the RNA helicase CrhB, RNase II and enolase were

shown to associate with RNase E via its NTD [38, 39]. Similarly, we found that the *P. aeruginosa*

RhlE2 RNA helicase, previously shown to associate with RNase E in an RNA-dependent manner [55],

interacts with the N-terminal domain of Pa RNase E. Although RhlE2 is not considered a core RNA

degradosome component based on our assumption of CTD-mediated interactions, this association

remains functionally significant [55, 56].

In the second part of our study, we evaluated the importance of RNA degradosome assembly at both

transcriptomic and phenotypic levels through an analysis of *P. aeruginosa* strains carrying RNase E

CTD truncations and mutations. Comparing the *rne588* and *rne529* strains allows us to discuss the

importance of RNA degradosome assembly and membrane attachment in *P. aeruginosa*. The two

mutants exhibited varying degrees of sensitivity to cold growth, with *rne529* being the most affected,

indicating that cold growth requires both RNase E membrane attachment and RNA degradosome

assembly. Loss of the CTD without delocalisation from the membrane (*rne588*) does not affect growth

in *P. aeruginosa* under standard laboratory condition (i.e., 37°C and rich medium), but impacts its

capacity to sustain growth in the cold. Transcripts levels were also generally affected in the mutants, as

well as global RNA degradation, as shown in the 5' read ends analysis. Decreased degradation of

cleaved transcripts could lead to an upregulation of their levels. Therefore, direct and indirect effects

remain to be distinguished and those changes remain to be correlated with proteins levels.

Surprisingly, we did not find any phenotypic changes associated to the loss of RhlB or PNPase binding,

questioning the importance of the RNA degradosome protein association for the complex functionality.

Competition assays could be useful in identifying any fitness defects in our mutants, as done in *E. coli*

for RNase E CTD truncations [19, 63], or different phenotypes or stresses could be tested.

On the other hand, our phenotypic analysis highlights the crucial role of CTD RNA binding for the

functionality of the RNA degradosome complex. Mutations in the SLiMs responsible for RNA binding

(AR1, AR4, and the REER-repeats) not only impair in vivo RNA degradosome foci assembly and/or

stability but also lead to decreased *P. aeruginosa* virulence and reduced growth at both 37°C and cold

temperatures. The effect of these mutations on foci formation mirrors the effect observed upon cell

exposure to rifampicin, which decreases RNA levels in the cell, and aligns with previous studies

showing that Ec and Cc RNase E orthologs also assemble foci in a CTD-dependent and RNA-dependent

manner [61-63]. Our data thus suggest that impairing foci formation has a major effect on the activity

of the RNA degradosome in *P. aeruginosa*.

Overall, our study identified core RNA degradosome components in *P. aeruginosa* and underscores the

role of this complex in bacterial adaptation to environmental changes and pathogenesis. By revealing

P. aeruginosa-specific interactions and their functional significance, our findings provide valuable

insights into the molecular mechanisms of adaptation in this pathogen and highlight the potential of the
RNA degradosome as a drug target.
484
485 **Methods**
486
Bacterial strains and growing conditions
A list of bacterial strains used can be found in Table S1. Unless otherwise specified, *P. aeruginosa* PAO1
was grown at 37°C in NYB (25 g of Difco nutrient broth, 5g of Difco yeast agar per liter) with shaking
at 180 rpm or on Nutrient Agar (NA) (40 g Oxoid blood agar base, 5 g Difco yeast agar per liter). When
required, antibiotics were added to these media at the following concentrations: 100 μg/ml ampicillin,
50 μg/mL kanamycin, 20 μg/mL tetracycline and 10 μg/ml gentamicin for *E. coli*; and 50 μg/mL or 100
μg/mL tetracycline and 50 μg/ml gentamicin for *P. aeruginosa*.
494
Construction of plasmids used in this study
Information about plasmid construction and primers used in this study can be found in supplementary
tables S2 and S3, respectively. Plasmids pUT18C and pKT25 were generally cloned by restriction
enzyme digestion using EcoRI-HF and BamHI-HF (NEB). When a gene to be inserted could be
internally cleaved by EcoRI and/or BamHI, the enzymes MfeI and/or BglII, respectively, were chosen
instead during primer design to allow ligation of the insert to the vector digested with EcoRI/BamHI
thanks to compatible cohesive ends. For cloning of *gdhB* into pKT25, NEB assembly® was used instead,
due to the presence of many restriction enzymes internal cleavage sites. Ligation reactions were carried
our using T4 DNA ligase (Roche or NEB) at 16°C overnight, or at room temperature for 2-3 hours,
following standard protocols. When cloning inserts into the suicide vectors pME3087 or pEXG2, the
NEBuilder HiFi DNA assembly Master Mix (NEB) was used according to the manufacturer's
recommendations. To constructs RNase E CTD mutants, the DNA was synthesized by GeneArt
(ThermoFisher Scientific) and cloned into appropriate vectors for protein purification or strain
construction as described in Table S4. To construct pME6032-based expression vectors, restriction
enzyme digestion was carried out using EcoRI and KpnI for both vector and inserts.
Ligation mixtures or the assembly reactions were transformed into chemically competent *E. coli* DH5α
cells by heat-shock, plating on appropriate selective LB agar medium. The resulting colonies were
screened by PCR for the presence of the insert in the plasmid of interest. After plasmid extraction
(Miniprep kit, Sigma-Aldrich/Merck), plasmids were verified by Sanger sequencing at Microsynth.
514
Construction of *P. aeruginosa* mutant strains
To construct chromosomal gene truncations or insert a Strep-tag or fluorescent protein into the
chromosome, in frame with a gene of interest, the suicide vectors pEXG2 [80] or pME3087 [81] were
used. Briefly, approximately 500 bp regions located immediately upstream or downstream of the region

to be edited, and, when applicable, the tag or fluorescent protein of interest, were amplified by PCR and
inserted together into the vector backbone using NEB assembly®. After sequencing verification of the
constructed plasmid quality, the plasmids were introduced into *P. aeruginosa* PAO1 strain by triparental
mating using the *E. coli* strain HB101 (pRK2013). After selection with the appropriate antibiotic
(tetracycline for pME3087 or gentamicin for pEXG2), merodiploids were resolved as described in [82]
for pME3087 or [80] for pEXG2, and strains with the desired insertion/deletion were identified by
colony PCR.

RNase E sequence analysis
Group-specific homologs of RNase E in *P. aeruginosa*, *P. fluorescens*, *P. protegens*, *P. putida* and *P.*
stutzeri were retrieved from NCBI databases using blastp [92] with *P. aeruginosa* PAO1 RNase E as a
query; only sequences with query coverage > 0.5 were conserved (total of 5000 proteins). The number
of RE[E/D]R motifs in each protein sequence was determined using the vcountPattern function from
the "biostrings" R package [93]. The similarity of the protein sequences within a taxonomic group was
determined using the conserv function (method="similarity") from the bio3D R package on a muscle
alignment of the protein sequences[94, 95].

Electrophoretic motility shift assay
The *malEF* transcript was synthesized with a T7 transcription kit (Invitrogen) on linear DNA template
(172 bp) obtained by PCR with primers p253-p254 (Table Sx). The electrophoretic mobility shift assay
(EMSA) was performed following the protocol of [86]. Briefly, the reaction mixtures (15 μl) contained
35 nM of RNA, 50mM Tris-HCl, 100mM NaCl, 5mM DTT, 1.0 units/μl RNasin (Promega), and
increasing concentrations of the His10Smt3-RNase E proteins (see Figure legends). The reactions were
incubated at 4 °C for 30 min before loading on 2% agarose gel and run at 150 V for 15 min. RNA bands
were visualized by gel staining with SybrGold (Invitrogen).

Epifluorescence microscopy imaging
Phase contrast and fluorescence microscopy were performed on a Zeiss Axio Imager 2 equipped with
an objective alpha Plan-Apochromat 100x/1,46 Oil Ph3 M27 and a ZEISS Axiocam 305 mono CCD
camera. Bacteria were grown until OD_{600nm}= 0.2-0.6 (approx. 2.5 hours) in 3 ml of NYB and were then
placed on an agarose pad containing 1% agarose dissolved in 0.5X PBS. Images were processed with
Fiji (NIH, USA) [84].

Image analysis
All image treatments and analysis were performed using Fiji (NIH, USA)[84]. Representative images
shown had contrast adjusted for display by opening the images in Fiji and using the Reset function in
the Adjust Brightness/Contrast menu, after average background intensity subtraction using the Math ->

Subtract function. These average background intensity values were obtained by measuring a square area
devoid of any cells. For quantification of signal intensity, a line was drawn alongside the long axis of
bacterial cells and signal was measured using Plot Profile. Results of this function were saved as .csv
files. For each strain, at least 50 bacterial cells were measured from the raw images, using at least two
different images for quantification. Average background values were subsequently subtracted from each
results file using a .csv file containing the Plot Profile measurements of a linear area devoid of cells for
each image used. Fluorescence intensity was measured along the longitudinal axis of the cells in 50-
100 cells using Fiji. Boxplots of fluorescence intensity and signal variance were plotted using R [85].
The signal variance was obtained by calculating the variance of the $\log_2$ fold-change between each
measured value and the average of all values for a given measured cell. This was done to normalize the
data and obtain comparable variances without introducing a bias due to differences in overall signal
intensity.
568

Pull-down identification of RNase E partners
Fresh cultures of PAO1 WT, PAO1 RNase E_{1-529}-Strep, or PAO1 RNase E_{1-1057}-Strep were inoculated to
OD=0.05 from overnight cultures in 500 mL Erlenmeyer flasks. Cultures were grown until approx.
OD_{600nm}=0.8-0.9. 50 mL of each culture was harvested by centrifugation at 4500 $x\ g$ for 10 minutes.
Pellets were stored at -80°C until needed.
Each thawed pellet was then resuspended in 3,25 mL of lysis buffer (for 10 mL of buffer, prepared just
before use: 20 mM Tris (pH 10), 250 mM NaCl, 10% v/v glycerol, 1 EDTA-free protease inhibitor
tablet, 50 μg/mL lysozyme). Samples were incubated the samples on ice for about 30 minutes,
transferred into 15-mL Falcon tubes and sonicated using a thin probe for 6x30s, with 15 seconds break
in between. A pulse of 70% and a power output of 50% were used. After sonication, 32.5 μL of Triton
X100 10% were added to each sample (~0.1% final) before incubation on a rotary wheel at 4°C for 8
minutes. The samples were transferred into 2 mL Eppendorf tubes and centrifuged for 1 hour at 10500
$x\ g$ at 4°C. MagStrep® Strep-Tactin®XT beads (IBA, cat# 2-5090) were washed in wash buffer (20 mM
Tris (pH 10), 250 mM NaCl, 10% v/v glycerol) according to the manufacturer's instructions. One mix
of beads was prepared for all the samples with a volume of beads suspension of 45 μL per sample. Final
resuspension was done in 200 μL wash buffer per sample, plus an additional 10 μL. After centrifugation
of the samples, the supernatant of each sample was regrouped into a 5 mL Eppendorf tube, and 200 μL
of washed beads suspension were added per sample. Incubation on a rotary wheel was done for 2h15 at
4°C. The 5 mL tubes were then placed against a magnetic rack and all the supernatant was carefully
removed. A first wash volume of 500 μL was added gently and agitated by horizontal rotations of the
tubes. Tubes were placed back against the magnetic rack and the wash volume was removed. A second
wash step was performed identically with a volume of 225 μL instead of 500 μL. A third wash step of
225 μL was also performed, and the resuspended beads were transferred into a clean 1.5 mL Eppendorf
tube by gentle pipetting with a 1000 μL tip, before removing the wash volume. The beads were covered

with 10 μL of sterile PBS and stored at 4°C before LC-MS/MS analysis at the Proteomics Core Facility

(University of Geneva). For plots in Fig. 3A-B, the log2 fold-change (log2FC) was calculated using the

following formula: log2(average peptide count of protein x identified in the RNase E^{1-xxx}-Strep / average

peptide count of protein x identified in the untagged sample).When an average count of 0 peptide was

identified in the untagged sample, the log2FC value was instead calculated using the following formula:

log2FC(average peptide count of protein x identified in the RNase E^{1-xxx}-Strep/1).

Bacterial adenylate cyclase two-hybrid assay

Cloned plasmids were introduced into chemically competent BTH101 cells by heat shock, and the cells

were plated in LB agar supplemented with ampicillin and kanamycin. Isolated colonies were then

streaked as a line of about 1-2 cm on freshly prepared Mac Conkey agar medium supplemented with

ampicillin, kanamycin, 0.5 mM isopropyl β-d-thiogalactoside (IPTG), and maltose 1%. The

MacConkey Petri dishes were then incubated at 30°C for 24h followed by an incubation at room

temperature (~23°C) for 24h. The bacteria were imaged by scanning the plates, after which

quantification of β-galactosidase activity was done using ONPG as described in [83].

Protein purification

$His_{10}Smt3$-tagged PNPase and $His_{10}Smt3$-tagged RhlB proteins were purified from soluble bacterial

lysates by nickel-agarose affinity chromatography as described previously in the supplementary

information of Hausmann *et al.* 2021 [55]. The imidazole elution profiles were monitored by SDS-

PAGE. The recombinant $His_{10}Smt3$-tagged proteins were recovered predominantly in the 250 mM

imidazole fraction, shown in Figure 4 (input lane I). Protein concentration was determined using the

Bio-Rad dye reagent, with BSA as the standard, and calculated by interpolation against the BSA

standard curve. The His_{10}-Smt3 tag remained uncleaved in the protein-protein interaction assays. Note

that the $His_{10}Smt3$-tagged proteins do not bind to RNase E fusion proteins (data not shown).

RNase E_{1-529}, RNase $E_{572-1057}$, RNase $E_{572-1057\_NDPRmut}$, and RNase $E_{572-1057\_AR1mut}$ proteins, each carrying

a $His_{10}Smt3$ tag at the N-terminus and a Twin-Strep-tag®-Flag at the C-terminus, were purified as

described above. The 250 mM imidazole elution fractions were used directly in the Protein-Protein

interaction assays. For the EMSA experiment, the RNase E proteins were further purified as follows:

the 250 mM imidazole fraction were applied to a 1 ml of Strep-Tactin Sepharose (Iba) column that had

been equilibrated with buffer C (50 mM Tris pH8, 200 mM NaCl, 10 % glycerol). The column was

washed with 5 ml of buffer C and then eluted stepwise with 1 ml aliquots of buffer C containing, 2.5

mM D-Desthiobiotin respectively. The elution profiles were monitored by SDS-PAGE (not shown).

Protein-protein interaction assays

Purified RNase E_{1-529}-Flag, RNase $E_{572-1057}$-Flag, RNase $E_{572-1057\_NDPRmut}$- Flag, or RNase $E_{572-1057\_AR1mut}$-

Flag fusion proteins (20-25 µg each) were incubated for 1 hour at 4 °C on a rotating device with 25 µl

of Anti-Flag M2 magnetic beads (Sigma M8823) in 100 µl of binding buffer H (50 mM Tris-HCl, pH

8.0, 150 mM NaCl, 0.01% Triton X-100). The beads were then collected using a magnetic separator

and washed three times with 0.25 ml of binding buffer H to remove any unbound protein.

For affinity chromatography, the anti-Flag magnetic beads (Sigma-Aldrich) with bound RNase E-Flag

fusion proteins, or beads alone, were incubated with either 5 µg of PNPase (Figure 4A) or 5 µg of RhlB

(Figure 4B) in 100 µl of binding buffer H for 1 hour at 4 °C on a rotating device. After incubation, the

beads were collected using the magnet, and the supernatant (free fraction F) was withdrawn. The anti-

Flag beads were then resuspended in 100 µl of binding buffer H and subjected to three rounds of

washing. Following the final wash, the protein bound to the beads was eluted with 100 µl of 2X LDS

sample buffer (Invitrogen). Aliquots (15 µl) of the input protein sample (PNPase or RhlB), the first

supernatant fraction (free fraction F), and the bead-bound fraction were mixed with 5 µl of 4X LDS

sample buffer (Invitrogen), heated at 80 °C for 5 minutes, and analysed by SDS-PAGE. Polypeptides

were visualized using Coomassie Blue staining.

Growth assays on solid or liquid media

To assess the growth of the mutant strains in liquid medium, 200 µL of NYB broth was inoculated to

OD_{600nm}=0.05 in a 96-well plate, using an overnight culture washed twice with fresh medium. The wells

on the edge of the plate were not used. The plate was incubated at 37°C using a Biotek Synergy H1

Hybrid Plate reader. OD_{600nm} values were measured every 30 minutes. Before each OD_{600nm}

measurement, orbital shaking with 282 cpm (3 mm) was performed for 1 minute. Data was retrieved

the following day and plotted after substraction of blank OD_{600nm} values using GraphPad Prism.

When growing strains carrying the pME6032 vector for complementation assays, NYB medium was

supplemented with 50 µg/mL of Tetracycline and 70 µM of IPTG for induction. Cells were resuspended

in supplemented NYB after washing and before inoculating the 96-well plate.

To assess the growth of the mutant strains on solid medium, square Petri dishes (12x12cm) were filled

with 50 mL of warm NA. When testing growth at low vs high pH, either 75 µL of 37% fuming HCl or

400 µL of 5N NaOH was added per 50 mL of NA agar to achieve the desired conditions. A fresh culture

(OD_{600nm}= 0.4-0.8) of each strain was adjusted to OD_{600nm}=1 by centrifugation and serially diluted in a

96-well plate using a multipipette. 5 µL of each dilution series was then spotted using a multipipette

onto pre-dried NA plates (1 hour under a sterile flowhood). After the spots had dried, plates were

incubated upside down at 37°C or 16°C.

Galleria killing assays

Galleria mellonella larvae were used to assess in vivo virulence and/or survival of several RNase E

mutant strains. Fresh cultures were adjusted to a suspension of OD_{600nm}= 1 and serially diluted to 10^{-6}

in Gibco Dulbecco's Phosphate Buffered Saline (dPBS) (Thermo Fisher Scientific). 10 µL of the 10^{-6}

dilution were spotted onto NYB agar to verify that the dilutions were similar for the different strains.

10 μL spots were then put onto a clean parafilm and injected using Omnican 50 sterile insulin syringe

(Braun) into the second to last leg of the larvae, facing towards the head. As a control, sterile dPBS was

injected similarly into larvae to assess background levels of death. For every 10 larvae injected, the time

of injection was recorded, and they were placed inside a dark incubator at 37°C. The next day, deaths

were recorded every hour. A larva was considered dead when it completely stopped responding to

stimuli (touched on the legs or the head with a pipette tip). Survival curves were plotted and analysed

with GraphPad Prism (GraphPad Software, Prism10.2).

RNA-seq sample collection

PAO1 WT, PAO1 RNase E$_{1-588}$, (*rne588*) and PAO1 RNase E$_{1-529}$ (*rne529*) strains were grown in 50 mL

Erlenmeyer flasks (10 mL NYB broth) until OD$_{600nm}$= 1.0-1.2. 500 μL of culture were immediately

added into four separate 1.5 mL tubes, each containing 1 mL of RNAlater (Sigma-Aldrich, cat). After

harvesting of all three strains, the tubes were centrifuged for 30 minutes at 9600 x *g*. Supernatant was

discarded and pellets were stored at -80°C until RNA extraction. The RNA extraction was performed

using two tubes of pellets for each strain, with the Monarch Total RNA Miniprep kit. For lysis of the

bacterial cells, the two pellets were resuspended in 250 μL of 1X Tris-EDTA (TE) buffer supplemented

with 1 mg/mL of lysozyme and regrouped into one tube for each strain, incubated for 5 minutes at RT,

and RNA was subsequently extracted following the manufacturer's instructions. After RNA

purification, an additional DNA removal treatment was done using TURBO DNase (Thermo Fisher)

according to the manufacturer's instructions. RNA was purified again using the RNA purification

columns (dark blue) from the Monarch Total RNA Miniprep kit, and following the priming, washing

and elution steps described by the manufacturer. The absence of residual gDNA contamination was

verified by performing a PCR for 35 cycles using primer pairs targeting *rpoD* [55]. Biological triplicate

samples for each strain were prepared. Quality check, ribosomal RNA depletion and RNA sequencing

were conducted by Novogene.

RNA sequencing analysis

Reads were mapped using Bowtie2 [87] to *Pseudomonas aeruginosa* PAO1 genome assembly

ASM676v1 and counts were aggregated using summarize Overlaps in the package GenomicAlignments

[88] in R using the ASM676v1 gtf version 2.2 (NCBI accession: GCF_000006765.1). Differential

expression was performed using DESeq2 [89] and transcripts were considered differentially expressed

between two conditions if the absolute value of log2FC between conditions was greater than 1 (i.e. 2-

fold up or down) with a p-value less than 0.05. To find functional enrichment among differentially

expressed genes groups, the Kyoto Encyclopedia of Genes and Genomes (KEGG) was used for gene

set definitions, downloaded using the package KEGGREST [90] in R. Enrichment was calculated using

a hypergeometric test. All correlations presented are Pearson's product moment correlation coefficient.

Pairwise correlations of log2 Reads Per Kilobase per Million mapped reads (RPKMs) and hierarchical
unsupervised clustering analysis of fold changes relative to WT demonstrate the similarity in the profile
of the biological replicates for each strain. Analysis of 5' read ends per loci was performed as described
by de Araújo et al. [91] except that each dataset read number was down-sampled to 8 million.
707

708 **Acknowledgments**
709   We thank Sylvain Guex-Crosier for his assistance with RNA extraction and Nathanaël Guggenheim for
710   protein purification and assistance with cloning experiments. We also acknowledge the Proteomics Core
711   Facility and the Bioinformatics Support Platform from the University of Geneva for assistance with the
712   realisation of LC-MS-MS and bioinformatics analyses, respectively.

References

References

1.   Hui MP, Foley PL, Belasco JG. Messenger RNA Degradation in Bacterial Cells. Annual Review

of Genetics. 2014;48(1):537-59. doi: 10.1146/annurev-genet-120213-092340. PubMed PMID:

25292357.

2.   Celesnik H, Deana A, Belasco JG. Initiation of RNA decay in *Escherichia coli* by 5'

pyrophosphate removal. Mol Cell. 2007;27(1):79-90. doi: 10.1016/j.molcel.2007.05.038. PubMed

PMID: 17612492; PubMed Central PMCID: PMCPMC2196405.

3.   Deana A, Celesnik H, Belasco JG. The bacterial enzyme RppH triggers messenger RNA

degradation by 5' pyrophosphate removal. Nature. 2008;451(7176):355-8. doi: 10.1038/nature06475.

PubMed PMID: 18202662.

4.   Luciano DJ, Vasilyev N, Richards J, Serganov A, Belasco JG. Importance of a diphosphorylated

intermediate for RppH-dependent RNA degradation. RNA Biol. 2018;15(6):703-6. Epub 20180529.

doi: 10.1080/15476286.2018.1460995. PubMed PMID: 29619898; PubMed Central PMCID:

PMCPMC6152431.

5.   Tejada-Arranz A, de Crecy-Lagard V, de Reuse H. Bacterial RNA Degradosomes: Molecular

Machines under Tight Control. Trends Biochem Sci. 2020;45(1):42-57. Epub 20191101. doi:

10.1016/j.tibs.2019.10.002. PubMed PMID: 31679841; PubMed Central PMCID: PMCPMC6958999.

6.   Dressaire C, Picard F, Redon E, Loubière P, Queinnec I, Girbal L, Cocaign-Bousquet M. Role

of mRNA stability during bacterial adaptation. PLoS One. 2013;8(3):e59059. PubMed PMID:

23516597.

7.   Broglia L, Le Rhun A, Charpentier E. Methodologies for bacterial ribonuclease characterization

using RNA-seq. FEMS Microbiol Rev. 2023;47(5). PubMed PMID: 37656885.

8.   Ono M, Kuwano M. A conditional lethal mutation in an *Escherichia coli* strain with a longer

chemical lifetime of messenger RNA. J Mol Biol. 1979;129(3):343-57. Epub 1979/04/15. doi:

10.1016/0022-2836(79)90500-x. PubMed PMID: 110942.

9.   Apirion D. Isolation, genetic mapping and some characterization of a mutation in *Escherichia*

coli that affects the processing of ribonuleic acid. Genetics. 1978;90(4):659-71. PubMed PMID:

369943.

10.   Jain C, Deana A, Belasco JG. Consequences of RNase E scarcity in *Escherichia coli*. Molecular

Microbiology. 2002;43(4):1053-64. doi: https://doi.org/10.1046/j.1365-2958.2002.02808.x.

11.   Caruthers JM, Feng Y, McKay DB, Cohen SN. Retention of core catalytic functions by a

conserved minimal ribonuclease E peptide that lacks the domain required for tetramer formation. J Biol

Chem. 2006;281(37):27046-51. Epub 20060719. doi: 10.1074/jbc.M602467200. PubMed PMID:

16854990.

12.   Callaghan AJ, Aurikko JP, Ilag LL, Günter Grossmann J, Chandran V, Kühnel K, et al. Studies
of the RNA degradosome-organizing domain of the *Escherichia coli* ribonuclease RNase E. J Mol Biol.
2004;340(5):965-79. doi: 10.1016/j.jmb.2004.05.046. PubMed PMID: 15236960.

13.   Carpousis AJ, Van Houwe G, Ehretsmann C, Krisch HM. Copurification of E. coli RNAase E
and PNPase: evidence for a specific association between two enzymes important in RNA processing
and degradation. Cell. 1994;76(5):889-900. doi: 10.1016/0092-8674(94)90363-8. PubMed PMID:
7510217.

14.   Py B, Higgins CF, Krisch HM, Carpousis AJ. A DEAD-box RNA helicase in the *Escherichia
coli* RNA degradosome. Nature. 1996;381(6578):169-72. doi: 10.1038/381169a0. PubMed PMID:
8610017.

15.   Chandran V, Luisi BF. Recognition of enolase in the Escherichia coli RNA degradosome. J Mol
Biol. 2006;358(1):8-15. Epub 20060221. doi: 10.1016/j.jmb.2006.02.012. PubMed PMID: 16516921.

16.   Worrall JAR, Górna M, Crump NT, Phillips LG, Tuck AC, Price AJ, et al. Reconstitution and
Analysis of the Multienzyme Escherichia coli RNA Degradosome. Journal of Molecular Biology.
2008;382(4):870-83. doi: https://doi.org/10.1016/j.jmb.2008.07.059.

17.   Bandyra KJ, Bouvier M, Carpousis AJ, Luisi BF. The social fabric of the RNA degradosome.
Biochim Biophys Acta. 2013;1829(6-7):514-22. Epub 20130228. doi: 10.1016/j.bbagrm.2013.02.011.
PubMed PMID: 23459248; PubMed Central PMCID: PMCPMC3991390.

18.   Tsai YC, Du D, Domínguez-Malfavón L, Dimastrogiovanni D, Cross J, Callaghan AJ, et al.
Recognition of the 70S ribosome and polysome by the RNA degradosome in *Escherichia coli*. Nucleic
Acids Res. 2012;40(20):10417-31. Epub 20120825. doi: 10.1093/nar/gks739. PubMed PMID:
22923520; PubMed Central PMCID: PMCPMC3488216.

19.   Leroy A, Vanzo NF, Sousa S, Dreyfus M, Carpousis AJ. Function in *Escherichia coli* of the
non-catalytic part of RNase E: role in the degradation of ribosome-free mRNA. Mol Microbiol.
2002;45(5):1231-43. doi: 10.1046/j.1365-2958.2002.03104.x. PubMed PMID: 12207692.

20.   Dendooven T, Paris G, Shkumatov AV, Islam MS, Burt A, Kubańska MA, et al. Multi-scale
ensemble properties of the *Escherichia coli* RNA degradosome. Mol Microbiol. 2022;117(1):102-20.
Epub 20210925. doi: 10.1111/mmi.14800. PubMed PMID: 34415624; PubMed Central PMCID:
PMCPMC7613265.

21.   Sinha D, De Lay NR. Target recognition by RNase E RNA-binding domain AR2 drives sRNA
decay in the absence of PNPase. Proc Natl Acad Sci U S A. 2022;119(48):e2208022119. Epub
20221121. doi: 10.1073/pnas.2208022119. PubMed PMID: 36409892; PubMed Central PMCID:
PMCPMC9860253.

22.   Lee K, Zhan X, Gao J, Qiu J, Feng Y, Meganathan R, et al. RraA. a protein inhibitor of RNase
E activity that globally modulates RNA abundance in *E. coli*. Cell. 2003;114(5):623-34. PubMed PMID:
13678585.

23.   Górna MW, Pietras Z, Tsai YC, Callaghan AJ, Hernández H, Robinson CV, Luisi BF. The

regulatory protein RraA modulates RNA-binding and helicase activities of the E. coli RNA

degradosome. Rna. 2010;16(3):553-62. Epub 20100127. doi: 10.1261/rna.1858010. PubMed PMID:

20106955; PubMed Central PMCID: PMCPMC2822920.

24.   Gao J, Lee K, Zhao M, Qiu J, Zhan X, Saxena A, et al. Differential modulation of E. coli mRNA

abundance by inhibitory proteins that alter the composition of the degradosome. Mol Microbiol.

2006;61(2):394-406. Epub 20060612. doi: 10.1111/j.1365-2958.2006.05246.x. PubMed PMID:

16771842.

25.   Prud'homme-Généreux A, Beran RK, Iost I, Ramey CS, Mackie GA, Simons RW. Physical and

functional interactions among RNase E, polynucleotide phosphorylase and the cold-shock protein,

CsdA: evidence for a 'cold shock degradosome'. Molecular Microbiology. 2004;54(5):1409-21. doi:

https://doi.org/10.1111/j.1365-2958.2004.04360.x.

26.   Blum E, Py B, Carpousis AJ, Higgins CF. Polyphosphate kinase is a component of the

Escherichia coli RNA degradosome. Molecular Microbiology. 1997;26(2):387-98. doi:

https://doi.org/10.1046/j.1365-2958.1997.5901947.x.

27.   Kannaiah S, Goldberger O, Alam N, Barnabas G, Pozniak Y, Nussbaum-Shochat A, et al.

MinD-RNase E interplay controls localization of polar mRNAs in E. coli. Embo j. 2024;43(4):637-62.

Epub 20240119. doi: 10.1038/s44318-023-00026-9. PubMed PMID: 38243117; PubMed Central

PMCID: PMCPMC10897333.

28.   Taghbalout A, Rothfield L. RNaseE and the other constituents of the RNA degradosome are

components of the bacterial cytoskeleton. Proceedings of the National Academy of Sciences.

2007;104(5):1667-72. doi: 10.1073/pnas.0610491104.

29.   Ikeda Y, Yagi M, Morita T, Aiba H. Hfq binding at RhlB-recognition region of RNase E is

crucial for the rapid degradation of target mRNAs mediated by sRNAs in Escherichia coli. Mol

Microbiol. 2011;79(2):419-32. Epub 20101118. doi: 10.1111/j.1365-2958.2010.07454.x. PubMed

PMID: 21219461.

30.   Aït-Bara S, Carpousis AJ, Quentin Y. RNase E in the γ-Proteobacteria: conservation of

intrinsically disordered noncatalytic region and molecular evolution of microdomains. Mol Genet

Genomics. 2015;290(3):847-62. Epub 20141129. doi: 10.1007/s00438-014-0959-5. PubMed PMID:

25432321; PubMed Central PMCID: PMCPMC4435900.

31.   Goldblum K, Apririon D. Inactivation of the ribonucleic acid-processing enzyme ribonuclease

E blocks cell division. J Bacteriol. 1981;146(1):128-32. doi: 10.1128/jb.146.1.128-132.1981. PubMed

PMID: 6163761; PubMed Central PMCID: PMCPMC217061.

32.   Bernstein JA, Lin PH, Cohen SN, Lin-Chao S. Global analysis of *Escherichia coli* RNA

degradosome function using DNA microarrays. Proc Natl Acad Sci U S A. 2004;101(9):2758-63. Epub

20040223. doi: 10.1073/pnas.0308747101. PubMed PMID: 14981237; PubMed Central PMCID:

PMCPMC365694.

33.   Kido M, Yamanaka K, Mitani T, Niki H, Ogura T, Hiraga S. RNase E polypeptides lacking a
carboxyl-terminal half suppress a mukB mutation in *Escherichia coli*. J Bacteriol. 1996;178(13):3917-
25. doi: 10.1128/jb.178.13.3917-3925.1996. PubMed PMID: 8682798; PubMed Central PMCID:
PMCPMC232654.

34.   Lopez PJ, Marchand I, Joyce SA, Dreyfus M. The C-terminal half of RNase E, which organizes
the *Escherichia coli* degradosome, participates in mRNA degradation but not rRNA processing in vivo.
Mol Microbiol. 1999;33(1):188-99. doi: 10.1046/j.1365-2958.1999.01465.x. PubMed PMID:
10411735.

35.   Hardwick SW, Chan VS, Broadhurst RW, Luisi BF. An RNA degradosome assembly in
Caulobacter crescentus. Nucleic Acids Res. 2011;39(4):1449-59. Epub 20101015. doi:
10.1093/nar/gkq928. PubMed PMID: 20952404; PubMed Central PMCID: PMCPMC3045602.

36.   Voss JE, Luisi BF, Hardwick SW. Molecular recognition of RhlB and RNase D in the
Caulobacter crescentus RNA degradosome. Nucleic Acids Res. 2014;42(21):13294-305. Epub
20141111. doi: 10.1093/nar/gku1134. PubMed PMID: 25389270; PubMed Central PMCID:
PMCPMC4245959.

37.   Zhang JY, Deng XM, Li FP, Wang L, Huang QY, Zhang CC, Chen WL. RNase E forms a
complex with polynucleotide phosphorylase in cyanobacteria via a cyanobacterial-specific nonapeptide
in the noncatalytic region. RNA. 2014;20(4):568-79. Epub 20140221. doi: 10.1261/rna.043513.113.
PubMed PMID: 24563514; PubMed Central PMCID: PMCPMC3964918.

38.   Yan HD, Qin XX, Wang L, Chen WL. Both Enolase and the DEAD-Box RNA Helicase CrhB
Can Form Complexes with RNase E in *Anabaena* sp. Strain PCC 7120. Applied and environmental
microbiology. 2020;86(13). doi: 10.1128/AEM.00425-20. PubMed PMID: WOS:000546650600011.

39.   Zhou C, Zhang JY, Hu XY, Li CC, Wang L, Huang QY, Chen WL. RNase II binds to RNase E
and modulates its endoribonucleolytic activity in the cyanobacterium *Anabaena* PCC 7120. Nuceic
Acid Res. 2020;48(7):3922-34. doi: 10.1093/nar/gkaa092. PubMed PMID: WOS:000525957700040.

40.   Purusharth RI, Klein F, Sulthana S, Jäger S, Jagannadham MV, Evguenieva-Hackenberg E, et
al. Exoribonuclease R Interacts with Endoribonuclease E and an RNA Helicase in the Psychrotrophic
Bacterium Pseudomonas syringae Lz4W*. Journal of Biological Chemistry. 2005;280(15):14572-8.
doi: https://doi.org/10.1074/jbc.M413507200.

41.   Jäger S, Fuhrmann O, Heck C, Hebermehl M, Schiltz E, Rauhut R, Klug G. An mRNA
degrading complex in *Rhodobacter capsulatus*. Nucleic Acids Res. 2001;29(22):4581-8. doi:
10.1093/nar/29.22.4581. PubMed PMID: 11713307; PubMed Central PMCID: PMCPMC92556.

42.   Aït-Bara S, Carpousis Agamemnon J. Characterization of the RNA Degradosome of
Pseudoalteromonas haloplanktis: Conservation of the RNase E-RhlB Interaction in the
Gammaproteobacteria. Journal of Bacteriology. 2010;192(20):5413-23. doi: 10.1128/jb.00592-10.

43. Hardwick SW, Gubbey T, Hug I, Jenal U, Luisi BF. Crystal structure of *Caulobacter crescentus*
polynucleotide phosphorylase reveals a mechanism of RNA substrate channelling and RNA
degradosome assembly. Open Biology. 2012;2(4):120028. doi: 10.1098/rsob.120028.

44. Nurmohamed S, Vaidialingam B, Callaghan AJ, Luisi BF. Crystal Structure of Escherichia coli
Polynucleotide Phosphorylase Core Bound to RNase E, RNA and Manganese: Implications for
Catalytic Mechanism and RNA Degradosome Assembly. Journal of Molecular Biology.
2009;389(1):17-33. doi: https://doi.org/10.1016/j.jmb.2009.03.051.

45. Green SK, Schroth MN, Cho JJ, Kominos SK, Vitanza-jack VB. Agricultural plants and soil as
a reservoir for *Pseudomonas aeruginosa*. Applied microbiology. 1974;28(6):987-91. Epub 1974/12/01.
PubMed PMID: 4217591; PubMed Central PMCID: PMCPMC186868.

46. Hardalo C, Edberg SC. *Pseudomonas aeruginosa*: assessment of risk from drinking water.
Critical reviews in microbiology. 1997;23(1):47-75. Epub 1997/01/01. doi:
10.3109/10408419709115130. PubMed PMID: 9097014.

47. Pendleton JN, Gorman SP, Gilmore BF. Clinical relevance of the ESKAPE pathogens. Expert
review of anti-infective therapy. 2013;11(3):297-308. Epub 2013/03/06. doi: 10.1586/eri.13.12.
PubMed PMID: 23458769.

48. https://www.who.int/news/item/27-02-2017-who-publishes-list-of-bacteria-for-which-new-
antibiotics-are-urgently-needed.

49. Garcia-Clemente M, de la Rosa D, Máiz L, Girón R, Blanco M, Olveira C, et al. Impact of
Pseudomonas aeruginosa Infection on Patients with Chronic Inflammatory Airway Diseases. Journal of
clinical medicine. 2020;9(12). Epub 2020/12/02. doi: 10.3390/jcm9123800. PubMed PMID: 33255354.

50. Urwin L, Okurowska K, Crowther G, Roy S, Garg P, Karunakaran E, et al. Corneal Infection
Models: Tools to Investigate the Role of Biofilms in Bacterial Keratitis. Cells. 2020;9(11). Epub
2020/11/14. doi: 10.3390/cells9112450. PubMed PMID: 33182687; PubMed Central PMCID:
PMCPMC7696224.

51. Costerton JW, Stewart PS, Greenberg EP. Bacterial biofilms: a common cause of persistent
infections. Science (New York, NY). 1999;284(5418):1318-22. Epub 1999/05/21. doi:
10.1126/science.284.5418.1318. PubMed PMID: 10334980.

52. Lu Q, Eggimann P, Luyt CE, Wolff M, Tamm M, François B, et al. Pseudomonas aeruginosa
serotypes in nosocomial pneumonia: prevalence and clinical outcomes. Critical care (London, England).
2014;18(1):R17. Epub 2014/01/17. doi: 10.1186/cc13697. PubMed PMID: 24428878; PubMed Central
PMCID: PMCPMC4057348.

53. Hancock RE. Resistance mechanisms in *Pseudomonas aeruginosa* and other nonfermentative
gram-negative bacteria. Clin Infect Dis. 1998;27 Suppl 1:S93-9. Epub 1998/08/26. PubMed PMID:
9710677.

54. Van den Bossche A, Hardwick SW, Ceyssens PJ, Hendrix H, Voet M, Dendooven T, et al.
Structural elucidation of a novel mechanism for the bacteriophage-based inhibition of the RNA

893 degradosome. Elife. 2016;5. Epub 20160722. doi: 10.7554/eLife.16413. PubMed PMID: 27447594;
894 PubMed Central PMCID: PMCPMC4980113.

895 55. Hausmann S, Gonzalez D, Geiser J, Valentini M. The DEAD-box RNA helicase RhlE2 is a
896 global regulator of *Pseudomonas aeruginosa* lifestyle and pathogenesis. Nucleic Acids Res.
897 2021;49(12):6925-40. doi: 10.1093/nar/gkab503. PubMed PMID: 34151378; PubMed Central PMCID:
898 PMCPMC8266600.

899 56. Hausmann S, Geiser J, Allen GE, Geslain SAM, Valentini M. Intrinsically disordered regions
900 regulate RhlE RNA helicase functions in bacteria. Nucleic Acids Res. 2024. Epub 20240614. doi:
901 10.1093/nar/gkae511. PubMed PMID: 38874491.

902 57. Sonnleitner E, Romeo A, Bläsi U. Small regulatory RNAs in *Pseudomonas aeruginosa*. RNA
903 Biol. 2012;9(4):364-71. Epub 2012/02/18. doi: 10.4161/rna.19231. PubMed PMID: 22336763.

904 58. Valentini M, Gonzalez D, Mavridou DA, Filloux A. Lifestyle transitions and adaptive
905 pathogenesis of *Pseudomonas aeruginosa*. Current opinion in microbiology. 2018;41:15-20. Epub
906 2017/11/23. doi: 10.1016/j.mib.2017.11.006. PubMed PMID: 29166621.

907 59. Ait-Bara S, Carpousis AJ, Quentin Y. RNase E in the gamma-Proteobacteria: conservation of
908 intrinsically disordered noncatalytic region and molecular evolution of microdomains. Mol Genet
909 Genomics. 2015;290(3):847-62. Epub 20141129. doi: 10.1007/s00438-014-0959-5. PubMed PMID:
910 25432321; PubMed Central PMCID: PMCPMC4435900.

911 60. Nishio S-y, Itoh T. Arginine-rich RNA binding domain and protein scaffold domain of RNase
912 E are important for degradation of RNAI but not for that of the Rep mRNA of the ColE2 plasmid.
913 Plasmid. 2009;62(2):83-7. doi: https://doi.org/10.1016/j.plasmid.2009.04.002.

914 61. Hamouche L, Poljak L, Carpousis AJ. Polyribosome-Dependent Clustering of Membrane-
915 Anchored RNA Degradosomes To Form Sites of mRNA Degradation in *Escherichia coli*. mBio.
916 2021;12(5):e0193221. Epub 20210907. doi: 10.1128/mBio.01932-21. PubMed PMID: 34488454;
917 PubMed Central PMCID: PMCPMC8546579.

918 62. Al-Husini N, Tomares DT, Bitar O, Childers WS, Schrader JM. α-Proteobacterial RNA
919 Degradosomes Assemble Liquid-Liquid Phase-Separated RNP Bodies. Mol Cell. 2018;71(6):1027-
920 39.e14. Epub 20180906. doi: 10.1016/j.molcel.2018.08.003. PubMed PMID: 30197298; PubMed
921 Central PMCID: PMCPMC6151146.

922 63. Khemici V, Poljak L, Luisi BF, Carpousis AJ. The RNase E of *Escherichia coli* is a membrane-
923 binding protein. Mol Microbiol. 2008;70(4):799-813. Epub 20081002. doi: 10.1111/j.1365-
924 2958.2008.06454.x. PubMed PMID: 18976283; PubMed Central PMCID: PMCPMC7610891.

925 64. Schuck A, Diwa A, Belasco JG. RNase E autoregulates its synthesis in Escherichia coli by
926 binding directly to a stem-loop in the rne 5' untranslated region. Mol Microbiol. 2009;72(2):470-8. Epub
927 20090306. doi: 10.1111/j.1365-2958.2009.06662.x. PubMed PMID: 19320830; PubMed Central
928 PMCID: PMCPMC2857391.

65. Liu S-J, Lin G-M, Yuan Y-Q, Chen W, Zhang J-Y, Zhang C-C. A conserved protein inhibitor brings under check the activity of RNase E in cyanobacteria. Nucleic Acids Research. 2024;52(1):404-19. doi: 10.1093/nar/gkad1094.

66. Karimova G, Ullmann A, Ladant D. [5] A bacterial two-hybrid system that exploits a cAMP signaling cascade in Escherichia coli. In: Thorner J, Emr SD, Abelson JN, editors. Methods in Enzymology. 328: Academic Press; 2000. p. 59-73.

67. Jander G, Rahme Laurence G, Ausubel Frederick M. Positive Correlation between Virulence ofPseudomonas aeruginosa Mutants in Mice and Insects. Journal of Bacteriology. 2000;182(13):3843-5. doi: 10.1128/jb.182.13.3843-3845.2000.

68. Lin J, Cheng J, Wang Y, Shen X. The Pseudomonas Quinolone Signal (PQS): Not Just for Quorum Sensing Anymore. Front Cell Infect Microbiol. 2018;8:230. Epub 20180704. doi: 10.3389/fcimb.2018.00230. PubMed PMID: 30023354; PubMed Central PMCID: PMCPMC6039570.

69. Gonçalves T, Vasconcelos U. Colour Me Blue: The History and the Biotechnological Potential of Pyocyanin. Molecules. 2021;26(4). Epub 20210210. doi: 10.3390/molecules26040927. PubMed PMID: 33578646; PubMed Central PMCID: PMCPMC7916356.

70. Chou HT, Li J-Y, Lu C-D. Functional Characterization of the agtABCD and agtSR Operons for 4-Aminobutyrate and 5-Aminovalerate Uptake and Regulation in Pseudomonas aeruginosa PAO1. Current Microbiology. 2014;68(1):59-63. doi: 10.1007/s00284-013-0446-y.

71. Hauser AR. The type III secretion system of Pseudomonas aeruginosa: infection by injection. Nat Rev Microbiol. 2009;7(9):654-65. doi: 10.1038/nrmicro2199. PubMed PMID: 19680249; PubMed Central PMCID: PMCPMC2766515.

72. Nandana V, Al-Husini N, Vaishnav A, Dilrangi KH, Schrader JM. Caulobacter crescentus RNase E condensation contributes to autoregulation and fitness. Mol Biol Cell. 2024;35(8):ar104. Epub 20240612. doi: 10.1091/mbc.E23-12-0493. PubMed PMID: 38865176; PubMed Central PMCID: PMCPMC11321048.

73. Wurtzel O, Yoder-Himes DR, Han K, Dandekar AA, Edelheit S, Greenberg EP, et al. The single-nucleotide resolution transcriptome of Pseudomonas aeruginosa grown in body temperature. PLoS Pathog. 2012;8(9):e1002945. Epub 20120927. doi: 10.1371/journal.ppat.1002945. PubMed PMID: 23028334; PubMed Central PMCID: PMCPMC3460626.

74. de Araújo Hugo L, Picinato Beatriz A, Lorenzetti Alan PR, Muthunayake Nisansala S, Rathnayaka-Mudiyanselage IW, dos Santos Naara M, et al. The DEAD-box RNA helicase RhlB is required for efficient RNA processing at low temperature in Caulobacter. Microbiology Spectrum. 2023;11(6):e01934-23. doi: 10.1128/spectrum.01934-23.

75. Vaillancourt M, Galdino ACM, Limsuwannarot SP, Celedonio D, Dimitrova E, Broerman M, et al. A compensatory RNase E variation increases Iron Piracy and Virulence in multidrug-resistant *Pseudomonas aeruginosa* during Macrophage infection. PLoS Pathog. 2023;19(4):e1010942. Epub

20230407. doi: 10.1371/journal.ppat.1010942. PubMed PMID: 37027441; PubMed Central PMCID:

PMCPMC10115287.

76. Khemici V, Toesca I, Poljak L, Vanzo NF, Carpousis AJ. The RNase E of *Escherichia coli* has

at least two binding sites for DEAD-box RNA helicases: functional replacement of RhlB by RhlE.

Molecular Microbiology. 2004;54(5):1422-30. doi: https://doi.org/10.1111/j.1365-2958.2004.04361.x.

77. Worrall JA, Howe FS, McKay AR, Robinson CV, Luisi BF. Allosteric activation of the ATPase

activity of the *Escherichia coli* RhlB RNA helicase. J Biol Chem. 2008;283(9):5567-76. PubMed

PMID: 18165229.

78. Lüthi E, Mercenier A, Haas D. The arcABC Operon Required for Fermentative Growth of

Pseudomonas aeruginosa on Arginine: Tn5-751-assisted Cloning and Localization of Structural Genes.

Microbiology. 1986;132(10):2667-75. doi: https://doi.org/10.1099/00221287-132-10-2667.

79. Gamper M, Ganter B, Polito MR, Haas D. RNA processing modulates the expression of the

arcDABC operon in *Pseudomonas aeruginosa*. Journal of Molecular Biology. 1992;226(4):943-57. doi:

https://doi.org/10.1016/0022-2836(92)91044-P.

80. Rietsch A, Vallet-Gely I, Dove SL, Mekalanos JJ. ExsE, a secreted regulator of type III

secretion genes in Pseudomonas aeruginosa. Proc Natl Acad Sci U S A. 2005;102(22):8006-11. Epub

20050523. doi: 10.1073/pnas.0503005102. PubMed PMID: 15911752; PubMed Central PMCID:

PMCPMC1142391.

81. Laville J, Voisard C, Keel C, Maurhofer M, Défago G, Haas D. Global control in Pseudomonas

fluorescens mediating antibiotic synthesis and suppression of black root rot of tobacco. Proc Natl Acad

Sci U S A. 1992;89(5):1562-6. doi: 10.1073/pnas.89.5.1562. PubMed PMID: 1311842; PubMed Central

PMCID: PMCPMC48492.

82. Ye RW, Haas D, Ka JO, Krishnapillai V, Zimmermann A, Baird C, Tiedje JM. Anaerobic

activation of the entire denitrification pathway in Pseudomonas aeruginosa requires Anr, an analog of

Fnr. J Bacteriol. 1995;177(12):3606-9. doi: 10.1128/jb.177.12.3606-3609.1995. PubMed PMID:

7768875; PubMed Central PMCID: PMCPMC177071.

83. Bordi C, Lamy MC, Ventre I, Termine E, Hachani A, Fillet S, et al. Regulatory RNAs and the

HptB/RetS signalling pathways fine-tune Pseudomonas aeruginosa pathogenesis. Mol Microbiol.

2010;76(6):1427-43. Epub 20100401. doi: 10.1111/j.1365-2958.2010.07146.x. PubMed PMID:

20398205; PubMed Central PMCID: PMCPMC2904497.

84. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al. Fiji: an open-

source platform for biological-image analysis. Nat Methods. 2012;9(7):676-82. Epub 20120628. doi:

10.1038/nmeth.2019. PubMed PMID: 22743772; PubMed Central PMCID: PMCPMC3855844.

85. Team RC. R: A Language and Environment for Statistical Computing. R Foundation for

Statistical Computing. Vienna, Austria2023.

86. Ream JA, Lewis LK, Lewis KA. Rapid agarose gel electrophoretic mobility shift assay for quantitating protein: RNA interactions. Anal Biochem. 2016;511:36-41. Epub 20160802. doi: 10.1016/j.ab.2016.07.027. PubMed PMID: 27495142; PubMed Central PMCID: PMCPMC5002362.

87. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nature Methods. 2012;9(4):357-9. doi: 10.1038/nmeth.1923.

88. Lawrence M, Huber W, Pagès H, Aboyoun P, Carlson M, Gentleman R, et al. Software for Computing and Annotating Genomic Ranges. PLOS Computational Biology. 2013;9(8):e1003118. doi: 10.1371/journal.pcbi.1003118.

89. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biology. 2014;15(12):550. doi: 10.1186/s13059-014-0550-8.

90. B. TDM. KEGGREST: Client-side REST access to the Kyoto Encyclopedia of Genes and Genomes (KEGG). doi:10.18129/B9.bioc.KEGGREST. R package version 1420, https://bioconductororg/packages/KEGGREST2023.

91. de Araújo HL, Picinato BA, Lorenzetti APR, Muthunayake NS, Rathnayaka-Mudiyanselage IW, Dos Santos NM, et al. The DEAD-box RNA helicase RhlB is required for efficient RNA processing at low temperature in Caulobacter. Microbiol Spectr. 2023:e0193423. Epub 2023/10/18. doi: 10.1128/spectrum.01934-23. PubMed PMID: 37850787.

92. Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J Mol Biol. 1990;215(3):403-10. doi: 10.1016/s0022-2836(05)80360-2. PubMed PMID: 2231712.

93. Pagès HA, P.; Gentleman, R.; DebRoy, S. Biostrings: Efficient manipulation of biological strings. 2024.

94. Grant BJ, Rodrigues AP, ElSawy KM, McCammon JA, Caves LS. Bio3d: an R package for the comparative analysis of protein structures. Bioinformatics. 2006;22(21):2695-6. Epub 20060829. doi: 10.1093/bioinformatics/btl461. PubMed PMID: 16940322.

95. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res. 2004;32(5):1792-7. Epub 20040319. doi: 10.1093/nar/gkh340. PubMed PMID: 15034147; PubMed Central PMCID: PMCPMC390337.

96. Erdős G, Pajkos M, Dosztányi Z. IUPred3: prediction of protein disorder enhanced with unambiguous experimental annotation and visualization of evolutionary conservation. Nucleic Acids Research. 2021;49(W1):W297-W303. doi: 10.1093/nar/gkab408.

97. Xue B, Dunbrack RL, Williams RW, Dunker AK, Uversky VN. PONDR-FIT: a meta-predictor of intrinsically disordered amino acids. Biochim Biophys Acta. 2010;1804(4):996-1010. Epub 20100125. doi: 10.1016/j.bbapap.2010.01.011. PubMed PMID: 20100603; PubMed Central PMCID: PMCPMC2882806.

98. McLaren RS, Newbury SF, Dance GS, Causton HC, Higgins CF. mRNA degradation by processive 3'-5' exoribonucleases in vitro and the implications for prokaryotic mRNA decay in vivo. J Mol Biol. 1991;221(1):81-95. PubMed PMID: 1920421.

1037   99.   Bolte S, Cordelières FP. A guided tour into subcellular colocalization analysis in light
1038   microscopy. J Microsc. 2006;224(Pt 3):213-32. doi: 10.1111/j.1365-2818.2006.01706.x. PubMed
1039   PMID: 17210054.

1040

     Figure Legends

     Figure 1: Comparison of *E. coli* (Ec) and *P. aeruginosa* (Pa) RNase E architecture. (A) Ec RNase

     E architecture, adapted from [17]. The NTD is shown is grey, and experimentally defined regions

     necessary for interaction with the membrane, with RNA, or with the three core protein partners are

     highlighted in red, dark blue, and yellow/purple/pink, respectively. **(B)** Pa RNase E architecture based

     on in-silico analyses performed in Fig S1-S2 and in [59]. The SLiM nomenclature was taken from [59],

     while the consensus sequence characterising each SLiM was specifically adapted based on sequence

     conservation within the Pseudomonas genus (Fig S1). MTS: membrane-targeting sequence, AR4:

     arginine-rich motif 4, REER-repeats: region with high frequency of occurrence of "RE[E/D]R" motif,

     M29: motif 29, AR1: arginine-rich motif 1, M20: motif 20, NDPR: motif with a highly conserved NDPR

     sequence. All diagrams were drawn to scale. IUPRED3 [96] and PONDR [97] predictions of disorder

     propensity (probability score) are shown for Ec RNase E **(A)** and Pa RNase E **(B)**.

     Figure 2: RNA binding of Pa RNase E CTD via the AR1, AR4, and REER SLiMs. (A) EMSA with

     purified His-Smt3 tagged RNase E CTD$^{572\text{-}1057}$ variants having either the native sequence (WT) or

     systematic alanine mutation of the arginine residues found within the REER-repeats region (REERmut),

     the AR4 SLiM (AR4mut) or the AR1 SLiM (AR1mut). The *malEF* mRNA intergenic region was used

     as a substrate [98]. Ratios of RNA:protein ranging from 1:10 to 1:80 were tested. See Fig S2 for

     additional EMSA with additional RNA:protein ratios. **(B)** Representative microscopy images of a PAO1

     strain expressing the RNase E-msfGFP fusion or mutated or truncated variants as indicated. **(C)**

     Quantification of signal intensity variance in the strains of panel B as indication of foci disappearance

     (see methods). **(D)** Representative microscopy images of a PAO1 strain expressing the RNase E-

     msfGFP fusion after exposure to rifampicin (+ Rif, exposure to a concentration of 100 μg/mL for 30

     minutes) or to DMSO as a negative control (- Rif). Corrected mean fluorescent signal intensity **(E)** and

     signal intensity variance **(F)** is showed in the graphs (see methods). Each quantification analysis

     involved at least 50 individual cells. The scale is 2 μm. Statistical significance was assessed by

     performing t-tests using R. p-values for comparisons tested are indicated as follows: *:<0.05; **:<0.01;

     :<0.001; *:<0.0001.

     Figure 3: **Identification of scaffold domain-dependent RNase E protein partners by pull-down**

     assays and mapping of the interactions by bacterial two-hybrid assays (A) Volcano plot

     highlighting the relative enrichment of proteins co-eluting with the RNase E$^{1\text{-}1057}$-Strep bait protein

     compared to the untagged negative control. The plot was generated from the average peptide count

     obtained in duplicate pull-down experiments. An arbitrary threshold of 5.5-fold change was chosen for

     selection of candidate (c) proteins. **(B)** Scatterplot highlighting comparative enrichment of proteins co-

     eluting with the RNase E$^{1\text{-}1057}$-Strep bait protein (x axis) or with the RNase E$^{1\text{-}529}$-Strep bait protein (y

axis) compared to the untagged negative control. **(C)** Quantification of β-galactosidase activity in *E.*
coli BTH101 cells co-expressing T18-RNase E with T25-PNPase, T25-RhlB, T25-ArcB, T25-GdhB or
T25-RpoB. Interaction between T18-RNase E and a T25-tagged protein partner results in a significant
restoration of β-galactosidase activity compared with empty vector (e.v.) negative controls. Unpaired
Welch's t-tests was performed using GraphPad Prism to assess differences between every tested protein-
protein interaction and the respective negative control. ***: p-value <0.0001; n.s: not significant (99%
confidence level). **(D-F)** Mapping of the interaction between RNase E and PNPase **(D)**, RhlB **(E)**, or
ArcB **(F)** by quantification of β-galactosidase activity in *E. coli* BTH101 cells co-expressing T25-
PNPase **(D)**, T25-RhlB **(E)**, or T25-ArcB **(F)** and either T18-RNase E or one of eight to ten truncated
T18-RNase E fragments. Empty vector (e.v.) negative controls are included into each separate graph
for comparison.

1089

Figure 4: PNPase and RhlB directly interact with RNase E scaffold domain via NDPR and AR1
SLiMs, respectively. Purified Flag-tagged RNase E$^{1-529}$ or RNase E$^{572-1057}$ proteins having either native
or mutated sequence were used as baits, immobilized to anti-flag magnetic beads and incubated with
purified His-Smt3 tagged PNPase **(A)** or RhlB **(B)** proteins to assess direct protein binding. I: input, F:
free fraction, B: bound fraction. NDPRmut, AR1mut: mutations of NDPR and AR1 SLiMs, respectively
(see Table S4). Binding assays were repeated at least 3-times and representative gels are shown.

1096

Figure 5: **Colocalisation of PNPase or RhlB with RNase E foci is dependent on their interaction**
with NDPR and AR1 RNase E SLiMs, respectively. (A) Visualisation of PNPase-msfGFP (upper
panels) or RhlB-msfGFP (lower panels) fusion and RNase E-mCherry fusion co-expressed in the same
strain. Pearson's and M coefficients were calculated in Fiji using the JaCoP plugin [99]. Three
independent images were used, each containing at least 30 cells, and an average coefficient was
calculated from the three values obtained with each image. **(B)** Visualisation of PNPase-msfGFP (upper
panels) and RhlB-msfGFP (lower panels) in strains expressing either the native RNase E protein (full
length) or truncated variants, as indicated above each image. Scale is 2 μm.

1105

Figure 6: Phenotypes affected by deletion or mutation of RNase E scaffolding domain. (A) Growth
on NA agar plates at 37°C or 16°C, or under low vs high pH conditions of wild-type (WT), *rne588*,
rne529, *rne*AR1^{mut}, *rne*NDPR^{mut}, and *rne*AR1+AR4+REER^{mut} strains, as indicated. Experiments were
performed in biological independent triplicates; a representative plate is shown. **(B)** Growth in NYB
broth at 37°C in 96-well plates of wild-type (WT), *rne588*, *rne529*, *rne*AR1^{mut}, *rne*NDPR^{mut}, and
*rne*AR1+AR4+REER^{mut} strains, as indicated. Independent biological triplicates, each with four
technical replicates, were used to generate the average growth curve shown. **(C)** Killing curves of *G.*
mellonella larvae injected with 10 μL of WT or mutant *P. aeruginosa* strains (as in panel A). The control
group was injected with 10 μL of sterile PBS instead. Larvae were incubated at 37°C and larval death

was monitored over 13-26 hours post-injection. A larva was considered dead when it completely

stopped responding to repeated touching stimuli. Two independent replicates were performed using a

total of at least 50 larvae for each strain tested. A Kaplan-Meier test for survival analysis was performed

using GraphPad Prism and comparing the survival curve of each mutant strain to that of the WT strain.

***: p-value <0.0001; n.s: not significant (99% confidence interval).

1120

Figure 7: RNA-seq analysis of RNase E CTD truncated mutants reveals broad transcriptome

changes. (A-B) Volcano plots highlighting genes differentially regulated in the **(A)** rne^{1-529} (R529) or

(B) rne $^{1-588}$ (R588) mutant strains compared with the WT strain. Genes significantly up- or down-

regulated (P-value < 0.05, fold change |FC| ≥ 2) are highlighted in red or blue, respectively. **(C)**

Scatterplot of genes differentially regulated in rne $^{1-529}$ strain versus genes differentially regulated in

rne^{1-588} strain, highlighting commonalities and differences between the two mutant strains. **(D-E)**

Clustered heat maps of type 3 secretion system **(D)** or ribosomal proteins **(E)** transcript levels in the

RNase E^{1-529} and by about 2-fold in the RNase E^{1-588} strains as compared to the WT strain (fold change).

(F) Differential analyses (log 2-fold change) of normalized total reads counts corresponding to the 5'

UTR region of the PA2976 (rne) transcript or RNase E NTD coding sequence in the RNase E^{1-529} and

by about 2-fold in the RNase E^{1-588} strains as compared to the WT strain. **(G)** Proportion of loci with 5'

read ends (sequenced fragments) counted only in a strain (nominator) relative to the 5' read ends counted

in that strain and in a WT replica (denominator). A total of 8 million reads was analysed in each strain

to exclude an effect due to the library depth (see methods). Histograms of individual values calculated

per each strain and replicate is shown in Figure S8.

1136

1137   **Figure 8: Model of RNA degradosome in _P. aeruginosa_.** Numbers corresponds to RNase E SLiM

1138   start-end residues involved in membrane, RNA, RhlB or PNPase interaction. RNA is represented as a

1139   dotted line. See discussion for details.

A

B

Figure 1

Figure 2

Figure 3

Figure 4

A

RNase E-mCherry   PNPase-msfGFP   Merged

Pearson's = 0.880
M1 = 0.910
M2 = 0.943

RNase E-mCherry   RhlB-msfGFP   Merged

Pearson's = 0.773
M1 = 0.937
M2 = 0,924

B

PAO1 WT   PAO1 *rne529*   PAO1 *rne*AR1^{mut} PAO1 *rne*NDPR^{mut}

PNPase-msfGFP

RhlB-msfGFP

Figure 5

Figure 6

Figure 7

Figure 8