1 Identification and characterization of a wet adhesive protein extracted
2 from *Dreissena bugensis*, the freshwater quagga mussel

Angelico R. Obille[1], Rida Hasan[1], David J. Rees[1], Judith Ng[1], Karina M. M. Carneiro[1,2],
Eli D. Sone*[1,2,3]

[1] Institute of Biomedical Engineering, University of Toronto, Toronto, ON, Canada, M5S 3G9
[2] Faculty of Dentistry, University of Toronto, Toronto, ON, Canada, M5G 1G6
[3] Department of Materials Science & Engineering, University of Toronto, Toronto, ON, Canada, M5S 3E4

* **Corresponding author**: Eli D. Sone (eli.sone@utoronto.ca; 416-978-7422; 124 Edward Street, Toronto,
ON, Canada, M5G 1G6)

Classification: Biological Sciences/Biochemistry
Keywords: quagga mussels, byssus, mussel adhesive protein, freshwater bioadhesion, atomic force
microscopy

13 *Abstract*

Mechanisms of wet adhesion have been developed by several aquatic organisms

over millions of years of evolutionary processes. Yet, the repertoire of synthetic

biocompatible wet adhesive materials is still limited. In most marine bioadhesive

proteins, 3,4-dihydroxyphenylalanine (DOPA) plays a significant role in strong interfacial

interactions. The bioadhesive proteins in freshwater organisms are less well

understood. The quagga mussel (*Dreissena bugensis*) is a notorious freshwater

invasive species in the Great Lakes that attaches to a plethora of surfaces via a byssus.

To determine the adhesive proteins in the quagga mussel byssus, we utilized

quantitative proteomics to identify the proteins enriched at the byssus-substrate

interface. Among the identified proteins was the Dbfp7 protein family. Dbfp7 is a small,

polymorphic, and mostly disordered protein that lacks significant amounts of DOPA.

Atomic force microscopy measurements of Dbfp7 confirm that this protein has similar

adhesion energy to marine mussel adhesive proteins in aqueous conditions despite

lacking DOPA. These results suggest that freshwater mussels may employ different

mechanisms of adhesion compared to marine byssates. The inclusion of Dbfp7 to the

library of known wet bioadhesive proteins – the first functionally characterized

freshwater bioadhesive protein to our knowledge – will allow for a better understanding

of the fundamental properties required to achieve biocompatible wet adhesion, a crucial

step for the development of bio-inspired wet adhesive materials, such as improved

medical adhesives.

Significance Statement

While many aquatic organisms evolved strategies to adhere to surfaces

underwater, current synthetic biocompatible adhesives lack reliable adhesive strength in

varying aqueous conditions. Investigating freshwater mussel adhesive proteins expands

the current understanding of aquatic bioadhesion and offers potential advancements in

wet adhesive technology. The discovery of a DOPA-deficient wet adhesive protein

derived from the invasive quagga mussels presents a paradigm shift to the currently

known mechanisms of byssal bioadhesion, thereby expanding the repertoire of wet

adhesive strategies. Further, understanding the mechanism of adhesion employed by

this biofouling species can help with the design of anti-fouling strategies to mitigate the

impacts of these animals on the infrastructure and ecosystems in the Great Lakes and

connected freshwater systems in North America.

Introduction

Adhesives found in nature perform in ways that synthetic adhesives have yet to

match. For example, even though many aquatic organisms have adapted to adhere to

surfaces underwater, it remains an engineering challenge to synthesize strong,

biocompatible adhesives that are effective in wet conditions. One strategy that has been

developed in nature over millions of years of natural selection is adherence via a

structure called the byssus. The byssus is utilized by several bivalves, including marine

mussels, oysters, and a few freshwater mussels, to permanently attach themselves and

support their sessile lifestyle (1). The byssus is composed of individual proteinaceous

threads extending from the body of the animal, with each thread ending distally in a

flattened plaque. The bulk of the plaque material connects the thread to the adhesive

proteins found at the plaque-substrate interface. These interfacial proteins bind so

strongly that when plaques are detached, a residue called the footprint usually remains

(2–5).

Originating from the Dnieper River in Eastern Europe and found more recently in

the Great Lakes basin and connected waterways in North America, quagga mussels

(*Dreissena bugensis*) and zebra mussels (*Dreissena polymorpha*) are invasive species

that are among only a handful of freshwater byssates (6). Although superficially similar

to that of the well-studied marine byssates, the biochemical composition of the

Dreissenid byssus is unique; *Mytilid* byssal proteins and *Dreissenid* byssal proteins

share very little homology (7, 8). Moreover, amounts of peptidyl 3,4-

dihydroxyphenylalanine (DOPA) in *Dreissenid* byssal proteins are considerably lower

than in marine mussels and are distributed evenly between the byssal thread and

plaque rather than concentrated in the plaque (9). This suggests that in the context of

protein-substrate interactions, *Dreissenids* may have evolved to employ a unique

mechanism of adhesion that relies less heavily on DOPA-based interfacial interactions

than their marine counterparts.

Byssal adhesive proteins that have been characterized in the current literature

are predominantly in the marine context, i.e. in ocean environments with high ionic

composition (1). The best studied system is the byssus of marine mussels of the genus

Mytilus (10–14). Following the discovery of the significant role of DOPA in interfacial

adhesion in marine mussels and other marine bioadhesive systems, DOPA-based

adhesion has been established as a central paradigm for solutions to the problem of

biocompatible wet adhesion (15–18). Subsequently, several materials utilizing DOPA-

based chemistry for adhesion and cross-linking have been developed (19–24, 13, 25–

29). Few of these materials, however, have translated to the clinical setting for medical

adhesives, in part due to the propensity for catechol to oxidize into the non-adhesive

DOPA-quinone form when exposed to pH values higher than 6.8 (30–32, 25). Aside

from translational considerations, the direct effect of ionic strength in the adhesive

mechanism of DOPA-based bioadhesion is still unclear (33–35). Studying bioadhesives

from freshwater systems can help expand the current repertoire of biocompatible wet

adhesives.

Sequencing of the quagga mussel foot transcriptome combined with bottom-up

proteomics led to the determination of fourteen novel *Dreissena bugensis* foot protein

sequences (Dbfp4 to Dbfp17), and more complete sequences of known Dbfp1 and

Dbfp2 (8, 9, 36). The location and function of each protein, however, remains elusive. Of

particular interest are the proteins at the plaque-substrate interface, where a 10 to 20

nm thick electron-dense footprint protein layer is present, as observed by transmission

electron microscopy (TEM) (2). Here, we report the relative enrichment of the proteins in

the bulk plaque and footprint via quantitative proteomics. This method allowed us to

identify the proteins significantly enriched in the footprint, signifying potential adhesive

ability. Among this set of footprint proteins is Dbfp7, a family of proteins that: matches

the molecular weights of proteins detected by matrix-assisted laser desorption/ionization

time-of-flight (MALDI-TOF) mass spectrometry (37), has been shown to be a significant

portion of the byssus (8), and herein is shown to be adhesive in aqueous conditions

despite lacking DOPA.

Results

Quantitative liquid chromatography tandem mass spectrometry (LC-MS/MS) was

used to identify the byssal proteins that are relatively more abundant in the footprint

compared to the bulk plaque extracts. Quagga mussels collected from Lake Ontario

were allowed to lay fresh byssi onto glass slides overnight. Footprint (FP) and bulk

plaque (BP) proteins were separately solubilized in extraction buffer following scraping

of the bulk plaque, which leaves behind a "footprint" residue on the slide (**Figure 1A**)

(3–5, 8, 9). The footprint and bulk plaque solubilized proteins were digested with trypsin

and the peptides were subsequently functionalized with tandem mass tag (TMT) labels.

The tagged peptides were pooled then sequenced via LC-MS/MS (**Figure 1B**). The

peptide sequences were fingerprinted against the quagga mussel foot transcriptome (8),

searched against protein databases using the basic local alignment search tool

(BLAST) algorithm for homology assessment and identification of proteins, and

compared to proteins identified from whole-thread/plaque byssal material that was

induced to be secreted by potassium chloride (KCl) injection in the pedal ganglion

(IND); the induced byssus dataset has been previously reported (8).

Identification of plaque proteins in the naturally secreted quagga mussel byssus

The induced byssus includes proteins composing the entire byssus, i.e. thread

proteins, bulk plaque proteins, and footprint proteins. In the naturally secreted samples,

bulk plaque and footprint samples were collected in separate groups, and the threads

were discarded. The plaque protein collection method cannot precisely separate the

bulk plaque from the footprint material. As a result, footprint proteins are expected to be

present in the bulk plaque group (albeit to a lesser extent) and vice versa. For this

reason, the 'Footprint + Bulk Plaque' group and the 'ALL' group were considered to most

reliably contain proteins localized to the plaque (**Figure 1D**). These proteins were

identified by the BLASTp algorithm and include (full list in **Table S1**): Dbfp's and

proteins with high homology to known enzymes, structural proteins, matrix proteins,

protease inhibitors, mucin-related proteins, immune/defense proteins, calcium-binding

protein, and rhamnose-binding protein.

Quantitative proteomics classifies 21 footprint-enriched byssal proteins

Several proteins were found to be enriched in the footprint relative to the bulk

plaque (**Figure 2**). The proteins that were measured to have a footprint-to-bulk plaque

fold change ratio (FCR footprint : bulk plaque) > 2 that is statistically significant (p-value

< 0.05) are likely to play direct roles in the adhesive mechanism since they are enriched

at the interface between the byssus and the substrate.

There were 22 significantly enriched footprint proteins identified (**Table 1**). Among

these are three variants of Dbfp7 and one variant of Dbfp8. The other proteins are

weakly (e-score > e^{-6}), moderately (e^{-30} < e-score < e^{-6}), or strongly (e-score < e^{-30})

homologous to known proteins, identified by utilizing the BLASTp algorithm against a

clustered non-redundant database (38, 39). Overall, the proteins relatively enriched at

the adhesive interface are: Dbfp7, Dbfp8, kielin/chordin-like protein, zonadhesin/IgGFc-

binding protein, neurotrypsin-like protein, mucin-like protein, peroxidase, plasminogen-

like protein, galaxin-like protein, and several homologues to uncharacterized proteins

from other aquatic organisms. Interestingly, some of the proteins contained at least one

possible hydroxylated tyrosine residue (i.e. DOPA). None of these proteins, however,

were notably abundant with DOPA, corroborating the overall scarcity of DOPA in

Dreissenid byssus (9) or the limited ability to solubilize DOPA-containing proteins, which

can cross-link post-secretion (16, 40–43). Dbfp1 and Dbfp2, previously identified to

have the most DOPA (9), were not present in the footprint nor bulk plaque. One

transcript (comp50166_c1_seq2-2R) did not have a clear protein-coding open reading

frame and was thus excluded as a footprint protein. Parts of the identified peptides of

this transcript are probably components of a different protein with a similar transcript

sequence.

Selection of Dbfp7 as the first footprint protein to characterize

Previously, three quagga mussel byssal proteins were identified via a DOPA-

specific stain on protein extracted from quagga mussel feet and named Dbfp1, Dbfp2,

and Dbfp3 (9). Subsequent characterization of Dbfp1, one of the higher molecular

weight DOPA-proteins, has classified it as a cuticle protein or a structural cohesive

component (9, 44, 36). The role of Dbfp2 remains unclear (8, 9, 45) and no investigation

of Dbfp3 has been done to date.

Proteins of similar molecular weight as Dbfp3 were detected in the plaque by

MALDI-TOF mass spectrometry and were inferred to contain little to no DOPA (37).

Dbfp3 was reclassified as Dbfp7 upon the construction of the quagga mussel foot

transcriptome, which revealed significant homology of this protein to the known zebra

mussel protein Dpfp7 (8, 46). Herein, the Dbfp7 protein family was identified to be

among the proteins that are significantly enriched in the footprint. Taken together this

suggests that Dbfp7 plays an important role in byssal adhesion, potentially having a role

in bonding to a surface. Dbfp7 was thus selected as the first adhesive protein candidate

to be extracted and further characterized.

<u>Dbfp7 is polymorphic and DOPA-deficient</u>

Dbfp7 was extracted from quagga mussel phenol glands (~ 5 µg per gland) via

several steps of selective precipitation and reversed-phase high-pressure liquid

chromatography (HPLC) (**Figure S1**) (9). The most abundant residue of Dbfp7 is

glycine, followed by significant amounts of proline, asparagine, and tyrosine. There are

13 known variants of Dbfp7 detected in naturally secreted byssi, induced byssi, and

purified Dbfp7, with lengths ranging from 82 to 132 residues (**Figure 3A**). These

variants can be broadly classified as either acidic or basic variants, determined by the

theoretical isoelectric point of the sequence. The acidic variants all contain a 28-residue

sequence in the C-terminal half of the protein sequence, which is missing from the basic

variants. Most of the variation of the Dbfp7 sequence occurs within the first 50 residues;

the C-terminal half of Dbfp7 is highly conserved (aside from the presence or absence of

the 28-residue sequence).

The amount of DOPA in Dbfp7 was found to be notably scarce. Only 1-2 LC-

MS/MS sequences of Dbfp7 peptides (in either purified, naturally secreted, or induced

instances) confidently contained DOPA, depending on the variant and the extracted

pool. The same site of tyrosine hydroxylation was not conserved between induced,

purified (pre-secreted), and naturally secreted forms. Most detected hydroxylation

modifications were ambiguous (**Figure S4**). Specifically, the identified putative DOPA

residues were in spectra of peptides that were not fully fragmented; thus, it is possible

the detected hydroxylation was on a different residue within the peptide (e.g.

hydroxylysine). Further, purified Dbfp7 is only slightly positively stained with nitroblue

tetrazolium (NBT) (**Figure 3B**). The amount of DOPA in Dbfp7 was calculated to be 0.11

± 0.02 mol% (n = 3, s.d.), about 1 DOPA residue per 10 Dbfp7 molecules (**Figure 3C**). It

is possible that Dbfp7 and other plaque proteins are hydroxylated extracellularly by

tyrosinase, which has been found in the bulk plaque (**Table S1**). Interestingly, the

chromatographic fraction of the S2 mixture (**Figure S5C**) eluted before the Dbfp7

elution (40-45% acetonitrile), i.e. "pre-Dbfp7 HPLC fraction", is NBT-positive (**Figure 3B**

and **Figure S5A**). This signifies that DOPA can be detected following the extraction

method via HPLC. Gel electrophoretic separation of this fraction reveals the presence of

low molecular weight proteins with similar weights of purified Dbfp7 (**Figure S5B**). Thus,

it is possible that the proteins in the strong low molecular weight NBT-positive band

detected by Rzepecki and Waite (9), so-called Dbfp3, was not primarily Dbfp7, but

rather was mostly a different protein in the extracted mixture. In any case, the

population of Dbfp7 extracted from phenol glands (pre-secretion) characterized herein is

deficient in DOPA.

Purified forms of Dbfp7 from phenol glands are significantly lower in molecular

weight than that of the full theoretical length of the sequence despite the detection of

peptides from the full sequence within the purified sample (**Figure S1** and **Figure S2**).

Curiously, the localization method of Dbfp7 reveals that although it is overall relatively

more abundant in the footprint, the peptides of Dbfp7 detected in the bulk plaque

consistently map to the C-terminal portion and the peptides in the footprint mostly map

to the N-terminal portion (**Figure S3**). This suggests that Dbfp7 is post-translationally or

post-secretionally cleaved. The population of Dbfp7 analyzed herein, which is extracted

pre-secretion, also does not contain the full-length molecule (**Figure S2**), therefore,

Dbfp7 is likely to be cleaved intracellularly. This raises the possibility that the two

portions of Dbfp7 have differing functions, where the N-terminal region interacts with

substrates while the C-terminal region is responsible for protein-protein interactions

resulting in cohesion between the adhesive and the bulk plaque material. Alternatively, it

is possible that C-terminal peptides interacting with the substrate were unable to be

extracted from the footprint.

Preliminary structural investigations of Dbfp7 reveal both ordered and disordered

structure. Due to the presence of multiple variants, structural measurements are an

average of the Dbfp7 population extracted from phenol glands. To characterize the

structure of Dbfp7, circular dichroism (CD) spectra of Dbfp7 dissolved in 5% acetic acid

(**Figure 4A**) were deconvoluted by Contin-LL (47). The proportions of structural features

were $3.3 \pm 0.3\%$ α-helix, $33 \pm 5\%$ β-strand, $20 \pm 3\%$ β-turns, and $44 \pm 5\%$ disordered

(**Figure 4B**). While almost half of the protein was found to be disordered, significant

portions of the sample adopted β-strand and β-turn structures. To predict the structure,

the AlphaFold protein folding algorithm was used. Due to caveats imposed by the lack

of homology of Dbfp7 to other known proteins, pLDDT scores used to determine model

confidence were very low (<0.5). The structure is predicted to be entirely disordered;

however, the low homology of Dbfp7 to known proteins may result in the prediction

algorithm missing ordered regions. Qualitative observations of the model show a C-

terminal region that is slightly more compact than the N-terminal portion, which looks

relatively extended (**Figure 4C**).

<u>Dbfp7 is adhesive in aqueous conditions</u>

To assess morphology on a surface and adhesive ability, Dbfp7 was drop casted

onto freshly cleaved mica and imaged in aqueous conditions using an atomic force

microscope (AFM) in PeakForce tapping mode with quantitative nanomechanical

mapping (PFQNM). At 100 μg/mL, a dense layer of round and flat globules (ranging

from 3.5 to 35 nm in height) was found throughout the surface (**Figure 4D**).

The surface containing Dbfp7 was adhesive, with some portions of relatively

lower measured adhesion force that correspond to larger globules in the height map

(**Figure 4D** and **Figure 5A**). Overall, Dbfp7 is relatively more adhesive than the control

protein, bovine serum albumin (BSA), known to be non-adhesive (**Figure 5C**) (48–51).

The average forces of adhesion for Dbfp7 and BSA were measured to be 470 ± 160 pN

and 160 ± 140 pN, respectively (s.d. and n = 262,144). Interestingly, the space between

Dbfp7 globules (so-called interglobular space) is more adhesive than the globules

themselves (**Figure 5B**). This may be due to the inconsistent indentation pressure

during scanning, since the setpoint was kept constant during data acquisition.

Alternatively, the interglobular space may contain single molecules of Dbfp7 that are

more readily available for adhesion to the AFM tip compared to self-assembled

globules. This trend is not present in BSA samples (**Figure S6**).

The DMT (Derjaguin-Muller-Toporov) and JKR (Johnson-Kendall-Roberts)

models of adhesion predict different contact areas between two surfaces depending on

the deformation of the indenting probe (52–54). Depending on the model used, the

adhesion energy from the measured adhesion force may vary, and therefore, the

appropriate model is required to convert the measured adhesion force to adhesion

energy (55–57). The Tabor parameter approximates the appropriate model based on the

radius and moduli of the surfaces allowing for the determination of the appropriate

theory for the materials in the analyzed system. The Tabor parameter for this

experimental set-up was calculated to be ~0.12 (see Supplemental Information for full

calculation details). This value means that the contact in this system lies between the

characteristic contact regimes described by the two models. Currently, there is no model

to adequately describe a system that is characteristic of contact partially described by

either the DMT or JKR contact models. Therefore, we report the converted adhesion

energy using both the DMT and the JKR models (**Figure 5D**) to compare to adhesive

proteins in current literature. The calculated adhesion energy of Dbfp7 is 4.4 ± 1.8

mJ/m^2 (JKR) or 3.3 ± 1.3 mJ/m^2 (DMT) which is ~5 times higher than BSA (0.90 ± 0.89

mJ/m^2 (JKR) or 0.67 ± 0.66 mJ/m^2 (DMT)) (s.d., N = 2,859,269 and 793,084 for Dbfp7

and BSA, respectively). Interestingly, the adhesion energy value for Dbfp7 is in a similar

range to the adhesion energy of Mfp3, one of the *Mytilus* adhesive proteins (~3-6 mJ/m^2

determined by surface forces apparatus (SFA) measurements) (14). However, the other

known *Mytilus* interfacial protein, Mfp5, remains the strongest known byssal adhesive

protein with the most relative amount of DOPA (15 mJ/m^2 and 30 mol% DOPA) (14).

Discussion

Several plaque proteins were identified to be relatively more enriched in the

footprint than in the bulk plaque material, suggesting they may play direct roles in

plaque-substrate adhesion. Among these, Dbfp7 stood out due to its detection in

previous studies on quagga mussel byssal proteins. When extracted from phenol glands

(pre-secretion), Dbfp7 adopts a disordered structure with significant proportions of β-

strand and β-turn features and contains little to no DOPA. Despite this lack of DOPA,

adhesion measurements by AFM performed in aqueous conditions revealed that Dbfp7

exhibits wet adhesion similar to Mfp3, one of the marine mussel adhesive proteins.

Dbfp7 is thus classified as a DOPA-deficient wet adhesive protein.

The marine *Mytilids* and the freshwater *Dreissenids* evolutionarily diverged at the

class level; *Dreissenids* have since evolved in brackish water originally and eventually

adapted to freshwater habitats (58). It is evident that the significant amount of time

between common ancestors gives rise to a variety of differences in the byssal adhesion

mechanism. For example, another freshwater byssate *Limnoperna fortunei* (golden

mussel) – found in the freshwater systems of Japan, Korea, and Eastern China –

diverges from *Mytilus sp*. at the family level and is thus genetically more similar. Indeed,

it has been found that the golden mussel byssus contains typical marine mussel

adhesive protein domains such as EGF domains, vWFA domains, and GXX motifs (59).

Further, DOPA was found in the outer sheath of golden mussel byssal threads and at

the bottom of adhesive plaques by NBT staining (59). The presence of DOPA at the

adhesive interface suggests DOPA-based interfacial chemistry, however the specific

golden mussel footprint proteins have not been characterized yet. In the zebra mussel

byssus, DOPA and catechol oxidase was found throughout the plaque, suggesting that

some interfacial *Dreissenid* footprint proteins may employ DOPA interfacial chemistry

while bulk plaque and thread DOPA-proteins contribute to cross-linking (60). However,

the adhesion mechanism of Dbfp7 may be less dependent on DOPA, given the even

distribution of DOPA between the thread and plaque.

In the marine mussel, the proteins found at the plaque-substrate interface have

significantly higher relative amounts of DOPA compared to the bulk plaque and thread

proteins. In *Dreissenids*, it was found that peptidyl-DOPA is present in low amounts

throughout the byssus, including the thread (9). This suggests that the role of DOPA in

the freshwater system is primarily for cross-linking purposes (16, 40–43). Dbfp7 purified

from phenol glands of quagga mussels has herein been shown to lack significant

amounts of DOPA while being more adhesive than BSA under the same conditions

(**Figure 5**). Conversion of the adhesive forces to adhesion energy reveals similar

adhesion energy of Dbfp7 compared to one of the marine mussel interfacial proteins,

Mfp3. Although similar in molecular weight with abundant glycine, tyrosine, proline, and

asparagine residue proportions, Dbfp7 and Mfp3 share very little sequence homology

and Mfp3 is more enriched with DOPA (around 10-20 mol%) compared to Dbfp7 (~0.1

mol%). The other known interfacial *Mytilus* protein, Mfp5, contains even more DOPA (30

mol%) than Mfp3 and is significantly more adhesive than both Mfp3 and Dbfp7 (15

mJ/m^2 measured by SFA) (14). It appears that some tyrosine residues in Dbfp7 may be

post-secretionally hydroxylated within the distal depression and secreted byssus, since

tyrosinases were identified in the bulk plaque (**Table S1**). Still, the DOPA-deficient pre-

secretion population of Dbfp7 is adhesive.

The mechanism of adhesion for Dbfp7 remains unknown and further studies into

specific peptide sequences and post-translational modifications are required to develop

a model for the adhesion mechanism. It is possible that in environments with lower ionic

strength, as in freshwater habitats, lower amounts of DOPA can achieve similar

adhesive strength. Further, several efforts are being made to understand the role of

lysine and other cationic moieties in conjunction with DOPA-based adhesion,

particularly in cation-pi interactions for cross-linking and complex coacervation (34, 61–

67). It is unclear whether the dihydroxyl moiety in catechol is necessary for this

interaction or if the phenyl moiety in residues such as tyrosine, phenylalanine, and

tryptophan is sufficient (68, 69). Charged residues may also play a role in dislodging

surface-bound water molecules and surface ions to allow for direct contact with a

surface (34). Indeed, Dbfp7 contains significant amounts of aromatic residues tyrosine

and tryptophan, as well as positively charged lysine (**Figure 3A**).

Dbfp7 is polymorphic with up to 13 detected variants. Several genes of the zebra

mussel homologue of Dbfp7, Dpfp7, have been identified in multiple regions of the

zebra mussel genome (70). Mfp3 is similarly polymorphic, with up to 35 known variants

encoded throughout the *Mytilus sp.* genome (5). The existence of multiple variants of a

gene that is preserved several times in the genome suggests vital importance.

Particularly, polymorphism of an interfacial protein may enhance the versatility of

adhesion to different kinds of surfaces (3, 5, 8, 16). Further, Mfp3 is known to exist as

two populations, Mfp3S and Mfp3F, so-called the slow and fast variants due to their

elution time in reversed-phase HPLC, thereby differing in hydrophobicity. Mfp3S was

found to be more resistant to oxidation and retains adhesion when pH is equilibrated to

seawater levels (7, 71). The role of the two variant types of Dbfp7 (acidic and basic) is

currently unclear and may: (i) function similarly to the Mfp3 variants, (ii) have different

roles in the footprint-bulk plaque and footprint-substrate interfaces, or (iii) contribute to

spreading behaviour on a surface by the formation of complex coacervates during

secretion (72–74). Encoding and synthesizing these two peptide components in the

same transcript may have the advantage of maintaining stoichiometric proportions.

The quagga mussel footprint contains 20 proteins in addition to Dbfp7 that

contribute to byssal adhesion, possibly in interfacial interactions (**Table 1**). Further

investigation is needed to understand the interactions between these proteins with each

other, with bulk plaque proteins, and with adherend substrates. Of particular interest are

the zonadhesin/IgGFc-binding proteins, homologues of which were found in the sea star

(*Asteria rubens*) adhesive footprint (75). Further, kielin/chordin-like proteins match the

molecular weights of interfacial plaque proteins identified by MALDI-TOF (37). With the

advent of high-throughput sequencing techniques, the library of proteins found in

bioadhesive systems is vast. Little has been done to localize these proteins at the direct

adhesive-substrate interface *in situ*, with a larger current focus on bulk cohesive

properties than on interfacial adhesion chemistry. Localization of proteins at direct

adhesive-substrate interfaces will provide a smaller subset to analyze for understanding

the fundamental mechanisms of bioadhesion in aqueous environments.

It is difficult to directly compare adhesive measurements of Dbfp7 to other studies

since the methods of wet adhesive protein characterization are varied. SFA developed

by Israelachvili and applied by Waite and others to marine byssal proteins has been the

predominant method of adhesive measurements (14, 34, 69, 76–79). This unique

instrument is not as widely implemented compared to AFM. Although similar in

fundamental operation (i.e. generation and analysis of submicron contact force-distance

curves), the planar scale of these two systems is different, with SFA analyzing at the

squared milli- to centimetre contact area range and AFM analyzing at the squared nano-

to micrometre contact area range. The classical models of contact (Hertz, DMT, JKR,

etc.) may allow for the comparison of adhesion energies obtained from different

systems; however, the physics at these two scales may be fundamentally different and

thus may not be directly comparable. To date, the adhesion in systems that lie between

DMT and JKR contact models is yet to be fully understood (55, 56, 80, 81). Further

complications arise when the two contacting surfaces are not entirely flat or smooth (57,

82). Advancements in theoretical contact physics at the nanoscale are needed to

harmonize any discrepancies that may arise from measuring adhesion at differing size

scales. The problem of scale may be circumvented with the use of colloidal probe

technique, allowing for the analysis of adhesion at the micrometre contact area scale in

AFM (83, 84). This technique, however, cannot be used for concurrent imaging and

characterization of mechanical properties. Further, the adhesion measurements

obtained herein presumably represent adhesion of Dbfp7 with the silicon nitride AFM tip.

It is possible that some adhesion measurements were from Dbfp7 bound to the tip,

resulting in a measurement of the interactions between Dbfp7-to-Dbfp7 or Dbfp7-to-

mica. Since the protein seems to aggregate resulting in a non-uniform molecular layer

and due to the limited lateral resolution of AFM imaging, these possible interactions may

contribute to the observed adhesion average.

This study highlights the importance of studying aqueous bioadhesive proteins

from a diversity of organisms. To our knowledge, Dbfp7 is the first freshwater wet

adhesive protein to be functionally characterized. The lack of DOPA in this protein is

interesting but not shocking when considering the evolutionary history of *Dreissenids*

compared to the other bioadhesive systems studied so far. The inclusion of Dbfp7 (and

potentially others from the 20 footprint proteins herein localized) to the repertoire of wet

bioadhesive proteins will allow for the identification of the fundamental properties

required to achieve biocompatible wet adhesion for the development of bio-inspired wet

adhesive materials. Further, understanding the specific mechanism of adhesion

employed by *Dreissenids* that are invasive species to the Great Lakes basin and

connected waterways in North America will aid in the development of anti-fouling

strategies (85).

Materials and Methods

Quagga mussel collection and maintenance

Freshwater mussels were harvested from Lake Ontario at Port Credit Harbour

Marina in Mississauga, Ontario. Live mussels were cleaned and sorted by species

(quagga mussels and zebra mussels) and transferred to tanks with circulating artificial

lake water (0.30 mM sodium bicarbonate, 0.13 mM potassium bicarbonate, 0.35 mM

sodium chloride, 0.67 mM magnesium sulfate, 0.08 mM magnesium chloride, 1.47 mM

calcium carbonate) at 12ºC and fed powdered green algae.

Plaque protein collection

Quagga mussels were first detached from their original substrates (tank walls or

other mussels), secured into individual cages with glass slides, and allowed to lay byssi

overnight. Mussels were gently detached from the substrates by cutting the byssal

threads, leaving the plaque intact on the glass slide. Bulk plaque material was scraped

off with a razor blade and collected in a pool of extraction buffer (composed of 200 mM

sodium borate, 4 M urea, 1 mM potassium cyanide, 1 mM ethylenediaminetetraacetic

acid, and 10 mM ascorbic acid, at pH 8.0). Extraction buffer was deposited onto the

footprint residue remaining on the slide, incubated for 1-2 minutes, then collected.

Samples were homogenized in a 1 mL tissue grinder on ice, centrifuged at 17,000 x g

for eight minutes at 4ºC, then flash frozen for storage until analysis. Amounts of bulk

plaque and footprint pools were normalized by weight (resulting in 25 bulk plaques and

400 footprints in total for comparative analysis).

Tandem mass spectrometry on digested peptides extracted from freshly secreted byssus

429
430

The bulk plaque and the footprint pools were digested with 13 ng/µL trypsin

(Porcine, sequencing grade, Promega) overnight at 37ºC. Following digestion, the bulk

plaque peptides were functionalized with TMT-2plex TMT-131 and footprint peptides

were functionalized with TMT-2plex TMT-126. The peptide pools were then pooled and

loaded onto a 100 µm internal diameter (ID) pre-column (Dionex) at 4 µL/min and

separated over a 50 µm ID analytical column (C18 2 µm, Dionex). The peptides were

eluted on a 0 to 35% acetonitrile gradient with an EASY-nLC 1000 nano-

chromatography pump (Thermo Fisher, Odense Denmark). Data was acquired at

70,000 FWHM resolution in the MS mode and 17,500 FWHM resolution in the MS/MS

mode. A total of 10 MS/MS scans were obtained per MS cycle.

In silico methods for protein identification, selection, and sequence analysis

LC-MS/MS data was analyzed with PEAKS Studio 10.5 (Bioinformatics Solutions

Inc.) and Scaffold 5.0 (Proteome Software Inc.). *De novo* sequences derived from

MS/MS data were matched against the quagga mussel foot transcriptome (8) with

tyrosine hydroxylation to DOPA, deamidation, oxidation, and carbamidomethylation set

as variable modifications. Parent ion and fragment ion mass tolerances were set to 5

PPM and 0.01 Da, respectively. Acceptance criteria for identified proteins were peptide

logP ≥ 15, protein –10 logP ≥ 50, and *de novo* average local confidence (ALC) score ≥

80%, with at least two peptide spectra per identification.

Scaffold Q+ in Scaffold 5.0 was used to quantitate label-based quantitation

peptide and protein identifications. Peptide identifications were accepted if they could be

established at ≥ 75.0% probability to achieve an FDR < 0.1% and contained at least two

identified peptides. Normalization against the reference (bulk plaque) were performed

iteratively across samples and spectra on intensities, as described in (86). Spectra data

were log-transformed and weighted by an adaptive intensity weighting algorithm.

Differentially expressed proteins were determined by applying the permutation test with

unadjusted significance level $p < 0.05$ corrected by the Benjamini-Hochberg procedure.

The Protein Basic Local Alignment Search Tool (BLASTp) algorithm was applied

on the identified proteins of interest against a clustered non-redundant database (38,

39) for homology assessment. Alignment of sequences were visualized in Jalview

2.11.3.3 using the Clustal Omega multiple sequence alignment program. The

ProtParam online tool (Expasy) was used to calculate residue frequency, molecular

weight, and isoelectric point of full-length protein sequences obtained from virtual

translation of the open-reading frame of identified transcripts.

The method for analyzing tyrosine hydroxylation was categorized based on the

level of confidence in the identified modification. A modification was considered

confident if the tyrosine residue (Y) was highlighted with a specific residue number by

the PEAKS PTM algorithm. It was classified as semi-confident if no number was

indicated, but the spectrum contained a Y+16.99 m/z modification in both the b- and y-

ions, indicating hydroxylation at that residue. Finally, a non-confident classification was

assigned if a number was not indicated and the spectrum only showed the Y+16.99 m/z

within an unfragmented peptide, suggesting that the tyrosine residue was not

individually sequenced.

Purification of Dbfp7

Dbfp7 was purified from quagga mussel phenol glands (**Figure S1**). The crude

protein extraction method was adapted from (9). Phenol glands were dissected out of

freshly harvested quagga mussels and homogenized in 5% acetic acid with 10 µM

pepstatin and leupeptin protease inhibitors. The mixture was centrifuged at 20,000 x g

for 40 minutes at 4ºC. The pellet (P1) was then homogenized in 5% acetic acid and 8 M

urea. The mixture is then centrifuged at 20,000 x g for 40 minutes at 4ºC. The

supernatant (S2), i.e. the urea-soluble proteins, was then mixed with mobile phase

(0.1% trifluoroacetic acid (TFA)) and injected into a C4 reversed-phase column (XBridge

Protein BEH C4, 300 Å). The eluent at 40 to 45% acetonitrile (with 0.1% TFA) was

collected, lyophilized, and stored at -80ºC until analysis.

Structure Prediction with AlphaFold

The structure of Dbfp7 variant 1 was predicted with AlphaFold2 protein structure

prediction (87). The query sequence was input in the ColabFold v1.5.5 python notebook

and the resulting structures were visualized with ChimeraX v1.8.

Gel electrophoresis

Samples were separated by molecular weight via gel electrophoresis in Bolt 12%

Bis-Tris precast gels, using Bolt MES-SDS running buffer and Bolt LDS sample buffer.

Gels were run at constant 165 V voltage for 40 minutes. The gels were fixed in 40%

methanol + 10% acetic acid and stained with Coomassie Brilliant Blue R-250 for 1 hour.

Gels were destained and visualized with a digital camera.

ESI mass spectrometry

Dbfp7 was dissolved in 5% acetic acid at 400 µg/mL and ionized by electrospray

ionization. The sample was analyzed with an Agilent 6538 UHD hybrid quadrupole time-

of-flight mass spectrometer. The mass spectrum was processed using the Maximum

Entropy algorithm implemented in the MassHunter BioConfirm software package.

<u>Nitroblue tetrazolium staining for DOPA detection</u>

NBT was used to detect and quantify DOPA amounts. This protocol was adapted

from (9, 88). Samples were prepared by adding 20 µL to 0.17 mM NBT in 2 M K-

glycinate, then incubating at room temperature in the dark for 1 hour. The absorbance at

530 nm was measured and compared to a calibration curve generated using samples

prepared with standard amounts of peptidyl DOPA. Peptidyl DOPA standards were

synthesized by Biomatik and include the first 13 residues of Dbfp7, with the penultimate

tyrosine replaced with DOPA.

Dot blots were prepared by depositing 10 µL of sample onto a nitrocellulose

membrane and allowing it to dry in air for 30 minutes. The membrane was then rinsed

with potassium glycinate for 30 minutes, then stained with 0.24 mM NBT for 1.5 hours in

the dark, then rinsed with 0.2 M sodium borate for 1 hour.

<u>Sample preparation and spectra analysis for circular dichroism</u>

To characterize the secondary structure of purified Dbfp7, far-UV circular

dichroism was performed. The CD spectra was recorded using a Jasco

spectropolarimeter, with a 2-nm bandwidth, scanning speed of 20 nm/min at a spectral

resolution of 1.0 nm. The scans were averaged over ten scans, and the data was

smoothed. The CD spectra of the buffer was recorded and manually subtracted from

sample spectra. The secondary structure was characterized by computational analysis

of the spectra using the Contin-LL program from the Dichroweb package (47).

Measured ellipticity was normalized to the molecular weight, pathlength, and

concentration of the sample to get the mean residue ellipticity (deg*cm^2/dmol). The

average molecular weight of 7.2 kDa, as determined by mass spectrometry, was used.

<u>Sample preparation for atomic force microscopy</u>

10 µL of sample was deposited onto freshly cleaved mica (V1 high grade) and

incubated in a humidified chamber for 20 minutes. Residual protein was thoroughly

washed away with blank solution (e.g. 5% acetic acid for Dbfp7 samples dissolved in

5% acetic acid). A droplet of deionized water was dropped onto the sample and placed

in a fluid cell for AFM imaging in aqueous conditions.

<u>Imaging and adhesion force mapping via AFM PFQNM</u>

Samples were imaged with a MultiMode Atomic Force Microscope equipped with

a Nanoscope 6 controller operated via Nanoscope 10.0 software in PeakForce tapping

mode with quantitative nanomechanical mapping (PFQNM). SCANASYST-FLUID

probes with 20 nm (nominal) silicon nitride tips were calibrated on blank mica in water,

resulting in calibrated spring constants around 0.7 N/m. The images were taken at 512-

line resolution at 1.0 Hz line rate. ScanAsyst mode allows for automatic control of

setpoint, gain, and scan rate. Initial images were taken with ScanAsyst auto-control on

to determine the ideal setpoint for the sample. ScanAsyst auto-control was

subsequently turned off to allow for direct pixel-to-pixel comparison of nanomechanical

properties derived from the collected force-distance curves. PeakForce data was

captured at a rate such that each height pixel has a corresponding force-distance curve

saved. Images were analyzed using the Nanoscope Analysis 3.0 software. The

individual curves for the entire map were baseline-corrected and moduli were fitted

using the JKR contact model. Adhesion was calculated using the positive-slope

algorithm.

Author Contributions

A.R.O., R.H., D.J.R., J.N., and E.D.S. designed research; A.R.O., R.H., D.J.R., and J.N.

performed research; A.R.O., R.H., D.J.R., J.N., K.M.M.C., and E.D.S. analyzed data; and A.R.O., R.H.,

K.M.M.C., and E.D.S. wrote the paper.

Acknowledgements

This research was supported by Discovery Grant #342455 to E.D.S. from the Natural Sciences

and Engineering Research Council of Canada (NSERC) (RGPIN-2019-06210), Post-Graduate

Scholarship – Doctoral (PGS-D) to A.R.O. from NSERC, and the Queen Elizabeth II Graduate

Scholarship in Science and Technology (QEII-GSST) to A.R.O. from the Government of Ontario. The

authors thank the SickKids Proteomics, Analytics, Robotics & Chemical Biology Centre for assistance with

LC-MS/MS data collection and interpretation, the Advanced Instrumentation for Molecular Structure

Laboratory for ESI MS, and the SickKids Structural and Biophysical Core Facility for CD and HPLC.

Special thanks are extended to Ryan Lee Chan for his assistance with AFM. The authors also gratefully

acknowledge Dr. Magdalena Wojtas for her valuable insights and advice throughout the project.

Competing Interest Statement

The authors declare no competing interest.

Abbreviations/Acronyms/Initialisms

DOPA = 3,4-dihydroxyphenylalanine
Dbfp = *Dreissena bugensis* foot protein
Dpfp = *Dreissena polymorpha* foot protein
MALDI-TOF = matrix-assisted laser desorption/ionization time-of-flight
LC-MS/MS = liquid chromatography tandem mass spectrometry
FP = footprint
BP = bulk plaque
IND = induced byssus
TMT = tandem mass tag
BLAST = basic local alignment search tool
KCl = potassium chloride
FCR = fold change ratio
HPLC = high-pressure liquid chromatography
NBT = nitroblue tetrazolium
pLDDT = predicted local distance difference test
AFM = atomic force microscope
CD = circular dichroism
BSA = bovine serum albumin
DMT = Derjaguin-Muller-Toporov
JKR = Johnson-Kendall-Roberts
SFA = surface forces apparatus
ID = internal diameter
TFA = trifluoroacetic acid

References

1. C. Liu, Comparative Proteomics for an In-Depth Understanding of Bioadhesion Mechanisms and Evolution across Metazoans. *Journal of Proteomics*, **256**, 104506 (2022). https://doi.org/10.1016/j.jprot.2022.104506.

2. N. Farsad, E. D. Sone, Zebra Mussel Adhesion: Structure of the Byssal Adhesive Apparatus in the Freshwater Mussel, *Dreissena Polymorpha*. *Journal of Structural Biology*, **177**(3), 613–620 (2012). https://doi.org/10.1016/j.jsb.2012.01.011.

3. H. Zhao, J. H. Waite, Linking Adhesive and Structural Proteins in the Attachment Plaque of *Mytilus Californianus*. *Journal of Biological Chemistry*, **281**(36), 26150–26158 (2006). https://doi.org/10.1074/jbc.M604357200.

4. J. H. Waite, Adhesion a La Moule. *Integrative and Comparative Biology*, **42**(6), 1172–1180 (2002). https://doi.org/10.1093/icb/42.6.1172.

5. R. Y. Floriolli, J. Von Langen, J. H. Waite, Marine Surfaces and the Expression of Specific Byssal Adhesive Protein Variants in *Mytilus*. *Mar. Biotechnol.*, **2**(4), 352–363 (2000). https://doi.org/10.1007/s101269900032.

6. M. A. McCartney, Structure, Function and Parallel Evolution of the Bivalve Byssus, with Insights from Proteomes and the Zebra Mussel Genome. *Phil. Trans. R. Soc. B*, **376**(1825), 20200155 (2021). https://doi.org/10.1098/rstb.2020.0155.

7. D. G. DeMartini, J. M. Errico, S. Sjoestroem, A. Fenster, J. H. Waite, A Cohort of New Adhesive Proteins Identified from Transcriptomic Analysis of Mussel Foot Glands. *J. R. Soc. Interface*, **14**(131), 20170151 (2017). https://doi.org/10.1098/rsif.2017.0151.

8. D. J. Rees, A. Hanifi, A. Obille, R. Alexander, E. D. Sone, Fingerprinting of Proteins That Mediate Quagga Mussel Adhesion Using a De Novo Assembled Foot Transcriptome. *Sci Rep*, **9**(1), 6305 (2019). https://doi.org/10.1038/s41598-019-41976-7.

9. L. M. Rzepecki, J. H. Waite, The Byssus of the Zebra Mussel, *Dreissena Polymorpha*. II: Structure and Polymorphism of Byssal Polyphenolic Protein Families. *Mol Mar Biol Biotechnol.* 2(5), 267–279 (1993).

10. J. H. Waite, M. L. Tanzer, Polyphenolic Substance of *Mytilus Edulis*: Novel Adhesive Containing L-Dopa and Hydroxyproline. *Science*, **212**(4498), 1038–1040 (1981). https://doi.org/10.1126/science.212.4498.1038.

11. M. Yu, J. Hwang, T. J. Deming, Role of l -3,4-Dihydroxyphenylalanine in Mussel Adhesive Proteins. *J. Am. Chem. Soc.*, **121**(24), 5825–5826 (1999). https://doi.org/10.1021/ja990469y.

12. H. Lee, N. F. Scherer, P. B. Messersmith, Single-Molecule Mechanics of Mussel Adhesion. *Proc. Natl. Acad. Sci. U.S.A.*, **103**(35), 12999–13003 (2006). https://doi.org/10.1073/pnas.0605552103.

13. B. K. Ahn, Perspectives on Mussel-Inspired Wet Adhesion. *J. Am. Chem. Soc.*, **139**(30), 10166–10171 (2017). https://doi.org/10.1021/jacs.6b13149.

14. J. H. Waite, Mussel Adhesion – Essential Footwork. *Journal of Experimental Biology*, **220**(4), 517–530 (2017). https://doi.org/10.1242/jeb.134056.

15. T. H. Anderson *et al*., The Contribution of DOPA to Substrate–Peptide Adhesion and Internal Cohesion of Mussel-Inspired Synthetic Peptide Films. *Adv Funct Materials*, **20**(23), 4196–4205 (2010). https://doi.org/10.1002/adfm.201000932.

16. R. J. Stewart, T. C. Ransom, V. Hlady, Natural Underwater Adhesives. *J Polym Sci B*
Polym Phys, **49**(11), 757–771 (2011). https://doi.org/10.1002/polb.22256.
17. M. L. B. Palacio, B. Bhushan, Bioadhesion: A Review of Concepts and Applications.
Phil. Trans. R. Soc. A., **370**(1967), 2321–2347 (2012).
https://doi.org/10.1098/rsta.2011.0483.
18. Y. Kan, Z. Wei, Q. Tan, Y. Chen, Inter- and Intramolecular Adhesion Mechanisms of
Mussel Foot Proteins. *Sci. China Technol. Sci.*, **63**(9), 1675–1698 (2020).
https://doi.org/10.1007/s11431-019-1541-8.
19. B. P. Lee, P. B. Messersmith, J. N. Israelachvili, J. H. Waite, Mussel-Inspired
Adhesives and Coatings. *Annu. Rev. Mater. Res.*, **41**(1), 99–132 (2011).
https://doi.org/10.1146/annurev-matsci-062910-100429.
20. M. Mehdizadeh, J. Yang, Design Strategies and Applications of Tissue Bioadhesives.
Macromolecular Bioscience, **13**(3), 271–288 (2013).
https://doi.org/10.1002/mabi.201200332.
21. S. Moulay, Dopa/Catechol-Tethered Polymers: Bioadhesives and Biomimetic
Adhesive Materials. *Polymer Reviews*, **54**(3), 436–513 (2014).
https://doi.org/10.1080/15583724.2014.881373.
22. P. Kord Forooshani, B. P. Lee, Recent Approaches in Designing Bioadhesive
Materials Inspired by Mussel Adhesive Protein. *J. Polym. Sci. Part A: Polym. Chem.*,
55(1), 9–33 (2017). https://doi.org/10.1002/pola.28368.
23. L. Li, H. Zeng, Marine Mussel Adhesion and Bio-Inspired Wet Adhesives.
Biotribology, **5**, 44–51 (2016). https://doi.org/10.1016/j.biotri.2015.09.004.
24. A. J. Vernengo, "Adhesive Materials for Biomedical Applications" in *Adhesives -*
Applications and Properties, A. Rudawska, Ed., (InTech, 2016).
https://doi.org/10.5772/64958.
25. G. P. Maier, C. M. Bernt, A. Butler, Catechol Oxidation: Considerations in the Design
of Wet Adhesive Materials. *Biomater. Sci.*, **6**(2), 332–339 (2018).
https://doi.org/10.1039/C7BM00884H.
26. S. Basak, Co-Evolving with Nature: The Recent Trends on the Mussel-Inspired
Polymers in Medical Adhesion. *Biotechnol Bioproc E*, **26**(1), 10–24 (2021).
https://doi.org/10.1007/s12257-020-0234-z.
27. G. M. Taboada *et al.*, Overcoming the Translational Barriers of Tissue Adhesives.
Nat Rev Mater, **5**(4), 310–329 (2020). https://doi.org/10.1038/s41578-019-0171-7.
28. J. H. Waite, M. J. Harrington, Following the Thread: *Mytilus* Mussel Byssus as an
Inspired Multi-Functional Biomaterial. *Can. J. Chem.*, **100**(3), 197–211 (2022).
https://doi.org/10.1139/cjc-2021-0191.
29. X. Ma *et al.*, Hydrogels for Underwater Adhesion: Adhesion Mechanism, Design
Strategies and Applications. *J. Mater. Chem. A*, **10**(22), 11823–11853 (2022).
https://doi.org/10.1039/D2TA01960D.
30. J. Yu, W. Wei, E. Danner, J. N. Israelachvili, J. H. Waite, Effects of Interfacial Redox
in Mussel Adhesive Protein Films on Mica. *Advanced Materials*, **23**(20), 2362–2366
(2011). https://doi.org/10.1002/adma.201003580.
31. J. Yu *et al.*, Mussel Protein Adhesion Depends on Interprotein Thiol-Mediated Redox
Modulation. *Nat Chem Biol*, **7**(9), 588–590 (2011).
https://doi.org/10.1038/nchembio.630.

32. R. Mirshafian, W. Wei, J. N. Israelachvili, J. H. Waite, α,β-Dehydro-Dopa: A Hidden Participant in Mussel Adhesion. *Biochemistry*, **55**(5), 743–750 (2016). https://doi.org/10.1021/acs.biochem.5b01177.

33. R. Chollakup, J. B. Beck, K. Dirnberger, M. Tirrell, C. D. Eisenbach, Polyelectrolyte Molecular Weight and Salt Effects on the Phase Behavior and Coacervation of Aqueous Solutions of Poly(Acrylic Acid) Sodium Salt and Poly(Allylamine) Hydrochloride. *Macromolecules*, **46**(6), 2376–2390 (2013). https://doi.org/10.1021/ma202172q.

34. G. P. Maier, M. V. Rapp, J. H. Waite, J. N. Israelachvili, A. Butler, Adaptive Synergy between Catechol and Lysine Promotes Wet Adhesion by Surface Salt Displacement. *Science* **349**, 628–632 (2015).

35. X. Ou *et al*., Structure and Sequence Features of Mussel Adhesive Protein Lead to Its Salt-Tolerant Adhesion Ability. *Sci. Adv.*, **6**(39), eabb7620 (2020). https://doi.org/10.1126/sciadv.abb7620.

36. K. E. Anderson, J. H. Waite, Immunolocalization of Dpfp1, a Byssal Protein of the Zebra Mussel *Dreissena Polymorpha*. *Journal of Experimental Biology*, **203**(20), 3065–3076 (2000). https://doi.org/10.1242/jeb.203.20.3065.

37. D. J. Rees, A. Hanifi, J. Manion, A. Gantayet, E. D. Sone, Spatial Distribution of Proteins in the Quagga Mussel Adhesive Apparatus. *Biofouling*, **32**(2), 205–213 (2016). https://doi.org/10.1080/08927014.2015.1135426.

38. M. Steinegger, J. Söding, MMseqs2 Enables Sensitive Protein Sequence Searching for the Analysis of Massive Data Sets. *Nat Biotechnol*, **35**(11), 1026–1028 (2017). https://doi.org/10.1038/nbt.3988.

39. S. F. Altschul, W. Gish, W. Miller, E. W. Myers, D. J. Lipman, Basic Local Alignment Search Tool.

40. B. Liu, L. Burdine, T. Kodadek, Chemistry of Periodate-Mediated Cross-Linking of 3,4-Dihydroxylphenylalanine-Containing Molecules to Proteins. *J. Am. Chem. Soc.*, **128**(47), 15228–15235 (2006). https://doi.org/10.1021/ja065794h.

41. L. A. Burzio, J. H. Waite, Reactivity of Peptidyl-tyrosine to Hydroxylation and Cross-linking. *Protein Science*, **10**(4), 735–740 (2001). https://doi.org/10.1110/ps.44201.

42. N. Holten-Andersen *et al*., pH-Induced Metal-Ligand Cross-Links Inspired by Mussel Yield Self-Healing Polymer Networks with near-Covalent Elastic Moduli. *Proc. Natl. Acad. Sci. U.S.A.*, **108**(7), 2651–2655 (2011). https://doi.org/10.1073/pnas.1015862108.

43. M. Mesko *et al*., Catechol-Vanadium Binding Enhances Cross-Linking and Mechanics of a Mussel Byssus Coating Protein. *Chem. Mater.*, **33**(16), 6530–6540 (2021). https://doi.org/10.1021/acs.chemmater.1c02063.

44. K. E. Anderson, J. H. Waite, Biochemical Characterization of a Byssal Protein from *Dreissena Bugensis* (Andrusov). *Biofouling*, **18**(1), 37–45 (2002). https://doi.org/10.1080/08927010290017716.

45. A. Gantayet, L. Ohana, E. D. Sone, Byssal Proteins of the Freshwater Zebra Mussel, *Dreissena Polymorpha*. *Biofouling*, **29**(1), 77–85 (2013). https://doi.org/10.1080/08927014.2012.746672.

46. A. Gantayet, D. J. Rees, E. D. Sone, Novel Proteins Identified in the Insoluble Byssal Matrix of the Freshwater Zebra Mussel. *Mar Biotechnol*, **16**(2), 144–155 (2014). https://doi.org/10.1007/s10126-013-9537-9.

47. A. J. Miles, S. G. Ramalli, B. A. Wallace, DichroWeb, a Website for Calculating Protein Secondary Structure from Circular Dichroism Spectroscopic Data. *Protein Science*, **31**(1), 37–46 (2022). https://doi.org/10.1002/pro.4153.

48. J. J. Valle-Delgado *et al*., Interactions between Bovine Serum Albumin Layers Adsorbed on Different Substrates Measured with an Atomic Force Microscope. *Phys. Chem. Chem. Phys.*, **6**(7), 1482–1486 (2004). https://doi.org/10.1039/B314060C.

49. M. S. Wang, L. B. Palmer, J. D. Schwartz, A. Razatos, Evaluating Protein Attraction and Adhesion to Biomaterials with the Atomic Force Microscope. *Langmuir*, **20**(18), 7753–7759 (2004). https://doi.org/10.1021/la049849+.

50. Y. Yu, P.Q. Ying, G. Jin, Competitive Adsorption between Bovine Serum Albumin and Collagen Observed by Atomic Force Microscope. *Chinese Chemical Letters*, **15**(12), 1465–1468 (2004).

51. A. P. Serro, K. Degiampietro, R. Colaço, B. Saramago, Adsorption of Albumin and Sodium Hyaluronate on UHMWPE: A QCM-D and AFM Study. *Colloids and Surfaces B: Biointerfaces*, **78**(1), 1–7 (2010). https://doi.org/10.1016/j.colsurfb.2010.01.022.

52. B.V. Derjaguin, V.M. Muller, Y.P. Toporov, Effect of Contact Deformations on the Adhesion of Particles. *Journal of Colloid and Interface Science*, **53**(2), 314–326 (1974).

53. V. M. Muller, B.V. Derjaguin, Y. P. Toporov, On Two Methods of Calculation of the Force of Sticking of an Elastic Sphere to a Rigid Plane. *Colloids and Surfaces*, **7**, 251–259 (1983).

54. K. L. Johnson, K. Kendall, A. D. Roberts, Surface Energy and the Contact of Elastic Solids. *Proc. R. Soc. Lond. A*, **324**, 301–313 (1971).

55. D. S. Grierson, E. E. Flater, R. W. Carpick, Accounting for the JKR–DMT Transition in Adhesion and Friction Measurements with Atomic Force Microscopy. *Journal of Adhesion Science and Technology*, **19**(3–5), 291–311 (2005). https://doi.org/10.1163/1568561054352685.

56. X. Gao, F. Hao, Z. Huang, D. Fang, Mechanics of Adhesive Contact at the Nanoscale: The Effect of Surface Stress. *International Journal of Solids and Structures*, **51**(3–4), 566–574 (2014). https://doi.org/10.1016/j.ijsolstr.2013.10.017.

57. S.-H. Kim *et al*., An Experimental and Numerical Study on Adhesion Force at the Nanoscale. *Nanoscale Adv.*, **6**(8), 2013–2025 (2024). https://doi.org/10.1039/D3NA01044A.

58. A. Y. Karatayev, D. Boltovskoy, L. E. Burlakova, D. K. Padilla, Parallels and Contrasts Between *Limnoperna Fortunei* and Species of *Dreissena*. In *Limnoperna Fortunei*, D. Boltovskoy, Ed. (Springer, Cham, 2015), pp. 261–297. https://doi.org/10.1007/978-3-319-13494-9_15.

59. S. Li, Z. Xia, Y. Chen, Y. Gao, A. Zhan, Byssus Structure and Protein Composition in the Highly Invasive Fouling Mussel *Limnoperna Fortunei*. *Front. Physiol.*, **9**, 418 (2018). https://doi.org/10.3389/fphys.2018.00418.

60. N. Farsad, T. W. Gilbert, E. D. Sone, Adhesive Structure of the Freshwater Zebra Mussel, *Dreissena polymorpha*. *Mater. Res. Soc. Symp. Proc.*, *1187-KK02-03* (2009).

61. S. Kim *et al*., Cation–π Interaction in DOPA-Deficient Mussel Adhesive Protein Mfp-1. *J. Mater. Chem. B*, **3**(5), 738–743 (2015). https://doi.org/10.1039/C4TB01646G.

62. M. A. Gebbie *et al*., Tuning Underwater Adhesion with Cation–π Interactions. *Nature Chem*, **9**(5), 473–479 (2017). https://doi.org/10.1038/nchem.2720.

63. L. Fischer *et al*., Sequence-Defined Positioning of Amine and Amide Residues to Control Catechol Driven Wet Adhesion. *Chem. Sci.*, **11**(36), 9919–9924 (2020). https://doi.org/10.1039/D0SC03457F.

64. Y. Li *et al*., Molecular Design Principles of Lysine-DOPA Wet Adhesion. *Nat Commun*, **11**(1), 3895 (2020). https://doi.org/10.1038/s41467-020-17597-4.

65. Y. Mu, P. Mu, X. Wu, X. Wan, The Two Facets of the Synergic Effect of Amine Cation and Catechol on the Adhesion of Catechol in Underwater Conditions. *Applied Surface Science*, **530**, 146973 (2020). https://doi.org/10.1016/j.apsusc.2020.146973.

66. M. Shin, J. Y. Shin, K. Kim, B. Yang, J. W. Han, N.-K. Kim, H. J. Cha, The Position of Lysine Controls the Catechol-Mediated Surface Adhesion and Cohesion in Underwater Mussel Adhesion. *Journal of Colloid and Interface Science*, **563**, 168–176 (2020). https://doi.org/10.1016/j.jcis.2019.12.082.

67. B. D. B. Tiu, P. Delparastan, M. R. Ney, M. Gerst, P. B. Messersmith, Cooperativity of Catechols and Amines in High-Performance Dry/Wet Adhesives. *Angew Chem Int Ed*, **59**(38), 16616–16624 (2020). https://doi.org/10.1002/anie.202005946.

68. X. Zhu, C. Wei, H. Chen, C. Zhang, H. Peng, D. Wang, J. Yuan, J. H. Waite, Q. Zhao, A Cation-Methylene-Phenyl Sequence Encodes Programmable Poly(Ionic Liquid) Coacervation and Robust Underwater Adhesion. *Adv Funct Materials*, **32**(2), 2105464 (2022). https://doi.org/10.1002/adfm.202105464.

69. W. R. Wonderly, T. R. Cristiani, K. C. Cunha, G. D. Degen, J.-E. Shea, J. H. Waite, Dueling Backbones: Comparing Peptoid and Peptide Analogues of a Mussel Adhesive Protein. *Macromolecules*, **53**(16), 6767–6779 (2020). https://doi.org/10.1021/acs.macromol.9b02715.

70. M. A. McCartney *et al*., The Genome of the Zebra Mussel, *Dreissena Polymorpha*: A Resource for Comparative Genomics, Invasion Genetics, and Biocontrol. *G3 Genes|Genomes|Genetics*, **12**(2), jkab423 (2022). https://doi.org/10.1093/g3journal/jkab423.

71. W. Wei, J. Yu, C. Broomell, J. N. Israelachvili, J. H. Waite, Hydrophobic Enhancement of Dopa-Mediated Adhesion in a Mussel Foot Protein. *J. Am. Chem. Soc.*, **135**(1), 377–383 (2013). https://doi.org/10.1021/ja309590f.

72. B. Gabryelczyk *et al*., Hydrogen Bond Guidance and Aromatic Stacking Drive Liquid-Liquid Phase Separation of Intrinsically Disordered Histidine-Rich Peptides. *Nat Commun*, **10**(1), 5465 (2019). https://doi.org/10.1038/s41467-019-13469-8.

73. Y. Sun, Z. W. Lim, Q. Guo, J. Yu, A. Miserez, Liquid–Liquid Phase Separation of Proteins and Peptides Derived from Biological Materials: Discovery, Protein Engineering, and Emerging Applications. *MRS Bull*., **45**(12), 1039–1047 (2020). https://doi.org/10.1557/mrs.2020.301.

74. M. Renner-Rao *et al*., Mussels Fabricate Porous Glues via Multiphase Liquid–Liquid Phase Separation of Multiprotein Condensates. *ACS Nano*, **16**(12), 20877–20890 (2022). https://doi.org/10.1021/acsnano.2c08410.

75. E. Hennebert, B. Leroy, R. Wattiez, P. Ladurner, An Integrated Transcriptomic and Proteomic Analysis of Sea Star Epidermal Secretions Identifies Proteins Involved in

Defense and Adhesion. *Journal of Proteomics*, **128**, 83–91 (2015).
https://doi.org/10.1016/j.jprot.2015.07.002.
76. J. N. Israelachvili, P. M. McGuiggan, Adhesion and short-range forces between
surfaces. Part I: New apparatus for surface forces measurements. *J. Mater. Res.*,
5(10), 2223-2231 (1990).
77. E. W. Danner, Y. Kan, M. U. Hammer, J. N. Israelachvili, J. H. Waite, Adhesion of
Mussel Foot Protein Mefp-5 to Mica: An Underwater Superglue. *Biochemistry*,
51(33), 6511–6518 (2012). https://doi.org/10.1021/bi3002538.
78. G. D. Degen, K. C. Cunha, Z. A. Levine, J. H. Waite, J.-E. Shea, Molecular Context
of Dopa Influences Adhesion of Mussel-Inspired Peptides. *J. Phys. Chem. B*,
125(35), 9999–10008 (2021). https://doi.org/10.1021/acs.jpcb.1c05218.
79. G. D. Degen, P. Delparastan, B. D. B. Tiu, P. B. Messersmith, Surface Force
Measurements of Mussel-Inspired Pressure-Sensitive Adhesives. *ACS Appl. Mater.*
Interfaces, **14**(4), 6212–6220 (2022). https://doi.org/10.1021/acsami.1c22295.
80. G. Violano, G. P. Demelio, L. Afferrante, On the DMT Adhesion Theory: From the
First Studies to the Modern Applications in Rough Contacts. *Procedia Structural*
Integrity, **12**, 58–70 (2018). https://doi.org/10.1016/j.prostr.2018.11.106.
81. N. Jia, Y. Yao, Z. Peng, S. Chen, Surface Effect in Nanoscale Adhesive Contact. *The*
Journal of Adhesion, **97**(4), 380–398 (2021).
https://doi.org/10.1080/00218464.2019.1660649.
82. M. Zakeri, J. Faraji, Modeling the Adhesion of Spherical Particles on Rough Surfaces
at Nanoscale. *International Journal of Adhesion and Adhesives*, **124**, 103385 (2023).
https://doi.org/10.1016/j.ijadhadh.2023.103385.
83. F. L. Leite, P. S. P. Herrmann, Application of Atomic Force Spectroscopy (AFS) to
Studies of Adhesion Phenomena: A Review. *Journal of Adhesion Science and*
Technology, **19**(3–5), 365–405 (2005). https://doi.org/10.1163/1568561054352667.
84. S. Kim, W. Moon, J. Jeon, A Reliable Method to Measure the Adhesive Force with a
Tiny Amount of Adhesive Material. *The Journal of Adhesion*, **84**(1), 60–77 (2008).
https://doi.org/10.1080/00218460801888417.
85. K. M. Kimmins, B. D. James, M. T. Nguyen, D. Asker, B. D. Hatton, E. D. Sone, Oil-
infused Silicone Prevents Zebra Mussel Adhesion. *ACS Appl. Bio. Mater.*, **2**, 5841–
5847 (2019). https://doi.org/10.1021/acsabm.9b00832
86. A. L. Oberg *et al.*, Statistical Analysis of Relative Labeled Mass Spectrometry Data
from Complex Samples Using ANOVA. *J. Proteome Res.*, **7**(1), 225–233 (2008).
https://doi.org/10.1021/pr700734f.
87. M. Mirdita *et al.*, ColabFold: Making Protein Folding Accessible to All. *Nat Methods*,
19(6), 679–682 (2022). https://doi.org/10.1038/s41592-022-01488-1.
88. M. A. Paz *et al.*, Specific Detection of Quinoproteins by Redox-Cycling Staining.
Journal of Biological Chemistry, 266(2), 689–692 (1991).
https://doi.org/10.1016/S0021-9258(17)35225-0.

Figures and Tables

Figure 1: Overview of plaque proteomics investigation. (A) Schematic of sample collection. (B) Summary
of workflow. (B) Summary of workflow. (C) Number of unique proteins found in each group of extractions. Insets indicate the
proportion of Dbfp's in each group. (D) Distribution of proteins in the intersecting 'footprint + bulk plaque'
group and the 'ALL' group, highlighting the proportion of proteins that are Dbfp's or show strong homology
(BLAST score $\leq$ e^{-30}), low homology (BLAST score $\leq$ e^{-6}), or no homology to known proteins. †(8)

Figure 2: Volcano plot of quagga mussel plaque proteins. Each point represents an identified protein.
Proteins with a $\log_2$(fold change ratio between footprint and bulk plaque) > 1 (i.e. > 2 FCR) and a
$-\log_{10}$(p-value) > 1.3 (i.e. p-value < 0.05) are significantly enriched in the footprint. Dbfp7 variants and
Dbfp8 are indicated as the red and blue points, respectively.

Figure 3: The Dbfp7 protein family is rich in glycine, proline, asparagine, and tyrosine and lacks
significant amounts of DOPA. (A) Sequences of Dbfp7 variants. Variants are sorted by length,
demonstrating the 28-residue region that is present only in the C-terminal half of the acidic variants. The
consensus chart (generated by Jalview 2.11.3.3) below the alignment indicates the residues that are
highly conserved, particularly in the C-terminal portion. (B) Dot blot with known amounts of peptidyl
DOPA, lysozyme negative control, pre-Dbfp7 HPLC fraction, and Dbfp7 stained with NBT. (C) Calibration
curve obtained with known amounts of peptidyl DOPA; the blue point indicates the absorbance of Dbfp7,
corresponding to 0.11 ± 0.02 mol% DOPA. Error bars indicate standard deviation (n = 3).

Figure 4: Dbfp7 surface morphology and structure. (A) CD spectrum of Dbfp7 dissolved in 5% acetic acid. (B) Proportion of secondary structure features via deconvolution of CD spectrum by Contin-LL. (C) Predicted structure of Dbfp7 (variant 1) visualized by ChimeraX. Labelled residues indicate the 28-residue C-terminal region found only in acidic variants. (D) Atomic force microscopy height images of Dbfp7 on mica in aqueous conditions. Scalebars each represent 500 nm.

Figure 5: Dbfp7 is adhesive in an aqueous environment. (A) Adhesion map of Dbfp7 (100 µg/mL) coated
on mica, imaged in aqueous conditions. Scale bar is 1 µm. (B) The mean height of each bin in the
adhesion histogram demonstrates that the areas of low adhesion correspond to regions with taller
features. Error bars indicate the standard error of the mean (s.e.m.). (C) Dbfp7 is overall more adhesive
than samples prepared with BSA (n = 262,144). (D) Adhesion energy of Dbfp7 and BSA calculated by the
DMT and JKR contact models from adhesion force measurements obtained by PFQNM AFM (s.d., N =
2,859,269 and 793,084 for Dbfp7 and BSA, respectively). *** indicates p-value < 0.001.

Table 1: The quagga mussel footprint proteome in order of relative enrichment in the footprint. Accession
#: transcript label and reading frame (0F, 1F, 2F, 0R, 1R, or 2R). $\log_2$(FCR FP:BP): log-transformed fold
change ratio between footprint and bulk plaque. $-\log_{10}$(p-value): log-transformed significance value. SP:
signal peptide. DOPA: confident (Y), possible (M), or no (N) detection of at least one hydroxylated
tyrosine (peptidyl 3,4-dihydroxyphenylalanine) residue via LC-MS/MS sequencing. pI: isoelectric point
calculated from the translated transcript sequence. MW: molecular weight calculated from the translated
transcript sequence.

Identification	Accession #	$\log_2$(FCR FP:BP)	$-\log_{10}$(p-value)	SP? (Y/N)	DOPA? (Y/M/N)	pI	MW (kDa)
Kielin/Chordin-like	comp50443_c0_seq4-0F	3.14	2.10	Y	M	7.99	15.2
Mucin/Spondin-like	comp52859_c0_seq1-1R	2.67	2.70	N	N	4.59	113
Hornerin	comp41116_c0_seq1-2F	2.66	3.00	Y	N	8.06	8.26
Zonadhesin/IgGFc-binding	comp53519_c2_seq1-2R	2.56	20.0	N	M	5.31	125
Zonadhesin/IgGFc-binding	comp53519_c2_seq2-0R	2.53	20.0	N	N	5.25	128
Hornerin	comp36226_c0_seq1-1F	2.33	3.00	Y	N	8.48	16.3
CD109 antigen-like	comp30100_c0_seq2-2F	2.20	3.00	Y	N	4.88	28.2
TPF1III	comp56420_c11_seq1-2F	2.07	2.10	N	N	8.01	8.67
Kielin/Chordin-like	comp31275_c0_seq2-2R	1.99	9.94	N	M	8.16	5.51
TTN1	comp53519_c2_seq3-0R	1.98	2.10	N	N	4.76	35.7
Peroxidase	comp51774_c0_seq1-2F	1.94	4.20	N	N	8.37	83.0
Kielin/Chordin-like	comp31275_c0_seq1-2R	1.82	6.70	Y	M	5.18	13.9
Kielin/Chordin-like	comp30932_c0_seq1-0F	1.81	1.55	N	N	7.53	9.08
Dbfp7	comp52765_c1_seq12-1F	1.79	1.66	Y	Y	9.22	12.0
Fibrillin/Galaxin	comp56544_c0_seq1-2R	1.77	20.0	N	M	5.95	149
Fibrillin/Galaxin	comp56544_c3_seq1-2R	1.71	20.0	N	M	6.91	231
Unknown	comp57807_c0_seq1-0R	1.68	1.55	N	N	8.82	12.4
Galaxin	comp56544_c8_seq1-1R	1.67	3.00	N	N	5.84	29.7
Dbfp7	comp52765_c1_seq11-1F	1.42	2.52	Y	Y	8.83	9.44
Dbfp7	comp52765_c1_seq14-1F	1.14	1.31	Y	Y	8.80	9.90
Dbfp8	comp44862_c1_seq1-2F	1.03	3.45	Y	M	9.99	16.4