Transcriptomic basis of within- and trans-generational predator-
induced plasticity in the freshwater snail *Physa acuta*

Authors:

Léo Dejeux[1], Nathanaëlle Saclier[2], Juliette Tariel-Adam[3], Maxime Hoareau[1],
Tristan Lefébure[1], Lara Konecny[1], Sandrine Plénet[1], Emilien Luquet[1, 4]

[1] Université Claude Bernard Lyon 1, LEHNA UMR 5023, CNRS, ENTPE, F-
69622, Villeurbanne, France

[2] ISEM, Univ Montpellier, CNRS, IRD, Montpellier, France

[3] Fish Lab, Department of Biological Sciences, Macquarie University,
Sydney, NSW, Australia

[4] Institut universitaire de France (IUF)

Abstract

Inducible defences in response to predation risk are a well-known example of adaptive phenotypic plasticity. Although inducible defences have been studied mainly within a generation (within-generational plasticity), there is now clear evidence that ancestral exposure to predation risk can influence the defences expressed by offspring, even if they have not been exposed themselves (transgenerational plasticity). The molecular mechanisms allowing the transmission of environmental information across generations are not well understood. In this study, we combined measures of antipredator responses (behavioural and morphological) with transcriptomic investigations across two generations in the freshwater snail *Physa acuta*. We hypothesised that both within- and transgenerational plasticity would induce phenotypic changes associated with differential gene expression. Our results confirmed within- and transgenerational plasticity: F1 snails respond to predator-cue exposure by increasing escape behaviour, reducing shell length, and developing thicker and slenderer shells, whereas F2 snails from exposed parents have longer and thicker shells with narrower apertures. Within- and transgenerational plasticity were accompanied by the differential expression of 112 genes (101 up- and 11 downregulated) and 23 differentially expressed genes (17 up- and 6 downregulated), respectively. Within- and transgenerational plasticity did not share common differentially expressed genes, but the associated molecular functions, involving metabolism and transcription regulation, were similar. These results suggest that predator-induced within-

generational plasticity and transgenerational plasticity may result from

different genomic pathways and may evolve independently.

INTRODUCTION

Phenotypic plasticity is the ability of a genotype to produce alternative

phenotypes under different environmental conditions (Pigliucci, 2005).

Plastic responses are widespread across the phylogenetic spectrum (*e.g.*

Galloway & Etterson, 2007; Luquet et al., 2011; Segers & Taborsky, 2011;

Walsh et al., 2016) and are a major concern in evolutionary biology as they

can be adaptive, i.e., increasing the fitness of organisms facing

environmental changes (Pigliucci, 2005). Phenotypic plasticity classically

refers to within-generational plasticity (WGP), meaning that it occurs within

one generation. Recently, plasticity has also been found to occur across

generations (transgenerational plasticity [TGP]), where the phenotype of an

organism is influenced by the environment experienced by previous

generations (Bell & Hellmann, 2019). Interest in TGP has grown rapidly in

recent years with evidence that TGP is at least as widespread as WGP

(Jablonka & Raz, 2009; Tariel, Plénet, et al., 2020b). Although there are

numerous examples of TGP generating seemingly adaptive offspring

phenotypes, its adaptive role is expected to be more challenging than that

of WGP because of patterns of environmental variation across generations

(Colicchio & Herman, 2020). Indeed, the selection of a strong temporal

autocorrelation between environments of distinct generations (Ezard et al.,

2014; Kuijper & Hoyle, 2015; Prizak et al., 2014) and a transmission system

that allows the induction of adaptive phenotypes across generations is

required (Fallet et al., 2020). The inheritance mechanisms appear diverse,

ranging from epigenetic changes to parental transmission of cytoplasmic

elements (small RNAs, nutrients, hormones, proteins), parental care and

cultural transmission (Bell & Hellmann, 2019; Danchin et al., 2011).

Regardless of the transmission mechanism, the influence of the parental

environment on the offspring phenotype should be linked to changes in

gene expression, at least in some specific tissues and developmental

stages. Surprisingly, the transgenerational plastic response at the molecular

level has been overlooked. This is indeed a necessary first step to unravel

the molecular mechanisms of plasticity in general and their

transgenerational inheritance. The most compelling explorations of

changes in gene expression profiles in response to both developmental

(WGP) and parental (TGP) environments have been conducted by Hales et

al. (2017) on *Daphnia ambigua* and (Stein et al. (2018) on the three-spined

stickleback (*Gasterosteus aculeatus*). These studies compared gene

expression between individuals that had been exposed to predator cues

relative to nonexposed individuals within a generation (WGP genes) and

among their offspring (from non-exposed and exposed parents; TGP

genes). Stein et al. (2018) reported that the expression of the same group

of genes changed their expression in offspring regardless of wether

predation risk was experienced by the offspring (TGP), the father (WGP), or

both. This suggests that a core set of genes is consistently affected by

predation risk regardless of who experiences it. Conversely, Hales et al.

(2017) reported that changes in the expression of genes that respond to

TGP differed depending on whether exposure to predation was

experienced by parents or grandparents. They also revealed that very few

genes were common to both WGP and TGP. Notably, Clark et al. (2019)

also reported that the TGP gene expression profiles of larvae of sea urchins

(*Psammechinus miliaris*) spawned at low pH from preacclimated adults

differed from those of larvae produced by adults exposed to ambient pH,

but the authors did not explore the WGP genes. These few results highlight

the necessity of investigating WGP and TGP together at the molecular

level. In addition to the inheritance mechanisms allowing organisms to

retain information about past environmental conditions to adjust their future

phenotype, we need to know whether the two forms of plasticity are distinct

at the molecular level and potentially free to evolve independently of one

another (Bell & Stein, 2017).

Predation risk is a major selection pressure for most species and results in

the evolution of defences in prey (Kikuchi et al., 2023). As predation risk

often varies over time and space, it favours the evolution of phenotypic

plasticity in defences (Reger et al., 2018; Tariel, Plénet, et al., 2020b; Viney

& Diaz, 2012). Indeed, although some defences against predators are

constitutive, plastic defences are only induced when prey detect some

predator cues in their environment (*e.g.* visual cues, chemical cues such as

predator odours, or alarm signals released by conspecifics). These induced

defences are classic examples of adaptive WGP: prey can increase their

fitness by producing defences only when predators are present. The links

between prey-predator interactions and WGP have long been studied (*e.g.*

Tariel et al., 2020b; Tollrian, 1995). Plastic defences can also be induced in

offspring after their parents or more distant ancestors have detected

predator cues. This transgenerational induction of defences was first

highlighted in Daphnia morphology by Agrawal et al. (1999) and has since

been repeatedly shown in multiple species and defences (review in

MacLeod et al., 2022; Tariel, Plénet, et al., 2020b).

Consequently, this work aims to elucidate the molecular mechanisms

underlying the expression of WGP and TGP in defences in the context of

predator-prey interactions. We used *Physa acuta* (also referred to as

Physella acuta), a freshwater snail known for its within- and

transgenerational antipredator defences, such as escape behaviour

e.g.,snails crawl out of the water when they or their parents detect predator

cues (Luquet & Tariel, 2016; Tariel, Luquet, et al., 2020; Tariel, Plénet, et

al., 2020a; Turner et al., 1999), changes in shell size and shape, increases

in shell thickness and overall increases in shell-crush resistance (*e.g.*, shell

are harder to handle or to crush by predators (Auld & Relyea, 2011; Beaty

et al., 2016; DeWitt, 1998; Tariel, Plénet, et al., 2020c; Tariel-Adam et al.,

2023). We combined a two-generation experiment to generate predator-

induced WGP and then TGP of defences (only in a nonexposed

environment) with a transcriptomic investigation. We used transcriptome

sequencing and assembly to investigate transcriptome mantle skirt gene

expression specifically (the mantle skirt is the tissue that synthesises the

shell). We expected that gene expression would be affected by current

(WGP) and parental (TGP) exposure to predator cues. More specifically, as

WGP and TGP induce similar phenotypic responses (*e.g.* increases in shell

thickness), we expected the two processes to share the same core set of

genes.

MATERIALS & METHODS

Animal collection, rearing conditions and experimental design

Adult *P. acuta* snails were collected from a wild population in the lentic backwater of the Rhône River (Lyon, France, 45° 48'6"N, 4° 55'33"E) in September 2018. Individuals from this population have been repeatedly used in previous studies on predator-induced plasticity (Luquet & Tariel, 2016) and are naturally exposed to predation by the crayfish *Faxonius limosus*. The experiment was conducted in our laboratory in a temperature-controlled chamber at 25 °C with a photoperiod of 12/12 h. The wild snails constituting the F0 generation were pooled in a 10 L plastic box filled with dechlorinated tap water (control water hereafter) and interbred overnight. They were then isolated in 80 mL plastic boxes (4.5 × 6 cm) filled with tap control (i.e., without predator cues) water to lay eggs for 3 days. Eggs within boxes constitute F1 matrilines. Eggs hatched seven days later, and siblings subsequently grew together for an additional ten days in water without predator cues. Then, 10 siblings of 20 F1 matrilines were randomly selected and split into two developmental environments: five siblings in water without predator cues (nonexposure treatment) and five siblings in water with predator cues (exposure treatment). F1 siblings individually developed for the next 31 days in 80 mL plastic boxes. On this date, one F1 snail per matriline and treatment was frozen at –80 °C for molecular analysis and all remaining F1 snails were measured (see below).

To generate the F2 generation, we pooled F1 snails from the same

treatment in an aquarium filled with control water and allowed them to

copulate for 24 h. As previously described, F1 snails were then isolated in

80 mL boxes and laid eggs for 3 days to produce F2 matrilines. F2 snails

from the two parental environments (nonexposed and exposed to predator

cues) were reared only in control water to investigate the effect of the

parental environment (TGP) alone, i.e., without the interaction between

parental and developmental environments. Our previous study revealed

that TGP is stronger in a mismatch situation (i.e., when F2 from exposed

parents are not exposed (Luquet & Tariel, 2016)). We followed the same

protocol steps as previously described for F1 snails for F2 generation,

except that eight siblings per matriline were reared and measurements

were made at an older age because of slower growth. For F1, one F2 snail

per matriline and treatment was frozen for molecular analysis, and the

others were measured.

Snails were fed *ad libitum* with boiled and mixed lettuce. The water

(according to the treatment) and food were renewed twice a week. The

water with predator cues (i.e., exposed treatment) was the water in which *F.*

limosus crayfish were raised (one crayfish/5 L), for 3 days and were

subsequently fed with snails. Additionally, to ensure the presence of fresh

alarm cues, we smashed several *P. acuta* adult snails in predator-cue water

(one snail/5 L) for 1 h before use.

Behavioural and morphological responses

Behavioural and morphological responses

In this study, to check that exposure to predator cues triggered WGP and

TGP, escape behaviour and shell morphology were measured in F1

(nonexposed snails = 76, exposed snails = 73) and F2 (snails from

nonexposed parents = 89, snails from exposed parents = 101).Some

mortality events led to uneven sample sizes, but the mortality rates

remained consistent across families (see Table S1). Crawling out of the

water (positioning themselves above the water surface) allows snails to

escape benthic predators such as crayfish (DeWitt, 1998; Tariel, Plénet, et

al., 2020a). Shorter and narrower shell and aperture dimensions and

thicker shells are adaptive antipredator responses (Auld & Relyea, 2011).

These phenotypic traits have consistently been implicated in the

transgenerational responses of *P. acuta* to predator cues (Beaty et al.,

2016; Luquet & Tariel, 2016).

Escape behaviour: One week before the measurements and four hours

after the water renewal, we recorded the position above or below the water

surface of each snail in their rearing boxes. Consequently, the behaviour of

F1 snails was recorded in water without or with predator cues according to

their respective treatments, whereas that of F2 snails was recorded only in

water without predator cues.

Shell thickness, shell length and shape: Shell thickness was directly

measured with an electronic calliper at the nearest 0.01 mm at the edge of

the aperture. A photograph of each snail (aperture upwards) was

subsequently taken with an Olympus SC50 camera installed on an

Olympus SZX9 binocular and an Olympus DF PLAPO 1X-2 objective at a

8× magnification. Snail shell length, shell width, aperture length and

aperture width were measured using ImageJ (Schneider et al., 2012). To

estimate potential changes in shell shape according to treatment, we used

geometric methods (Gustafson et al., 2014). For each snail shell, we

digitised 19 landmarks (Fig. 1) using ImageJ software. Snails that lacked

homologous landmarks because of broken shells (typically broken

apertures and apexes) or small sizes (lack of the third shell spire) were

removed from the analyses (nonexposed F1 snails = 76, exposed F1 snails

= 72, F2 snails from nonexposed parents = 89, F2 snails from exposed

parents = 99). To compare shell shapes according to treatment, we scaled

the shapes using the size metric defined below (Eqn. 1). The rotational

components were removed by translating all shapes so that the centroids

lie at the origin. Rotational components were removed by rotating all the

shapes until the Procrustes distance was minimised, with the first shell

fixed as a reference.

$$s_k = \sqrt{\sum_{i=1}^{19} ((x_{k,i} - \bar{x}_k)^2 + (y_{k,i} - {}_k{}^2)}$$

Equation 1: Procustes equation used to scale the shapes using the size metric , which is

the size of the -th shell; and are the x and y components of the -th landmark;

( ) is the centroid of the -th shell.

Figure 1: Locations of the different landmarks (circle symbols) and pseudo-landmarks (positioned using

a geometric method, square symbols) on *P. acuta* shell. Pseudo-landmarks were created using a

geometric method: first, two tangents (T1 and T2) were drawn from landmark 1 (apex). T1 touches the

curve of the habitation spire (5) and T2 touches the external edge of the aperture (10). From the two

points of tangency (points 5 and 10), two lines perpendicular to the tangents were traced (P1 and P2).

The intersections with the columellar edge of the aperture, the external edge of the aperture and the

curve of the habitation spire give additional pseudo-landmarks (6, 9, 18 and 19). The length of the shell

L (straight line passing by 1 and 8) gives pseudo-landmark 18 at the intersection with the columellar

edge of the aperture).

Statistical analyses

The same statistical analyses were separately performed to test for WGP on F1 snails and TGP on F2 snails. The treatment factor (i.e., nonexposed or exposed to predator cues) was the developmental environment for WGP (F1), whereas it was the parental environment for TGP (F2).

Escape behaviour: The effects of predator cues on escape behaviour (i.e. snail position above/on or below the water surface) were analysed using generalised linear mixed models (GLMMs) assuming a binomial distribution (logit link function). The treatment was a fixed effect, and matriline was a random intercept effect. We tested the significance of fixed and random effects with likelihood ratio tests.

Shell thickness and shell length: To analyse the effects of predator cues on shell length and shell thickness, we performed linear mixed models with treatment as a fixed effect and matriline as a random intercept. We used restricted maximum likelihood estimation and Kenward and Roger's approximation for degrees of freedom. We tested the significance of fixed effects with type II F-tests (Kuznetsova et al., 2017) and the significance of random effects with likelihood ratio tests.

Linear mixed models and generalised linear mixed models were generated using the lme4 package (Bates et al., 2014) in R software (version R-4.4.1 for Windows, (RStudio Team, 2020).

Shell shape: The effects of predator cues on shell shape were analysed

using principal component analysis (PCA) and visualised with the 2D

landmark relative warp analysis functionality. All the standardised shapes

obtained via the Procrustes method were analysed using PAST software

with a Student test (Hammer et al., 2001).

RNA Extraction and Sequencing The mantle skirt was dissected for 5

nonexposed F1 snails and 5 exposed F1 snails, and for 10 F2 snails from

nonexposed parents and 10 F2 snails from exposed parents. Total RNA

from the mantle was extracted by adding TRI Reagent (Molecular Research

Center MRC, TR118) to the samples and homogenising the tissues on ice

with pistons and an agitator (Motor Mixer Clearline ref.045011 Dutscher). All

remaining steps were carried out according to the manufacturer's protocols

(Molecular Research Center MRC, TR118). Then, the RNA was treated with

Turbo DNase enzyme (Turbo DNA free kit, Invitrogen, AM1907) and

assayed by fluorescence using a Qubit fluorometer (Qubit® RNA HS Assay

Kits Molecular Probes, Invitrogen, Q32855). Library construction was

carried out using an NEBNext Ultra II RNA Library Prep Kit for Illumina

(Biolabs New England, NEB E7770, E7490, E7600). Libraries were

randomly split into three groups and sequenced on three lanes with an

Illumina® HiSeq4000™ machine on the GenomEast platform hosted at the

IGBMC (GenomEast Platform – Institut de Génétique et de la Biologie

Moléculaire et Cellulaire IGBMC, Illkirch, France), resulting in 50 bp long

single-end reads.

Differential gene expression analysis

FASTQC (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/)) was used to assess the quality of the reads. We used a reference genome ORF *P. acuta* available on NCBI (Isolate: Inbredline_101_S28. RefSeq: GCF_028476545.12). Gene expression levels were assessed on annotated transcripts using Kallisto (-l 50 -s 10 options; Bray et al., 2016). We aggregated effective counts at the gene level and then applied rounding to the nearest whole number to convert noninteger values into integer counts. We also applied a minimum count filter of 1 (cpm) where we removed all the genes with reads below 1 (i.e. removing 5634 genes from the total of 46,965). The rounded effective counts were used as inputs for differential gene expression using the DESeq2 package (Love et al., 2014), which performs differential gene expression analysis using negative binomial distribution and shrinkage estimation for dispersions ("apeglm" type shrinkage, Zhu et al. 2019), and outputs expression fold changes to improve the interpretability of the estimates. We specified nonexposed or exposed treatment in the model (design formula: "design = ~treatment") within each generation independently, meaning that we compared F1 snails from nonexposed and exposed treatments, and F2 snails from nonexposed and exposed parents. First, the Wald test (DESeq2 package) was used to test the significance of differentially expressed genes (DEGs) in all samples between the nonexposure and exposure groups. Second, we checked for any structures in the normalised count table by performing a principal

component analysis (PCA) of the 500 most DEGs from DESeq2 analysis

(pval < 0.05) based on Log2FoldChange (|LFC|, i.e., the logarithm base 2

of the fold change in gene expression between two experimental

conditions, providing a measure of both the magnitude and direction of

differential expression). Third, we filtered the significant DEGs by an |LFC|

greater than 1 in each generation. Consequently, only differential

expression levels with adjusted p-values < 0.05 and |LFC| > 1 were

considered thereafter. We subsequently investigated whether the DEGs in

F1 were also differentially expressed in F2 by comparing specific gene IDs

across the datasets from the two generations. Finally, as the quality of

annotations of the reference genome did not allow complex functional

analyses, we determined the potential gene functions and the associated

protein families of all significant DEGs using the scan procedure of

InterPro, a tool provided by the European Bioinformatics Institute from the

European Molecular Biology Laboratory (EMBL-EBI; (Blum et al., 2024;

Jones et al., 2014). InterPro integrates signatures from 13 member

databases to classify proteins into families, to predict functional domains,

and to identify key sites, enabling comprehensive functional annotation.

RESULTS

<u>Predator-induced behaviour and shell morphology</u>

Within-generational Plasticity

In the F1 generation, the proportion of crawling-out behaviour increased

significantly in snails exposed to predator cues (+85.25%, $F_{[1,19.43]} =$

321.82, $p < 0.001$; Fig. 2A). Shell length significantly decreased in snails

exposed to predator cues (-6.3%, $F_{[1,128.71]} = 15.46$, $p < 0.001$; Fig. 2B).

Shell thickness significantly increased in exposed snails (+18.8%,

$F_{[1,129.58]} = 4.02$, $p = 0.047$; Fig. 2C). The matriline random effect was

significant for shell length ($\chi^2[1] = 19.022$, $p < 0.001$) and nonsignificant for

behaviour and shell thickness ($\chi^2[1] = 0.690$, $p = 0.406$ and $\chi^2[1] = 2.6116$,

$p = 0.106$, respectively).

342
Figure 2: Effects of developmental exposure to predator cues (WGP) on crawling-out behaviour, shell

length and shell thickness in the F1 generation. A: Proportion of crawling-out behaviour; B: mean shell

length (mm) ; C: mean shell thickness (mm). Error bars represent SE. Nonexposed and exposed to

predator-cue treatments are symbolised with blue circles and red triangles, respectively.

Exposure to predator cues induced changes in shell shape. The first two

axes of the PCA accounted for 59.2% of the shell shape variation (Fig. 3),

and PCA axes 1–4 (74.2% cumulative shape variation) were the only axes

accounted for more than 5% of the shape variation. PC1 showed a change

in the columellar area (close to landmarks 18–19) from shells with relatively

wide apertures (dilated columellar area; Fig. 3C) to shells with relatively

narrow apertures (compressed columellar area; Fig. 3D). PC2 changed in

the slenderness of the shell from shells with relatively short spires

(compressed apex; Fig. 3B) and relatively wide apertures (dilated aperture)

to shells with relatively short spires (dilated apex) and narrow aperture

widths (compressed aperture; Fig.3A).

Compared to exposed snails, nonexposed snails presented the same range

of deformation in terms of the columellar area (PC2, t-test: t = 1.07, df =

144, p = 0.28). However, the shells of snails exposed to predator cues were

slenderer, with longer spires and narrower aperture widths than those of

nonexposed snails (PC1, t-test: t = 4.803, df = 144, p < 0.001). This

elongation was associated with the dilatation of the last three spires and a

narrow aperture at the base of the external edge. In contrast, the shells of

nonexposed snails had a wider aperture at the base of the external edge

and smaller spires.

Figure 3: Shape variation of nonexposed and exposed snails from the F1 generation (blue circle and

red triangle symbols, respectively. Panels A, B, C, and D represent deformation grids with shell shapes

on the extreme ends of the PCA axes. The 95% confidence ellipses are represented for nonexposed

and exposed to predator-cue treatments. The dilation□compression gradient bar shows the deformation

of the shell landmarks relative to the points of an average shell (which would be located at the centre of

the PCA).

Transgenerational plasticity

In the F2 generation, the proportion of crawled snails did not differ between

snails from nonexposed and those from exposed parents ($F_{[1,20.31]}$ =

3.929, p = 0.061; Fig. 4A). Snails from parents exposed to predator cues

were significantly longer and had thicker shells than those from

nonexposed parents (+4.9%, $F_{[1,170.87]}$ = 7.55, p = 0.006 and +16.9%,

$F_{[1,174.11]}$ = 9.858, p = 0.002, respectively; Fig. 4B-C). The matriline

random effect was significant for shell length ($\chi^2[1]$ = 31.685, p < 0.001) and

nonsignificant for behaviour and shell thickness ($\chi^2[1]$ = 0.022, p = 0.882

and $\chi^2[1]$ = 1.882, p = 0.17, respectively).

Figure 4: Effects of parental exposure (TGP) on crawling-out behaviour, shell length and shell thickness

in the F2 generation. A: Proportion of crawling-out behaviour; B: mean shell length (mm) ; C: mean shell

thickness (mm). Error bars represent SE. Non-exposed and exposed to predator cues are symbolised

with blue circles and red triangles, respectively.

Parental exposure to predator cues induced changes in shell shape. The

first two axes of the PCA accounted for 63.8% of the shell shape variation,

and the cumulative percentage accounting for PCA axes 1–4 was 80.14%

(Fig. 5). PC1 displayed a change in shell aperture width; the shells with

high PC1 scores (Fig. 5C) showed a dilatation of the shell where the outer

lip meets the parietal wall and a small compression of the columellar area

(i.e., a wider aperture). Shells with low PC1 scores (Fig. 5D) presented a

narrower aperture, a consequence of a dilated columellar area but

compression of the parietal wall and basal lip in comparison to shells with

high PC1 scores. PC2 showed a change in shell aperture width but also in

shell length, ranging from shells (Fig. 5C) with a dilated outer lip and

compression of the apex (which tends to reduce spire length, i.e., overall

shell length) to shells (Fig. 5D) showing elongated spires from the dilated

apex, with a narrower aperture due to outer lip compression.

Snails from predator-exposed parents had a dilatation of the shell in the

columellar area accompanied by a small compression of the parietal wall

(PC1: $t = 2.067$, $p = 0.04$). Conversely, there was no difference in

deformation of the outer lip or dilatation or compression of the apex

between snails from exposed and nonexposed parental environments

(PC2: $t = 0.59$, $p = 0.55$).

Figure 5: Shape variation of F2 snails from nonexposed and exposed parents (blue circles and red triangles symbols, respectively). Panels A, B, C, and D represent deformation grids with shell shapes on the extreme ends of the PCA axes. The 95% confidence ellipses are represented for nonexposed and exposed to predator-cue treatments. The dilation☐compression gradient bar shows the deformation of the shell landmarks relative to the points of an average shell (which would be located at the centre of the PCA).

Differential gene expression induced by predator-cues

A principal component analysis (PCA) was performed to explore the clustering of read counts on the first 500 most DEGs. The first and second principal components (PCs 1 and 2) explained 18% and 11% of the variance respectively (Fig. 6, PCA loadings available in Fig. S1). The plot did not show clustering between the nonexposed and exposed treatments or between the F1 and F2 generations (Fig. 6).In the F1 generation, differential expression analysis revealed a total of 112 DEGs between

nonexposed and exposed snails (101 upregulated genes and 11

downregulated gene; Fig. S2). In the F2 generation, the differential

expression analysis revealed a total of 23 DEGs between snails from

nonexposed and exposed parents (17 upregulated and 6 downregulated

gene; Fig. S2). A comparison of DEGs in F1 and F2 revealed that they did

not overlap (Fig. 7; Fig. S3).

Figure 6: PCA plot of the 30 transcriptome samples of *P. acuta* snails showing the divergence in

differential expression among the first 500 most differentially expressed genes (selected by greatest

|LFC|). Nonexposed and exposed to predator cue individuals are represented in blue and red,

respectively. F1 individuals are represented by circles and plain ellipses, and F2 individuals are

represented by triangles and dotted ellipses.

The DEGs in F1 were involved mainly in metabolism, signal transduction, extracellular matrix organisation and protein and transcriptional regulation (Fig. 8, Tables S2 and S3 for further details). The DEGs in F2 were involved mainly in metabolism and transcriptional regulation, including DNA methylation regulation (Fig. 8, Tables S2 and S3 for further details). In both F1 and F2, some DEGs were associated with calcium signalling (Fig. 8, Tables S2 and S3 for further details), and the majority of DEGs were genes with unknown functions.

Figure 8: Proportions of classes of functions of the differentially expressed genes in F1 (A) and F2 (B)

based on the InterPro EMBL-EBI database (Jones et al., 2014; Blum et al., 2024).

DISCUSSION

This work aimed to elucidate the molecular mechanisms underlying the

expression of within- and trans-generational plasticity of antipredator

defences in *P. acuta*. We expected that gene expression would be affected

by developmental (WGP) and parental (TGP) exposure to predator cues

and that WGP and TGP would share the same core set of genes. Exposure

to predator cues resulted in changes in escape behaviour, shell growth,

shell thickness and shell shape. The parental exposure to predator cues

(TGP) also influenced the offspring phenotype, particularly morphological

traits as previously shown in this system (*e.g.*, Luquet & Tariel 2016).

Accordingly, DEGs were detected between nonexposed and exposed

snails in F1 (WGP) and between F2 snails according to parental exposure

to predator cues (TGP). However, the number of DEGs detected for TGP

was lower than that detected for WGP. Finally, our results revealed that

DEGs associated with WGP and TGP did not overlap.

Phenotypic changes and gene expression induced by within-

generational exposure to predator cues

As expected, snails exposed to predator cues crawled more out of the

water, had thicker shells, were smaller and developed slenderer shells with

longer spires and narrower aperture widths than nonexposed snails did.

These behavioural and morphological responses represent defences and

are supposed to be adaptive as they are consistent with escape behaviour

(Tariel, Luquet, et al., 2020) and an increase in shell crush resistance

(Tariel, Plénet, et al., 2020a, 2020c), increasing snail survival in the face of

crayfish predation (Auld & Relyea, 2011). The decreased shell size

observed in exposed snails may result from physiological stress caused by

exposure to predator cues, including costs to produce antipredator

defences (*e.g.*, crawling-out behaviour occurs at the expense of foraging;

DeWitt, 1998).

These predator-induced WGP changes were associated with a distinct

pattern of gene expression with 112 DEGs (101 upregulated and 11

downregulated genes). Overall, this pattern is consistent with several

studies that explored the changes in gene expression induced by the

environment (*e.g.*, Aubin-Horth & Renn, 2009; Herman & Sultan, 2011). In

the context of predator☐prey interactions, the main knowledge about the

transcriptomic responses of prey to predators comes from studies on

Daphnia species. For example, Hales et al. (2017) in *Daphnia ambigua* and

Rozenberg et al. (2015) in *Daphnia pulex* reported that antipredator

defences, mainly those related to morphological traits, were associated with

48 and 230 DEGs respectively. In contrast, no DEGs were identified in

Daphnia galeata, a species known to adjust life history traits (such as

reproduction) but not morphological traits in response to predator cues.

Similarly, Orsini et al. (2018) revealed no changes in gene expression

associated with short-term exposure to fish kairomones in *Daphnia magna.*

These contrasting results suggest that the patterns of gene expression may

depend on several factors such as the traits considered and the strength of

predator signals (*e.g.* cue concentration and duration of exposure).

The functional analysis of DEGs in snails exposed to predator cues

suggested that these genes are linked mainly to metabolic functions and

transcription regulation, both of which are closely associated with stress

responses. Notably, 26% of the DEGs had unknown functions. A

substantial portion of these genes are involved in generic metabolic

processes (e.g., complex I intermediate-associated protein 30, which plays

a role in respiratory metabolism; Walker et al., 1992). These functions are

essential for maintaining cellular stability and enabling adaptation to

fluctuating environmental conditions. However, some genes are implicated

in more stress-specific responses that may be directly triggered by predator

exposure (*e.g.*, involvement in the TOR protein complex, which is well

known to influence the response to abiotic stressors; Blackwell et al., 2019;

Katewa & Kapahi, 2011). Similarly, some DEGs were linked to calcium ion

metabolism (*e.g.*, by coding for the annexin and peptidase C2 protein

families; Khorchid & Ikura, 2002) and could be involved in shell synthesis.

Calcium is the main component of shell gastropods (Bukowski & Auld,

2014), and modulations of general calcium pathways could influence the

expression of antipredator defences, such as increases in shell thickness

and changes in shell shape, as observed in our study. Our findings align

with previous research investigating the functions of DEGs in predator cue

environments and showing that genes are associated with defensive

structures, metabolism and transcription regulation. Indeed, Hales et al.

(2017) and Rozenberg et al. (2015) reported that the gene functions of *D.*

ambigua and *D. pulex* mostly explained the observed defensive phenotypic

changes (*e.g.*, the expression of proteins involved in the formation of

protective structures and in cuticle strengthening), as well as the metabolic

pathways involved in resource allocation and stress responses. Similar

results have been reported in larvae of the damselfly *Enallagma*

cyathigerum and in the marine copepod *Calanus finmarchicus,* where

predation risk activated genes encoding for stress proteins, reduced

antioxidant defence, and altered lipid metabolism (Skottene et al., 2020;

Slos & Stoks, 2008). Although the relationship between the functions of

DEGs and phenotypic traits is highly speculative in nonmodel organisms,

our study may highlight some interesting molecular pathways underlying

predator-induced WGP in *P. acuta* that would be interesting to study in

further detail.

Phenotypic changes and gene expression induced by

transgenerational exposure to predator cues.

Our results confirmed predator-induced TGP. The parental exposure to

predator cues significantly influenced the offspring phenotype, particularly

the morphological traits. F2 snails from exposed parents had thicker and

longer shells with narrower apertures than snails from nonexposed parents

while they had similar antipredator behaviour. These transgenerational

effects are consistent with previous studies on this system which

demonstrated that parental environments shape the antipredator traits of

offspring (Beaty et al., 2016; Tariel, Plénet, et al., 2020c). A thicker shell

and a larger size increase the shell-crush resistance, and a narrower

aperture limits body snail extirpation by crayfish (Auld & Relyea, 2011;

DeWitt, 1998). The longer shell of F2 snails from exposed parents was

opposed to the shorter shell of exposed snails in F1. This contrasting

pattern aligns with our previous study (Tariel, Plénet, et al., 2020c),

suggesting that offspring of exposed parents, having received information

about predation risk at an early developmental stage, activate pathways

that enable simultaneous investment in shell length to reach refuge size

(Auld & Houser, 2015) and shell thickness for increased crush-resistance

(Beaty et al., 2016; Tariel-Adam et al., 2023). The antipredator behaviour

was similar between snails from nonexposed and exposed parents, which

is in line with some contrasting results reported in other studies (Tariel,

Plénet, et al., 2020a, 2020c). Overall, the expression of antipredator traits

by offspring from exposed parents, even if they are not themselves

exposed to predator cues, may allow them to anticipate future predation

risk and increase their survival (MacLeod et al., 2022; Tariel, Plénet, et al.,

2020b).

At the molecular level, predator-induced TGP was associated with 23 DEGs

(17 upregulated and 6 downregulated) according to the parental

environment. The low number of DEGs in the F2 generation contrasts with

studies on factors other than predation (*e.g.*, pH in Clark et al., 2019;

temperature in Ledón-Rettig, 2023) and with parental exposure to predator

cues (Hales et al., 2017; Mommer & Bell, 2014; Stein et al., 2018). For

example, Stein et al., (2018) found 322 DEGs in three-spined stickleback

G. aculeatus offspring when predation risk was experienced by the father.

In *D. ambigua*, individuals from predator-exposed parents and grand-

parents presented 233 and 170 DEGs, respectively, compared with

individuals from nonexposed parents (Hales et al., 2017). Although this

contrast may result from a less stringent method of DEG selection (no LFC

filtering, *e.g.* Stein et al. 2018), it also raises the question of whether our

experiment captured the entire transgenerational transcriptomic response.

First, gene expression has been investigated in only one tissue (i.e., the

mantle skirt that synthesises the shell) in this study, whereas in many

studies it has been studied in whole organs (*e.g.*, Stein et al. 2018) or

whole individuals (*e.g.*, Hales et al. 2017). This may drastically decrease

the number of DEGs detected, capturing only a specific part of the gene

expression changes caused by parental environments. However, this

tissue-specific approach is particularly relevant in systems such as ours as

it allows the specific study of gene expression in the tissue at the origin of

phenotypic traits (i.e. shell modifications in this study).

Second, this study investigated only the gene expression in offspring that

were not themselves exposed to predator cues, which did not allow us to

determine whether the gene expression associated with parental

environments is modulated by developmental cues. In *P. acuta*, there is

evidence that parental and developmental environments interact to shape

the offspring phenotype: the resulting phenotypic patterns showed much

stronger TGP in offspring that had not experienced predator cues

themselves (WGP masking TGP for most traits; (Luquet & Tariel, 2016)).

Although it would be relevant to provide an integrative view of gene

expression patterns in all environmental scenarios, our study of DEGs

associated with TGP focused on the environmental situation where TGP is

the strongest.

Third, the lower number of DEGs in F2 could also result from a time lag

between changes in gene expression and antipredator defence induction

(Gilbert, 2001). Indeed, the environmental conditions experienced in early

life can have important consequences for phenotypes later in life (Gilbert,

2001; Lee et al., 2013) as well as for offspring phenotypes (Burton &

Metcalfe, 2014; Tariel-Adam et al., 2023; Yin et al., 2019). For example,

maternal provisioning plays a crucial role in transmitting environmental

information to offspring by influencing cytoplasmic components, including

mRNAs and proteins, during early embryogenesis (Burton & Metcalfe,

2014). This process can impact gene regulation during the maternal-to-

zygotic transition, potentially shaping phenotypic responses before they are

detectable in adults. In this study, both differential gene expression and

antipredator defences in offspring were investigated at the adult stage, and

offspring may have integrated the environmental information from their

parents early in development engaging them in specific predation risk

developmental trajectories. Consequently, the antipredator defence

observed in offspring from exposed parents may result from gene

expression changes early in development that no longer exist at the adult

stage, which could also explain why we did not find a clear common core

set of genes between WGP and TGP. For example, Mommer & Bell (2014)

showed that maternal experience with predation risk influences genome-

wide embryonic expression in *G. aculeatus*, an early expression that may

have phenotypic consequences later (Sharda et al., 2021). This complex

interplay among the timing of environmental cue perception, gene

expression and phenotypic consequences deserves to be explored in more

detail, for example by tracking DEGs throughout development.

The functional analysis of DEGs between snails from nonexposed parents

and those from exposed parents suggested the same main functions as

WGP, i.e., metabolism and transcription regulation. Genes associated with

metabolism are mainly involved in the response to oxidative stress and

carbohydrate metabolism. One gene might be involved in the regulation of

the cytosolic calcium ion concentration, which might influence the shell

synthesis as previously mentioned for WGP. Most genes linked with

transcription regulation in F2 are involved in methyltransferase activity,

suggesting that DNA methylation, a well-known bearer of epigenetic

information, may be involved in predator-induced TGP (Fallet et al., 2020).

Epigenetic mechanisms, such as DNA methylation and histone

modifications, are known to contribute to the transgenerational inheritance

of phenotypic traits (Duncan et al., 2014; Fallet et al., 2020 for a review on

molluscs). Recent studies on the freshwater gastropod *Potamopyrgus*

antipodarum have demonstrated that differential DNA methylation patterns

are associated with heritable morphological differentiations among

genetically uniform populations (Smithson et al., 2020; Thorson et al., 2017,

2019). These results suggest that the transgenerational inheritance of

antipredator traits in *P. acuta* may be mediated by bearers of epigenetic

information and call for more investigations of epigenetic inheritance in the

context of predator-induced plasticity.

Comparison of differentially expressed genes associated with WGP

and TGP

The number of DEGs associated with TGP was lower than that associated

with WGP. As mentioned above, this may have occured because we only

investigated DEGs at the adult stage or because F2 snails did not

experience direct exposure to predator cues. Surprisingly, the DEGs

associated with WGP and TGP did not overlap, suggesting that predator-

induced TGP and WGP do not share common genes. This finding is

consistent with Hales et al. (2017), who reported that the DEGs for

predator-induced WGP and TGP in *D. ambigua* were largely distinct. In

contrast, Stein et al. (2018) reported that there were unique sets of genes

related to the different forms of plasticity but also identified a common core

set of genes in *G. aculeatus*. Despite the absence of common genes in our

study, the DEGs in F2 seemed to be involved in the same main functions as

those in F1, i.e., metabolism and transcriptional regulation. Consequently,

these results suggest that common functions are used for predator-induced

WGP and TGP via different genomic pathways. This finding has important

implications for the evolution of predator-induced plasticity as it suggests

that WGP and TGP can evolve independently.

Conclusion

This study confirmed predator-induced WGP and TGP in *P. acuta*, showing

that antipredator traits (escape behaviour, shell thickness, shell length and

shell shape) were induced when snails were exposed during their own

development or when they originated from parents exposed to predator

cues (but not exposed themselves). These within- and transgenerational

phenotypic responses were both associated with differential patterns of

gene expression. However, the transcriptomic signal was lower for TGP (a

lower number of DEGs), and surprisingly no common genes were shared

between WGP and TGP, although the main functions were similar

(metabolism and transcription regulation). Expression analysis revealed

that only a small set of genes (112 in F1 and 23 in F2) exhibited strong

differential expression responses consistently in both generations. Several

hypotheses have been proposed to explain these distinct transcriptomic

patterns between predator-induced within-generational and

transgenerational plasticity. Further research is needed to overcome the

limitations of this study by examining transcriptomic signals throughout

development, across multiple tissues, and under diverse environmental

conditions. This study provides a more integrative view of the underlying

molecular mechanisms of WGP and TGP, addressing an important gap in

knowledge concerning important processes for adaptation to changing

environments.

Acknowledgements. We are grateful to Cyril Degletagne who helped in
experimental works and to the Rhône-Alpes bioinformatic centre (PRABI)
that has supported bioinformatics analyses. This work was financially
supported by the BioEnviS research federation (FR3728) of Lyon University
and the TEATIME (ANR-21-CE02-0005) grants from the French National
Research Agency (ANR).

Authors Contributions. E.L., S.P. and T.L. conceived the project. J.T., S.P.
and E.L. performed the experiments. L.K. performed RNA extractions and
libraries. L.D., J.T. and M.H. analysed phenotypic data. L.D., N.S and T.L.
performed bioinformatic molecular analyses. L.D. wrote the first version of
the manuscript. All authors revised the manuscript.

Ethics. This work did not require ethical approval from a human subject or
animal welfare committee.

Conflict of interest declaration. We declare we have no competing
interests.

Funding. This work was financially supported by the TEATIME (ANR-21-
CE02-0005) grants from the French National Research Agency (ANR).

Data accessibility. RNA-seq data have been deposited in the European
Nucleotide Archive (ENA) under the accession number [PRJEB87216].
The raw sequencing data are publicly available at [
https://www.ebi.ac.uk/ena/browser/view/PRJEB87216]. Additionally, scripts
used for the analysis, along with the non-genomic data, have been made

available on Zenodo. The repository can be accessed at [

10.5281/zenodo.15043538].

References

Agrawal, A. A., Laforsch, C., & Tollrian, R. (1999). Transgenerational

induction of defences in animals and plants. *Nature*, *401*(6748),

Article 6748. https://doi.org/10.1038/43425

Aubin-Horth, N., & Renn, S. C. P. (2009). Genomic reaction norms: Using

integrative biology to understand molecular mechanisms of

phenotypic plasticity. *Molecular Ecology*, *18*(18), 3763–3780.

https://doi.org/10.1111/j.1365-294X.2009.04313.x

Auld, J. R., & Houser, R. (2015). Age-dependent effects of predation risk on

reproductive success in a freshwater snail. *Evolution*, *69*(10), 2793–

2798. https://doi.org/10.1111/evo.12769

Auld, J. R., & Relyea, R. A. (2011). Adaptive plasticity in predator-induced

defenses in a common freshwater snail: Altered selection and mode

of predation due to prey phenotype. *Evolutionary Ecology*, *25*(1),

189–202. https://doi.org/10.1007/s10682-010-9394-1

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). *Fitting Linear Mixed-*

Effects Models using lme4 (arXiv:1406.5823). arXiv.

https://doi.org/10.48550/arXiv.1406.5823

Beaty, L. E., Wormington, J. D., Kensinger, B. J., Bayley, K. N., Goeppner,

S. R., Gustafson, K. D., & Luttbeg, B. (2016). Shaped by the past,

acting in the present: Transgenerational plasticity of anti-predatory

traits. *Oikos*, *125*(11), 1570–1576. https://doi.org/10.1111/oik.03114

Bell, A. M., & Hellmann, J. K. (2019). An Integrative Framework for

Understanding the Mechanisms and Multigenerational

Consequences of Transgenerational Plasticity. *Annual Review of*

Ecology, Evolution, and Systematics, *50*(1), 97–118.

https://doi.org/10.1146/annurev-ecolsys-110218-024613

Bell, A. M., & Stein, L. R. (2017). Transgenerational and developmental

plasticity at the molecular level: Lessons from Daphnia. *Molecular*

Ecology, *26*(19), 4859–4861. https://doi.org/10.1111/mec.14327

Blackwell, T. K., Sewell, A. K., Wu, Z., & Han, M. (2019). TOR Signaling in

Caenorhabditis elegans Development, Metabolism, and Aging.

Genetics, *213*(2), 329–360.

https://doi.org/10.1534/genetics.119.302504

Blum, M., Andreeva, A., Florentino, L. C., Chuguransky, S. R., Grego, T.,

Hobbs, E., Pinto, B. L., Orr, A., Paysan-Lafosse, T., Ponamareva, I.,

Salazar, G. A., Bordin, N., Bork, P., Bridge, A., Colwell, L., Gough,

J., Haft, D. H., Letunic, I., Llinares-López, F., … Bateman, A. (2024).

InterPro: The protein sequence classification resource in 2025.

Nucleic Acids Research, *53*(D1), D444–D456.

https://doi.org/10.1093/nar/gkae1082

Bray, N. L., Pimentel, H., Melsted, P., & Pachter, L. (2016). Near-optimal

probabilistic RNA-seq quantification. *Nature Biotechnology*, *34*(5),

Article 5. https://doi.org/10.1038/nbt.3519

Bukowski, S. J., & Auld, J. R. (2014). The effects of calcium in mediating

the inducible morphological defenses of a freshwater snail, Physa

acuta. *Aquatic Ecology*, *48*(1), 85–90.

https://doi.org/10.1007/s10452-013-9468-6

Burton, T., & Metcalfe, N. B. (2014). Can environmental conditions

experienced in early life influence future generations? *Proceedings*

of the Royal Society B: Biological Sciences, *281*(1785), 20140311.

https://doi.org/10.1098/rspb.2014.0311

Clark, M. S., Suckling, C. C., Cavallo, A., Mackenzie, C. L., Thorne, M. A.

S., Davies, A. J., & Peck, L. S. (2019). Molecular mechanisms

underpinning transgenerational plasticity in the green sea urchin

Psammechinus miliaris | Scientific Reports. *Scientific Reports*.

https://www-nature-com.docelec.univ-lyon1.fr/articles/s41598-018-

37255-6

Colicchio, J. M., & Herman, J. (2020). Empirical patterns of environmental

variation favor adaptive transgenerational plasticity. *Ecology and*

Evolution, *10*(3), 1648–1665. https://doi.org/10.1002/ece3.6022

Danchin, É., Charmantier, A., Champagne, F. A., Mesoudi, A., Pujol, B., &

Blanchet, S. (2011). Beyond DNA: Integrating inclusive inheritance

into an extended theory of evolution. *Nature Reviews Genetics*,

12(7), Article 7. https://doi.org/10.1038/nrg3028

DeWitt, T. J. (1998). Costs and limits of phenotypic plasticity: Tests with

predator-induced morphology and life history in a freshwater snail.

Journal of Evolutionary Biology, *11*(4), 465–480.

https://doi.org/10.1046/j.1420-9101.1998.11040465.x

Duncan, E. J., Gluckman, P. D., & Dearden, P. K. (2014). Epigenetics,

plasticity, and evolution: How do we link epigenetic change to

phenotype? *Journal of Experimental Zoology Part B: Molecular and*

Developmental Evolution, *322*(4), 208–220.

https://doi.org/10.1002/jez.b.22571

Ezard, T. H. G., Prizak, R., & Hoyle, R. B. (2014). The fitness costs of

adaptation via phenotypic plasticity and maternal effects. *Functional*

Ecology, *28*(3), 693–701.

Fallet, M., Luquet, E., David, P., & Cosseau, C. (2020). Epigenetic

inheritance and intergenerational effects in mollusks. *Gene*, *729*,

144166. https://doi.org/10.1016/j.gene.2019.144166

Galloway, L. F., & Etterson, J. R. (2007). Transgenerational Plasticity Is

Adaptive in the Wild. *Science*, *318*(5853), 1134–1136.

https://doi.org/10.1126/science.1148766

Gilbert, S. F. (2001). Ecological Developmental Biology: Developmental

Biology Meets the Real World. *Developmental Biology*, *233*(1), 1–

12. https://doi.org/10.1006/dbio.2001.0210

Gustafson, K., Kensinger, B., Bolek, M., & Luttbeg, B. (2014). Distinct snail

(Physa) morphotypes from different habitats converge in shell shape

and size under common garden conditions. *Evolutionary Ecology*

Research, *16*, 77–89.

Hales, N. R., Schield, D. R., Andrew, A. L., Card, D. C., Walsh, M. R., &

Castoe, T. A. (2017). Contrasting gene expression programs

correspond with predator-induced phenotypic plasticity within and

across generations in Daphnia. *Molecular Ecology*, *26*(19), 5003–

5015. https://doi.org/10.1111/mec.14213

Hammer, O., Harper, D. A. T., & Ryan, P. D. (2001). PAST: Paleontological

Statistics Software Package for Education and Data Analysis.

Palaeontologia Electronica.

Herman, J., & Sultan, S. (2011). Adaptive Transgenerational Plasticity in

Plants: Case Studies, Mechanisms, and Implications for Natural

Populations. *Frontiers in Plant Science*, *2*.

https://www.frontiersin.org/articles/10.3389/fpls.2011.00102

Jablonka, E., & Raz, G. (2009). Transgenerational epigenetic inheritance:

Prevalence, mechanisms, and implications for the study of heredity

and evolution. *The Quarterly Review of Biology*, *84*(2), 131–176.

https://doi.org/10.1086/598822

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C.,

McWilliam, H., Maslen, J., Mitchell, A., Nuka, G., Pesseat, S.,

Quinn, A. F., Sangrador-Vegas, A., Scheremetjew, M., Yong, S.-Y.,

Lopez, R., & Hunter, S. (2014). InterProScan 5: Genome-scale

protein function classification. *Bioinformatics*, *30*(9), 1236–1240.

https://doi.org/10.1093/bioinformatics/btu031

Katewa, S. D., & Kapahi, P. (2011). Role of TOR signaling in aging and

related biological processes in Drosophila melanogaster.

Experimental Gerontology, *46*(5), 382.

https://doi.org/10.1016/j.exger.2010.11.036

Khorchid, A., & Ikura, M. (2002). How calpain is activated by calcium.

Nature Structural Biology, *9*(4), 239–241.

https://doi.org/10.1038/nsb0402-239

Kikuchi, D. W., Allen, W. L., Arbuckle, K., Aubier, T. G., Briolat, E. S.,

Burdfield-Steel, E. R., Cheney, K. L., Daňková, K., Elias, M.,

Hämäläinen, L., Herberstein, M. E., Hossie, T. J., Joron, M., Kunte,

K., Leavell, B. C., Lindstedt, C., Lorioux-Chevalier, U., McClure, M.,

McLellan, C. F., … Exnerová, A. (2023). The evolution and ecology

of multiple antipredator defences. *Journal of Evolutionary Biology*,

36(7), 975–991. https://doi.org/10.1111/jeb.14192

Kuijper, B., & Hoyle, R. B. (2015). When to rely on maternal effects and

when on phenotypic plasticity? *Evolution*, *69*(4), 950–968.

https://doi.org/10.1111/evo.12635

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest

Package: Tests in Linear Mixed Effects Models. *Journal of Statistical*

Software, *82*(13). https://doi.org/10.18637/jss.v082.i13

Ledón-Rettig, C. C. (2023). A transcriptomic investigation of heat-induced

transgenerational plasticity in beetles. *Biological Journal of the*

Linnean Society, *138*(3), 318–327.

https://doi.org/10.1093/biolinnean/blac151

Lee, P. C., Bussière, L. F., Webber, C. E., Poole, J. H., & Moss, C. J.

(2013). Enduring consequences of early experiences: 40 year

effects on survival and success among African elephants

(Loxodonta africana). *Biology Letters*, *9*(2), 20130011.

https://doi.org/10.1098/rsbl.2013.0011

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold

change and dispersion for RNA-seq data with DESeq2. *Genome*

Biology, *15*(12), 550. https://doi.org/10.1186/s13059-014-0550-8

Luquet, E., Léna, J.-P., David, P., Joly, P., Lengagne, T., Perrin, N., &

Plénet, S. (2011). Consequences of genetic erosion on fitness and

phenotypic plasticity in European tree frog populations (Hyla

arborea). *Journal of Evolutionary Biology*, *24*(1), 99–110.

https://doi.org/10.1111/j.1420-9101.2010.02138.x

Luquet, E., & Tariel, J. (2016). Offspring reaction norms shaped by parental

environment: Interaction between within- and trans-generational

plasticity of inducible defenses. *BMC Evolutionary Biology*, *16*(1),

Article 1. https://doi.org/10.1186/s12862-016-0795-9

MacLeod, K. J., Monestier, C., Ferrari, M. C. O., McGhee, K. E., Sheriff, M.

J., & Bell, A. M. (2022). Predator-induced transgenerational

plasticity in animals: A meta-analysis. *Oecologia*, *200*(3), 371–383.

https://doi.org/10.1007/s00442-022-05274-w

Mommer, B. C., & Bell, A. M. (2014). Maternal experience with predation

risk influences genome-wide embryonic gene expression in

threespined sticklebacks (Gasterosteus aculeatus). *PloS One*, *9*(6),

e98564. https://doi.org/10.1371/journal.pone.0098564

Orsini, L., Brown, J. B., Shams Solari, O., Li, D., He, S., Podicheti, R.,

Stoiber, M. H., Spanier, K. I., Gilbert, D., Jansen, M., Rusch, D. B.,

Pfrender, M. E., Colbourne, J. K., Frilander, M. J., Kvist, J.,

Decaestecker, E., De Schamphelaere, K. A. C., & De Meester, L.

(2018). Early transcriptional response pathways in Daphnia magna

are coordinated in networks of crustacean-specific genes. *Molecular*

Ecology, *27*(4), 886–897. https://doi.org/10.1111/mec.14261

Pigliucci, M. (2005). Evolution of phenotypic plasticity: Where are we going

now? *Trends in Ecology & Evolution*, *20*(9), 481–486.

https://doi.org/10.1016/j.tree.2005.06.001

Prizak, R., Ezard, T. H. G., & Hoyle, R. B. (2014). Fitness consequences of

maternal and grandmaternal effects. *Ecology and Evolution*, *4*(15),

3139–3145. https://doi.org/10.1002/ece3.1150

Reger, J., Lind, M. I., Robinson, M. R., & Beckerman, A. P. (2018).

Predation drives local adaptation of phenotypic plasticity. *Nature*

Ecology & Evolution, *2*(1), 100–107. https://doi.org/10.1038/s41559-

017-0373-6

Rozenberg, A., Parida, M., Leese, F., Weiss, L. C., Tollrian, R., & Manak, J.

R. (2015). Transcriptional profiling of predator-induced phenotypic

plasticity in Daphnia pulex. *Frontiers in Zoology*, *12*(1), 18.

https://doi.org/10.1186/s12983-015-0109-x

RStudio Team. (2020). *RStudio: Integrated Development Environment for*

R.

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to

ImageJ: 25 years of Image Analysis. *Nature Methods*, *9*(7), 671–

675.

Segers, F. H. I. D., & Taborsky, B. (2011). Juvenile exposure to predator

cues induces a larger egg size in fish. *Proceedings of the Royal*

Society B: Biological Sciences, *279*(1731), 1241–1248.

https://doi.org/10.1098/rspb.2011.1290

Sharda, S., Zuest, T., Erb, M., & Taborsky, B. (2021). Predator-induced

maternal effects determine adaptive antipredator behaviors via egg

composition. *Proceedings of the National Academy of Sciences*,

118(37), e2017063118. https://doi.org/10.1073/pnas.2017063118

Skottene, E., Tarrant, A. M., Altin, D., Olsen, R. E., Choquet, M., & Kvile, K.

Ø. (2020). Lipid metabolism in Calanus finmarchicus is sensitive to

variations in predation risk and food availability. *Scientific Reports*,

10(1), 22322. https://doi.org/10.1038/s41598-020-79165-6

Slos, S., & Stoks, R. (2008). Predation risk induces stress proteins and

reduces antioxidant defense. *Functional Ecology*, *22*(4), 637–642.

https://doi.org/10.1111/j.1365-2435.2008.01424.x

Smithson, M., Thorson, J. L. M., Sadler-Riggleman, I., Beck, D., Skinner, M.

K., & Dybdahl, M. (2020). Between-Generation Phenotypic and

Epigenetic Stability in a Clonal Snail. *Genome Biology and*

Evolution, *12*(9), 1604–1615. https://doi.org/10.1093/gbe/evaa181

Stein, L. R., Bukhari, S. A., & Bell, A. M. (2018). Personal and

transgenerational cues are nonadditive at the phenotypic and

molecular level. *Nature Ecology & Evolution*, *2*(8), 1306–1311.

https://doi.org/10.1038/s41559-018-0605-4

Tariel, J., Luquet, É., & Plénet, S. (2020). Interactions Between Maternal,

Paternal, Developmental, and Immediate Environmental Effects on

Anti-predator Behavior of the Snail Physa acuta. *Frontiers in*

Ecology and Evolution, *8*.

https://www.frontiersin.org/articles/10.3389/fevo.2020.591074

Tariel, J., Plénet, S., & Luquet, E. (2020a). How do developmental and

parental exposures to predation affect personality and immediate

behavioural plasticity in the snail Physa acuta? *Proceedings of the*

Royal Society B: Biological Sciences, *287*(1941), 20201761.

https://doi.org/10.1098/rspb.2020.1761

Tariel, J., Plénet, S., & Luquet, É. (2020b). Transgenerational Plasticity in

the Context of Predator-Prey Interactions. *Frontiers in Ecology and*

Evolution, *8*. https://doi.org/10.3389/fevo.2020.548660

Tariel, J., Plénet, S., & Luquet, É. (2020c). Transgenerational plasticity of

inducible defences: Combined effects of grand-parental, parental

and current environments. *Ecology and Evolution*, *10*(5), 2367–

2376. https://doi.org/10.1002/ece3.6046

Tariel-Adam, J., Luquet, É., & Plénet, S. (2023). Sensitive windows for

within- and trans-generational plasticity of anti-predator defences.

Peer Community Journal, *3*. https://doi.org/10.24072/pcjournal.304

Thorson, J. L. M., Smithson, M., Beck, D., Sadler-Riggleman, I., Nilsson,

E., Dybdahl, M., & Skinner, M. K. (2017). Epigenetics and adaptive

phenotypic variation between habitats in an asexual snail. *Scientific*

Reports, *7*(1), 14139. https://doi.org/10.1038/s41598-017-14673-6

Thorson, J. L. M., Smithson, M., Sadler-Riggleman, I., Beck, D., Dybdahl,

M., & Skinner, M. K. (2019). Regional epigenetic variation in asexual

snail populations among urban and rural lakes. *Environmental*

Epigenetics, *5*(4), dvz020. https://doi.org/10.1093/eep/dvz020

Tollrian, R. (1995). Predator-Induced Morphological Defenses: Costs, Life

History Shifts, and Maternal Effects in Daphnia Pulex. *Ecology*,

76(6), 1691–1705. https://doi.org/10.2307/1940703

Turner, A. M., Fetterolf, S. A., & Bernot, R. J. (1999). Predator identity and

consumer behavior: Differential effects of fish and crayfish on the

habitat use of a freshwater snail. *Oecologia*, *118*(2), 242–247.

https://doi.org/10.1007/s004420050724

Viney, M., & Diaz, A. (2012). Phenotypic plasticity in nematodes:

Evolutionary and ecological significance. *Worm*, *1*(2), 98–106.

https://doi.org/10.4161/worm.21086

Walsh, M. R., Castoe, T., Holmes, J., Packer, M., Biles, K., Walsh, M.,

Munch, S. B., & Post, D. M. (2016). Local adaptation in

transgenerational responses to predators. *Proceedings of the Royal*

Society B: Biological Sciences, *283*(1823), 20152271.

https://doi.org/10.1098/rspb.2015.2271

Yin, J., Zhou, M., Lin, Z., Li, Q. Q., & Zhang, Y.-Y. (2019). Transgenerational

effects benefit offspring across diverse environments: A meta-

analysis in plants and animals. *Ecology Letters*, *22*(11), 1976–1986.

https://doi.org/10.1111/ele.13373