Transcriptomic bases of within- and trans-generational predator-induced plasticity in the
freshwater snail *Physa acuta*

Authors:

Léo Dejeux[1], Nathanaëlle Saclier[2], Juliette Tariel-Adam[3], Maxime Hoareau[1], Tristan Lefébure[1], Lara
Konecny[1], Sandrine Plénet[1], Emilien Luquet[1, 4]

[1] Université Claude Bernard Lyon 1, LEHNA UMR 5023, CNRS, ENTPE, F-69622, Villeurbanne, France

[2] ISEM, Univ Montpellier, CNRS, IRD, Montpellier, France

[3] Fish Lab, Department of Biological Sciences, Macquarie University, Sydney, NSW, Australia

[4] Institut universitaire de France (IUF)

Abstract

Inducible defences in response to predation risk are a well-known example of adaptive phenotypic

plasticity. While inducible defences have been mainly studied within a generation (within-generational

plasticity), there is now clear evidence that the ancestral exposure to predation risk can influence the

defences expressed by the offspring, even if they have not been exposed themselves (transgenerational

plasticity). The molecular mechanisms allowing the transmission of environmental information across

generations are still misunderstood. In this study, we combined measures of anti-predator responses

(behavioural and morphological) with transcriptomic investigations across two generations in the

freshwater snail *Physa acuta*. We hypothesised that both within- and transgenerational plasticity would

induce phenotypic changes associated with differential gene expression. Our results confirmed within-

and transgenerational plasticity: F1 snails have responded to predator-cue exposure by increasing

escape behaviour, reducing shell length, and developing thicker and slender shells and F2 from exposed

parents have shown similar responses. Within- and transgenerational plasticity were accompanied by

the differential expression of 78 genes (77 up- and 1 down-regulated) and 14 differentially expressed

genes (13 up- and 1 down-regulated), respectively. Within- and transgenerational plasticity did not

share common differentially expressed genes but some differentially expressed genes in F1 are also

highly expressed in F2 and overall, the molecular functions were similar, involving metabolism and

transcription regulation. The quantitatively low molecular response associated with transgenerational

plasticity suggests that gene expression changes induced by parental exposure may have occurred early

in development and have decreased at adult stage.

INTRODUCTION

Phenotypic plasticity is the ability of a genotype to produce alternative phenotypes under different environmental conditions (Pigliucci, 2005). Plastic responses are widespread across the phylogenetic spectrum (e.g. in plants: Galloway & Etterson, 2007; in amphibians: Luquet et al., 2015; in fishes: Segers & Taborsky, 2011; in ostracods: Walsh et al., 2016), and are a major concern in evolutionary biology as they can be adaptive, i.e. increasing fitness of organisms facing environmental changes (Pigliucci, 2005). Phenotypic plasticity classically refers to within-generational plasticity (WGP), meaning that it occurs within one generation. Recently, plasticity has also been found to occur across generations (transgenerational plasticity, TGP) where the phenotype of an organism is influenced by the environment experienced by the previous generations (Bell & Hellmann, 2019). Interest in TGP has grown rapidly in recent years with evidence that TGP is at least as widespread as WGP (Jablonka & Raz, 2009; Tariel, Plénet, et al., 2020b). Although there are numerous examples of TGP generating seemingly adaptive offspring phenotypes, its adaptive role is expected to be more challenging than that of WGP due to patterns of environmental variation across generations (Colicchio & Herman, 2020). Indeed, it requires a strong temporal autocorrelation between environments of distinct generations to be selected (Ezard et al., 2014; Kuijper & Hoyle, 2015; Prizak et al., 2014) and a transmission system that allows the induction of adaptive phenotypes across generations (Fallet et al., 2020). The inheritance mechanisms appear diverse, from epigenetic changes to parental transmission of cytoplasmic elements (small RNAs, nutrients, hormones, proteins), parental care and cultural transmission (Bell & Hellmann, 2019; Danchin et al., 2011).

Regardless of the transmission mechanisms, they should lead in the offspring to the modification of the expression of some of the genes involved in the phenotypes, at least in some tissues and at some stages during the development. While transgenerational plasticity studies encompass all aspects of phenotype (including physiological states, morphological traits, behaviour and life-history traits),

transgenerational plastic response at the molecular level is certainly the least studied. This is yet a

necessary first step to unravel the molecular mechanisms of plasticity in general and their

transgenerational inheritance. The most compelling explorations of changes in gene expression profiles

in response to both developmental (WGP) and parental (TGP) environments come from the studies of

Hales et al. (2017) on *Daphnia ambigua* and Stein et al. (2018) on three spined stickleback

(*Gasterosteus aculeatus*). These studies compared gene expression between individuals that had been

exposed to predator cues relative to unexposed individuals within a generation (WGP genes) and

among their offspring (descendants of unexposed or exposed parents; TGP genes). Stein et al. (2018)

found that the same group of genes changed their expression in offspring whether the predation risk

was experienced by the offspring (TGP), the father (WGP), or both. This suggests that a core set of

genes is consistently affected by predation risk regardless of who experiences it. On the other hand,

Hales et al. (2017) found that changes in the expression of genes that respond to TGP differed

depending on whether exposure to predation was experienced by parents or grandparents. They also

showed that very few genes were common to both WGP and TGP. Note that Clark et al. (2019) also

showed that TGP gene expression profiles of larvae of sea urchin (*Psammechinus miliaris*) spawned at

low pH from pre-acclimated adults differed to those of larvae produced from adults exposed to ambient

pH, but the authors did not explore the WGP genes. These few results highlight the necessity to

investigate together WGP and TGP at the molecular level. Beyond the inheritance mechanisms allowing

organisms to memorise past environmental conditions to adjust their future phenotype, we need to

know whether the two forms of plasticity are distinct at the molecular level and potentially free to

evolve independently of one another (Bell & Stein, 2017).

Predation risk is a major selection pressure for most species and results in the evolution of defences in

prey (Kikuchi et al., 2023). As predation risk often varies over time and space, it favours the evolution

of phenotypic plasticity in defences (Reger et al., 2018; Tariel, Luquet, et al., 2020; Viney & Diaz, 2012).

Indeed, while some defences against predators are constitutive, plastic defences are only induced when

prey detect some predator cues in their environment (e.g. visual cues, chemical cues such as predator

odours, or alarm signals released by conspecifics). These induced defences are classic examples of

adaptive WGP: prey can increase their fitness by producing defences only when predators are present.

The links between prey-predator interactions and WGP have been studied for a long time (e.g Tariel et

al., 2020; Tollrian, 1995). Plastic defences can also be induced in offspring after their parents or more

distant ancestors have detected predator cues. This transgenerational induction of defences was first

highlighted in Daphnia morphology by Agrawal et al., (1999) and has since been repeatedly shown in

multiple species and defences (review in MacLeod et al., 2022; Tariel et al., 2020).

In this context, this work aims to unravel the molecular mechanisms underlying the expression of WGP

and TGP of defences in the context of predator-prey interactions. We used *Physa acuta*, a freshwater

snail known for their within- and transgenerational anti-predator defences such as escape behaviour

(snails crawl out of the water when they or their parents detect predator cues; Luquet & Tariel, 2016;

Tariel et al., 2020a; Turner et al., 1999), change in shell size and shape, increase in shell thickness and

overall increase in shell-crush resistance (*e.g.* shell harder to handle or to crush by predators;(Auld &

Relyea, 2011; Beaty et al., 2016; DeWitt, 1998; Tariel et al., 2020c). We combined a two-generation

experiment to generate WGP and TGP of defences (behavioural and morphological responses as

previously done in e.g. Luquet & Tariel, 2016) with a transcriptomic investigation. We used

transcriptome sequencing and assembly to specifically investigate the transcriptome mantle skirt gene

expression (mantle skirt is the tissue that synthesises the shell). We expected that gene expression

should be affected by current (WGP) and parental (TGP) exposure to predator cues. More specifically,

as WGP and TGP induce similar phenotypic responses (e.g. increase in shell thickness, (Tariel et al.,

2020), we expected the two processes share the same core set of genes.

MATERIALS & METHODS

Animal collection, rearing conditions and experimental design

Adult *P. acuta* snails were collected from a wild population in a lentic backwater of the Rhône River (Lyon, France, 45° 48'6"N, 4° 55'33"E) in September 2018. Individuals from this population have been repeatedly used in previous studies on predator-induced plasticity (Luquet & Tariel, 2016) and are naturally exposed to predation by the crayfish *Faxonius limosus*. The experiment was conducted in our laboratory in a temperature-controlled chamber at 25 °C with a photoperiod of 12/12h. The wild snails constituting the F0 generation were pooled in a 10 L plastic box filled with dechlorinated tap water (control water hereafter) and interbreed overnight. They were then isolated in 80 mL plastic boxes (4.5 x 6 cm) filled with control (without predator cues) water to lay eggs for 3 days. Eggs within boxes constitute F1 matrilines. They developed until hatching at 25°C (ca. seven days). Siblings of each F1 matriline developed for 10 days in control water. Then, 10 siblings of 20 F1 matrilines were split into two developmental environments: five siblings remained in control (F1C) water and five siblings moved to predator-cue (F1P) water. F1 siblings developed together in 80 mL plastic boxes for seven days before being isolated in the same environment (control or predator-cue water) and kind of plastic boxes for 24 days (i.e. 41 days old snails from hatching to measurements). At the end, one F1 snail per matriline and treatment was frozen at -80°C for molecular analysis and all remaining F1 snails were measured (see § Trait measurements).

To generate the F2 generation, we pooled F1 snails from the same treatment in an aquarium filled with control water and let them copulate for 24h. As previously, F1 snails were then isolated and laid eggs for 3 days to produce the F2 matrilines. F2 snails from the two parental environments (C and P) were all reared in control water (noted F2CC and F2PC thereafter) with the same protocol steps as previously for F1 snails, except that we measured F2 snails at an older age than the F1 generation because of a

slower growth. As for F1, one F2 snail per matriline and treatment was frozen for molecular analysis

and the others were measured.

Snails have been fed *ad libitum* with boiled and mixed lettuce. Water (according to treatment) and food

was renewed twice a week. The predator-cue water was water in which *F. limosus* crayfish have been

raised (one crayfish/5L), fed with snails, for 3 days and in which we smashed *P. acuta* adult snails (one

snail/5L) for 1h before using the predator-cue water (these smashed snails were then used as crayfish

food).

Behavioural and morphological responses

In order to check that predator cue exposition triggered WGP and TGP, we measured escape behaviour

and shell morphology on F1 and F2 snails (339 snails: F1C=76, F1P=73, F2CC=89, F2PC=101). Crawling

out of the water (position above the water surface) is considered as allowing snails to escape from

benthic predators such as crayfishes (DeWitt, 1998; Tariel et al., 2020a). Shorter and narrower shell and

aperture dimensions and thicker shell are adaptive anti-predator responses (Auld & Relyea, 2011).

These phenotypic traits have been recurrently involved in transgenerational responses of *P. acuta* to

predator cues (Beaty et al., 2016; Luquet & Tariel, 2016).

Escape behaviour - The position above on or below the water surface of each individual was recorded

in their rearing vials one week before measurements for each generation, four hours after water

change.

Shell thickness, shell length and morphology - Shell thickness was directly measured with an electronic

calliper at the nearest 0.01 mm at the edge of the aperture. After, a photograph of each snail (aperture

upwards) was taken with an Olympus SC50 camera installed on an Olympus SZX9 binocular and its

Olympus DF PLAPO 1X-2 objective at a x8 magnification. Snail shell length, shell width, aperture length

and aperture width were measured using ImageJ (Schneider et al., 2012). To estimate potential changes

in global snail shell morphology according to treatment, we used geometric methods (Gustafson et al.,

2014). On each snail shell, we digitised 19 landmarks (Fig. 1) using the software ImageJ. Snails that

lacked homologous landmarks because of broken shells (typically broken apertures and apexes) or

small size (lack of the third shell spire) were removed from the analyses, resulting in 336 snails (F1C=76,

F1P=72, F2CC=89, F2PC=99). To compare shell shapes according to treatments, we scaled the shapes

using the size metric defined below (eqn. 1). Translational components were removed by translating all

shapes so that the centroids lie at the origin. Rotational components were removed by rotating all

shapes until the Procrustes distance was minimised with the first shell fixed as reference.

$$s_k = \sqrt{\sum_{i=1}^{19} \left((x_{k,i} - \bar{x}_k)^2 + (y_{k,i} - \bar{y}_k{}^2) \right)}$$

185   **Equation 1:** Procuste's equation used to scale the shapes using the size metric s_k is the size of the k-th shell, $x_{k,i}$

186   and $y_{k,i}$ is the x and y component of the i-th landmark, and $(\bar{x}_k, \bar{y}_k)$, is the centroid of the k-th shell.

Figure 1: Locations of the different landmarks (circle symbol) and pseudo-landmarks (positioned using a geometric method,

square symbol) on *P. acuta* shell. Pseudo-landmarks have been created using a geometric method: first, two tangents (T1 and

T2) have been drawn from landmark 1 (apex). T1 touches the curve of the habitation spire (5) and T2 touches the external

edge of the aperture (10). From the two points of tangency (points 5 and 10), two perpendicular lines to the tangents have

been traced (P1 and P2). The intersections with the columellar edge of the aperture, the external edge of the aperture and

the curve of the habitation spire give additional pseudo-landmarks (6, 9, 18 and 19). The length of the shell L (straight line

passing by 1 and 8) gives the pseudo-landmark 18 at the intersection with the columellar edge of the aperture).

Statistical analyses

Escape behaviour - The effects of predator cues on escape behaviour (i.e. snail position above/on or

below the water surface) were analysed using generalised linear mixed models (GLMM) assuming a

binomial distribution (logit link function). To test for WGP and TGP in F1 and F2 snails respectively,

treatment (parental treatment) was a fixed effect; family was considered as a random intercept. We

tested significance of fixed and random effects with likelihood ratio tests.

Shell thickness, shell length - To analyse the effects of predator cues on shell length and shell thickness,

we performed linear mixed models with treatment (predation; parental treatment for F2) as fixed effect

and family as a random intercept. We used restricted maximum likelihood estimation and Kenward and

Roger's approximation for degrees of freedom. We tested significance of fixed effects with type II F-

tests (Kuznetsova et al., 2017) and significance of random effect with likelihood ratio test.

Linear mixed models were performed using the lme4 package (Bates et al., 2014) on R software (version

R-4.4.1 for Windows, RStudio Team, 2020).

Shell morphology - The effects of predator cues on shell shape were analysed using principal

component analyses (PCA) and were visualised with the 2D landmarks relative wraps analysis

functionality. All standardised shapes obtained by Procustes method were analysed using the PAST

software with a Student test (Hammer et al., 2001).

Molecular biology analysis

We dissected 30 snails (F1C = 5, F1P = 5, F2CC = 10, F2PC = 10) to extract the mantel skirt, the tissue

that synthesises the shell. Total RNA of the mantle was extracted by adding TRI Reagent (Molecular

Research Center MRC, TR118) to the samples and by homogenising tissues on ice with pistons and

agitator (Motor Mixer Clearline ref.045011 Dutscher). All remaining steps were carried out according

to the manufacturer's protocols (Molecular Research Center MRC, TR118). Then, the RNA was treated

with Turbo DNase enzyme (Turbo DNA free kit, Invitrogen, AM1907) and assayed by fluorescence using

a Qubit fluorometer (Qubit® RNA HS Assay Kits Molecular Probes, Invitrogen, Q32855). Library

construction was carried out using the NEBNext Ultra II RNA Library Prep Kit for Illumina (Biolabs New

England, NEB E7770, E7490, E7600). Libraries were randomly splitted in three groups and sequenced

on three lanes with an Illumina® HiSeq4000™ machine on the GenomEast platform hosted at the

IGBMC (GenomEast Platform – Institut de Génétique et de la Biologie Moléculaire et Cellulaire IGBMC,

Illkirch, France), resulting in 50 bp long single-end reads.

Differential gene expression analysis

FASTQC (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/)) was used to assess the quality

of the reads. We used a reference genome ORF *P. acuta* available on NCBI (Isolate: Inbredline_101_S28.

RefSeq: GCF_028476545.12). Gene expression levels were assessed on annotated transcripts using

Kallisto (-l 50 -s 10 options; (Bray et al., 2016). We used rounded effective counts as input for

differential gene expression using the DESeq2 package (Love et al., 2014), which performs differential

gene expression analysis using negative binomial distribution and shrinkage estimation for dispersions,

and outputs expression fold changes to improve the interpretability of estimates. We specified control

or predation treatment in the model (design formula: "design = ~treatment") within each generation

independently. Firstly, we checked for any structures in the dataset (normalised count table) by

performing a principal component analysis (PCA; DESeq2) on the 500 most differentially expressed

genes) across samples. Secondly, we used the Wald test (DESeq2 package) to test the significance of

differentially expressed genes between the control and predation treatment; we have considered a

significant differential expression when adjusted p-value < 0.05 and absolute Log2FoldChange (|LFC|)

greater than 3 (LFC is the logarithm base 2 of the fold change in gene expression between two

experimental conditions, providing a measure of both the magnitude and direction of differential

expression). In addition, we have compared the observed numbers of genes differentially expressed in

F1 and F2 to an expected distribution (generated by 1000 permutations) under the null hypothesis (i.e.

the differential expressions observed are due to the stochasticity of gene expression between

individuals). Thirdly, we determined the potential gene functions and the associated protein families

of all significant differentially expressed genes using the InterPro EMBL-EMI database (Paysan-Lafosse

et al., 2023). Finally, we investigated whether the differentially expressed genes (|LFC| > 3) in F1

parental generation were also differentially expressed in F2 offspring generation by comparing

heatmaps. We also compared the mean LFC of differentially expressed genes in F1 versus all the genes

expressed in F2 using a Wilcoxon test.

RESULTS

Predator-induced behaviour and shell morphology

Within-generational Plasticity - In F1 parental generation, the proportion of crawling out increased

significantly in predator-cue treatment from 7.9 % of snails out of the water in control treatment to

93.15 % in predator-cues (F[1,19.43] = 321.82, p < 0.001; Fig. 2A). Shell length significantly decreased

(-6.3%) in predator-cue treatment in F1 generation (F[1,128.71] = 15.46, p < 0.001; Fig. 2B). Shell

thickness significantly increased in predator-cue treatment (+18.8%; F[1,129.58] = 4.02, p = 0.047; Fig.

2C). The family random effect was always non-significant (behaviour: $\chi^2[1]$ = 0.690, p = 0.406; shell

length: $\chi^2[1]$ = 19.022, p<0.001; shell thickness:$\chi^2[1]$ = 2.6116, p = 0.106).

Predator-cues exposure induced changes in shell shape. The two first axes of PCA analysis accounted

for 59.2% of the shell shape variation (Fig. 3) and PCA axes 1-4 (74.2% cumulative shape variation) were

the only axes to each account for more than 5% of shape variation. PC1 showed a change in the

columellar area (close to landmark 18-19) from shells with relatively wide aperture (dilated columellar

area; Fig. 3C) to shells with relatively narrow aperture (compressed columellar area; Fig. 3D). PC2

showed a change in the slenderness of the shell from shells with relatively short spires (compressed

apex; Fig. 3B) and relatively wide apertures (dilated aperture) to shells with relatively short spires

(dilated apex) and narrow aperture widths (compressed aperture; Fig.3A).

Non-exposed snails showed the same range of deformation of columellar area than exposed snails

(PC2, t-test: t = 1.07, df = 144, p = 0.28). However, exposed shells are slender, with longer spires and

narrow aperture widths than non-exposed ones (PC1, t-test: t = 4.803, df = 144, p < 0.001). This

elongation is associated with the dilatation of the three last spires and a narrow aperture at the base

of the external edge. In contrast, non-exposed shells have a wider aperture at the base of the external

edge and smaller spires.

Figure 2: Effect of current exposure to predator cues (WGP) on crawling-out behaviour, shell length and shell thickness in the

F1 parental generation of *P. acuta*. Non-exposed and exposed to predator-cue treatments are symbolised with blue circles

and red triangles, respectively.

 Figure 3: Shape variation of *P. acuta* snails from generation F1C (blue circle symbol) and F1P (red triangle symbol) to predator

cues. Panels A, B ,C, D represent deformation grids modelizing shell shape on the extreme ends of the PCA axes. The 95%

confidence ellipses are represented for non-exposed and exposed to predator-cue treatments.

Transgenerational plasticity - In F2 offspring generation, the proportion of crawling-out marginally

increased in snails from predator-exposed parents, from 3.3% of snails out of the water from parental

non-exposed treatment to 8.75% from predator-exposed parents ($F_{[1,20.31]}$ = 3.929, p = 0.061; Fig.

4A). Snails from predator-exposed parents were significantly longer (+4.9%) and had a thicker shell

(+16.9%) than those from non-exposed parents ($F_{[1,170.87]}$ = 7.55, p = 0.006 and $F_{[1,174.11]}$ = 9.858,

p = 0.002, respectively; Fig. 4B-C). The family random effect was always non-significant (behaviour:

$\chi^2_{[1]}$ = 0.022, p = 0.882; shell length: $\chi^2_{[1]}$ = 31.685, p < 0.001; shell thickness: $\chi^2_{[1]}$ = 1.882, p = 0.17).

Parental exposure to predator cues induced changes in shell shape. The two first axes of PCA analysis

accounted for 63.8% of the shell shape variation and cumulative accounting for PCA axes 1-4 is 80.14%

(Fig. 5). PC1 displayed a change in shell aperture width; the shells with high PC1 scores (Fig. 5C) showed

a dilatation of the shell where the outer lip meets the parietal wall and a small compression of the

columellar area (i.e. a wider aperture). Shells with low PC1 scores (Fig. 5D) presented a narrower

aperture, a consequence of a dilated columellar area but compressed parietal wall and basal lip. PC2

showed a change in shell aperture width but also in shell length, from shells (Fig. 5C) with a dilated

outer lip and a compression of the apex (which tends to reduce spire length i.e. overall shell length) to

shells (Fig. 5D) showing elongated spires from dilated apex, with a narrower aperture due to outer lip

compression.

Snails from predator-exposed parents had a dilatation of the shell in the columellar area accompanied

by a small compression of the parietal wall (PC1: t = 2.067, p = 0.04). Conversely, there is no difference

in deformation of the outer lip, nor dilatation or compression of the apex between snails from both

parental environments (PC2: t = 0.59, p = 0.55).

Figure 4: Effect of parental exposure (TGP) on crawling-out behaviour, shell length and shell thickness in the F2 offspring

generation of *P. acuta*. Non-exposed and exposed to predator-cue treatments are symbolised with blue circles and red

triangles, respectively.

Figure 5: Shape variation of *P. acuta* snails from generation F2C (blue circle symbol) and F2P (red triangle symbol) to predator

cues. Panels A, B ,C, D represent deformation grids modelizing shell shape on the extreme ends of the PCA axes. The 95%

confidence ellipses are represented for non-exposed and exposed to predator-cue treatments.

Differential gene expression induced by predator-cues

A principal component analysis (PCA) was performed to visualise the grouping of read counts and to

help identify patterns on the first 500 most differentially expressed genes. The first and second principal

components (PC1) explained 18% and 11% of the variance respectively (Fig. 6). The plot did not indicate

an evident clustering between the control and predator-cue groups or between the F1 parental and F2

offspring generations (Fig. 5).

The differential expression analysis considering the effect of predator-cue exposure in the F1 parental

generation revealed a total of 78 genes with a differential expression (77 up-regulated genes and 1

down-regulated gene; Fig. S1). At the F2 offspring generation, the differential expression analysis

considering the parental effect of predator-cue exposure showed a total of 14 genes with a differential

expression (13 up-regulated and 1 down-regulated gene; Fig. S1). The permutation analyses confirmed

that the observed number of up-regulated genes in F1 and F2 are higher than under the null hypothesis

(Fig. S2).

The differentially expressed genes in F1 and in F2 were mainly involved in metabolism processes (e.g.

carbohydrate metabolism, calcium metabolism, fatty acid metabolism) and transcriptional regulation

(Fig. 7; see Tables S1 and S2 for details).

The comparison of the differentially expressed genes in F1 and in F2 revealed that the 77 up-regulated

genes in F1 were not differentially expressed in F2 offspring generation (Fig. 8, the differentially

expressed genes in F2 were not expressed in F1 see Fig. S3, an overall comparison of LFC F1 against LFC

F2 is shown in Fig. S4). However, we showed that the mean LFC of the 77 up-regulated genes in F1 was

significantly higher than the mean LFC of expressed genes in F2 (mean LFC = 0.172 vs 0.023; W =

748633, p < 0.001). Among the 77 up-regulated genes in F1, 7 were not expressed in F2, 64 were not

differentially expressed between non-exposed and exposed parental treatments with a very low LFC

(LFC = 0.05) and 6 were not differentially expressed (adjust p-value > 0.35) but with a high LFC (LFC

from 1.13 to 3.47; see Table S1).

Figure 6: PCA plot of the 30 transcriptome samples of *P. acuta* snails showing the divergence in differential expression

among the first 500 most DE genes. Non-exposed and exposed to predator cue individuals are represented in blue and red,

respectively. F1 individuals are represented by circles and plain ellipses and F2 individuals by triangles and dotted ellipses.

Figure 7: Proportions of classes of functions of the differentially expressed genes in F1 (A) and in F2 (B) based on the

InterPro EMBL-EMI database (Paysan-Lafosse et al. 2022).

Figure 8: Heatmap of read counts of 77 differentially expressed genes (|LFC| > 3) between non-exposed and exposed to

predator cues F1 snails (A) and the same genes between F2 snails from non-exposed and exposed parentals (B).

DISCUSSION

This work aimed to unravel the molecular mechanisms underlying the expression of within- and trans-

generational plasticity of anti-predator defences in *P. acuta*. We expected that gene expression should

be affected by current (WGP) and parental (TGP) exposure to predator cues, and that both WGP and

TGP share the same core set of genes. Exposure to predator cues results in changes in behaviour,

growth and morphology confirming that snails perceived predator cues and responded accordingly in

F1 (WGP) and that parental exposure influenced the F2 offspring phenotype (TGP). Our study

consistently revealed differential gene expression profiles between F1 non-exposed snails and exposed

to predator cues (WGP) and also between F2 snails according to parental exposure to predator cues

(TGP). However, the number of differentially expressed genes detected for TGP was lower than for WGP

and were different.

Gene expression induced by within-generational exposure to predator cues

As expected, snails exposed to predator cues crawled 85.25% more out of the water, had shells 18.8%

thicker, were 6.3% smaller and developed slender shells with longer spires and narrower aperture

width than unexposed snails. These behavioural and morphological responses are supposed to be

adaptive as they are consistent with an escape behaviour and an increase in shell crush resistance

(Tariel et al., 2020c; Tariel-Adam et al., 2023), increasing snail survival in face of crayfish predation

(Auld & Relyea, 2011). These predator-induced changes were associated with distinct patterns of gene

expression with 78 differentially regulated genes (77 up-regulated and 1 down-regulated genes).

Overall, this is consistent with several studies that explored the changes in gene expression induced by

the environment (e.g. Aubin-Horth & Renn, 2009; Herman & Sultan, 2011). In the context of predator-

prey interaction, the main knowledge about the transcriptomic responses of preys to predators comes

from studies on Daphnia species. Hales et al., (2017)) in *Daphnia ambigua* and Rozenberg et al. (2015)

in *Daphnia pulex* showed that anti-predator defences, including morphological traits, were associated

with 48 and 230 differentially regulated genes respectively. In *Daphnia galeata*, a species only

displaying shifts in life history traits when exposed to predator cues, no differentially expressed genes

between environmental groups was revealed (but some differences were detected at the scale of

genotypes; Tams et al., 2020, suggesting that traits such as morphological traits could be caused by or

correlated to much stronger differential gene expression. In *Daphnia magna*, Orsini et al. (2018)

revealed no changes in gene expression associated with a short-term exposure to fish kairomones.

Consequently, the number of differentially regulated genes involved in predator-induced responses

may depend on the defensive trait type and the strength of predator signals (e.g. cue concentration,

duration of exposure).

Our results showed that the most differentially up-regulated genes in snails exposed to predator cues

were mainly linked to metabolism, including metabolic stress and calcium metabolism. Firstly, 35.9%

of the up-regulated genes were coding for generic metabolic processes. For example, Complex I

intermediate-associated protein 30 is involved in respiratory metabolism, while a protein from AMP-

dependent synthetase/ligase domain is involved in the fatty acid metabolism (DiRusso et al., 2005;

Walker et al., 1992). Secondly, four up-regulated genes are involved in metabolic stress as for example

the gene coding for a protein belonging to the Regulatory Associated Protein of TOR (Target of

Rapamycin) family. This protein complex is well-known for its influence on various metabolic pathways

(Loewith et al., 2002; Saxton & Sabatini, 2017; Wu & Storey, 2021) and in particular in response to

abiotic stressors such as temperature and oxygen availability (Blackwell et al., 2019; Katewa & Kapahi,

2011). Since predation is known to generate metabolic stress in preys (Slos & Stoks, 2008), TOR proteins

could represent an important molecular mechanism involved in the anti-predator responses of preys;

however, to our knowledge, no studies have yet shown direct relationships between TOR proteins in

predator-prey interaction. Consistently with a metabolic stress induced by predator-cue exposure,

other metabolic genes (e.g. coding for Sulfhydryl Oxidase ALR/ERV) involved in oxidoreductase activity

and oxidative stress response have been identified (Braun et al., 1998). In addition, we found that

12,8% of up-regulated genes were linked to the regulation of transcription and gene expression; both

molecular processes are known to be intimately associated with metabolic stress (de Nadal et al., 2011;

Westfall et al., 2004). For example, a gene was coding for proteins from ARID DNA-binding and Lysine-

specific demethylase-like domains, both participating in chromatin remodelling and regulation of gene

expression (Kim et al., 2004; Lu et al., 2014). Thirdly, we interestingly found two up-regulated genes

coding for Annexin and Peptidase C2 protein families that are playing a role in calcium ion metabolism

(Khorchid & Ikura, 2002). Calcium being the main component of shell gastropods (i.e. calcium

carbonate in the form of aragonite; Heller & Magaritz, 1983; McKillop & Harrison, 1972), the

concentration of calcium in the water is known to influence the shell morphology of *P. acuta* and in

particular, the predator-induced plastic response (Bukowski & Auld, 2014). Thus, modulations of

general calcium pathways could influence the shell synthesis and be directly linked to anti-predator

defences like increase of shell thickness and change in shell morphology observed in this study.

Altogether, these results are consistent with previous studies investigating the functions of

differentially expressed genes in predator cue environments and showing that genes are associated

with defensive structures, metabolism and transcription regulation. Indeed, Hales et al. (2017) and

Rozenberg et al. (2015) showed on *D. ambigua* and *D. pulex* that the gene functions mostly explained

the observed defensive phenotypic changes (e.g. expression of proteins involved in formation of

protective structures and in cuticle strengthening) and were also involved in metabolism pathways as

for example metabolic proteins required for resource allocation or stress responses. Similar results

have been found in larvae of the damselfly *Enallagma cyathigerum* and in the marine copepod *Calanus*

finmarchicus where predation risk triggered different gene expression coding for stress proteins and

reduced antioxidant defence and for an alteration of lipid metabolism, respectively (Skottene et al.,

2020; Slos & Stoks, 2008). Although to relate the differentially gene expression profiles with phenotypic

traits is always largely speculative in non-model organisms, our study highlights some interesting

molecular pathways to study the underlying mechanisms of predator-induced defences in *P. acuta*.

Gene expression induced by trans-generational exposure to predator cues.

As observed in several previous studies on *P. acuta*, we confirmed the predator-induced

transgenerational plasticity as the parental exposure to predator cues significantly influenced

some offspring traits but to a lesser extent than WGP (Beaty et al., 2016; Tariel et al., 2020c). Indeed,

F2 snails from exposed parents were 5.5% more crawling-out the water and had a 16.9% thicker and

4.9% longer shell than snails from non-exposed parents while they did not differ in shell morphology.

At the molecular level, we have consistently detected different patterns of gene expression according

to the parental environments but based on a low number of differentially expressed genes. Indeed,

there were only 13 up-regulated and 1 down-regulated genes. Interestingly, our study showed that the

differentially expressed genes in response to parental predator-cue exposure (TGP) were not

differentially expressed for WGP in F1, suggesting that predator-induced TGP and WGP do not share

common genes. This is consistent with Hales et al. (2017) showing that the genes differentially

regulated for predator-induced WGP and TGP in *D. ambigua* were largely distinct. Stein et al. (2018)

also showed that there were unique sets of genes to the different forms of plasticity but also found a

common core set of genes in *G. aculeatus*. We consistently found that 6 differentially expressed genes

in F1 were more expressed in F2 than the average of all other genes in F2, suggesting that some genes

important for WGP may remain active in offspring but at a lower level than genes specific to TGP. In

particular, one differentially expressed gene in F1 and highly expressed in F2 is involved in the calcium

metabolism and potentially in shell construction (transcript XM_059293345.1). We also showed that

the differentially expressed genes in F2 are involved in the same functions that those in F1, i.e.

metabolism, including calcium metabolism and response to oxidative stress, and transcriptional

regulation. Consequently, these results suggest that common functions are used for predator-induced

WGP and TGP via both common and different genomic pathways. Thus, we can speculate as Stein et al.

(2018) that the shared genes are involved in the similar responses to personal and parental exposure

to predator cues (higher escape behaviour and thicker shell) while the specific genes reflect differences

in response to cues from different sources (growth difference: shorter shell in parents versus longer

shell in offspring). This would have important implications on the evolution of predator-induced

plasticity as it suggests that WGP and TGP can evolve independently for some specific responses, while

maintaining to some extent a common genetic basis.

A small number of genes can have an important impact on the phenotype (e.g. Meltzer, 2005),

especially when differentially expressed genes are involved in transcriptional regulation (e.g. Gill, 2003)

as observed for TGP in this study (e.g. cytosine-specific C5-methyltransferases, C2H2-type zinc-finger

family). However, the low number of differentially expressed genes evidenced for predator-induced

TGP may question whether our experiment has captured the entire transgenerational transcriptomic

response. Indeed, such a low number of differentially expressed genes in F2 generation strongly

contrasts with other studies on various factors (e.g. Clark et al., 2019; Ledón-Rettig, 2023) and on the

parental exposure to predator cues (e.g. Brass et al., 2020; Hales et al., 2017; Mommer & Bell, 2014;

Stein et al., 2018). For example, Stein et al., (2018) found 322 genes that were differentially regulated

in threespined stickleback *G. aculeatus* offspring when predation risk was experienced by the father. In

D. ambigua, individuals from predator-exposed parents and grand-parents had 233 and 170

differentially regulated genes respectively compared to individuals from non-exposed parents (Hales

et al., 2017). The quantitatively low molecular response induced by parental exposure to predator cues

observed in our study could result from a time lag between changes in gene expression and anti-

predator defence induction (Gilbert, 2001). Indeed, the environmental conditions experienced in early

life can have important consequences for phenotype later in life (Gilbert, 2001; Lee et al., 2013) but

also for offspring phenotype (Burton & Metcalfe, 2014; Tariel-Adam et al., 2023; Yin et al., 2019). In

this study, both differential gene expression and anti-predator defences in offspring have been

investigated at the adult stage while offspring may have integrated the environmental information from

their parents early in development engaging them in specific predation risk developmental trajectories.

As a consequence, the anti-predator defences observed in offspring from exposed parents may result

from gene expression changes early in development that no longer exist at the adult stage, which could

also explain why we did not find a clear common core set of genes between WGP and TGP. For example,

Mommer & Bell (2014) showed that maternal experience with predation risk influences genome-wide

embryonic expression in *G. aculeatus*, an early expression that may have phenotypic consequences

later (e.g. Sharda et al., 2021). Such a complex interplay between the timing of environmental cue

perception, gene expression and phenotypic consequences deserves to be explored in more details,

for example by tracking the differential gene expression throughout the development.

Conclusion

Both WGP and TGP are triggered by exposure to predator cues, involving changes in behaviour and

morphology associated with differential patterns of gene expression in F1 and F2 generations of *P.*

acuta. The differential profiles involved genes implicated in metabolism and transcriptional regulation.

In the F2 generation, the low number of differentially expressed genes suggests that either the gene

expression change induced by parental environment occurs early in development and decreases at

adult stage. Our study evidenced that unravelling the underlying molecular mechanisms of predator-

induced TGP requires further investigations and particularly focusing on the early developmental

period, a period during which differential gene expression induced by the environment may have a

delayed impact on the phenotype.

Acknowledgements

We are grateful to Cyril Degletagne who helped in experimental works and to the Rhône-Alpes

bioinformatic centre (PRABI) that has supported bioinformatics analyses. This work was financially

supported by the BioEnviS research federation (FR3728) of Lyon University and the TEATIME (ANR-21-

CE02-0005) grants from the French National Research Agency (ANR).

Authors Contributions

E.L., S.P. and T.L. conceived the project. J.T., S.P. and E.L. performed the experiments. L.K. performed

RNA extractions and libraries. L.D., J.T and M.H. analysed phenotypic data. L.D., N.S and T.L. performed

bioinformatic molecular analyses. L.D. wrote the first version of the manuscript. All authors revised the

manuscript.

Competing interests

No competing interests.

References

References

Agrawal, A. A., Laforsch, C., & Tollrian, R. (1999). Transgenerational induction of defences in animals
and plants. *Nature*, *401*(6748), Article 6748. https://doi.org/10.1038/43425

Aubin-Horth, N., & Renn, S. C. P. (2009). Genomic reaction norms: Using integrative biology to
understand molecular mechanisms of phenotypic plasticity. *Molecular Ecology*, *18*(18), 3763–3780.
https://doi.org/10.1111/j.1365-294X.2009.04313.x

Auld, J. R., & Relyea, R. A. (2011). Adaptive plasticity in predator-induced defenses in a common
freshwater snail: Altered selection and mode of predation due to prey phenotype. *Evolutionary*
Ecology, *25*(1), 189–202. https://doi.org/10.1007/s10682-010-9394-1

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). *Fitting Linear Mixed-Effects Models using lme4*
(arXiv:1406.5823). arXiv. https://doi.org/10.48550/arXiv.1406.5823

Beaty, L. E., Wormington, J. D., Kensinger, B. J., Bayley, K. N., Goeppner, S. R., Gustafson, K. D., &
Luttbeg, B. (2016). Shaped by the past, acting in the present: Transgenerational plasticity of anti-
predatory traits. *Oikos*, *125*(11), 1570–1576. https://doi.org/10.1111/oik.03114

Bell, A. M., & Hellmann, J. K. (2019). An Integrative Framework for Understanding the Mechanisms
and Multigenerational Consequences of Transgenerational Plasticity. *Annual Review of Ecology,*
Evolution, and Systematics, *50*(1), 97–118. https://doi.org/10.1146/annurev-ecolsys-110218-024613

Bell, A. M., & Stein, L. R. (2017). Transgenerational and developmental plasticity at the molecular
level: Lessons from Daphnia. *Molecular Ecology*, *26*(19), 4859–4861.
https://doi.org/10.1111/mec.14327

Blackwell, T. K., Sewell, A. K., Wu, Z., & Han, M. (2019). TOR Signaling in Caenorhabditis elegans
Development, Metabolism, and Aging. *Genetics*, *213*(2), 329–360.
https://doi.org/10.1534/genetics.119.302504

Brass, K. E., Herndon, N., Gardner, S. A., Grindstaff, J. L., & Campbell, P. (2020). Intergenerational
effects of paternal predator cue exposure on behavior, stress reactivity, and neural gene expression.
Hormones and Behavior, *124*, 104806. https://doi.org/10.1016/j.yhbeh.2020.104806

Braun, E. L., Kang, S., Nelson, M. A., & Natvig, D. O. (1998). Identification of the first fungal annexin:
Analysis of annexin gene duplications and implications for eukaryotic evolution. *Journal of Molecular*
Evolution, *47*(5), 531–543. https://doi.org/10.1007/pl00006409

Bray, N. L., Pimentel, H., Melsted, P., & Pachter, L. (2016). Near-optimal probabilistic RNA-seq
quantification. *Nature Biotechnology*, *34*(5), Article 5. https://doi.org/10.1038/nbt.3519

Bukowski, S. J., & Auld, J. R. (2014). The effects of calcium in mediating the inducible morphological
defenses of a freshwater snail, Physa acuta. *Aquatic Ecology*, *48*(1), 85–90.
https://doi.org/10.1007/s10452-013-9468-6

Burton, T., & Metcalfe, N. B. (2014). Can environmental conditions experienced in early life influence
future generations? *Proceedings of the Royal Society B: Biological Sciences*, *281*(1785), 20140311.
https://doi.org/10.1098/rspb.2014.0311

Clark, M. S., Suckling, C. C., Cavallo, A., Mackenzie, C. L., Thorne, M. A. S., Davies, A. J., & Peck, L. S.
(2019). Molecular mechanisms underpinning transgenerational plasticity in the green sea urchin

Psammechinus miliaris | Scientific Reports. *Scientific Reports*. https://www-nature-com.docelec.univ-lyon1.fr/articles/s41598-018-37255-6

Colicchio, J. M., & Herman, J. (2020). Empirical patterns of environmental variation favor adaptive transgenerational plasticity. *Ecology and Evolution*, *10*(3), 1648–1665. https://doi.org/10.1002/ece3.6022

Danchin, É., Charmantier, A., Champagne, F. A., Mesoudi, A., Pujol, B., & Blanchet, S. (2011). Beyond DNA: Integrating inclusive inheritance into an extended theory of evolution. *Nature Reviews Genetics*, *12*(7), Article 7. https://doi.org/10.1038/nrg3028

de Nadal, E., Ammerer, G., & Posas, F. (2011). Controlling gene expression in response to stress. *Nature Reviews Genetics*, *12*(12), 833–845. https://doi.org/10.1038/nrg3055

DeWitt, T. J. (1998). Costs and limits of phenotypic plasticity: Tests with predator-induced morphology and life history in a freshwater snail. *Journal of Evolutionary Biology*, *11*(4), 465–480. https://doi.org/10.1046/j.1420-9101.1998.11040465.x

DiRusso, C. C., Li, H., Darwis, D., Watkins, P. A., Berger, J., & Black, P. N. (2005). Comparative biochemical studies of the murine fatty acid transport proteins (FATP) expressed in yeast. *The Journal of Biological Chemistry*, *280*(17), 16829–16837. https://doi.org/10.1074/jbc.m409598200

Ezard, T. H. G., Prizak, R., & Hoyle, R. B. (2014). The fitness costs of adaptation via phenotypic plasticity and maternal effects. *Functional Ecology*, *28*(3), 693–701.

Fallet, M., Luquet, E., David, P., & Cosseau, C. (2020). Epigenetic inheritance and intergenerational effects in mollusks. *Gene*, *729*, 144166. https://doi.org/10.1016/j.gene.2019.144166

Galloway, L. F., & Etterson, J. R. (2007). Transgenerational Plasticity Is Adaptive in the Wild. *Science*, *318*(5853), 1134–1136. https://doi.org/10.1126/science.1148766

Gilbert, S. F. (2001). Ecological Developmental Biology: Developmental Biology Meets the Real World. *Developmental Biology*, *233*(1), 1–12. https://doi.org/10.1006/dbio.2001.0210

Gill, G. (2003). Post-translational modification by the small ubiquitin-related modifier SUMO has big effects on transcription factor activity. *Current Opinion in Genetics & Development*, *13*(2), 108–113. https://doi.org/10.1016/S0959-437X(03)00021-2

Gustafson, K., Kensinger, B., Bolek, M., & Luttbeg, B. (2014). Distinct snail (Physa) morphotypes from different habitats converge in shell shape and size under common garden conditions. *Evolutionary Ecology Research*, *16*, 77–89.

Hales, N. R., Schield, D. R., Andrew, A. L., Card, D. C., Walsh, M. R., & Castoe, T. A. (2017). Contrasting gene expression programs correspond with predator-induced phenotypic plasticity within and across generations in Daphnia. *Molecular Ecology*, *26*(19), 5003–5015. https://doi.org/10.1111/mec.14213

HELLER, J., & MAGARITZ, M. (1983). FROM WHERE DO LAND SNAILS OBTAIN THE CHEMICALS TO BUILD THEIR SHELLS? *Journal of Molluscan Studies*, *49*(2), 116–121. https://doi.org/10.1093/oxfordjournals.mollus.a065701

Herman, J., & Sultan, S. (2011). Adaptive Transgenerational Plasticity in Plants: Case Studies, Mechanisms, and Implications for Natural Populations. *Frontiers in Plant Science*, *2*. https://www.frontiersin.org/articles/10.3389/fpls.2011.00102

Jablonka, E., & Raz, G. (2009). Transgenerational epigenetic inheritance: Prevalence, mechanisms, and
implications for the study of heredity and evolution. *The Quarterly Review of Biology*, *84*(2), 131–176.
https://doi.org/10.1086/598822

Katewa, S. D., & Kapahi, P. (2011). Role of TOR signaling in aging and related biological processes in
Drosophila melanogaster. *Experimental Gerontology*, *46*(5), 382.
https://doi.org/10.1016/j.exger.2010.11.036

Khorchid, A., & Ikura, M. (2002). How calpain is activated by calcium. *Nature Structural Biology*, *9*(4),
239–241. https://doi.org/10.1038/nsb0402-239

Kikuchi, D. W., Allen, W. L., Arbuckle, K., Aubier, T. G., Briolat, E. S., Burdfield-Steel, E. R., Cheney, K. L.,
Daňková, K., Elias, M., Hämäläinen, L., Herberstein, M. E., Hossie, T. J., Joron, M., Kunte, K., Leavell, B.
C., Lindstedt, C., Lorioux-Chevalier, U., McClure, M., McLellan, C. F., … Exnerová, A. (2023). The
evolution and ecology of multiple antipredator defences. *Journal of Evolutionary Biology*, *36*(7), 975–
991. https://doi.org/10.1111/jeb.14192

Kim, S., Zhang, Z., Upchurch, S., Isern, N., & Chen, Y. (2004). Structure and DNA-binding Sites of the
SWI1 AT-rich Interaction Domain (ARID) Suggest Determinants for Sequence-specific DNA Recognition
*. *Journal of Biological Chemistry*, *279*(16), 16670–16676. https://doi.org/10.1074/jbc.M312115200

Kuijper, B., & Hoyle, R. B. (2015). When to rely on maternal effects and when on phenotypic
plasticity? *Evolution*, *69*(4), 950–968. https://doi.org/10.1111/evo.12635

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest Package: Tests in Linear Mixed
Effects Models. *Journal of Statistical Software*, *82*(13). https://doi.org/10.18637/jss.v082.i13

Ledón-Rettig, C. C. (2023). A transcriptomic investigation of heat-induced transgenerational plasticity
in beetles. *Biological Journal of the Linnean Society*, *138*(3), 318–327.
https://doi.org/10.1093/biolinnean/blac151

Lee, P. C., Bussière, L. F., Webber, C. E., Poole, J. H., & Moss, C. J. (2013). Enduring consequences of
early experiences: 40 year effects on survival and success among African elephants (Loxodonta
africana). *Biology Letters*, *9*(2), 20130011. https://doi.org/10.1098/rsbl.2013.0011

Loewith, R., Jacinto, E., Wullschleger, S., Lorberg, A., Crespo, J. L., Bonenfant, D., Oppliger, W., Jenoe,
P., & Hall, M. N. (2002). Two TOR complexes, only one of which is rapamycin sensitive, have distinct
roles in cell growth control. *Molecular Cell*, *10*(3), 457–468. https://doi.org/10.1016/s1097-
2765(02)00636-6

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550. https://doi.org/10.1186/s13059-014-0550-
8

Lu, Y., Wajapeyee, N., Turker, M. S., & Glazer, P. M. (2014). Silencing of the DNA Mismatch Repair
Gene MLH1 Induced by Hypoxic Stress in a Pathway Dependent on the Histone Demethylase LSD1.
Cell Reports, *8*(2), 501–513. https://doi.org/10.1016/j.celrep.2014.06.035

Luquet, E., Léna, J.-P., Miaud, C., & Plénet, S. (2015). Phenotypic divergence of the common toad
(Bufo bufo) along an altitudinal gradient: Evidence for local adaptation. *Heredity*, *114*(1), Article 1.
https://doi.org/10.1038/hdy.2014.71

Luquet, E., & Tariel, J. (2016). Offspring reaction norms shaped by parental environment: Interaction
between within- and trans-generational plasticity of inducible defenses. *BMC Evolutionary Biology*,
16(1), 209. https://doi.org/10.1186/s12862-016-0795-9

MacLeod, K. J., Monestier, C., Ferrari, M. C. O., McGhee, K. E., Sheriff, M. J., & Bell, A. M. (2022).
Predator-induced transgenerational plasticity in animals: A meta-analysis. *Oecologia*, *200*(3), 371–
383. https://doi.org/10.1007/s00442-022-05274-w

McKillop, W. B., & Harrison, A. D. (1972). Distribution of aquatic gastropods across an interface
between the Canadian Shield and limestone formations. *Canadian Journal of Zoology*, *50*(11), 1433–
1445. https://doi.org/10.1139/z72-192

Meltzer, P. S. (2005). Small RNAs with big impacts. *Nature*, *435*(7043), 745–746.
https://doi.org/10.1038/435745a

Mommer, B. C., & Bell, A. M. (2014). Maternal experience with predation risk influences genome-
wide embryonic gene expression in threespined sticklebacks (Gasterosteus aculeatus). *PloS One*, *9*(6),
e98564. https://doi.org/10.1371/journal.pone.0098564

O, H. (2001). PAST: Paleontological statistics software package for education and data analysis.
Palaeontol Electron, *4*, 9.

Orsini, L., Brown, J. B., Shams Solari, O., Li, D., He, S., Podicheti, R., Stoiber, M. H., Spanier, K. I.,
Gilbert, D., Jansen, M., Rusch, D. B., Pfrender, M. E., Colbourne, J. K., Frilander, M. J., Kvist, J.,
Decaestecker, E., De Schamphelaere, K. A. C., & De Meester, L. (2018). Early transcriptional response
pathways in Daphnia magna are coordinated in networks of crustacean-specific genes. *Molecular*
Ecology, *27*(4), 886–897. https://doi.org/10.1111/mec.14261

Paysan-Lafosse, T., Blum, M., Chuguransky, S., Grego, T., Pinto, B. L., Salazar, G. A., Bileschi, M. L., Bork,
P., Bridge, A., Colwell, L., Gough, J., Haft, D. H., Letunić, I., Marchler-Bauer, A., Mi, H., Natale, D. A.,
Orengo, C. A., Pandurangan, A. P., Rivoire, C., … Bateman, A. (2023). InterPro in 2022. *Nucleic Acids*
Research, *51*(D1), D418–D427. https://doi.org/10.1093/nar/gkac993

Pigliucci, M. (2005). Evolution of phenotypic plasticity: Where are we going now? *Trends in Ecology &*
Evolution, *20*(9), 481–486. https://doi.org/10.1016/j.tree.2005.06.001

Prizak, R., Ezard, T. H. G., & Hoyle, R. B. (2014). Fitness consequences of maternal and grandmaternal
effects. *Ecology and Evolution*, *4*(15), 3139–3145. https://doi.org/10.1002/ece3.1150

Reger, J., Lind, M. I., Robinson, M. R., & Beckerman, A. P. (2018). Predation drives local adaptation of
phenotypic plasticity. *Nature Ecology & Evolution*, *2*(1), 100–107. https://doi.org/10.1038/s41559-
017-0373-6

Rozenberg, A., Parida, M., Leese, F., Weiss, L. C., Tollrian, R., & Manak, J. R. (2015). Transcriptional
profiling of predator-induced phenotypic plasticity in Daphnia pulex. *Frontiers in Zoology*, *12*(1), 18.
https://doi.org/10.1186/s12983-015-0109-x

RStudio Team. (2020). *RStudio: Integrated Development Environment for R*.

Saxton, R. A., & Sabatini, D. M. (2017). mTOR Signaling in Growth, Metabolism, and Disease. *Cell*,
168(6), 960–976. https://doi.org/10.1016/j.cell.2017.02.004

Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ: 25 years of Image
Analysis. *Nature Methods*, *9*(7), 671–675.

Segers, F. H. I. D., & Taborsky, B. (2011). Juvenile exposure to predator cues induces a larger egg size in fish. *Proceedings of the Royal Society B: Biological Sciences*, *279*(1731), 1241–1248. https://doi.org/10.1098/rspb.2011.1290

Sharda, S., Zuest, T., Erb, M., & Taborsky, B. (2021). Predator-induced maternal effects determine adaptive antipredator behaviors via egg composition. *Proceedings of the National Academy of Sciences*, *118*(37), e2017063118. https://doi.org/10.1073/pnas.2017063118

Skottene, E., Tarrant, A. M., Altin, D., Olsen, R. E., Choquet, M., & Kvile, K. Ø. (2020). Lipid metabolism in Calanus finmarchicus is sensitive to variations in predation risk and food availability. *Scientific Reports*, *10*(1), 22322. https://doi.org/10.1038/s41598-020-79165-6

Slos, S., & Stoks, R. (2008). Predation risk induces stress proteins and reduces antioxidant defense. *Functional Ecology*, *22*(4), 637–642. https://doi.org/10.1111/j.1365-2435.2008.01424.x

Stein, L. R., Bukhari, S. A., & Bell, A. M. (2018). Personal and transgenerational cues are nonadditive at the phenotypic and molecular level. *Nature Ecology & Evolution*, *2*(8), 1306. https://doi.org/10.1038/s41559-018-0605-4

Tams, V., Nickel, J. H., Ehring, A., & Cordellier, M. (2020). Insights into the genetic basis of predator-induced response in Daphnia galeata. *Ecology and Evolution*, *10*(23), 13095–13108. https://doi.org/10.1002/ece3.6899

Tariel, J., Luquet, É., & Plénet, S. (2020). Interactions Between Maternal, Paternal, Developmental, and Immediate Environmental Effects on Anti-predator Behavior of the Snail Physa acuta. *Frontiers in Ecology and Evolution*, *8*. https://www.frontiersin.org/articles/10.3389/fevo.2020.591074

Tariel, J., Plénet, S., & Luquet, E. (2020a). How do developmental and parental exposures to predation affect personality and immediate behavioural plasticity in the snail *Physa acuta* ? *Proceedings of the Royal Society B: Biological Sciences*, *287*(1941), 20201761. https://doi.org/10.1098/rspb.2020.1761

Tariel, J., Plénet, S., & Luquet, É. (2020b). Transgenerational Plasticity in the Context of Predator-Prey Interactions. *Frontiers in Ecology and Evolution*, *8*. https://www.frontiersin.org/articles/10.3389/fevo.2020.548660

Tariel, J., Plénet, S., & Luquet, É. (2020c). Transgenerational plasticity of inducible defences: Combined effects of grand-parental, parental and current environments. *Ecology and Evolution*, *10*(5), 2367–2376. https://doi.org/10.1002/ece3.6046

Tariel-Adam, J., Luquet, É., & Plénet, S. (2023). Sensitive windows for within- and trans-generational plasticity of anti-predator defences. *Peer Community Journal*, *3*. https://doi.org/10.24072/pcjournal.304

Tollrian, R. (1995). Predator-Induced Morphological Defenses: Costs, Life History Shifts, and Maternal Effects in Daphnia Pulex. *Ecology*, *76*(6), 1691–1705. https://doi.org/10.2307/1940703

Turner, A. M., Fetterolf, S. A., & Bernot, R. J. (1999). Predator identity and consumer behavior: Differential effects of fish and crayfish on the habitat use of a freshwater snail. *Oecologia*, *118*(2), 242–247. https://doi.org/10.1007/s004420050724

Viney, M., & Diaz, A. (2012). Phenotypic plasticity in nematodes: Evolutionary and ecological significance. *Worm*, *1*(2), 98–106. https://doi.org/10.4161/worm.21086

Walker, J. E., Arizmendi, J. M., Dupuis, A., Fearnley, I. M., Finel, M., Medd, S. M., Pilkington, S. J.,

Runswick, M. J., & Skehel, J. M. (1992). Sequences of 20 subunits of NADH:ubiquinone

oxidoreductase from bovine heart mitochondria. Application of a novel strategy for sequencing

proteins using the polymerase chain reaction. *Journal of Molecular Biology*, *226*(4), 1051–1072.

https://doi.org/10.1016/0022-2836(92)91052-q

Walsh, M. R., Castoe, T., Holmes, J., Packer, M., Biles, K., Walsh, M., Munch, S. B., & Post, D. M. (2016).

Local adaptation in transgenerational responses to predators. *Proceedings of the Royal Society B:*

Biological Sciences, *283*(1823), 20152271. https://doi.org/10.1098/rspb.2015.2271

Westfall, P. J., Ballon, D. R., & Thorner, J. (2004). When the Stress of Your Environment Makes You Go

HOG Wild. *Science*, *306*(5701), 1511–1512. https://doi.org/10.1126/science.1104879

Wu, C.-W., & Storey, K. B. (2021). mTOR Signaling in Metabolic Stress Adaptation. *Biomolecules*, *11*(5).

https://doi.org/10.3390/biom11050681

Yin, J., Zhou, M., Lin, Z., Li, Q. Q., & Zhang, Y.-Y. (2019). Transgenerational effects benefit offspring

across diverse environments: A meta-analysis in plants and animals. *Ecology Letters*, *22*(11), 1976–

1986. https://doi.org/10.1111/ele.13373

727