Transcriptomic bases of within- and trans-generational predator-induced
plasticity in the freshwater snail *Physa acuta*

Authors:

Léo Dejeux[1], Nathanaëlle Saclier[2], Juliette Tariel-Adam[3], Maxime Hoareau[1],
Tristan Lefébure[1], Lara Konecny[1], Sandrine Plénet[1], Emilien Luquet[1, 4]

[1] Université Claude Bernard Lyon 1, LEHNA UMR 5023, CNRS, ENTPE, F-
69622, Villeurbanne, France

[2] ISEM, Université Montpellier, CNRS, IRD, Montpellier, France

[3] Fish Lab, Department of Biological Sciences, Macquarie University, Sydney,
NSW, Australia

[4] Institut universitaire de France (IUF)

Abstract: 223 words

Main body of text: 6040 words

Figures/tables: 8

References: 66

Abstract

Inducible defences in response to predation risk are a well-known example

of adaptive phenotypic plasticity. While inducible defences have been mainly

studied within a generation (within-generational plasticity), there is now clear

evidence that the ancestral exposure to predation risk can influence the

defences expressed by the offspring, even if they have not been exposed

themselves (transgenerational plasticity). The molecular mechanisms

allowing the transmission of environmental information across generations

are not well understood. In this study, we combined measures of anti-

predator responses (behavioural and morphological) with transcriptomic

investigations across two generations in the freshwater snail *Physa acuta*.

We hypothesised that both within- and transgenerational plasticity would

induce phenotypic changes associated with differential gene expression. Our

results confirmed within- and transgenerational plasticity: F1 snails have

responded to predator-cue exposure by increasing escape behaviour,

reducing shell length, and developing thicker and slender shells and F2 from

exposed parents have longer and thicker shells with a narrower aperture.

Within- and transgenerational plasticity were accompanied by the differential

expression of 112 genes (101 up- and 11 down-regulated) and 23

differentially expressed genes (17 up- and 6 down-regulated), respectively.

Within- and transgenerational plasticity did not share common differentially

expressed genes but the molecular functions were similar, involving

metabolism and transcription regulation. These results suggest that

predator-induced WGP and TGP may result from different genomic pathways

and may evolve independently.

INTRODUCTION

Phenotypic plasticity is the ability of a genotype to produce alternative

phenotypes under different environmental conditions (Pigliucci, 2005).

Plastic responses are widespread across the phylogenetic spectrum (*e.g.*

Galloway & Etterson, 2007; Luquet et al., 2011; Segers & Taborsky, 2011;

Walsh et al., 2016), and are a major concern in evolutionary biology as they

can be adaptive, i.e. increasing fitness of organisms facing environmental

changes (Pigliucci, 2005). Phenotypic plasticity classically refers to within-

generational plasticity (WGP), meaning that it occurs within one generation.

Recently, plasticity has also been found to occur across generations

(transgenerational plasticity, TGP) where the phenotype of an organism is

influenced by the environment experienced by the previous generations (Bell

& Hellmann, 2019). Interest in TGP has grown rapidly in recent years with

evidence that TGP is at least as widespread as WGP (Jablonka & Raz, 2009;

Tariel, Plénet, et al., 2020b). Although there are numerous examples of TGP

generating seemingly adaptive offspring phenotypes, its adaptive role is

expected to be more challenging than that of WGP due to patterns of

environmental variation across generations (Colicchio & Herman, 2020).

Indeed, it requires a strong temporal autocorrelation between environments

of distinct generations to be selected (Ezard et al., 2014; Kuijper & Hoyle,

2015; Prizak et al., 2014) and a transmission system that allows the induction

of adaptive phenotypes across generations (Fallet et al., 2020). The

inheritance mechanisms appear diverse, from epigenetic changes to

parental transmission of cytoplasmic elements (small RNAs, nutrients,

hormones, proteins), parental care and cultural transmission (Bell &

Hellmann, 2019; Danchin et al., 2011).

Regardless of the transmission mechanisms, these should lead in the

offspring to the modification of the expression of some of the genes involved

in the phenotypes, at least in some tissues and at some stages during the

development. Surprisingly, transgenerational plastic response at the

molecular level is overlooked. This is indeed a necessary first step to unravel

the molecular mechanisms of plasticity in general and their transgenerational

inheritance. The most compelling explorations of changes in gene

expression profiles in response to both developmental (WGP) and parental

(TGP) environments come from the studies of Hales et al. (2017) on *Daphnia*

ambigua and Stein et al. (2018) on three spined stickleback (*Gasterosteus*

aculeatus). These studies compared gene expression between individuals

that had been exposed to predator cues relative to non-exposed individuals

within a generation (WGP genes) and among their offspring (from non-

exposed and exposed parents; TGP genes). Stein et al. (2018) found that

the same group of genes changed their expression in offspring whether the

predation risk was experienced by the offspring (TGP), the father (WGP), or

both. This suggests that a core set of genes is consistently affected by

predation risk regardless of who experiences it. On the other hand, Hales et

al. (2017) found that changes in the expression of genes that respond to TGP

differed depending on whether exposure to predation was experienced by

parents or grandparents. They also showed that very few genes were

common to both WGP and TGP. Note that Clark et al. (2019) also showed

that TGP gene expression profiles of larvae of sea urchin (*Psammechinus*

miliaris) spawned at low pH from pre-acclimated adults differed from those

of larvae produced by adults exposed to ambient pH, but the authors did not

explore the WGP genes. These few results highlight the necessity to

investigate together WGP and TGP at the molecular level. Beyond the

inheritance mechanisms allowing organisms to memorise past

environmental conditions to adjust their future phenotype, we need to know

whether the two forms of plasticity are distinct at the molecular level and

potentially free to evolve independently of one another (Bell & Stein, 2017).

Predation risk is a major selection pressure for most species and results in

the evolution of defences in prey (Kikuchi et al., 2023). As predation risk often

varies over time and space, it favours the evolution of phenotypic plasticity

in defences (Reger et al., 2018; Tariel, Plénet, et al., 2020b; Viney & Diaz,

2012). Indeed, while some defences against predators are constitutive,

plastic defences are only induced when prey detect some predator cues in

their environment (*e.g*. visual cues, chemical cues such as predator odours,

or alarm signals released by conspecifics). These induced defences are

classic examples of adaptive WGP: prey can increase their fitness by

producing defences only when predators are present. The links between

prey-predator interactions and WGP have been studied for a long time (*e.g.*

Tariel et al., 2020b; Tollrian, 1995). Plastic defences can also be induced in

offspring after their parents or more distant ancestors have detected predator

cues. This transgenerational induction of defences was first highlighted in

Daphnia morphology by Agrawal et al. (1999) and has since been repeatedly

shown in multiple species and defences (review in MacLeod et al., 2022;

Tariel, Plénet, et al., 2020b).

Consequently, this work aims to unravel the molecular mechanisms

underlying the expression of WGP and TGP of defences in the context of

predator-prey interactions. We used *Physa acuta* (also referred to as

Physella acuta), a freshwater snail known for their within- and

transgenerational anti-predator defences such as escape behaviour (snails

crawl out of the water when they or their parents detect predator cues;

(Luquet & Tariel, 2016; Tariel, Luquet, et al., 2020; Tariel, Plénet, et al.,

2020a; Turner et al., 1999) change in shell size and shape, increase in shell

thickness and overall increase in shell-crush resistance (*e.g.* shell harder to

handle or to crush by predators; (Auld & Relyea, 2011; Beaty et al., 2016;

DeWitt, 1998; Tariel, Plénet, et al., 2020c; Tariel-Adam et al., 2023). We

combined a two-generation experiment to generate predator-induced WGP

and then TGP of defences (only in a non-exposed environment) with a

transcriptomic investigation. We used transcriptome sequencing and

assembly to specifically investigate the transcriptome mantle skirt gene

expression (mantle skirt is the tissue that synthesises the shell). We

expected that gene expression should be affected by current (WGP) and

parental (TGP) exposure to predator cues. More specifically, as WGP and

TGP induce similar phenotypic responses (*e.g.* increase in shell thickness,

we expected the two processes share the same core set of genes.

MATERIALS & METHODS

Animal collection, rearing conditions and experimental design

Adult *P. acuta* snails were collected from a wild population in a lentic backwater of the Rhône River (Lyon, France, 45° 48'6"N, 4° 55'33"E) in September 2018. Individuals from this population have been repeatedly used in previous studies on predator-induced plasticity (Luquet & Tariel, 2016) and are naturally exposed to predation by the crayfish *Faxonius limosus*. The experiment was conducted in our laboratory in a temperature-controlled chamber at 25 °C with a photoperiod of 12/12h. The wild snails constituting the F0 generation were pooled in a 10 L plastic box filled with dechlorinated tap water (control water hereafter) and interbreed overnight. They were then isolated in 80 mL plastic boxes (4.5 x 6 cm) filled with tap control (*i.e.*, without predator cues) water to lay eggs for 3 days. Eggs within boxes constitute F1 matrilines. Eggs hatched seven days later and then siblings grew together for a further ten days in water without predator cues. Then, 10 siblings of 20 F1 matrilines were randomly selected and split into two developmental environments: five siblings in water without predator cues (non-exposed treatment) and five siblings in water with predator cues (exposed treatment). F1 siblings individually developed for the next 31 days in 80 mL plastic boxes. On this date, one F1 snail per matriline and treatment was frozen at -80°C for molecular analysis and all remaining F1 snails were measured (see below).

To generate the F2 generation, we pooled F1 snails from the same treatment

in an aquarium filled with control water and let them copulate for 24h. As

previously, F1 snails were then isolated in 80 mL boxes and laid eggs for 3

days to produce the F2 matrilines. F2 snails from the two parental

environments (non-exposed and exposed to predator cues) were only reared

in control water in order to investigate the effect of parental environment

(TGP) alone, i.e. without the interaction between parental and developmental

environments. The same protocol steps as previously for F1 snails were

followed, except that eight siblings per matriline were reared and

measurements were made at an older age because of a slower growth. As

for F1, one F2 snail per matriline and treatment was frozen for molecular

analysis and the others were measured.

Snails have been fed *ad libitum* with boiled and mixed lettuce. Water

(according to treatment) and food was renewed twice a week. The water with

predator cues (i.e. exposed treatment) was the was water in which *F. limosus*

crayfish have been raised (one crayfish/5L), for 3 days and fed with snails.

Additionally, to ensure fresh alarm cues, we smashed several *P. acuta* adult

snails in the predator-cue water (one snail/5L) for 1h before using.

Behavioural and morphological responses

In this study, in order to check that exposure to predator cues triggered WGP

and TGP, escape behaviour and shell morphology were measured on F1

(non-exposed snails = 76, exposed snails = 73) and F2 (snails from non-

exposed parents = 89, snails from exposed parents = 101). A few mortalities

have led to uneven sample sizes but the mortality rate was consistent across

families (see Table S1). Crawling out of the water (position above the water

surface) is considered as allowing snails to escape from benthic predators

such as crayfishes (DeWitt, 1998; Tariel, Plénet, et al., 2020a). Shorter and

narrower shell and aperture dimensions and thicker shell are adaptive anti-

predator responses (Auld & Relyea, 2011). These phenotypic traits have

been recurrently involved in transgenerational responses of *P. acuta* to

predator cues (Beaty et al., 2016; Luquet & Tariel, 2016).

Escape behaviour - One week before measurements and four hours after the

water renewal, we recorded the position above or below the water surface of

each snail in their rearing boxes. Consequently, the behaviour of F1 snails

was recorded in water without or with predator cues according to their

respective treatments while that of F2 snails was only recorded in water

without predator cues.

Shell thickness, shell length and shape - Shell thickness was directly

measured with an electronic calliper at the nearest 0.01 mm at the edge of

the aperture. After, a photograph of each snail (aperture upwards) was taken

with an Olympus SC50 camera installed on an Olympus SZX9 binocular and

its Olympus DF PLAPO 1X-2 objective at a x8 magnification. Snail shell

length, shell width, aperture length and aperture width were measured using

ImageJ (Schneider et al., 2012). To estimate potential changes in shell shape

according to treatment, we used geometric methods (Gustafson et al., 2014).

On each snail shell, we digitised 19 landmarks (Fig. 1) using the software

ImageJ. Snails that lacked homologous landmarks because of broken shells

(typically broken apertures and apexes) or small size (lack of the third shell

spire) were removed from the analyses (non-exposed F1 snails = 76,

exposed F1 snails = 72, F2 snails from non-exposed parents = 89, F2 snails

from exposed parents = 99). To compare shell shapes according to

treatments, we scaled the shapes using the size metric defined below (Eqn.

1). Translational components were removed by translating all shapes so that

the centroids lie at the origin. Rotational components were removed by

rotating all shapes until the Procrustes distance was minimised with the first

shell fixed as reference.

$$s_k = \sqrt{\sum_{i=1}^{19}((x_{k,i} - \bar{x}_k)^2 + (y_{k,i} - \bar{y}_k{}^2)}$$

234   **Equation 1:** Procuste's equation used to scale the shapes using the size metric s_k is the size

235   of the k-th shell, $x_{k,i}$ and $y_{k,i}$ is the x and y component of the i-th landmark, and $(\bar{x}_k, \bar{y}_k)$, is the

236   centroid of the k-th shell.

Figure 1: Locations of the different landmarks (circle symbol) and pseudo-landmarks (positioned using a geometric method, square symbol) on *P. acuta* shell. Pseudo-landmarks have been created using a geometric method: first, two tangents (T1 and T2) have been drawn from landmark 1 (apex). T1 touches the curve of the habitation spire (5) and T2 touches the external edge of the aperture (10). From the two points of tangency (points 5 and 10), two perpendicular lines to the tangents have been traced (P1 and P2). The intersections with the columellar edge of the aperture, the external edge of the aperture and the curve of the habitation spire give additional pseudo-landmarks (6, 9, 18 and 19). The length of the shell L (straight line passing by 1 and 8) gives the pseudo-landmark 18 at the intersection with the columellar edge of the aperture).

Statistical analyses

The same statistical analyses were separately performed to test for WGP on

F1 snails and TGP on F2. The treatment factor (i.e. non-exposed or exposed

to predator cues) was the developmental environment for WGP (F1) while it

was the parental one for TGP (F2).

Escape behaviour - The effects of predator cues on escape behaviour (i.e.

snail position above/on or below the water surface) were analysed using

generalised linear mixed models (GLMM) assuming a binomial distribution

(logit link function). The treatment was a fixed effect and matriline was a

random intercept effect. We tested significance of fixed and random effects

with likelihood ratio tests.

Shell thickness, shell length - To analyse the effects of predator cues on shell

length and shell thickness, we performed linear mixed models with treatment

as fixed effect and matriline as a random intercept. We used restricted

maximum likelihood estimation and Kenward and Roger's approximation for

degrees of freedom. We tested significance of fixed effects with type II F-

tests (Kuznetsova et al., 2017) and significance of random effect with

likelihood ratio tests.

Linear mixed models and generalised linear mixed models were performed

using the lme4 package (Bates et al., 2014) on R software (version R-4.4.1

for Windows, (RStudio Team, 2020).

Shell shape - The effects of predator cues on shell shape were analysed

using principal component analyses (PCA) and were visualised with the 2D

landmarks relative warps analysis functionality. All standardised shapes

obtained by Procustes method were analysed using the PAST software with

a Student test (Hammer et al., 2001).

Molecular biology analysis

The mantle skirt was dissected for 5 non-exposed F1 snails and 5 exposed

F1 snails, and for 10 F2 snails from non-exposed parents and 10 F2 from

exposed parents. Total RNA of the mantle was extracted by adding TRI

Reagent (Molecular Research Center MRC, TR118) to the samples and by

homogenising tissues on ice with pistons and agitator (Motor Mixer Clearline

ref.045011 Dutscher). All remaining steps were carried out according to the

manufacturer's protocols (Molecular Research Center MRC, TR118). Then,

the RNA was treated with Turbo DNase enzyme (Turbo DNA free kit,

Invitrogen, AM1907) and assayed by fluorescence using a Qubit fluorometer

(Qubit® RNA HS Assay Kits Molecular Probes, Invitrogen, Q32855). Library

construction was carried out using the NEBNext Ultra II RNA Library Prep Kit

for Illumina (Biolabs New England, NEB E7770, E7490, E7600). Libraries

were randomly split in three groups and sequenced on three lanes with an

Illumina® HiSeq4000™ machine on the GenomEast platform hosted at the

IGBMC (GenomEast Platform – Institut de Génétique et de la Biologie

Moléculaire et Cellulaire IGBMC, Illkirch, France), resulting in 50 bp long

single-end reads.

306   Differential gene expression analysis

FASTQC (http://www.bioinformatics.babraham.ac.uk/projects/fastqc/)) was

used to assess the quality of the reads. We used a reference genome ORF

P. acuta available on NCBI (Isolate: Inbredline_101_S28. RefSeq:

GCF_028476545.12). Gene expression levels were assessed on annotated

transcripts using Kallisto (-l 50 -s 10 options; Bray et al., 2016). We

aggregated effective counts at the gene level and then applied a rounding to

the nearest whole number to convert non-integer values into integer counts.

We also applied a minimum count filter of 1 (cpm) where we removed all the

genes with reads below 1 (i.e. removing 5634 genes from the total of 46965).

The rounded effective counts was used as input for differential gene

expression using the DESeq2 package (Love et al., 2014), which performs

differential gene expression analysis using negative binomial distribution and

shrinkage estimation for dispersions ("apeglm" type shrinkage, Zhu et al.

2019), and outputs expression fold changes to improve the interpretability of

estimates. We specified non-exposed or exposed treatment in the model

(design formula: "design = ~treatment") within each generation

independently, meaning we compared F1 snails from non-exposed and

exposed treatments, and F2 snails from non-exposed and exposed parents.

Firstly, the Wald test (DESeq2 package) was used to test the significance of

differentially expressed genes (DEGs) in all samples between the non-

exposed and exposed treatments. Secondly, we checked for any structures

in the normalised count table by performing a principal component analysis

(PCA) in all samples on the 500 most DEGs from DESeq2 analysis (pval <

0.05) and based on Log2FoldChange (|LFC|, i.e. the logarithm base 2 of the

fold change in gene expression between two experimental conditions,

providing a measure of both the magnitude and direction of differential

expression). Thirdly, we filtered the significant DEGs by a |LFC| greater than

1 in each generation. Consequently, only differential expression with

an adjusted p-value < 0.05 and |LFC| > 1 was considered thereafter. Then,

we investigated whether the DEGs in F1 were also differentially expressed

in F2 by comparing specific gene IDs across the two generations datasets.

Finally, as the quality of annotations of the reference genome did not allow

complex functional analyses, we determined the potential gene functions and

the associated protein families of all significant DEGs by using the scan

procedure of InterPro, a tool provided by the European Bioinformatics

Institute from the European Molecular Biology Laboratory (EMBL-EBI; (Blum

et al., 2024; Jones et al., 2014). InterPro integrates signatures from 13

member databases to classify proteins into families, to predict functional

domains, and to identify key sites, enabling comprehensive functional

annotation.

RESULTS

Predator-induced behaviour and shell morphology

Within-generational Plasticity

In F1 generation, the proportion of crawling-out behaviour increased

significantly in snails exposed to predator-cues (+85.25%, F[1,19.43] =

321.82, p < 0.001; Fig. 2A). Shell length significantly decreased in snails

exposed to predator cues (-6.3%, F[1,128.71] = 15.46, p < 0.001; Fig. 2B).

Shell thickness significantly increased in exposed snails (+18.8%,

F[1,129.58] = 4.02, p = 0.047; Fig. 2C). The matriline random effect was

significant for shell length (χ²[1] = 19.022, p < 0.001) and non-significant for

behaviour and shell thickness (χ²[1] = 0.690, p = 0.406 and χ²[1] = 2.6116, p

= 0.106, respectively).

Figure 2: Effect of developmental exposure to predator cues (WGP) on crawling-out behaviour, shell

length and shell thickness in the F1 generation. A: Proportion of Crawling-out behaviour; B: Mean shell

length (mm); C: Mean shell thickness (mm). Error bars represent SE. Non-exposed and exposed to

predator-cue treatments are symbolised with blue circles and red triangles, respectively.

Predator-cues exposure induced changes in shell shape. The two first axes

of PCA analysis accounted for 59.2% of the shell shape variation (Fig. 3) and

PCA axes 1-4 (74.2% cumulative shape variation) were the only axes to each

account for more than 5% of shape variation. PC1 showed a change in the

columellar area (close to landmark 18-19) from shells with relatively wide

aperture (dilated columellar area; Fig. 3C) to shells with relatively narrow

aperture (compressed columellar area; Fig. 3D). PC2 showed a change in

the slenderness of the shell from shells with relatively short spires

(compressed apex; Fig. 3B) and relatively wide apertures (dilated aperture)

to shells with relatively short spires (dilated apex) and narrow aperture widths

(compressed aperture; Fig.3A).

Figure 3: Shape variation of non-exposed and exposed snails from F1 generation (blue circle and red

triangle symbols, respectively. Panels A, B, C, D represent deformation grids showing shell shape on the

extreme ends of the PCA axes. The 95% confidence ellipses are represented for non-exposed and

exposed to predator-cue treatments. The dilation-compression gradient bar shows the deformation of the

shell landmarks relative to the points of an average shell (which would be located at the centre of the

PCA).

Non-exposed snails showed the same range of deformation of columellar

area than exposed snails (PC2, t-test: t = 1.07, df = 144, p = 0.28). However,

shells of snails exposed to predator cues are slender, with longer spires and

narrow aperture widths than non-exposed ones (PC1, t-test: t = 4.803, df =

144, p < 0.001). This elongation is associated with the dilatation of the three

last spires and a narrow aperture at the base of the external edge. In contrast,

shells of non-exposed snails have a wider aperture at the base of the external

edge and smaller spires.

Transgenerational plasticity

In F2 generation, the proportion of crawling-out did not differ between snails

from non-exposed and exposed parents ($F_{[1,20.31]}$ = 3.929, p = 0.061; Fig.

4A). Snails from parents exposed to predator cues were significantly longer

and had a thicker shell than those from non-exposed parents (+4.9%,

$F_{[1,170.87]}$ = 7.55, p = 0.006 and +16.9%, $F_{[1,174.11]}$ = 9.858, p = 0.002,

respectively; Fig. 4B-C). The matriline random effect was significant for shell

length ($\chi^2_{[1]}$ = 31.685, p < 0.001) and non-significant for behaviour and shell

thickness ($\chi^2_{[1]}$ = 0.022, p = 0.882 and $\chi^2_{[1]}$ = 1.882, p = 0.17, respectively).

Figure 4: Effect of parental exposure (TGP) on crawling-out behaviour, shell length and shell thickness

in the F2 generation. A: Proportion of Crawling-out behaviour; B: Mean shell length (mm); C: Mean shell

thickness (mm). Error bars represent SE. Non-exposed and exposed to predator cues treatments are

symbolised with blue circles and red triangles, respectively.

Parental exposure to predator cues induced changes in shell shape. The two

first axes of PCA analysis accounted for 63.8% of the shell shape variation

and cumulative accounting for PCA axes 1-4 is 80.14% (Fig. 5). PC1

displayed a change in shell aperture width; the shells with high PC1 scores

(Fig. 5C) showed a dilatation of the shell where the outer lip meets the

parietal wall and a small compression of the columellar area (i.e. a wider

aperture). Shells with low PC1 scores (Fig. 5D) presented a narrower

aperture, a consequence of a dilated columellar area but compressed

parietal wall and basal lip in comparison to shells with high PC1 score. PC2

showed a change in shell aperture width but also in shell length, from shells

(Fig. 5C) with a dilated outer lip and a compression of the apex (which tends

to reduce spire length i.e. overall shell length) to shells (Fig. 5D) showing

elongated spires from dilated apex, with a narrower aperture due to outer lip

compression.

Figure 5: Shape variation of F2 snails from non-exposed and exposed parents (blue circle and red triangle symbols, respectively). Panels A, B, C, D represent deformation grids showing shell shape on the extreme ends of the PCA axes. The 95% confidence ellipses are represented for non-exposed and exposed to predator-cue treatments. The dilation-compression gradient bar shows the deformation of the shell landmarks relative to the points of an average shell (which would be located at the centre of the PCA).

Snails from predator-exposed parents had a dilatation of the shell in the columellar area accompanied by a small compression of the parietal wall (PC1: $t = 2.067$, $p = 0.04$). Conversely, there is no difference in deformation of the outer lip, nor dilatation or compression of the apex between snails from exposed and non-exposed parental environments (PC2: $t = 0.59$, $p = 0.55$).

Differential gene expression induced by predator-cues

Figure 6: PCA plot of the 30 transcriptome samples of *P. acuta* snails showing the divergence in differential expression among the first 500 most differentially expressed genes (selected by greatest |LFC|). Non-exposed and exposed to predator cue individuals are represented in blue and red, respectively. F1 individuals are represented by circles and plain ellipses and F2 individuals by triangles and dotted ellipses.

A principal component analysis (PCA) was performed to explore the clustering of read counts on the first 500 most DEGs. The first and second principal components (PC1 and 2) explained 18% and 11% of the variance respectively (Fig. 6, PCA loadings available in Fig. S1). The plot did not show clustering neither between the non-exposed and exposed treatments nor between the F1 and F2 generations (Fig. 6). In the F1 generation, the differential expression analysis revealed a total of 112 DEGs between non-exposed and exposed snails (101 up-regulated genes and 11 down-regulated gene; Fig. S2). In the F2 generation, the differential expression

analysis showed a total of 23 DEGs between snails from non-exposed and

exposed parents (17 up-regulated and 6 down-regulated gene; Fig. S2). The

comparison of DEGs in F1 and F2 revealed that they were not overlapped

(Fig. 7; FigS3).

Figure 7: Venn diagram (obtained with DESeq2 package) illustrating the number and overlap of

differentially expressed genes between F1 and F2 groups. Genes specific to each group are shown in

individual circles, while the intersection represents the shared genes.

The DEGs in F1 are mainly involved in metabolism, signal transduction,

extracellular matrix organization, protein and transcriptional regulations (Fig.

8, Tables S2 and S3 for further details). The DEGs in F2 are mainly involved

in metabolism and transcriptional regulation, including DNA methylation

regulation (Fig. 8, Tables S2 and S3 for further details). In both F1 and F2,

some DEGs were associated with calcium signalling (Fig. 8, Tables S2 and

S3 for further details) and a main proportion of DEGs was genes with

unknown functions.

Figure 8: Proportions of classes of functions of the differentially expressed genes in F1 (A) and in F2 (B)
based on the InterPro EMBL-EBI database (Jones et al., 2014; Blum et al., 2024).

DISCUSSION

This work aimed to unravel the molecular mechanisms underlying the

expression of within- and trans-generational plasticity of anti-predator

defences in *P. acuta*. We expected that gene expression should be affected

by developmental (WGP) and parental (TGP) exposure to predator cues, and

that both WGP and TGP share the same core set of genes. Exposure to

predator cues results in changes in escape behaviour, shell growth, shell

thickness and shell shape. The parental exposure to predator cues (TGP)

also influenced the offspring phenotype, in particular morphological traits as

previously shown in this system (*e.g.* Luquet & Tariel 2016). Accordingly,

there were DEGs between non-exposed and exposed snails in F1 (WGP)

and between F2 snails according to parental exposure to predator cues

(TGP). However, the number of DEGs detected for TGP were lower than for

WGP. Finally, our results showed that DEGs for WGP and TGP did not

overlap.

Phenotypic changes and gene expression induced by within-

generational exposure to predator cues

As expected, snails exposed to predator cues crawled more out of the water,

had thicker shells, were smaller and developed slender shells with longer

spires and narrower aperture width than non-exposed snails. These

behavioural and morphological responses represent defences and are

supposed to be adaptive as they are consistent with an escape behaviour

(Tariel, Luquet, et al., 2020) and an increase in shell crush resistance (Tariel,

Plénet, et al., 2020a, 2020c), increasing snail survival in face of crayfish

predation (Auld & Relyea, 2011). The decreased shell size observed in

exposed snails may result from a physiological stress caused by exposure

to predator cues, including costs to produce anti-predator defences (*e.g.*

crawling-out behaviour is at the expense of foraging; DeWitt, 1998).

These predator-induced WGP was associated with a distinct pattern of gene

expression with 112 DEGs (101 up-regulated and 11 down-regulated genes).

Overall, this pattern is consistent with several studies that explored the

changes in gene expression induced by the environment (*e.g.* Aubin-Horth

& Renn, 2009; Herman & Sultan, 2011). In the context of predator-prey

interaction, the main knowledge about the transcriptomic responses of preys

to predators comes from studies on Daphnia species. For example, Hales et

al. (2017) in *Daphnia ambigua* and Rozenberg et al. (2015) in *Daphnia pulex*

showed that anti-predator defences, mainly morphological traits, were

associated with 48 and 230 DEGs respectively. On the contrary, no DEGs

were identified in *Daphnia galeata,* a species known to adjust life history traits

(such as reproduction) but not morphological ones in response to predator

cues. Similarly, Orsini et al. (2018) revealed no changes in gene expression

associated with a short-term exposure to fish kairomones in *Daphnia magna*.

These contrasted results suggest that the patterns of gene expression may

depend on several factors such as the traits considered and the strength of

predator signals (*e.g.* cue concentration, duration of exposure).

The functional analysis of DEGs in snails exposed to predator cues

suggested that these genes would be mainly linked with metabolism

functions and transcription regulation, both processes known to be intimately

associated with stress responses (note however that 26% of DEGs had

unknown functions). A major part of genes would code for generic metabolic

processes (*e.g.* Complex I intermediate-associated protein 30 involved in

respiratory metabolism, Walker et al., 1992). These functions are essential

for cellular stability and adaptation to fluctuating conditions. However, some

genes would be involved in more stress-specific responses that might be

triggered by exposure to predators (*e.g.* a gene would be involved in the TOR

protein complex, well-known to influence response to abiotic stressors;

Blackwell et al., 2019; Katewa & Kapahi, 2011). In the same vein, some

DEGs were linked to the calcium ion metabolism (*e.g.* by coding for Annexin

and Peptidase C2 protein families; Khorchid & Ikura, 2002) and would be

involved in the shell synthesis. Calcium being the main component of shell

gastropods (Bukowski & Auld, 2014), the modulations of general calcium

pathways could influence the expression of anti-predator defences like

increase of shell thickness and change in shell shape observed in our study.

Our results are consistent with previous studies investigating the functions of

DEGs in predator cue environments and showing that genes are associated

with defensive structures, metabolism and transcription regulation. Indeed,

Hales et al. (2017) and Rozenberg et al. (2015) showed on *D. ambigua* and

D. pulex that the gene functions mostly explained the observed defensive

phenotypic changes (*e.g.* expression of proteins involved in formation of

protective structures and in cuticle strengthening) and were also involved in

metabolism pathways as for example metabolic proteins required for

resource allocation or stress responses. Similar results have been found in

larvae of the damselfly *Enallagma cyathigerum* and in the marine copepod

Calanus finmarchicus where predation risk triggered different gene expression coding for stress proteins and reduced antioxidant defence and for an alteration of lipid metabolism, respectively (Skottene et al., 2020; Slos & Stoks, 2008). Although to relate the functions of DEGs with phenotypic traits is always highly speculative in non-model organisms, our study may highlight some interesting molecular pathways underlying predator-induced WGP in *P. acuta* that would be interesting to study in further detail.

Phenotypic changes and gene expression induced by trans-generational exposure to predator cues.

Our results confirmed the predator-induced TGP. The parental exposure to predator cues significantly influenced the offspring phenotype, in particular morphological traits. F2 snails from exposed parents had thicker and longer shells with narrower apertures than snails from non-exposed parents while they had a similar anti-predator behaviour. These transgenerational effects are consistent with previous studies on this system demonstrating that parental environments shape anti-predator traits of offspring (Beaty et al., 2016; Tariel, Plénet, et al., 2020c). A thicker shell and a larger size increase the sell-crush resistance, and a narrower aperture limits the body snail extirpation by crayfish (Auld & Relyea, 2011; DeWitt, 1998). The longer shell of F2 snails from exposed parents was opposed to the shorter shell of exposed snails in F1. This opposite response is consistent with previous

studies (Tariel, Plénet, et al., 2020c) and suggest that snails from exposed

parents, by being informed of the risk at a very early stage, may use

developmental pathways allowing them to invest in both shell growth to reach

a refuge size (Auld & Houser, 2015) and shell thickness to increase crush-

resistance (Beaty et al., 2016; Tariel-Adam et al., 2023). The anti-predator

behaviour was similar between snails from non-exposed and exposed

parents, which is in line with some contrasted results observed in other

studies (Tariel, Plénet, et al., 2020a, 2020c). Overall, the expression of anti-

predator traits by offspring from exposed parents, even if they are not

themselves exposed to predator cues, may allow them to anticipate future

predation risk and to increase their survival (MacLeod et al., 2022; Tariel,

Plénet, et al., 2020b).

At the molecular level, the predator-induced TGP was associated with 23

DEGs (17 up-regulated and 6 down-regulated) according to the parental

environments. The low number of DEGs in F2 generation contrasts with other

studies on various factors (Clark et al., 2019; Ledón-Rettig, 2023) and on the

parental exposure to predator cues (Hales et al., 2017; Mommer & Bell,

2014; Stein et al., 2018). For example, Stein et al., (2018) found 322 DEGs

in threespined stickleback *G. aculeatus* offspring when predation risk was

experienced by the father. In *D. ambigua*, individuals from predator-exposed

parents and grand-parents had 233 and 170 DEGs respectively compared

to individuals from non-exposed parents (Hales et al., 2017). Although this

contrast may result from a less stringent method of DEGs selection (no LFC

filtering, *e.g*. Stein et al. 2018), it may also question whether our experiment

has captured the entire transgenerational transcriptomic response.

Firstly, gene expression is investigated in only one tissue (i.e. the mantle skirt

that synthesizes the shell) in this study while in many studies it was studied

in whole organs (*e.g*. Stein et al. 2018) or whole individuals (*e.g*. Hales et al.

2017). This may drastically decrease the number of DEGs detected,

capturing only a specific part of gene expression changes caused by parental

environments. However, this tissue-specific approach is particularly relevant

in systems like ours as it allows to specifically study gene expression in the

tissue at the origin of phenotypic traits (i.e. shell modifications in this study).

Secondly, this study only investigated the gene expression in offspring that

were not themselves exposed to predator cues, which did not allow to gain

information whether the gene expression associated with parental

environments is modulated by developmental one. In *P. acuta*, there is

evidence that parental and developmental environments interact to shape

the offspring phenotype: the resulting phenotypic patterns showed much

stronger TGP in offspring that had not experienced predator cues themselves

(WGP masking TGP for most traits; (Luquet & Tariel, 2016). Although it would

be relevant to provide an integrative view of gene expression patterns in all

environmental scenarios, our study of DEGs associated with TGP focused

on the environmental situation where TGP is the strongest.

Thirdly, the lower number of DEGs in F2 could also result from a time lag

between changes in gene expression and anti-predator defence induction

(Gilbert, 2001). Indeed, the environmental conditions experienced in early

life can have important consequences for phenotype later in life (Gilbert,

2001; Lee et al., 2013) but also for offspring phenotype (Burton & Metcalfe,

2014; Tariel-Adam et al., 2023; Yin et al., 2019). For example, maternal

provisioning plays a crucial role in transmitting environmental information to

offspring by influencing cytoplasmic components, including mRNAs and

proteins, during early embryogenesis (Burton & Metcalfe, 2014). This

process can impact gene regulation during the maternal-to-zygotic transition,

potentially shaping phenotypic responses before they are detectable in

adults. In this study, both differential gene expression and anti-predator

defences in offspring have been investigated at the adult stage while

offspring may have integrated the environmental information from their

parents early in development engaging them in specific predation risk

developmental trajectories. As a consequence, the anti-predator defences

observed in offspring from exposed parents may result from gene expression

changes early in development that no longer exist at the adult stage, which

could also explain why we did not find a clear common core set of genes

between WGP and TGP. For example, Mommer & Bell (2014) showed that

maternal experience with predation risk influences genome-wide embryonic

expression in *G. aculeatus*, an early expression that may have phenotypic

consequences later (Sharda et al., 2021). Such a complex interplay between

the timing of environmental cue perception, gene expression and phenotypic

consequences deserves to be explored in more details, for example by

tracking the DEGs throughout the development.

The functional analysis of DEGs between snails from non-exposed parents

and those from exposed parents suggested the same main functions as for

WGP, i.e. metabolism and transcription regulation. Genes linked with

metabolisms would be mainly involved in response to oxidative stress and

carbohydrate metabolism. One gene would be involved in the regulation of

cytosolic calcium ion concentration, which might influence the shell synthesis

as previously mentioned for WGP. Most genes linked with transcription

regulation in F2 would be involved in methyltransferase activity, suggesting

that DNA methylations, a well-known bearer of epigenetic information, might

be involved in predator-induced TGP (Fallet et al., 2020). Epigenetic

mechanisms, such as DNA methylation and histone modifications, are known

to contribute to transgenerational inheritance of phenotypic traits (Duncan et

al., 2014; Fallet et al., 2020). Recent studies on the freshwater gastropod

Potamopyrgus antipodarum have demonstrated that differential DNA

methylation patterns were associated with heritable morphological

differentiations among genetically uniform populations (Smithson et al.,

2020; Thorson et al., 2017, 2019). These results suggest that

transgenerational inheritance of anti-predator traits in *P. acuta* might be

mediated by bearers of epigenetic information and call for more investigation

of epigenetic inheritance in the context of predator-induced plasticity.

Comparison of differentially expressed genes associated with WGP and TGP

The number of DEGs associated with TGP was lower than those for WGP. As mentioned above, this may occur because we only investigated DEGs at the adult stage or because F2 snails did not experiment direct exposure to predator cues. Surprisingly, the DEGs associated with WGP and TGP were not overlapped, suggesting that predator-induced TGP and WGP do not share common genes. This is consistent with Hales et al. (2017) showing that the DEGs for predator-induced WGP and TGP in *D. ambigua* were largely distinct. On the contrary, Stein et al. (2018) also showed that there were unique sets of genes to the different forms of plasticity but also found a common core set of genes in *G. aculeatus*. Despite the absence of common genes in our study, the DEGs in F2 seem involved in the same main functions that those in F1, i.e. metabolism and transcriptional regulation. Consequently, these results could suggest that common functions are used for predator-induced WGP and TGP via different genomic pathways. This would have important implications on the evolution of predator-induced plasticity as it suggests that WGP and TGP can evolve independently.

Conclusion

This study confirmed the predator-induced WGP and TGP in *P. acuta*, showing that anti-predator traits (escape behaviour, shell thickness, shell length and shell shape) are induced when snails were exposed during their

own development or when they originated from parents exposed to predator

cues (but not exposed themselves). These within- and transgenerational

phenotypic responses were both associated with differential patterns of gene

expression. However, the transcriptomic signal was lower for TGP (a lower

number of DEGs) and surprisingly no common genes were shared between

WGP and TGP although the main functions would be similar (metabolism

and transcription regulation). Expression analysis showed that only a small

set of genes (112 in F1 and 23 in F2) had a great response with high

differential expression levels in both generations. Different hypotheses have

been proposed to explain the different transcriptomic patterns observed

between predator-induced WGP and TGP and called for further

investigations allowing to address the limitations of this study in order to track

the transcriptomic signal throughout the development, across different

tissues and environmental scenarios. This will provide a more integrative

view of underlying molecular mechanisms of WGP and TGP addressing an

important gap of knowledge on important processes for adaptation to

changing environments.

Acknowledgements. We are grateful to Cyril Degletagne who helped in
experimental works and to the Rhône-Alpes bioinformatic centre (PRABI)
that has supported bioinformatics analyses. This work was financially
supported by the BioEnviS research federation (FR3728) of Lyon University
and the TEATIME (ANR-21-CE02-0005) grants from the French National
Research Agency (ANR).

Authors Contributions. E.L., S.P. and T.L. conceived the project. J.T., S.P.
and E.L. performed the experiments. L.K. performed RNA extractions and
libraries. L.D., J.T. and M.H. analysed phenotypic data. L.D., N.S and T.L.
performed bioinformatic molecular analyses. L.D. wrote the first version of
the manuscript. All authors revised the manuscript.

Ethics. This work did not require ethical approval from a human subject or
animal welfare committee.

Conflict of interest declaration. We declare we have no competing
interests.

Funding. This work was financially supported by the TEATIME (ANR-21-
CE02-0005) grants from the French National Research Agency (ANR).

Data accessibility. RNA-seq data have been deposited in the European
Nucleotide Archive (ENA) under the accession number [PRJEB87216]. The
raw sequencing data are publicly available at [
https://www.ebi.ac.uk/ena/browser/view/PRJEB87216]. Additionally, scripts
used for the analysis, along with the non-genomic data, have been made

available on Zenodo. The repository can be accessed at [

<u>10.5281/zenodo.15043538</u>].

References

Agrawal, A. A., Laforsch, C., & Tollrian, R. (1999). Transgenerational induction

of defences in animals and plants. *Nature*, *401*(6748), Article 6748.

https://doi.org/10.1038/43425

Aubin-Horth, N., & Renn, S. C. P. (2009). Genomic reaction norms: Using

integrative biology to understand molecular mechanisms of

phenotypic plasticity. *Molecular Ecology*, *18*(18), 3763–3780.

https://doi.org/10.1111/j.1365-294X.2009.04313.x

Auld, J. R., & Houser, R. (2015). Age-dependent effects of predation risk on

reproductive success in a freshwater snail. *Evolution*, *69*(10), 2793–

2798. https://doi.org/10.1111/evo.12769

Auld, J. R., & Relyea, R. A. (2011). Adaptive plasticity in predator-induced

defenses in a common freshwater snail: Altered selection and mode of

predation due to prey phenotype. *Evolutionary Ecology*, *25*(1), 189–

202. https://doi.org/10.1007/s10682-010-9394-1

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). *Fitting Linear Mixed-*

Effects Models using lme4 (arXiv:1406.5823). arXiv.

https://doi.org/10.48550/arXiv.1406.5823

Beaty, L. E., Wormington, J. D., Kensinger, B. J., Bayley, K. N., Goeppner, S. R.,

Gustafson, K. D., & Luttbeg, B. (2016). Shaped by the past, acting in

the present: Transgenerational plasticity of anti-predatory traits. *Oikos*,

125(11), 1570–1576. https://doi.org/10.1111/oik.03114

Bell, A. M., & Hellmann, J. K. (2019). An Integrative Framework for

Understanding the Mechanisms and Multigenerational Consequences

of Transgenerational Plasticity. *Annual Review of Ecology, Evolution,*

and Systematics, *50*(1), 97–118. https://doi.org/10.1146/annurev-

ecolsys-110218-024613

Bell, A. M., & Stein, L. R. (2017). Transgenerational and developmental

plasticity at the molecular level: Lessons from Daphnia. *Molecular*

Ecology, *26*(19), 4859–4861. https://doi.org/10.1111/mec.14327

Blackwell, T. K., Sewell, A. K., Wu, Z., & Han, M. (2019). TOR Signaling in

Caenorhabditis elegans Development, Metabolism, and Aging.

Genetics, *213*(2), 329–360.

https://doi.org/10.1534/genetics.119.302504

Blum, M., Andreeva, A., Florentino, L. C., Chuguransky, S. R., Grego, T., Hobbs,

E., Pinto, B. L., Orr, A., Paysan-Lafosse, T., Ponamareva, I., Salazar, G.

A., Bordin, N., Bork, P., Bridge, A., Colwell, L., Gough, J., Haft, D. H.,

Letunic, I., Llinares-López, F., ... Bateman, A. (2024). InterPro: The

protein sequence classification resource in 2025. *Nucleic Acids*

Research, *53*(D1), D444–D456. https://doi.org/10.1093/nar/gkae1082

Bray, N. L., Pimentel, H., Melsted, P., & Pachter, L. (2016). Near-optimal

probabilistic RNA-seq quantification. *Nature Biotechnology*, *34*(5),

Article 5. https://doi.org/10.1038/nbt.3519

Bukowski, S. J., & Auld, J. R. (2014). The effects of calcium in mediating the

inducible morphological defenses of a freshwater snail, Physa acuta.

Aquatic Ecology, *48*(1), 85–90. https://doi.org/10.1007/s10452-013-

9468-6

Burton, T., & Metcalfe, N. B. (2014). Can environmental conditions

experienced in early life influence future generations? *Proceedings of*

the Royal Society B: Biological Sciences, *281*(1785), 20140311.

https://doi.org/10.1098/rspb.2014.0311

Clark, M. S., Suckling, C. C., Cavallo, A., Mackenzie, C. L., Thorne, M. A. S.,

Davies, A. J., & Peck, L. S. (2019). Molecular mechanisms

underpinning transgenerational plasticity in the green sea urchin

Psammechinus miliaris | Scientific Reports. *Scientific Reports*.

https://www-nature-com.docelec.univ-lyon1.fr/articles/s41598-018-

37255-6

Colicchio, J. M., & Herman, J. (2020). Empirical patterns of environmental

variation favor adaptive transgenerational plasticity. *Ecology and*

Evolution, *10*(3), 1648–1665. https://doi.org/10.1002/ece3.6022

Danchin, É., Charmantier, A., Champagne, F. A., Mesoudi, A., Pujol, B., &

Blanchet, S. (2011). Beyond DNA: Integrating inclusive inheritance into

an extended theory of evolution. *Nature Reviews Genetics*, *12*(7),

Article 7. https://doi.org/10.1038/nrg3028

DeWitt, T. J. (1998). Costs and limits of phenotypic plasticity: Tests with

predator-induced morphology and life history in a freshwater snail.

Journal of Evolutionary Biology, *11*(4), 465–480.

https://doi.org/10.1046/j.1420-9101.1998.11040465.x

Duncan, E. J., Gluckman, P. D., & Dearden, P. K. (2014). Epigenetics, plasticity,

and evolution: How do we link epigenetic change to phenotype?

Journal of Experimental Zoology Part B: Molecular and Developmental

Evolution, *322*(4), 208–220. https://doi.org/10.1002/jez.b.22571

Ezard, T. H. G., Prizak, R., & Hoyle, R. B. (2014). The fitness costs of adaptation

via phenotypic plasticity and maternal effects. *Functional Ecology*,

28(3), 693–701.

Fallet, M., Luquet, E., David, P., & Cosseau, C. (2020). Epigenetic inheritance

and intergenerational effects in mollusks. *Gene*, *729*, 144166.

https://doi.org/10.1016/j.gene.2019.144166

Galloway, L. F., & Etterson, J. R. (2007). Transgenerational Plasticity Is Adaptive

in the Wild. *Science*, *318*(5853), 1134–1136.

https://doi.org/10.1126/science.1148766

Gilbert, S. F. (2001). Ecological Developmental Biology: Developmental

Biology Meets the Real World. *Developmental Biology*, *233*(1), 1–12.

https://doi.org/10.1006/dbio.2001.0210

Gustafson, K., Kensinger, B., Bolek, M., & Luttbeg, B. (2014). Distinct snail

(Physa) morphotypes from different habitats converge in shell shape

and size under common garden conditions. *Evolutionary Ecology*

Research, *16*, 77–89.

Hales, N. R., Schield, D. R., Andrew, A. L., Card, D. C., Walsh, M. R., & Castoe,

T. A. (2017). Contrasting gene expression programs correspond with

predator-induced phenotypic plasticity within and across generations

in Daphnia. *Molecular Ecology*, *26*(19), 5003–5015.

https://doi.org/10.1111/mec.14213

Hammer, O., Harper, D. A. T., & Ryan, P. D. (2001). PAST: Paleontological

Statistics Software Package for Education and Data Analysis.

Palaeontologia Electronica.

Herman, J., & Sultan, S. (2011). Adaptive Transgenerational Plasticity in Plants:

Case Studies, Mechanisms, and Implications for Natural Populations.

Frontiers in Plant Science, *2*.

https://www.frontiersin.org/articles/10.3389/fpls.2011.00102

Jablonka, E., & Raz, G. (2009). Transgenerational epigenetic inheritance:

Prevalence, mechanisms, and implications for the study of heredity

and evolution. *The Quarterly Review of Biology*, *84*(2), 131–176.

https://doi.org/10.1086/598822

Jones, P., Binns, D., Chang, H.-Y., Fraser, M., Li, W., McAnulla, C., McWilliam,

H., Maslen, J., Mitchell, A., Nuka, G., Pesseat, S., Quinn, A. F.,

Sangrador-Vegas, A., Scheremetjew, M., Yong, S.-Y., Lopez, R., &

Hunter, S. (2014). InterProScan 5: Genome-scale protein function

classification. *Bioinformatics*, *30*(9), 1236–1240.

https://doi.org/10.1093/bioinformatics/btu031

Katewa, S. D., & Kapahi, P. (2011). Role of TOR signaling in aging and related

biological processes in Drosophila melanogaster. *Experimental*

Gerontology, *46*(5), 382. https://doi.org/10.1016/j.exger.2010.11.036

Khorchid, A., & Ikura, M. (2002). How calpain is activated by calcium. *Nature*

Structural Biology, *9*(4), 239–241. https://doi.org/10.1038/nsb0402-

239

Kikuchi, D. W., Allen, W. L., Arbuckle, K., Aubier, T. G., Briolat, E. S., Burdfield-

Steel, E. R., Cheney, K. L., Daňková, K., Elias, M., Hämäläinen, L.,

Herberstein, M. E., Hossie, T. J., Joron, M., Kunte, K., Leavell, B. C.,

Lindstedt, C., Lorioux-Chevalier, U., McClure, M., McLellan, C. F., …

Exnerová, A. (2023). The evolution and ecology of multiple

antipredator defences. *Journal of Evolutionary Biology*, *36*(7), 975–991.

https://doi.org/10.1111/jeb.14192

Kuijper, B., & Hoyle, R. B. (2015). When to rely on maternal effects and when

on phenotypic plasticity? *Evolution*, *69*(4), 950–968.

https://doi.org/10.1111/evo.12635

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). lmerTest

Package: Tests in Linear Mixed Effects Models. *Journal of Statistical*

Software, *82*(13). https://doi.org/10.18637/jss.v082.i13

Ledón-Rettig, C. C. (2023). A transcriptomic investigation of heat-induced

transgenerational plasticity in beetles. *Biological Journal of the*

Linnean Society, *138*(3), 318–327.

https://doi.org/10.1093/biolinnean/blac151

Lee, P. C., Bussière, L. F., Webber, C. E., Poole, J. H., & Moss, C. J. (2013).

Enduring consequences of early experiences: 40 year effects on

survival and success among African elephants (Loxodonta africana).

Biology Letters, *9*(2), 20130011.

https://doi.org/10.1098/rsbl.2013.0011

Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold

change and dispersion for RNA-seq data with DESeq2. *Genome*

Biology, *15*(12), 550. https://doi.org/10.1186/s13059-014-0550-8

Luquet, E., Léna, J.-P., David, P., Joly, P., Lengagne, T., Perrin, N., & Plénet, S.

(2011). Consequences of genetic erosion on fitness and phenotypic

plasticity in European tree frog populations (Hyla arborea). *Journal of*

Evolutionary Biology, *24*(1), 99–110. https://doi.org/10.1111/j.1420-

9101.2010.02138.x

Luquet, E., & Tariel, J. (2016). Offspring reaction norms shaped by parental

environment: Interaction between within- and trans-generational

plasticity of inducible defenses. *BMC Evolutionary Biology*, *16*(1),

Article 1. https://doi.org/10.1186/s12862-016-0795-9

MacLeod, K. J., Monestier, C., Ferrari, M. C. O., McGhee, K. E., Sheriff, M. J., &

Bell, A. M. (2022). Predator-induced transgenerational plasticity in

animals: A meta-analysis. *Oecologia*, *200*(3), 371–383.

https://doi.org/10.1007/s00442-022-05274-w

Mommer, B. C., & Bell, A. M. (2014). Maternal experience with predation risk

influences genome-wide embryonic gene expression in threespined

sticklebacks (Gasterosteus aculeatus). *PloS One*, *9*(6), e98564.

https://doi.org/10.1371/journal.pone.0098564

Orsini, L., Brown, J. B., Shams Solari, O., Li, D., He, S., Podicheti, R., Stoiber,

M. H., Spanier, K. I., Gilbert, D., Jansen, M., Rusch, D. B., Pfrender, M.

E., Colbourne, J. K., Frilander, M. J., Kvist, J., Decaestecker, E., De

Schamphelaere, K. A. C., & De Meester, L. (2018). Early transcriptional

response pathways in Daphnia magna are coordinated in networks of

crustacean-specific genes. *Molecular Ecology*, *27*(4), 886–897.

https://doi.org/10.1111/mec.14261

Pigliucci, M. (2005). Evolution of phenotypic plasticity: Where are we going

now? *Trends in Ecology & Evolution*, *20*(9), 481–486.

https://doi.org/10.1016/j.tree.2005.06.001

Prizak, R., Ezard, T. H. G., & Hoyle, R. B. (2014). Fitness consequences of

maternal and grandmaternal effects. *Ecology and Evolution*, *4*(15),

3139–3145. https://doi.org/10.1002/ece3.1150

Reger, J., Lind, M. I., Robinson, M. R., & Beckerman, A. P. (2018). Predation

drives local adaptation of phenotypic plasticity. *Nature Ecology &*

Evolution, *2*(1), 100–107. https://doi.org/10.1038/s41559-017-0373-6

Rozenberg, A., Parida, M., Leese, F., Weiss, L. C., Tollrian, R., & Manak, J. R.

915   (2015). Transcriptional profiling of predator-induced phenotypic

916   plasticity in Daphnia pulex. *Frontiers in Zoology*, *12*(1), 18.

917   https://doi.org/10.1186/s12983-015-0109-x

918 RStudio Team. (2020). *RStudio: Integrated Development Environment for R*.

919 Schneider, C. A., Rasband, W. S., & Eliceiri, K. W. (2012). NIH Image to ImageJ:

920   25 years of Image Analysis. *Nature Methods*, *9*(7), 671–675.

921 Segers, F. H. I. D., & Taborsky, B. (2011). Juvenile exposure to predator cues

922   induces a larger egg size in fish. *Proceedings of the Royal Society B:*

923   *Biological Sciences*, *279*(1731), 1241–1248.

924   https://doi.org/10.1098/rspb.2011.1290

925 Sharda, S., Zuest, T., Erb, M., & Taborsky, B. (2021). Predator-induced maternal

926   effects determine adaptive antipredator behaviors via egg

927   composition. *Proceedings of the National Academy of Sciences*,

928   *118*(37), e2017063118. https://doi.org/10.1073/pnas.2017063118

929 Skottene, E., Tarrant, A. M., Altin, D., Olsen, R. E., Choquet, M., & Kvile, K. Ø.

930   (2020). Lipid metabolism in Calanus finmarchicus is sensitive to

931   variations in predation risk and food availability. *Scientific Reports*,

932   *10*(1), 22322. https://doi.org/10.1038/s41598-020-79165-6

933 Slos, S., & Stoks, R. (2008). Predation risk induces stress proteins and reduces

934   antioxidant defense. *Functional Ecology*, *22*(4), 637–642.

935   https://doi.org/10.1111/j.1365-2435.2008.01424.x

Smithson, M., Thorson, J. L. M., Sadler-Riggleman, I., Beck, D., Skinner, M. K.,

& Dybdahl, M. (2020). Between-Generation Phenotypic and Epigenetic

Stability in a Clonal Snail. *Genome Biology and Evolution*, *12*(9), 1604–

1615. https://doi.org/10.1093/gbe/evaa181

Stein, L. R., Bukhari, S. A., & Bell, A. M. (2018). Personal and transgenerational

cues are nonadditive at the phenotypic and molecular level. *Nature*

Ecology & Evolution, *2*(8), 1306–1311. https://doi.org/10.1038/s41559-

018-0605-4

Tariel, J., Luquet, É., & Plénet, S. (2020). Interactions Between Maternal,

Paternal, Developmental, and Immediate Environmental Effects on

Anti-predator Behavior of the Snail Physa acuta. *Frontiers in Ecology*

and Evolution, *8*.

https://www.frontiersin.org/articles/10.3389/fevo.2020.591074

Tariel, J., Plénet, S., & Luquet, E. (2020a). How do developmental and parental

exposures to predation affect personality and immediate behavioural

plasticity in the snail Physa acuta? *Proceedings of the Royal Society B:*

Biological Sciences, *287*(1941), 20201761.

https://doi.org/10.1098/rspb.2020.1761

Tariel, J., Plénet, S., & Luquet, É. (2020b). Transgenerational Plasticity in the

Context of Predator-Prey Interactions. *Frontiers in Ecology and*

Evolution, *8*. https://doi.org/10.3389/fevo.2020.548660

Tariel, J., Plénet, S., & Luquet, É. (2020c). Transgenerational plasticity of

inducible defences: Combined effects of grand-parental, parental and

current environments. *Ecology and Evolution*, *10*(5), 2367–2376.

https://doi.org/10.1002/ece3.6046

Tariel-Adam, J., Luquet, É., & Plénet, S. (2023). Sensitive windows for within-

and trans-generational plasticity of anti-predator defences. *Peer*

Community Journal, *3*. https://doi.org/10.24072/pcjournal.304

Thorson, J. L. M., Smithson, M., Beck, D., Sadler-Riggleman, I., Nilsson, E.,

Dybdahl, M., & Skinner, M. K. (2017). Epigenetics and adaptive

phenotypic variation between habitats in an asexual snail. *Scientific*

Reports, *7*(1), 14139. https://doi.org/10.1038/s41598-017-14673-6

Thorson, J. L. M., Smithson, M., Sadler-Riggleman, I., Beck, D., Dybdahl, M., &

Skinner, M. K. (2019). Regional epigenetic variation in asexual snail

populations among urban and rural lakes. *Environmental Epigenetics*,

5(4), dvz020. https://doi.org/10.1093/eep/dvz020

Tollrian, R. (1995). Predator-Induced Morphological Defenses: Costs, Life

History Shifts, and Maternal Effects in Daphnia Pulex. *Ecology*, *76*(6),

1691–1705. https://doi.org/10.2307/1940703

Turner, A. M., Fetterolf, S. A., & Bernot, R. J. (1999). Predator identity and

consumer behavior: Differential effects of fish and crayfish on the

habitat use of a freshwater snail. *Oecologia*, *118*(2), 242–247.

https://doi.org/10.1007/s004420050724

Viney, M., & Diaz, A. (2012). Phenotypic plasticity in nematodes: Evolutionary

and ecological significance. *Worm*, *1*(2), 98–106.

https://doi.org/10.4161/worm.21086

Walker, J. E., Arizmendi, J. M., Dupuis, A., Fearnley, I. M., Finel, M., Medd, S.

M., Pilkington, S. J., Runswick, M. J., & Skehel, J. M. (1992). Sequences

of 20 subunits of NADH:ubiquinone oxidoreductase from bovine heart

mitochondria. Application of a novel strategy for sequencing proteins

using the polymerase chain reaction. *Journal of Molecular Biology*,

226(4), 1051–1072. https://doi.org/10.1016/0022-2836(92)91052-q

Walsh, M. R., Castoe, T., Holmes, J., Packer, M., Biles, K., Walsh, M., Munch, S.

B., & Post, D. M. (2016). Local adaptation in transgenerational

responses to predators. *Proceedings of the Royal Society B: Biological*

Sciences, *283*(1823), 20152271.

https://doi.org/10.1098/rspb.2015.2271

Yin, J., Zhou, M., Lin, Z., Li, Q. Q., & Zhang, Y.-Y. (2019). Transgenerational

effects benefit offspring across diverse environments: A meta-analysis

in plants and animals. *Ecology Letters*, *22*(11), 1976–1986.

https://doi.org/10.1111/ele.13373