B cell expression of the enzyme PexRAP, an intermediary in ether lipid biosynthesis, promotes antibody responses and germinal center size

Sung Hoon Cho[†@&*], Marissa A. Jones, Kaylor Meyer[†],

David M. Anderson[+], Sergei Chetyrkin[+], M. Wade Calcutt[+], Richard M. Caprioli[+],

Clay F. Semenkovich[~], Mark R. Boothby[†@&] *

Author affiliations [†]Department of Pathology-Microbiology-Immunology, [@]Vanderbilt Center for Immunobiology, and [&]Vanderbilt Institute for Infection, Inflammation, & Immunology, [+]Vanderbilt University Medical Center, Nashville, TN 37232; Mass Spectrometry Research Center and Department of Biochemistry, Vanderbilt University Medical School, and [^]Cancer Biology Program, Vanderbilt University, Nashville TN 37212; [~] Division of Endocrinology. Metabolism & Lipid Research, Dept. of Medicine, Washington University, St. Louis MO 63110

* Correspondence: Sunghoon.Cho@vumc.org and Mark.Boothby@vumc.org

Dept. of Pathology, Microbiology & Immunology, Vanderbilt University Medical Center,

AA-4214 MCN Building, 1161 21st Ave. S., Nashville, TN 37232-2363

ABSTRACT

The qualities of antibody (Ab) responses provided by B lymphocytes and their plasma cell (PC) descendants are crucial facets of responses to vaccines and microbes. Metabolic processes and products regulate aspects of B cell proliferation and differentiation into germinal center (GC) and PC states as well as Ab diversification. However, there is little information about lymphoid cell-intrinsic functions of enzymes that mediate ether lipid biosynthesis, including a major class of membrane phospholipids. Imaging mass spectrometry (IMS) results had indicated that concentrations of a number of these phospholipids were substantially enhanced in GC compared to the background average in spleens. However, it was not clear if biosynthesis in B cells was a basis for this finding, or whether such cell-intrinsic biosynthesis contributes to B cell physiology or Ab responses. Ether lipid biosynthesis can involve the enzyme PexRAP, the product of the *Dhrs7b* gene. Using combinations of IMS and immunization experiments in mouse models with inducible *Dhrs7b* loss-of-function, we now show that B lineage-intrinsic expression of PexRAP promotes the magnitude and affinity maturation of a serological response. Moreover, the data revealed a *Dhrs7b*-dependent increase in ether phospholipids in primary follicles with a more prominent increase in GC. Mechanistically, PexRAP impacted B cell proliferation via enhanced survival associated with controlling levels of ROS and membrane peroxidation. These findings reveal a vital role of this peroxisomal enzyme in B cell homeostasis and the physiology of humoral immunity.

Abbreviations used:

Ag, antigen; Ab, antibody; WT, wildtype; KO, knockout; MZB, marginal zone B cell; FoB, follicular B cell; PC, plasma cell; Ig, immunoglobulin; GC, germinal center; ETC, electron transport chain; ROS, reactive oxygen species; mtROS, mitochondrial ROS; FAO, fatty acid oxidation; EL, ether lipid(s); 2D, 2-dimensional; MALDI, matrix-assisted laser desorption/ionization; IMS, imaging mass spectrometry; ER, Endoplasmic Reticulum; PexRAP, Peroxisomal Reductase Activating PPARγ; DHAP, dihydroxyacetonephosphate; G3P, glycerol-3-phosphate; CTV, CellTrace Violet; SRBC, sheep red blood corpuscle; TCR, T cell receptor; CD, Cluster of differentiation; AID, Activation-induced deaminase; GFP, Green fluorescent protein; NP-OVA, 3-Nitrophenylacetyl (NP)-ovalbumin (OVA); ASCs, Ab-secreting cells; CSR, class-switch recombination; MBC, memory B cell; NADPH, nicotinamide adenine dinucleotide phosphate; NAC, N-acetyl-L-cysteine; OCR, oxygen consumption rate; ECAR, extracellular acidification rate; RCDP, rhizomelic chondrodysplasia punctata; FAS, fatty acid synthase; Tmx, tamoxifen; 4OHT, 4-hydroxytamoxifen; PUFA, poly-unsaturated fatty acids; 2-DG, 2-deoxyglucose; FCCP, carbonyl cyanide 4-phenylhydrazone; BSA, bovine serum albumin; PSA, porcine serum albumin; H2DCFDA, dichlorodihydrofluorescein diacetate; ELISA, enzyme-linked immunosorbent assay; ELISpot, enzyme-linked immunosorbent spot; DHA, dihydroxyacetophenone; DAN, 1,5-diaminonaphthalene; timsTOF, trapped ion mobility spectrometry time-of-flight; SDS-PAGE, sodium dodecyl-sulfate polyacrylamide gel electrophoresis; SEM, standard error of means.

INTRODUCTION

The qualities and concentrations of antigen (Ag)-specific antibodies (Ab) are key elements of immunity and the pathophysiology of diverse conditions involving inflammation (1-3). Ab are secreted by cells in the B lymphocyte lineage, but the exact sources are diverse (4-7). Many derive from a subset termed B1 B cells, which are thought to having characteristics of innate responses (4, 5), but progeny of follicular and marginal zone B cells (FoB and MZB, respectively) can differentiate into Ab-secreting plasma cells (PC) after activation and proliferation (6-8). In addition to Ab concentrations and affinity for Ag, the efficacy of pathogen clearance can depend on the isotype to which B lymphocytes may transfer the Ag-combining variable region from an IgM heavy chain constant region to a new immunoglobulin (Ig) heavy chain (9-12). Importantly, the affinities of serum Ab for target Ag can increase over time after an immune exposure, especially after recurrent encounter(s) (8, 9, 13, 14).

Many sources and forms of Ab can be protective, though in several autoimmune diseases these molecules and the somatic mutations that diversify an initial BCR repertoire drive pathology (15, 16). A micro-anatomic structure termed the secondary follicle or germinal center (GC) provides one mechanism to increase spectrum of Ab affinities over time and repeated Ag exposure (17-19). Although diversification can, like PC production, occur independent from GC, results with different forms of vaccination support the practical importance of T cell help (20) and of this micro-anatomic process (21). Current evidence holds that most Ig class-switching is executed before GC entry of proliferating B cells (22). GC reactions are initiated after B cells in a primary lymphoid follicle encounter Ag, migrate to interact with activated helper T cells after extensive proliferation (23). GC formation, size, and function depend on factors that include both the efficiency of population growth for B cells in a phase before they enter and take on characteristics of GC B cells, and on homeostatic and differentiative processes while in the secondary follicle [reviewed in (24)]. Proliferation and selection end in development as Ab-secreting plasma cells or memory B cells [reviewed in (14, 18, 19)]. In line with a selection

process, substantial rates of B cell death have been measured in GC (25), so that determinants of

survival efficiency affect their outputs. As such, the qualities and quantities of Ab elicited by

immunization (or vaccination) depend on B cell proliferation and survival both prior to GC entry

as well as during GC reactions.

Accumulating evidence indicates that metabolic reprogramming and intermediary

metabolism play pivotal roles in survival, proliferation, differentiation and function in the pre-

immune populations that provide immune protection (26, 27), including B cells (28, 29). Among

core metabolic processes, several converge on oxidative metabolism fed by glycolytic generation

of pyruvate, anaplerotic use of amino acids such as glutamine, and fatty acid oxidation [reviewed

in (29, 30)]. Compared to resting populations such as naive B cells, rates of such metabolism in

B cells appear to be increased after activation, including in GC B cells [31-36; reviewed in (29)]

although in one case at a rate that may be less heightened in this specialized subset than after T-

independent B cell activation (33). Such increases support the generation of substrates for

anabolic processes in the dividing cell population and perhaps the energy demands thereof.

Mitochondria have been the focus of investigations exploring molecular mechanisms for

oxidative metabolism to generate energy and substrates in lymphocytes during immune

responses [reviewed in (37, 38)]. Oxidative phosphorylation in and dynamics of this organelle

appear to be important in determining rates of somatic hypermutation that characterize GC B cell

biology (33-35), albeit by a molecular mechanism that has not yet been identified. Use of the

Krebs TCA cycle to feed an electron transport chain (ETC) and efficient conversion of ADP to

ATP as a high-energy phosphate donor is among specialized functions of this highly dynamic

subcellular organelle. Along with other sources, mitochondria generate reactive oxygen species

(ROS) via the ETC at a rate determined by structural and biochemical features (39, 40). Rates of

ROS production, release, and resolution require regulation at multiple levels because these labile

species and their reactions with cellular molecules are important for both signaling and cell

proliferation but undermine cell function or survival when excessive.

B cells in most GC exhibit inadequacy of oxygen delivery relative to demands (31, 41-43).

Of note, genome-wide screening for genes that help cells adapt to hypoxia revealed a

requirement for peroxisomes and an ER enzyme involved in synthesizing plasmalogen ether

lipids (EL) (44). Using an unbiased lipidomic analysis that combined imaging mass spectrometry

(IMS) and techniques for identifying GC in sections of spleen from immunized mice, we had

found that after immunization, local concentrations of a subset of EL were heightened in GC

relative to other portions of the spleen (45). Ether phospholipids (EPL) have a fatty acid linked to

the glycerol backbone with an alkyl or a vinyl ether bond at the sn-1 position (46, 47). Synthesis

of ether phospholipids depends on the generation of precursors in peroxisomes and final

processing in the ER (48, 49). However, EL biosynthesis is prominent in the liver, and in theory

these species or their precursors might distribute to tissues from a primary site of biosynthesis or

from the diet (49-53). Thus, the finding that secondary follicles had greater densities of some -

but not all - ether phospholipids left several key questions unanswered – (1) Does the capacity to

synthesize and increase ether lipids in B cell follicles – primary or secondary (i.e., GC) - affect

antibody responses? (2) Are any of the EL dependent on B lineage-intrinsic metabolism to

generate precursors or final products? Moreover, there is debate whether EL or their plasmalogen

subset promote cell death or resistance, and whether inactivation of biosynthesis causes defects

in establishment or maintenance of hematopoietic cells (46, 53-55).

Peroxisomal Reductase Activating PPARγ (PexRAP, encoded by the *Dhrs7b* gene) catalyzes

the reduction of alkyl-dihydroxyacetonephosphate (DHAP) to 1-alkyl-G3P (46, 48, 56), which

then is transferred into endoplasmic reticulum (ER) and further metabolized to generate EL.

Recent work notes that this enzyme may substantially function in the ER in addition to the

peroxisome (56). Widespread inactivation of *Dhrs7b* in mature mice decreased hepatic

production of ether lipid species and reduced the population of erythrocytes and neutrophils (46).

Although questioned based on inborn errors of metabolism affecting other aspects of ether lipid

biosynthesis (54), this body of work indicated that PexRAP can be essential in some aspects of

ether lipid metabolism and may function in hematopoietic cells. Diverse functions of ether lipids

in general, and the plasmalogen subset in particular, have been proposed or supported by prior

work [reviewed in (57-61)]. The ether linkage may provide a sink for oxygen radicals - thereby

acting directly to resolve the reactivity and contribute to ROS homeostasis, although there is

debate as to the physiological impact of this chemical property (62). Whether or not this lipid

subset influences adaptive immunity or conventional lymphocyte lineages, however, is unclear.

Herein, we tested the hypotheses that (i) the ether lipid profiles in activated B cells depend

on their expression of the biosynthetic enzyme PexRAP [alternatively referred to as acyl/alkyl-

DHAP reductase (ADHAPR) (56)], and (ii) this enzyme promotes B cell physiology. To do so,

we combined conventional lipidomics, Imaging Mass Spectrometry (IMS), and immunization

experiments with genetic models in which loss-of-function for *Dhrs7b* is induced in adult mice.

We provide evidence that PexRAP impacts B cell proliferation via enhanced survival associated

with controlling levels of ROS and membrane peroxidation. The results also indicate that beyond

an impact on B cell homeostasis, GC are reduced in its absence, and the magnitude and affinity

maturation of a serological response depend on B cell expression of PexRAP. Taken together,

these finding support a function of B cell-intrinsic biosynthesis of ether lipids in B lymphocyte

homeostasis, the production of Ab, and in promoting GC reactions.

103
104
105

RESULTS

PexRAP promotes homeostatic maintenance and proliferation of B cells. By use of IMS, we had reported that at least a dozen ether phospholipids - including plasmalogens - were enriched in splenic germinal centers of immunized mice when compared to the rest of the tissue (45). While previous work provided evidence that these lipid molecules were important for the homeostasis of short-lived neutrophils (46), this effect was questioned based on inborn errors of metabolism (54). Moreover, the impact of these molecules on adaptive immune cells or functions is not known. Accordingly, we sought to test for effects of an intervention that interferes with a key biosynthetic pathway used for endogenous synthesis of these molecules. To do so, we started with analyses using induced loss-of-function for $Dhrs7b^{f/f}$ and the widely expressed $Rosa26$-$CreER^{T2}$ transgene (63, 64) used in (46), which reduced hepatic generation of a subset of ether lipids and acts in all hematopoietic cells as well.

We modeled initial experiments on the published work with imaging mass spectrometry (45). Mice were treated with tamoxifen using conditions less intense than a regimen shown to have no effect on pre-immune splenic populations or B lineage (65, 66), immunized with SRBC [akin to the analyses in (31, 36, 41, 45)], and analyzed 7 d after immunization (Fig. 1A). Functional inactivation of $Dhrs7b$ in T and B lymphocytes was confirmed by Western blot analysis using anti-PexRAP Ab (Fig. 1B). The frequencies and numbers of CD19$^+$ B220$^+$ B cells in $Dhrs7b^{f/f}$; $Rosa26$-$CreER^{T2}$ - only a small minority of which would have been activated by the immunization - were approximately 0.6 the values for controls ($Dhrs7b^{+/+}$ $Rosa26$-$CreER^{T2}$, i.e., wild-type locus) (Fig. 1C, D). In contrast to B cell populations, TCRβ$^+$ CD4$^+$ T cells and TCRβ$^+$ CD4$^-$ CD8 T cells were intact in tamoxifen-treated and SRBC-immunized $Dhrs7b^{f/f}$; $Rosa26$-$CreER^{T2}$ mice (Supplemental Fig. 1A-C). These results suggest that $Dhrs7b$ is non-redundantly required for fully functional B cell maintenance but indicate that the enzyme is dispensable for the main populations of conventional T cells. In light of the earlier finding that a

subset of ether lipids was more prevalent in secondary follicles, we analyzed GC in these

samples. Microscopy and flow cytometry analyses after immunofluorescent staining found that

less GC B cells were present after immunization of *Dhrs7b^{f/f}*; *Rosa26-CreER^{T2}* mice, i.e., less

IgD^- CD95^+ GL7^+ (GC-phenotype) B cells (Fig. 1E-I; Supplemental Fig. 1D). Quantitation of the

immune fluorescence micrography with spleens from immunized mice revealed that the numbers

of GC per spleen were halved in *Dhrs7b^{Δ/Δ}* mice compared with controls (Fig. 1I; Supplemental

Fig. 1E). The sizes of those GC that did form were substantially reduced as well (Fig. 1I;

Supplemental Fig. 1F). The lower number of B cells found in immunized mice - while a less

profound effect than the reduction in GC - tempers the result, and it was possible that the

function of helper CD4 T cells was impaired even their though numbers were not affected. While

not addressing whether or not B cells are part of the requirement for PexRAP in promoting GC,

these results indicate that *Dhrs7b* is necessary for a normal-sized geminal center response.

To determine if there are B lineage-specific functions of PexRAP, we used a conditionally

active Cre transgene that is expressed specifically in mature B cells (67). *Dhrs7b* f/f , *huCD20-*

CreER^{T2} mice and *huCD20-CreER^{T2}* were treated with tamoxifen to induce Cre-mediated

recombination of the floxed alleles after initial establishment of normal B cell populations (Fig.

2A). Western blot analyses showed an almost complete absence of PexRAP protein from B cells

purified from tamoxifen-treated *Dhrs7b^{f/f}*; *huCD20-CreER^{T2}* (hereinafter, *Dhrs7b^{Δ/Δ-B}*) mice

compared to similarly treated *huCD20*-CreER^{T2} controls (Fig. 2B). The enzyme encoded by this

gene catalyzes the synthesis of a lipid precursor to many - but not all - plasmalogens and other

ether phospholipids (46-48), and *Dhrs7b* deficiency impacted neutrophil survival (46). When we

tested if *Dhrs7b* inactivation affected the steady-state B cell population, the frequencies and total

numbers of CD19^+ B220^+ B cells in viable lymphocyte gates were about 20% lower in

tamoxifen-treated *Dhrs7b^{Δ/Δ-B}* mice compared to tamoxifen-injected, *huCD20*-CreER^{T2},

Dhrs7b^{+/+} controls (Fig. 2C). With loss-of-function after production of mature B cells, i.e.,

inactivation of *Dhrs7b* at ages over 6 wk, we observed balanced decreases in the numbers of

multiple subsets within the B lineage without evidence of selectivity among main sub-classes of

B cell (Supplemental Fig 1G-I). Thus, the frequencies of MZB and FOB amidst the smaller

population of splenic B cells in tamoxifen-treated *Dhrs7b^{f/f}; huCD20-CreER^{T2}* mice were similar

(Supplemental Fig 1H), and frequencies of B1 B cells in the peritoneal cavity also were

unaffected (Supplemental Fig 1I). Prior work documents normal frequencies and numbers of

developing and mature B cells under harsher conditions of tamoxifen-induced deletion with

CreER^{T2} (65). De novo B cell production rates are too low to yield the ~20% decrease observed

one week after gene disruption, and few splenic B cells are in cell cycle. Accordingly, we infer

that after maturation of a B cell, *Dhrs7b* promotes its survival, albeit to a modest degree.

B cell population growth can be impacted by increased cell death. To test if PexRAP affects

proliferation in vivo, *Dhrs7b^{Δ/Δ}* B cells were stained with CellTrace Violet (CTV) and adoptively

transferred into B cell deficient μMT recipient mice. Strikingly fewer PexRAP-depleted B cells

were recovered (Fig. 2D), and the frequencies of *Dhrs7b^{Δ/Δ}* B cells that were IgD⁻ (i.e., had been

activated) were half those of controls (Fig. 2E; supplemental Fig. S1J). As compared to control B

cells, CTV partitioning analyses suggested that division rates in vivo were modestly lower for

the *Dhrs7b^{Δ/Δ}* B cells (Fig. 2F, G). Mitogen-stimulated *Dhrs7b^{Δ/Δ}* B cells that proliferated in

culture exhibited substantially less robust division: frequencies of viable cells that had undergone

>3 divisions were halved in *Dhrs7b^{Δ/Δ}* B cells) and yielded ~1/3 as large a progeny population

(Fig. 2G, H). Collectively, these data indicate that the *Dhrs7b* gene product supports homeostatic

maintenance of a quiescent (pre-immune) B cell population *in vivo* and effective proliferation of

B cells.

B cell expression of PexRAP is required for achieving normally distributed concentrations

of ether phospholipids. As noted, lipidomic analyses that used 2-dimensional image mass-

spectrometry (2D-IMS) discovered that local concentrations of a subset of ether lipids are

enriched in GC compared with the area outside of GC after immunization (45). To measure the

extent to which PexRAP expression within mature B cells can affect their lipid content and

composition, including their ether- and plasmalogen phospholipids, LC-MS-MS analyses were

performed with B cells directly isolated from spleens as well as those cultured after mitogenic

activation. These analyses showed substantial reductions in many ether phospholipids - as well

as lysophosphatidylethanolamines (LPE) generated by phospholipase cleavage of the R2 fatty

acid sidechain of an ether phospholipid - in PexRAP-deficient B cells (Fig. 3A, B; Supplemental

Fig. 2A, B). While beyond the scope of this work, this evidence suggests that generation of

potential signaling molecules via phospholipase(s) such as PLA2 may be reduced. Of note,

whereas differences were modest or absent in the naive population, the greatest impact of

PexRAP on quantitative amounts of phospholipids was found in the activated B cells, for which

tamoxifen treatment and CreERT2 made minimal differences (Supplemental Fig. 2C, D). Several

species, skewed towards those with the most detected ions, were at less than 1/5th the level of

non-deleted control B cells (Fig. 3C, D). There is no in vitro surrogate that faithfully represents

GC or the GC B cell, but collectively these data provide strong evidence that the capacity to

rapidly produce increased levels of many ether lipid molecules after lymphocyte activation

depends on a functional *Dhrs7b* gene in B cells. Although additional uptake or biosynthesis

pathways may add to the overall pool, as previously noted (46, 47) - and especially in resting B

cells that are out of cycle - the data indicate that *Dhrs7b* in B cells enhances cell-autonomous

biosynthesis. The straightforward inference is that PexRAP catalyzes the generation of ether lipid

precursors in amounts crucial for regulating the overall ether phospholipid pools, especially after

B cell activation.

Our earlier work (45) reported only ion mode features in negative mode and did not analyze

the primary follicle (the vast majority of which are resting B cells) relative to the extrafollicular

white pulp, red pulp, and GC. To investigate these issues and ultimately the requirement for

specific expression of PexRAP, we started with AID-GFP mice to enhance spatial localization.

These measurements confirmed that a substantial number of both positive and negatively

charged ions whose exact masses identified them as plasmalogens were substantially

concentrated in GC as compared to the rest of the B cell follicle (Table 1; Supplemental Fig. 3A,

B). Accordingly, we tested if the differential enrichment of any ether lipid species in primary or

secondary follicles depends on biosynthesis within a specific lymphoid lineage. Mice with

conditional mutations were tamoxifen-treated, immunized with SRBC, harvested seven days

thereafter, and analyzed by IMS (Fig. 4A, B). With deletion driven by the widely expressed

Rosa26-CreER^{T2} transgene, as in earlier work (46) and Fig. 1, substantial reductions of the GC

enrichment pattern were observed (Fig. 4C). For instance, two representative ions identified in

negative ion mode (i.e. *m/z* 752.5545 and *m/z* 776.5556) were again enriched in GC regions of

spleens from immunized WT mice. These features barely increased in splenic samples of

immunized *Dhrs7b*$^{Δ/Δ}$ (tamoxifen-injected *Dhrs7b*$^{f/f}$; *Rosa26-CreER*^{T2}) mice (Fig. 4C).

To determine if the levels of particular ether phospholipids in GC regions were affected by

PexRAP expression in B cells, *Dhrs7b*$^{f/f}$; *huCD20*-CreER^{T2} mice and *huCD20*-CreER^{T2} controls

were analyzed after tamoxifen injections followed by immunization, and compared to samples

from unimmunized mice. IMS using both positive and negative ion modes identified at least

eight ions - including *m/z* 752.5545, *m/z* 776.5556, and *m/z* 872.5749 - much more substantially

(~2-3-fold) concentrated in GC of immunized control mice (Fig. 4D, E; Table 2) than in the

primary follicle of unimmunized mice. Notably, these increased signals in secondary follicles

(GC) were reduced or almost completely eliminated in GCs of mice immunized after B cell-

specific PexRAP depletion (*Dhrs7b*$^{Δ/Δ-B}$) (Fig. 4D, E; Table 2; Supplemental Fig. 3C).

Collectively, these results indicate that *Dhrs7b* gene expression in activated B cells regulates the

spectrum of ether lipids in the micro-anatomic locale of a secondary follicle. Interestingly,

immunization also led to modest increases in the levels of some ether and conventional

phospholipids in the B cell-rich primary follicle regions (Table 2). This finding extends the

previous report (45), which focused on the relationship between the AID-GFP^{hi} region (i.e., GC)

and lipid features. Moreover, the absence of PexRAP blunted most of the immunization-induced

increases observed in the primary follicles. These findings and those of Fig. 3 suggest that there

are secondary consequences of *Dhrs7b* inactivation (e.g., some di-acyl phospholipids are

affected), but provide direct evidence that the enhancement of many ether phospholipid signals

in the GC depends on PexRAP in B cells.

B cell-intrinsic role of *Dhrs7b* in Ab affinity and quantity.

As key components of adaptive immunity, progeny of an activated B cell can differentiate

into Ab-secreting plasma cells (PC) or into GC B cells - whose physiology diversifies the affinity

for and breadth of antigen recognized by the original clone and improves properties of memory

[(14, 17-19, 68). PC that develop after a second encounter with antigen yield affinity maturation,

and the higher quality and proper quantity of Ag-specific Abs from plasma cells are integral to

humoral immune responses (68). An increasing body of evidence indicates that metabolic

reprogramming and intermediary metabolites modulate immune cell differentiation and function

(26-36). The observed B cell-intrinsic function of *Dhrs7b* in the accumulation of many ether

lipid species in both primary and secondary follicles prompted us to test the effect of B cell type-

restricted PexRAP depletion on GC responses and Ag-specific Ab production. Tamoxifen-

injected *Dhrs7b^{f/f}; huCD20-CreER^{T2}* and control mice were immunized with NP-ovalbumin (NP-

OVA), then boosted with NP-OVA to elicit affinity-matured Ab, and harvested 1 week thereafter

(Fig. 5A). Although PexRAP-deficient B cell numbers were reduced less than 20% a week after

starting gene inactivation prior to immunization (i.e., were over 0.8-fold those of controls) (Fig

2A-C), frequencies and numbers of IgD^{neg} B cells were halved in *Dhrs7b^{Δ/Δ-B}* mice at harvest (4

wk after completion of the induced deletion) (Fig. 5B, C). Moreover, the GL7^{+} CD95^{+} fraction

of the B cell population in the dump / IgD^{neg} gate was substantially reduced (Fig. 5D), such that

numbers of GL7^{+} CD95^{+} GC B cells were dramatically decreased in *Dhrs7b^{Δ/Δ-B}* mice (Fig. 5E).

Consistent with these findings, GC in *Dhrs7b^{Δ/Δ-B}* mice were reduced after SRBC immunization,

with a modest change in the DZ/LZ ratio (Supplemental Fig. 4A, B).

Ab class-switch recombination (CSR) and affinity maturation can occur through

extrafollicular response, but the GC reaction significantly increases Ab diversification and high-

affinity Ab production, in part later in a primary response but also upon secondary exposure to

antigens (6, 8, 9, 17-19). ELISA on sera at 3 wk post-immunization (Fig. 6A-D), just before the

boost, showed that B cell-specific depletion of PexRAP prior to immunization reduced NP-

specific IgM and the switched isotype IgG1 (Fig. 6A, C). Of note, the capacity to generate high-

affinity Ab, detected with low valency NP2, was even more severely undermined than the overall

response - especially for IgG1 (Fig. 6B, D). A week after a second immunization, Ab-secreting

cells (ASCs) in the spleen (Fig. 6E) and circulating anti-NP IgM concentrations were diminished

in $Dhrs7b^{\Delta/\Delta-B}$ mice compared with controls that were $huCD20$-CreER^{T2}, $Dhrs7b^{+/+}$ treated with

tamoxifen in parallel to the mice with induced loss of PexRAP (Fig. 6F). The ratio of high-

affinity (NP2-binding) to all-affinity (NP20) IgM Ab also was substantially lower in $Dhrs7b^{\Delta/\Delta-B}$

mice (Fig. 6G), as were the serum levels of high-affinity IgG1 and IgG2c, class-switched

isotypes (Supplemental Fig. S4C, D). These data reinforce and extend the conclusion that the

expression of PexRAP in mature B cells is important for their physiology and function.

Consistent with *in vivo* results finding reduced Ag-specific ASCs in $Dhrs7b^{\Delta/\Delta-B}$ mice

compared with controls (Fig. 6E), the attenuated population of PexRAP-deficient B lineage cells

recovered after transfer into recipient mice (Fig. 2D) yielded lower frequencies of CD138[+]

progeny (Fig. 6H, I) along with evidence of reduced activation [i.e., lower frequencies of IgD[-]

progeny (Supplemental Fig. 1I). Mitogen-stimulated $Dhrs7b^{\Delta/\Delta}$ B cells also yielded lower

frequencies of CD138[+] cells in vitro, but the division-specific frequencies of CD138[+] cells

showed only modest decreases in $Dhrs7b^{\Delta/\Delta}$ B cells (Supplemental Fig. 4E). Thus, the reduction

of CD138[+] cell differentiation appears to be due mostly to its dependence on survival to a

sufficient division count.

Dhrs7b inactivation was initiated prior to immunization in the preceding experiments.

Therefore, the impact of PexRAP deficiency on GC and the Ab response in such a setting might

be due exclusively to impairment of B cells, e.g., their reduced population expansion, prior to

their entry into GC. Alternatively, the effects could also in part involve a requirement for

PexRAP-dependent metabolites within GC B cells. To test if *Dhrs7b* functions within GC, we

used conditional deletion of *Dhrs7b* driven by the *S1pr2-CreER^{T2}* transgene whose expression at

high levels marks GC B cells (69). Of note, experiments with a fate-marking reporter allele

showed that activated B cells that lack the GC B phenotype were not marked by this conditional

Cre after immunization with the NP-carrier approach (69). *Dhrs7b^{f/f}*; *S1pr2-CreER^{T2}* and control

S1pr2-CreER^{T2} (control) mice were immunized with SRBC, and tamoxifen was injected at a time

point after the initiation of GC to assure more fully that the inactivation of *Dhrs7b* would be in

GC B cells rather than pre-GC blasts (Figure 7A). Frequencies of GC B cells were substantially

lower in tamoxifen-treated *Dhrs7b^{f/f}*; *S1pr2*-CreER^{T2} mice (Figure 7B, C), whereas the overall

populations of $CD138^+$ cells, and $CD19^+$ $B220^+$ plasmablasts (PBs) among them - almost all of

which would have developed prior to the induced loss-of-function - were normal (Figure 7D, E).

These data provide evidence that *Dhrs7b* expression within GC B cells influences them during

an Ab response. Collectively, these data indicate that an optimal GC response requires PexRAP

function in B cells, and the B cell-intrinsic functions of *Dhrs7b* promote the quantity and affinity

maturation of Ab responses, in part via net proliferation of B cells.

***Dhrs7b* contributes to modulation of ROS and their impact on B cell population growth.**

Increased susceptibility to cell death after activation would be a mechanism that could

cause the reduced population expansion, GC, and Ab production. Consistent with earlier analyses

of LPS-stimulated B cells (70), we had noted that ROS levels were higher in GC B cells,

memory B cells (MBCs) and ASCs than in the naive B2 B cell pool (36). ROS can be produced

in activated B cells via diverse sources that include mitochondrial electron transport chain

function, NADPH oxidase activated by BCR engagement, and fatty acid oxidation in

peroxisomes (71-73). ROS can positively mediate signal transduction, but their steady-state

concentrations need to be calibrated because overproduction can have deleterious effect on cells,

for instance via lipid peroxidation at excessive rates (74-76). Endogenous antioxidant properties

have been imputed to plasmalogens because the vinyl ether bond is susceptible to reaction with

reactive oxygen, thereby scavenging ROS (48, 57-59). When we tested if *Dhrs7b* function is

critical for redox control in B cells, mitogen-stimulated *Dhrs7b*$^{\Delta/\Delta}$ B cells exhibited ~3 fold

higher signal and about a doubling when stained with fluorescent sensors of cellular and

mitochondrial ROS, respectively, compared to WT B cells *in vitro* (Fig. 8A-D). Iron- and copper-

dependent lipid peroxidation is considered to be a biological mechanism for ROS-mediated cell

death (75, 76). Lipid peroxidation measured by a fluorescent indicator (C11-Bodipy) in vitro was

enhanced in mitogen-activated *Dhrs7b*$^{\Delta/\Delta}$ B cells compared with WT controls (Fig. 8E). One

independent means of driving increased ROS and lipid peroxidation in B cells (supplemental Fig.

S5A) rapidly led to reduced cell viability and B cell numbers (Fig. 8F). Furthermore, in vitro

analyses indicated that B cells lacking PexRAP also exhibited increases in the activated

executioner caspase, cleaved caspase-3 (CC3) along with early apoptotic cells that are annexin

V^{+} but exclude 7-aminoactinomycin D (Fig. 8G, H). These data indicate that *Dhrs7b* function

supports not only protection against cell death, but also control of cellular and mitochondrial

ROS levels and their impact on lipid modification. Of note, while H_2O_2 did not increase this

early apoptotic population (Supplemental Fig. SC), increasing ROS with menadione drove B cell

death preceded by annexin V^{+} state without an increase in C11-Bodipy signal (Supplemental Fig.

S5D-G). We infer that rather than a single mode of death, increased ROS resulting from PexRAP

depletion stimulates at least two distinct modes of B cell death.

Consistent with the increased steady-state ROS and death, activated *Dhrs7b*$^{\Delta/\Delta}$ B cells

exhibited a defect in B cell population growth *in vivo* and *in vitro* (Fig. 2; Fig. 8A-H). The well-

established ROS scavenger, N-acetyl-L-cysteine (NAC), exerted a concentration-dependent

effect that improved the survival and population growth of PexRAP-deficient B cells (Fig. 8I).

We exploited the sub-maximal rescue by a lower concentration of NAC (1 mM) to explore if

peroxisomal oxidative metabolism might contribute to the toxicity observed when B cells are

PexRAP-depleted. Although thioridazine on its own provided no increase in the B cell

population growth, its combination with 1 mM NAC treatment improved growth of the

Dhrs7b$^{\Delta/\Delta}$ B cell population compared to either agent on its own (Fig. 8J). These data indicate

that sufficient generation of PexRAP-dependent products is crucial for B cell survival. Moreover,

this enzyme contributes - directly or indirectly - to a resolution or detoxification of ROS that is

critical for population growth of activated B cells.

***Dhrs7b* in B cells affects their oxidative metabolism and ER mass.**

Our finding that mitochondrial ROS were increased prompted us to investigate how PexRAP

affects physiological functions by measuring respiration in activated B cells. Mitochondrial

stress tests with in vitro activated B lymphoblasts measured small but definite decreases in both

basal and maximal respiration (oxygen consumption rates, or OCR) (Fig. 9A-C). The altered

performance of mitochondria was associated with reductions in calculated ATP generation and

proton leak as well as a small decrease in spare respiratory capacity (SRC) (Fig. 9D-F,

respectively). In contrast, the rates of glucose-stimulated extracellular acidification - cytosolic

reactions and a surrogate that can approximate glycolytic activity - were unaffected (Fig. 9G).

Thus, the altered respiration did not reflect a decrease in all cell metabolism. Inasmuch as

peroxisomes and mitochondria form contacts and functionally interact with the ER, we measured

relative ER mass and found that the signal of the fluorophore ERTracker was consistently

reduced in PexRAP-depleted B cells (Fig. 9H, I). We infer that both mitochondrial function and

ER mass in B cells are promoted by their expression of PexRAP.

DISCUSSION

We have shown herein that B lymphocyte expression of an enzyme essential for biosynthesis

of a subset of ether phospholipids is crucial for part of the increases in their levels localized to

GC. Moreover, the genetic approach used to deplete B cells of PexRAP allowed us to show that

the B cell type-restricted gene, *Dhrs7b*, promotes B cell survival, proliferation, and GC size

along with affinity maturation and serum concentrations of Ag-specific Ab after immunization.

Taken together, this work establishes the predictive value of the discovery approach that

identified an unexpected feature of cellular biochemistry in GC and indicates that B cell-

autonomous generation of ether lipid species is crucial for B cell survival and the qualities of the

Ab elicited by immunization.

Using imaging mass spectrometry for discovery-based hypothesis generation, we had

reported that the concentrations of at least a dozen ether phospholipids are increased in splenic

GC relative to the remainder of the tissue (45). The analyses presented here confirm and extend

the observations, and provide evidence that after immunization, these concentrations increase

even in the primary B cell follicle, albeit to a modest degree when compared to the magnitude of

increases in GC, and in B cells activated ex vivo. The cell type-specific gene inactivation after

establishment of a pre-immune population establishes that an enzyme crucial for generation of a

number of plasmalogens has substantial effects on B lymphocyte physiology and function.

Plasmalogens and other ether lipids and phospholipids have long been shown to be major

constituents of lipid bilayers and cell membranes (48, 49, 57). However, remarkably little is

known about whether or not cell-intrinsic synthesis of ether lipids or the balance among their

specific molecular species matters for hematopoietic cells or in immunity. Several inborn errors

of metabolism in humans are attributable to loss-of-function mutations of genes encoding

peroxisomal proteins that impact ether lipid (and hence plasmalogen) synthesis (58) but affect

other critical processes as well [reviewed in (48, 58-62)]. Several of these - such as Zellweger

Syndrome and rhizomelic chondrodysplasia punctata (RCDP) - and the mouse models generated

to study mechanisms in these human disease, lead to severe neurological defects and early post-

partum death (57, 58). Whether or not lymphocyte development, homeostasis or function were

affected in these studies is not clear. Deficient generation of ether lipids is among many

abnormalities caused by elimination of fatty acid synthase (FAS) (46, 47). Acute inactivation of

Fasn, the gene encoding this enzyme, in young mature mice caused a decrease in spleen size

disproportionate to the reduction in neutrophils, which was the most profound hematological

consequence (46). In addition, circulating lymphocytes were reduced despite normal steady-state

representation in the marrow. In-trans cell-extrinsic functions of ether lipids have been noted

previously - specifically, as ligands for the invariant natural killer T cell receptor (77) or for a G-

protein coupled receptor on natural killer cells (78). As such, the widespread loss of FAS

function and the potential for indirect and pleiotropic effects of this enzyme deficiency left

unresolved what cells actually depend on their own synthesis of ether lipids.

PexRAP, the enzyme encoded by the *Dhrs7b* gene, generates intermediates vital for the

synthesis of some - but not all - ether lipids in cells such as neutrophils (46). Analyses of

neutrophil numbers and survival provided evidence that this enzyme can be crucial for a normal

membrane composition. Moreover, PexRAP promoted viability of this very short-lived cell type

whose membranes have a high plasmalogen content (46). The bloodborne population of

lymphocytes was reduced in the *Rosa*26-CreER^{T2}, *Dhrs7b*$^{f/f}$ mice (46), but the specific types of

lymphoid cell were not reported and the circulating cell number does not necessarily correlate

with that in spleen or other organs. Of note, although subject to points articulated in (53), the

interpretation that the neutrophil phenotype is due to insufficiency of ether lipids or

plasmalogens has been questioned based on alternative gene disruption results (54). Several

models of unconditional gene inactivation or mutation that drastically reduced the phospholipid

end-products were noted to have normal neutrophils [summarized in (54), albeit at older ages (4 -

7 mo) than those used in (46). As noted, however, there may be age-related changes in the

processes analyzed herein using young (age ~ 2 mo) mice (53). Moreover, cellular and

organismal selection for adaptation variants makes the results of acute loss-of-function inherently

different from unconditional loss and heterozygote parentage (53). The potential for toxicity

induced by 4OHT activation of CreER^{T2} was among concerns mooted about the earlier study of

neutrophils (54). This concern may have been extrapolated from papers that noted impairment in

earlier hematopoiesis when *Rosa26*-CreER^{T2} was combined with intensive regimens of

tamoxifen administration to infant mice (65, 79). Of note, such studies used individual doses

over 4-fold higher that those used herein, and delivered by gavaged five times on a daily basis,

yet the IgM$^+$ B cell population sizes in spleen and marrow were normal even in the face of

reduced erythropoiesis and thymopoiesis (65). Moreover, the analyses herein compared B

lineage-specific deletion to CreER$^{T2+/-}$, *Dhrs7b*$^{+/+}$ mice identically dosed in parallel with the test

subjects. Thus, the work herein indicate that PexRAP is essential for normal physiology and

function of mature B lymphocytes, in particular after their activation.

The findings are most consistent with a capacity for PexRAP function to contribute towards

B lymphocyte survival and effective proliferation by restraining both executioner caspase

activation and lipid peroxidation due to excessive ROS levels. These findings are consistent with

those obtained with neutrophils (46). On the surface such results might appear to present a

conundrum relative to work indicating that both fatty acyl-CoA reductase 1 and the enzyme that

later generates an alkenyl bond at a terminal step in plasmalogen biosynthesis are crucial for the

susceptibility of HT-1080 and 786-O tumor cell lines to ferroptosis (55). However, our data are

consistent with published evidence that PexRAP is needed for a narrower subset of ether lipids

than the broad requirement for FAS or fatty acyl-CoA reductases (46-49, 80, 81). Moreover,

molecular dissection of mechanisms in various non-hematopoietic cell types (including the

cancer line 786-O) showed that ether lipid biosynthesis can either promote susceptibility or

resistance to ferroptosis (82). Collectively, our findings suggest that for B cells the inactivation

of PexRAP tilts the balance toward death and reduced Ab - perhaps involving lower fractions of

poly-unsaturated fatty acids (PUFA) (82), or a lower ratio of plasmalogens (alkenyl linkages)

relative to alkyl-linked ether lipids.

PexRAP localizes to peroxisomes (35-37), and prior work has suggested both that

peroxisomes are increased in GC B cells and likely contribute to FAO by these cells (26). In line

with the possibility that this organelle may affect PexRAP-dependent survival, our data indicated

that specific inhibition of peroxisomal FAO could collaborate with sub-optimal ROS scavenging

to mitigate the reduced growth of activated B cells. However, recent findings show that PexRAP

also is expressed in the ER and is capable of contributing to ether lipid biosynthesis via catalysis

in this organelle in addition to or instead of the peroxisome (56). Thus, although ROS generated

in peroxisomes likely contribute to the stress experienced by PexRAP-deficient B cells, the

phenotypes observed herein may not reflect the need for the peroxisomal fraction of the enzyme.

Our findings with an intermediary in ether lipid biosynthesis add to an increasing body of

evidence that metabolism in B lymphocytes affects their differentiation or function (28-36).

Many sources and forms of Ab can be protective, while in several autoimmune diseases these

molecules can drive pathology (1, 2). The capacity to generate increased affinities and circulating

concentrations of Ab is an important factor both in protective immunity and auto-immune

disease. Antigen-independent, TLR-driven PC differentiation requires increased oxidative

phosphorylation (70), while oxidative metabolism in GC B cells promotes affinity maturation by

unknown mechanisms (33-35). While not the exclusive determinant of such properties, GC are

major sources of higher affinity Ab with a broader spectrum of specificities (9, 13, 14). GC

formation, size, and function depend on the efficiency of population growth for B cells in a phase

before they enter and take on characteristics of GC B cells, and then homeostatic and

differentiative processes while in the secondary follicle. Thus, while the findings with the post-

immunization inactivation of *Dhrs7b* using *S1pr2*-CreER[T2] provide evidence that the expression

of PexRAP is important within GC B cells, the decreases of GC and affinity-matured Ab are

likely also to involve a pre-GC phase of clonal expansion. In any case, the rapidity of the effects

suggests that pharmacological inhibition of PexRAP may be a target in inflammatory disease in

which B cells participate, particularly in light of the potential for impacts on pro-inflammatory

neutrophils.

486
487

MATERIALS & METHODS

Reagents Monoclonal antibodies (mAb) against mouse CD40, CD90.2, B220 (CD45R), and other mAbs (purified, biotinylated, or fluorophore-conjugated) were from BD Biosciences or Tonbo Biosciences (San Diego CA) unless otherwise indicated. BAFF was from AdipoGen (San Diego, CA). Recombinant mouse IL-4 and recombinant mouse IL-5 were from Peprotech (Rocky Hill NJ). Glucose, 2-deoxyglucose, oligomycin, rotenone, antimycin A, carbonyl cyanide 4-phenylhydrazone (FCCP), N-acetyl-cysteine (NAC), thioridazine hydrochloride, H_2O_2, and 4-hydroxytamoxifen (4-OHT) were from Sigma-Aldrich Chemicals (St. Louis MO), and menadione from Cayman Chemicals (Ann Arbor, MI). Tamoxifen (Tmx) was from APExBio Technology (Houston TX). NP-BSA and NP-PSA (for capture in ELISA) as well as NP-OVA (for immunization) were obtained from Biosearch Technology (Novato CA). SRBCs (sheep red blood cells) were from Thermo Fisher Scientific (Waltham MA). CellTrace™ Violet cell proliferation kit, CM-H_2DCFDA, MitoSOX™ Red, and BODIPY™ 581/591 C11 were obtained from Invitrogen (Waltham, MA).

Mice, immunizations, ELISA and ELISpot All animal protocols were reviewed and approved by the Vanderbilt University Institutional Animal Care and Use Committee. Mice were housed in ventilated micro-isolators under Specified Pathogen-Free conditions in a Vanderbilt University Medical Center mouse facility and used both male and female mice at 6–8 weeks of age. For cell type specific inactivation of *Dhrs7b* genes, *Dhrs7b*[f/f] mice (46) were crossed with *Rosa26*-CreER[T2] (46), *huCD20*-CerER[T2] (30), or *S1pr2*-CreER[T2] strains (51, 52), all on C57/BL6 genetic background. Tamoxifen was administered as previously reported (25, 30). To control for potential Cre toxicity, CreER[T2] mice were similarly injected to use as wild-type (WT) controls. Mice (ages ~ 8wk) were immunized with SRBCs (2×10^8 cells per mouse) and analyzed 1 week after immunization as described (25, 45). Alternatively, mice were immunized and boosted by i.p. injections [each of 100 µg NP[16]-OVA (Biosearch Technologies, Novato, CA) emulsified in

100 µL of Imject^{TM} Alum (Thermo Fisher Scientific, Pittsburgh, PA)] as described previously

(25, 30), and harvested for analyses 1 wk after the 2^{nd} injection. Isotype-specific relative levels of

Ag-specific Ab were quantitated by capture ELISA using Ag (NP$_{20}$-BSA or NP$_2$-PSA for all- or

high-affinity hapten-specific Abs, respectively) followed by SBA Clonotyping System (Southern

Biotech, Birmingham AL), as described previously (25, 30). As previously described (30),

frequencies of Ab-secreting cells (ASCs) were analyzed by ELISpot and quantitated using an

ImmunoSpot Analyzer (Cellular Technology, Shaker Heights OH).

Cell cultures, proliferation assay, and reversion of population growth   Splenic B

cells were purified (90-95%) using negative selection as previously described (30) or by positive

selection with anti-mouse B220 microbeads (Miltenyi Biotech, Auburn, CA). B cells (5×10^5

cells in 1 ml) were activated with anti-CD40 (1 µg/mL) and cultured with BAFF (10 ng/mL),

IL-4 (10 ng/mL), IL-5 (10 ng/mL) and 4-OHT (50 nM) in Iscoves Modified Dulbecco's Medium

(IMDM) supplemented with 10% Fetal Bovine Serum (FBS), 100 U/mL penicillin 100 µg/mL

streptomycin (Invitrogen), 3 mM L-glutamine, nonessential amino acids (Invitrogen), 0.1 mM 2-

ME (Sigma-Aldrich). To analyze the effect of ROS scavenger and inhibition of peroxisomal lipid

oxidation on population growth, bead-purified B cells from tamoxifen-treated *huCD20*-CerER^{T2}

or *Dhrs7b$^{f/f}$*; *huCD20*-CerER^{T2} mice were activated with anti-CD40, cultured with BAFF, IL-4

and IL-5 in the presence or absence of NAC (1 mM and 5 mM) and/or thioridazine·HCl (100

µM) for 5 days. For enhancement of ROS in WT B cells, purified B cells were analyzed 3 d after

activation as for ROS scavenging, but vehicle, H$_2$O$_2$ (200 µM) or menadione (8 µM) was added

at three different time points (overnight, 3 hours, and 1 hour prior to processing for flow

cytometry to counting and measurements of DCFDA, annexin V, and C11 Bodipy signals.

For the analysis of proliferation in vitro, B cells were purified by depleting CD90.2$^+$ T

cells and CD138$^+$ cells using biotinylated anti-CD90.2 Ab and biotinylated anti-CD138 Ab

followed by streptavidin-conjugated microbeads (iMag^{TM}; BD Bioscience), labeled with 5 µM

CellTrace^{TM} Violet (CTV, Invitrogen), activated and cultured as above. To analyze proliferation

in vivo, B cells were purified, labeled with CTV, and injected intravenously into B cell deficient

μMT recipient mice. Mice were harvested at 4 days after adoptive transfer.

Flow cytometry - general and measurements of ROS, lipids peroxidation, and ER   GC B

cells were identified as $GL7^+ CD95^+$ events in the viable $B220^+ dump^-$ gate (dump channel

consisting of one fluorophore for CD11b, CD11c, F4/80, Gr1, TCRβ, and 7AAD) , while

plasmablasts were defined as $B220^+ CD138^+ TACI^+ dump^-$ cells. For flow analyses of total

intracellular ROS and mitochondrial superoxide, B cells (1×10^6 cells) were washed with PBS

and stained with 1.25 μM CM-H2DCFDA or 5 μM MitoSOX^{TM} Red in PBS (20 min at 37°C),

respectively, then washed with 1% BSA containing PBS, and further stained with anti-B220,

anti-CD19, anti-CD138, and Ghost-BV510. To measure the lipid peroxidation, cultured B cells

(1×10^6) were washed with PBS and stained with 1.25 μM C11-BODIPY^{TM} in PBS (20 min at

37°C), and washed with 1% BSA containing PBS followed by surface staining as above. To

analyze the level of endoplasmic reticulum (ER), B cells (1×10^6) were stained with 0.5 μM ER-

Tracker^{TM} Green (Molecular Probes, Eugene OR) in Hank's Balanced Salt Solution with calcium

and magnesium (HBSS/Ca/Mg; Gibco) for 30 min at 37°C.

For measurements of cell death (78), cultured B cells (1×10^6) were washed twice with PBS,

once with Annexin V binding buffer (10 mM HEPES, pH 7.4, 140 mM NaCl, 2.5 mM CaCl2),

and then incubated (15 min at 20 C in the dark) with Annexin V-rPE, 7AAD, APC-conjugated

CD138, APC-Cy7-conjugated B220, e450-CD19. Cells were then washed with Annexin V

binding buffer and analyzed on the flow cytometer. To measure levels of cleaved caspase-3 (25),

an activated apoptosis executor, the cultured cells were rinsed with PBS after harvest, stained (15

min at 4 C) with 7AAD, APC-conjugated CD138, APC-Cy7-conjugated B220, e450-CD19,

washed (FBS, 1% v/v in PBS), then fixed with paraformaldehyde (4% w/v, 10 min at 20 C),

permeabilized with permeabilization buffer (saponin, 0.2% w/v with FBS 1% in PBS), stained

with FITC-conjugated Ab specific for cleaved caspase-3 (BD Biosciences), rinsed, and analyzed

by flow cytometry.

Immunohistochemistry, MALDI-IMS, and IMS data analysis   Mice were immunized with

SRBC and spleens were harvested at 1 week after immunization. Spleens were snap frozen on

dry ice, and mounted to a cryostat chuck with a minimal amount of OCT (Thermo Fisher

Scientific, San Jose, CA). Frozen spleens were cut at 12 μm thickness using a Leica CM3050S

(Leica, IL, USA) at -20 ºC, fixed with 1% paraformaldehyde for 10 min, washed twice with PBS,

and blocked with M.O.M^{TM} (Vector Lab) followed by incubation with GL7-FITC, anti-IgD-PE,

and anti-CD35-biotin Ab followed by streptavidin-conjugated Alex647at 4 °C. After the tile

scanning of spleen sections, GC size and numbers were quantified with FIJI Image J software.

For Matrix-Assisted Laser Desorption/Ionization (MALDI) Imaging Mass Spectrometry

(IMS), frozen spleens were cut as above, and mounting onto indium tin oxide (ITO) coated

microscope slides (Delta Technologies ETC). Slides were then vacuum desiccated for at least 30

minutes before matrix application. Pre-IMS autofluorescence (AF) images were acquired prior to

matrix application on a Zeiss Axio Scan Z1 Slide Scanner using the brightfield channel and the

DAPI (ex. 340-380 nm; em. 435-485 nm, blue), EGFP (ex. 450-490 nm; em. 500-550 nm,

green), DsRed (ex. 538-562 nm; em. 570-640 nm, red) filter cubes. After taking AF images, a

custom in-house developed sublimation device was used to apply the matrices 2,5-

dihydroxyacetophenone (DHA) and 1,5-diaminonaphthalene (DAN) (Sigma Aldrich, St. Louis,

MO, USA) to tissue sections for positive and negative ion mode analyses, respectively (78). For

immunized AID-GFP mice, the fluorescence emissions identified GC localization within the

section used next for IMS. Alternatively, two serial sections of immunized mice spleens were

used, one for 2D-IMS and the contiguous section for immunohistochemistry (IHC). Fluorescence

data were registered with the 2D-IMS visualizations of phospholipids to align images (45),

allowing quantitation of the ion intensities in specific m/z peaks within defined micro-anatomic

portions of the spleen (Fig. 4B). MALDI IMS data were acquired with a 10 μm pixel size (laser

spot size 8 μm) in full scan mode using a Bruker trapped ion mobility time-of-flight (timsTOF)

Flex mass spectrometer (Bruker Daltonics Billerica, MA, USA). Data were acquired with 250

shots per pixel and a mass range of *m/z* 200-2000. SCiLS (software Bruker Daltonics) was used

to process the data, normalize ion intensity, visualize ion images, and merge the images.

LC-MS for lipidomics   To prepare samples, a one-phase method was used to extract lipids

(85). Briefly, 0.5 mL of MeOH/MTBE/CHCl$_3$ mix (1.3:1:1) was added to a frozen pellet of B-

cells (1x10^6 cells total), spiked with 10 uL EquiSPLASH-lipidomics internal standard mix

(Avanti Research), briefly vortexed and shaken gently for 20 min, followed by centrifugation at

20,000 x *g* for 15 min at 10°C. The supernatant was transferred to a clean Eppendorf tube,

evaporated under a gentle stream of N$_2$ gas, and resuspended in 100 uL methanol/CHCl$_3$ (9:1)

and 2uL were used for LC-HRMS (high-resolution MS) analysis. Each sample was injected two

times - one injection in positive ESI mode followed by one in negative mode. Pooled QCs were

injected to assess the performance of the LC and MS instruments at the beginning, in the middle

and at the end of each sequence. Discovery lipidomics data were acquired using a Vanquish

UHPLC (ultrahigh performance liquid chromatography) system interfaced to a Q Exactive HF

quadrupole/orbitrap mass spectrometer (Thermo Fisher Scientific).

Chromatographic separation was performed with a reverse-phase Acquity BEH C18 column

(1.7 mm, 2.1x150mm, Waters, Milford, MA) at a flow rate of 250 ul/min. Mobile phases were

made up of 10 mM ammonium formate and 0.1% formic acid in (A) H$_2$O/CH$_3$CN (40:60) and

in (B) CH$_3$CN/ iPrOH (10:90). Gradient conditions were as follows: 0–1 min, B = 20 %; 1–8

min, B = 20– 100 %; 8–10 min, B = 100 %; 10–10.5 min, B = 100–20 %; 10.5–15 min, B =

20%. The total chromatographic run time was 15 min. Mass spectra were acquired over a

precursor ion scan range of m/z 200 to 1,600 at a resolving power of 60,000 using the

following HESI-II source parameters: spray voltage 4 kV (3 kV in negative mode); capillary

temperature 250°C; S-lens RF level 60 V; N$_2$ sheath gas 40; N$_2$ auxiliary gas 10; auxiliary gas

temperature 350°C. MS/MS spectra were acquired for the top-seven most abundant precursor

ions with an MS/MS AGC target of 1e5, a maximum MS/MS injection time of 100 ms, and a

normalized collision energy of 15, 30, 40. High resolution mass spectrometry data were

processed with MS-DIAL version 4.70 in lipidomics mode (86). MS1, and MS2 tolerances

were set to 0.01 and 0.025 Da respectively. Minimum peak height was set to 30,000 to

decrease the number of false positive hits. Peaks were aligned on a quality control (QC)

reference file with RT tolerance of 0.1 min and mass tolerance of 0.015 Da. Default lipid

library was used (Msp20210527163602_converted.lbm2), solvent type was set to $HCOONH_4$

to match the solvent used for separation, and the identification score cut off was set to 80%.

All lipid classes were made available for the search. After lipid identification was completed,

MS-DIAL results were exported into Excel and cleaned using minimum RSD for QC samples

set to 20% and minimum ratio of QC to Blank set to 10. For species identified in both

PexRAP-depleted (*Dhrs7b* Δ/Δ) B cells and controls, mean levels (areas under peak curves)

and their variance were analyzed in Excel.

Metabolic flux analyses   Oxygen Consumption Rate (OCR) were measured using Seahorse

XF96 extracellular flux analyzer (Agilent Technology, Santa Clara, CA) as described previously

(30). Briefly, in vitro activated B cells were washed twice, resuspended in XF Base Media

(Agilent Technologies) supplemented with 2 mM L-glutamine, and equal numbers of B cells (2

$\times 10^5$) were plated on extracellular flux assay plates (Agilent Technologies) coated with 2.5

µg/mL CellTak (Corning) according to the manufacturer's protocol. Before extracellular flux

analysis, B cells were rested (25 minutes at 37°C, atmospheric CO_2) in XF Base Media. OCR

and ECAR were measured before and after the sequential addition of 1.5 µM oligomycin, 0.5

µM FCCP and 0.5 µM rotenone/antimycin A.

647
648

ACKNOWLEDGEMENTS

The experimental work was funded by NIH Grants to VUMC [R01 AI113292 (M.R.B), R01

HL106812 (M.R.B.), followed by R21 AI164760 (M.R.B.) and departmental funds]. Mass

spectrometry imaging was supported in part by NIH grant P41 GM103391 to Vanderbilt

University (R.M.C.). Additional support for M. A. J. was provided by the National Science

Foundation, NSF DGE-1445197. NIH Shared Instrumentation Grant 1S10OD018015 as well

as scholarships via the Cancer Center Support Grant (CA068485) and Diabetes Research Center

(DK0205930) helped defray costs of Vanderbilt Cores. We thank J. Cyster for generously

expediting shipment of *S1pr2*-CreERT2 breeding stock, and Vanderbilt institutional cores (High-

Throughput Screening; Flow Cytometry Shared Resource; Mass Spectrometry Shared Resource;

Cell & Developmental Biology) for equipment, expertise, and assistance.

661
662
References

1. Geyer CE, Mes L, Newling M, den Dunnen J, Hoepel W. 2021. Physiological and

pathological inflammation induced by antibodies and pentraxins. *Cells* **10:** 1175.

2. McGettigan SE, Debes GF. 2021. Immunoregulation by antibody secreting cells

in inflammation, infection, and cancer. *Immunol Rev*. **303:** 103-118.

3. Lu LL, Suscovich TJ, Fortune SM, Alter G. 2018. Beyond binding: antibody effector

functions in infectious diseases. *Nat Rev Immunol.* **18:** 46-61.

4. Allman D, Pillai S. 2008. Peripheral B cell subsets. *Curr Opin Immunol.* **20**: 149-157.

5. Martin F, Kearney JF. 2000. B-cell subsets and the mature preimmune repertoire. Marginal

zone and B1 B cells as part of a "natural immune memory". *Immunol Rev.* **175:** 70-79.

6. MacLennan, IC, Toellner KM, Cunningham AF, Serre K, Sze DM, Zuniga E, Cook MC,

Vinuesa CG. 2003. Extrafollicular antibody responses. *Immunol Rev* **194:** 8-18.

7. Hsu, M. C., K. M. Toellner, C. G. Vinuesa, and I. C. Maclennan. 2006. B cell clones that

sustain long-term plasmablast growth in T-independent extrafollicular antibody

responses. *Proc Natl Acad Sci USA* **103:** 5905-5510.

8. Rajewsky K. 1996. Clonal selection and learning in the antibody system. *Nature* **81:** 751-

758.

9. Boothby MR, Hodges E, Thomas JW. 2019. Molecular regulation of peripheral B cells and

their progeny in immunity. *Genes Dev.* **33:** 26-48.

10. Rubtsova K, Rubtsov AV, van Dyk LF, Kappler JW, Marrack P. 2013. T-box transcription

factor T-bet, a key player in a unique type of B-cell activation essential for effective viral

clearance. *Proc Natl Acad Sci USA* 110: E3216-24.

11. Alemán OR, Rosales C. 2023. Human neutrophil Fc gamma receptors: different buttons for

different responses. *J Leukoc Biol.* 114: 571-584.

12. Xu Z, Zan H, Pone EJ, Mai T, Casali P. 2012.Immunoglobulin class-switch recombination:

induction, targeting, and beyond. *Nat Rev Immunol.* **12:** 517-531.

13. Elsner RA, Shlomchik MJ. 2020. Germinal Center and extrafollicular B cell responses in

vaccination, immunity, and autoimmunity. *Immunity* **53:** 1136-1150.

14. Good-Jacobson, K. L., and M. J. Shlomchik. 2010. Plasticity and heterogeneity in the

generation of memory B cells and long-lived plasma cells: the influence of germinal

center interactions and dynamics. *J Immunol* **185:** 3117-3125.

15. Shlomchik M, Mascelli M, Shan H, Radic MZ, Pisetsky D, Marshak-Rothstein A, Weigert

M. 1990. Anti-DNA antibodies from autoimmune mice arise by clonal expansion and

somatic mutation. *J Exp Med.* **171:** 265-292.

16. Thomas JW, Hulbert C. 1996. Somatically mutated B cell pool provides precursors for

insulin antibodies. *J Immunol.* **157:** 763-771.

17. De Silva NS, Klein U. 2015. Dynamics of B cells in germinal centres. *Nat Rev Immunol.* **15:**

137-148.

18. Cyster JG, Allen CDC. 2019. B cell responses: Cell interaction dynamics and decisions. *Cell*

177: 524-540.

19. Victora, G. D., and M. C. Nussenzweig. 2022. Germinal Centers. *Annu Rev Immunol* **40:**

413-442.

20. Guttormsen HK, Sharpe AH, Chandraker AK, Brigtsen AK, Sayegh MH, Kasper DL. 1999.

Cognate stimulatory B-cell-T-cell interactions are critical for T-cell help recruited by

glyco-conjugate vaccines. *Infect Immun.* **67:** 6375-6384.

21. Kim W, Zhou JQ, Horvath SC, Schmitz AJ, Sturtz AJ, Lei T, Liu Z, Kalaidina E, Thapa M,

Alsoussi WB, Haile A, Klebert MK, Suessen T, Parra-Rodriguez L, Mudd PA, Whelan

SPJ, Middleton WD, Teefey SA, Pusic I, O'Halloran JA, Presti RM, Turner JS, Ellebedy

AH. 2022. Germinal centre-driven maturation of B cell response to mRNA vaccination.

Nature **604:** 141-145.

22. Roco JA, Mesin L, Binder SC, Nefzger C, Gonzalez-Figueroa P, Canete PF, Ellyard J, Shen

Q, Robert PA, Cappello J, Vohra H, Zhang Y, Nowosad CR, Schiepers A, Corcoran LM,

Toellner KM, Polo JM, Meyer-Hermann M, Victora GD, Vinuesa CG. 2019. Class-

switch recombination occurs infrequently in Germinal Centers. *Immunity* **51:** 337-350.

23. Gitlin AD, Shulman Z, Nussenzweig MC 2014. Clonal selection in the germinal centre by

regulated proliferation and hypermutation. *Nature* **509:** 637-40.

24. Vinuesa CG, Linterman MA, Yu D, MacLennan IC. 2016. Follicular helper T cells. *Annu*

Rev Immunol. **34:** 335-368.

25. Mayer CT, Gazumyan A, Kara EE, Gitlin AD, Golijanin J, Viant C, Pai J, Oliveira TY,

Wang Q, Escolano A, Medina-Ramirez M, Sanders RW, Nussenzweig MC. 2017. The

micro-anatomic segregation of selection by apoptosis in the germinal center. *Science* **358:**

eaao2602.

26. Chapman NM, Chi H. 2022. Metabolic adaptation of lymphocytes in immunity and disease.

Immunity **55:** 14-30.

27. Bacigalupa ZA, Landis MD, Rathmell JC. 2024. Nutrient inputs and social metabolic control

of T cell fate. *Cell Metab.* **36:** 10-20.

28. Jellusova J. 2020. Metabolic control of B cell immune responses. *Curr Opin Immunol*. **63:**

21-28.

29. Boothby MR, Brookens SK, Raybuck AL, Cho SH. 2022. Supplying the trip to antibody

production-nutrients, signaling, and the programming of cellular metabolism in the

mature B lineage. *Cell Mol Immunol.* **19:** 352-369.

30. O'Neill LA, Kishton RJ, Rathmell J. 2016. A guide to immunometabolism for

immunologists. *Nat Rev Immunol*. **16:** 553-565.

31. Cho, S. H., A. L. Raybuck, K. Stengel, M. Wei, T. C. Beck, E. Volanaki, J. W. Thomas, S.

Hiebert, V. H. Haase, and M. R. Boothby. 2016. Germinal centre hypoxia and regulation

of antibody qualities by a hypoxia response system. *Nature* **537:** 234-237.

32. Weisel, F. J., S. J. Mullett, R. A. Elsner, A. V. Menk, N. Trivedi, W. Luo, D. Wikenheiser,

W. F. Hawse, M. Chikina, S. Smita, L. J. Conter, S. M. Joachim, S. G. Wendell, M. J.

Jurczak, T. H. Winkler, G. M. Delgoffe, and M. J. Shlomchik. 2020. Germinal center B

cells selectively oxidize fatty acids for energy while conducting minimal glycolysis. *Nat*

Immunol **21:** 331-342.

33. Chen D, Wang Y, Manakkat Vijay GK, Fu S, Nash CW, Xu D, He D, Salomonis N, Singh H,

Xu H. 2021. Coupled analysis of transcriptome and BCR mutations reveals role of

OXPHOS in affinity maturation. *Nat Immunol.* 22: 904-913.

34. Urbanczyk S, Baris OR, Hofmann J, Taudte RV, Guegen N, Golombek F, Castiglione K,

Meng X, Bozec A, Thomas J, Weckwerth L, Mougiakakos D, Schulz SR, Schuh W,

Schlotzer-Schrerhardt U, Steinmetz TD, Brodesser S, Wiesner RJ, Mielenz D. 2022.

Mitochondrial respiration in B lymphocytes is essential for humoral immunity by

controlling the flux of the TCA cycle. *Cell Rep.* **39:** 110912.

35. Yazicioglu Y. F., E. Marin, C. Sandhu, S. Galiani, I. G. A. Raza, M. Ali, B. Kronsteiner, E.

B. Compeer, M. Attar, S. J. Dunachie, M. L. Dustin, and A. J. Clarke. 2023. Dynamic

mitochondrial transcription and translation in B cells control germinal center entry and

lymphomagenesis. *Nat Immunol.* **24:** 991-1006.

36. Brookens SK, Cho SH, Paik Y, Meyer K, Raybuck AL, Park C, Greenwood DL, Rathmell

JC, Boothby MR. 2024. Plasma cell differentiation, antibody quality, and initial germinal

center B cell population depend on glucose influx rate. *J Immunol.* **212:** 43-56.

37. Olenchock BA, Rathmell JC, Vander Heiden MG. 2017. Biochemical underpinnings of

immune cell metabolic phenotypes. *Immunity* **46:** 703-713.

38. Kong H, Chandel NS. 2018. Regulation of redox balance in cancer and T cells. *J Biol Chem*.

293: 7499-7507.

39. Addabbo F, Montagnani M, Goligorsky MS. 2009. Mitochondria and reactive oxygen

species. *Hypertension* **53:** 885-892.

40. Zorov DB, Juhaszova M, Sollott SJ. 2014. Mitochondrial reactive oxygen species (ROS) and

ROS-induced ROS release. *Physiol Rev.* **94:** 909-950.

41. Jellusova, J., M. H. Cato, J. R. Apgar, P. Ramezani-Rad, C. R. Leung, C. Chen, A. D.

Richardson, E. M. Conner, R. J. Benschop, J. R. Woodgett, and R. C. Rickert. 2017.

Gsk3 is a metabolic checkpoint regulator in B cells. *Nat Immunol* **18:** 303-312.

42. Li L, Feng C, Qin J, Li D, Liu M, Han S, Zheng B. 2021. Regulation of humoral immune

response by HIF-1α-dependent metabolic reprogramming of the germinal center reaction.

Cell Immunol. **367:** 104409.

43. Nakagawa R, Llorian M, Varsani-Brown S, Chakravarty P, Camarillo JM, Barry D, George

R, Blackledge NP, Duddy G, Kelleher NL, Klose RJ, Turner M, Calado DP. 2023. Epi-

microRNA mediated metabolic reprogramming ensures affinity maturation. *bioRxiv*

[Preprint]. 2023 Oct 31:2023.07.31.551250.

44. Jain IH, Calvo SE, Markhard AL, Skinner OS, To TL, Ast T, Mootha VK. 2020. Genetic

778   screen for cell fitness in high or low oxygen highlights mitochondrial and lipid

779   metabolism. *Cell* **181:** 716-727.e11.

780   45. Jones MA, Cho SH, Patterson NH, Van de Plas R, Spraggins JM, Boothby MR, Caprioli

781   RM. 2020. Discovering new lipidomic features using cell type-specific fluorophore

782   expression to provide spatial and biological specificity in a multimodal workflow with

783   MALDI Imaging Mass Spectrometry. *Anal Chem.* **92:** 7079-7086.

784   46. Lodhi IJ, Wei X, Yin L, Feng C, Adak S, Abou-Ezzi G, Hsu FF, Link DC, Semenkovich CF.

785   2015. Peroxisomal lipid synthesis regulates inflammation by sustaining neutrophil

786   membrane phospholipid composition and viability. *Cell Metab.* **21:** 51-64.

787   47. Lodhi IJ, Yin L, Jensen-Urstad AP, Funai K, Coleman T, Baird JH, El Ramahi MK, Razani

788   B, Song H, Fu-Hsu F, Turk J, Semenkovich CF. 2012. Inhibiting adipose tissue

789   lipogenesis reprograms thermogenesis and PPARγ activation to decrease diet-induced

790   obesity. *Cell Metab.* **16:** 189-201.

791   48. Dean JM, Lodhi IJ. 2018. Structural and functional roles of ether lipids. *Protein Cell* **9:** 196-

792   206.

793   49. Honsho M, Fujiki Y. 2023. Regulation of plasmalogen biosynthesis in mammalian cells and

794   tissues. *Brain Res Bull.* **194:** 118-123.

795   50. Werner ER, Swinkels D, Juric V, Dorninger F, Baes M, Keller MA, Berger J, Watschinger

796   K. 2023. Normal plasmalogen levels are maintained in tissues from mice with

797   hepatocyte-specific deletion in peroxin 5. *Brain Res Bull.* **193:** 158-165.

798   51. Dorninger F, Werner ER, Berger J, Watschinger K. 2022. Regulation of plasmalogen

799   metabolism and traffic in mammals: The fog begins to lift. *Front Cell Dev Biol.* **10:**

800   946393.

52. Kuerschner L, Richter D, Hannibal-Bach HK, Gaebler A, Shevchenko A, Ejsing CS, Thiele

C. 2012. Exogenous ether lipids predominantly target mitochondria. *PLoS One* **7:**

e31342.

53. Lodhi IJ, Link DC, Semenkovich CF. 2015. Acute ether lipid deficiency affects neutrophil

biology in mice. *Cell Metab.* **21:** 652-653.

54. Dorninger F, Wiesinger C, Braverman NE, Forss-Petter S, Berger J. 2015. Ether lipid

deficiency does not cause neutropenia or leukopenia in mice and men. *Cell Metab.* **21:**

650-651.

55. Cui W, Liu D, Gu W, Chu B. 2021. Peroxisome-driven ether-linked phospholipids

biosynthesis is essential for ferroptosis. *Cell Death Differ.* **28:** 2536-2551.

56. Honsho M, Tanaka M, Zoeller RA, Fujiki Y. 2020. Distinct functions of acyl/alkyl

dihydroxyacetonephosphate reductase in peroxisomes and endoplasmic reticulum. *Front*

Cell Dev Biol. **8:** 855.

57. Braverman NE, Moser AB. 2012. Functions of plasmalogen lipids in health and disease.

Biochim Biophys Acta **1822:** 1442-1452.

58. Gorgas K, Teigler A, Komljenovic D, Just WW. 2006. The ether lipid-deficient mouse:

tracking down plasmalogen functions. *Biochim Biophys Acta* **1763:** 1511-1526.

59. Jiménez-Rojo N, Riezman H. 2019. On the road to unraveling the molecular functions of

ether lipids. *FEBS Lett.* **93:** 2378-2389.

60. Rangholia N, Leisner TM, Holly SP. 2021. Bioactive ether lipids: Primordial modulators of

cellular signaling. *Metabolites* **11:** 41.

61. Bozelli JC Jr, Azher S, Epand RM. 2021. Plasmalogens and chronic inflammatory diseases.

Front Physiol. **12:** 730829.

62. Wallner S, Schmitz G. 2011. Plasmalogens the neglected regulatory and scavenging lipid

species. *Chem Phys Lipids* **164:** 573-589.

63. Friedrich, G, Soriano, P. 1991. Promoter traps in embryonic stem cells: A genetic screen to

827   identify and mutate developmental genes in mice. *Genes Devel*. **5:** 1513–1523.

828   64. Shapiro-Shelef M, Lin KI, Savitsky D, Liao J, Calame K. 2005. Blimp-1 is required for

829   maintenance of long-lived plasma cells in the bone marrow. *J Exp Med.* **202:** 1471-1476.

830   65. Higashi AY, Ikawa T, Muramatsu M, Economides AN, Niwa A, Okuda T, Murphy AJ, Rojas

831   J, Heike T, Nakahata T, Kawamoto H, Kita T, Yanagita M. 2009. Direct hematological

832   toxicity and illegitimate chromosomal recombination caused by the systemic activation of

833   CreERT2. *J Immunol.* **182:** 5633-5640.

834   66. Uhmann A, Dittmann K, Wienands J, Hahn H. 2009. Comment on "Direct hematological

835   toxicity and illegitimate recombination caused by the systemic activation of CreERT2". *J*

836   *Immunol.* **183:**2891.

837   67. Khalil AM, Cambier JC, Shlomchik MJ. 2012. B cell receptor signal transduction in the GC

838   is short-circuited by high phosphatase activity. *Science* **336:** 1178-1181.

839   68. Inoue T, Kurosaki T. 2024. Memory B cells. *Nat Rev Immunol.* **24:** 5-17.

840   69. Shinnakasu R, Inoue T, Kometani K, Moriyama S, Adachi Y, Nakayama M,

841   Takahashi Y, Fukuyama H, Okada T, Kurosaki T. 2016. Regulated selection of germinal-

842   center cells into the memory B cell compartment. *Nat Immunol.* **17:** 861-869.

843   70. Price MJ, Patterson DG, Scharer CD, Boss JM. 2018. Progressive upregulation of oxidative

844   metabolism facilitates plasmablast differentiation to a T-independent antigen.

845   *Cell Rep.* **23:** 3152-3159.

846   71. Capasso M, Bhamrah MK, Henley T, Boyd RS, Langlais C, Cain K, Dinsdale D, Pulford K,

847   Khan M, Musset B, Cherny VV, Morgan D, Gascoyne RD, Vigorito E, DeCoursey TE,

848   MacLennan IC, Dyer MJ. 2010. MacLennan IC, Dyer MJ. 2010. HVCN1 modulates

849   BCR signal strength via regulation of BCR-dependent generation of reactive oxygen

850   species. *Nat Immunol*. **11:** 265-272.

851   72. Wheeler ML, Defranco AL. 2012. Prolonged production of reactive oxygen species in

852   response to B cell receptor stimulation promotes B cell activation and proliferation. *J*

853   *Immunol.* **189:** 4405-4416.

854   73. Tsubata T. 2020. Involvement of reactive oxygen species (ROS) in BCR signaling as a

855   second messenger. *Adv Exp Med Biol.* **1254:** 37-46.

856   74. Jiang X, Stockwell BR, Conrad M. 2021. Ferroptosis: mechanisms, biology and role in

857   disease. *Nat Rev Mol Cell Biol.* **22:** 266-282.

858   75. Nordgren M, Fransen M. 2014. Peroxisomal metabolism and oxidative stress. *Biochimie* **98:**

859   56-62.

860   76. Liang D, Minikes AM, Jiang X. 2022. Ferroptosis at the intersection of lipid metabolism and

861   cellular signaling. *Mol Cell* **82:** 2215-2227.

862   77. Facciotti F, Ramanjaneyulu GS, Lepore M, Sansano S, Cavallari M, Kistowska M, Forss-

863   Petter S, Ni G, Colone A, Singhal A, Berger J, Xia C, Mori L, De Libero G. 2012.

864   Peroxisome-derived lipids are self antigens that stimulate invariant natural killer T cells

865   in the thymus. *Nat Immunol.* 3: 474-480.

866   78. Hossain MS, Mawatari S, Fujino T. 2022. Plasmalogen-mediated activation of GPCR21

867   regulates cytolytic activity of NK cells against the target cells. *J Immunol.* **209:** 310-325.

868   79. Rossi M, Salomon A, Chaumontel N, Molet J, Bailly S, Tillet E, Bouvard C. 2023. Warning

869   regarding direct hematological toxicity of tamoxifen-activated CreERT2 in young

870   Rosa26-CreERT2 mice. *Sci Rep.* **13:** 5976.

871   80. Lodhi IJ, Semenkovich CF. 2014. Peroxisomes: a nexus for lipid metabolism and cellular

872   signaling. *Cell Metab.* **19:** 380-392.

873   81. He A, Dean JM, Lodhi IJ. 2021. Peroxisomes as cellular adaptors to metabolic and

874   environmental stress. *Trends Cell Biol.* **31:** 656-670.

82. Zou Y, Henry WS, Ricq EL, Graham ET, Phadnis VV, Maretich P, Paradkar S, Boehnke N,

Deik AA, Reinhardt F, Eaton JK, Ferguson B, Wang W, Fairman J, Keys HR, Dančík V,

Clish CB, Clemons PA, Hammond PT, Boyer LA, Weinberg RA, Schreiber SL. 2020.

Plasticity of ether lipids promotes ferroptosis susceptibility and evasion. *Nature* **585:**

603-608.

83. Esselman AB, Patterson NH, Migas LG, Dufresne M, Djambazova KV, Colley ME, Van de

Plas R, Spraggins JM. 2023. Microscopy-Directed Imaging Mass Spectrometry for Rapid

High Spatial Resolution Molecular Imaging of Glomeruli. *J Am Soc Mass Spectrom.* **34:**

1305-1314.

84. Lee K, Heffington L, Jellusova J, Nam KT, Raybuck A, Cho SH, Thomas JW, Rickert

RC, Boothby M. 2013. Requirement for Rictor in homeostasis and function of mature B

lymphoid cells. *Blood* **122:** 2369-2379.

85. Pellegrino RM, Di Veroli A, Valeri A, Goracci L, Cruciani G. 2014. LC/MS lipid profiling

from human serum: a new method for global lipid extraction. *Anal Bioanal Chem.* **406:**

7937-7948.

86. Tsugawa H, Ikeda K, Takahashi M, Satoh A, Mori Y, Uchino H, Okahashi N, Yamada Y,

Tada I, Bonini P, Higashi Y, Okazaki Y, Zhou Z, Zhu Z-J, Koelmel J, Cajka T, Fiehn O,

Saito K, Arita M, Arita M. 2020. A lipidome atlas in MS-DIAL 4. *Nat Biotechnol.* **38:**

1159-1163.

87. Tao L, Lemoff A, Wang G, Zarek C, Lowe A, Yan N, Reese TA. 2020. Reactive oxygen

species oxidize STING and suppress interferon production. *Elife* **9:** e57837. doi:

10.7554/eLife.57837.

     Figure Legends

Figure 1. PexRAP promotes GC response. (A) Schematic of immunization with SRBC after

inactivation of *Dhrs7b* in adult mice. Mice (*Rosa26*-CreER^{T2}; *Dhrs7b$^{+/+}$, or Rosa26*-CreER^{T2};

Dhrs7b$^{f/f}$) were treated with tamoxifen, immunized with SRBC, and harvested 1 wk after

immunization, as described in *Materials and Methods*. (B) Deletion efficiency of *Dhrs7b*

conditional alleles. B and T lymphocytes were isolated from spleens of *Rosa26*-CreER^{T2} or

Dhrs7b$^{f/f}$;Rosa26-CreER^{T2} mice after in vivo tamoxifen injections followed by immunization

with SRBC. WT and *Dhrs7b$^{\Delta/\Delta}$* were analyzed by immunoblotting with antibodies directed

against PexRAP (protein product of *Dhrs7b*) and actin (internal loading control). (C, D) PexRAP

acutely regulates B cell numbers. Representative flow plots of viable splenic B cells (C) and

aggregate data for three biologically independent replicate experiments (D) (n = 8 WT and 7

cKO). For data on T cells, see supplemental data Fig. S1A-C. (E, F) Effect of PexRAP on GC B

cell response. Flow plots of GL7^{+} CD95^{+} GC B cells among viable B cells (E), and aggregated

frequencies and numbers of GC B cells, as indicated, in the three replicate experiments (F). (G,

H) PexRAP impact on GC response. After tamoxifen injections, *Dhrs7b* deficient mice [*Rosa26*-

CreER^{T2}; *Dhrs7b$^{f/f}$* (46)] and WT controls (*Rosa26*-CreER^{T2}) were immunized with SRBC and

analyzed 7 d later, as in Fig. 1A-F. Data on gating strategy, GC counts per unit area and LZ area

are in Supplemental data Fig. 1D-F. (G, H) Shown are representative images from

immunofluorescence staining of spleens with the indicated Ab in two independent experiments

(5 WT vs 4 *Dhrs7b* cKO), showing a low-power overview with many follicles (G) and a

representative higher-magnification image to better delineate primary and secondary follicles

(H). (I) Quantitation of number (left panel) and size (right panel) of GC in spleen sections, with

GL7-positive, IgD-negative areas that include both CD35+ (LZ) and CD35- (DZ) areas. Mann-

Whitney U test was used to calculate p values.

Figure 2. PexRAP promotes proliferation of B cells. (A) Schematic of *Dhrs7b* inactivation in

mature mice via tamoxifen treatment and a B cell type-specific conditional allele. Mice

(*huCD20*-CreER^{T2}; *Dhrs7b$^{+/+}$, or huCD20*-CreER^{T2}; *Dhrs7b$^{f/f}$*) were injected with tamoxifen (d1,

3, 5) and harvested at day 10. (B) Deletion efficiency of conditional *Dhrs7b* alleles. After in vivo

tamoxifen injections, B cells were isolated from spleens of *hCD20*-CreER^{T2} or *Dhrs7b$^{f/f}$;*

hCD20-CreER^{T2} mice, as indicated, and analyzed by immunoblotting as in Fig. 1B. (C) PexRAP

and the maintenance of B cells. Shown are the frequencies and the numbers of CD19$^+$ B220$^+$ B

cells among viable lymphocytes in spleen (left and right panels, respectively). Data are pooled

from four independent replicate experiments (n = 12 WT and 12 cKO). Shown in feather plots

are the means, with whiskers that extend to the minimum and maximum values and boxes that

outline upper and lower quartile values with the midline identifying the median.(D-F) PexRAP

regulates B cell proliferation in vivo. CTV-labeled B cells were adoptively transferred into μMT

recipient mice and analyzed 4 days thereafter. Shown are the frequencies of B220$^+$ CD19$^+$ events

among splenocytes in the viable cell gate (D), along with representative flow plots of CTV

partitioning and surface IgD in B cell gates (E) and aggregated frequencies of divided B cells

from five independent replicate experiments (F) (n =10 WT and 11 cKO). Additional data on the

IgD^{neg} population are in supplemental Fig.1J. (G-I) PexRAP promotes B cell proliferation in

vitro. After in vivo tamoxifen injections, bead-purified B cells from spleens of CreER^{T2} mice

(WT and *Dhrs7b$^{Δ/Δ}$, i.e., cKO*) were stained with Cell Trace Violet (CTV), activated and

cultured 4 days in anti-CD40, BAFF, IL-4, IL-5 and 4-hydroxytamoxifen in three biologically

independent experiments totaling 6 WT and 7 cKO mice). (G) Representative flow-cytometric

analysis of CTV partitioning. (H) Quantified frequencies of divided >3 times. (I) Numbers of B

cells recovered at the end of the cultures.

Figure 3. B cell expression of PexRAP is essential for normal concentrations of most ether

phospholipids. Splenic B cells (CreER^{T2+} or CreER^{T2+}, *Dhrs7b* Δ/Δ) from tamoxifen-injected

mice were analyzed by LC-MS-MS either directly ex vivo ("Naive") or after activation and

culture (48 h) ("Activated") as in Fig. 2. Shown are z-scored heat maps for the relative

abundance of subsets of ether and plasmalogen phospholipids and lysophospholipids identified

by exact mass and secondary fragmentation in (A) positive and/or (B) negative ion modes. More

species are displayed in Supplemental Fig. 2. (C) Quantitated peak areas for activation-induced

increases ('fold-induction' ratios of activated/resting) for the indicated ether, plasmalogen, and

diacyl (non-ether) phospholipids in the PexRAP-sufficient and -depleted B cells. * denotes

species for which P<0.05 for the effect of *Dhrs7b* inactivation. (D) Shown are the raw values of

mean peak areas for the indicated species in the freshly purified and the activated B cells of the

indicated genotypes, as indicated. (Three independent samples from individual mice of each

genotype were analyzed.) * p<0.05, ** p<0.01 by unpaired Student's. t-test. Additional data

including controls comparing B cells of unmanipulated B6 mice to those expressing CreER^{T2}

and injected with tamoxifen are in Supplemental Fig. 2.

Figure 4. PexRAP is essential for normal concentrations and distributions of some ether

phospholipids in splenic follicles. (A) Schematic of immunization for IMS analyses. Mice of

the indicated genotypes were treated with tamoxifen, immunized with SRBC, and harvested 1

wk after immunization. (B) Schematic diagram illustrating the 2D-IMS analysis work flow,

image merging and quantitation. Harvested spleens were used to generate serial tissue sections

followed by immunofluorescence (IF) staining of one section and IMS analysis with the adjacent

one. IF and IMS images were aligned to map ion intensity distributions to microanatomic regions

(B cell follicle and GC). The intensities of specific ions on B cell follicles and GC regions were

quantitated as described in *Materials and Methods.* (C) Identification of ether lipid species

localizing to lymphoid follicles. Representative ion images of two ions [*m/z* 752.5545, and *m/z*

776.5556] with spleens from immunized mice (WT and *Dhrs7b$^{Δ/Δ}$*). (D) Identification of ether

lipid species localizing to lymphoid follicles. Representative ion images of three ions [*m/z*

752.5545, *m/z* 872.5749, and *m/z* 700.4958] in mass spectrometry imaging of spleens from WT

and *Dhrs7b$^{Δ/Δ−B}$* mice. Immunofluorescent images at higher magnification, delineating LZ and

DZ marked by CD35 staining, along with quantification of sizes of GC and their LZ and DZ, are

shown in supplemental data Fig. S4A, B. (E) Shown in the bar graphs are the mean (±SEM) ion

intensities in primary lymphoid follicles (B cell zones) and GC from spleens of WT and

Dhrs7b$^{Δ/Δ−B}$ mice, immunized or not ("UI") as indicated. Median intensity of each ion was

obtained from 3 follicles / spleen and 3 GC / spleen from WT and *Dhrs7b$^{Δ/Δ−B}$* mice (three

biological replications comprising 7 WT and 6 cKO spleens). P values were calculated by Mann-

Whitney U test.

Figure 5. B cell intrinsic role of PexRAP in GC response. (A) Schematic of immunization

with NP-OVA in alum after inactivation of *Dhrs7b* in B lineage cells. Tamoxifen-treated WT

(*hCD20*-CreER^{T2}; *Dhrs7b$^{+/+}$*) or *Dhrs7b$^{Δ/Δ−B}$* mice *(hCD20*-CreER^{T2}; *Dhrs7b^{ff}*) were immunized

with NP-OVA, boosted with NP-OVA 3 wk thereafter, and harvested 1 wk after the 2nd

immunization. (B, C) Representative flow plots of splenic IgD^{neg} B cells (B) and aggregate data

(C) for two replicate experiments (n = 5 WT and 5 cKO). The cocktail of reagents for the dump

channel included anti-IgD. (D, E) B cell-intrinsic function of PexRAP in GC B cell response.

Representative flow plots of GL7$^+$ CD95$^+$ GC B cells among viable IgD^{neg} B cells (D), and

aggregated frequencies and numbers of GC B cells for two replicate experiments (E). Mann-

Whitney U test was used to calculate p values.

Figure 6. Ab response and affinity increase promoted by PexRAP in B cells. Tamoxifen-

treated mice (*hCD20*-CreER^{T2}; *Dhrs7b$^{+/+}$* or *hCD20*-CreER^{T2}; *Dhrs7b^{ff}* , i.e., *Dhrs7b$^{\Delta/\Delta-B}$*) were

immunized as in Fig. 5, with venous blood collected to measure Ag-specific Ab just prior to a

second immunization, followed by harvesting a week thereafter. (A-D) Ag-specific Ab in

primary response sera, prior to the boost. Shown are the all- (NP$_{20}$) and high- (NP$_2$) -affinity anti-

NP IgM (A, B) and IgG1 (C, D), as indicated, with NP$_{20}$ a high hapten density to detect both

low- and high-affinity Ab and NP$_2$ a low hapten density selective for high-affinity Ab.

(E, F) Levels of anti-NP IgM detected using NP$_{20}$ and NP$_2$ for ELISpot (F) and ELISA (G) as

described in *Materials and Methods*. Graphs show mean (± SEM) number of Ab-secreting cells

in spleen (E) and (F) mean (± SEM) OD$_{450nm}$ in measurement of NP-specific IgM in serial

dilutions of sera. P values were calculated by students' t-test. * indicates p<0.05, and **

indicates p<0.01. (G) PexRAP regulates Ab affinity maturation. The bar graph shows the mean

(± SEM) ratios of high-affinity to all-affinity NP-specific IgM Ab in sera of individual mice

(each dot representing one subject) using OD$_{450nm}$ values at the 1:2000 dilution, with data from

three independent experiments comprising 8 mice of each type (WT; *Dhrs7b$^{\Delta/\Delta-B}$*). P values were

calculated by Mann-Whitney U test. (H, I) PexRAP promotes CD138$^+$ cell differentiation in

vivo. B cells were adoptively transferred into μMT recipient mice and analyzed at 4 days after

transfer. Representative flow plot of CD138 and cD19 expression (H) and aggregated

frequencies of CD138$^+$ CD19$^+$ cells in the viable cell gate (I) from four independent replicate

experiments. To test for a potential distortion arising from outlier values, statistical testing was

performed both with and without their inclusion.

1018

Figure 7. Function of PexRAP in GC response. (A) Schematic of the time line of

immunization followed later by tamoxifen injections into *S1pr2*-CreER^{T2} mice (*Dhrs7b$^{+/+}$*or

Dhrs7b^{f/f}). Note that deletion is initiated after the germinal center reaction starts (~ 3.5 d post-

immunization). Mice (*S1pr2*-CreER^{T2}; *Dhrs7b^{+/+}* or *S1pr2*-CreER^{T2}; *Dhrs7b^{f/f}*) were immunized

with NP-OVA, treated with tamoxifen on days 3, 5, and 7 after NP-OVA immunization (36) and

harvested at day 9. (B) Representative flow plots of GL7$^+$ CD95$^+$ GC B cells among viable B

cells and (C) aggregated mean (±SEM) frequencies of GC B cells from three replicate

experiments (7 WT; 9 cKO mice). Mann-Whitney U test was used to calculate p values. (D, E)

Effect of PexRAP on the level of plasmablasts in spleen. Representative flow plots of B220$^+$

CD19$^+$ plasmablasts among the events in a dump$^-$ CD138$^+$ TACI$^+$ gate (D) and aggregated mean

(±SEM) frequencies of these plasmablasts (E).

1030

Figure 8. PexRAP contributes to ROS homeostasis and B cell population growth in vitro.

(A-D) PexRAP is critical for maintenance of normal ROS levels. Bead-purified B cells from

spleens of tamoxifen-treated *huCD20*-CreER^{T2} mice (*Dhrs7b^{Δ/Δ−B}* and *Dhrs7b^{+/+}*) were cultured

3 days in anti-CD40, BAFF, IL-4, IL-5 and 4-hydroxytamoxifen. Total cellular (A, B) and

mitochondrial ROS, mtROS (C, D) in B lymphoblasts were determined by flow cytometry after

staining with surface markers and H$_2$DCFDA and MitoSOX, as described in the Methods.

Representative histogram image of H$_2$DCFDA (A) and MitoSOX (C) in the B cell gate, and

aggregated mean (± SEM) geometric MFI of H$_2$DCFDA (B) and MitoSOX (D) from 3

independent experiments, each using two mice of each type (6 WT; 6 cKO). P values were

calculated by Mann-Whitney U test. (E) PexRAP restrains lipid peroxidation. B cells were

activated and cultured as in (A). A representative result of flow cytometric analyses of lipid

peroxidation assayed using C11-Bodipy is shown, based on three independent experiments.

(F, G) PexRAP promotes B cell survival. (F) Shown are the mean (±SEM) frequencies of total

viable 'events' (by FSC, SSC, and 7-AAD exclusion) in flow cytometry (filled circles) and MFI

of Bodipy-C11 (open squares) after exposure to H$_2$O$_2$ (200 µM). (*, ** - p=0.03 and 0.003,

respectively). (G-I) WT and *Dhrs7b^{Δ/Δ}* B cells were cultured (2 d) in anti-CD40, BAFF, IL-4, IL-

5 and 4-hydroxytamoxifen. (G) Increased cleaved caspase 3 (CC3) in PexRAP-deficient cells

generated in vitro. Cleaved caspase 3 in B cells was detected by intracellular staining and flow

cytometry. Shown are a representative pair of histograms for WT and *Dhrs7b^{Δ/Δ}* B cells as

indicated (left panel) and a dot graph aggregating all experiments' outcomes (right panel).

(H) Frequencies of apoptotic B cells in cultures as in (A-D) were scored for annexin V and

7AAD as described (84). Shown are representative data of flow plots in the lymphocyte gate

(left panel) and a dot graph aggregating all experiments' outcomes (right panel). (I, J) PexRAP

and ROS control promote B cell population growth in vitro. WT and *Dhrs7b*$^{\Delta/\Delta}$ B cells were

cultured 5 days in anti-CD40, BAFF, IL-4, IL-5 and 4-hydroxytamoxifen in the presence or

absence of ROS scavenger NAC (1 mM vs 5 mM) (H). (I) B cells activated as in (A-G) were

cultured 5 d in the presence or absence of NAC (1 mM) or thioridazine (100 nM). The bar graphs

show the mean (± SEM) recovered cell number from three independent experiments, each with

two independent samples of each genotype (WT; *Dhrs7b*$^{\Delta/\Delta}$).

1060

Figure 9. PexRAP deficiency in activated B cells reduces mitochondrial metabolism and ER

mass. (A-G) Purified B cells were activated with and cultured (2 d) in αCD40, BAFF, IL-4, IL-5,

and 4-OHT, then divided and analyzed using a Seahorse XFe96. (A) Shown are aggregated

results from three independent experiments measuring oxygen consumption rate (OCR)

quantified via metabolic flux analysis during mitochondrial stress testing. (B) Basal OCR, (C)

maximal OCR, (D) ATP-production coupled respiration, (E) proton leak, and (F) spare

respiratory capacity of WT and *Dhrs7b*$^{\Delta/\Delta}$ B cells assayed in (A). Data points of individual

samples are show (each individual dot), as well as bars to display mean (±SEM) values. T-tests

with Welch's correction were used to calculate p-values. *, $p<0.05$; ***, $p<0.001$. (G)

Extracellular acidification rate (ECAR) of B cells cultured as described previously (36) and

assessed using the glycolytic stress test. (H, I) PexRAP influences ER mass. WT and *Dhrs7b*$^{\Delta/\Delta}$

B cells were cultured 5 days in anti-CD40, BAFF, IL-4, IL-5 and 4-hydroxytamoxifen followed

by the staining with ER-Tracker Green. Representative histogram of ER-Tracker Green in viable

cells (H) and aggregated MFI (±SEM) of ER-Tracker Green from three replicate experiments (I).

Mann-Whitney U test was used to calculate p values.

1076

1077

 SUPPLEMENTAL INFORMATION

 Supplemental Figures

 Supplemental Figure 1. PexRAP is dispensable for T cell homeostasis and does not affect

 balance among pre-immune B cell subsets. Mice ($Rosa26$-CreER^{T2}; $Dhrs7b^{+/+}$, or $Rosa26$-

 CreER^{T2}; $Dhrs7b^{f/f}$) were treated with tamoxifen and immunized with SRBC as described in

 Materials and Methods, and 1 wk after immunization. (A-C) PexRAP is dispensable for

 homeostatic maintenance of T cells. Shown are the representative flow plots of splenic T cells

 (A), mean (± SEM) frequencies of TCRβ^{+} CD4^{+} T cells (B), and CD8 (TCRβ^{+} CD4$^{-)}$ T cells (C)

 from WT and cKO mice in three replicate experiments totaling 8 WT and 7 cKO mice. (D) A

 representative gating scheme illustrating the flow-cytometric determination of the GC B cell

 population in mice. (E) Numbers of GC identified by microscopy of immunofluorescently

 stained sections of immunized mice as in Fig. 1. (F) A representative gating scheme illustrating

 the flow-cytometric estimation of the MZ, FO, and B1 B cell populations in mice. (G) Shown are

 the mean (±SEM) frequencies of CD19^{+} IgM^{+} CD23^{-} MZB cells (left panel) and CD19^{+} IgM^{-}

 CD23^{+} FOB cells (righthand panel) in spleen, and (H) CD19^{+} CD23^{lo} CD43^{+} CD5^{+} B1B cells in

 peritoneal cavity. (I) In vivo generation of IgD^{-} progeny from four independent replicate

 experiments (n =8 WT and 9 cKO). Shown are mean (±SEM) frequencies of IgD^{-} B cells as

 aggregate data for the frequencies of IgD- events in the gate for viable B cells recovered from

 μMT recipient mice after transfer of WT or PexRAP-depleted B cells (experiments of Fig. 2D-

 F), as indicated, with representative flow plots in Fig. 2E.

 Supplemental Figure 2. Activation of B cells induces PexRAP-dependent increases in their

 ether and plasmalogen P-lipids. As in Fig. 3, B lymphocytes of the indicated *Dhrs7b* genotypes

 were prepared from individual tamoxifen-injected mice and divided to analyze without activation

 ("naive") or after activation and culture (48 h) ("activated"). Shown are heat maps with more

 extensive lists of ether phospholipids and plasmalogens identified in the LC-MS-MS analyses for

 both genotypes in positive (A) and negative (B) ion modes. (C, D) LC-MS-MS results for lipids

 in B cells, activated or not, as indicated, of conventional non-transgenic B6 mice not injected

 with tamoxifen and those of tamoxifen-injected CreER^{T2+} mice (the "+/+" samples of panels A,

 B and of Figure 3). Bar graphs in (C) and (D) show results of these control comparisons for some

 of the phospholipids analyzed in prior figures panels (Fig. 3A, B).

Supplemental Figure 3. Immunization induces increases in P-lipids of both primary and

secondary follicles dependent on PexRAP in B cells. (A, B) Selected representative m/z

features Spleens of AID-GFP were harvested at 7 days after immunization with SRBC and

analyzed for exact mass signatures of ether lipid species in (A) negative and (B) positive ion

modes. (C) Spleens of immunized mice (tamoxifen-treated $hCD20$-CreER^{T2}; $Dhrs7b^{f/f}$ and

$hCD20$-CreER^{T2} controls) were analyzed by IMS. Shown are the positive ions presented in (B)

for GC vs follicles of immunized AID-GFP mice. The bar graph shows the mean ($\pm$ SEM) ion

intensities in lymphoid follicles and GC from WT and $Dhrs7b^{\Delta/\Delta-B}$ spleens. Median intensity of

each ion was obtained from three follicles per spleen and three GC per spleen from WT and

$Dhrs7b^{\Delta/\Delta-B}$ mice (three biological replication experiments totaling seven WT and six cKO

subjects). P values were calculated by Mann-Whitney U test.

1121

Supplemental Figure 4. A subset of ether phospholipids in primary follicles and GC

(secondary follicles) depend on PexRAP in B cells. (A) As in Fig. 4, $Dhrs7b$ was inactivated

by tamoxifen injections into mice bearing the hCD20-CreER^{T2} transgene, followed by

immunization with SRBC and analysis. Shown are representative higher-magnification images

defining the primary (IgD$^+$) and secondary (IgD$^{lo/neg}$, GL7$^+$) follicles and the identification of

light (CD35$^+$) and dark (CD35^{neg}) zones. (B) The sizes of GC and their light zones (LZ) after

immunization were quantified. Spleens of immunized mice (tamoxifen-treated $hCD20$-CreER^{T2};

$Dhrs7b^{f/f}$ and $hCD20$-CreER^{T2} controls, as indicated) were analyzed by immunofluorescence

staining as in (A). Each dot represents one follicle in a given mouse. Totals of three GC per

spleen from immunized mice were used were used, with seven WT and five $Dhrs7b^{\Delta/\Delta-B}$.

(C, D) Mean ($\pm$SEM) absorbances of ELISA performed across the indicated dilutions of sera

from the tamoxifen-treated mice ($hCD20$-CreER^{T2}; $Dhrs7b^{+/+}$ or $hCD20$-CreER^{T2}; $Dhrs7b^{f/f}$,

i.e., $Dhrs7b^{\Delta/\Delta-B}$) of Fig. 6. Shown are the results for (C) all- (NP$_{20}$ captured) and (D) high- (NP$_2$

captured) -affinity anti-NP Ab of the IgG1 and IgG2c isotypes, as indicated. (E) Proliferation and

development of CD138$^+$ progeny. Shown are representative flow plots from three independent

experiments with six WT and seven $Dhrs7b^{\Delta/\Delta-B}$ mice.

1138

Supplemental Figure 5. Distinct outcomes of heightened ROS elicited by H$_2$O$_2$ versus

menadione. (A-B) Induction of ROS independent from B cell activation. Shown are

representative flow plots of DCFDA in the viable B cell gate after exposure (1 h) (A) to H$_2$O$_2$

(200 μM) or (B) menadione (8 μM), as in (87). (C-D) Representative flow plots of annexin V vs

7-AAD for identification of early-apoptotic B cells. Cultured B cells were analyzed 3 hr after

treatment with (C) 200 μM H_2O_2 or (D) 8 μM menadione. Inset numbers represent the fraction

of B cells that are Annexin V^+ 7-AAD^{neg} or Annexin V^{neg} 7-AAD$^+$. (E-G) Shown are the results

of three independent experiments to measure (E) total cell viability, (F) BODIPY C11 MFI and

(G) frequencies of Annexin V^+ 7-AAD^{neg} frequency in B cells treated (8 μM) menadione

overnight (E) or for 3 h (D). P-values were calculated using the unpaired Student's t-test.

1149

1150

Supplemental Table 1. Immunization-induced increases of selected ether lipids in GC.

Shown are (a) a sample of m/z features characteristic of the indicated phospholipid species

identified by the LIPIDMAPS database, and (b) the mean ±SEM ion intensity / counts in IMS

data generated from spleens of AID-GFP mice, immunized or not (UI), after mapping to the

indicated regions of interest (primary follicles or GC, i.e. secondary follicles) using fluorescent

images. Shown are ions more accumulated in GC regions compared to primary follicles (8 ions

from negative ion mode, and 5 ions from positive ion mode, all p<0.05 for comparison of 1^0

follicle to UI (c), or GC to 1^0 follicle (d). P values were calculated by Mann-Whitney U test.

1159

Supplemental Table 2. Quantitation of PexRAP-dependent ether lipid species. As in Table 1

except that B cells of immunized mice were either WT or lacked PexRAP (cKO, i.e., *Dhrs7b*$^{Δ/Δ\text{-}}$

^{B}). Shown are (a) a sample of m/z features characteristic of the indicated phospholipid species.

(b) the mean ±SEM ion intensity counts in IMS data after mapping to the indicated regions of

interest as in Table 1. (c-f) indicate that p<0.05 for the null hypothesis in considering a difference

between WT mouse spleens after immunization versus UI controls (c), 1^0 B cell follicles in

spleens of immunized WT vs *Dhrs7b*$^{Δ/Δ\text{-}B}$ mice (d), GC vs in 1^0 B cell follicles in spleens of

immunized WT mice (e), and GC of immunized WT vs *Dhrs7b*$^{Δ/Δ\text{-}B}$ mice (f) in the ions counts

for designated m/z features. P values were calculated by Mann-Whitney U test.

1169

SIGNIFICANCE STATEMENT: The qualities and quantities of antibodies produced after B

cell activation are key elements of protection elicited by vaccines, the interplay of microbes with

our bodies, and some autoimmune reactions. Effective proliferation of B cells, i.e., the capacity

to increase the population of activated blasts, is a determinant of Ab response magnitude. In

addition, the size and function of germinal centers (GC) - micro-anatomic structures that form

after immunization and help diversify Ab responses and increase their affinity for antigens - are

influenced both by B cell physiology prior to entry into GC and by that of the GC B cell subset.

This work shows that a peroxisomal and ER-resident enzyme that generates intermediates for

ether lipid biosynthesis must be expressed in mature B cells in order for a subset of such

phospholipids to increase locally in lymphoid follicles and especially GC (secondary follicles).

Moreover, lack of this enzyme undermines B cell ROS control, survival, proliferation, GC

homeostasis, affinity maturation, and serum concentrations of antigen-specific antibodies.

Together, these findings reveal an essential function of B cell-autonomous synthesis of ether

lipids in adaptive immunity.

1184

1185 **********   ***********   ***********   ***********

1186

A

R26-CreER^{T2}

Dhrs7b^{f/f};
R26-CreER^{T2}

Tmx

SRBC   *Analysis*

Wk: 0   1

B

WT   *Dhrs7b^{Δ/Δ}*

B T   B T

◄ PexRAP

◄ Actin

C

WT   *Dhrs7b^{Δ/Δ}*

CD19

37   24

B220

D

p<0.001

% B cells among live cells

WT   Dhrs7b^{Δ/Δ}

p=0.006

# of B cells (× 10^6)

WT   Dhrs7b^{Δ/Δ}

E

WT   *Dhrs7b^{Δ/Δ}*

GL7

9.6   3.6

CD95

F

% GCB cells

WT   Dhrs7b^{Δ/Δ}

p=0.01

# of GCB cells (× 10^6)

WT   Dhrs7b^{Δ/Δ}

G

WT

KO

■ GL7 ■ IgD ■ CD35

H

WT

Dhrs7b^{Δ/Δ−B}

■ IgD^+
■ LZ (GL7^+ CD35^-)
■ DZ (GL7^+ CD35^+)

500 μm

I

p=0.02

GC number per spleen section

WT   Dhrs7b^{Δ/Δ}

p=0.047

GC area (mm^2)

WT   Dhrs7b^{Δ/Δ}

Fig. 1

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. 8

Fig. 9

Supplemental Fig. 1

Supplemental Fig. 2

Supplemental Fig. 3

Supplemental Fig. 4

Supplemental Fig. 5

Table 1.
    Immunization-induced increases of selected lipids in GC

Negative Ions			
m/z [a] Lipid ID	UI	1° B cell Follicle	GC
716.5216 PE(18:0_16:1)	618 ± 22 [b]	2400 ± 83 [c]	3674 ± 154 [d]
752.5545 PE(O-18:0_20:4)	301 ± 14	1739 ± 33 [c]	1960 ± 52 [d]
760.5056 PE(16:0_22:6)	203 ± 4	1816 ± 30 [c]	2031 ± 23 [d]
772.5242 PE(P-18:1_22:6)	234 ± 5	2334 ± 47 [c]	2578 ± 63 [d]
776.5556 PE(O-18:0_22:6)	277 ± 9	1638 ± 51 [c]	2074 ± 90 [d]
869.5526 PI(O-18:0_20:5)	242 ± 7	1611 ± 24 [c]	1789 ± 15 [d]
872.576 PC(18:0_24:0)	171 ± 6	1167 ± 16 [c]	1369 ± 18 [d]
888.5612 PS(22:1_22:6)	506 ± 13	2684 ± 54 [c]	3386 ± 72 [d]

Positive Ions			
m/z [a] Lipid ID	UI	1° B cell Follicle	GC
739.4587 PG(12:0_20:3)	3042 ± 102	3733 ± 116 [c]	4778 ± 274 [d]
740.465 PC(16:0_15:1)	1310 ± 13	1478 ± 66 [c]	1933 ± 142 [d]
784.5564 PC(10:0_24:0)	2840 ± 24	7956 ± 147 [c]	9856 ± 270 [d]
798.5347 PC(10:0_25:0)	9937 ± 269	27589 ± 574 [c]	34733 ± 1644 [d]
799.5375 PG(18:0_18:1)	5665 ± 65	12078 ± 282 [c]	15311 ± 790 [d]

Table 2. Quantitation of PexRAP-dependent ether lipid species.

Negative Ions					
m/z [a] Lipid ID	UI	WT follicle	PexRap cKO follicle	WT GC	PexRap cKO GC
716.5216 PE(18:0_16:1)	754 ± 43 [b]	1608 ± 182 [c]	1263 ± 112	2729 ± 247 [e]	1770 ± 137 [f]
752.5545 PE(O-18:0_20:4)	459 ± 49	657 ± 51 [c]	548 ± 46 [d]	1191 ± 61 [e]	734 ± 25 [f]
760.5056 PE(16:0_22:6)	229 ± 8	386 ± 25 [c]	312 ± 16 [d]	569 ± 27 [e]	408 ± 15 [f]
772.5242 PE(P-18:1_22:6)	388 ± 47	617 ± 46 [c]	499 ± 33 [d]	918 ± 21 [e]	646 ± 9 [f]
776.5556 PE(O-18:0_22:6)	449 ± 53	661 ± 50 [c]	562 ± 46	1130 ± 49 [e]	789 ± 13 [f]
869.5526 PI(O-18:0_20:5)	509 ± 81	641 ± 50 [c]	494 ± 40 [d]	1014 ± 21 [e]	690 ± 12 [f]
872.576 PC(18:0_24:0)	230 ± 19	285 ± 12 [c]	247 ± 9 [d]	522 ± 17 [e]	336 ± 9 [f]
888.5612 PS(22:1_22:6)	1423 ± 277	2051 ± 199 [c]	1904 ± 199	3260 ± 156 [e]	2629 ± 80 [f]

Positive Ions					
m/z [a] Lipid ID	UI	WT follicle	PexRap cKO follicle	WT GC	PexRap cKO GC
739.4587 PG(12:0_20:3)	3042 ± 102	3094 ± 40	2944 ± 31 [d]	3506 ± 140 [e]	3011 ± 46 [f]
740.465 PC(16:0_15:1)	1310 ± 13	1549 ± 14 [c]	1478 ± 15 [d]	1672 ± 53	1461 ± 14 [f]
784.5564 PC(10:0_24:0)	2840 ± 24	2917 ± 74	2617 ± 46 [d]	3522 ± 102 [e]	2561 ± 42 [f]
798.5347 PC(10:0_25:0)	9937 ± 269	10928 ± 214 [c]	11372 ± 155 [d]	15378 ± 116 [e]	14694 ± 186 [f]
799.5375 PG(18:0_18:1)	5665 ± 65	6411 ± 76 [c]	5822 ± 74 [d]	6772 ± 139 [e]	5983 ± 85 [f]
801.5509 PG(18:0_18:0)	1540 ± 37	1591 ± 46	1633 ± 26	1444 ± 27 [e]	1533 ± 22 [f]
847.5361 PI(15:0_18:0)	1277 ± 24	1339 ± 54	1217 ± 35	1150 ± 46 [e]	1044 ±44
874.5623 PC(20:5_22:6)	754 ± 39	688 ± 48	675 ± 25	469 ± 27 [e]	556 ± 37