B cell expression of the enzyme PexRAP, an intermediary in ether lipid biosynthesis, promotes antibody responses and germinal center size

Sung Hoon Cho[†@&][*], Marissa A. Jones, Kaylor Meyer[†],

David M. Anderson[+], Sergei Chetyrkin[+], M. Wade Calcutt[+], Richard M. Caprioli[+],

Clay F. Semenkovich[~], Mark R. Boothby[†@&] [*]

Author affiliations [†]Department of Pathology-Microbiology-Immunology, [@]Vanderbilt Center for Immunobiology, and [&]Vanderbilt Institute for Infection, Inflammation, & Immunology, [+]Vanderbilt University Medical Center, Nashville, TN 37232; Mass Spectrometry Research Center and Department of Biochemistry, Vanderbilt University Medical School, and ^Cancer Biology Program, Vanderbilt University, Nashville TN 37212; [~] Division of Endocrinology. Metabolism & Lipid Research, Dept. of Medicine, Washington University, St. Louis MO 63110

[*] Correspondence: Sunghoon.Cho@vumc.org and Mark.Boothby@vumc.org

Dept. of Pathology, Microbiology & Immunology, Vanderbilt University Medical Center,

AA-4214 MCN Building, 1161 21[st] Ave. S., Nashville, TN 37232-2363

ABSTRACT

The qualities of antibody (Ab) responses provided by B lymphocytes and their plasma cell (PC) descendants are crucial facets of responses to vaccines and microbes. Metabolic processes and products regulate aspects of B cell proliferation and differentiation into germinal center (GC) and PC states as well as Ab diversification. However, there is little information about lymphoid cell-intrinsic functions of enzymes that mediate ether lipid biosynthesis, including a major class of membrane phospholipids. Imaging mass spectrometry (IMS) results had indicated that concentrations of a number of these phospholipids were substantially enhanced in GC compared to the background average in spleens. However, it was not clear if biosynthesis in B cells was a basis for this finding, or whether such cell-intrinsic biosynthesis contributes to B cell physiology or Ab responses. Ether lipid biosynthesis can involve the enzyme PexRAP, the product of the *Dhrs7b* gene. Using combinations of IMS and immunization experiments in mouse models with inducible *Dhrs7b* loss-of-function, we now show that B lineage-intrinsic expression of PexRAP promotes the magnitude and affinity maturation of a serological response. Moreover, the data revealed a *Dhrs7b*-dependent increase in ether phospholipids in primary follicles with a more prominent increase in GC. Mechanistically, PexRAP impacted B cell proliferation via enhanced survival associated with controlling levels of ROS and membrane peroxidation. These findings reveal a vital role of this peroxisomal enzyme in B cell homeostasis and the physiology of humoral immunity.

Abbreviations used:

Ag, antigen; Ab, antibody; WT, wildtype; KO, knockout; MZB, marginal zone B cell; FoB, follicular B cell; PC, plasma cell; Ig, immunoglobulin; GC, germinal center; ETC, electron transport chain; ROS, reactive oxygen species; mtROS, mitochondrial ROS; FAO, fatty acid oxidation; EL, ether lipid(s); 2D, 2-dimensional; MALDI, matrix-assisted laser desorption/ionization; IMS, imaging mass spectrometry; ER, Endoplasmic Reticulum; PexRAP, Peroxisomal Reductase Activating PPARγ; DHAP, dihydroxyacetonephosphate; G3P, glycerol-3-phosphate; CTV, CellTrace Violet; SRBC, sheep red blood corpuscle; TCR, T cell receptor; CD, Cluster of differentiation; AID, Activation-induced deaminase; GFP, Green fluorescent protein; NP-OVA, 3-Nitrophenylacetyl (NP)-ovalbumin (OVA); ASCs, Ab-secreting cells; CSR, class-switch recombination; MBC, memory B cell; NADPH, nicotinamide adenine dinucleotide phosphate; NAC, N-acetyl-L-cysteine; OCR, oxygen consumption rate; ECAR, extracellular acidification rate; RCDP, rhizomelic chondrodysplasia punctata; FAS, fatty acid synthase; Tmx, tamoxifen; 4OHT, 4-hydroxytamoxifen; PUFA, poly-unsaturated fatty acids; 2-DG, 2-deoxyglucose; FCCP, carbonyl cyanide 4-phenylhydrazone; BSA, bovine serum albumin; PSA, porcine serum albumin; H2DCFDA, dichlorodihydrofluorescein diacetate; ELISA, enzyme-linked immunosorbent assay; ELISpot, enzyme-linked immunosorbent spot; DHA, dihydroxyacetophenone; DAN, 1,5-diaminonaphthalene; timsTOF, trapped ion mobility spectrometry time-of-flight; SDS-PAGE, sodium dodecyl-sulfate polyacrylamide gel electrophoresis; SEM, standard error of means.

INTRODUCTION

The qualities and concentrations of antigen (Ag)-specific antibodies (Ab) are key elements of immunity and the pathophysiology of diverse conditions involving inflammation (1-3). Ab are secreted by cells in the B lymphocyte lineage, but the exact sources are diverse (4-7). Many derive from a subset termed B1 B cells, which are thought to having characteristics of innate responses (4, 5), but progeny of follicular and marginal zone B cells (FoB and MZB, respectively) can differentiate into Ab-secreting plasma cells (PC) after activation and proliferation (6-8). Ab concentrations and affinity for Ag, as well as Ab isotype after class switching in a B cell, determine the efficacy of pathogen clearance (9-12). Importantly, the affinities of serum Ab for target Ag can increase over time after an immune exposure, especially after recurrent encounter(s) (8, 9, 13, 14).

Many sources and forms of Ab can be protective, though in several autoimmune diseases these molecules and the somatic mutations that diversify an initial BCR repertoire drive pathology (15, 16). The micro-anatomic structure termed the germinal center (GC) provides one mechanism to increase spectrum of Ab affinities over time and repeated Ag exposure (17-19). Although T cell help-dependent diversification and PC production can occur independent from GC, results with different forms of vaccination support the practical importance of GC (20, 21). Current evidence holds that most Ig class-switching is executed before GC entry of proliferating B cells (22). GC reactions are initiated after B cells in a primary lymphoid follicle encounter Ag, migrate to interact with activated helper T cells after extensive proliferation (23). GC formation, size, and function depend on multiple factors. These include the efficiency of population growth for B cells in a phase before they enter and take on characteristics of GC B cells, and on homeostatic and differentiative processes while in the secondary follicle [reviewed in (24)]. Once in the GC, B cells cycle between light and dark zones, undergoing iterative cycles of selection. Proliferation and selection end in development as Ab-secreting plasma cells or memory B cells [reviewed in (14, 18, 19)]. In line with a selection process, substantial rates of B cell death have

been measured in GC (25). As such, the qualities and quantities of Ab elicited by immunization

(or vaccination) depend on B cell proliferation and survival both before and during GC reactions.

Accumulating evidence indicates that metabolic reprogramming and intermediary

metabolism play pivotal roles in survival, proliferation, differentiation and function in the pre-

immune populations (26, 27), including B cells (28, 29). Among core metabolic processes,

several converge on oxidative metabolism fed by glycolytic generation of pyruvate, anaplerotic

use of amino acids such as glutamine, and fatty acid oxidation [reviewed in (29, 30)]. Compared

to resting populations such as naive B cells, rates of such metabolism in B cells appear to be

increased after activation, including in GC B cells [31-36; reviewed in (29)]. Such increases

support the generation of substrates for anabolic processes in the dividing cell population and

perhaps the energy demands thereof.

Mitochondria have been the focus of investigations exploring molecular mechanisms for

oxidative metabolism to generate energy and substrates in lymphocytes during immune

responses [reviewed in (37, 38)]. Oxidative phosphorylation in and dynamics of this organelle

appear to be important in determining rates of somatic hypermutation that characterize GC B cell

biology (33-35), albeit by a molecular mechanism that has not yet been identified. Among

specialized functions of this highly dynamic subcellular organelle, mitochondria use the Krebs

TCA cycle to feed an electron transport chain (ETC) and efficiently generate ATP. Along with

other sources, mitochondria generate reactive oxygen species (ROS) via the ETC at a rate

determined by structural and biochemical features (39, 40). Rates of ROS production, release,

and resolution require regulation at multiple levels because these labile species and their

reactions with cellular molecules are important for both signaling and cell proliferation but

undermine cell function or survival when excessive.

B cells in most GC exhibit inadequacy of oxygen delivery relative to demands (31, 41-43).

Of note, genome-wide screening for genes that help cells adapt to such hypoxia revealed that

peroxisomes and an ER enzyme involved in synthesizing plasmalogen ether lipids (EL) decrease

the death of cells due to insufficient oxygen (44). Using an unbiased lipidomic analysis that

combined imaging mass spectrometry (IMS) and techniques for identifying GC in sections of

spleen from immunized mice, we had found that local concentrations of a subset of EL were

heightened in GC relative to other portions of the spleen (45). Ether phospholipids (EPL) have a

fatty acid linked to the glycerol backbone with an alkyl or a vinyl ether bond at the sn-1 position

(46, 47). Synthesis of ether phospholipids depends on the generation of precursors in

peroxisomes and final processing in the ER (48, 49). However, EL biosynthesis is prominent in

the liver, and in theory these species or their precursors might distribute to tissues from a primary

site of biosynthesis or from the diet (49-53). Thus, the finding that secondary follicles had greater

densities of some - but not all - ether phospholipids left several key questions unanswered – (1)

Does the capacity to synthesize and increase ether lipids in B cell follicles – primary or

secondary (i.e., GC) - affect antibody responses? (2) Are any of the EL dependent on B lineage-

intrinsic metabolism to generate precursors or final products? Moreover, there is debate whether

EL or their plasmalogen subset promote cell death or resistance, and whether inactivation of

biosynthesis causes defects in establishment or maintenance of hematopoietic cells (46, 53-55).

Peroxisomal Reductase Activating PPARγ (PexRAP, encoded by the *Dhrs7b* gene) catalyzes

the reduction of alkyl-dihydroxyacetonephosphate (DHAP) to 1-alkyl-G3P (46, 48, 56), which

then is transferred into endoplasmic reticulum (ER) and further metabolized to generate EL.

Recent work notes that this enzyme may substantially function in the ER in addition to the

peroxisome (56). Widespread inactivation of *Dhrs7b* in mature mice decreased hepatic

production of ether lipid species and reduced the population of erythrocytes and neutrophils (46).

This body of work indicated that PexRAP can be essential in some aspects of ether lipid

metabolism and may function in hematopoietic cells, but was questioned based on inborn errors

of metabolism affecting other aspects of ether lipid biosynthesis (54). Diverse functions of ether

lipids in general, and the plasmalogen subset in particular, have been proposed or supported by

prior work [reviewed in (57-61)]. The ether linkage may provide a sink for oxygen radicals -

thereby acting directly to resolve the reactivity and contribute to ROS homeostasis, although

there is debate as to the physiological impact of this chemical property (62). Whether or not this

lipid subset influences adaptive immunity or conventional lymphocyte lineages, however, is

unclear.

Herein, we tested the hypotheses that (i) the ether lipid profiles in activated B cells depend

on their expression of the biosynthetic enzyme PexRAP [alternatively referred to as acyl/alkyl-

DHAP reductase (ADHAPR) (56)], and (ii) this enzyme promotes B cell physiology. To do so,

we combined conventional lipidomics, Imaging Mass Spectrometry (IMS), and immunization

experiments with genetic models in which loss-of-function for *Dhrs7b* is induced in adult mice.

We provide evidence that PexRAP impacts B cell proliferation via enhanced survival associated

with controlling levels of ROS and membrane peroxidation. The results also indicate that beyond

an impact on B cell homeostasis, GC are reduced in its absence, and the magnitude and affinity

maturation of a serological response depend on B cell expression of PexRAP. Taken together,

these findings support a function of B cell-intrinsic biosynthesis of ether lipids in B lymphocyte

homeostasis, the production of Ab, and in promoting GC reactions.

RESULTS

PexRAP promotes homeostatic maintenance and proliferation of B cells. By use of IMS, we

had reported that at least a dozen ether phospholipids - including plasmalogens - were enriched

in splenic germinal centers of immunized mice when compared to the rest of the tissue (45).

Previous work provided evidence that these lipid molecules were important for the homeostasis

of short-lived neutrophils (46). However, this effect was questioned based on hematological data

from patients inborn errors of metabolism that affect ether lipid biosynthesis (54). Moreover, the

impact of these molecules on adaptive immune cells or functions is not known. Accordingly, we

sought to test for effects of an intervention that interferes with a key biosynthetic pathway used

for endogenous synthesis of these molecules. To do so, we started with analyses using induced

loss-of-function for $Dhrs7b^{f/f}$ and the widely expressed $Rosa26\text{-}CreER^{T2}$ transgene (63, 64) used

in (46), which reduced hepatic generation of a subset of ether lipids and acts in all hematopoietic

cells as well.

We modeled initial experiments on the published work with imaging mass spectrometry

(45). Mice were treated with tamoxifen using conditions less intense than a regimen shown to

have no effect on pre-immune splenic and B lineage populations (65, 66), immunized with

SRBC [akin to the analyses in (31, 36, 41, 45)], and analyzed 7 d after immunization (Fig. 1A).

Functional inactivation of $Dhrs7b$ in T and B lymphocytes was confirmed by Western blot

analysis using anti-PexRAP Ab (Fig. 1B). The frequencies and numbers of CD19$^+$ B220$^+$ B cells

in $Dhrs7b^{f/f}$; $Rosa26\text{-}CreER^{T2}$ - only a small minority of which would have been activated by the

immunization - were approximately 0.6 the values for controls ($Dhrs7b^{+/+} Rosa26\text{-}CreER^{T2}$,

i.e., wild-type locus) (Fig. 1C, D). In contrast to B cell populations, TCRβ$^+$ CD4$^+$ T cells and

TCRβ$^+$ CD4$^-$ CD8 T cells were intact in tamoxifen-treated and SRBC-immunized $Dhrs7b^{f/f}$;

$Rosa26\text{-}CreER^{T2}$ mice (Supplemental Fig. 1A-C). These results suggest that $Dhrs7b$ is required

for fully maintaining the B cell population but indicate that the enzyme is dispensable for the

main sets of conventional T cells. In light of the earlier finding that a subset of ether lipids was

more prevalent in secondary follicles, we analyzed GC in these samples. Microscopy and flow

cytometry analyses after immunofluorescent staining found that less GC B cells were present

after immunization of *Dhrs7b^{f/f}*; *Rosa26-CreER^{T2}* mice, i.e., less IgD⁻ CD95⁺ GL7⁺ (GC-

phenotype) B cells (Fig. 1E-I; Supplemental Fig. 1D). Quantitation of the immune fluorescence

micrography with spleens from immunized mice revealed that the numbers of GC per spleen

were halved in *Dhrs7b^{Δ/Δ}* mice compared with controls (Fig. 1I; Supplemental Fig. 1E). The sizes

of those GC that did form were substantially reduced as well (Fig. 1I; Supplemental Fig. 1F).

The lower number of B cells found in immunized mice - while a less profound effect than the

reduction in GC - tempers the result, and it was possible that the function of helper CD4 T cells

was impaired even their though numbers were not affected. Consistent with this hypothesis, the

frequencies of Tfh and GC-Tfh cells were lower in *Dhrs7b^{Δ/Δ}* mice compared with controls

(Supplemental Fig 1G-I). While not addressing whether or not B cells are part of the requirement

for PexRAP in promoting GC, these results indicate that *Dhrs7b* is necessary for a normal-sized

geminal center response.

To determine if there are B lineage-specific functions of PexRAP, we used a conditionally

active Cre transgene that is expressed specifically in mature B cells (67). *Dhrs7b* f/f , *huCD20-*

CreER^{T2} mice and *huCD20-CreER^{T2}* were treated with tamoxifen to induce Cre-mediated

recombination of the floxed alleles after initial establishment of normal B cell populations (Fig.

2A). Western blot analyses showed an almost complete absence of PexRAP protein from B cells

purified from tamoxifen-treated *Dhrs7b^{f/f}*; *huCD20-CreER^{T2}* (hereinafter, *Dhrs7b^{Δ/Δ-B}*) mice

compared to similarly treated *huCD20*-CreER^{T2} controls (Fig. 2B). The enzyme encoded by this

gene catalyzes the synthesis of a lipid precursor to many - but not all - plasmalogens and other

ether phospholipids (46-48), and *Dhrs7b* deficiency impacted neutrophil survival (46). When we

tested if *Dhrs7b* inactivation affected the steady-state B cell population, the frequencies and total

numbers of CD19⁺ B220⁺ B cells in viable lymphocyte gates were about 20% lower in

tamoxifen-treated $Dhrs7b^{\Delta/\Delta-B}$ mice compared to tamoxifen-injected, $huCD20$-CreER^{T2},

$Dhrs7b^{+/+}$ controls (Fig. 2C). With loss-of-function after production of mature B cells, i.e.,

inactivation of $Dhrs7b$ at ages over 6 wk, we observed balanced decreases in the numbers of

multiple subsets within the B lineage without evidence of selectivity among main sub-classes of

B cell. Moreover, the surface expression of IgM was comparable in $Dhrs7b^{\Delta/\Delta}$ mice versus WT

controls (Supplemental Fig 2. A-F). Thus, the frequencies of MZB and FOB amidst the smaller

population of splenic B cells in tamoxifen-treated $Dhrs7b^{f/f}$; $huCD20$-CreER^{T2} mice were similar

(Supplemental Fig 2B), and frequencies of B1 B cells in the peritoneal cavity also were

unaffected (Supplemental Fig 2E). Prior work documents normal frequencies and numbers of

developing and mature B cells under harsher conditions of tamoxifen-induced deletion with

CreER^{T2} (65). De novo B cell production rates are too low to yield the ~20% decrease observed

one week after gene disruption, and few splenic B cells are in cell cycle. Accordingly, we infer

that after maturation of a B cell, $Dhrs7b$ promotes its survival, albeit to a modest degree.

B cell population growth can be impacted by increased cell death and/or decreased

proliferation. To test if PexRAP affects proliferation in vivo, $Dhrs7b^{\Delta/\Delta}$ B cells were stained with

CellTrace Violet (CTV) and adoptively transferred into B cell deficient μMT recipient mice.

Strikingly fewer PexRAP-depleted B cells were recovered (Fig. 2D), and the frequencies of

$Dhrs7b^{\Delta/\Delta}$ B cells that were IgD^{neg} (i.e., had been activated) were half those of controls (Fig. 2E;

Supplemental Fig. 2G). As compared to control B cells, CTV partitioning analyses suggested that

division rates in vivo were modestly lower for the $Dhrs7b^{\Delta/\Delta}$ B cells (Fig. 2E, F). Mitogen-

stimulated $Dhrs7b^{\Delta/\Delta}$ B cells that proliferated in culture exhibited substantially less robust

division: frequencies of viable cells that had undergone ≥3 divisions were halved in $Dhrs7b^{\Delta/\Delta}$ B

cells) and yielded ~1/3 as large a progeny population (Fig. 2G-I; Supplemental Fig 2H).

Collectively, these data indicate that the $Dhrs7b$ gene product supports homeostatic maintenance

of a quiescent (pre-immune) B cell population *in vivo* and effective proliferation of B cells.

B cell expression of PexRAP is required for achieving normal concentrations

of ether phospholipids. As noted, lipidomic analyses that used 2-dimensional image mass-

spectrometry (2D-IMS) discovered that local concentrations of a subset of ether lipids are

enriched in GC compared with the area outside of GC after immunization (45). To measure the

extent to which PexRAP expression within mature B cells can affect their lipid content and

composition, including their ether- and plasmalogen phospholipids, LC-MS-MS analyses were

performed with B cells directly isolated from spleens as well as those cultured after mitogenic

activation. These analyses showed substantial reductions in many ether phospholipids - as well

as lysophosphatidylethanolamines (LPE) generated by phospholipase cleavage of the R2 fatty

acid sidechain of an ether phospholipid - in PexRAP-deficient B cells (Fig. 3A, B; Supplemental

Fig. 3A, B). While beyond the scope of this work, this evidence suggests that generation of

potential signaling molecules via phospholipase(s) such as PLA2 may be reduced. Of note,

Dhrs7b inactivation reduced levels of a subset of plasmalogens in naive B cells (Fig. 3A, B, D;

Supplemental Fig 3A) though changes of others were modest or absent in the naive population. A

greater impact of PexRAP on quantitative amounts of phospholipids was found in the activated B

cells. Several species, skewed towards those with the most detected ions, a number of which

were at less than 1/5th the level of non-deleted control B cells (Fig. 3C, D). Phospholipid results

observed with naive and activated B cells from tamoxifen-treated CreER^{T2}-expressing mice were

similar to those of unmanipulated controls (Supplemental Fig. 3C, D). There is no in vitro

surrogate that faithfully represents GC or the GC B cell, but collectively these data provide

strong evidence that the capacity to rapidly produce increased levels of many ether lipid

molecules after lymphocyte activation depends on a functional *Dhrs7b* gene in B cells. The data

provide evidence that *Dhrs7b* in B cells enhances cell-autonomous biosynthesis of some

plasmalogens and other ether lipid species. Especially in resting B cells that are out of cycle,

additional uptake or biosynthesis pathways may add to the overall pool (46, 47).

Notwithstanding these potential contributions, however, the straightforward inference is that

PexRAP catalyzes the generation of ether lipid precursors in amounts crucial for regulating the

overall ether phospholipid pools, especially after B cell activation.

Our earlier work (45) reported only ion mode features in negative mode and did not analyze

the primary follicle (the vast majority of which are resting B cells) relative to the extrafollicular

white pulp, red pulp, and GC. To investigate the requirement for specific expression of PexRAP,

we started with AID-GFP mice to enhance spatial localization. These measurements confirmed

that a substantial number of both positive and negatively charged ions whose exact masses

identified them as plasmalogens were substantially concentrated in GC as compared to the rest of

the B cell follicle (Table 1; Supplemental Fig. 4A, B). Accordingly, we tested if the differential

enrichment of any ether lipid species in primary or secondary follicles depends on biosynthesis

within a specific lymphoid lineage. Mice with conditional mutations were tamoxifen-treated,

immunized with SRBC, harvested seven days thereafter, and analyzed by IMS (Fig. 4A, B). With

deletion driven by the widely expressed *Rosa26-CreER^{T2}* transgene, as in earlier work (46) and

Fig. 1, substantial reductions of the GC enrichment pattern were observed (Fig. 4C). For

instance, two representative ions identified in negative ion mode (i.e. *m/z* 752.5545 and *m/z*

776.5556) were again enriched in GC regions of spleens from immunized WT mice. These

features barely increased in splenic samples of immunized *Dhrs7b$^{\Delta/\Delta}$* (tamoxifen-injected

Dhrs7b$^{f/f}$; *Rosa26-CreER^{T2})* mice (Fig. 4C).

To determine if the levels of particular ether phospholipids in GC regions were affected by

PexRAP expression in B cells, *Dhrs7b$^{f/f}$*; *huCD20*-CreER^{T2} mice and *huCD20*-CreER^{T2} controls

were analyzed after tamoxifen injections followed by immunization, and compared to samples

from unimmunized mice. IMS using both positive and negative ion modes identified at least

eight ions - including *m/z* 752.5545, *m/z* 776.5556, and *m/z* 872.5749 - much more substantially

(~2-3-fold) concentrated in GC of immunized control mice (Fig. 4D, E; Table 2) than in the

primary follicle of unimmunized mice. Notably, these increased signals in secondary follicles

(GC) were reduced or almost completely eliminated in GCs of mice immunized after B cell-

specific PexRAP depletion ($Dhrs7b^{\Delta/\Delta.B}$) (Fig. 4D, E; Table 2; Supplemental Fig. 4C).

Collectively, these results indicate that *Dhrs7b* gene expression in activated B cells regulates the

spectrum of ether lipids in the micro-anatomic locale of a secondary follicle. Interestingly,

immunization also led to modest increases in the levels of some ether and conventional

phospholipids in the B cell-rich primary follicle regions (Table 2). This finding extends the

previous report (45), which focused on the relationship between the AID-GFP^{hi} region (i.e., GC)

and lipid features. Moreover, the absence of PexRAP blunted most of the immunization-induced

increases observed in the primary follicles. These findings and those of Fig. 3 suggest that there

are secondary consequences of *Dhrs7b* inactivation (e.g., some di-acyl phospholipids are

affected), but provide direct evidence that the enhancement of many ether phospholipid signals

in the GC depends on PexRAP in B cells.

B cell-intrinsic role of *Dhrs7b* in Ab affinity and quantity.

As key components of adaptive immunity, progeny of an activated B cell can differentiate

into Ab-secreting plasma cells (PC) or into GC B cells. Over the course of an immune response,

the GC reaction diversifies the affinity for and breadth of antigen recognized by the original

BCR, and improves properties of memory [(14, 17-19, 68). PC that develop after a second

encounter with antigen increase the amounts of high-affinity Ab. The higher quality and quantity

of Ag-specific Abs from plasma cells are integral to humoral immune responses (68). An

increasing body of evidence indicates that metabolic reprogramming and intermediary

metabolites modulate immune cell differentiation and function (26-36). The observed B cell-

intrinsic function of *Dhrs7b* in the accumulation of many ether lipid species in both primary and

secondary follicles prompted us to test the effect of B cell type-restricted PexRAP depletion on

GC responses and Ag-specific Ab production. Tamoxifen-injected $Dhrs7b^{f/f}$; $huCD20\text{-}CreER^{T2}$

and control mice were immunized with NP-ovalbumin (NP-OVA), then boosted with NP-OVA to

elicit affinity-matured Ab, and harvested 1 week thereafter (Fig. 5A). Although PexRAP-

deficient B cell numbers were reduced less than 20% a week after starting gene inactivation prior

to immunization (i.e., were over 0.8-fold those of controls) (Fig 2A-C), frequencies and

numbers of IgD^{neg} B cells were halved in $Dhrs7b^{\Delta/\Delta\text{-}B}$ mice at harvest (4 wk after completion of

the induced deletion) (Fig. 5B, C). Moreover, the GL7$^+$ CD95$^+$ fraction of the B cell population

in the dump / IgD^{neg} gate was substantially reduced (Fig. 5D), such that numbers of GL7$^+$ CD95$^+$

GC B cells were dramatically decreased in $Dhrs7b^{\Delta/\Delta\text{-}B}$ mice (Fig. 5E). Consistent with these

findings, GC in $Dhrs7b^{\Delta/\Delta\text{-}B}$ mice were reduced after SRBC immunization (Fig 5 F, G;

Supplemental Fig. 5A, B), and the numbers of centrocytes and centroblasts were dramatically

lower in $Dhrs7b^{\Delta/\Delta\text{-}B}$ mice compared with WT controls (Fig 5H). LZ B cells (CD86$^+$ CXCR4^{lo})

trended toward being at lower prevalence in $Dhrs7b^{\Delta/\Delta\text{-}B}$ mice, whereas their DZ counterparts

(CD86^{neg} CXCR4$^+$) were comparable to wildtype controls, resulting in a modest decrease in the

LZ/DZ ratio when B cells lacked PexRAP (Supplemental Fig. 5C). The frequencies of CD138$^+$

GL7$^+$ IgD$^-$ early plasmablasts were comparable between $Dhrs7b^{\Delta/\Delta\text{-}B}$ mice and control mice

(Supplemental Fig 5D). Although the loss-of-function was B cell specific, the frequencies of Tfh

and GC-Tfh cells were decreased in $Dhrs7b^{\Delta/\Delta\text{-}B}$ mice compared with control mice

(Supplemental Fig 5E). We infer that support for the Tfh populations was reduced due to

impact(s) on GC B cell numbers and / or function.

Ab class-switch recombination (CSR) and affinity maturation can occur through

extrafollicular responses. However, the GC reaction significantly increases Ab diversification

and high-affinity Ab production, in part later in a primary response but also upon secondary

exposure to antigens (6, 8, 9, 17-19). ELISA performed with sera at 3 wk post-immunization

(Fig. 6A-D), just before the boost, showed that B cell-specific depletion of PexRAP prior to

immunization reduced NP-specific IgM and the switched isotype IgG1 (Fig. 6A, C). Of note, the

capacity to generate high-affinity Ab, detected with low valency NP$_2$, was even more severely

undermined than the overall response - especially for IgG1 (Fig. 6B, D). A week after a second

immunization - an interval similar to that which followed immunization with SRBC - Ab-

secreting cells (ASCs) in the spleen (Fig. 6E) and circulating anti-NP IgM concentrations were

diminished in $Dhrs7b^{\Delta/\Delta-B}$ mice compared with controls ($huCD20$-CreER^{T2}, $Dhrs7b^{+/+}$ treated

with tamoxifen in parallel to the mice with induced loss of PexRAP) (Fig. 6F). The ratio of high-

affinity (NP$_2$-binding) to all-affinity (NP$_{20}$) IgM Ab also was substantially lower in $Dhrs7b^{\Delta/\Delta-B}$

mice (Fig. 6G), as were the serum levels of high-affinity IgG1 and IgG2c, class-switched

isotypes (Supplemental Fig. 5F, G). These data reinforce and extend the conclusion that the

expression of PexRAP in mature B cells is important for their physiology and function.

Consistent with *in vivo* results finding reduced Ag-specific ASCs in $Dhrs7b^{\Delta/\Delta-B}$ mice

compared with controls (Fig. 6E), the attenuated population of PexRAP-deficient B lineage cells

recovered after transfer into recipient mice (Fig. 2D) yielded lower frequencies of CD138$^+$

progeny (Fig. 6H, I) along with evidence of reduced activation [i.e., lower frequencies of IgD$^-$

progeny (Supplemental Fig. 2G). Mitogen-stimulated $Dhrs7b^{\Delta/\Delta}$ B cells also yielded lower

frequencies of CD138$^+$ cells in vitro, but the division-specific frequencies of CD138$^+$ cells

showed only modest decreases in $Dhrs7b^{\Delta/\Delta}$ B cells (Supplemental Fig. 6 A, B). Thus, the

reduction of CD138$^+$ cell differentiation appears to be due mostly to its dependence on survival

to a sufficient division count.

Dhrs7b inactivation was initiated prior to immunization in the preceding experiments.

Therefore, the impact of PexRAP deficiency on GC and the Ab response in such a setting might

be due exclusively to impairment of B cells, e.g., their reduced population expansion, prior to

their entry into GC. Alternatively, the effects could also in part involve a requirement for

PexRAP-dependent metabolites within GC B cells. To test if *Dhrs7b* functions within GC, we

used conditional deletion of *Dhrs7b* driven by the *S1pr2-CreER^{T2}* transgene whose expression at

high levels marks GC B cells (69). Of note, experiments with a fate-marking reporter allele

showed that activated B cells that lack the GC B phenotype were not marked by this conditional

Cre after immunization with the NP-carrier approach (69). *Dhrs7b$^{f/f}$*; *S1pr2-CreER^{T2}* and control

S1pr2-CreER^{T2} (control) mice were immunized with SRBC, and tamoxifen was injected at a time

point after the initiation of GC to test more directly that the inactivation of *Dhrs7b* would be in

GC B cells rather than pre-GC blasts (Figure 7A). Frequencies of GC B cells were substantially

lower in tamoxifen-treated *Dhrs7b$^{f/f}$*; *S1pr2*-CreER^{T2} mice (Figure 7B, C). While this finding

does not exclude that there may be an additional effect of impaired clonal expansion in B cells

activated by immunization but not yet resident in the GC, the results indicate that PexRAP

functions within GC B cells to support a full population of this subset. All together, these data

provide evidence that *Dhrs7b* expression influences the lipidome and function of GC B cells .

Notably, an optimal GC response requires PexRAP function in B cells, and the B cell-intrinsic

functions of *Dhrs7b* promote the quantity and affinity maturation of Ab responses, in part via net

proliferation of B cells.

***Dhrs7b* contributes to modulation of ROS and their impact on B cell population growth.**

329   Increased susceptibility to cell death after activation would be a mechanism that could

cause the reduced population expansion, GC, and Ab production. Consistent with earlier analyses

of LPS-stimulated B cells (70), we had noted that ROS levels were higher in GC B cells,

memory B cells (MBCs) and ASCs than in the naive B2 B cell pool (36). ROS can be produced

in activated B cells via diverse sources that include mitochondrial electron transport chain

function, NADPH oxidase activated by BCR engagement, and fatty acid oxidation in

peroxisomes (71-73). ROS can positively mediate signal transduction, but their steady-state

concentrations need to be calibrated because overproduction can have deleterious effect on cells,

for instance via lipid peroxidation at excessive rates (74-76). Endogenous antioxidant properties

have been imputed to plasmalogens because the vinyl ether bond is susceptible to reaction with

reactive oxygen, thereby scavenging ROS (48, 57-59), so we tested if *Dhrs7b* function

contributes to redox control in B cells. ROS levels were ~2-fold higher in GC B cells from

immunized *Dhrs7b$^{\Delta/\Delta\text{-}B}$* mice compared to those from immunized WT controls (Fig 7D).

Dhrs7b$^{\Delta/\Delta}$ GC B cells also showed higher cell death and attenuated BrdU incorporation (Fig 7E,

F; Supplemental Fig 6C.) Moreover, CD40-stimulated *Dhrs7b$^{\Delta/\Delta}$* B cells exhibited ~3 fold higher

signal and about a doubling when stained with fluorescent sensors of cellular and mitochondrial

ROS, respectively, compared to WT B cells *in vitro* (Fig. 8A-D). Substantially increased ROS

also were measured after activation by BCR and CD40 cross-linking (Supplemental Fig 6D).

Iron- and copper-dependent lipid peroxidation is considered to be a biological mechanism for

ROS-mediated cell death (75, 76). Lipid peroxidation measured by a fluorescent indicator (C11-

Bodipy) in vitro was higher in $Dhrs7b^{\Delta/\Delta}$ B cells compared with WT controls after each type of

mitogen activation (Fig. 8E; Supplemental Fig 6E). Chemical elicitation of increased ROS and

lipid peroxidation in B cells, independent from B cell activation or proliferation, rapidly led to

reduced cell viability and B cell numbers (Fig. 8F; Supplemental Fig 7A). Furthermore, in vitro

analyses indicated that B cells lacking PexRAP also exhibited increases in the activated

executioner caspase, cleaved caspase-3 (CC3) along with early apoptotic cells that are annexin

V$^+$ but exclude 7-aminoactinomycin D (Fig. 8G, H). These data indicate that $Dhrs7b$ function

supports not only protection against cell death, but also control of cellular and mitochondrial

ROS levels and their impact on lipid modification (Supplemental Fig. 7). Of note, while H_2O_2

did not increase this early apoptotic population (Supplemental Fig. 7C), increasing ROS with

menadione drove B cell death preceded by annexin V$^+$ state without an increase in C11-Bodipy

signal (Supplemental Fig. 7D-G). We infer that rather than a single mode of death, increased

ROS resulting from PexRAP depletion probably stimulates at least two distinct modes of B cell

death.

Consistent with the increased steady-state ROS and death, activated $Dhrs7b^{\Delta/\Delta}$ B cells

exhibited a defect in B cell population growth *in vivo* and *in vitro* (Fig. 2; Fig. 8A-H). The well-

established ROS scavenger, N-acetyl-L-cysteine (NAC), exerted a concentration-dependent

effect that improved the survival and population growth of PexRAP-deficient B cells (Fig. 8I).

We exploited the sub-maximal rescue by a lower concentration of NAC (1 mM) to explore if

peroxisomal oxidative metabolism might contribute to the toxicity observed when B cells are

PexRAP-depleted. Although thioridazine on its own provided no increase in the B cell

population growth, its combination with 1 mM NAC treatment improved growth of the

Dhrs7b$^{\Delta/\Delta}$ B cell population compared to either agent on its own (Fig. 8J). Analyses of B cells

activated by BCR crosslinking along with anti-CD40 also found that PexRAP expression in B

cells supports their population increase (Supplemental Fig 6F). Moreover, inclusion of NAC in

the cultures partially mitigated the inactivation of *Dhrs7b* (Supplemental Fig 6F). Of note,

measurements of CTV partitioning identified a modest but statistically significant decrease in

division counts of PexRAP-deficient B cells under these conditions, and found that NAC did not

increase this component of proliferation (Supplemental Fig 6G, H). These data support the

conclusion that sufficient generation of PexRAP-dependent products is crucial for B cell

survival. Moreover, this enzyme contributes - directly or indirectly - to a resolution or

detoxification of ROS that is critical for population growth of activated B cells.

Dhrs7b in B cells affects their oxidative metabolism and ER mass.

Our finding that mitochondrial ROS were increased prompted us to investigate how PexRAP

affects physiological functions by measuring respiration in activated B cells. Mitochondrial

stress tests with in vitro activated B lymphoblasts measured small but definite decreases in both

basal and maximal respiration (oxygen consumption rates, or OCR) (Fig. 9A-C). The altered

performance of mitochondria was associated with reductions in calculated ATP generation and

proton leak as well as a small decrease in spare respiratory capacity (SRC) (Fig. 9D-F,

respectively). In contrast, the rates of glucose-stimulated extracellular acidification - cytosolic

reactions and a surrogate that can approximate glycolytic activity - were unaffected (Fig. 9G).

Thus, the effects of PexRAP on B cell metabolism are not global, in that altered respiration did

not reflect a decrease in glucose-stimulated acidification. Inasmuch as peroxisomes and

mitochondria form contacts and functionally interact with the ER, we measured relative ER mass

and found that the signal of the fluorophore ERTracker was consistently reduced in PexRAP-

depleted B cells (Fig. 9H, I). We infer that both mitochondrial function and ER mass in B cells

are promoted by the product of the *Dhrs7b* gene.

DISCUSSION

We have shown herein that B lymphocyte expression of an enzyme essential for biosynthesis

of a subset of ether phospholipids is crucial for part of the increases in their levels localized to

GC. Moreover, the genetic approach used to deplete B cells of PexRAP allowed us to show that

the B cell type-restricted gene, *Dhrs7b*, promotes B cell survival, proliferation, and GC size

along with affinity maturation and serum concentrations of Ag-specific Ab after immunization.

Taken together, this work (i) establishes the predictive value of the discovery approach that

identified an unexpected feature of cellular biochemistry in GC, (ii) indicates that B cell-

autonomous generation of ether lipid species is crucial for B cell survival, and (iii) regulates the

qualities of the Ab elicited by immunization.

Using imaging mass spectrometry for discovery-based hypothesis generation, we had

reported that the concentrations of at least a dozen ether phospholipids are increased in splenic

GC relative to the remainder of the tissue (45). The analyses presented here confirm and extend

the observations, and provide evidence that after immunization, these concentrations increase

even in the primary B cell follicle, albeit to a modest degree when compared to the magnitude of

increases in GC, and in B cells activated ex vivo. The cell type-specific gene inactivation after

establishment of a pre-immune population establishes that an enzyme crucial for generation of a

number of plasmalogens has substantial effects on B lymphocyte physiology and function.

Plasmalogens and other ether lipids and phospholipids have long been shown to be major

constituents of lipid bilayers and cell membranes (48, 49, 57). However, remarkably little is

known about whether or not cell-intrinsic synthesis of ether lipids or the balance among their

specific molecular species matters for hematopoietic cells or in immunity. Several inborn errors

of metabolism in humans are attributable to loss-of-function mutations of genes encoding

peroxisomal proteins that impact ether lipid (and hence plasmalogen) synthesis (58) but affect

other critical processes as well [reviewed in (48, 58-62)]. Several of these - such as Zellweger

Syndrome and rhizomelic chondrodysplasia punctata (RCDP) - and the mouse models generated

to study mechanisms in these human disease, lead to severe neurological defects and early post-

partum death (57, 58). Whether or not lymphocyte development, homeostasis or function were

affected in these studies is not clear. Deficient generation of ether lipids is among many

abnormalities caused by elimination of fatty acid synthase (FAS) (46, 47). Acute inactivation of

Fasn, the gene encoding this enzyme, in young mature mice caused a decrease in spleen size

disproportionate to the reduction in neutrophils, which was the most profound hematological

consequence (46). In addition, circulating lymphocytes were reduced despite normal steady-state

representation in the marrow. In-trans cell-extrinsic functions of ether lipids have been noted

previously - specifically, as ligands for the invariant natural killer T cell receptor (77) or for a G-

protein coupled receptor on natural killer cells (78). As such, the widespread loss of FAS

function and the potential for indirect and pleiotropic effects of this enzyme deficiency left

unresolved what cells actually depend on their own synthesis of ether lipids.

PexRAP, the enzyme encoded by the *Dhrs7b* gene, generates intermediates vital for the

synthesis of some - but not all - ether lipids in cells such as neutrophils (46). We interpret our

lipidomic data as indicating that this is the case for B lymphocytes. Specifically, although a

number of plasmalogen or other ether lipid species in the naive, activated, or, in IMS, GC B cells

were lower as a consequence of cell type-specific gene inactivation, other phospholipids in these

classes were unaffected. Analyses of neutrophil numbers and survival provided evidence that this

enzyme can be crucial for a normal membrane composition. Moreover, PexRAP promoted

viability of this very short-lived cell type whose membranes have a high plasmalogen content

(46). The bloodborne population of lymphocytes was reduced in the *Rosa*26-CreER^{T2}, *Dhrs7b*$^{f/f}$

mice (46), but the specific cell types were not reported and cells in circulation may not reflect the

spleen or other organs. Of note, the interpretation that the neutrophil phenotype is due to

insufficiency of ether lipids or plasmalogens has been questioned based on alternative gene

disruption results (54). Several models of unconditional gene inactivation or mutation that

drastically reduced the phospholipid end-products were noted to have normal neutrophils

[summarized in (54), albeit at older ages (4 - 7 mo) than those used in (46). However, these

caveats are subject to points articulated in (53). There may be age-related changes in the

processes analyzed herein using young (age ~ 2 mo) mice (53). Moreover, cellular and

organismal selection for adaptation variants makes the results of acute loss-of-function inherently

different from unconditional loss and heterozygote parentage (53). The potential for toxicity

induced by 4OHT activation of CreER^{T2} was among concerns mooted about the earlier study of

neutrophils (54). This concern may have been extrapolated from papers that noted impairment in

earlier hematopoiesis when *Rosa26*-CreER^{T2} was combined with very intense regimens of

tamoxifen administration to infant mice (65, 79). Such studies used individual doses over 4-fold

higher that those used herein, and delivered by gavage five times on a daily basis instead of

three. Nonetheless, the IgM$^+$ B cell population sizes in spleen and marrow were normal even in

the face of reduced erythropoiesis and thymopoiesis (65). Moreover, the analyses herein

compared B lineage-specific deletion to CreER$^{T2+/-}$, *Dhrs7b*$^{+/+}$ mice identically dosed in parallel

with the test subjects. Thus, our work indicates that PexRAP is essential for normal physiology

and function of mature B lymphocytes, in particular after their activation.

The findings are most consistent with a capacity for PexRAP function to contribute towards

B lymphocyte survival and effective proliferation by restraining both executioner caspase

activation and lipid peroxidation due to excessive ROS levels. For instance, population

expansion of *Dhrs7b* Δ/Δ B cells was severely undermined after mitogenic stimulation in vitro

with little difference in the division history but higher frequencies of annexin V-positive B cells.

These findings are consistent with those obtained with neutrophils (46). On the surface such

results might appear to present a conundrum relative to work indicating that both fatty acyl-CoA

reductase 1 and the enzyme that later generates an alkenyl bond at a terminal step in plasmalogen

biosynthesis are crucial for the susceptibility of HT-1080 and 786-O tumor cell lines to

ferroptosis (55). However, our data are consistent with published evidence that PexRAP is

needed for a narrower subset of ether lipids than the broad requirement for FAS or fatty acyl-

CoA reductases (46-49, 80, 81). Moreover, molecular dissection of mechanisms in various non-

hematopoietic cell types (including the cancer line 786-O) showed that ether lipid biosynthesis

can either promote susceptibility or resistance to ferroptosis (82). Collectively, our findings

suggest that for B cells the inactivation of PexRAP tilts the balance toward death and reduced Ab

- perhaps involving lower fractions of poly-unsaturated fatty acids (PUFA) (82), or a lower ratio

of plasmalogens (alkenyl linkages) relative to alkyl-linked ether lipids.

PexRAP localizes to peroxisomes (35-37), and prior work has suggested both that

peroxisomes are increased in GC B cells and likely contribute to FAO by these cells (26). In line

with the possibility that this organelle may affect PexRAP-dependent survival, our data indicated

that specific inhibition of peroxisomal FAO could collaborate with sub-optimal ROS scavenging

to mitigate the reduced growth of activated B cells. However, recent findings indicate that

PexRAP can also be expressed in the ER and is capable of contributing to ether lipid

biosynthesis via catalysis in this organelle in addition to or instead of the peroxisome (56). Thus,

although ROS generated in peroxisomes likely contribute to the stress experienced by PexRAP-

deficient B cells, the phenotypes observed herein may not exclusively reflect a need for the

peroxisomal fraction of the enzyme but instead might include functions of ER-localized protein.
495

Our findings with an intermediary in ether lipid biosynthesis add to an increasing body of

evidence that metabolism in B lymphocytes affects their differentiation or function (28-36).

Many sources and forms of Ab can be protective, while in several autoimmune diseases these

molecules can drive pathology (1, 2). The capacity to generate increased affinities and circulating

concentrations of Ab is an important factor both in protective immunity and auto-immune

disease. Antigen-independent, TLR-driven PC differentiation requires increased oxidative

phosphorylation (70), while oxidative metabolism in GC B cells promotes affinity maturation by

unknown mechanisms (33-35). While not the exclusive determinant of such properties, GC are

major sources of higher affinity Ab with a broader spectrum of specificities (9, 13, 14). Several

lines of evidence suggest that PexRAP affects the physiology of GC B cells during their

residence. Rates of cell cycling as measured with BrdU incorporation appeared reduced, while

ROS levels within GC B cells were higher along with the frequency of death. Because GC Tfh

cells provide new stimulation to GC B cells that present peptide recognized by the TCR, some of

these effects may be secondary consequences of a substantially lower GC-Tfh population even

when B cell-specific loss-of-function was tested. A speculative corollary is that the observed

reduction in ER mass might be functionally notable. For instance, cognate B-T interaction in the

GC involves BCR-mediated internalization of Ag followed by processing and MHC-II

presentation of T cell epitope peptide. Moreover, ether lipids and plasmalogens are presented to

innate-like T cells via CD1d (77). In addition to the GC-localized abnormalities, GC formation,

size, and function also depend on the efficiency of population growth for B cells in a phase

before they enter and take on characteristics of GC B cells, and then homeostatic and

differentiative processes while in the secondary follicle. Thus, while the findings with the post-

immunization inactivation of *Dhrs7b* using *S1pr2*-CreER[T2] provide evidence that the expression

of PexRAP is important within GC B cells, the decreases of GC and affinity-matured Ab are

likely also to involve a pre-GC phase of clonal expansion. In any case, the rapidity of the effects

suggests that pharmacological inhibition of PexRAP may be a target in inflammatory disease in

which B cells participate, particularly in light of the potential for impacts on pro-inflammatory

neutrophils.

MATERIALS & METHODS

Reagents Monoclonal antibodies (mAb) against mouse CD40, CD90.2, B220 (CD45R), and other mAbs (purified, biotinylated, or fluorophore-conjugated) were from BD Biosciences or Tonbo Biosciences (San Diego CA) unless otherwise indicated. BAFF was from AdipoGen (San Diego, CA). Recombinant mouse IL-4 and recombinant mouse IL-5 were from Peprotech (Rocky Hill NJ). Glucose, 2-deoxyglucose, oligomycin, rotenone, antimycin A, carbonyl cyanide 4-phenylhydrazone (FCCP), N-acetyl-cysteine (NAC), thioridazine hydrochloride, H_2O_2, and 4-hydroxytamoxifen (4-OHT) were from Sigma-Aldrich Chemicals (St. Louis MO), and menadione from Cayman Chemicals (Ann Arbor, MI). Tamoxifen (Tmx) was from APExBio Technology (Houston TX). NP-BSA and NP-PSA (for capture in ELISA) as well as NP-OVA (for immunization) were obtained from Biosearch Technology (Novato CA). SRBCs (sheep red blood cells) were from Thermo Fisher Scientific (Waltham MA). CellTrace^{TM} Violet cell proliferation kit, CM-H$_2$DCFDA, MitoSOX^{TM} Red, and BODIPY^{TM} 581/591 C11 were obtained from Invitrogen (Waltham, MA).

Mice, immunizations, ELISA and ELISpot All animal protocols were reviewed and approved by the Vanderbilt University Institutional Animal Care and Use Committee. Mice were housed in ventilated micro-isolators under Specified Pathogen-Free conditions in a Vanderbilt University Medical Center mouse facility and used both male and female mice at 6–8 weeks of age. For cell type specific inactivation of *Dhrs7b* genes, *Dhrs7b*$^{f/f}$ mice (46) were crossed with *Rosa26-*CreER^{T2} (46), *huCD20-*CerER^{T2} (30), or *S1pr2-*CreER^{T2} strains (51, 52), all on C57/BL6 genetic background. Tamoxifen was administered as previously reported (25, 30). To control for potential Cre toxicity, CreER^{T2} mice were similarly injected to use as wild-type (WT) controls. Mice (ages ~ 8wk) were immunized with SRBCs (2×10^8 cells per mouse) and analyzed 1 week after immunization as described (25, 45). Alternatively, mice were immunized and boosted by i.p. injections [each of 100 µg NP$_{16}$-OVA (Biosearch Technologies, Novato, CA) emulsified in

100 µL of Imject^{TM} Alum (Thermo Fisher Scientific, Pittsburgh, PA)] as described previously

(25, 30), and harvested for analyses 1 wk after the 2nd injection. Isotype-specific relative levels of

Ag-specific Ab were quantitated by capture ELISA using Ag (NP$_{20}$-BSA or NP$_2$-PSA for all- or

high-affinity hapten-specific Abs, respectively) followed by SBA Clonotyping System (Southern

Biotech, Birmingham AL), as described previously (25, 30). As previously described (30),

frequencies of Ab-secreting cells (ASCs) were analyzed by ELISpot and quantitated using an

ImmunoSpot Analyzer (Cellular Technology, Shaker Heights OH).

Cell cultures, proliferation assay, and reversion of population growth   Splenic B

cells were purified (90-95%) using negative selection as previously described (30) or by positive

selection with anti-mouse B220 microbeads (Miltenyi Biotech, Auburn, CA). B cells (5×10^5

cells in 1 ml) were activated with anti-CD40 (1 µg/mL) and cultured with BAFF (10 ng/mL),

IL-4 (10 ng/mL), IL-5 (10 ng/mL) and 4-OHT (50 nM) in Iscoves Modified Dulbecco's Medium

(IMDM) supplemented with 10% Fetal Bovine Serum (FBS), 100 U/mL penicillin 100 µg/mL

streptomycin (Invitrogen), 3 mM L-glutamine, nonessential amino acids (Invitrogen), 0.1 mM 2-

ME (Sigma-Aldrich). To analyze the effect of ROS scavenger and inhibition of peroxisomal lipid

oxidation on population growth, bead-purified B cells from tamoxifen-treated *huCD20*-CerER^{T2}

or *Dhrs7b$^{f/f}$*; *huCD20*-CerER^{T2} mice were activated with anti-CD40, cultured with BAFF, IL-4

and IL-5 in the presence or absence of NAC (1 mM and 5 mM) and/or thioridazine·HCl (100

µM) for 5 days. For enhancement of ROS in WT B cells, purified B cells were analyzed 3 d after

activation as for ROS scavenging, but vehicle, H$_2$O$_2$ (200 µM) or menadione (8 µM) was added

at three different time points (overnight, 3 hours, and 1 hour prior to processing for flow

cytometry to counting and measurements of DCFDA, annexin V, and C11 Bodipy signals.

For the analysis of proliferation in vitro, B cells were purified by depleting CD90.2$^+$ T

cells and CD138$^+$ cells using biotinylated anti-CD90.2 Ab and biotinylated anti-CD138 Ab

followed by streptavidin-conjugated microbeads (iMag^{TM}; BD Bioscience), labeled with 5 µM

CellTrace^{TM} Violet (CTV, Invitrogen), activated and cultured as above. To analyze proliferation

in vivo, B cells were purified, labeled with CTV, and injected intravenously into B cell deficient

μMT recipient mice. Mice were harvested at 4 days after adoptive transfer. To measure

proliferation rates of GC B cells, in vivo BrdU incorporation was performed by injecting mice

with BrdU (2 mg/mouse; BD-Pharmingen, San Jose, CA) at 16 h and 4 h before harvesting

spleens from SRBC-immunized mice. Cells were stained for surface markers (B220, IgD, GL7,

CD38) followed by fixation, permeabilization, and staining with anti-BrdU-APC Ab using the

BrdU Flow Kit (BD-Pharmingen) according to manufacturer's protocol.

Flow cytometry - general and measurements of ROS, lipids peroxidation, and ER   GC B

cells were identified as $GL7^+$ $CD95^+$ events in the viable $B220^+$ $dump^-$ gate (dump channel

consisting of one fluorophore for CD11b, CD11c, F4/80, Gr1, TCRβ, and 7AAD) , while

plasmablasts were defined as $B220^+$ $CD138^+$ $TACI^+$ $dump^-$ cells. For flow analyses of total

intracellular ROS and mitochondrial superoxide, B cells (1×10^6 cells) were washed with PBS

and stained with 1.25 μM CM-H$_2$DCFDA or 5 μM MitoSOX^{TM} Red in PBS (20 min at 37°C),

respectively, then washed with 1% BSA containing PBS, and further stained with anti-B220,

anti-CD19, anti-CD138, and Ghost-BV510. To measure the lipid peroxidation, cultured B cells

(1×10^6) were washed with PBS and stained with 1.25 μM C11-BODIPY^{TM} in PBS (20 min at

37°C), and washed with 1% BSA containing PBS followed by surface staining as above. To

analyze the level of endoplasmic reticulum (ER), B cells (1×10^6) were stained with 0.5 μM ER-

Tracker^{TM} Green (Molecular Probes, Eugene OR) in Hank's Balanced Salt Solution with calcium

and magnesium (HBSS/Ca/Mg; Gibco) for 30 min at 37°C.

For measurements of cell death (78), cultured B cells (1×10^6) were washed twice with PBS,

once with Annexin V binding buffer (10 mM HEPES, pH 7.4, 140 mM NaCl, 2.5 mM CaCl$_2$),

and then incubated (15 min at 20 C in the dark) with Annexin V-rPE, 7AAD, APC-conjugated

CD138, APC-Cy7-conjugated B220, e450-CD19. Cells were then washed with Annexin V

binding buffer and analyzed on the flow cytometer. To measure levels of cleaved caspase-3 (25),

an activated apoptosis executor, the cultured cells were rinsed with PBS after harvest, stained (15

min at 4 C) with 7AAD, APC-conjugated CD138, APC-Cy7-conjugated B220, e450-CD19,

washed (FBS, 1% v/v in PBS), then fixed with paraformaldehyde (4% w/v, 10 min at 20 C),

permeabilized with permeabilization buffer (saponin, 0.2% w/v with FBS 1% in PBS), stained

with FITC-conjugated Ab specific for cleaved caspase-3 (BD Biosciences), rinsed, and analyzed

by flow cytometry.

Immunohistochemistry, MALDI-IMS, and IMS data analysis   Mice were immunized with

SRBC and spleens were harvested at 1 week after immunization. Spleens were snap frozen on

dry ice, and mounted to a cryostat chuck with a minimal amount of OCT (Thermo Fisher

Scientific, San Jose, CA). Frozen spleens were cut at 12 µm thickness using a Leica CM3050S

(Leica, IL, USA) at -20 ℃, fixed with 1% paraformaldehyde for 10 min, washed twice with PBS,

and blocked with M.O.M™ (Vector Lab) followed by incubation with GL7-FITC, anti-IgD-PE,

and anti-CD35-biotin Ab followed by streptavidin-conjugated Alex647at 4 ℃. After the tile

scanning of spleen sections, GC size and numbers were quantified with FIJI Image J software.

622   For Matrix-Assisted Laser Desorption/Ionization (MALDI) Imaging Mass Spectrometry

(IMS), frozen spleens were cut as above, and mounting onto indium tin oxide (ITO) coated

microscope slides (Delta Technologies ETC). Slides were then vacuum desiccated for at least 30

minutes before matrix application. Pre-IMS autofluorescence (AF) images were acquired prior to

matrix application on a Zeiss Axio Scan Z1 Slide Scanner using the brightfield channel and the

DAPI (ex. 340-380 nm; em. 435-485 nm, blue), EGFP (ex. 450-490 nm; em. 500-550 nm,

green), DsRed (ex. 538-562 nm; em. 570-640 nm, red) filter cubes. After taking AF images, a

custom in-house developed sublimation device was used to apply the matrices 2,5-

dihydroxyacetophenone (DHA) and 1,5-diaminonaphthalene (DAN) (Sigma Aldrich, St. Louis,

MO, USA) to tissue sections for positive and negative ion mode analyses, respectively (78). For

immunized AID-GFP mice, the fluorescence emissions identified GC localization within the

section used next for IMS. Alternatively, two serial sections of immunized mice spleens were

used, one for 2D-IMS and the contiguous section for immunohistochemistry (IHC). Fluorescence

data were registered with the 2D-IMS visualizations of phospholipids to align images (45),

allowing quantitation of the ion intensities in specific m/z peaks within defined micro-anatomic

portions of the spleen (Fig. 4B). MALDI IMS data were acquired with a 10 µm pixel size (laser

spot size 8 µm) in full scan mode using a Bruker trapped ion mobility time-of-flight (timsTOF)

Flex mass spectrometer (Bruker Daltonics Billerica, MA, USA). Data were acquired with 250

shots per pixel and a mass range of *m/z* 200-2000. SCiLS (software Bruker Daltonics) was used

to process the data, normalize ion intensity, visualize ion images, and merge the images.

LC-MS for lipidomics   To prepare samples, a one-phase method was used to extract lipids

(85). Briefly, 0.5 mL of MeOH/MTBE/CHCl$_3$ mix (1.3:1:1) was added to a frozen pellet of B-

cells (1x10^6 cells total), spiked with 10 uL EquiSPLASH-lipidomics internal standard mix

(Avanti Research), briefly vortexed and shaken gently for 20 min, followed by centrifugation at

20,000 x *g* for 15 min at 10°C. The supernatant was transferred to a clean Eppendorf tube,

evaporated under a gentle stream of N$_2$ gas, and resuspended in 100 uL methanol/CHCl$_3$ (9:1)

and 2uL were used for LC-HRMS (high-resolution MS) analysis. Each sample was injected two

times - one injection in positive ESI mode followed by one in negative mode. Pooled QCs were

injected to assess the performance of the LC and MS instruments at the beginning, in the middle

and at the end of each sequence. Discovery lipidomics data were acquired using a Vanquish

UHPLC (ultrahigh performance liquid chromatography) system interfaced to a Q Exactive HF

quadrupole/orbitrap mass spectrometer (Thermo Fisher Scientific).

Chromatographic separation was performed with a reverse-phase Acquity BEH C18 column

(1.7 mm, 2.1x150mm, Waters, Milford, MA) at a flow rate of 250 ul/min. Mobile phases were

made up of 10 mM ammonium formate and 0.1% formic acid in (A) H$_2$O/CH$_3$CN (40:60) and

in (B) CH$_3$CN/ iPrOH (10:90). Gradient conditions were as follows: 0–1 min, B = 20 %; 1–8

min, B = 20– 100 %; 8–10 min, B = 100 %; 10–10.5 min, B = 100–20 %; 10.5–15 min, B =

20%. The total chromatographic run time was 15 min. Mass spectra were acquired over a

precursor ion scan range of m/z 200 to 1,600 at a resolving power of 60,000 using the

following HESI-II source parameters: spray voltage 4 kV (3 kV in negative mode); capillary

temperature 250°C; S-lens RF level 60 V; N_2 sheath gas 40; N_2 auxiliary gas 10; auxiliary gas

temperature 350°C. MS/MS spectra were acquired for the top-seven most abundant precursor

ions with an MS/MS AGC target of 1e5, a maximum MS/MS injection time of 100 ms, and a

normalized collision energy of 15, 30, 40. High resolution mass spectrometry data were

processed with MS-DIAL version 4.70 in lipidomics mode (86). MS1, and MS2 tolerances

were set to 0.01 and 0.025 Da respectively. Minimum peak height was set to 30,000 to

decrease the number of false positive hits. Peaks were aligned on a quality control (QC)

reference file with RT tolerance of 0.1 min and mass tolerance of 0.015 Da. Default lipid

library was used (Msp20210527163602_converted.lbm2), solvent type was set to $HCOONH_4$

to match the solvent used for separation, and the identification score cut off was set to 80%.

All lipid classes were made available for the search. After lipid identification was completed,

MS-DIAL results were exported into Excel and cleaned using minimum RSD for QC samples

set to 20% and minimum ratio of QC to Blank set to 10. For species identified in both

PexRAP-depleted (*Dhrs7b* Δ/Δ) B cells and controls, mean levels (areas under peak curves)

and their variance were analyzed in Excel.

Metabolic flus analyses   Oxygen Consumption Rate (OCR) were measured using Seahorse

XF96 extracellular flux analyzer (Agilent Technology, Santa Clara, CA) as described previously

(30). Briefly, in vitro activated B cells were washed twice, resuspended in XF Base Media

(Agilent Technologies) supplemented with 2 mM L-glutamine, and equal numbers of B cells (2

$\times 10^5$) were plated on extracellular flux assay plates (Agilent Technologies) coated with 2.5

µg/mL CellTak (Corning) according to the manufacturer's protocol. Before extracellular flux

analysis, B cells were rested (25 minutes at 37°C, atmospheric CO_2) in XF Base Media. OCR

and ECAR were measured before and after the sequential addition of 1.5 µM oligomycin, 0.5

µM FCCP and 0.5 µM rotenone/antimycin A.

689
690

ACKNOWLEDGEMENTS

The experimental work was funded by NIH Grants to VUMC [R01 AI113292 (M.R.B), R01

HL106812 (M.R.B.), followed by R21 AI164760 (M.R.B.) and Vanderbilt University Medical

Center Pathology-Microbiology-Immunology departmental funds]. Mass spectrometry imaging

was supported in part by NIH grant P41 GM103391 to Vanderbilt University (R.M.C.).

Additional support for M. A. J. was provided by the National Science Foundation, NSF DGE-

1445197. NIH Shared Instrumentation Grant 1S10OD018015 as well as scholarships via the

Cancer Center Support Grant (CA068485) and Diabetes Research Center (DK0205930) helped

defray costs of Vanderbilt Cores. We thank L. Clark for a critical reading and comments on the

revised manuscript, J. Cyster for generously expediting shipment of *S1pr2*-CreERT2 breeding

stock, and Vanderbilt institutional cores (High-Throughput Screening; Flow Cytometry Shared

Resource; Mass Spectrometry Shared Resource; Cell & Developmental Biology) for equipment,

expertise, and assistance.

705
706
References

1. Geyer CE, Mes L, Newling M, den Dunnen J, Hoepel W. 2021. Physiological and

pathological inflammation induced by antibodies and pentraxins. *Cells* **10:** 1175.

2. McGettigan SE, Debes GF. 2021. Immunoregulation by antibody secreting cells

in inflammation, infection, and cancer. *Immunol Rev*. **303:** 103-118.

3. Lu LL, Suscovich TJ, Fortune SM, Alter G. 2018. Beyond binding: antibody effector

functions in infectious diseases. *Nat Rev Immunol.* **18:** 46-61.

4. Allman D, Pillai S. 2008. Peripheral B cell subsets. *Curr Opin Immunol.* **20**: 149-157.

5. Martin F, Kearney JF. 2000. B-cell subsets and the mature preimmune repertoire. Marginal

zone and B1 B cells as part of a "natural immune memory". *Immunol Rev*. **175:** 70-79.

6. MacLennan, IC, Toellner KM, Cunningham AF, Serre K, Sze DM, Zuniga E, Cook MC,

Vinuesa CG. 2003. Extrafollicular antibody responses. *Immunol Rev* **194:** 8-18.

7. Hsu, M. C., K. M. Toellner, C. G. Vinuesa, and I. C. Maclennan. 2006. B cell clones that

sustain long-term plasmablast growth in T-independent extrafollicular antibody

responses. *Proc Natl Acad Sci USA* **103:** 5905-5510.

8. Rajewsky K. 1996. Clonal selection and learning in the antibody system. *Nature* **81:** 751-

758.

9. Boothby MR, Hodges E, Thomas JW. 2019. Molecular regulation of peripheral B cells and

their progeny in immunity. *Genes Dev.* **33:** 26-48.

10. Rubtsova K, Rubtsov AV, van Dyk LF, Kappler JW, Marrack P. 2013. T-box transcription

factor T-bet, a key player in a unique type of B-cell activation essential for effective viral

clearance. *Proc Natl Acad Sci USA* 110: E3216-24.

11. Alemán OR, Rosales C. 2023. Human neutrophil Fc gamma receptors: different buttons for

different responses. *J Leukoc Biol.* 114: 571-584.

12. Xu Z, Zan H, Pone EJ, Mai T, Casali P. 2012.Immunoglobulin class-switch recombination:

induction, targeting, and beyond. *Nat Rev Immunol.* **12:** 517-531.

13. Elsner RA, Shlomchik MJ. 2020. Germinal Center and extrafollicular B cell responses in

vaccination, immunity, and autoimmunity. *Immunity* **53:** 1136-1150.

14. Good-Jacobson, K. L., and M. J. Shlomchik. 2010. Plasticity and heterogeneity in the

generation of memory B cells and long-lived plasma cells: the influence of germinal

center interactions and dynamics. *J Immunol* **185:** 3117-3125.

15. Shlomchik M, Mascelli M, Shan H, Radic MZ, Pisetsky D, Marshak-Rothstein A, Weigert

M. 1990. Anti-DNA antibodies from autoimmune mice arise by clonal expansion and

somatic mutation. *J Exp Med*. **171:** 265-292.

16. Thomas JW, Hulbert C. 1996. Somatically mutated B cell pool provides precursors for

742   insulin antibodies. *J Immunol.* **157:** 763-771.

743   17. De Silva NS, Klein U. 2015. Dynamics of B cells in germinal centres. *Nat Rev Immunol.* **15:**

744   137-148.

745   18. Cyster JG, Allen CDC. 2019. B cell responses: Cell interaction dynamics and decisions. *Cell*

746   **177:** 524-540.

747   19. Victora, G. D., and M. C. Nussenzweig. 2022. Germinal Centers. *Annu Rev Immunol* **40:**

748   413-442.

749   20. Guttormsen HK, Sharpe AH, Chandraker AK, Brigtsen AK, Sayegh MH, Kasper DL. 1999.

750   Cognate stimulatory B-cell-T-cell interactions are critical for T-cell help recruited by

751   glyco-conjugate vaccines. *Infect Immun.* **67:** 6375-6384.

752   21. Kim W, Zhou JQ, Horvath SC, Schmitz AJ, Sturtz AJ, Lei T, Liu Z, Kalaidina E, Thapa M,

753   Alsoussi WB, Haile A, Klebert MK, Suessen T, Parra-Rodriguez L, Mudd PA, Whelan

754   SPJ, Middleton WD, Teefey SA, Pusic I, O'Halloran JA, Presti RM, Turner JS, Ellebedy

755   AH. 2022. Germinal centre-driven maturation of B cell response to mRNA vaccination.

756   *Nature* **604:** 141-145.

757   22. Roco JA, Mesin L, Binder SC, Nefzger C, Gonzalez-Figueroa P, Canete PF, Ellyard J, Shen

758   Q, Robert PA, Cappello J, Vohra H, Zhang Y, Nowosad CR, Schiepers A, Corcoran LM,

759   Toellner KM, Polo JM, Meyer-Hermann M, Victora GD, Vinuesa CG. 2019. Class-

760   switch recombination occurs infrequently in Germinal Centers. *Immunity* **51:** 337-350.

761   23. Gitlin AD, Shulman Z, Nussenzweig MC 2014. Clonal selection in the germinal centre by

762   regulated proliferation and hypermutation. *Nature* **509:** 637-40.

763   24. Vinuesa CG, Linterman MA, Yu D, MacLennan IC. 2016. Follicular helper T cells. *Annu*

764   *Rev Immunol.* **34:** 335-368.

25. Mayer CT, Gazumyan A, Kara EE, Gitlin AD, Golijanin J, Viant C, Pai J, Oliveira TY,

Wang Q, Escolano A, Medina-Ramirez M, Sanders RW, Nussenzweig MC. 2017. The

micro-anatomic segregation of selection by apoptosis in the germinal center. *Science* **358:**

eaao2602.

26. Chapman NM, Chi H. 2022. Metabolic adaptation of lymphocytes in immunity and disease.

Immunity **55:** 14-30.

27. Bacigalupa ZA, Landis MD, Rathmell JC. 2024. Nutrient inputs and social metabolic control

of T cell fate. *Cell Metab.* **36:** 10-20.

28. Jellusova J. 2020. Metabolic control of B cell immune responses. *Curr Opin Immunol.* **63:**

21-28.

29. Boothby MR, Brookens SK, Raybuck AL, Cho SH. 2022. Supplying the trip to antibody

production-nutrients, signaling, and the programming of cellular metabolism in the

mature B lineage. *Cell Mol Immunol.* **19:** 352-369.

30. O'Neill LA, Kishton RJ, Rathmell J. 2016. A guide to immunometabolism for

immunologists. *Nat Rev Immunol.* **16:** 553-565.

31. Cho, S. H., A. L. Raybuck, K. Stengel, M. Wei, T. C. Beck, E. Volanaki, J. W. Thomas, S.

Hiebert, V. H. Haase, and M. R. Boothby. 2016. Germinal centre hypoxia and regulation

of antibody qualities by a hypoxia response system. *Nature* **537:** 234-237.

32. Weisel, F. J., S. J. Mullett, R. A. Elsner, A. V. Menk, N. Trivedi, W. Luo, D. Wikenheiser,

W. F. Hawse, M. Chikina, S. Smita, L. J. Conter, S. M. Joachim, S. G. Wendell, M. J.

Jurczak, T. H. Winkler, G. M. Delgoffe, and M. J. Shlomchik. 2020. Germinal center B

cells selectively oxidize fatty acids for energy while conducting minimal glycolysis. *Nat*

Immunol **21:** 331-342.

33. Chen D, Wang Y, Manakkat Vijay GK, Fu S, Nash CW, Xu D, He D, Salomonis N, Singh H,

Xu H. 2021. Coupled analysis of transcriptome and BCR mutations reveals role of

OXPHOS in affinity maturation. *Nat Immunol.* 22: 904-913.

34. Urbanczyk S, Baris OR, Hofmann J, Taudte RV, Guegen N, Golombek F, Castiglione K,

Meng X, Bozec A, Thomas J, Weckwerth L, Mougiakakos D, Schulz SR, Schuh W,

Schlotzer-Schrerhardt U, Steinmetz TD, Brodesser S, Wiesner RJ, Mielenz D. 2022.

Mitochondrial respiration in B lymphocytes is essential for humoral immunity by

controlling the flux of the TCA cycle. *Cell Rep.* **39:** 110912.

35. Yazicioglu Y. F., E. Marin, C. Sandhu, S. Galiani, I. G. A. Raza, M. Ali, B. Kronsteiner, E.

B. Compeer, M. Attar, S. J. Dunachie, M. L. Dustin, and A. J. Clarke. 2023. Dynamic

mitochondrial transcription and translation in B cells control germinal center entry and

lymphomagenesis. *Nat Immunol.* **24:** 991-1006.

36. Brookens SK, Cho SH, Paik Y, Meyer K, Raybuck AL, Park C, Greenwood DL, Rathmell

JC, Boothby MR. 2024. Plasma cell differentiation, antibody quality, and initial germinal

center B cell population depend on glucose influx rate. *J Immunol.* **212:** 43-56.

37. Olenchock BA, Rathmell JC, Vander Heiden MG. 2017. Biochemical underpinnings of

immune cell metabolic phenotypes. *Immunity* **46:** 703-713.

38. Kong H, Chandel NS. 2018. Regulation of redox balance in cancer and T cells. *J Biol Chem.*

293: 7499-7507.

39. Addabbo F, Montagnani M, Goligorsky MS. 2009. Mitochondria and reactive oxygen

species. *Hypertension* **53:** 885-892.

40. Zorov DB, Juhaszova M, Sollott SJ. 2014. Mitochondrial reactive oxygen species (ROS) and

ROS-induced ROS release. *Physiol Rev.* **94:** 909-950.

41. Jellusova, J., M. H. Cato, J. R. Apgar, P. Ramezani-Rad, C. R. Leung, C. Chen, A. D.

Richardson, E. M. Conner, R. J. Benschop, J. R. Woodgett, and R. C. Rickert. 2017.

Gsk3 is a metabolic checkpoint regulator in B cells. *Nat Immunol* **18:** 303-312.

42. Li L, Feng C, Qin J, Li D, Liu M, Han S, Zheng B. 2021. Regulation of humoral immune

response by HIF-1α-dependent metabolic reprogramming of the germinal center reaction.

Cell Immunol. **367:** 104409.

43. Nakagawa R, Llorian M, Varsani-Brown S, Chakravarty P, Camarillo JM, Barry D, George

R, Blackledge NP, Duddy G, Kelleher NL, Klose RJ, Turner M, Calado DP. 2023. Epi-

microRNA mediated metabolic reprogramming ensures affinity maturation. *bioRxiv*

[Preprint]. 2023 Oct 31:2023.07.31.551250.

44. Jain IH, Calvo SE, Markhard AL, Skinner OS, To TL, Ast T, Mootha VK. 2020. Genetic

screen for cell fitness in high or low oxygen highlights mitochondrial and lipid

metabolism. *Cell* **181:** 716-727.e11.

45. Jones MA, Cho SH, Patterson NH, Van de Plas R, Spraggins JM, Boothby MR, Caprioli

RM. 2020. Discovering new lipidomic features using cell type-specific fluorophore

expression to provide spatial and biological specificity in a multimodal workflow with

MALDI Imaging Mass Spectrometry. *Anal Chem.* **92:** 7079-7086.

46. Lodhi IJ, Wei X, Yin L, Feng C, Adak S, Abou-Ezzi G, Hsu FF, Link DC, Semenkovich CF.

2015. Peroxisomal lipid synthesis regulates inflammation by sustaining neutrophil

membrane phospholipid composition and viability. *Cell Metab.* **21:** 51-64.

47. Lodhi IJ, Yin L, Jensen-Urstad AP, Funai K, Coleman T, Baird JH, El Ramahi MK, Razani

B, Song H, Fu-Hsu F, Turk J, Semenkovich CF. 2012. Inhibiting adipose tissue

lipogenesis reprograms thermogenesis and PPARγ activation to decrease diet-induced

obesity. *Cell Metab.* **16:** 189-201.

48. Dean JM, Lodhi IJ. 2018. Structural and functional roles of ether lipids. *Protein Cell* **9:** 196-

206.

49. Honsho M, Fujiki Y. 2023. Regulation of plasmalogen biosynthesis in mammalian cells and

tissues. *Brain Res Bull.* **194:** 118-123.

50. Werner ER, Swinkels D, Juric V, Dorninger F, Baes M, Keller MA, Berger J, Watschinger

K. 2023. Normal plasmalogen levels are maintained in tissues from mice with

hepatocyte-specific deletion in peroxin 5. *Brain Res Bull.* **193:** 158-165.

51. Dorninger F, Werner ER, Berger J, Watschinger K. 2022. Regulation of plasmalogen

metabolism and traffic in mammals: The fog begins to lift. *Front Cell Dev Biol.* **10:**

946393.

52. Kuerschner L, Richter D, Hannibal-Bach HK, Gaebler A, Shevchenko A, Ejsing CS, Thiele

C. 2012. Exogenous ether lipids predominantly target mitochondria. *PLoS One* **7:**

e31342.

53. Lodhi IJ, Link DC, Semenkovich CF. 2015. Acute ether lipid deficiency affects neutrophil

biology in mice. *Cell Metab.* **21:** 652-653.

54. Dorninger F, Wiesinger C, Braverman NE, Forss-Petter S, Berger J. 2015. Ether lipid

deficiency does not cause neutropenia or leukopenia in mice and men. *Cell Metab.* **21:**

650-651.

55. Cui W, Liu D, Gu W, Chu B. 2021. Peroxisome-driven ether-linked phospholipids

biosynthesis is essential for ferroptosis. *Cell Death Differ.* **28:** 2536-2551.

56. Honsho M, Tanaka M, Zoeller RA, Fujiki Y. 2020. Distinct functions of acyl/alkyl

dihydroxyacetonephosphate reductase in peroxisomes and endoplasmic reticulum. *Front*

Cell Dev Biol. **8:** 855.

57. Braverman NE, Moser AB. 2012. Functions of plasmalogen lipids in health and disease.

Biochim Biophys Acta **1822:** 1442-1452.

58. Gorgas K, Teigler A, Komljenovic D, Just WW. 2006. The ether lipid-deficient mouse:

tracking down plasmalogen functions. *Biochim Biophys Acta* **1763:** 1511-1526.

59. Jiménez-Rojo N, Riezman H. 2019. On the road to unraveling the molecular functions of

ether lipids. *FEBS Lett.* **93:** 2378-2389.

60. Rangholia N, Leisner TM, Holly SP. 2021. Bioactive ether lipids: Primordial modulators of

cellular signaling. *Metabolites* **11:** 41.

61. Bozelli JC Jr, Azher S, Epand RM. 2021. Plasmalogens and chronic inflammatory diseases.

Front Physiol. **12:** 730829.

62. Wallner S, Schmitz G. 2011. Plasmalogens the neglected regulatory and scavenging lipid

species. *Chem Phys Lipids* **164:** 573-589.

63. Friedrich, G, Soriano, P. 1991. Promoter traps in embryonic stem cells: A genetic screen to

identify and mutate developmental genes in mice. *Genes Devel.* **5:** 1513–1523.

64. Shapiro-Shelef M, Lin KI, Savitsky D, Liao J, Calame K. 2005. Blimp-1 is required for

maintenance of long-lived plasma cells in the bone marrow. *J Exp Med.* **202:** 1471-1476.

65. Higashi AY, Ikawa T, Muramatsu M, Economides AN, Niwa A, Okuda T, Murphy AJ, Rojas

J, Heike T, Nakahata T, Kawamoto H, Kita T, Yanagita M. 2009. Direct hematological

toxicity and illegitimate chromosomal recombination caused by the systemic activation of

CreERT2. *J Immunol.* **182:** 5633-5640.

66. Uhmann A, Dittmann K, Wienands J, Hahn H. 2009. Comment on "Direct hematological

toxicity and illegitimate recombination caused by the systemic activation of CreERT2". *J

Immunol.* **183:**2891.

67. Khalil AM, Cambier JC, Shlomchik MJ. 2012. B cell receptor signal transduction in the GC

is short-circuited by high phosphatase activity. *Science* **336:** 1178-1181.

68. Inoue T, Kurosaki T. 2024. Memory B cells. *Nat Rev Immunol.* **24:** 5-17.

69. Shinnakasu R, Inoue T, Kometani K, Moriyama S, Adachi Y, Nakayama M,

Takahashi Y, Fukuyama H, Okada T, Kurosaki T. 2016. Regulated selection of germinal-

center cells into the memory B cell compartment. *Nat Immunol.* **17:** 861-869.

70. Price MJ, Patterson DG, Scharer CD, Boss JM. 2018. Progressive upregulation of oxidative

metabolism facilitates plasmablast differentiation to a T-independent antigen.

Cell Rep. **23:** 3152-3159.

71. Capasso M, Bhamrah MK, Henley T, Boyd RS, Langlais C, Cain K, Dinsdale D, Pulford K,

Khan M, Musset B, Cherny VV, Morgan D, Gascoyne RD, Vigorito E, DeCoursey TE,

MacLennan IC, Dyer MJ. 2010. MacLennan IC, Dyer MJ. 2010. HVCN1 modulates

BCR signal strength via regulation of BCR-dependent generation of reactive oxygen

species. *Nat Immunol.* **11:** 265-272.

72. Wheeler ML, Defranco AL. 2012. Prolonged production of reactive oxygen species in

response to B cell receptor stimulation promotes B cell activation and proliferation. *J*

Immunol. **189:** 4405-4416.

73. Tsubata T. 2020. Involvement of reactive oxygen species (ROS) in BCR signaling as a

second messenger. *Adv Exp Med Biol.* **1254:** 37-46.

74. Jiang X, Stockwell BR, Conrad M. 2021. Ferroptosis: mechanisms, biology and role in

disease. *Nat Rev Mol Cell Biol.* **22:** 266-282.

75. Nordgren M, Fransen M. 2014. Peroxisomal metabolism and oxidative stress. *Biochimie* **98:**

56-62.

76. Liang D, Minikes AM, Jiang X. 2022. Ferroptosis at the intersection of lipid metabolism and

cellular signaling. *Mol Cell* **82:** 2215-2227.

77. Facciotti F, Ramanjaneyulu GS, Lepore M, Sansano S, Cavallari M, Kistowska M, Forss-

Petter S, Ni G, Colone A, Singhal A, Berger J, Xia C, Mori L, De Libero G. 2012.

Peroxisome-derived lipids are self antigens that stimulate invariant natural killer T cells

in the thymus. *Nat Immunol.* 3: 474-480.

78. Hossain MS, Mawatari S, Fujino T. 2022. Plasmalogen-mediated activation of GPCR21

regulates cytolytic activity of NK cells against the target cells. *J Immunol.* **209:** 310-325.

79. Rossi M, Salomon A, Chaumontel N, Molet J, Bailly S, Tillet E, Bouvard C. 2023. Warning

regarding direct hematological toxicity of tamoxifen-activated CreERT2 in young

Rosa26-CreERT2 mice. *Sci Rep*. **13:** 5976.

80. Lodhi IJ, Semenkovich CF. 2014. Peroxisomes: a nexus for lipid metabolism and cellular

916   signaling. *Cell Metab.* **19:** 380-392.

917   81. He A, Dean JM, Lodhi IJ. 2021. Peroxisomes as cellular adaptors to metabolic and

918   environmental stress. *Trends Cell Biol*. **31:** 656-670.

919   82. Zou Y, Henry WS, Ricq EL, Graham ET, Phadnis VV, Maretich P, Paradkar S, Boehnke N,

920   Deik AA, Reinhardt F, Eaton JK, Ferguson B, Wang W, Fairman J, Keys HR, Dančík V,

921   Clish CB, Clemons PA, Hammond PT, Boyer LA, Weinberg RA, Schreiber SL. 2020.

922   Plasticity of ether lipids promotes ferroptosis susceptibility and evasion. *Nature* **585:**

923   603-608.

924   83. Esselman AB, Patterson NH, Migas LG, Dufresne M, Djambazova KV, Colley ME, Van de

925   Plas R, Spraggins JM. 2023. Microscopy-Directed Imaging Mass Spectrometry for Rapid

926   High Spatial Resolution Molecular Imaging of Glomeruli. *J Am Soc Mass Spectrom.* **34:**

927   1305-1314.

928   84. Lee K, Heffington L, Jellusova J, Nam KT, Raybuck A, Cho SH, Thomas JW, Rickert

929   RC, Boothby M. 2013. Requirement for Rictor in homeostasis and function of mature B

930   lymphoid cells. *Blood* **122:** 2369-2379.

931   85. Pellegrino RM, Di Veroli A, Valeri A, Goracci L, Cruciani G. 2014. LC/MS lipid profiling

932   from human serum: a new method for global lipid extraction. *Anal Bioanal Chem*. **406:**

933   7937-7948.

934   86. Tsugawa H, Ikeda K, Takahashi M, Satoh A, Mori Y, Uchino H, Okahashi N, Yamada Y,

935   Tada I, Bonini P, Higashi Y, Okazaki Y, Zhou Z, Zhu Z-J, Koelmel J, Cajka T, Fiehn O,

936   Saito K, Arita M, Arita M. 2020. A lipidome atlas in MS-DIAL 4. *Nat Biotechnol*. **38:**

937   1159-1163.

938   87. Tao L, Lemoff A, Wang G, Zarek C, Lowe A, Yan N, Reese TA. 2020. Reactive oxygen

939   species oxidize STING and suppress interferon production. *Elife* **9:** e57837. doi:

940   10.7554/eLife.57837.

    Figure Legends

    Figure 1. PexRAP promotes GC response. (A) Schematic of immunization with SRBC after

    inactivation of *Dhrs7b* in adult mice. Mice (*Rosa26*-CreER^{T2}; *Dhrs7b$^{+/+}$, or Rosa26*-CreER^{T2};

    Dhrs7b^{ff}) were treated with tamoxifen, immunized with SRBC, and harvested 1 wk after

    immunization, as described in *Materials and Methods*. (B) Deletion efficiency of *Dhrs7b*

    conditional alleles. B and T lymphocytes were isolated from spleens of *Rosa26*-CreER^{T2} or

    Dhrs7b$^{f/f}$;Rosa26-CreER^{T2} mice after in vivo tamoxifen injections followed by immunization

    with SRBC. WT and *Dhrs7b$^{Δ/Δ}$* were analyzed by immunoblotting with antibodies directed

    against PexRAP (protein product of *Dhrs7b*) and actin (internal loading control). (C, D) PexRAP

    acutely regulates B cell numbers. Representative flow plots of viable splenic B cells (C) and

    aggregate data for three biologically independent replicate experiments (D) (n = 8 WT and 7

    cKO). For data on T cells, see supplemental data Fig. S1A-C. (E, F) Effect of PexRAP on GC B

    cell response. Flow plots of GL7^{+} CD95^{+} GC B cells in the gate for viable IgD-negative (IgD^{neg}),

    dump-negative B cells (E), and aggregated frequencies and numbers of GC B cells, as indicated,

    in the three replicate experiments (F). (G, H) PexRAP impact on GC response. After tamoxifen

    injections, *Dhrs7b* deficient mice [*Rosa26*-CreER^{T2}; *Dhrs7b^{ff}* (46)] and WT controls (*Rosa26*-

    CreER^{T2}) were immunized with SRBC and analyzed 7 d later, as in Fig. 1A-F. Data on gating

    strategy, GC counts per unit area and LZ area are in Supplemental Fig. 1D-F, and on Tfh cells in

    Supplemental Fig. 1G-I. (G, H) Shown are representative images from immunofluorescence

    staining of spleens with the indicated Ab in two independent experiments (5 WT vs 4 *Dhrs7b*

    cKO, i.e., *Dhrs7b$^{Δ/Δ}$*), showing a low-power overview with many follicles (G) and a

    representative higher-magnification image to better delineate primary and secondary follicles

    (H). (I) Quantitation of number (left panel) and size (right panel) of GC in spleen sections, with

    GL7^{+}, IgD^{neg} areas that include both CD35^{+} (LZ) and CD35^{neg} (DZ) areas. Mann-Whitney U test

    was used to calculate p values.

    Figure 2. PexRAP promotes proliferation of B cells. (A) Schematic of *Dhrs7b* inactivation in

    mature mice via tamoxifen treatment and a B cell type-specific conditional allele. Mice

    (*huCD20*-CreER^{T2}; *Dhrs7b$^{+/+}$, or huCD20*-CreER^{T2}; *Dhrs7b$^{f/f}$*) were injected with tamoxifen

    (d1, 3, 5) and harvested at day 10. (B) Deletion efficiency of conditional *Dhrs7b* alleles. After in

    vivo tamoxifen injections, B cells were isolated from spleens of *huCD20*-CreER^{T2} or *Dhrs7b$^{f/f}$;*

    huCD20-CreER^{T2} mice, as indicated, and analyzed by immunoblotting as in Fig. 1B. (C)

    PexRAP and the maintenance of B cells. Shown are the frequencies and the numbers of CD19^{+}

    B220^{+} B cells among viable lymphocytes in spleen (left and right panels, respectively). Data are

    pooled from four independent replicate experiments (n = 12 WT and 12 cKO). Shown in box and

whisker plots are the means, with whiskers that extend to the minimum and maximum values and

boxes that outline upper and lower quartile values with the midline identifying the median. (D-F)

PexRAP regulates B cell proliferation in vivo. CTV-labeled B cells were adoptively transferred

into µMT recipient mice and analyzed 4 days thereafter. Shown are the frequencies of B220$^+$

CD19$^+$ events among splenocytes in the viable cell gate (D), along with representative flow plots

of CTV partitioning and surface IgD in B cell gates, with rectangles defining the gating for

divided > or ≤ 3x, and IgD$^+$ vs IgD^{neg}. A similar difference was observed in analyzing divided vs

undivided. (E). (F) Aggregated frequencies of divided B cells from five independent replicate

experiments, as defined in (E) (n =10 WT and 11 cKO). Additional data on the IgD^{neg} population

are in Supplemental Fig.2G. (G-I) PexRAP promotes B cell proliferation in vitro. After in vivo

tamoxifen injections, bead-purified B cells from spleens of CreER^{T2} mice (WT and *Dhrs7b$^{Δ/Δ}$*,

i.e., cKO) were stained with Cell Trace Violet (CTV), activated and cultured 4 days in anti-

CD40, BAFF, IL-4, IL-5 and 4-hydroxytamoxifen in three biologically independent experiments

totaling 6 WT and 7 cKO mice). (G) Representative flow-cytometric analysis of CTV

partitioning, with the gating line denoting multiply divided cells, with inset numbers representing

the frequencies of such B cells in each of the two plots shown. (H) Quantified frequencies of

divided ≥3 times. (I) Numbers of B cells recovered at the end of the cultures.

Figure 3. B cell expression of PexRAP is essential for normal concentrations of most ether

phospholipids. Splenic B cells (CreER^{T2+} or CreER^{T2+}, *Dhrs7b* Δ/Δ) from tamoxifen-injected

mice were analyzed by LC-MS-MS either directly ex vivo ("Naive") or after activation and

culture (48 h) ("Activated") as in Fig. 2. Shown are z-scored heat maps for the relative

abundance of subsets of ether and plasmalogen phospholipids and lysophospholipids identified

by exact mass and secondary fragmentation in (A) positive and/or (B) negative ion modes. PE,

phosphatidylethanolamine; PC, phosphatidylcholine; LPE, lysophosphatidylethanolamine. More

species are displayed in Supplemental Fig. 2. (C) Quantitated peak areas for activation-induced

increases ('fold-induction' ratios of activated/resting, plotted on log$_{10}$ scale) for the indicated

ether, plasmalogen, and diacyl (non-ether) phospholipids in the PexRAP-sufficient and -depleted

B cells. * denotes species for which P<0.05 for the effect of *Dhrs7b* inactivation. (D) Shown are

the raw values of mean peak areas for the indicated species in the freshly purified and the

activated B cells of the indicated genotypes, as indicated. (Three independent samples from

individual mice of each genotype were analyzed.) * p<0.05, ** p<0.01 by unpaired Student's. t-

test. Additional data including controls comparing B cells of unmanipulated B6 mice to those

expressing CreER^{T2} and injected with tamoxifen are in Supplemental Fig. 3.

1012

Figure 4. PexRAP is essential for normal concentrations and distributions of some ether
phospholipids in splenic follicles. (A) Schematic diagram illustrating the 2D-IMS analysis
work flow, image merging and quantitation. Spleens harvested 1 wk after immunization with
SRBC were used to generate serial tissue sections followed by immunofluorescence (IF) staining
of one section and IMS analysis with the adjacent one. IF and IMS images were aligned to map
ion intensity distributions to microanatomic regions (B cell follicle and GC). The intensities of
specific ions on B cell follicles and GC regions were quantitated as described in *Materials and*
Methods. (B) Identification of ether lipid species localizing to lymphoid follicles. Representative
ion images of two ions [*m/z* 752.5545, and *m/z* 776.5556] with spleens from immunized mice
(WT and *Dhrs7b*$^{\Delta/\Delta}$) as shown in Fig. 1A. (C) Schematic of immunization for IMS analyses of B
cell-specific PexRAP loss. Mice of the indicated genotypes (huCD20-CreER^{T2} ± *Dhrs7b* f/f)
were treated with tamoxifen, immunized with SRBC, and harvested 1 wk after immunization.
(D) Identification of ether lipid species localizing to lymphoid follicles. Representative ion
images of three ions [*m/z* 752.5545, *m/z* 872.5749, and *m/z* 700.4958] in mass spectrometry
imaging of spleens from WT and *Dhrs7b*$^{\Delta/\Delta-B}$ mice. Immunofluorescent images at higher
magnification, delineating LZ and DZ marked by CD35 staining, along with quantification of
sizes of GC and their LZ and DZ, are shown in Supplemental Fig. 5A, B. (E) Shown in the bar
graphs are the mean (±SEM) ion intensities in primary lymphoid follicles (B cell zones) and GC
from spleens of WT and *Dhrs7b*$^{\Delta/\Delta-B}$ mice, immunized or not ("UI") as indicated. Median
intensity of each ion was obtained from 3 follicles / spleen and 3 GC / spleen from WT and
Dhrs7b$^{\Delta/\Delta-B}$ mice (three biological replications comprising 7 WT and 6 cKO spleens). P values
were calculated by Mann-Whitney U test.
1035
Figure 5. B cell intrinsic role of PexRAP in GC response. (A) Schematic of immunization
with NP-OVA in alum after inactivation of *Dhrs7b* in B lineage cells. Tamoxifen-treated WT
(*huCD20*-CreER^{T2}; *Dhrs7b*$^{+/+}$) or *Dhrs7b*$^{\Delta/\Delta-B}$ mice (*huCD20*-CreER^{T2}; *Dhrs7b*^{ff}) were
immunized with NP-OVA, with sera collected 3 wk thereafter ("1^0 response"), followed by
boosting with NP-OVA and harvest 1 wk after the 2nd immunization. (B, C) Representative flow
plots of splenic IgD^{neg} B cells (B) at the time of harvest (1 wk after 2nd immunization), and
aggregate data (C) for two replicate experiments (n = 5 WT and 5 cKO). The cocktail of reagents
for the dump channel included anti-IgD. (D, E) B cell-intrinsic function of PexRAP in GC B cell
response. Representative flow plots of GL7$^+$ CD95$^+$ GC B cells among viable IgD^{neg} B cells (D),
and aggregated frequencies and numbers of GC B cells for two replicate experiments (E). Mann-
Whitney U test was used to calculate p values. (F, G) WT (*huCD20*-CreER^{T2}; *Dhrs7b*$^{+/+}$) or
Dhrs7b$^{\Delta/\Delta-B}$ mice (*huCD20*-CreER^{T2}; *Dhrs7b*^{ff}) were injected with tamoxifen and immunized
with SRBC as in Fig 4A. Shown are the representative flow plots of GL7$^+$ CD38$^-$ GC B cells

among IgD⁻ CD138⁻ viable B cells (F), and aggregate data (G) for two replicate experiments (n =

6 WT and 6 cKO). (H) PexRAP is dispensable for the balance of LZ and DZ B cells. The graph

shows the mean (± SEM) frequencies (left) and the numbers (right) of $CD86^+$ $CXCR4^{lo}$ LZ B

cells and $CD86^{neg}$ $CXCR4^+$ DZ B cells among IgD^{neg} $CD38^{neg}$ $CD138^{neg}$ $GL7^+$ GC cells. Mann-

Whitney U test was used to calculate p values.

1054

Figure 6. Ab response and affinity increase promoted by PexRAP in B cells. Tamoxifen-

treated mice (*huCD20*-CreER^{T2}; *Dhrs7b$^{+/+}$* or *huCD20*-CreER^{T2}; *Dhrs7b$^{f/f}$* , i.e., *Dhrs7b$^{Δ/Δ-B}$*)

were immunized as in Fig. 5, with venous blood collected to measure Ag-specific Ab in the 1^0

response just prior to a second immunization, followed by harvesting a week thereafter (2^0

response). (A-D) Ag-specific Ab in primary response sera, prior to the boost. Shown are the all-

(NP_{20}) and high- (NP_2) -affinity anti-NP IgM (A, B) and IgG1 (C, D), as indicated, with NP_{20} a

high hapten density to detect both low- and high-affinity Ab and NP_2 a low hapten density

selective for high-affinity Ab. (E, F) Levels of anti-NP IgM detected using NP_{20} and NP_2 for

ELISpot (E) and ELISA (F) as described in *Materials and Methods*. Graphs show mean (± SEM)

number of Ab-secreting cells in spleen (E) and (F) mean (± SEM) OD_{450nm} in measurement of

NP-specific IgM in serial dilutions of sera. P values were calculated by students' t-test. *

indicates p<0.05, and ** indicates p<0.01. (G) PexRAP in B cells promotes Ab affinity

maturation. The bar graph shows the mean (± SEM) ratios of high-affinity to all-affinity NP-

specific IgM Ab in sera of individual mice (each dot representing one subject) using OD_{450nm}

values at the 1:2000 dilution, with data from three independent experiments comprising 8 mice

of each type (WT; *Dhrs7b$^{Δ/Δ-B}$*). P values were calculated by Mann-Whitney U test. (H, I)

PexRAP promotes $CD138^+$ cell differentiation in vivo. B cells were adoptively transferred into

μMT recipient mice and analyzed at 4 days after transfer. Representative flow plot of CD138 and

cD19 expression (H) and aggregated frequencies of $CD138^+$ $CD19^+$ cells in the viable cell gate

(I) from four independent replicate experiments. To test for a potential distortion arising from

outlier values, statistical testing was performed both with and without their inclusion.

1076

Figure 7. Function of PexRAP in GC response. (A) Schematic of the time line, with

immunization followed later by tamoxifen injections into *S1pr2*-CreER^{T2} mice (*Dhrs7b$^{+/+}$*or

Dhrs7b$^{f/f}$). Note that deletion is initiated only just as the germinal center reaction starts (~ 3.5 d

post-immunization). Mice (*S1pr2*-CreER^{T2}; *Dhrs7b$^{+/+}$* or *S1pr2*-CreER^{T2}; *Dhrs7b$^{f/f}$*) were

immunized with NP-OVA, treated with tamoxifen on days 3, 5, and 7 after NP-OVA

immunization (36) and harvested at day 9. (B) Representative flow plots of $GL7^+$ $CD95^+$ GC B

cells among viable B cells and (C) aggregated mean (±SEM) frequencies of GC B cells from

three replicate experiments (7 WT; 9 cKO mice). Mann-Whitney U test was used to calculate p

values.-(D-F) Effect of PexRAP on the levels of ROS (D), cell death (E), and proliferation (F) of

GC B cells. (D) Total cellular ROS in IgD^{neg} CD38^{neg} GL7$^+$ GC B cells were determined by flow

cytometry after staining with surface markers and H$_2$DCFDA as described in the Methods. The

graph shows the mean (±SEM) geometric MFI of H$_2$DCFDA from two independent replicate

experiments (n = 6 WT and 6 cKO). (E) PexRAP promotes GC B cell survival. Shown are the

representative flow plot (left), and a dot graph aggregating all experiments' outcomes for the

frequencies of annexin V$^+$ 7AAD$^+$ cells in GC B gated cells as in Fig 7D (right panel). (F)

PexRAP regulates proliferation of GC B cells. Tamoxifen-treated mice (*huCD20*-CreER^{T2};

Dhrs7b$^{+/+}$ or *huCD20*-CreER^{T2}; *Dhrs7b^{ff}* , i.e., *Dhrs7b$^{Δ/Δ−B}$*) were immunized with SRBC, and

the mice were injected with BrdU as described in Methods. Shown are a representative

histograms for WT and *Dhrs7b$^{Δ/Δ−B}$* GC B cells as indicated (left panel) and a graph indicating

the aggregated result of each independent experiments (right panel) (x=2; n = 6 WT and 6 cKO).

1097

Figure 8. PexRAP contributes to ROS homeostasis and B cell population growth in vitro.

(A-D) PexRAP is critical for maintenance of normal ROS levels. Bead-purified B cells from

spleens of tamoxifen-treated *huCD20*-CreER^{T2} mice (*Dhrs7b$^{Δ/Δ−B}$* and *Dhrs7b$^{+/+}$*) were cultured

3 days in anti-CD40, BAFF, IL-4, IL-5 and 4-hydroxytamoxifen. Total cellular (A, B) and

mitochondrial ROS, mtROS (C, D) in B lymphoblasts were then determined by flow cytometry

after staining with surface markers and H$_2$DCFDA and MitoSOX, as described in the Methods.

Representative histogram image of H$_2$DCFDA (A) and MitoSOX (C) in the B cell gate, and

aggregated mean (± SEM) geometric MFI of H$_2$DCFDA (B) and MitoSOX (D) from 3

independent experiments, each using two mice of each type (6 WT; 6 cKO). P values were

calculated by Mann-Whitney U test. (E) PexRAP restrains lipid peroxidation. B cells were

activated and cultured as in (A). A representative result of flow cytometric analyses of lipid

peroxidation assayed using C11-Bodipy is shown, based on three independent experiments.

(F, G) PexRAP promotes B cell survival. (F) Shown are the mean (±SEM) frequencies of total

viable 'events' (by FSC, SSC, and 7-AAD exclusion) in flow cytometry (filled circles) and MFI

of Bodipy-C11 (open squares) after exposure to H$_2$O$_2$ (200 µM). (*, ** - p=0.03 and 0.003,

respectively). (G-I) WT and *Dhrs7b$^{Δ/Δ}$* B cells were cultured (2 d) in anti-CD40, BAFF, IL-4, IL-

5 and 4-hydroxytamoxifen. (G) Increased cleaved caspase 3 (CC3) in PexRAP-deficient cells

generated in vitro. Cleaved caspase 3 in B cells was detected by intracellular staining and flow

cytometry. Shown are a representative pair of histograms for WT and *Dhrs7b$^{Δ/Δ}$* B cells as

indicated (left panel) and a dot graph aggregating all experiments' outcomes (right panel).

(H) Frequencies of apoptotic B cells in cultures as in (A-D) were scored for annexin V and

7AAD as described (84). Shown are representative data of flow plots in the lymphocyte gate

(left panel) and a dot graph aggregating all experiments' outcomes (right panel). (I, J) PexRAP

and ROS control promote B cell population growth in vitro. WT and *Dhrs7b$^{\Delta/\Delta}$* B cells were
cultured 5 days in anti-CD40, BAFF, IL-4, IL-5 and 4-hydroxytamoxifen in the presence or
absence of ROS scavenger NAC (1 mM vs 5 mM) (H). (I) B cells activated as in (A-G) were
cultured 5 d in the presence or absence of NAC (1 mM) or thioridazine (100 nM). The bar graphs
show the mean (± SEM) recovered cell number from three independent experiments, each with
two independent samples of each genotype (WT; *Dhrs7b$^{\Delta/\Delta}$*). Complementary results of
experiments including anti-IgM for BCR cross-linking are shown in Supplemental Figs 2 and 6.
1128
Figure 9. PexRAP deficiency in activated B cells reduces mitochondrial metabolism and ER
mass. (A-G) Purified B cells were activated with and cultured (2 d) in anti-CD40, BAFF, IL-4,
IL-5, and 4-OHT, then analyzed using a Seahorse XFe96 after harvest and division in equal
portions. (A) Shown are aggregated results from three independent experiments measuring
oxygen consumption rate (OCR) quantified via metabolic flux analysis during mitochondrial
stress testing. (B) Basal OCR, (C) maximal OCR, (D) ATP-production coupled respiration, (E)
proton leak, and (F) spare respiratory capacity of WT and *Dhrs7b$^{\Delta/\Delta}$* B cells assayed in (A). Data
points of individual samples are show (each individual dot), as well as bars to display mean
(±SEM) values. T-tests with Welch's correction were used to calculate p-values. *, $p<0.05$; ***,
$p<0.001$. (G) Extracellular acidification rate (ECAR) of B cells cultured as described previously
(36) and assessed using the glycolytic stress test. (H, I) PexRAP influences ER mass. WT and
Dhrs7b$^{\Delta/\Delta}$ B cells were cultured 5 days in anti-CD40, BAFF, IL-4, IL-5 and 4-hydroxytamoxifen
followed by the staining with ER-Tracker Green. Representative histogram of ER-Tracker Green
in viable cells (H) and aggregated MFI (±SEM) of ER-Tracker Green from three replicate
experiments (I). Mann-Whitney U test was used to calculate p values.

 SUPPLEMENTAL INFORMATION

 Supplemental Figures

 Supplemental Figure 1. PexRAP is dispensable for steady-state T cell numbers, but impacts

 Tfh cell population. Results from flow cytometry analyses of splenocytes from mice (*Rosa26-*

 CreER^{T2}, *Dhrs7b$^{+/+}$,* "WT" or *Rosa26*-CreER^{T2}; *Dhrs7b$^{f/f}$,* "*Dhrs7b* Δ/Δ") harvested after

 sequential tamoxifen treatments followed by immunization with SRBC and harvest 1 wk

 thereafter, all as described in *Materials and Methods* and illustrated in Fig. 1A. (A-C) Normal

 steady-state numbers of T cells. Shown are representative flow plots of splenic T cells (A), mean

 (± SEM) frequencies of TCRβ^{+} CD4^{+} T cells (B), and CD8 (TCRβ^{+} CD4$^{-)}$ T cells (C) from WT

 and *Dhrs7b* Δ/Δ (cKO) mice based on three replicate experiments totaling 8 WT and 7 cKO

 mice. (D) Linking to Fig. 1C-F, representative gating scheme illustrating the flow-cytometric

 determination of splenic B cell frequencies and humbers (overall and GC B phenotype) in mice

 immunized as in A-C, Fig. 1A. (E, F) Linking to Fig. 1G-I, frequencies of GC, plotted as

 GC/mm^{2} (E) and sizes of their LZ (plotted as area of each GC in the microscopic sections) (F)

 identified by immunofluorescent staining and microscopy of spleen sections from immunized

 mice. (G) Representative gating scheme illustrating the flow cytometric determination of Tfh and

 GC-Tfh cells in splenocyte suspensions of mice prepared and immunized as in Fig. 1. (H, I)

 PexRAP promotes Tfh cell population size. (H) Shown are representative flow plots of Tfh and

 GC-Tfh cells, determined using the gating scheme of Supplemental Fig. 1G. (I) Shown are mean

 (±SEM) frequencies of PD-1^{med} CXCR5^{med} Tfh cells (left) and PD-1^{hi} CXCR5^{hi} GC-Tfh cells

 (right) from WT and cKO mice in two independent replicate experiments (n = 6 WT and 6 cKO).

 Supplemental Figure 2. PexRAP is dispensable for pre-immune B cell subset balance and

 surface IgM expression, but contribute to B cell population growth.

 (A) A representative gating scheme illustrating the flow-cytometric estimation of the MZ and FO

 B cell populations in mice. (B, C) Shown are the mean (±SEM) frequencies of B220^{+} CD43^{neg}

 CD23^{neg} CD21^{+} MZB cells (left panel) and B220^{+} CD43^{neg} CD23^{+} CD21^{lo} FOB cells (right

 panel) in spleen (B), and aggregated mean (±SEM) geometric MFI of surface IgM on MZB and

 FOB subsets (C). (D) Gating strategy for identification of B1-b cells and measurement of their

 surface IgM levels. (E, F) Shown are the mean (±SEM) frequencies of B220^{+} CD23^{lo} CD43^{+}

 CD5^{neg} B1-b cells in peritoneal cavity (E), and aggregated mean (±SEM) geometric MFI of

 surface IgM on B1-b cells (F). (G) In vivo generation of IgD^{neg} progeny from transferred B cells

 of the indicated genotypes, as measured in four independent replicate experiments (n =8 WT and

 9 cKO). Shown are mean (±SEM) frequencies of IgD^{neg} B cells as aggregate data for the

 frequencies of IgD^{neg} events in the gate for viable B cells recovered from μMT recipient mice

after transfer of WT or PexRAP-depleted B cells (experiments of Fig. 2D-F), as indicated, with

representative flow plots in Fig. 2E. (H) PexRAP promotes B cell proliferation in vitro. B cells

were purified and stained with CTV as in Fig 2G. WT and $Dhrs7b^{\Delta/\Delta}$ B cells were stained with

Cell Trace Violet (CTV), activated and cultured 4 days in anti-IgM, anti-CD40, BAFF, IL-4, IL-

5, and 4-OHT. Shown are the representative flow-cytometric analysis of CTV partitioning (left),

and aggregated frequencies of divided B cells from two independent replicate experiment, each

using two separately sourced pools of B cells (n = 4 WT and 4 cKO). P values were calculated

by Mann-Whitney U test.

1188

Supplemental Figure 3. Activation of B cells induces PexRAP-dependent increases in their

ether and plasmalogen P-lipids. As in Fig. 3, B lymphocytes of the indicated $Dhrs7b$ genotypes

(WT and Δ/Δ) were prepared from individual tamoxifen-injected mice and divided to analyze

without activation ("naive") or after activation and culture (48 h) ("activated"). Shown are heat

maps with more extensive lists of ether phospholipids and plasmalogens identified in the LC-

MS-MS analyses for both genotypes in positive (A) and negative (B) ion modes. (C, D)

Controlling for transgene and tamoxifen effects. Shown are LC-MS-MS results for lipids in B

cells, activated or not, as indicated, of conventional non-transgenic B6 mice not injected with

tamoxifen and those of tamoxifen-injected CreER^{T2+} mice (the "+/+" samples of panels A, B and

of Figure 3). Bar graphs in (C) and (D) show results of these control comparisons for some of the

phospholipids analyzed in prior figures panels (Fig. 3A, B).

1200

Supplemental Figure 4. Immunization induces increases in P-lipids of both primary and

secondary follicles dependent on PexRAP in B cells. (A, B) Selected representative m/z

features Spleens of AID-GFP were harvested at 7 days after immunization with SRBC and

analyzed for exact mass signatures of ether lipid species in (A) negative and (B) positive ion

modes. (C) Spleens of immunized mice (tamoxifen-treated $huCD20$-CreER^{T2}; $Dhrs7b^{f/f}$, i.e.,

$Dhrs7b^{\Delta/\Delta\text{-B}}$, and $huCD20$-CreER^{T2}, i.e., "WT", controls) were analyzed by IMS. Shown are the

positive ions presented in (B) for GC vs follicles of immunized AID-GFP mice. The bar graph

shows the mean (± SEM) ion intensities in lymphoid follicles and GC from WT and $Dhrs7b^{\Delta/\Delta-\text{B}}$

spleens. Median intensity of each ion was obtained from three follicles per spleen and three GC

per spleen from WT and $Dhrs7b^{\Delta/\Delta-B}$ mice (three biological replication experiments totaling

seven WT and six cKO subjects). P values were calculated by Mann-Whitney U test.

1212

1213   **Supplemental Figure 5. A subset of ether phospholipids in primary follicles and GC**

1214   **(secondary follicles) depend on PexRAP in B cells.** (A) As in Fig. 4, $Dhrs7b$ was inactivated

1215   by tamoxifen injections into mice bearing the huCD20-CreER^{T2} transgene, followed by

immunization with SRBC and analysis. Shown are representative higher-magnification images defining the primary (IgD^+) and secondary ($IgD^{lo/neg}$, $GL7^+$) follicles and the identification of light ($CD35^+$) and dark ($CD35^{neg}$) zones. (B-C) After immunization and harvest of the mice as shown in Fig. 4, the sizes of GC, light zones (LZ), and dark zones (DZ) in photomicrographs of sectioned and immunofluorescently stained spleens of the immunized mice (using three GC per spleen, with seven WT and five $Dhrs7b^{\Delta/\Delta-B}$.) were quantified. (B) Spleens of immunized mice (tamoxifen-treated $huCD20$-CreER^{T2}; $Dhrs7b^{f/f}$ and $huCD20$-CreER^{T2} controls, as indicated) were analyzed by immunofluorescence staining as in (A). Each dot represents one follicle in a given mouse. (C) Dot graph showing the ratio of LZ/DZ using the data of panel B. (D, E) Mice ($huCD20$-CreER^{T2}; $Dhrs7b^{+/+}$, or $huCD20$-CreER^{T2}; $Dhrs7b^{f/f}$) were treated with tamoxifen and immunized with SRBC as in Fig 4A, and were analyzed 7 d after immunization. (D) Shown are aggregated frequencies of $CD138^+$ $GL7^+$ early plasmablasts in $B220^+$ IgD^{neg} dump^{neg} gate (dump channel = IgD, CD11b, CD11c, F4/80, Gr1, TCRβ, and 7AAD). (E) The aggregated frequencies of GC-Tfh (PD-1^{hi}, CXCR5^{hi}) and Tfh (PD-1^{int}, CXCR5^{int}) among $CD4^+$ T cells with the gating strategy as shown in Supplemental Fig. 1G. (x=2; 6 WT vs 6 cKO). P value was calculated by Mann-Whitney U test. (F, G) Mean (±SEM) absorbances of ELISA performed across the indicated dilutions of sera from the tamoxifen-treated mice ($huCD20$-CreER^{T2}; $Dhrs7b^{+/+}$ or $huCD20$-CreER^{T2}; $Dhrs7b^{f/f}$, i.e., $Dhrs7b^{\Delta/\Delta-B}$) of Fig. 6. Shown are the results for (F) all- (NP_{20} captured) and (G) high- (NP_2 captured) -affinity anti-NP Ab of the IgG1 and IgG2c isotypes, as indicated.

Supplemental Figure 6. PexRAP contributes to GC B cell proliferation, ROS homeostasis and B cell population growth. (A, B) Proliferation and development of $CD138^+$ progeny derived from B cells activated and cultured with anti-CD40, BAFF, IL-4, and IL-5. Shown are representative flow plots (A), and aggregated mean (±SEM) frequencies of $CD138^+$ cells in viable lymphocyte from three independent experiments with 6 WT and 7 $Dhrs7b^{\Delta/\Delta-B}$ mice at t = 5dq. (C) PexRAP regulates proliferation of GC B cells. Tamoxifen-treated mice ($huCD20$-CreER^{T2}; $Dhrs7b^{+/+}$ or $huCD20$-CreER^{T2}; $Dhrs7b^{ff}$, i.e., $Dhrs7b^{\Delta/\Delta-B}$) were immunized with SRBC, and the mice were injected with BrdU as described in Methods. Shown are a dot graph aggregating all experiments' outcomes for the frequencies of $BrdU^+$ cells in GCB cells (x=2; n = 6 WT and 6 cKO). (D) WT and $Dhrs7b^{\Delta/\Delta}$ B cells were activated and cultured 3 days in anti-IgM, anti-CD40, BAFF, IL-4, IL-5, and 4-OHT. Shown are representative histogram image of H_2DCFDA in the B cell gate (left panel), and aggregated mean (± SEM) geometric MFI of H_2DCFDA (right panel) from two independent replicate experiment (n = 4 WT and 4 cKO). P values were calculated by Mann-Whitney U test. (E) PexRAP restrains lipid peroxidation. B

cells were activated and cultured as in Fig 8E. Shown are the aggregated means (± SEM) of
geometric MFI of C11-Bodipy from two independent replicate experiments (n = 5 WT and 5
cKO). (F-H) NAC mitigated the impairment of population increase for PexRAP-deficient B
cells, but failed to enhance cell division. B cells of the indicated genotypes were activated with
combined BCR and CD40 cross-linking as in (D), cultured 5 d in the presence or absence of
NAC (5 mM), counted and analyzed by flow cytometry. (F) The dot graph shows the mean (±
SEM) recovered cell number from two independent experiments. P values were calculated by
Mann-Whitney U test. (G, H) Prior to activation, B cells of the each genotype were stained with
CellTrace Violet (CTV). (G) Shown are flow cytometry ouputs of CTV fluorescence in the
viable B cell gate for the indicated samples from one of the independent replicate pools. A
measurement bar denotes the cut-off between 0-2 divisions and ≥3 divisions, with the inset
numbers representing the percentage of cells that divided at least three times. (H) A dot graph
displaying the individual as well as mean percentages of B cells that divided at least three times
for samples of the indicated genotypes and treatment conditions. P values were calculated by
Mann-Whitney U test.
1267
Supplemental Figure 7. Distinct outcomes of heightened ROS elicited by H_2O_2 versus
menadione. (A-B) Induction of ROS independent from B cell activation. Shown are
representative flow plots of DCFDA in the viable B cell gate after exposure (1 h) (A) to H_2O_2
(200 μM) or (B) menadione (8 μM), as in (87). (C-D) Representative flow plots of annexin V vs
7-AAD for identification of early-apoptotic B cells. Cultured B cells were analyzed 3 hr after
treatment with (C) 200 μM H_2O_2 or (D) 8 μM menadione. Inset numbers represent the fraction
of B cells that are Annexin V[+] 7-AAD[neg] or Annexin V[neg] 7-AAD[+]. (E-G) Shown are the results
of three independent experiments to measure (E) total cell viability, (F) BODIPY C11 MFI and
(G) frequencies of Annexin V[+] 7-AAD[neg] frequency in B cells treated (8 μM) menadione
overnight (E) or for 3 h (D). P-values were calculated using the unpaired Student's t-test.
1278
1279
Supplemental Table 1. Immunization-induced increases of selected ether lipids in GC.
Shown are (a) a sample of *m/z* features characteristic of the indicated phospholipid species
identified by the LIPIDMAPS database, and (b) the mean (±SEM) ion intensity / counts in IMS
data generated from spleens of AID-GFP mice, immunized or not (UI), after mapping to the
indicated regions of interest (primary follicles or GC, i.e. secondary follicles) using fluorescent
images. Shown are ions more accumulated in GC regions compared to primary follicles (8 ions
from negative ion mode, and 5 ions from positive ion mode, all p<0.05 for comparison of 1^0
follicle to UI (c), or GC to 1^0 follicle (d). P values were calculated by Mann-Whitney U test.

1288

Supplemental Table 2. . Impact of PexRAP on relative quantities of selected lipid species in

primary and secondary follicles (GC). As in Table 1 except that B cells of immunized mice

were either WT or lacked PexRAP (cKO, i.e., $Dhrs7b^{\Delta/\Delta-B}$). Shown are (a) a sample

of m/z features characteristic of the indicated phospholipid species. (b) the mean ±SEM ion

intensity counts in IMS data after mapping to the indicated regions of interest as in Table 1. (c-f)

indicate that p<0.05 for the null hypothesis in considering a difference between WT mouse

spleens after immunization versus UI controls (c), 1^0 B cell follicles in spleens of immunized

WT vs $Dhrs7b^{\Delta/\Delta-B}$ mice (d), GC vs in 1^0 B cell follicles in spleens of immunized WT mice (e),

and GC of immunized WT vs $Dhrs7b^{\Delta/\Delta-B}$ mice (f) in the ions counts for designated m/z features.

P values were calculated by Mann-Whitney U test.

1299

A

R26-CreER^{T2}

Dhrs7b^{f/f};
R26-CreER^{T2}

Tmx → SRBC   Analysis
         ↓   ↑
Wk: 0   1

B   WT   *Dhrs7b^{Δ/Δ}*
         B T   B T

◀ PexRAP

◀ Actin

C   WT   *Dhrs7b^{Δ/Δ}*
CD19 / B220   37   24

D
p<0.001
% B cells among live cells
WT   Dhrs7b^{Δ/Δ}

p=0.006
# of B cells (× 10^6)
WT   Dhrs7b^{Δ/Δ}

E   WT   *Dhrs7b^{Δ/Δ}*
GL7 / CD95   9.6   3.6

F
% GCB cells
WT   Dhrs7b^{Δ/Δ}

p=0.01
# of GCB cells (× 10^6)
WT   Dhrs7b^{Δ/Δ}

G
WT
KO
GL7 IgD CD35

H   WT
         Dhrs7b^{Δ/Δ}
500 µm
IgD^+
DZ (GL7^+ CD35^-)
LZ (GL7^+ CD35^+)

I
p=0.02
GC number per spleen section
WT   Dhrs7b^{Δ/Δ}

p=0.047
GC area (mm^2)
WT   Dhrs7b^{Δ/Δ}

Fig. 1

A

huCD20-CreER^{T2}
or
Dhrs7b$^{f/f}$;
huCD20-CreER^{T2}

TAM   *Analysis*

Day: 1 - 5   10

B

WT   *Dhrs7b$^{Δ/Δ–B}$*

◄ PexRAP

◄ Actin

C

p=0.002

% B cells among live cells

WT   Δ/Δ-B

p=0.007

# of B cells (× 10^6)

WT   Δ/Δ-B

D

p<0.001

% B cells

WT   Δ/Δ-B

E

WT   *Dhrs7b$^{Δ/Δ-B}$*

25.2   13.4

IgD

CTV

12.6   5.4

IgD

CD19

F

p=0.02

% CTV of ≥3 divisions

WT   Δ/Δ-B

G

WT   *Dhrs7b$^{Δ/Δ-B}$*

65.5   45.5

CTV

H

p<0.01

% CTV of ≥3 divisions

WT   Δ/Δ-B

I

p=0.002

# of B cells (× 10^6)

WT   Δ/Δ-B

Fig. 2

Fig. 3

Fig. 4

Fig. 5

Fig. 6

Fig. 7

Fig. 8

Fig. 9

Supplemental Fig. 1

Supplemental Fig. 2

Supplemental Fig. 3

Supplemental Fig. 4

A

WT

Dhrs7b$^{\Delta/\Delta-B}$

500 μm

■ IgD$^+$
■ DZ (GL7$^+$ CD35$^-$)
■ LZ (GL7$^+$ CD35$^+$)

B

GC area (mm^2)   *p=0.01*   WT   Δ/Δ-B

LZ area (mm^2)   *p=0.02*   WT   Δ/Δ-B

DZ area (mm^2)   0.0916   WT   Δ/Δ-B

C

LZ/DZ ratio   *p=0.02*   WT   Δ/Δ-B

D

Early PB

% CD138$^+$ GL7$^+$ among dump$^-$ B cells   WT   Δ/Δ-B

E

GC-Tfh   p<0.01   WT   Δ/Δ-B

Tfh   p<0.01   WT   Δ/Δ-B

% among CD4$^+$ cells

F

IgG1_NP$_{20}$ IgG2c_NP$_{20}$

■ WT   □ *Dhrs7b*$^{\Delta/\Delta-B}$

OD$_{450nm}$

1:10000 1:20000 1:40000 1:80000 1:160000 1:320000

1:2000 1:4000 1:8000 1:16000 1:32000 1:64000

G

IgG1_NP$_2$ IgG2c_NP$_2$

■ WT   □ *Dhrs7b*$^{\Delta/\Delta-B}$

OD$_{450nm}$

1:10000 1:20000 1:40000 1:80000 1:160000 1:320000

1:2000 1:4000 1:8000 1:16000 1:32000 1:64000

Supplemental Fig. 5

Supplemental Fig. 6

Supplemental Fig. 7

Table 1.
   Immunization-induced increases of selected lipids in GC

Negative Ions			
m/z [a] Lipid ID	UI	1° B cell Follicle	GC
716.5216 PE(18:0_16:1)	618 ± 22 [b]	2400 ± 83 [c]	3674 ± 154 [d]
752.5545 PE(O-18:0_20:4)	301 ± 14	1739 ± 33 [c]	1960 ± 52 [d]
760.5056 PE(16:0_22:6)	203 ± 4	1816 ± 30 [c]	2031 ± 23 [d]
772.5242 PE(P-18:1_22:6)	234 ± 5	2334 ± 47 [c]	2578 ± 63 [d]
776.5556 PE(O-18:0_22:6)	277 ± 9	1638 ± 51 [c]	2074 ± 90 [d]
869.5526 PI(O-18:0_20:5)	242 ± 7	1611 ± 24 [c]	1789 ± 15 [d]
872.576 PC(18:0_24:0)	171 ± 6	1167 ± 16 [c]	1369 ± 18 [d]
888.5612 PS(22:1_22:6)	506 ± 13	2684 ± 54 [c]	3386 ± 72 [d]

Positive Ions			
m/z [a] Lipid ID	UI	1° B cell Follicle	GC
739.4587 PG(12:0_20:3)	3042 ± 102	3733 ± 116 [c]	4778 ± 274 [d]
740.465 PC(16:0_15:1)	1310 ± 13	1478 ± 66 [c]	1933 ± 142 [d]
784.5564 PC(10:0_24:0)	2840 ± 24	7956 ± 147 [c]	9856 ± 270 [d]
798.5347 PC(10:0_25:0)	9937 ± 269	27589 ± 574 [c]	34733 ± 1644 [d]
799.5375 PG(18:0_18:1)	5665 ± 65	12078 ± 282 [c]	15311 ± 790 [d]

Table 2. Impact of PexRAP on relative quantities of selected lipid species in primary and secondary follicles (GC).

Negative Ions					
m/z [a] Lipid ID	UI	WT follicle	PexRap cKO follicle	WT GC	PexRap cKO GC
436.2801 LPE (O-16:1)	315 ± 4	369 ± 7	374 ± 11	610 ± 20 [e]	475 ± 19
716.5216 PE (18:0_16:1)	754 ± 43 [b]	1608 ± 182 [c]	1263 ± 112	2729 ± 247 [e]	1770 ± 137 [f]
746.5098 PE (O-16:1_22:6)	622 ± 19	672 ± 35	609 ± 23	1066 ± 39 [e]	836 ± 39
752.5545 PE (O-18:0_20:4)	459 ± 49	657 ± 51 [c]	548 ± 46 [d]	1191 ± 61 [e]	734 ± 25 [f]
760.5056 PE (16:0_22:6)	229 ± 8	386 ± 25 [c]	312 ± 16 [d]	569 ± 27 [e]	408 ± 15 [f]
772.5242 PE (P-18:1_22:6)	388 ± 47	617 ± 46 [c]	499 ± 33 [d]	918 ± 21 [e]	646 ± 9 [f]
776.5556 PE (O-18:0_22:6)	449 ± 53	661 ± 50 [c]	562 ± 46	1130 ± 49 [e]	789 ± 13 [f]
869.5526 PI (O-18:0_20:5)	509 ± 81	641 ± 50 [c]	494 ± 40 [d]	1014 ± 21 [e]	690 ± 12 [f]
872.576 PC (18:0_24:0)	230 ± 19	285 ± 12 [c]	247 ± 9 [d]	522 ± 17 [e]	336 ± 9 [f]
888.5612 PS (22:1_22:6)	1423 ± 277	2051 ± 199 [c]	1904 ± 199	3260 ± 156 [e]	2629 ± 80 [f]

Positive Ions					
m/z [a] Lipid ID	UI	WT follicle	PexRap cKO follicle	WT GC	PexRap cKO GC
730.5767 PE (P-18:0_18:1)	91 ± 3	97 ± 3	85 ± 3	130 ± 7 [e]	102 ± 4
739.4587 PG (12:0_20:3)	3042 ± 102	3094 ± 40	2944 ± 31 [d]	3506 ± 140 [e]	3011 ± 46 [f]
784.5564 PC (10:0_24:0)	2840 ± 24	2917 ± 74	2617 ± 46 [d]	3522 ± 102 [e]	2561 ± 42 [f]
798.5347 PC (10:0_25:0)	9937 ± 269	10928 ± 214 [c]	11372 ± 155 [d]	15378 ± 116 [e]	14694 ± 186 [f]
801.5509 PG (18:0_18:0)	1540 ± 37	1591 ± 46	1633 ± 26	1444 ± 27 [e]	1533 ± 22 [f]
847.5361 PI (15:0_18:0)	1277 ± 24	1339 ± 54	1217 ± 35	1150 ± 46 [e]	1044 ±44
874.5623 PC (20:5_22:6)	754 ± 39	688 ± 48	675 ± 25	469 ± 27 [e]	556 ± 37