Multi-omics investigation of spontaneous T2DM macaque emphasizes gut microbiota could up-regulate the absorption of excess palmitic acid in the T2DM progression

Xu Liu[1#], Yuchen Xie[1#], Shengzhi Yang[1], Cong Jiang[1], Ke Shang[1], Jinxia Luo[1,3], Lin Zhang[1], Gang Hu[2], Qinghua Liu[2], Bisong Yue[1], Zhenxin Fan[1], Zhanlong He[4], Jing Li[1*]

[1] Key Laboratory of Bio-resources and Eco-environment (Ministry of Education), College of Life Sciences, Sichuan University, Chengdu 610065, China.

[2] SCU-SGHB Joint Laboratory on Non-human Primates Research, Meishan, Sichuan, China.

[3] Sichuan Key Laboratory of Development and Application of Monkey Models for Human Major Disease, Sichuan, China.

[4] Institute of Medical Biology, Chinese Academy of Medical Sciences and Peking Union Medical College, Yunnan, China.

[#] These authors contributed equally to this work.

* Correspondence:

Jing Li (ljtjf@126.com)

College of Life Sciences, Sichuan University

Abstract

Although gut microbiota and lipid metabolites have been suggested to be closely associated with type 2 diabetes mellitus (T2DM), the interactions between gut microbiota, lipid metabolites and the host in T2DM development remains unclear. Rhesus macaques may be the best animal model to investigate these relationships given their spontaneous development of T2DM. We identified eight spontaneous T2DM macaques and conducted a comprehensive study investigating the relationships using multi-omics sequencing technology. Our results from 16S rRNA, metagenome, metabolome and transcriptome analyses identified that gut microbiota imbalance, tryptophan metabolism and fatty acid β oxidation disorders, long-chain fatty acid (LCFA) accumulation, and inflammation occurred in T2DM macaques. We verified the accumulation of palmitic acid (PA) and activation of inflammation in T2DM macaques. Importantly, mice transplanted with spontaneous T2DM macaque fecal microbiota and fed a high PA diet developed prediabetes within 120 days. We determined that gut microbiota mediated the absorption of excess PA in the ileum, resulting in the accumulation of PA in the serum consequently leading to T2DM in mice. In particular, we demonstrated that the specific microbiota composition was probably involved in the process. This study provides new insight into interactions between microbiota and metabolites and confirms causative effect of gut microbiota on T2DM development.

Key words: spontaneous T2DM macaques, multi-omics, gut microbiota, palmitic acid, fecal microbiota transplantation

Introduction

Diabetes mellitus is considered to be a refractory disease causing a significant socio-economic burden. Type 2 diabetes mellitus (T2DM) is the dominant type of diabetes mellitus characterized by metabolic disorders, insulin resistance, and deficiency of insulin secretion (1, 2). The pathogenesis of this chronic disease is complex and genetic and environmental factors, such as sugar and lipid intake, gut microbiota, many metabolites, and even air pollutants, contribute to its increase in prevalence (3, 4). Accumulating evidence has linked gut microbiota with T2DM development by variety ways. The gut microbiota can impact the integrity of the intestinal epithelial barrier, mediate insulin resistance, as well as regulate the function of mitochondria (5, 6, 7). They can regulate local or systemic immunity and inflammation, which also contributes to the development of T2DM (8). Moreover, various gut microbial metabolites, such as short-chain fatty acids, bile acid, and tryptophan-derived metabolites, have been reported to be closely related to the pathogenesis of T2DM (5, 6, 9, 10). However, the interactions between gut microbiota and its host with T2DM have not yet been fully characterized. In pathological conditions, the dysregulation of the host can lead to changes in gut microbiota composition. In turn, the microbiota plays a regulatory role to participate in the development of T2DM. Despite the complicate interactions between host and microbiota in the context of T2DM, some studies suggest antidiabetic interventions targeting the gut microbiota such as fecal microbiota transplanting (FMT) can be applied as a clinical treatment of T2DM (11, 12, 13).

Meanwhile, dysfunction of lipid metabolism contributes to T2DM development by inducing lipotoxicity in humans and animal models (14, 15). The long-chain fatty acids (LCFAs) are the principal lipid components naturally occurring in animal fats and vegetable oil, as well as the main metabolites of fat. LCFAs such as palmitic acid (PA, C16:0), palmitoleic acid (C16:1N7), and oleic acid (C18:1N9), are reported to have strong association with T2DM. In the last few decades, there is increasing

evidence that frequent consumption of LCFAs contributes to metabolic diseases such

as obesity and T2DM due to the high PA content (16, 17). PA is a saturated fatty acid

and its increase in serum is a significant contributing factor in T2DM development

(18, 19, 20). The suggested mechanisms by which PA mediates T2DM include

increasing diacylglycerol and ceramide synthesis (21), mitochondrial and

endoplasmic reticulum stress (22, 23), and activation of pro-inflammatory pathways

(24). Nevertheless, whether gut microbiota involving in PA mediated T2DM and the

interactions between gut microbiota and LCFA metabolites in T2DM development are

still unclear.

Spontaneous development of T2DM in non-human primates (e.g. macaques) is

highly similar to human T2DM, such as insulin resistance in early stages and later

abnormal glucose tolerance and T2DM development follows the same pathological

changes of pancreatic islets and complications. In fact, T2DM macaques avoid

medication interference and environmental heterogeneity under controlled

experimental conditions, and share key pathological features with humans, such as

amyloidosis of pancreatic islets, which is absent in mouse models (25, 26), suggesting

that T2DM macaques are the optimal animal model for simulating human T2DM and

its complications (27). However, previous studies indicated that naturally occurring

spontaneous T2DM macaques in captive populations were rare even if individuals

were given a high lipid and high sugar diet (28, 29). After nasally fed cynomolgus

macaques with a high-fat dietary emulsion for 12 months, the macaques did not

experience significant increases in fasting blood glucose and glycosylated hemoglobin

(29). The controversial effects of high fat on the T2DM development are worthy of

further investigations to better understand this complex disease.

Here, this study identified spontaneous development of T2DM in individuals

(hereafter spontaneous T2DM macaques) from a large group of rhesus macaques.

These spontaneous T2DM macaques have never been treated with anti-diabetic drugs

and therefore provide valuable models for pathogenesis investigation of T2DM.

Based on the macaque model, we used multi-omics techniques to address the

interactions between gut microbiota, host gene expression, and fecal metabolites and the development of T2DM. Our results demonstrated that gut microbiota and LCFA metabolites played important roles in the pathogenesis of T2DM. We validated the increased content of plasma PA and activation of inflammation in the T2DM macaques. In addition, we successfully induced prediabetes in mice by transplanting fecal microbiota from T2DM macaques into mice in conjunction with a diet high in PA. We also revealed the specific structure of gut microbiota that promoted T2DM development by regulating the absorption of excess PA in mice, providing experimental evidence for the functional role of gut microbiota in T2DM pathogenesis.

Results

Metagenome and 16S sequencing demonstrate alterations on gut microbiota in spontaneous T2DM macaques

We identified eight spontaneous T2DM macaques out of 1698 individuals from long-term glucose monitoring in a captive population (Table S1). The fasting plasma glucose (FPG), fasting plasma insulin (FPI), and HOMA-IR levels in the T2DM macaques were significantly higher than in the control group ($p<0.01$), suggesting insulin resistance in T2DM macaques. However, glycosylated hemoglobin A1c (HbA1c), triglycerides (TG), total cholesterol (TC), high-density lipoprotein cholesterol (HDL), low-density lipoprotein cholesterol (LDL), and BMI did not significantly differ from the controls ($p>0.05$, Table 1). Comparison of gut microbiota between the two groups based on 16S rRNA amplicon found that both Shannon and Simpson index decreased in T2DM macaques but it was not significantly different from controls ($p>0.05$; Figures 1A and B). PCoA indicated that the microbiota composition of spontaneous T2DM macaques was different from the control group (Figure 1C). As one of the most dominant families in both groups, Lachnospiraceae showed significantly higher abundance in the T2DM group (14.48%) than in the control group (6.66%). While the abundance of the Lactobacillaceae family was

significantly greater in the control group (25.95%) compared to spontaneous T2DM

macaques (20.88%) (Figure 1D). A total of 21 microbes were identified as differential

microbes between the T2DM group and control group, where ten microbes including

Erysipelotrichaceae and five members in Lachnospiraceae family (*Ruminococcus*

gnavus (current name: *Mediterraneibacter gnavus*), *Lachnospira*, *Coprococcus* sp.

Dorea longicatena, and *Roseburia*) were significantly greater in the T2DM group

(Figure 1E).

The metagenome results showed that the T2DM group had a higher abundance

of Erysipelorictchaeae, *Eubacterium rectale*, Lachnospiraceae, Negativicutes, *Blautia*,

and Coriobacteriia than the control group (Figure S1A). Functional enrichment

demonstrated a total of 74 KEGG pathways with significant differences between the

two groups. These pathways were mainly associated with T2DM and inflammation,

including type II diabetes mellitus, glucagon signaling pathway, starch and sucrose

metabolism, lipid metabolism, and toll-like receptor signaling pathways (Figure 1F).

Among the six CAZy enzyme families, the Carbohydrate-binding module (CBMs),

Glycosyl Transferases (GTs), Carbohydrate esterases (CEs), and Polysaccharide

Lyases (PLs) families were significantly upregulated in the T2DM group, indicating

significant changes in carbohydrate metabolism (Figure 1G). The results of

metagenome and 16S sequencing demonstrated significant alterations in the

composition and function of gut microbiota in spontaneous T2DM macaques, with a

greater abundance of microbes associated with T2DM and fewer beneficial microbes.

Fecal metabolome and blood transcriptome reveals dysfunction of

fatty acid β oxidation and tryptophan metabolism in T2DM

macaques

The UHPLC-MS based metabolome analysis on fecal samples of T2DM and control

macaques identified 1564 metabolites belonging to various types of secondary

metabolites, with lipids and lipid-like molecules being most abundant (31.1%) (Figure

S1B). We found 64 significantly differential metabolites between the two groups

using a combined multidimensional statistical analysis (OPLS-DA) and univariate

statistical analysis (T-test) (VIP>1, $p<0.05$) (Figures 2A and B; Table S2). Among

them, muscon, indole-3-acetaldehyde, and serotonin were significantly lower in the

T2DM group (Figure 2C) and are associated with anti-inflammatory activity (34, 35,

36). Notably, 64 differential metabolites were significantly enriched in one pathway:

tryptophan metabolism (Figure 2D), and we identified two significantly different

metabolites, indole-3-acetaldehyde and serotonin in this pathway. Both metabolites

are microbiota-derived tryptophan metabolites and the ligands for AhR (35, 36). The

lower concentration of AhR ligands may lead to the development of inflammation and

metabolic syndromes in humans (37). In addition, we found the contents of many

acylcarnitine metabolites were significantly higher in the T2DM group (VIP>1,

$p<0.1$), including l-propionylcarnitine, hexanoyl-l-carnitine, (r)-butyrylcarnitine, and

isovaleryl-l-carnitine (Table S2). Among them, l-propionylcarnitine, a kind of

acylcarnitines, was the most upregulated metabolite in the T2DM macaques (FC:

16.19) (Figure 2C). Acylcarnitines are products of incomplete oxidation of LCFAs,

which can activate pro-inflammatory signaling pathways and ultimately inhibit insulin

activity (38). Our results indicated incomplete LCFAs β oxidation in spontaneous

T2DM macaques, while similar characteristics were also observed in insulin resistant

and T2DM humans (39, 40).

Blood transcriptome analysis was consistent with metabolome results, indicating

dysfunction of fatty acid β oxidation and inflammation in T2DM macaques. Gene

expression in T2DM macaques exhibited significant differences from the controls

(Figure 2E), and a total of 161 differentially expressed genes (DEGs) (26 upregulated

and 135 downregulated) were identified in T2DM macaques at a FDR level of 0.05

(Figure 2F). Enrichment analysis of the DEGs was linked to diabetes, fatty acid

metabolism, and inflammation, such as diabetic cardiomyopathy, glucose homeostasis,

fatty acid metabolic process, and chemokine production (Figure 2G). We also

identified 26 differential enrichment pathways between the two groups by aggregate

fold change (AFC), and most were associated with insulin resistance and

inflammation, including insulin resistance, PI3K-Akt signaling pathway, bacterial invasion of epithelial cells, and NOD-like receptor signaling pathway (Figure S2A). As shown in the insulin resistance pathway, expression of *IL6* and *IRS1* were upregulated, while *INSR* was downregulated in T2DM macaques (Figure S2B). The WGCNA analysis identified four modules that were related to T2DM. Darkgreen module and brown module were significantly positively correlated with T2DM, while green module and darkred module were negatively correlated with T2DM (Figure 2H). Genes in the darkgreen module and brown module related to lipolysis and inflammation were significantly upregulated in T2DM macaques, and genes in the green module and darkred module related to fatty acid metabolism and insulin secretion were significantly downregulated (Figure S2C). With the cut-off (|kME>0.8|), a total of 546 genes in the four modules were identified as hub genes (Table S3). Of these 546, 102 hub genes were also DEGs (Figure 2I). Several of these genes have been reported as correlated with T2DM in humans, including *IGF2BP2*, *LEPR*, *RAP1A*, *SESTRIN 3*, and *ITLN1* (41, 42, 43, 44, 45). In addition, *ACSM3*, *HADHB*, and *EFHB* are involved in fatty acid β oxidation and inflammation (46, 47, 48).

Validation of LCFAs accumulation and inflammation in T2DM macaques

Collectively, metabolome and transcriptome results indicated dysfunction in fatty acid β oxidation and tryptophan metabolism in T2DM macaques, which may lead to LCFA accumulation and inflammation. To support this conclusion, we performed targeted medium- and long-chain fatty acid mass spectrometry of plasma and examined serum inflammatory cytokines in the macaques. A total of 34 fatty acids were detected, and among the five types of fatty acids, the concentration of saturated fatty acid (SFA) was significantly greater in the T2DM group ($p<0.05$, Figure 3A), while other types of fatty acids were not significantly different between the two groups ($p>0.05$, Figures 3B-E). In particular, concentrations of PA, palmitoleic acid, and oleic acid were

significantly higher in the T2DM group than control group ($p<0.05$ and VIP>1). The

concentration of PA in the plasma of T2DM macaques increased, while the

concentration of palmitic acid in the stool decreased (Figures 3F and G, Table S2).

Increased content of PA, palmitoleic acid, and oleic acid in the plasma was also found

in human T2DM (49, 50). PA was an important metabolite mediating insulin

resistance through three main mechanisms, being increased diacylglycerol and

ceramide synthesis, mitochondrial and endoplasmic reticulum stress, and activation of

pro-inflammatory pathways through membrane receptors (21, 22, 23, 24). Analysis on

the serum inflammatory cytokines found that IL-1β was significantly higher in the

T2DM group ($p<0.05$, Figure 3H), but TNF-α and IL-6 levels showed no significant

difference between the two groups ($p>0.05$, Figures 3I and J). IL-1β is a major player

in a variety of autoinflammatory diseases and a key promoter of T2DM systemic and

tissue inflammation (51). Moreover, blood routine examination showed an increase of

white blood cell (WBC) number, neutrophil (NEU) percentage and NEU number and

a decrease of lymphocyte (LYM) percentage and LYM number, also indicating the

inflammation in the T2DM macaques ($p<0.05$, Table 2).

To investigate effect of gut microbiota on the LCFAs accumulation and

inflammation in T2DM macaques, we performed a correlation analysis among the

DEGs, differential metabolites, and differential microbes using spearman rank

correlation. Four differential microbes in Lachnospiraceae family (*Coprococcus*,

Lachnospira, *Roseburia* and *Dorea longicatena*) were significantly associated with

three differential metabolites of PA, palmitoleic acid and oleic acid (|r|>0.5, adj

$p<0.05$), suggesting the participation of Lachnospiraceae microbes in LCFAs

accumulation in T2DM macaques (Figure 3K). In addition, the bacteria in class

Coriobacteriia was also associated with the three LCFAs (|r|>0.5, adj $p<0.05$, Figures

S1C and D).

255 Fecal microbiota transplantation (FMT) with high content PA food

256 induce prediabetes in mice

To determine the causative effect of gut microbiota and PA on T2DM development, we collected feces from the spontaneous T2DM macaques and performed fecal microbiota transplantation (FMT) in antibiotic-pretreated mice. Mice were either administrated by FMT (FT), fed with high concentration PA diet (PA), or were combined FMT and PA diet (FTPA). A control group was used and they were fed with normal commercial food and lacked FMT (Figure 4A). FPG monitoring found that FPG levels in the FTPA group and FT group increased continuously from day 60 (Figure 4B), while the control group maintained stable FPG levels throughout the 120 days. The FTPA group showed the highest FPG of 6.7 mmol/L at day 120, which was significantly higher than the control group and indicated FTPA-treated mice had developed prediabetes (Figure 4C). The oral glucose tolerance test (OGTT) showed obvious glucose intolerance in the FTPA group (Figures 4D and E), and the FPI values and insulin tolerance test (ITT) were both significantly elevated in the FTPA group, indicating insulin resistance (Figures 4F and G). Meanwhile, body weight (BW), TC, and TG significantly increased in the FTPA group (Figures 4H-J). While mice in the FT group and PA group had significant weight gain and mild insulin resistance, they did not exhibit significant glucose intolerance and no significant elevation in TC and TG levels compared to controls. Histopathological changes in the pancreas and liver of mice were investigated using H-E staining. Hepatocytes focal necrosis with inflammatory cell infiltration was commonly observed in the FTPA mice, but not frequent in hepatocytes in FT and PA groups (Figure 4K). Furthermore, decreased volume and area in pancreatic islets and inflammatory cell infiltration were detected in the FTPA mice (Figure 4L), while such pathological changes were not found in the control group. To eliminate interference from host species divergence in gut microbiota composition, we supplemented mouse experiments using FMT from control macaques (HFT group) (Figure S4A). By day 30, the HFT group exhibited significantly lower body weight than the untreated control group ($p<0.05$) (Figure S4B). Throughout the experimental period, FPG levels in both HFT and control groups remained within the normal range (<6 mmol/L) without significant differences,

indicating that transplantation of control macaque microbiota did not induce glycemic alterations (Figure S4C).

Specific structure of gut microbiota mediates the absorption of excess PA in the ileum

Given that the content of plasma PA significantly increased in the T2DM macaques (Figures 3F and G), we examined PA content in the feces, ileum, and serum in mice to compare the prediabetes mice and the spontaneous T2DM macaques. PA content in the serum and ileum of the FTPA group was significantly higher than the control group ($p<0.05$, Figures 5A and B), but its content in feces was significantly lower than the control group ($p<0.05$, Figure 5C). This suggested that the absorption of PA was significantly enhanced in the ileum leading to the increase of PA in serum. To verify this conclusion, expression of the *Cd36* gene, a gene involved in the uptake and oxidation of LCFAs (52), was examined in the ileum of mice. Ileums showed a significantly upregulated expression of *Cd36* in the FTPA group compared to the control group ($p<0.05$, Figure 5D). In contrast, the level of IL-17A, a protein inhibiting the expression of *Cd36* (53), was significantly reduced in the ileum of the FTPA group ($p<0.05$, Figure 5E). Interestingly, the PA group's content of PA, *Cd36* expression and IL-17A was not significantly different from the control group. This indicated that without FMT, the ileum could not absorb PA effectively even when fed in high concentrations. Consequently, gut microbiota mediated the absorption of excess PA in the ileum.

We then compared diversity and composition of microbiota communities between the FTPA, FT, PA and control groups, and a fifth group consisting of the microbiota transplants (TP) from T2DM macaques. Shannon and Simpson indices were lower in the FTPA, FT and PA groups than the control group (Figures S3A and B). NMDS analysis also indicated a distinct microbiota composition from the control group compared to other groups (NMDS1). In addition, microbiota composition of TP from T2DM macaques was distinct from other groups but a little closer to the FTPA

and FT groups (NMDS2, Figure 5F). In particular, the Lachnospiraceae family showed the higher abundance in the TP, FTPA and FT groups than in the PA and control groups (Figure S3C). The abundances of three members of microbes in Lachnospiraceae (*R. gnavus* (current name: *M. gnavus*), *Coprococcus* sp. and *Clostridium*) in the FTPA and FT groups gradually increased over time after FMT (Figure S3D). The FTPA and FT groups shared many differential microbes compared to the control group, such as the significantly upregulated *R. gnavus* (current name: *M. gnavus*) and *Coprococcus* sp., and the significantly downregulated Christensenellaceae, F16, *Treponema* sp. and *Fibrobacter succinogenes* (Figures 5G and H). It is noteworthy that these microbes were also differential microbes between T2DM macaques and controls, and their abundances changed in the same way as in macaques (Figure 1E). However, mice in the PA group did not share differential microbes with the spontaneous T2DM macaques. Correspondingly, the change of serum PA content in the PA group was not significantly higher than the control group (Figure S3E). Integrating 16S rRNA sequencing data from the HFT, FT, and FTPA groups showed that the antibiotic treatment effectively depleted the gut microbiota, resulting in microbial diversity decreased sharply, with the dominant phyla shifting from Bacteroidota and Bacillota to Pseudomonadota (Figure S4D-G). The HFT group restored microbial diversity within 30 days, achieving community proportions comparable to untreated controls. Core functional phyla (Bacteroidota and Bacillota) stably colonized in HFT group (Figure S4D-I). Critically, FT and FTPA groups exhibited increased Lachnospiraceae (including genera *Ruminococcus* (current name: *Mediterraneibacter*), *Coprococcus*, and *Clostridium*) compared with the HFT group on day 30. In addition, LEfSe comparison identified significant *R. gnavus* (current name: *M. gnavus*) enrichment in the FT group (LDA>3, $p<0.01$) (Figure S4J-M). Our results suggested that the transplanted microbiota from spontaneous T2DM macaques, especially the increased abundance of *R. gnavus* (current name: *M. gnavus*) and *Coprococcus* sp. and decreased abundance of *Treponema*, *F. succinogenes*, Christensenellaceae and F16, promoted the absorption of excess PA by regulating the

expression of IL-17A and Cd36, leading to the LCFAs accumulation and insulin

resistance (Figures 5I).

Discussion

With a multi-omics technology, this study comprehensively characterizes the gut

microbiota, metabolites and gene expression of spontaneous T2DM macaques. The

gut microbiota diversity in T2DM macaques decreased. In particular, the abundance

of bacteria *R. gnavus* (current name: *M. gnavus*) and Erysipelotrichaceae were

upregulated while the abundance of Christensenellaceae was downregulated in the

T2DM macaques. Metabolome results demonstrated a decrease of microbiota-derived

tryptophan and anti-inflammatory metabolites, indicating that T2DM macaques were

prone to inflammation. Notably, the accumulation of acylcarnitine metabolites, the

suggested biomarkers for human T2DM (50), indicated incomplete mitochondrial

LCFA β oxidation in T2DM macaques. Transcriptome results identified many DEGs

linked to insulin resistance, fatty acid β oxidation and inflammation, including

IGF2BP2, *LEPR*, *RAP1A*, *SESTRIN 3*, and *ITLN1* that have also been reported in

human T2DM (41, 42, 43, 44, 45). Combining the multi-omics results we revealed the

complex pathological mechanisms in the spontaneous T2DM macaques (Figure 6),

which is comparable to T2DM humans. Firstly, expression of genes related to

lipolysis, fatty acid oxidation, LCFA accumulation, inflammation, and insulin seretion

are dysregulated, promoting the development of T2DM (Figure 6A). This is

particularly related to lipid metabolism, where the increase of lipolysis and the

downregulated expression of fatty acid metabolism-related genes *HADHB* and

ACSM3 cause the accumulation of acylcarnitine and LCFAs, which results in

incomplete LCFA oxidation in T2DM macaques (Figure 6B). We also suggest that the

decrease of *Lactobacillus* sp. cause the reduction of Aryl hydrocarbon receptor (AhR)

ligands (serotonin and indole-3-acetaldehyde) given that *Lactobacillus* is producer of

AhR ligands (54). This then promotes the expansion of the PA producer

Erysipelotrichacea (55, 56) and ultimately leads to the increase of PA levels. The

increase in the abundance of Erysipelotrichacea and *R. gnavus* (current name: *M.*

gnavus) also promotes the development of T2DM by activation of inflammation (57, 58, 59) (Figure 6C). The accumulation of PA was reported to lead to insulin resistance by affecting genes in the insulin signaling pathway (16). Our study demonstrates that the significantly changed expression of *RAP1A*, *SESTRIN3*, and *IRS1* in PA-mTORC1-Akt pathway causes insulin resistance in T2DM macaques. Moreover, the increase of PA can promote the development of T2DM by upregulating the NF-κB signaling pathway (Figure 6D).

We validated the observations from the multi-omics analysis finding the significantly higher inflammatory cytokines IL-1β and LCFA accumulation, especially significant PA accumulation in the T2DM macaques. And found LCFA metabolites were significantly correlated with bacteria in the Lachnospiraceae family. Numerous studies imply an association of gut microbiota with T2DM development. By transplanting gut microbiota from healthy individuals to T2DM individuals, symptoms such as insulin resistance or inflammation were improved (6, 60). However, there is no evidence to date suggesting gut microbiota have directly causative effects on T2DM development. Our study confirmed the causative effect of gut microbiota and PA on T2DM development by transplanting fecal microbiota from spontaneous T2DM macaques to antibiotic pretreated mice. We successfully induced prediabetes in mice after combining FMT administration and high PA ingestion. However, when the treatments were administered on their own, the mice did not develop prediabetes. We determined, for the first time, that gut microbiota mediated the absorption of excess PA in the ileum by quantitative examining PA contents in feces, ileum, and serum, and analysis of the expression of *Cd36* and IL-17A level in the ileum. Most notably, this then resulted in accumulation of PA in the serum and finally led to T2DM development. Without the transplanting gut microbiota, the ileum could not absorb the PA effectively even at a high concentration of ingested PA. Our study highlights the essential roles of gut microbiota in T2DM development, which may account for the inability of prior studies to induce T2DM in macaques through high-fat diet intervention alone (28, 29). Furthermore, applying this approach to induce T2DM in

macaques will enable deeper investigation into gut-microbiota-driven mechanisms
underlying disease pathogenesis.

We then determined the specific gut microbiota structure that related to T2DM
development in the prediabetes mice and spontaneous T2DM macaques. We found
that the increased abundance of *R. gnavus* (current name: *M. gnavus*) and
Coprococcus sp., and the decreased abundance of *Treponema*, *F. succinogenes*,
Christensenellaceae, and F16, were involved in the T2DM development. *R. gnavus*
(current name: *M. gnavus*) can promote insulin resistance by regulating the content of
tryptamine/phenethylamine (61). Moreover, *R. gnavus* (current name: *M. gnavus*) is a
mucin-degrading microbe that leads to an increase of inflammation (62, 63, 64). The
intestinal mucous layer is an important barrier separating intestinal tissue from
microbiota, and microbiota composition plays a major role in affecting the integrity of
intestinal mucous layer (62). The increase of *R. gnavus* (current name: *M. gnavus*)
suggested a higher risk of damage to the integrity of the intestinal mucous layer. This
is further supported by our results of the lower abundance of *Lactobacillus* sp., which
are AhR ligand producers, and the decreased content of AhR ligands (tryptophan
microbial metabolites) in spontaneous T2DM macaques. The deficiency of AhR
ligand reduced the production of intestinal mucus and increased the risk of microbial
invasion, which in turn affected the immune cell differentiation and cytokine
production (65, 66). The cytokine IL-17A is a regulator of the fatty acid transporter
Cd36 and lipid absorption can be promoted by reducing the inhibition of *Cd36*
expression by IL-17A (58). The decreased IL-17A level and increased *Cd36*
expression in prediabetes mice indicated that the specific gut microbiota promoted the
absorption of excess PA by disrupting the integrity of the intestinal mucous layer and
regulating the expression of IL-17A and *Cd36*. In addition, the beneficial bacteria
decreased in abundance in T2DM macaques and prediabetes mice. These beneficial
bacteria, such as *F. succinogenes*, Christensenellaceae, and F16, protect the mucosal
barrier and improved insulin resistance (67, 68, 69). We inferred the collectively
effects of these gut microbes determined the absorption of excess PA from ileum to

serum, which might contribute to the development of T2DM. Previous studies have shown that insulin-resistant patients exhibit increased fecal monosaccharides associated with microbial carbohydrate metabolism (70). Furthermore, commensal species of Lachnospiraceae actively overproduce long-chain fatty acids during metabolic dysfunction through altered bacterial lipid metabolism. The microbe-derived fatty acids impair intestinal epithelial integrity to exacerbate metabolic dysregulation (71). Given that microbial metabolic activity causally modulates host metabolic homeostasis, the content change of PA was potentially associated with a dynamic equilibrium between host absorption and microbial metabolism. Further integrative studies on the fecal fatty acid metabolome, microbial PA metabolism, and functional pathways will be crucial for delineating causal links between dysbiosis and lipid metabolic dysfunction in T2DM.

In conclusion, spontaneous T2DM macaques that have never been treated with diabetes-related drugs provide a valuable model for our understanding of the pathological characteristics and pathogenesis of T2DM. This study characterized changes in gene expression, metabolites, and gut microbiota levels of spontaneous T2DM macaques using multi-omics techniques. We found gut abnormal microbiota, tryptophan metabolism and fatty acid β oxidation disorders, inflammation, and PA accumulation. We also successfully induced prediabetes in mice by transplanting fecal microbiota from T2DM macaques into antibiotic pretreated mice fed a high PA diet. Our study confirms the functional role of gut microbiota and PA in the T2DM progression. The microbiota composition, specifically higher abundance of *R. gnavus* (current name: *M. gnavus*) and *Coprococcus* sp. and lower abundance of *Treponema*, *F. succinogenes*, Christensenellaceae, and F16, promoted the absorption of excess PA which is important for the development of T2DM. However, in this study, such microbial alterations were detected in macaques after they had developed the disease of T2DM instead of before or onset of T2DM, the causative effect of gut microbiota and their action mechanism on the development of T2DM is worth further investigation. This study provides new insights into the interaction of gut microbiota

and metabolites in the development of T2DM, which expands our understanding of

the pathogenesis of this metabolic disease and may provide novel insights for the

treatment of T2DM in the future.

Materials and methods

The screening of spontaneous T2DM macaques

The experimental macaques used in this study were all from Greenhouse

Biotechnology Co., LTD (Meishan, China). We obtained eight spontaneous T2DM

macaques with FPG≥7 mmol/L and eight heathy control macaques with FPG≤6.1

mmol/L (three consecutive detections, each detection interval of one month) from a

population of 1698 captive macaques. None of these 16 screened macaques received

any medical treatment for diabetes.

Sample collection

Each experimental macaque was kept in a single cage, and was fasted for 12 hours but

had free access to drinking water. We obtained serum, plasma and whole blood for the

detection of physiological and biochemical parameters, and metabolome and

transcriptome analysis. Fecal samples were collected within ten minutes after

deposition. During the sampling process, fecal samples were loaded into a 50 mL

sterile centrifuge tube were stored at -80□. We followed animal welfare guidelines

throughout the sample collection process, and all observations and samplings were

approved by the Sichuan University's Animal Care Committee.

Physiological and biochemical parameters

In this study, FPG, HbA1c and FPI were detected by hexokinase method, high

performance liquid chromatography and electro-chemiluminescence method,

respectively. TC, TG, HDL and LDL were detected by automatic biochemical

analyzer. HOMA-IR is one of the criteria for T2DM, calculated as: (FPI×FPG)/22.5.

Feces 16S rRNA amplicon sequencing and analysis

Total DNA from fecal samples was extracted using the QIAamp Fast DNA Stool Mini

Kit. The V3-V4 region of the 16S rRNA gene was amplified using the 341F/806R

primer (341F: 5'-CCTAYGGGRBGCASCAG -3', 806R:

5'-GGACTACNNGGGTATCTAAT -3'). The purification and quantification of the

amplified products were performed and followed by the sequencing library

preparation with TruSeq Nano DNA LT Library Prep Kit (Illumina, USA). The library

sequencing was performed on Illumina MiSeq platform and 250 bp paired-end reads

were generated. Raw sequencing reads were merged using FLASH (VI.2.7,

http://ccb.jhu.edu/software/FLASH/) (72) and analyzed by QIIME2 (version

2020.11.1) pipeline with default parameters. Reads after denoising by DADA2 were

clustered into OTUs at 99% similarity threshold. Taxonomy of OTUs was assigned

based on Greengenes reference database (73). The QIIME2 diversity plugin was used

to calculate alpha diversity (74). Principal Coordinate Analysis (PCoA) was

determined by using the R package vegan. Differentially microbes were determined

using linear discriminant analysis effect size (LEfSe) (75).

Feces shotgun metagenome sequencing and analysis

Total DNA from each sample was extracted using Tiangen DNA Stool Mini Kit

(Tiangen Biotech Co., Ltd., China). Metagenome sequencing was performed using the

Illumina NovaSeq 6000 platform with a paired-end sequencing length of 150 bp.

Trimmomatic was used for removing the adapters and low-quality raw reads based on

a four-base wide sliding window, with average quality per base>20 and minimum

length 90 bp (76). The rhesus macaque potential sequences were removed using

Bowtie2 (77) with the reference genome (assembly Mmul_10). Taxonomy of

remaining reads was assigned using Kraken2 (78) with the option "--use-mpa-style".

De novo assembly of remaining reads was performed using MEGAHIT (79) with the

option "-m 0.90 --min-contig-len 300". Prodigal (80) was used for gene prediction.

The construction of non-redundant gene catalogue was performed with CD-HIT (81)

with the option "-c 0.95 -aS 0.90". Quantification of the non-redundant genes in each

sample was performed using Salmon (82). The amino acid sequences translated from

non-redundant genes were aligned (--id 80% --query-cover 70% --evalue 1e-5) by

DIAMOND (83) in the Carbohydrate-Active enZYmes (CAZy) database (84). The

annotation metabolic pathway was performed using HUMANn3 (85).

RNA sequencing and DEG analysis

Total RNA was extracted using PAXgene Blood RNA kit. The cDNA Library was constructed following the NEBNext® UltraTM RNA Library Prep Kit for Illumina® (NEB, USA) manual, and index was added to each sample for sample differentiation. cBot Cluster Generation System was used to cluster the sequences with the same index. Illumina Hiseq 2500 sequencing platform was used to obtain the paired-end sequencing reads (PE150). NGS QC Toolkit v2.3.3 (86) was used to obtain high quality reads (high-quality paired reads with more than 90% of bases with Q-value≥20 were retained). Processed reads of each sample were mapped to the macaque reference genome using HISAT2 v2.1.0 (Kim, et al. 2015). Each alignment output file was assembled into a separate transcriptome using StringTie v1.3.6 (87), resulting in a transcript GTF file. To obtain the expression value of TPM (transcripts per million) and raw read counts of each gene, transcript GTF file was used as the reference annotation file. Differential expression analysis was performed using DESeq2 R package (88). DEGs were screened according to Foldchange value (FC) and p value corrected by FDR (Benjamini-Hochberg method was selected) (adj $p<0.05$, |log2FC|>1).

Weighted Gene Co-Expression Network Analysis (WGCNA) (89) was used to analyse the correlation between genes and phenotypes. We selected an appropriate "soft thresholding power" using the "picksoftthreshold" function in the WGCNA package (v1.61). Next, "blockwiseModules" function was used to construct the co-expression matrix with the option "checkMissingData=TRUE, power=16, TOMType='unsigned', minModuleSize=30, maxBlockSize=6000, mergeCutHeight=0.25". The modules with high correlation to T2DM phenotype and $p<0.05$ were selected for downstream analysis, and genes in these modules were used for GO and KEGG functional enrichment analysis. Hub genes were identified based on the module eigengene-based connectivity (kME), |kME>0.8| as the cut-off criteria.

Instead of identifying the differentially expressed genes within a pathway between the two groups, Aggregate Fold Change (AFC) calculated the average

multiple change for each gene and defined the pathway score as the average

difference multiple for all genes in that pathway. A null hypothesis test was performed

using the pathway scores of the gene expression dataset, and the significance of each

pathway was estimated by p-value (90). STRING (v11.0) provides a tool for

functional enrichment analysis based on AFC. GO and KEGG enrichment analysis

was performed using g: Profiler (91), AFC enrichment analysis was performed using

STRING

(https://www.string-db.org/cgi/input?sessionId=bfRWW6asD8S4&input_page_show_

search=on).

Untargeted metabolomics processing

The 200 μL homogenized fecal sample was mixed with 800 μL cold

methanol/acetonitrile (1:1, v/v) to remove the protein. The mixture was centrifuged

for 15 min (14000g, 4°C) followed by the drying of the supernatant in a vacuum

centrifuge. For LC-MS analysis, the samples were re-dissolved in 100 μL

acetonitrile/water (1:1, v/v) solvent. LC-MS/MS analysis was performed using an

UHPLC (1290 Infinity LC, Agilent Technologies) coupled to a quadrupole

time-of-flight (AB Sciex TripleTOF 6600) in Shanghai Applied Protein Technology

Co., Ltd. The samples were separated by Agilent 1290 Infinity LC ultra-high

performance liquid chromatography (UHPLC) HILIC column. In both ESI positive

and negative modes, the mobile phase contained A=25 mM ammonium acetate and 25

mM ammonium hydroxide in water and B=acetonitrile. The gradient was 85% B for 1

min and was linearly reduced to 65% in 11 min, and then was reduced to 40% in 0.1

min and kept for 4 min, and then increased to 85% in 0.1 min, with a 5 min

re-equilibration period employed. AB Triple TOF 6600 mass spectrometer was used

to collect the primary and secondary spectra of per sample. After separation by

UHPLC, the samples were analyzed by Triple TOF 6600 mass spectrometer (AB

SCIEX). Positive and negative modes of electrospray ionization (ESI) were

respectively detected.

Targeted medium-and long-chain fatty acid metabolomics processing

A total amount of 100 μL plasma per sample was taken in 2 mL glass centrifuge tubes

and 1mL chloroform methanol solution was added. After 30min of ultrasound, the

supernatant was taken and 2mL of 1% sulfuric acid-methanol solution was added. The

mixed solution was placed in a water bath at 80□, for 30min, and then 1mL of

N-hexane and 5 mL of pure water were added in turn. Next, 500 μL of supernatant

was absorbed, 25 μL of methyl N-nonaconate was added and mixed. The final sample

was detected by GC-MS. All samples were separated by Agilent DB-WAX capillary

column (30 m×0.25 mm ID×0.25 μm) gas chromatography. Agilent 7890/5975C

gas-mass spectrometer was used for mass spectrometry. The chromatographic peak

area and retention time were extracted by MSD ChemStation software. The content of

medium-and long-chain fatty acids was calculated by drawing a standard curve.

Metabolomics statistical processing

The screening of significant changed metabolites was performed using univariate and

multidimensional analysis. Student's t test was applied to determine the significance

of differences between two groups. The variable importance in the projection (VIP)

value of each variable was obtained from the Orthogonal Partial Least Squares

Discriminant Analysis (OPLS-DA) model, which was used to indicate its contribution

to the classification. The screening criteria were $p<0.05$ and VIP>1.

Animal treatments

Specific-pathogen-free C57BL/6J male mice at 6 weeks of age (vendor: Chengdu

dossy experimental animals Co., LTD) were randomly assigned to four groups (FTPA,

FT, PA and Control). All the mice lived in cages with the same conditions, including

12h light and 12h dark cycles, temperature 22-25□°C and humidity 40-60%.

Diets

HPAD was prepared by adding 12% PA to conventional forage. Both the conventional

forage and the HPAD were sterile and the fresh forage was renewed three times a

week. On day 0, the diets of FTPA-treated mice and PA-treated mice were switched to

HPAD, FT-treated mice and control mice were still fed with conventional forage.

Drinking water was sterile and renewed twice a day.

Transplant preparation and use

After single cage feeding, FPG detection and fecal collection were performed, and fecal samples of seven T2DM macaques were mixed for the preparation of transplants. The appropriate volume of diluent was added to the fecal sample (i.e. add 4 ml diluent per gram of feces) and the preparation of diluent can be found in (92). Sodium L-ascorbic acid and L-cysteine hydrochloride monohydrate were added to all suspensions at final concentrations of 5% (w/v) and 0.1% (w/v), respectively (The sterile diluent of control group was added with the same amount of reagent). The mixture was homogenized and filtered with a 200-mesh sterile mesh screen to remove large particles from the feces, and the filtrate was passed through 400 and 800 sterile mesh screens to remove undigested food and smaller particulate matter. The filtrate was divided into 50 ml centrifuge tubes, centrifuged at 600×g for 5 min, and the precipitation was discarded. Finally, the fecal supernatant was divided into new centrifuge tubes in equal parts (400 μL per tube) and frozen at -80°C. For use, the transplant was quickly thawed in a 37☐ water bath.

Fecal microbiota transplantation (FMT)

After one week of changed feeding regime, the FTPA and FT mice groups were pre-treated with 1 g/L neomycin sulfate, 1 g/L ampicillin and 1 g/L metronidazole in the drinking water for 14 days, and the control group was not treated. For FMT treatment, the gavage with 400 μL transplant, which were thawed ahead of time, was performed for 14 days in FTPA and FT-treated mice. At the same time, the gavage in control and PA-treated mice were performed with sterile diluent.

Metabolic measurements

Throughout the experiment, body weight and feces were collected every month, FPG was detected every half month under fasting at least 12 h. OGTT was performed on day 110 and ITT was performed on day 115. For OGTT, after a 12h overnight fast, oral glucose gavage (1.2 g/kg of 12% dextrose solution) was performed and followed by blood sample collection from the tail vein at 0, 15, 30, 60, 90 and 120 min. For ITT, after a 6h fast, intraperitoneal insulin injection was performed (0.75 U/kg, human

regular insulin), followed by the blood samples collection from the tail vein at 0, 20,

40, 60, 90 and 120 min. On day 120, after FPG detection, blood collection and issues

were collected for quantitative RT-PCR, H-E staining and ELISA.

Isolation of tissue for quantitative RT-PCR and ELISA

Tissue was homogenized for RNA extraction, after adjusting the final concentration of

RNA, and DNA reverse transcription was performed. Quantitative RT-PCR was

performed on CFX Connect Real-Time PCR Detection System (Bio-rad, USA) using

SYBR Green (Table 3-5). PA, IL-1β, IL-6, TNF-α and IL-17A were detected using

Jiangsu Meimian ELISA kit and followed operator instructions.

H-E staining

The isolated livers and pancreas were dehydrated and embedded after fixation with

formalin for 24 h. Paraffin-embedded livers and pancreas specimens were cut at a

thickness of 3 μm. All sections were stained with hematoxylin then eosin, and finally

microscopy and image acquisition were performed.

Statistical analysis

In this study, one-way ANOVA was used to determine statistical significance for

comparisons of more than three groups, and for comparisons of two groups,

two-tailed t-test was used. p values are represented on figures as follows: ns, not

significant, $*p<0.05$, $**p<0.01$.

Data availability

The raw data of transcriptomes, metagenomes, and 16S rRNA have been submitted to

the China National Center for Bioinformation/Beijing Institute of Genomics, Chinese

Academy of Sciences with BioProject accession no. PRJCA021499. We provide four

links that reviewers can use to access the raw data. Transcriptomes data:

https://ngdc.cncb.ac.cn/gsa/s/4xAVJ3Lk (GSA: CRA013604). 16S rRNA data

(macaque): https://ngdc.cncb.ac.cn/gsa/s/bsDz9qrN (GSA: CRA013637). 16S rRNA

data (mouse): https://ngdc.cncb.ac.cn/gsa/s/1YcuC464 (GSA: CRA013638).

metagenomes data: https://ngdc.cncb.ac.cn/gsa/s/3k38T849 (GSA: CRA013607). The

identified untargeted metabolites were listed in the Table S4. The identified targeted
medium-and long-chain fatty acid metabolites were listed in the Table S5.

Acknowledgements

This work was supported by the Science and Technology Foundation of Sichuan
Province (2021YJ0136) and the National Natural Science Foundation of China (No.
32171607). Special thanks to Sichuan Green-house Biotech Co., Ltd. for the sample
collection and thanks to Prof. Jinchuan Xing and Dr. Megan Price for revising the
manuscript.

Author contributions

X.L., G.H., Q.H.L., S.Z.Y. and C.J. collected the samples; X.L. and S.Z.Y. performed
the bioinformatics analyses; X.L., Y.C.X., K.S., C.J., J.X.L. and L.Z. performed the
experiments; X.L. wrote the manuscript; J.L., Z.L.H., Z.X.F. and B.S.Y. revised the
manuscript; J.L. conceived and designed the experiments.

Declaration of interests

The authors declare no competing interests.

Figure legends

Figure 1. The changes in gut microbiota in spontaneous T2DM macaques.

(A) Alpha diversity estimates (Shannon index) between T2DM and control groups (ns,
not significant, two-tailed t-test, n=8).

(B) Alpha diversity estimates (Simpson index) between T2DM and control groups (ns,
not significant, two-tailed t-test, n=8).

(C) Principal Coordinate Analysis (PCoA) (n=8).

(D) Differential analysis of gut microbial composition in T2DM and control groups
(n=8).

(E) LEfSe analysis between T2DM and control groups (n=8).

(F) Differential analysis of gut microbial function in T2DM and control groups (n=5).

The pathways with red color were associated with T2DM and inflammation. Error bar

is mean with ± standard deviation (s.d.).

(G) Differential analysis of gut microbial CAZy enzyme in T2DM and control groups

(n=5). CBMs: carbohydrate-binding module (p=0.022, two-tailed t-test); GTs:

Glycosyl Transferases (p=0.013, two-tailed t-test); PLs: Polysaccharide Lyases

(p=0.017, two-tailed t-test); AA: Auxiliary activity enzymes (ns, not significant,

two-tailed t-test); GH: Glycoside hydrolases (ns, not significant, two-tailed t-test); CE:

Carbohydrate esterases (p=0.039, two-tailed t-test). For all boxplots: centre lines, upper

and lower bounds show median values, 25th and 75th quantiles; upper and lower

whiskers show the largest and smallest non-outlier values. In c, ellipses represent the

95% confidence intervals.

Figure 2. The alterations of fecal metabolites and gene expression in spontaneous

T2DM macaques.

(A) Orthogonal partial least squares discriminant analysis (OPLS-DA) score plots

based on the metabolic profiles.

(B) Volcano plots of metabolomics (p<0.05, two-tailed t-test).

(C) Fecal metabolites with significant differences between T2DM and control groups

(VIP>1, p<0.05, two-tailed t-test).

(D) Enrichment analysis of the differentially abundant pathways between T2DM and

control groups (p<0.05, two-tailed t-test).

(E) Non-metric multidimensional scaling (NMDS) analysis between T2DM and

control groups (p=0.019, two-tailed t-test).

(F) Volcano plots of DEGs (log fold change≥1, p<0.05, two-tailed t-test).

(G) The GO and KEGG pathway enrichment analyses (p<0.05, two-tailed t-test).

(H) Weighted Gene Co-Expression Network Analysis (WGCNA).

(I) Venn analysis between hub genes and DEGs. In A and E, ellipses represent the 95%

confidence intervals. Data shown are from 8 individuals per group.

Figure 3. LCFAs accumulation and inflammation occurred in spontaneous T2DM

macaques.

(A-E) The contents of SFA (A, p=0.038), MUFA (B), PUFA (C), N3 (D), and N6 (E) in

plasma (ns, not significant, two-tailed t-test).

(F) The univariate analysis by two-tailed t-test, error bar is mean with $\pm$ s.d.

(G) The multidimensional analysis by VIP value (VIP>1).

(H-J) The contents of serum inflammatory cytokines, including IL-1β (H, p=0.032),

TNF-α (I) and IL-6 (J) (ns, not significant, two-tailed t-test).

(K) Correlation analysis between DEGs, differential metabolites, and differential

microbes using Spearman rank correlation (|r|>0.5, adj p<0.05). For all boxplots: centre

lines, upper and lower bounds show median values, 25th and 75th quantiles; upper and

lower whiskers show the largest and smallest non-outlier values. Data shown are from 7

individuals per group.

Figure 4. The FMT and high PA diet mice developed pre-T2DM characteristics.

(A) Experimental scheme of FMT and high PA diet treatment.

(B-H) Metabolic analysis, including the trend of FPG within 120 days (B), FPG (C,

p=0.0003), OGTT (D), AUC of OGTT (E, p=0.028), FPI (F, p=0.007), ITT (G), and

body weight change (H) on day 120.

(I-J) The contents of TC (I, p=0.005) and TG (J, p=0.041) in serum on day 120.

(K and L) Representative H-E staining images of liver (K) and pancreas (L). For all

boxplots: centre lines, upper and lower bounds show median values, 25th and 75th

quantiles; upper and lower whiskers show the largest and smallest non-outlier values.

Significance was determined using one-way ANOVA. In d, g, and h: *$p<0.05$,

**$p<0.01$. Data shown are from 4-6 individuals per group.

Figure 5. The PA accumulation required the specific gut microbiota.

(A-C) Total PA contents in serum (A, $p=0.013$), ileum (B, $p=0.016$) and feces (C,

$p=0.014$) on day 120.

(D) Quantitative RT-PCR for Cd36 transcripts in ileum on day 120 ($p=0.049$).

(E) The content of IL-17A in ileum on day 120 ($p=0.027$). For all boxplots: centre lines,

upper and lower bounds show median values, 25th and 75th quantiles; upper and lower

whiskers show the largest and smallest non-outlier values. Significance was determined

using one-way ANOVA. Data shown are from 3-4 individual macaques per group.

(F) NMDS analysis (p=0.001, one-way ANOVA), ellipses represent the 95%

confidence intervals.

(G and H) LEfSe analysis between FTPA and control groups (G), FT and control

groups (H). Data shown are from 4 individuals per group.

(I) Specific gut microbiota structure promoted the absorption of excess PA by

regulating the expression of IL-17A and *Cd36*, leading to the LCFAs accumulation

and insulin resistance.

Figure 6. Integration of multi-omics results.

(A) Insulin resistance, fatty acid oxidation disorders, LCFAs accumulation and

inflammation occurred in spontaneous T2DM macaques.

(B) Incomplete mitochondrial LCFAs β oxidation. The expression levels of fatty acid

metabolism-related genes HADHB and ACSM3 were downregulated in spontaneous

T2DM macaques, which could lead to accumulation of acylcarnitine, including

l-propionylcarnitine, hexanoyl-l-carnitine, (r)-butyrylcarnitine, and

isovaleryl-l-carnitine.

(C) Gut inflammation. The decrease of *Lactobacillus sp.* likely caused the reduction of

serotonin and indole-3-acetaldehyde, which promotes the expansion of PA producer

Erysipelotrichacea and ultimately led to PA accumulation. Both Erysipelotrichacea and

Ruminococcus gnavus (current name: *Mediterraneibacter gnavus*) promote the

development of inflammation. Accumulation of PA and inflammation are important

factors in the development of T2DM.

(D) Accumulation of PA promoted the development of insulin resistance. In the

PA-mTORC1-Akt pathway, the changes of *RAP1A*, *SESTRIN3*, and *IRS1* expression

promoted the development of insulin resistance in spontaneous T2DM macaques. The

increase of PA promoted the development of T2DM by up-regulating the NF-κB

signaling pathway.

Tables

Table 1 Physiological and biochemical parameters of Control and T2DM group

Index	T2DM (n=8)	Control (n=8)
FPG (mmol/L)	7.75±1	4.05±1.02[**]
FPI (μU/mL)	17.97±8.50	6.45±2.56[**]
HOMA-IR	6.24±3.06	1.20±0.65[**]
BMI	14.03±7.71	14.30±1.42
HbA1c (%)	3.78±0.70	3.26±0.62
TG (mmol/L)	0.97±0.53	0.68±0.42
TC (mmol/L)	3.30±0.98	3.52±0.86
HDL (mmol/L)	1.23±0.40	1.42±0.38
LDL (mmol/L)	1.36±0.44	1.47±0.57

 *p<0.05, ** p<0.01

FPG: fasting plasma glucose (normal range: ≤ 6.1 mmol/L); FPI: fasting plasma

insulin (normal range: ≤ 12 μU/mL); HOMA-IR: homeostasismodel assessment-

insulin resistance (normal range: ≤ 2.67); BMI: body mass index; HbA1c:

glycosylated hemoglobin A1c (normal range: < 6.5%); TG: triglycerides (normal

range: 0.95±0.47 mmol/L); TC: total cholesterol (normal range: 3.06±0.98 mmol/L);

HDL: high-density lipoprotein cholesterol (normal range: 1.62±0.46 mmol/L); LDL:

low-density lipoprotein cholesterol (normal range: 2.47±0.98 mmol/L). (30, 31, 32,

33).

Table 2 blood routine examination of Control and T2DM group

Index	T2DM (n=7)	Control (n=7)
WBC (10e9/L)	15.63±4.66	11.32±2.19*
RBC (10e12/L)	5.47±0.51	5.77±0.43
HGB (g/L)	129.57±15.48	136.71±9.60
HCT (%)	41.31±3.64	44.22±3.12
MCV (fl)	74.13±2.49	76.74±2.00
MCH (pg)	23.17±1.00	23.71±0.60
MCHC (g/L)	312.86±10.16	309±8.14
RDW (%)	15.23±2.12	14.69±1.74
PLT (10e9/L)	402±86.66	371.57±86.42
MPV (fl)	10.54±1.62	10.24±1.12
PCT (%)	0.42±0.08	0.38±0.09
PDW (%)	14.94±0.58	14.83±1.64
LYM% (%)	23.67±10.26	47.71±8.13**
LYM# (10e9/L)	3.46±1.66	5.39±1.52*
MON% (%)	4.18±3.34	6.18±2.29
MON# (10e9/L)	0.65±0.62	0.71±0.38
NEU% (%)	71.13±13.23	44.28±8.96**

NEU# (10e9/L)	11.36±4.91	5.00±1.38**
EOS% (%)	0.91±0.69	1.50±1.94
EOS# (10e9/L)	0.14±0.12	0.17±0.22
BAS% (%)	0.12±0.13	0.33±0.47
BAS# (10e9/L)	0.02±0.02	0.04±0.07
NRBC# (10e9/L)	0	0
NRBC% (%)	0	0

* $p<0.05$, ** $p<0.01$

WBC: white blood cell; RBC: red blood cell; HGB: hemoglobin; HCT: hematocrit;

MCV: mean corpuscular volume; MCH: mean corpuscular hemoglobin; MCHC:

mean corpuscular hemoglobin concentration; RDW: red blood cell distribution width;

PLT: platelet; MPV: mean platelet volume; PCT: procalcitonin; PDW: platelet volume

distribution width; LYM%: lymphocyte percentage; LYM#: lymphocyte value;

MON%: monocytes percentage; MON#: monocytes value; NEU%: neutrophil

percentage; NEU#: neutrophil value; EOS%: eoseosinophil percentage; EOS#:

eoseosinophil value; BAS%: basophil percentage; BAS#: basophil value; NRBC%:

nucleated red blood cell percentage; NRBC#: nucleated red blood cell value.

Table 3 Primers of RT-PCR

Gene name	Sequences of primers
CD36/F	5'-ATGGGCTGTGATCGGAACTG-3'
CD36/R	5'-GTCTTCCCAATAAGCATGTCTCC-3'
GADPH/F	5'-CCTCGTCCCGTAGACAAAATG-3'
GADPH/R	5'-TCTCCACTTTGCCACTGCAA-3'

804 **Table 4 RT-PCR reaction components**

Reagent	Volume
2×SG Fast qPCR Master Mix	10.0 μL
DNF Buffer	2.0 μL
F primer (10umol/L)	0.4 μL

Reagent	Volume
R primer (10umol/L)	0.4 μL
cDNA	1.0 μL
ddH$_2$O	6.2 μL

Table 5 RT-PCR cycle procotol

Step	Temperature	Time
1	95□	3 min
2	95□	1-3 s
3	60□	30 s
4		
5	72□	1 min

Supplementary Figures

Figure S1 Metagenome analysis of microbiota.

(A) Differential microbes screened by metagenome analysis ($p<0.05$, two-tailed t-test).

T2DM: type 2 diabetes mellitus

(B) The proportion of all metabolites.

(C) Correlation analysis between differential untargeted metabolites and differential

microbes (Spearman's Rho).

(D) Correlation analysis between differential targeted metabolites and differential

microbes (Spearman's Rho), *$p<0.05$. Data shown are from 5 individuals per group.

Figure S2 Pathway enrichment analysis by AFC and WGCNA.

(A) Total 26 differential pathways were enriched (FDR<0.05).

(B) The changes of gene expression in insulin resistance pathway compared to the

control group, red color illustrating the up-regulation of genes and blue showed the

down-regulation of genes in T2DM group, and the darker the color of the genes, the

greater the $|\log_{10}FC|$ value.

(C)WGCNA functional enrichment. Data shown are from 8 individuals per group.

Figure S3 The changes in gut microbiota in FTPA, FT, and PA-treated mice.

(A) Alpha diversity estimates (Shannon index) ($p>0.05$, one-way ANOVA, n=4).

(B) Alpha diversity estimates (Simpson index) between T2DM and control groups

($p>0.05$, one-way ANOVA, n=4).

(C) Family level taxonomy and relative abundance of five groups.

(D) The changes of members of Lachnospiraceae family from day -14 to day 120.

(E) LEfSe analysis between PA and control groups. Data shown are from 4

individuals per group.

Figure S4 The changes in gut microbiota in HFT-treated mice

(A) Experimental scheme of fecal FMT from control macaques.

(B and C) The physiological characteristic of mice with different treatments,

including body weight change (B), and trend of FPG within 30 days (C). * $p<0.05$.

(D-F) Changes in gut microbiota α-diversity and structure in mice, including Shannon

index ($p>0.05$, one-way ANOVA) (D), Simpson index ($p>0.05$, one-way ANOVA) (E),

and NMDS analysis ($p=0.001$, one-way ANOVA) (F).

(G-I) Composition of gut microbiota in mice at phylum level (G), family level (H),

and genus level (I).

(J and K) The relative abundance of members in Lachnospiraceae between HTP and

TP groups (J), control, HFT, FT, and FTPA groups on day 30 (K).

(L and M) LEfSe analysis between pre-antibiotic (-14D) and post-antibiotic (0D)

groups (L), HFT and FT groups on day 30 (M).

Supplementary Tables: Table. S1.xlsx- Table. S5.xlsx

Table S1. Information on the macaque samples.

Table S2. List of 64 differential metabolites.

Table S3. List of genes in four modules significantly correlated with T2DM.

Table S4. List of all identified untargeted metabolites.

Table S5. List of all targeted medium-and long-chain fatty acid metabolites.

References

1.Rayburn, W.F. (1997). Diagnosis and classification of diabetes mellitus: highlights

from the American Diabetes Association. J Reprod Med *42*, 585–586.

2.Zhang, L., Chu, J., Hao, W., Zhang, J., Li, H., Yang, C., Yang, J., Chen, X., and

Wang, H. (2021). Gut microbiota and type 2 diabetes mellitus: Association,

mechanism, and translational applications. Mediators Inflamm. *17*, e5110276.

https://doi.org/10.1155/2021/5110276.

3.Kahn, S.E., Cooper, M.E., and Del Prato, S. (2014). Pathophysiology and treatment

of type 2 diabetes: perspectives on the past, present, and future. Lancet *383*,

1068–1083. https://doi.org/10.1016/s0140-6736(13)62154-6.

4.Kang, N., Chen, G., Tu, R., Liao, W., Liu, X., Dong, X., Li, R., Pan, M., Yin, S., Hu,

K., et al. (2022). Adverse associations of different obesity measures and the

interactions with long-term exposure to air pollutants with prevalent type 2 diabetes

mellitus: The Henan Rural Cohort study. Environ. Res. *207*, 112640.

https://doi.org/10.1016/j.envres.2021.112640.

5.Qin, J., Li, Y., Cai, Z., Li, S., Zhu, J., Zhang, F., Liang, S., Zhang, W., Guan, Y.,

Shen, D., et al. (2012). A metagenome-wide association study of gut microbiota in

type 2 diabetes. Nature *490*, 55–60. https://doi.org/10.1038/nature11450.

6.Zhao, L., Zhang, F., Ding, X., Wu, G., Lam, Y.Y., Wang, X., Fu, H., Xue, X., Lu, C.,

Ma, J., et al. (2018). Gut bacteria selectively promoted by dietary fibers alleviate type

2 diabetes. Science *359*, 1151–1156. https://doi.org/10.1126/science.aao5774.

7.Karlsson, F.H., Tremaroli, V., Nookaew, I., Bergström, G., Behre, C.J., Fagerberg,

B., Nielsen, J., and Bäckhed, F. (2013). Gut metagenome in European women with

normal, impaired and diabetic glucose control. Nature *498*, 99–103.

https://doi.org/10.1038/nature12198.

8.Tilg, H., and Moschen, A.R. (2014). Microbiota and diabetes: an evolving

relationship. Gut *63*, 1513–1521. https://doi.org/10.1136/gutjnl-2014-306928.

9.Hendrikx, T., and Schnabl, B. (2019). Indoles: metabolites produced by intestinal

bacteria capable of controlling liver disease manifestation. J. Intern. Med. *286*, 32–40.

https://doi.org/10.1111/joim.12892.

10.Qi, Q., Li, J., Yu, B., Moon, J., Chai, J.C., Merino, J., Hu, J., Ruiz-Canela, M.,

Rebholz, C., Wang, Z., et al. (2022). Host and gut microbial tryptophan metabolism

and type 2 diabetes: an integrative analysis of host genetics, diet, gut microbiome and

circulating metabolites in cohort studies. Gut *71*, 1098–1105.

https://doi.org/10.1136/gutjnl-2021-324053.

11.Wu, Z., Zhang, B., Chen, F., Xia, R., Zhu, D., Chen, B., Lin, A., Zheng, C., Hou,

D., Li, X., et al. (2023). Fecal microbiota transplantation reverses insulin resistance in

type 2 diabetes: A randomized, controlled, prospective study. Frontiers in Cellular and

Infection Microbiology 12. https://doi.org/10.3389/fcimb.2022.1089991.

12.Ng, S.C., Xu, Z., Mak, J.W.Y., Yang, K., Liu, Q., Zuo, T., Tang, W., Lau, L., Lui,

R.N., Wong, S.H., et al. (2021). Microbiota engraftment after faecal microbiota

transplantation in obese subjects with type 2 diabetes: a 24-week, double-blind,

randomised controlled trial. Gut *71*, 716–723.

https://doi.org/10.1136/gutjnl-2020-323617.

13.Ding, D., Yong, H., Na, Y., Lu, W., Xu, Y., Ye, X., Wang, Y., Cai, T., Zheng, X.,

Chen, H., et al. (2022). Prospective study reveals host microbial determinants of

clinical response to fecal microbiota transplant therapy in type 2 diabetes patients.

Front. Cell. Infect. Microbiol. *12*, 820367.

https://doi.org/10.3389/fcimb.2022.820367.

14.Unger, R.H., and Zhou, Y. (2001). Lipotoxicity of beta-cells in obesity and in other

causes of fatty acid spillover. Diabetes *50*, S118–S121.

https://doi.org/10.2337/diabetes.50.2007.s118.

15.Poitout, V., and Robertson, R.P. (2008). Glucolipotoxicity: fuel excess and β-cell

dysfunction. Endocr. Rev. *29*, 351–366. https://doi.org/10.1210/er.2007-0023.

16.Palomer, X., Pizarro-Delgado, J., Barroso, E., and Vázquez-Carrera, M. (2018).

Palmitic and oleic acid: the yin and yang of fatty acids in type 2 diabetes mellitus.

Trends Endocrinol. Metab. *29*, 178–190. https://doi.org/10.1016/j.tem.2017.11.009.

17.Milanski, M., Degasperi, G., Coope, A., Morari, J., Denis, R., Cintra, D.E.,

Tsukumo, D.M.L., Anhe, G., Amaral, M.E., Takahashi, H.K., et al. (2009). Saturated

fatty acids produce an inflammatory response predominantly through the activation of

TLR4 signaling in hypothalamus: implications for the pathogenesis of obesity. J.

Neurosci. *29*, 359–370. https://doi.org/10.1523/jneurosci.2760-08.2009.

18.Fiehn, O., Garvey, W.T., Newman, J.W., Lok, K.H., Hoppel, C.L., and Adams, S.H.

(2010). Plasma metabolomic profiles reflective of glucose homeostasis in

non-diabetic and type 2 diabetic obese African-American women. PLoS One *5*,

e15234. https://doi.org/10.1371/journal.pone.0015234.

19.Li, X., Xu, Z., Lu, X., Yang, X., Yin, P., Kong, H., Yu, Y., and Xu, G. (2009).

Comprehensive two-dimensional gas chromatography/time-of-flight mass

spectrometry for metabonomics: Biomarker discovery for diabetes mellitus. Anal.

Chim. Acta *633*, 257–262. https://doi.org/10.1016/j.aca.2008.11.058.

20.Chandra, K., Jain, V., Azhar, M., Khan, W., Alam, O., Ahmad, S., and Jain, S.K.

(2020). Effect of augmented glycation in mobilization of plasma free fatty acids in

type 2 diabetes mellitus. Diabetes Metab Syndr *14*, 1385–1389.

https://doi.org/10.1016/j.dsx.2020.07.028.

21.Ertunc, M.E., and Hotamisligil, G.S. (2016). Lipid signaling and lipotoxicity in

metaflammation: indications for metabolic disease pathogenesis and treatment. J.

Lipid Res. *57*, 2099–2114. https://doi.org/10.1194/jlr.r066514.

22.Salvadó, L., Palomer, X., Barroso, E., and Vázquez-Carrera, M. (2015). Targeting

endoplasmic reticulum stress in insulin resistance. Trends Endocrinol. Metab. *26*,

438–448. https://doi.org/10.1016/j.tem.2015.05.007.

23.Tumova, J., Andel, M., and Trnka, J. (2016). Excess of free fatty acids as a cause

of metabolic dysfunction in skeletal muscle. Physiol Res *65*, 193–207.

https://doi.org/10.33549/physiolres.932993.

24.Cani, P.D., Amar, J., Iglesias, M.A., Poggi, M., Knauf, C., Bastelica, D., Neyrinck,

A.M., Fava, F., Tuohy, K.M., Chabo, C., et al. (2007). Metabolic endotoxemia

initiates obesity and insulin resistance. Diabetes *56*, 1761–1772.

https://doi.org/10.2337/db06-1491.

25.Birdwell, L., Levesque, D., Machiah, D., and Gumber, S. (2022).

Clinicopathologic characteristics of pancreatic islet amyloidosis in the rhesus

macaque (Macaca mulatta) and sooty mangabey (Cercocebus atys). J. Med. Primatol.

51, 155–164. https://doi.org/10.1111/jmp.12579.

26.Westermark, P., Andersson, A., and Westermark, G.T. (2011). Islet amyloid

polypeptide, islet amyloid, and diabetes mellitus. Physiol. Rev. *91*, 795–826.

https://doi.org/10.1152/physrev.00042.2009.

27.Cefalu, W.T. (2006). Animal models of type 2 diabetes: clinical presentation and

pathophysiological relevance to the human condition. ILAR J *47*, 186–198.

https://doi.org/10.1093/ilar.47.3.186.

28.Ji, F., Jin, L., Zeng, X., Zhang, X., Zhang, Y., Sun, Y., Gao, L., He, H., Rao, J., Liu,

X., et al. (2012). Comparison of gene expression between naturally occurring and

diet-induced T2DM in cynomolgus monkeys. Dongwuxue Yanjiu *33*, 79–84. doi:

10.3724/SP.J.1141.2012.01079.

29.Tang, MT. (2020). Study on the Role of Glucose and Lipid in the Establishment of

Type 2 Diabetic Cynomolgus Monkey Model. M.S. Thesis, Dept. Veterinary Med.,

South China Agricultural Univ. *2020*.

30.Matsuda, M., DeFronzo, R.A. (1999). Insulin sensitivity indices obtained from oral

glucose tolerance testing: comparison with the euglycemic insulin clamp. Diabetes

care. *22*, 1462-1470. https://doi.org/10.2337/diacare.22.9.1462.

31.Lorenzo, C., Hazuda, H.P., Haffner, S.M. (2012). Insulin resistance and excess risk

of diabetes in Mexican-Americans: the San Antonio Heart Study. J. Clin. Endocr.

Metab. *97*, 793-799. https://doi.org/10.1210/jc.2011-2272.

32.Cowie, C.C., Rust, K.F., Byrd-Holt, D.D., Gregg, E.W., Ford, E.S., Geiss, L.S.,

Bainbridge, K.E., Fradkin, J.E. (2010) Prevalence of diabetes and high risk for

diabetes using A1C criteria in the US population in 1988–2006. Diabetes care. *33*,

562-568. https://doi.org/10.2337/dc09-1524.

33.Yu, W., Hao, X., Yang, F., Ma, J., Zhao, Y., Li, Y., Wang, J., Xu, H., Chen, L., Liu,

Q., et al. (2019). Hematological and biochemical parameters for Chinese rhesus

macaque. PLoS One. *14*, e0222338. https://doi.org/10.1371/journal.pone.0222338.

34.Du, Y., Gu, X., Meng, H., Aa, N., Liu, S., Peng, C., Ge, Y., and Yang, Z. (2018).

Muscone improves cardiac function in mice after myocardial infarction by alleviating

cardiac macrophage-mediated chronic inflammation through inhibition of NF-κB and

NLRP3 inflammasome. Am. J. Transl. Res. *10*, 4235–4246.

35.Alexeev, E.E., Lanis, J.M., Kao, D.J., Campbell, E.L., Kelly, C.J., Battista, K.D.,

Gerich, M.E., Jenkins, B.R., Walk, S.T., Kominsky, D.J., et al. (2018).

Microbiota-derived indole metabolites promote human and murine intestinal

homeostasis through regulation of interleukin-10 receptor. Am. J. Pathol. *188*,

1183–1194. https://doi.org/10.1016/j.ajpath.2018.01.011.

36.Manzella, C., Singhal, M., Alrefai, W.A., Saksena, S., Dudeja, P.K., and Gill, R.K.

(2018). Serotonin is an endogenous regulator of intestinal CYP1A1 via AhR. Sci. Rep.

8, 6103. https://doi.org/10.1038/s41598-018-24213-5.

37.Natividad, J.M., Agus, A., Planchais, J., Lamas, B., Jarry, A.C., Martin, R., Michel,

M., Chong-Nguyen, C., Roussel, R., Straube, M., et al. (2018). Impaired aryl

hydrocarbon receptor ligand production by the gut microbiota is a key factor in

metabolic syndrome. Cell Metab. *28*, 737-749.e4.

https://doi.org/10.1016/j.cmet.2018.07.001.

38.Rutkowsky, J.M., Knotts, T.A., Ono-Moore, K.D., McCoin, C.S., Huang, S.,

Schneider, D., Singh, S., Adams, S.H., and Hwang, D.H. (2014). Acylcarnitines

activate proinflammatory signaling pathways. Am. J. Physiol. Endocrinol. Metab. *306*,

E1378–E1387. https://doi.org/10.1152/ajpendo.00656.2013.

39.Tilg, H., and Moschen, A.R. (2008). Inflammatory mechanisms in the regulation of

insulin resistance. Mol. Med. *14*, 222–231. https://doi.org/10.2119/2007-00119.tilg.

40.Koves, T.R., Ussher, J.R., Noland, R.C., Slentz, D., Mosedale, M., Ilkayeva, O.,

Bain, J., Stevens, R., Dyck, J.R.B., Newgard, C.B., et al. (2008). Mitochondrial

overload and incomplete fatty acid oxidation contribute to skeletal muscle insulin

resistance. Cell Metab. *7*, 45–56. https://doi.org/10.1016/j.cmet.2007.10.013.

41.Lee, Y., Kang, E.S., Kim, S.H., Han, S.J., Kim, C.H., Kim, H.J., Ahn, C.W., Cha,

B.S., Nam, M., Nam, C.M., et al. (2008). Association between polymorphisms in

SLC30A8, HHEX, CDKN2A/B, IGF2BP2, FTO, WFS1, CDKAL1, KCNQ1 and type

2 diabetes in the Korean population. J. Hum. Genet. *53*, 991–998.

https://doi.org/10.1007/s10038-008-0341-8.

42.Su, S., Zhang, C., Zhang, F., Li, H., Yang, X., and Tang, X. (2016). The association

between leptin receptor gene polymorphisms and type 2 diabetes mellitus: A

systematic review and meta-analysis. Diabetes Res. Clin. Pract. *121*, 49–58.

https://doi.org/10.1016/j.diabres.2016.08.008.

43.Zhao, J. (2014). Molecular mechanisms of AGE/RAGE-mediated fibrosis in the

diabetic heart. World J Diabetes *5*, 860. https://doi.org/10.4239/wjd.v5.i6.860.

44.Nascimento, E.B., Osler, M.E., and Zierath, J.R. (2013). Sestrin 3 regulation in

type 2 diabetic patients and its influence on metabolism and differentiation in skeletal

muscle. Am. J. Physiol. Endocrinol. Metab. *305*, E1408–E1414.

https://doi.org/10.1152/ajpendo.00212.2013.

45.Tabassum, R., Mahajan, A., Dwivedi, O.P., Chauhan, G., Spurgeon, C.J., Kumar, M.V.K., Ghosh, S., Madhu, S.V., Mathur, S.K., Chandak, G.R., et al. (2012). Common variants of SLAMF1 and ITLN1 on 1q21 are associated with type 2 diabetes in Indian population. Am. J. Hum. Genet. *57*, 184–190. https://doi.org/10.1038/jhg.2011.150.

46.Junková, K., Mirchi, L.F., Chylíková, B., Janků, M., Šilhavý, J., Hüttl, M., Marková, I., Miklánková, D., Včelák, J., Malínská, H., et al. (2021). Hepatic transcriptome profiling reveals lack of Acsm3 expression in polydactylous rats with high-fat diet-induced hypertriglyceridemia and visceral fat accumulation. Nutrients *13*, 1462–1462. https://doi.org/10.3390/nu13051462.

47.Sekine, Y., Yamamoto, K., Kurata, M., Honda, A., Onishi, I., Kinowaki, Y., Kawade, G., Watabe, S., Nomura, S., Fukuda, S., et al. (2022). HADHB, a fatty acid beta-oxidation enzyme, is a potential prognostic predictor in malignant lymphoma. Pathology *54*, 286–293. https://doi.org/10.1016/j.pathol.2021.06.119.

48. Howson, J.M.M., Walker, N.M., Smyth, D.J., and Todd, J.A. (2009). Analysis of 19 genes for association with type I diabetes in the Type I Diabetes Genetics Consortium families. Genes & Immunity *10*, S74–S84. https://doi.org/10.1038/gene.2009.96.

49.Xu, L., Yin, L., Jin, Y., Qi, Y., Han, X., Xu, Y., Liu, K.X., Zhao, Y., and Peng, J. (2020). Effect and possible mechanisms of dioscin on ameliorating metabolic glycolipid metabolic disorder in type-2-diabetes. Phytomedicine *67*, 153139. https://doi.org/10.1016/j.phymed.2019.153139.

50.Liu, L., Li, Y., Guan, C., Li, K., Wang, C., Feng, R., and Sun, C. (2010). Free fatty acid metabolic profile and biomarkers of isolated post-challenge diabetes and type 2 diabetes mellitus based on GC–MS and multivariate statistical analysis. J. Chromatogr. B *878*, 2817–2825. https://doi.org/10.1016/j.jchromb.2010.08.035.

51.Dinarello, C.A., Donath, M.Y., and Mandrup-Poulsen, T. (2010). Role of IL-1β in

type 2 diabetes. Curr Opin Endocrinol Diabetes Obes *17*, 1.

https://doi.org/10.1097/med.0b013e32833bf6dc.

52.Sun, J., Su, Y., Chen, J., Qin, D., Xu, Y., Chu, H., Lu, T., Dong, J., Qin, L., and Li,

W. (2023). Differential roles of CD36 in regulating muscle insulin response depend on

palmitic acid load. Biomedicines *11*, 729.

https://doi.org/10.3390/biomedicines11030729.

53.Kawano, Y., Edwards, M., Huang, Y., Bilate, A.M., Araujo, L.P., Tanoue, T.,

Atarashi, K., Ladinsky, M.S., Reiner, S.L., Wang, H.H., et al. (2022). Microbiota

imbalance induced by dietary sugar disrupts immune-mediated protection from

metabolic syndrome. Cell *185*, 3501-3519.e20.

https://doi.org/10.1016/j.cell.2022.08.005.

54.Hezaveh, K., Shinde, R.S., Klötgen, A., Halaby, M.J., Lamorte, S., Ciudad, M.T.,

Quevedo, R., Neufeld, L., Liu, Z.Q., Jin, R., et al. (2022). Tryptophan-derived

microbial metabolites activate the aryl hydrocarbon receptor in tumor-associated

macrophages to suppress anti-tumor immunity. Immunity *55*, 324-340.e8.

https://doi.org/10.1016/j.immuni.2022.01.006.

55.Brawner, K.M., Yeramilli, V.A., Duck, L.W., Van Der Pol, W., Smythies, L.E.,

Morrow, C.D., Elson, C.O., and Martin, C.A. (2019). Depletion of dietary aryl

hydrocarbon receptor ligands alters microbiota composition and function. Sci. Rep. *9*,

14724. https://doi.org/10.1038/s41598-019-51194-w.

56.Turpin, W., Bedrani, L., Espin-Garcia, O., Xu, W., Silverberg, M.S., Smith, M.I.,

Garay, J.A.R., Lee, S.H., Guttman, D.S., Griffiths, A., et al. (2020). Associations of

NOD2 polymorphisms with Erysipelotrichaceae in stool of in healthy first degree

relatives of Crohn's disease subjects. BMC Medical Genet. *21*, 204.

https://doi.org/10.1186/s12881-020-01115-w.

57.Turnbaugh, P.J., Bäckhed, F., Fulton, L., and Gordon, J.I. (2008). Diet-induced

obesity is linked to marked but reversible alterations in the mouse distal gut

microbiome. Cell Host Microbe *3*, 213–223.

https://doi.org/10.1016/j.chom.2008.02.015.

58.Zhang, H., DiBaise, J.K., Zuccolo, A., Kudrna, D., Braidotti, M., Yu, Y.,

Parameswaran, P., Crowell, M.D., Wing, R., Rittmann, B.E., et al. (2009). Human gut

microbiota in obesity and after gastric bypass. Proc. Natl. Acad. Sci. U.S.A. *106*,

2365–2370. https://doi.org/10.1073/pnas.0812600106.

59.López☐Almela, I., Romaní☐Pérez, M., Bullich☐Vilarrubias, C., Benítez-PáezA.,

Gómez, E.M., Francés, R., Liebisch, G., and Sanz, Y. (2021). Bacteroides uniformis

combined with fiber amplifies metabolic and immune benefits in obese mice. Gut

Microbes *13*, 1–20. https://doi.org/10.1080/19490976.2020.1865706.

60.Kootte, R.S., Levin, E., Salojärvi, J., Smits, L.P., Hartstra, A.V., Udayappan, S.D.,

Hermes, G., Bouter, K.E., Koopen, A.M., Holst, J.J., et al. (2017). Improvement of

insulin sensitivity after lean donor feces in metabolic syndrome is driven by baseline

intestinal microbiota composition. Cell Metab. *26*, 611-619.e6.

https://doi.org/10.1016/j.cmet.2017.09.008.

61.Zhai, L., Xiao, H., Lin, C., Wong, H.L.X., Lam, Y.Y., Gong, M., Wu, G., Ning, Z.,

Huang, C., Zhang, Y., et al. (2023). Gut microbiota-derived tryptamine and

phenethylamine impair insulin sensitivity in metabolic syndrome and irritable bowel

syndrome. Nat. Commun. *14*, 4986. https://doi.org/10.1038/s41467-023-40552-y.

62.Paone, P., and Cani, P.D. (2020). Mucus barrier, mucins and gut microbiota: the

expected slimy partners? Gut *69*, 2232–2243.

https://doi.org/10.1136/gutjnl-2020-322260.

63.Png, C.W., Lindén, S.K., Gilshenan, K.S., Zoetendal, E.G., McSweeney, C.S., Sly,

L.I., McGuckin, M.A., and Florin, T.H.J. (2010). Mucolytic bacteria with increased

prevalence in IBD mucosa augment in vitro utilization of mucin by other bacteria. Am.

J. Gastroenterol. *105*, 2420–2428. https://doi.org/10.1038/ajg.2010.281.

64.Henke, M.T., Kenny, D.J., Cassilly, C.D., Vlamakis, H., Xavier, R.J., and Clardy, J.

(2019). Ruminococcus gnavus, a member of the human gut microbiome associated

with Crohn's disease, produces an inflammatory polysaccharide. Proc. Natl. Acad. Sci.

U.S.A. *116*, 12672–12677. https://doi.org/10.1073/pnas.1904099116.

65.Yin, J., Yang, K., Zhou, C., Xu, P., Xiao, W., and Yang, H. (2019). Aryl

hydrocarbon receptor activation alleviates dextran sodium sulfate-induced colitis

through enhancing the differentiation of goblet cells. Biochem. Biophys. Res.

Commun. *514*, 180–186. https://doi.org/10.1016/j.bbrc.2019.04.136.

66.Ma, N., Guo, P., Zhang, J., He, T., Kim, S.W., Zhang, G., and Ma, X. (2018).

Nutrients mediate intestinal bacteria–mucosal immune crosstalk. Front. Immunol. *9*, 5.

https://doi.org/10.3389/fimmu.2018.00005.

67.Oki, K., Toyama, M., Banno, T., Chonan, O., Benno, Y., and Watanabe, K. (2016).

Comprehensive analysis of the fecal microbiota of healthy Japanese adults reveals a

new bacterial lineage associated with a phenotype characterized by a high frequency

of bowel movements and a lean body type. BMC Microbiol *16*, 284.

https://doi.org/10.1186/s12866-016-0898-x.

68.Miguel, M.A., Lee, S.S., Mamuad, L.L., Choi, Y.J., Jeong, C.D., Son, A., Cho,

K.K., Kim, E.T., Kim, S.B., and Lee, S.S. (2019). Enhancing butyrate production,

ruminal fermentation and microbial population through supplementation with

Clostridium saccharobutylicum. J. Microbiol. Biotechnol. *29*, 1083–1095.

https://doi.org/10.4014/jmb.1905.05016.

69.Zhong, H., Wang, J., Abdullah, Hafeez, M.A., Guan, R., and Feng, F. (2021).

Lactobacillus plantarum ZJUFB2 Prevents High Fat Diet-Induced Insulin Resistance

in Association With Modulation of the Gut Microbiota. Front. Nutr. *8*, 75422.

https://doi.org/10.3389/fnut.2021.754222.

70.Takeuchi, T., Kubota, T., Nakanishi, Y., Tsugawa, H., Suda, W., Kwon, A.T.,

Yazaki, J., Ikeda, K., Nemoto, S., Mochizuki, T., et al. (2023). Gut microbial

carbohydrate metabolism contributes to insulin resistance. Nature. *621*, 389-395.

https://doi.org/10.1038/s41586-023-06466-x.

71.Takeuchi, T., Kameyama, K., Miyauchi, E., Nakanishi, Y., Kanaya, T., Fuji, T.,

Kato, T., Sasaki, T., Tachibana, N., Negishi, H., et al. (2023). Fatty acid

overproduction by gut commensal microbiota exacerbates obesity. Cell metab. *35*, 361-375. https://doi.org/10.1016/j.cmet.2022.12.013.

72.Magoc, T., and Salzberg, S.L. (2011). FLASH: fast length adjustment of short reads to improve genome assemblies. Bioinformatics *27*, 2957–2963. https://doi.org/10.1093/bioinformatics/btr507.

73.McDonald, D., Price, M.N., Goodrich, J., Nawrocki, E.P., DeSantis, T.Z., Probst, A., Andersen, G.L., Knight, R., and Hugenholtz, P. (2011). An improved Greengenes taxonomy with explicit ranks for ecological and evolutionary analyses of bacteria and archaea. ISME J *6*, 610–618. https://doi.org/10.1038/ismej.2011.139.

74.Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C.C., Al-Ghalith, G.A., Alexander, H., Alm, E.J., Arumugam, M., Asnicar, F., et al. (2019). Reproducible, interactive, scalable and extensible microbiome data science using QIIME 2. Nat. Biotechnol. *37*, 852–857. https://doi.org/10.1038/s41587-019-0209-9.

75.Segata, N., Izard, J., Waldron, L., Gevers, D., Miropolsky, L., Garrett, W.S., and Huttenhower, C. (2011). Metagenomic biomarker discovery and explanation. Genome Biol *12*, R60. https://doi.org/10.1186/gb-2011-12-6-r60.

76.Bolger, A.M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. Bioinformatics *30*, 2114–2120. https://doi.org/10.1093/bioinformatics/btu170.

77.Langmead, B., and Salzberg, S.L. (2012). Fast gapped-read alignment with Bowtie 2. Nat. Methods *9*, 357–359. https://doi.org/10.1038/nmeth.1923.

78.Wood, D.E., and Salzberg, S.L. (2014). Kraken: ultrafast metagenomic sequence classification using exact alignments. Genome Biol. *15*, R46. https://doi.org/10.1186/gb-2014-15-3-r46.

79.Li, D., Liu, C.M., Luo, R., Sadakane, K., and Lam, T.W. (2015). MEGAHIT: an ultra-fast single-node solution for large and complex metagenomics assembly via succinct de Bruijn graph. Bioinformatics *31*, 1674–1676. https://doi.org/10.1093/bioinformatics/btv033.

80.82.Hyatt, D., Chen, G.L., LoCascio, P.F., Land, M.L., Larimer, F.W., and Hauser,

L.J. (2010). Prodigal: prokaryotic gene recognition and translation initiation site

identification. BMC Bioinform. *11*, 119. https://doi.org/10.1186/1471-2105-11-119.

81.Fu, L., Niu, B., Zhu, Z., Wu, S., and Li, W. (2012). CD-HIT: accelerated for

clustering the next-generation sequencing data. Bioinformatics *28*, 3150–3152.

https://doi.org/10.1093/bioinformatics/bts565.

82.Patro, R., Duggal, G., Love, M.I., Irizarry, R.A., and Kingsford, C. (2017). Salmon

provides fast and bias-aware quantification of transcript expression. Nat. Methods *14*,

417–419. https://doi.org/10.1038/nmeth.4197.

83.Buchfink, B., Xie, C., and Huson, D.H. (2014). Fast and sensitive protein

alignment using DIAMOND. Nat. Methods *12*, 59–60.

https://doi.org/10.1038/nmeth.3176.

84.Lombard, V., Golaconda Ramulu, H., Drula, E., Coutinho, P.M., and Henrissat, B.

(2013). The carbohydrate-active enzymes database (CAZy) in 2013. Nucleic Acids

Res. *42*, D490–D495. https://doi.org/10.1093/nar/gkt1178.

85.Franzosa, E.A., McIver, L.J., Rahnavard, G., Thompson, L.R., Schirmer, M.,

Weingart, G., Lipson, K.S., Knight, R., Caporaso, J.G., Segata, N., et al. (2018).

Species-level functional profiling of metagenomes and metatranscriptomes. Nat.

Methods *15*, 962–968. https://doi.org/10.1038/s41592-018-0176-y.

86.Patel, R.K., and Jain, M. (2012). NGS QC Toolkit: a toolkit for quality control of

next generation sequencing data. PLoS One *7*, e30619.

https://doi.org/10.1371/journal.pone.0030619.

87.Pertea, M., Pertea, G.M., Antonescu, C.M., Chang, T.C., Mendell, J.T., and

Salzberg, S.L. (2015). StringTie enables improved reconstruction of a transcriptome

from RNA-seq reads. Nat. Biotechnol. *33*, 290–295. https://doi.org/10.1038/nbt.3122.

88.Love, M.I., Huber, W., and Anders, S. (2014). Moderated estimation of fold change

and dispersion for RNA-seq data with DESeq2. Genome Biol. *15*.

https://doi.org/10.1186/s13059-014-0550-8.

89.Langfelder, P., and Horvath, S. (2008). WGCNA: an R package for weighted

correlation network analysis. BMC Bioinform *9*, 559.

https://doi.org/10.1186/1471-2105-9-559.

90.Yu, C., Woo, H.J., Yu, X., Oyama, T., Wallqvist, A., and Reifman, J. (2017). A

strategy for evaluating pathway analysis methods. BMC Bioinform *18*.

https://doi.org/10.1186/s12859-017-1866-7.

91.Raudvere, U., Kolberg, L., Kuzmin, I., Arak, T., Adler, P., Peterson, H., and Vilo, J.

(2019). g:Profiler: a web server for functional enrichment analysis and conversions of

gene lists (2019 update). Nucleic Acids Res. *47*, W191–W198.

https://doi.org/10.1093/nar/gkz369.

92.Berland, M., Cadiou, J., Levenez, F., Galleron, N., Quinquis, B., Thirion, F.,

Gauthier, F., Le Chatelier, E., Plaza Oñate, F., Schwintner, C., et al. (2021). High

engraftment capacity of frozen ready-to-use human fecal microbiota transplants

assessed in germ-free mice. Sci. Rep *11*, 4365.

https://doi.org/10.1038/s41598-021-83638-7.

A OPLS-DA

B Volcano Plot

D KEGG enrichment

C

E metaMDS: Stress = 0.106
P-value = 0.019

F Volcano Plot

G Functional enrichment

H Module−trait relationships

I

Metabolite1(2s,3r,4s,5s,6r)-2-[(2r,3r,4s,5s,6r)-2-[[(3s,6s,8r,10r,12r,13r,14r,17s)-3,12-dihydroxy-4,4,8,10,14-pentamethyl-17-[(2s)-6-methyl-2-[(2s,3r,4s,5s,6r)-3,4,5-trihydroxy-6-(hydroxymethyl)oxan-2-yl]oxyhept-5-en-2-yl]-2,3,5,6,7,9,11,12,13,15,16,17-dodecahydro-1h-cyclopenta[a]phenanthren-6-yl]oxy]-4,5-dihydroxy-6-(hydroxymethyl)oxan-3-yl]oxy-6-(hydroxymethyl)oxane-3,4,5-triol