Agent-Based Simulation of Cortical Microtubule Band Movement in the *Arabidopsis Zygote*

Short title:

Simulation of Microtubule Band Movement

Authors: Tomonobu Nonoyama[1], Zichen Kang[1], Hikari Matsumoto[2], Sakumi Nakagawa[2], Minako Ueda[2], Satoru Tsugawa[1]

1 Department of Mechanical Engineering, Faculty of Systems Science and Technology, Akita Prefectural University, Yurihonjo, Akita, 015-0055, Japan

2 Graduate School of Life Sciences, Tohoku University, Sendai, Miyagi, 980-8578, Japan

*Corresponding author: tsugawa@akita-pu.ac.jp

Keywords: Agent-based model, Arabidopsis zygote, Cortical microtubules, Self-organization, Orientational order

Abstract

Arabidopsis thaliana zygotes form a circumferential cortical microtubule (CMT) band that stays at a constant position from the growing tip. CMT bands support zygote anisotropic growth that is essential to determine the apical-basal axis of the plant body, but how stable CMT bands are maintained remains unclear. We hypothesized that cell surface tension induces CMT reorientation of which the strength was defined as the directional cue (DC). We constructed an agent-based simulation model using the DC and its spatial distribution (DC zone) to determine the range of molecular parameters and the width and movement speed of the DC zone required to form the CMT band. We then developed a data–model correspondence method to determine the range of the parameters which are difficult to observe experimentally. We determined that the DC zone must move with an appropriate speed of mechanical change may exist to form a moving CMT band.

Introduction

In the *Arabidopsis thaliana*, the anisotropic growth of the zygote, the single cell formed by the fusion of the egg and sperm cells in plant sexual reproduction, determines the longitudinal (apical–basal) axis of the resulting plant body because the zygotes undergo unequal division after the anisotropic growth to generate a dome-shaped apical cell that leads to the main body of the plant (Tanaka et al*.,* 2024). During zygote elongation, a cortical microtubule (CMT)

band is formed in the circumferential orientation near the subapical region of the cell guiding the future formed first unequal division plane (Kimata et al., 2016; Kang et al., 2024). Given that the CMT band is maintained within a specific subapical region during cell growth, the CMTs above the band may continuously form while the CMTs below the band are degraded to produce a macroscopic movement of the CMT band. Since the cell shape becomes more isotropic without CMT band with the CMT polymerization inhibitors and the CMT band becomes disappeared in the *yoda* mutant with growth arrest prematurely, the interactive feedback between cell growth and CMT organization must be bidirectional (Kang et al., 2024). Root hairs and pollen tubes also exhibit tip growth, but unlike the zygote, their microtubules are aligned in the longitudinal direction (Braun et al., 1999; Cheung et al., 2008). The elongation rate of root hairs is 18 μm/h, more than 10 times that of the zygote (Kang et al., 2023), which might explain why root hairs do not form a circumferential CMT band. These results suggest that the CMT band must move with an appropriate speed consistent with the cell growth, though the mechanisms involved remain unclear.

Many studies have focused on the mechanical feedback from (to) the microtubule to (from) mechanical stress, as introduced below. One pioneering study explored the similarity between the stress distribution derived from the cell geometry and the CMT orientation (Hamant et al., 2008). This was followed by the experimental finding that CMT-severing proteins cut the CMTs perpendicular to the stress orientation (Uyttewaal et al., 2012), and by many reports of the similarity between stress profiles on the cell surface and the distribution of the CMT orientations (Sampathkumar et al., 2014; Hervieux et al., 2017; Sapala et al., 2018). Among them, one important finding was that CMT increased polymerization in regions of increased tension (Inoue et al., 2017; Hamant et al., 2019), suggesting that CMTs may respond to tensile intensity. More interestingly, the property of default orientation in the absence of tension (Durand-Smet et al., 2020) and a hypothesis that the default formation can be overwritten by tensional cues (Colin et al., 2020) were proposed. The overriding of the CMTs by tension defaults back to normal after 10 h, strongly suggesting that CMTs respond not only to the tension intensity itself but also to the changes or fluctuation in tension (Colin et al., 2021; Moulia et al., 2021). In the present study, we therefore hypothesize that the cell surface tension induces the reorientation of the direction in which CMTs are newly formed. The strength of this reorientation is defined as the directional cue (DC), and its spatial distribution as the DC zone. In previous work, the systematic change of DC induced by cell surface tension changed the default CMT orientation (Mirabet et al., 2018); however, the width and the movement speed of the DC zone were not systematically investigated.

Cellulose synthase enzyme complexes move preferentially on the CMTs (Heath, 1974; Paradez et al., 2006), meaning that the CMTs guide the deposition of cellulose microfibrils and

thus cell growth anisotropy (Baluska et al., 2003; Lloyd & Chan, 2008). CMT dynamics are therefore essentially linked to cell or tissue morphogenesis. The growth speed of the CMTs is faster than their shrinkage speed, so the polymers can grow and thus move in an oriented direction, which is known as treadmilling (Mitchison & Kirschner, 1984; Shaw & Kamyar, 2003). The CMTs interact with each other by "zipping" together, inducing depolymerization (called "induced catastrophe"), or "crossover" one another, depending on their interaction angle (Dixit & Cyr, 2004a; 2004b. Based on these phenomenological observations, there were two main ways to model CMT self-organization (Dumond, 2012; Shi & Ma, 2010). Particle-based simulations assumed that all the CMTs act as agents that can grow or shrink while simultaneously interacting with each other (Tindemans et al., 2010; 2014; Chakrabortty et al., 2018; 2018-2), whereas the probability-based approach involved the computation of the density of growing CMTs (Hawkins et al., 2010). Both of these models revealed a qualitatively similar phase diagram (ordered or disordered states as functions of the model parameters of the CMTs) (Shi & Ma, 2010).

These pioneering studies cannot be directly applied to the single zygote, however. First, the studies were based on closed systems in which the CMTs are packed into a restricted zone, which is not applicable to the elongating zygote because it is an open-ended system due to cell growth. Second, the studies did not take the tensional effect into account, with the exception of the recent work by Mirabet et al. (2018). To address the first challenge, we aimed to determine the effect of various molecular parameters for the fixed DC zone on the open-ended cylinder. To overcome the second issue, we reconstructed an agent-based model of the CMTs with systematic changes in the width and the movement speed of the DC zone to assess the appropriateness of the DC movement, i.e., its ability to form a moving CMT band. Another important challenge was the correspondence between data and model, as the data only show the width and the movement speed of the CMT band while the simulation models often contain too many parameters at the molecular scale. A new research framework is therefore required to extract effective parameters that match the actual data.

In this study, we first defined a CMT dynamic space (CDS) with the width and the movement speed of the moving CMT band from actual data to ensure correspondence between data and model. We then reconstructed an agent-based model of CMTs with DCs derived from the tension of the cylindrical cell surface, with the aim of extracting the model parameter range that best explains the width and the movement speed of the CMT band. We tuned the strength of the tension-responsive ability of the CMTs and found that this parameter affects both the width and the orientational order of the resulting CMT band. Next, we analyzed the effects of the induced-catastrophe rate, the width of the DC zone, and the cell radius. Finally, we systematically modified the movement speed of the DC zone and found that the width and

movement speed of the tension band should be constrained to be consistent with the observed

data; otherwise, the CMT band falls behind the input speed. Our simulation suggests that an

appropriate speed of mechanical change may exist to form a moving CMT band for building up

the apical-basal body axis in the Arabidopsis zygote, which is not present in elongating cells

without a CMT band, such as pollen tubes and root hairs.

Methods

To understand the CMT dynamics beneath the membrane of the zygote, we

constructed an agent-based simulation, as described below. An analysis of the actual zygote data,

introduced later, revealed that the width and movement speed of the CMT band have

characteristic values at the zygote elongation stage with cylindrical cell geometry; thus, a

cylindrical cell surface was selected for use in this model. The agent-based model incorporated

the growth, shrinkage, and interaction events between agents (CMTs), the latter of which

included zippering, crossovers, and induced catastrophe, as detailed below. The DC derived

from tension and the finite tubulin-pool effect were both taken into consideration.

Agent-based model of CMTs on the cylindrical cell surface

The agent-based model of the CMTs was employed based on previous works

(Mirabert et al., 2018; Chakrabortty et al., 2018), as shown in Fig. 1. There are six crucial

parameters of CMT dynamics: (1) the growth speed of the plus end, v^+ (Fig. 1A); (2) the

shrinkage speed of the plus end, v^- (Fig. 1B); (3) the shrinkage speed of the minus end

(treadmilling speed), v^{tm}; (4) the switching rate from shrinkage to growth at the plus end

(rescue rate), r_r; (5) the switching rate from shrinkage to growth at the plus end (spontaneous

catastrophe rate), r_c; and (6) the nucleation rate of CMTs on the cylindrical surface, r_n. The

parameter values are summarized in Table 1. In most simulations, cylindrical cell geometry was

used with a radius R and height H. The simulation was iterated N times with the time

interval Δt for each step.

Interactions of growing CMTs

In addition to the dynamics of a single CMT, described above, the interactions

between growing CMTs were also implemented in the simulation (Tindemans et al., 2010, 2014;

Chakrabortty et al., 2018, 2018-2; Hawkins et al., 2010; Shi & Ma, 2010). Three interactions

were considered: (1) "zippering", when the growing CMT changes the direction of its growing

front to be parallel to an existing CMT without changing the direction of the existing CMT,

which happens when a growing CMT collides with an existing CMT at an angle smaller than

40° (Fig. 1C); (2) "induced catastrophe", when a CMT begins to depolymerize from the

growing front when it meets an existing CMT at an angle greater than 40° (Fig. 1D); and (3)

"crossover", when a growing CMT crosses an existing CMT at an angle greater than 40° (Fig.

1E). Induced catastrophes and crossovers are both equally probable (50% probability) when a

growing CMT collides with an existing CMT at an angle greater than 40°.

Time and space units

The typical spatial scale of a zygote is 5 μm in radius (Kang et al., 2023). The unit of

length of the CMT segment was considered to be 8 nm, the reported width of a ring of tubulin

(Mirabet et al., 2018). The previously reported speed of growth at the plus end was 4.8 μm/min

(Tindemans et al., 2010), and the simulation time step was approximately 0.2 s. Most of the

simulation iterations comprise 12,000 time steps, corresponding to about 40 min.

DC derived from tension

Owing to the growing and shrinking of the CMT plus and minus ends, the length of

the CMT agents can be changed, whereas the orientation of the CMTs does not change by

default. With this setup, the DC zone (as shown in brown in Fig. 1C) was defined as the region

where the newly adding vector is biased in a horizontal direction with weight b_d:

$$\boldsymbol{r_{n+1}} = \frac{(1 - b_d)\boldsymbol{r_n} + b_d\boldsymbol{u}}{\|(1 - b_{d)}\boldsymbol{r_n} + b_d\boldsymbol{u}\|} \#(1)$$

where $\boldsymbol{r_n}$ and $\boldsymbol{r_{n+1}}$ are the current and next unit vector of the CMT plus end, respectively, $\boldsymbol{u}$

is a circumferential unit vector, and $\|\cdot\|$ is the standard norm. The calculation of the unit vector

was almost the same as in a previous study (Mirabet et al., 2018), except that random

fluctuations in the direction of the CMT growth were not considered. The notations of position

μ_T and width σ_T of the DC zone were used. The direction of the CMT plus end depends on the

position of the plus end, as follows:

(i) The microtubule (MT) plus ends located outside of the DC zone do not change the direction.

(ii) The MT plus ends located within the DC zone change the direction toward the horizontal

direction.

Finite tubulin-pool effect

In order to observe the CMT dynamics effectively, a finite tubulin pool proposed by

Chakrabortty et al. (2018) was used. The pool limits the CMT length by L_{max}, such that the

speed of a growing CMT plus end at any time t is dependent on the total length $L(t)$ of all

CMTs in the system:

$$v_{tub}^+(t) = v^+\left(1 - \frac{L(t)}{L_{max}}\right)\#(2)$$

177 where $L_{max} = \rho_{tub}A$ is the length of all the existing CMTs on the surface area A and finite
178 tubulin density ρ_{tub}, and v^+ is the CMT plus-end growth speed without taking the pool into
179 consideration.
180

Quantification of the width of the CMT band in the data

Based on the work of Kang (2023), the coordinate normalization method was used to
fix the zygote position. With this normalization, the cell centerline analyzed in the previous
studies (Bastien et al., 2016; Kang et al., 2024) was extracted and the fluorescence intensity of
the CMTs within the cell was projected to the centerline. The fluorescence intensity was fitted
with the truncated Gaussian distribution as a function of position z on the centerline using the
parameters $(\alpha, \beta, \mu, s, b)$ as follows:

$$P(S) = \frac{1}{\sigma} \frac{\varphi\left(\frac{S-\mu}{s}\right)}{\Phi\left(\frac{\beta-\mu}{s}\right) - \Phi\left(\frac{\alpha-\mu}{s}\right)} + b \#(3)$$

188 where S is the curvilinear coordinate along the cell centerline, and the Gaussian probability density
189 function is $\varphi(S) = 1/\sqrt{2\pi} \exp(-1/2 S^2)$ with an average μ and standard deviation s. The
190 cumulative distribution function is $\Phi(S) = 1/2\left(1 + erf\left(S/\sqrt{2}\right)\right)$. The width of the CMT band
191 is defined as $\sigma_{MT}^{data} = 2s$. Furthermore, the CMT position is defined as $\mu_{MT}^{data} = \mu$.
192

Quantification of the width of the CMT band in the model

The width of the CMT band in the simulations was calculated by quantifying the
cumulative probability function P of the CMT segments larger than the threshold 3.85 μm (1/8
of the circumference); this threshold can omit unorganized CMTs. $P(S)$ was then fitted with a
sigmoid curve using five points around $S = 0.5$ (Fig. S1, gray dashed line). The width of the
CMT band was estimated by the 68th percentile of the corresponding probability density
function for the fitted sigmoidal cumulative distribution (Fig. S1).
200

Order parameter tensor quantifying the orientational order of the CMTs

202 To measure the degree of the orientational order of the CMT array in the simulations, a
203 local order parameter tensor $\boldsymbol{q}$ and a global order parameter tensor $\boldsymbol{Q}$ were introduced
204 (Chakrabortty et al., 2018). Using the orientation of the CMT segment at the position x,
205 $\hat{\omega} = \cos\alpha \, \hat{e}_x(x) + \sin\alpha \, \hat{e}_y(x)$ where $\alpha \in [0, 2\pi)$ is the declination angle between $\hat{\omega}$ and $\hat{e}_x$.
206 The local order parameter tensor $\boldsymbol{q}$ at the local area dA is defined as

$$q_{ij} = 2\langle\widehat{\omega_i}\widehat{\omega_j}\rangle - \delta_{ij}, \#(4)$$

207 with the matrix form

$$q = \begin{bmatrix} (2\cos^2\alpha - 1)e_x \otimes e_x & 2\cos\alpha \sin\alpha \, e_x \otimes e_y \\ 2\cos\alpha \sin\alpha \, e_y \otimes e_x & (2\sin^2\alpha - 1)e_y \otimes e_y \end{bmatrix}. \#(5)$$

208   Note that $<\cdot>$ means the sample average of $\cdot$ within the local area dA. The local

209   orientational order and the local orientation of the CMT segments within dA are estimated by

210   the first eigenvalue of q and by the orientation of the corresponding eigenvector of q,

211   respectively. For the characterization of the global orientation, the global order parameter tensor

212   was defined as

$$Q = \frac{\int dA(x)\rho(x)q(x)}{\int dA(x)\rho(x)} \#(6)$$

where $\rho(x)$ is the local areal density of the CMT segments within dA. The global orientational

order and the global orientation of all the CMTs within the whole region are estimated by the

first eigenvalue denoted by Λ of Q and by the orientation of the corresponding eigenvector of

Q, respectively. Λ becomes 0 for completely random CMT orientations, while for

well-organized CMTs, Λ becomes +1 for the horizontal array and −1 for the vertical array. The

number of local regions is n_{div} in this study, which divides the whole region into the parts with

height H/n_{div}.

Results

The CMT band moves upwards with a speed of 1.5 ± 1.2 μm/min and a width of 5.2 ± 2.4

μm

To determine the mechanism by which the CMT band moves to maintain its position

as the zygote expands, we first quantified the width and movement speed of the CMT band

using actual observation data (Fig. 2A and 2B). At the CMT-building stage, which occurs just

after fertilization, the CMTs would not be well organized yet; subsequently, the cell becomes

protruded in the apical direction with CMT band near the cell tip (Kimata et al*.,* 2016). The cell

maintains anisotropic elongation during the elongation stage, during which the organized CMT

band is localized about 5~7μm below the growing tip. To elucidate how the CMT band is

maintained, we focused on the elongating stage assuming a cylindrical cell geometry.

To standardize the dynamics of the CMT bands of different individuals to an

appropriate time scale, we used the characteristic time t_{RGS} at the rapid growth stage (RGS),

during which the cell length rapidly increased (Kang et al*.,* 2023). We defined the stage from t_0

(5 h before t_{RGS}) to t_{RGS} as the elongating stage. The CMT fluorescence intensity projected on

the cell centerline (Fig. 2C and 2D) was fitted by the truncated Gaussian distribution with mean

μ_{MT}^{data} and standard deviation σ_{MT}^{data} (Fig. 2E and 2F; see also Methods). The kymograph of the

CMT fluorescence intensity revealed that the CMT band is actually localized to the subapical

region, maintaining almost the exact same distance to the cell tip as the zygote elongates (Fig.

2G). The width of the CMT band also remained constant at approximately 5.2 ± 2.4 μm during

elongation. The movement speed of the CMT band was less than 2.7 μm/h (Fig. 2H).

Tension-responsive ability parameter b_d affects both the width and orientational order of

the CMT band

245   To establish correspondence between the observed data and the model parameters, we

introduced a CMT dynamic space (CDS) describing the speed $d\mu_{MT}^{data}/dt$ and the width σ_{MT}^{data}

of the CMT band (Fig. 2H). A total of 11 parameters were included in the model (Fig. 1): six

CMT dynamics parameters v^+, v^-, v^{tm}, r_r, r_c, and r_n; three mechanical parameters b_d, μ_T,

and σ_T; and two geometrical parameters R and H (Table 1). Based on previous studies and

the observed data (Chakrabortty et al., 2018-2, Shaw et al., 2003, Eren et al., 2010, Tindemans

et al., 2014-2), we set six parameters (v^+, v^-, v^{tm}, r_r, H, and r_n) as fixed and observed the

effects of changing the remaining five parameters (r_c, b_d, μ_T, σ_T, and R). We used the CDS to

make correspondence with these five parameters.

254   To observe the effects of parameter perturbations, it is necessary to identify the

255   parameters that can replicate a similar CMT band with sample-averaged data. We initially fixed

256   the parameter μ_T at the specific location $z = 10$ μm within the range of $z \in [0, 20]$ (Fig.

257   2H) and used specific parameters ($r_c = 0.001$, $b_d = 0.01$, $\sigma_T = 6.67$ [μm], and $R = 5$ [μm]).

258   Using this configuration, we examined the changes in the resulting CMT band by perturbing the

259   parameters (r_c, b_d, σ_T, and R).

260   To avoid misunderstanding, we defined the terms "single CMT formation" as a single

CMT oriented in the circumferential direction (a single ring of CMT) and "CMT band formation"

as a band formed in the circumferential direction (a bundled structure of CMTs). We noticed that

the CMT band can be formed for the parameter $b_d = 0$, but it was not specifically localized

(Fig. 3A). Next, we set $b_d = 0.001$ for the case of the weak DC, resulting in a CMT band

partially localized near the DC zone (dashed line) (Fig. 3B). When $b_d = 0.01$, for the case of

the strong DC, the formed CMT band covered the whole region of the DC zone (Fig. 3C). To

check the time scale of the reorientation during the single CMT formation, we confirmed that, in

the cases $b_d < 0.01$, the converging time to the horizontal orientation of the CMTs becomes

more than the time scale of single CMT formation roughly estimated as $2\pi R/v_+ \sim 6.5$ min for

$R = 5$ μm (Fig. 3D), therefore we set the case $b_d = 0.01$ as strong DC where the converging

time is fast enough than the single CMT formation.

272   To quantitatively describe the degree of CMT band formation, we evaluated the width

of the CMT band σ_{MT}^{model}, and the global order parameter Λ, measuring both the orientation

and density of the CMTs (see Methods). The order parameter Λ becomes 0 for the completely

random orientation of the CMTs and 1 for well-organized CMTs. As a result, we quantitatively

confirmed that the width σ_{MT}^{model} decreases as a function of b_d, consistent with the above

qualitative observations (Fig. 3F). Furthermore, we quantitatively confirmed that the order of

CMT organization increases as b_d increases, which was also consistent as well (Fig. 3G).

Catastrophe rate induces a phase transition for both strong and weak DC

Next, to confirm the robustness of the CMT band against CMT depolymerization, we

systematically increased the catastrophe rate. Since depolymerization shortens CMTs, it is

expected to decrease the probability of CMT collisions that may inhibit the formation of CMT

bands. Because the DC may compensate for the increased catastrophe rate by facilitating band

formation, we tested the changes for the weak and strong DC conditions. In the case of weak

DC, the CMT organization at $t = 20$ min becomes sparse as the catastrophe rate increases

because the induced catastrophe frequently cuts the CMTs (Fig. S2A). Moreover, the

quantitative evaluation of Λ revealed that the degree of orientational order was less than 0.75,

meaning that the CMTs are not strongly organized (Fig. S2B). As expected, our definition of the

width of the CMT band suddenly dropped to small values at $r_c = 0.002$, where the phase

transition in terms of CMT band width occurs (Fig. S2C). This sudden decrease in the width of

the CMT band occurs even in the case of the strong DC, where the horizontally aligned single

CMT formation was cut by the catastrophe (Fig. S2D). In this case, the index Λ was greater

than 0.75, meaning that the CMTs are strongly organized (Fig. S2E) and a similar phase

transition occurs (Fig. S2F).

The CMT band widens as the input DC widens under both weak and strong DC

To exlpore whether the width of the output CMT band σ_{MT}^{model} corresponds to the

width of the input, we focused on the perturbation of the input DC width σ_T. In the case of

weak DC, the CMTs at $t = 20$ min were widely distributed beyond the input DC zone (Fig.

3G). In the case of strong DC, the CMTs at $t = 20$ min were distributed with almost the same

width as that of the input DC zone (Fig. 3H). We quantitatively confirmed that these results

were preserved independent of the parameter σ_T (Fig. 3I).

Cylinder radius does not strongly affect the CMT band but emphasizes two different types

of CMT organization

Adimittingthat changes in the cylinder radius alter the circumference and surface area

of the cylinder, which is expected to increase the time required for CMT to form bands, we

tested the cases of $R = 5.0$ μm (typical zygote scale reported by Kang et al. (2023)) and $R = 15$

μm. In addition, we tested the weak and strong DC conditions because the DC may compensate

for a delay in CMT band formation. In the case of strong DC, the CMT band was rapidly

organized, and this behavior was not strongly affected by the radius (Fig. 4A); therefore, the
index Λ and the CMT band width σ_{MT} showed a similar trend (Fig. 4A and 4B). On the other
hand, in the case of weak DC, the CMTs were broadly distributed across the entire region, and
they gradually organized within the DC zone (Fig. 4D and 4E). We call this phenomenon a
gradual organization of CMTs and distinguish this from the rapid organization. As a result, the
index Λ values are lower than those in the case of strong DC, and the CMT band width σ_{MT}
with $R = 15$ μm gradually decreases, reflecting the gradual organization (Fig. 4E). The
mechanism behind this might be the different frequency of the CMT encounters. In the case of
strong DC, the horizontal CMTs are rapidly organized, preventing the diagonal organization of
the CMT band (Fig. 4C). In the case of weak DC, the horizontal CMTs are not well organized,
allowing the diagonal or spiral organization of the CMT band (Fig. 4F).
323
324

The moving CMT band can be formed when the DC zone displays an appropriate
movement speed and width
As the results above suggested that a desired CMT band can be reproduced with
$r_c \leq 0.002$, $b_d \geq 0.01$, σ_T slightly smaller than the desired band, and $R = 5.0$,
corresponding to the observed data, we are now able to move the designated CMT band by
changing the specific parameter $d\mu_T/dt$ (the movement speed of the DC zone). As
schematically illustrated in Fig. 5A, we searched for the best fitted parameter set ($d\mu_T/dt$, σ_T)
from the CDS in Fig. 2H using an exhaustive search of the model parameters. As the expected
DC parameters were unknown, we distributed the parameter set with $d\mu_T/dt \in [0, 4.0]$ μm/h
and $\sigma_T \in [0.5, 3.5]$ μm as a similar range to the data. As a result, σ_{MT}^{model} becomes wider than
σ_T while unexpectedly the resulting $d\mu_{MT}^{model}/dt$ becomes smaller or sometimes even a
negative value, which is below the expected movement speed of the CMT band (Fig. 5B, D, F,
H). To demonstrate what happens, we plotted the simulated CMTs in Fig. 5C. (i) shows CMT
behaviors with appropriate width and movement speed of the DC zone (Fig. 5D and Fig. 5E).
(ii) shows the behavior with slower DC movement (Fig. 5F, G). (iii) shows the behavior with
faster DC movement and a wider DC (Fig. 5H). These results suggest that when the DC zone
moves faster than the CMT growing speed, the organized CMT band is left behind the DC zone
(Fig. 5H and 5I).
In summary, we conclude that the moving CMT band could be achieved with an
appropriate width and movement speed of the DC zone.
345

346 **Discussion**
347 In this study, we developed an agent-based simulation based on the CMT

tension-responsive hypothesis to identify the parameters necessary for the formation of the moving CMT band, which is specifically formed in the plant zygote. Through a CDS analysis, we successfully compared the CMT bands between the model and the observed data to estimate the necessary parameters, and obtained three new main findings regarding the reproduction of the moving CMT band in *Arabidopsis* zygotes. (1) The width of the CMT band was about 5.2 ± 2.4 μm, and the movement speed was about 1.3 ± 1.5 μm/h. (2) There can be two types of CMT band formation: rapid organization and gradual organization. (3) The formation of the moving CMT band requires an appropriate width and movement speed in the input mechanical parameters too.

In addition to the CMT tension-responsive hypothesis, the organization of the CMT band has been hypothesized to involve other mechanisms. The edge catastrophe hypothesis, where the cell edge induces CMT catastrophe, was proposed to explain CMT organization in early embryonic apical cells in previous studies (Chakrabortty et al., 2018; Chakrabortty et al., 2018-2); however, this hypothesis seems unlikely to hold for the zygote, especially in the elongating term, when cell edges are lacking around the CMT band. The conceptual basis for the CMT tension-responsive hypothesis is an increase in the rate of CMT polymerization in the region of higher tension, based on previous studies (Inoue et al., 2016; Hamant et al., 2019), but subsequent experimental research on CMTs in protoplasts (Colin, 2021) has shown that tension changes over time may increase the rate of CMT polymerization. Thus, the driving force might be a time derivative of the tension (Kang et al., 2024). Additionally, we admit that the possibility that CMTs are severed in a direction perpendicular to the tension exerted by katanin outside the DC zone (Uyttewaal et al., 2012) is not ruled out.

For the case of the moving DC zone, in order to be consistent with the parameter ranges of CMT band in the actual observed data, the movement speed of the DC zone must be comparable to that of cell elongation (1.5 μm/h on average). A similar mechanism for a moving ordered structure may exist for the formation of CMT band structure in the tip growth mode in fern protonema, where the elongation rate is small (Murata & Wada, 1989). The elongation rate of root hairs is 18 μm/h, which is more than 10 times that of the zygote (Kang et al., 2023), so it is thought that they are unable to form CMT bands, similar to our numerical result. This is the "left-behind" phenomenon, in which the mechanical (tensional) change during elongation becomes so fast that CMT organization does not catch up with the input mechanical perturbation. As the elongation speed of the pollen tubes is 100–1000 times that of the zygote (Bedinger, 1992; Schiøtt et al., 2004), the reason that they do not have CMT bands is thought to be similar to the left-behind phenomenon. The question of when and how the organisms evolved a moving CMT band should be of great significance, not only for cell and molecular biology, but also for understanding the evolution and diversity of plant cells.

In this study, we developed a data–model correspondence method called CDS to
narrow down the range of molecular parameters such as the tension-responsive ability and
catastrophe rate of the CMTs, which are difficult to observe experimentally. This method can be
applied not only to other cylindrical plant cells (Hasezawa et al*., 1997), but also to animal cells
in general, such as the orientational order during the formation of the extracellular matrix of
animal cells (for example, in the *Drosophila* trachea; Öztürk-Çolak et al*., 2016). The
interdisciplinary research that links biological images and simulation models can thus provide
new opportunities to elucidate and understand the essential dynamics of plant and animal
cytoskeletons.

Acknowledgments:
The authors thank Koichi Fujimoto, Katsuyoshi Matsushita and Naoya Kamamoto
(Hiroshima Univerisity), Takumi Higaki, and Haruka Ono (Kumamoto University), Ishimoto
Yukitaka (Saga University) and Yusuke Kimata (Tohoku University) for helpful discussions.

Author Contributions:
T.N. and S.T. conceived and designed the study. T.N. and S.T. developed and
implemented models and algorithms. H.M., S.N and M.U. carried out data acquisition from live
cell imaging. T.N. and S.T. wrote the manuscript. T.N., Z.K., S.T., and M.U. edited and
reviewed the manuscript. S.T., H.M., and M.U. provided research funding. All authors have read
and approved the manuscript.

Financial Support
This work was supported by the Japan Society for the Promotion of Science [JSPS
KAKENHI (JP20K15832 to S.T.), a Grant-in-Aid for Early-Career Scientists (JP22K15135 to
H.M.), a Grant-in-Aid for Scientific Research on Innovative Areas (JP19H05670 and
JP19H05676 to M.U.), a Grant-in-Aid for Scientific Research (B) (JP23H02494 to M.U.),
International Leading Research (JP22K21352 to M.U.)], the Japan Science and Technology
Agency [CREST (JPMJCR2121)],, the Suntory Rising Stars Encouragement Program in Life
Sciences (SunRiSE; to M.U.), and the Toray Science Foundation (20-6102 to M.U.).

Data Availability Statement
The data supporting the findings of this study are available from the author, Tomonobu
Nonoyama, upon a reasonable request.

Competing Interest

The authors have no competing interest.

References:

1. Tanaka, S., Matsushita, Y., Hanaki, Y., Higaki, T., Kamamoto, N., Matsushita, K., Higashiyama, T., Fujimoto, K., Ueda, M. (2024) HD-ZIP IV genes are essential for embryo initial cell polarization in the radial axis initiation in *Arabidopsis*. *Current Biology*. 0960-9822. doi: 10.1016/j.cub.2024.08.03

2. Kimata, Y., Higaki, T., Kawashima, T., Kurihara, D., Sato, Y., Yamada, T., Hasezawa, S., Berger, F., Higashiyama, T., Ueda, M.（2016）. Cytoskeleton dynamics control the first asymmetric cell division in Arabidopsis zygote. *Proceedings of the National Academy of Sciences*, 113(49), 14157-14162.

3. Kang, Z. C., Matsumoto, H., Nonoyama, T., Nakagawa, S., Ishimoto, Y., Tsugawa, S., Ueda, M. (2023). Coordinate normalization of live-cell imaging data reveals growth dynamics of the Arabidopsis zygote. *Plant Cell Physiology,* pcad020.

4. Kang, Z. C., Nonoyama, T., Ishimoto, Y., Matsumoto, H., Nakagawa, S., Ueda, M., Tsugawa, S., An elasto-plastic deformation model of hemispherical tip growth in Arabidopsis zygotes. *Submitted*.

5. Kang, Z. C., Nakagawa, S., Matsumoto, H., Ishimoto, Y., Nonoyama, T., Hanaki, Y., Tsugawa, S., Ueda, M. (2024). Material derivative of tension ensures accurate asymmetric cell division of Arabidopsis zygotes, *bioRxiv*.

6. Braun, M., Baluska, F., Witsch, M. V., Menzel, D., (1999). Redistribution of actin, profilin and phosphatidylinositol-4, 5-bisphosphate in growing and maturing root hairs. *Planta,* 209, 435–443

7. Cheung, A. Y., Duan, Q. H., Costa, S. S., Barend, H. J. G., Stillo, V. S., Feijo, J., Wu, H. M., (2008). The Dynamic Pollen Tube Cytoskeleton: Live Cell Studies Using Actin-Binding and Microtubule-Binding Reporter Proteins. *Molecular Plant*, 1(4), 686 - 702

8. Hamant, O., Heisler, M. G., Jönsson, H., Krupinski, P., Uyttewaa, M., Bokov, P., Corson, F., Sahlin, P., Boudaoud, A., Meyerowitz, E. M., Couder, Y., Traas, J., (2008). Developmental patterning by mechanical signals in Arabidopsis. *Science*. 322(5908), 1650-1655.

9. Uyttewaal, M., Burian, A., Alim, K., Landrein, B., Borowska-Wykręt, D., Dedieu, A., Peaucelle, A., Ludynia, M., Traas, J., Boudaoud, A., Kwiatkowska, D., Hamant, O. (2012). Mechanical stress acts via katanin to amplify differences in growth rate between adjacent cells in Arabidopsis. *Cell*. 149(2), 439-451.

10. Sampathkumar, A., Krupinski, P., Wightman, R., Milani, P., Berquand, A., Boudaoud, A., Hamant, O., Jönsson, H., (2014). Meyerowitz. Subcellular and supracellular mechanical stress prescribes cytoskeleton behavior in Arabidopsis cotyledon pavement cells. *eLife*, 3:

456  e01967.

11. Hervieux, N., Tsugawa, S., Fruleux, A., Dumond, M., Routier-Kierzkowska, A-L., Komatsuzaki, T., Boudaoud, A., Larkin, J. C., Smith, R. S., Li, C-B., Hamant, O. (2017). Mechanical Shielding of Rapidly Growing Cells Buffers Growth Heterogeneity and Contributes to Organ Shape Reproducibility, *Current Biology*. 27(22), 3468-3479.

12. Sapala, A., Runions, A., Routier-Kierzkowska, A-L., Gupta, M. S., Hong, L., Hofhuis, H., Verger, S., Mosca, G., Li, C-B., Hay, A., Hamant, O., Roeder, A. H. K., Tsiantis, M., Prusinkiewicz, P., Smith, R. S. (2018). Why plants make puzzle cells, and how their shape emerges. *eLife*, 7, e32794.

13. Inoue, D., Nitta, T., Kabir, A. M.R., Sada, K., Gond, J. P., Konagaya, A., Kakugo, A., (2016). Sensing surface mechanical deformation using active probes driven by motor proteins. *Nature Communications,* 7, 12557.

14. Hamant, O., Inoue, D., Bouchez, D., Dumais, J., Mjolsness, E. (2019). Are microtubules tension sensors? *Nature Communications*, 10(1), 2360.

15. Durand-Smet, P., Spelman, T. A., Meyerowitz, E. M., Jönsson, H. (2020). Cytoskeletal organization in isolated plant cells under geometry control. *Proceedings of the National Academy of Sciences*, 117(29), 17399-17408.

16. Colin, L., Chevallier, A., Tsugawa, S., Gacon, F., Godin, C., Viasnoff, V., Saunders, T. E., Hamant, O. (2021). Cortical tension overrides geometrical cues to orient microtubules in confined protoplasts. *Proceedings of the National Academy of Sciences*, 117(51), 32731-32738.

17. Moulia, B., Douady, S., Hamant, O. (2021). Fluctuations shape plants through proprioception. *Science*. 372(6540), eabc6868.

18. Heath, I. B., (1974). A unified hypothesis for the role of membrane bound enzyme complexes and microtubules in plant cell wall synthesis. *Journal of Theoretical Biology*. 48(2), 445-449.

19. Paradez, A. R., Somerville, C. R., Ehrhardt, D. W., (2006). Visualization of cellulose synthase demonstrates functional association with microtubules. *Science*. 312(5779), 1491-1495.

20. Balsuka, F. B., Samaj, J., Wojtaszek, P., Volkmann, D., Menzel, D. (2003). Cytoskelen-plasma Membrane-cell wall continuum in plants. Emerging links revisited. *Plant physiology*, 133(2), 482-491.

21. Lloyd, C., Chan, J. (2008). The parallel lives of microtubules and cellulose microfibrils. *Current Opinion in Plant Biology*. 11(6), 641-646.

22. Mitchison, T., Kirschner, M. (1984). Dynamic instability of microtubule growth. *Nature*. 312, 237-242.

23.   Shaw, S. L., Kamyar, R., Ehrhardt, D. (2003) Sustained microtubule treadmilling in
Arabidopsis cortical arrays. *Science*. 300(5626), 1715-1718.

24.   Dixit, R., Cyr, R. (2004). Encounters between dynamic cortical microtubules promote
ordering of the cortical array through angle-dependent modifications of microtubule
behavior. *Plant Cell*. 16(12), 3274-3284.

25.   Dixit, R., Cyr, R. The cortical microtubule array: From Dynamics to Organization. *Plant*
Cell.   16(10), 2546-2552.

26.   Dumond, M. (2013) Two main ways to model the self-organization of cortical microtubules.
BioSciences Master Reviews in Ecole Normale Superieure de Lyon.

27.   Shi, X-Q., Ma, Y-Q., (2010). Understanding phase behavior of plant cell cortex
microtubule organization. *Proceedings of the National Academy of Sciences Sciences.* 107,
11709-11714.

28.   Tindemans, S. H., Hawkins, R. J., Mulder, B. M. (2010). Survival of the Aligned: Ordering
of the Plant Cortical Microtubule Array. *Physical Review Letters,* 104, 058103.

29.   Tindemans S. H., Deinum, E. E., Lindeboom, J. J., Mulder, B. M. (2014). Efficient
event-driven simulations shed new light on microtubule organization in the plant cortical
array. `Frontiers in Physics` 2, 1-15.

30.   Chakrabortty, B., Willemsen, V., Zeeuw, T., Liao, C-Y., Weijers, D., Mulder, B. M., Scheres,
B. (2018-1). *Current Biology*, 28(19): 3031-3043.e2.

31.   Chakrabortty, B., Blilou, I., Scheres, B., Mulder, B. M. (2018-2). A computational
framework for cortical microtubule dynamics in realistically shaped plant cells. *Plos*
Computational Biology, 14(2), e1005959.

32.   Hawkins, R. J., Tindemans, S. H., Mulder, B. M. (2010). Model for the orientational
ordering of the plant microtubule cortical array. *Physical Review E*, 82, 011911.

33.   Mirabet, V., Krupinski, P., Hamant, O., Meyerowitz, E., Jönsson, H., Boudaoud, A. (2018)
The self-organization of plant microtubules inside the cell volume yields their cortical
localization, stable alignment, and sensitivity to external cues. *Plos Computational Biology,*
14(2), 21006011.

34.   Bastien, R., Legland, D., Martin, M., Fregosi, L., Peaucelle, A., Douady, S., Moulia, B.,
Höfte, H. (2016). KymoRod: a method for automated kinematic analysis of rod-shaped
plant organs. *The Plant Journal,* 88, 468-475.

35.   Shaw, S. L., Kamyar, R., Ehrhardt, D. W. (2003). Sustained Microtubule Treadmilling in
Arabidopsis Cortical Arrays. *Science,*300, 1715-1718.

36.   Eren, E. C., Dixit, R., Gautam, N. (2010). A Three-Dimensional Computer Simulation
Model Reveals the Mechanisms for Self-Organization of Plant Cortical Microtubules into
Oblique Arrays, *Molecular Biology of the Cell.* 21:15, 2674-2684,

37. Hasezawa, S., Kumagai, F., Nagata, T., Sites of microtubule reorganization in tabacco BY-2
cells during cell-cycle progression. (1997). *Protoplasma*, 198, 202-209

38. Murata, T., Wada, M. (1989). Organization of cortical microtubules and microfibril
deposition in response to blue-light-induced apical swelling in a tip-growing *Adiantum*
protonema cell. *Planta.* 178, 334–341.

39. Bedinger, P. (1992). The remarkable biology of pollen. *Plant Cell.* 4(8), 879-87..

40. Schiøtt, M., Romanowsky, S. M., Bækgaard, L., Harper, J. F., (2004). A plant plasma
membrane Ca2+ pump is required for normal pollen tube growth and fertilization.
Proceedings of the National Academy of Sciences. 4101(25), 9502-9507.

41. Öztürk-Çolak, A., Moussian, B., Araújo, S. J., Casanova, J. (2016). A feedback mechanism
converts individual cell features into a supracellular ECM structure in Drosophila trachea,
eLife, 5, e09373

42. Robinson, S., Kuhlemeier, C. (2018). Global Compression Reorients Cortical Microtubules
in Arabidopsis Hypocotyl Epidermis and Promotes Growth, *Current Biology.* 28(11).

Tables:

Table 1. Simulation parameters and variables with their default values. The height and radius of

the simulation cylinders are based on the observed data in actual *Arabidopsis* zygotes (Kang et

al., 2023). The threshold angle of zippering and the probabilities of induced catastrophes and

crossover were based on Dixit & Cyr (2004a) and Tindemans et al. (2014b). The strength of the

directional cue (DC) was based on Mirabet et al. (2018).

parameter	Value	Meaning
H	20 [µm]	Height of the cylinder
R	5.0-15.0 [µm]	Radius of the cylinder
N	6000,12000	Simulation steps
Δt	0.02 [sec]	Time interval of the simulation steps
v^+	0.08 [µm/sec]	Growth speed of the plus end
v^-	0.16 [µm/sec]	Shrinkage speed of the plus end
v^{tm}	0.01 [µm/sec]	Shrinkage speed of the minus end
r_r	0.007 [1/sec]	Rescue rate
r_c	0.001-0.01[1/sec]	Catastrophe rate
r_n	0.001 [1/sec]	Nucleation rate
θ^*	40 [degree]	Threshold angle of CMT interactions
P_{cat}	0.5	Probability of induced catastrophe
P_{cro}	0.5	Probability of crossover
b_d	0.0001-0.01	Strength of the DC
μ_T	10.0 [µm]	Position of the below end of the DC zone
σ_T	2-10 [µm]	Width of the DC zone

548   Table1

 Figures:

Fig. 1. Schematic illustrations of cortical microtubule (CMT) dynamics, their interactions, and

the effect of directional cue (DC). (A) Microtubule (MT) growth, when the CMT plus end

grows at speed v^+. (B) MT shrinkage, when the CMT plus end shrinks at speed v^- while the

CMT minus end undergoes shrinkage at speed v^{tm}. Switching rate from growth to shrinkage at

the plus end, called the rescue rate r_r, and switching rate from shrinkage to growth at the plus

end, called the catastrophe rate r_c, are also shown. (C) CMT zippering, when one CMT

interacts with another within 40° and changes its direction to align with the other. At this angle

of interaction, zippering occurs at 100% frequency. (D) Induced CMT catastrophe, when one

CMT interacting with another at greater than 40° is induced to depolymerize. (E) CMT

crossover, when two CMTs interact with each other at greater than 40° and cross one another.

The induced catastrophes (D) and crossovers (E) each occur with a probability of 50% when

CMTs meet at an angle greater than 40°. (F) CMTs with DC, which causes CMTs within the DC

zone to change their direction to gradually align to a specific direction (circumferential in this

study).

Fig. 2. Quantification of the cortical microtubule (CMT) band from live cell imaging data of the

Arabidopsis zygote. (A) Two-photon excitation microscopy (2PEM) images of the time-lapse

observation of a zygote expressing the MT/nucleus marker. Numbers indicate the elapsed time

(h:min). Scale bar: 10 μm. (B) Schematic illustrations of zygote growth. MT, microtubule. (C)

Definition of the cell centerline *S* estimated from the cell contour. (D) Quantification of the cell

contour and the cell centerlines. The colors of the contours represent the time from the first

frame. (E) Quantification of the CMT band. The green line indicates the width of the CMT band

(defined below). (F) Truncated Gaussian fitting of the fluorescence intensity projected on the

centerline (black line). The red line represents the fitted truncated Gaussian curve. The position

of the CMT band μ is estimated by the median, and the width of the CMT band is estimated by

σ_{MT}^{data}.. (G) Kymograph of the fluorescence intensity projected on the centerline of the zygote.

The green and blue lines indicate the upper and lower limits of the CMT band, respectively,

where s is the standard deviation(H) MT dynamic space (MTD space, $n = 4$, indicating the

speed and width of the CMT bands with their frequencies (histograms)). The vertical axis

denotes σ_{MT}^{data} and the horizontal axis is $d\mu_{MT}^{data}/dt$. The 95[th] percentile of the MTD space is

shown as a pale orange region. Typical CMT patterns for the cases of the red circles are

schematically illustrated.

Fig. 3. Effect of the tension-responsive ability parameter (b_d) and the width of the directional

cue (DC) σ_T. (A–C) Simulation results when using no DC (A), a weak DC ($b_d = 0.001$; B),

and a strong DC ($b_d = 0.01$; C). The orientation angle of the CMT measured from the

horizontal axis is represented by the color used. The DC zone is indicated by black dashed lines.

(D) Orientational change of a single CMT with vertical growth as a function of b_d. The DC

zone is highlighted in orange. (E) Dependence of the CMT width σ_{MT}^{model} as a function of b_d.

(F) Dependence of the global order parameter Λ as a function of b_d. (G, H) Results of the

simulations at $t = 20$ [min] for different σ_T values in the case of a weak DC (G) and a strong

DC (H). (I) Dependence of the width σ_{MT}^{model} as a function of σ_T in the case of a weak or

strong DC.

Fig. 4. Effect of the cell radius R and two different types of cortical microtubule (CMT)

organization. (A, B) Simulation results for a strong directional cue (DC) with a temporal

dependence of Λ and σ_{MT}^{model} when $R = 5.0$ [μm] (A) and $R = 15.0$ [μm] (B). The green

lines represent the sigmoid function for Λ and the linear function for σ_{MT}^{model}. (C) Schematic

illustrations of rapid CMT organization. Owing to the horizontal CMTs that were rapidly

organized, diagonal CMTs may be disorganized, and only the horizontal CMT band remains.

The green dashed lines represent newly forming CMTs, where the red circles at the leading edge

indicate those that do not collide with pre-existing CMTs, making them likely to grow. In contrast,

the red crosses at the leading edge represent CMTs that collide with pre-existing CMTs, indicating a

reduced likelihood of growth. The colored band represents the region under strong DC influence. (D,

E) Simulation results for a weak DC with a temporal dependence of Λ and σ_{MT}^{model} when

$R = 5.0$ [μm] (D) and $R = 15.0$ [μm] (E). For cases where the CMT band cannot be

defined, the corresponding box plot is omitted. (F) Schematic illustrations of gradual CMT

organization. The green dashed lines represent newly forming CMTs, where the red circles at

the leading edge indicate those that do not collide with pre-existing CMTs, making them likely

to grow. In contrast, the red crosses at the leading edge represent CMTs that collide with

pre-existing CMTs, indicating a reduced likelihood of growth. The colored band represents the

region under weak DC influence.   Without horizontal CMTs, diagonal CMTs can be organized

so that the CMT band becomes widened.

Fig. 5. Parameter inference by mapping input directional cue (DC) parameters to output the

cortical microtubule (CMT) dynamic space (CDS). (A) Schematic illustrations of the

microtubule dynamic (MTD) space in the data, with expected input DC parameters (green) and

a similar range of input DC parameters to the MTD space (dashed line). (B) Simulated results in

the CDS with the data range from Fig. 2H (green). (C) The discrepancy between the output and

input CMT speed compared with the diagonal equivalent line (gray dashed line). Depending on
the output speed of the CMT band, there are static states, states that follow the input, and states
that lag behind the input. (D, F, H) CMT dynamics with appropriate speed and width for the
resulting CMT band (Case 01) (D), with a faster speed than the CMT band (Case 02) (F), and
with a broader width and faster speed than the CMT band (Case 03) (H). (E, G, I) Schematic
illustration of the obtained results. Depending on the input speed of the DC zone, the resulting
CMT band did not move (E, i), moved appropriately (G, ii), or lagged behind the DC zone (I,
iii).
628

(A) MT growth (B) MT shrinkage

(C) Zippering (<40°) (D) Catastrophe (>40°)

(E) Crossover (>40°) (F) Directional cue

v^{tm} v^{+} (A) r_c

v^{-} (B) r_r

(C) <40°

(D) >40°

(E) >40°

(F)

(A) $b_d = 0$

$t=8$ [min] $t=24$ [min] $t=40$ [min]

(B) $b_d = 0.001$

$t=8$ [min] $t=24$ [min] $t=40$ [min]

(C) $b_d = 0.01$

$t=8$ [min] $t=24$ [min] $t=40$ [min]

(D)

$b_d = 0.01$
$b_d = 0.005$
$b_d = 0.001$
$b_d = 0.0005$
$b_d = 0.0001$
DC zone

(E)

DC strength b_d

(F)

DC strength b_d

(G) $b_d = 0.001$

$\sigma_T = 2$ [µm] $\sigma_T = 6$ [µm] $\sigma_T = 10$ [µm]

(H) $b_d = 0.01$

$\sigma_T = 2$ [µm] $\sigma_T = 6$ [µm] $\sigma_T = 10$ [µm]

(I)

strong DC
weak DC

DC length σ_T [µm]

(A) *R*=5.0 [μm]

(B) *R*=15.0 [μm]

(C) rapid organization

CMT band

(D) *R*=5.0 [μm]

(E) *R*=15.0 [μm]

(F) gradual organization

CMT band

(A) Input cue parameters → Output MT space
DC area length, DC zone speed → CMT band length, CMT band speed

(B) CMT band length σ_{MT}^{model} [μm] vs CMT band speed $\frac{d\mu_{MT}^{model}}{dt}$ [$\mu m/hour$]
- simulation
- data

(C) CMT band speed in model $\frac{d\mu_{MT}^{model}}{dt}$ [$\mu m/hour$] vs DC zone speed $\frac{d\mu_T}{dt}$ [$\mu m/hour$]
(i), (ii), (iii)

(D) ⭕ DC zone speed= 2.0 [$\mu m/hour$], DC zone length= 3.0 [μm]
t=10 [min]   t= 20 [min]   t= 30 [min]   t= 40 [min]

(E) (i) DC zone speed : appropriate
input   output
CMT band speed ≈ input speed

(F) ⭕ DC zone speed = 1.5 [$\mu m/hour$], DC zone length= 4.0 [μm]
t=10 [min]   t= 20 [min]   t= 30 [min]   t= 40 [min]

(G) (ii) DC zone speed : slow
input   output
CMT band speed ≈ 0

(H) ⭕ DC zone speed = 3.0 [$\mu m/hour$], DC zone length= 4.0 [μm]
t=10 [min]   t= 20 [min]   t= 30 [min]   t= 40 [min]

(I) (iii) DC zone speed : fast
input   output
CMT band speed < input speed