Facial Coloration as a Social Signal in a Neotropical Cichlid: Testing the Facial Expression Hypothesis

Shun Satoh [1,2†*], Kazuya Fukuda [3†*], Sayaka Matsuo [3], Takezo Banda [3], Kota Kanbe [3], Alu Konno [4], Shiro Takei [5], Masanori Kohda [6], and Nobuyuki Kutsukake [7]

[1] The Hakubi Center for Advanced Research, Kyoto University, Yoshida-honmachi, Sakyo-ku, Kyoto, Japan

[2] Department of Zoology, Graduate School of Science, Kyoto University, Kitashirakawa-oiwake-cho, Sakyo-ku, Kyoto, Japan

[3] School of Marine Biosciences, Kitasato University, Kitazato, Minami-ku, Sagamihara city, Kanagawa, Japan

[4] Department of Virology and Parasitology, Hamamatsu University School of Medicine, Handayama, Higashi-ku, Hamamatsu city, Shizuoka, Japan

[5] Department of Environmental Biology, Chubu University, Matumoto-cho, Kasugai-shi, Aichi, Japan

[6] Department of Biology, Graduate School of Science, Osaka Metropolitan University, Sumiyoshi-ku, Osaka city, Osaka, Japan

[7] Research Center for Integrative Evolutionary Science & Department of Evolutionary Studies

of Biosystems, The Graduate University for Advanced Studies, Hayama, Kanagawa, Japan

[†] Equally contribution

Abstract

The face is a uniquely distinctive stimulus for humans, encapsulating a wealth of information.

Among the myriad of social cues conveyed by the face, emotional signals, known as facial

expressions, are paramount not only for humans but also for numerous social animals. The

evolution of facial expressions in these animals can also manifest in taxa other than mammals,

as suggested by various studies highlighting the socio-ecological benefits of facial expressions.

In this study, we elucidated the social function of facial coloration, determined by

melanophores, in the neotropical social cichlid *Symphysodon aequifasciatus*. In this species,

facial coloration exhibits flexibility and instantaneous changes in response to varying social

contexts. Through behavioral observations and experimental manipulation, we confirmed that

facial coloration in *S. aequifasciatus* serves to attenuate unnecessary aggressive competition

among conspecifics. Furthermore, we observed that the facial area subjected to coloration in this

species is innervated by the adenosine triphosphate- and noradrenaline-ergic nervous system.

These findings indicated that facial expression in *S. aequifasciatus* depends on the sympathetic

nervous system and has evolved independently of mammalian facial expressions. Our study

highlights teleost fishes as valuable animal models for exploring the universality of facial

expressions and their underlying cognitive mechanisms in vertebrates.

Introduction

Human faces, which are the most important stimuli in various social contexts, are remarkably unique as they encapsulate a large amount of information and social cues related to various aspects, including sex [1], identity [2], emotional states [3], and attentive states [4,5]. In particular, humans are capable of manifesting emotions through facial movements (commonly referred to as facial expressions), which comprise specific motions that support optimal emotional signaling and are controlled by specific neural systems [6]. Human expressions are primarily produced by the action of muscles below the facial skin, commonly known as facial muscles [6,7]. These muscles are innervated by the facial nerve, which has the largest motor nucleus among all motor nuclei in the brainstem [8]. Additionally, the involvement of at least five cortical regions in the upstream regulation of motor control highlights the crucial role of facial expressions in humans [8]. Facial expressions are appropriately interpreted by the receiver's specialized neural circuits, including regions such as the superior temporal sulcus and amygdala, thereby conveying complex social and emotional information [9-12]. These elaborate systems are universal among humans, irrespective of cultural differences [13]. It is undeniable that facial expressions hold a unique and significant role in the communication of human emotions. Thus, delving into the evolutionary history of facial expressions presents an intriguing challenge within the realm of evolutionary biology.

A valuable methodology for elucidating the evolutionary trajectory of human facial expressions involves juxtaposing them with patterns of emotional expression observed in animals. It was Charles Darwin who initially recognized the significance of comparing the facial expressions of animals and humans, along with their associated cognitive abilities [14, 15]. In his seminal work "*The Expression of the Emotions in Man and Animals*," Darwin postulated that human emotional expressions have evolved from those of our animal ancestors rather than being the product of our society, suggesting their comparability [16], a proposition that sparked

considerable controversy [14]. While the degree of comparability between human and animal

facial expressions remains a contentious subject, research in the 21st century has provided

evidence that non-human primates are able to convey many of the same facial expressions

observed in humans through intricate facial muscle movements [17,18]. These studies suggest that

facial expressions in primates are species-specific and that some are socially learned, while also

reporting many innate aspects [19]. However, regardless of the patterns of facial expression in

primates, the dominant theory is that the face is important as a signal for the communication of

emotional states to other individuals. Moreover, there has been some debate as to what extent

the use of specialized visual cues from the face and the consequent development of associated

specializations within the neural system are unique features found only in primates. For

instance, laboratory mice [20] and macaws [21] have been shown to display a variety of discernible

facial expressions.

Interestingly, as observed in humans, faces in these animals are recognized as vehicles

for conveying information and social cues related for example to identity, dominance rank, and

sex [19,22], alongside emotional signals. In these animals, multiple social cues are thought to be

centralized in the face because of the evolutionary consequences of attention to the eyes [19].

Moreover, eyes serve as important signals to gaze direction and objects of interest by other

individuals [23,24], resulting in social benefits for both signalers and signal receivers in social

animals. Importantly, according to these theories, we can predict that the centralization of social

cues, including facial expressions and the cognitive systems that process them, occur

convergently in social animals, regardless of the taxonomic group.

Social fishes are good candidates for testing this hypothesis regarding the evolution

and social benefits of facial expression. Teleost fishes lack the elaborated facial musculature of

mammals; therefore, it has been thought that emotional facial expressions are restricted in this

taxon [19]. However, recent studies have pointed to peculiarities in the faces of social fishes [25].

Specifically, numerous species spanning various taxonomic groups have been documented to

exhibit notable morphological or color variations localized to the face serving as social cues for

inter-specific or individual recognition [25-31]. Moreover, it has been posited that fish possess

specialized cognitive and neural processing mechanisms for facial recognition [28, 32] and that the

facial area may also play a pivotal role in self-recognition in cleaner wrasse [33]. As well as using

social signals related to individual recognition, social fishes are known to initially focus on the

face when presented with images depicting conspecifics [30], like primates do [34]. All these

investigations underscore the face-related peculiarities in fish. Studying facial expressions in

these animals, the vertebrates situated at the furthest phylogenetic distance from mammals

(which have traditionally been the primary subjects of such studies), contributes to a deeper

understanding of the general principles underlying the evolution of this phenomenon. However,

to date, no studies have been conducted on emotional signals in fish based on the perspective of

facial expressions.

To address this knowledge gap, we here focused on the social function of facial

coloration in the discus fish *Symphysodon aequifasciatus*, a neotropical cichlid species that

dwells in the Amazon River basin [35]. These fish reside in sunken trees alongside their

conspecifics during the non-breeding dry season and reproduce in flooded forests throughout

the breeding rainy season [36]. Both male and female members in reproductive pairs provide

parental care and nourishment in the form of mucus for their offspring [37-39]. Remarkably, *S.*

aequifasciatus individuals are able to differentiate between their breeding partner and unfamiliar

fish by analyzing the structural color pattern on their faces [25]. Moreover, we observed that the

vertical black bar located above their eyes, which is composed of melanophores [40], undergoes

frequent changes during social interactions with conspecifics (Fig. 1a; Supplemental movie S1).

Individuals have vertical black bars on their body, including the facial area. Despite population

and individual differences in vertical black bars along the body flank, facial black bars (the so-

called eye-patch pattern) are universal among discus fishes [41]. Therefore, we hypothesized that

this facial coloration may serve as a mode of emotional expression.

In this study, we aimed to clarify whether facial coloration in *S. aequifasciatus*

functions as an emotional signal by using two complementary approaches: behavioral

experiments and neuroscientific methods. In particular, we evaluated whether immediate

changes in facial coloration can be referred to as facial expression by testing 1) whether color

change was concentrated on the face in different social contexts and 2) whether variation in face

color suppressed unnecessary social costs. Through a combination of behavioral observations

and experimental manipulation of facial coloration, we seeked to elucidate the role of this

phenomenon as a social signal. Moreover, using experiments to determine which

neurotransmitters activate melanophores in the facial area [40] and the slightly modified DEEP-

Clear method [42], a clearing method for exploring the peripheral innervation of facial areas

subjected to coloration, we attempted to identify which peripheral nerves are responsible for

face color changes in *S. aequifasciatus* and examined whether these are analogous to facial

expressions in mammals.

Results and Discussion

Association between social behaviors and facial pigmentation patterns

To explore the association between distinct social behaviors and pigmentation patterns, we

conducted behavioral observations of free-ranging *S. aequifasciatus* individuals (n = 8 males

and 4 females) under simulated natural conditions. The pigmentation pattern, which

encompasses melanophores in discus fishes, was classified into three categories: 1) complete

absence of all vertical bars on the body, 2) manifestation of solely the eye-patch pattern, and 3)

manifestation of all vertical bars, including the eye-patch pattern (Fig. 1b). Specifically, when

exhibiting aggressive behavior toward conspecifics, individuals typically abstained from

displaying eye-patch patterns (Fig. 1b). Conversely, when under attack or during periods of rest

and foraging, they did show them. Interestingly, eye patch patterns disappeared during sexual

behavior as well as during aggression. With regard to the third pigmentation pattern, in which

all vertical bars are manifested, individuals did not show it frequently, regardless of social

behaviors.

The pigmentation patterns in *S. aequifasciatus* were clearly correlated with certain

social behaviors and were subject to rapid alteration, as shown in Supplementary movie S1.

Numerous instances of body color change serving as a social signal have been documented in

various social fishes. For example, in the group-living cichlid *Neolamprologus pulche*r [43] and

mouth-brooding cichlid *Astatotilapia burtoni* [40,44], the intricate pigmentation pattern in the facial

area serves as a social signal indicating aggressiveness, as similarly observed in discus fishes.

However, there is no conclusive evidence that these changes in pigmentation pattern occur

immediately in conjunction with the specific display, which would reflect a motive for

aggression associated with social rank over a longer term, as suggested by Rodrigues et al.

(2009) [45]. Color change in the facial area in discus fishes is more akin to facial expression

reported in primates [19] than to variation in facial coloration reported in other fish species.

The expression of distinct emotional states on the human face is characterized by the

subtle movements of facial musculature [13]. Conversely, in discus fishes, it is the alteration of

pigmentation patterns in the facial area generated by the contraction of chromatophores, rather

than facial muscles, that leads to the manifestation of specific facial expressions that correspond

to specific emotional states. In particular, in this study, *S. aequifasciatus* individuals showed the

eye-patch pattern when they were being fed or had received aggression but did not display it

when engaged in aggressive or partner-specific behaviors. This suggested that the presence of

the eye-patch pattern may be indicative of normalcy (submissive behavior) while its absence

may denote excitement (aggressive behavior).

Benefits of facial coloration patterns as social signals

To examine whether specific facial coloration patterns influence the degree of same-sex sexual

harassment perpetrated by dominant individuals, we observed a series of aggressive behaviors

between subordinate and dominant individuals for 20 minutes. We specifically focused on

whether a subordinate that was under attack could decrease the possibility of being subjected to

continuous harassment (defined as an attack occurring within 5 seconds of the first attack) from

a dominant individual by modifying its eye-patch pattern.

When a subordinate individual was attacked, alterations in facial coloration

significantly affected the occurrence of persistent harassment (binomial generalized linear

mixed models [GLMM], df = 3, χ^2 = 63.589, p < 0.001) (Fig. 2). Subordinate individuals who

did not exhibit the eye-patch pattern and did not modify their facial coloration upon being

attacked by dominant individuals were significantly more likely to experience continuous

harassment compared with those who did (df = 1, χ^2 = 54.186, p < 0.001). Similarly, in

subordinate individuals who maintained the eye-patch pattern before and during harassment, the

possibility of continuous attacks also decreased (df = 1, χ^2 = 10.05, p = 0.002). Conversely, the

likelihood of continuous harassment remained comparable when subordinate individuals

without the eye-patch pattern failed to alter their facial coloration during aggression, regardless

of whether those initially displaying the pattern modified their facial coloration (df = 1, χ^2 =

2.093, p = 0.148). Our behavioral observations suggested that alterations in facial coloration

serve to deter harassment from dominant individuals in *S. aequifasciatus*.

However, because facial coloration in discus fishes may be correlated with a variety of

social behaviors and psychological motivations [25], it is impossible to say whether it has a

specific function in inhibiting harassment from dominant individuals. Therefore, we conducted

experiments to manipulate facial coloration in *S. aequifasciatus* using anecdotal methods based

on light exposure commonly employed by ornamental fish keepers. As shown in several studies

conducted in the field of aquaculture science [46], exposing juvenile fish to intense light can affect

the melanophore-derived body coloration. Therefore, we obtained *S. aequifasciatus* individuals

in which the function of melanophores was abolished by exposing them to intense light during

the juvenile stage. The manipulated discus fish were unable to exhibit melanophore patterns

regardless of social context (Supplementary S2), and thus were considered to be unable to

convey proper social communication signals. In the behavioral experiment, we observed

changes in aggressive interaction over time in experimental pairs in which 1) both individuals

could properly display the eye-patch pattern (WW pairs, n = 8 pairs), 2) one individual was

light-treated (WL pairs, n = 6 pairs), and 3) both individuals were light-treated (LL pairs, n = 7

pairs).

210   We detected a significant two-way interaction between the first day of the behavioral

experiment and the type of experimental pair in terms of the frequency of aggressive

interactions over a period of 20 minutes (negative binomial GLMM, df = 2, χ^2 = 67.968, p <

0.001). In the WW group, the frequency of aggressive interactions between same-sex

individuals decreased day by day, but in the LL group, which was composed of pairs in which

both individuals who could not express optimal emotional signals, the frequency of aggressive

interactions did not decrease within days (Fig. 3). In the LW group, the variation in the

frequency of aggressive interaction was in-between the values observed for the WW and LL

groups (Fig. 3). In this experiment, the manipulation inevitably affected not only facial

coloration but also melanophore-derived pigmentation patterns in the rest of the body. However,

considering that pigmentation patterns other than those in the face were largely unchanged in

social displays between conspecifics (Fig. 1), the obtained results can be safely interpreted as

being caused by the inability of facial melanophores to produce patterns, which therefore

prevented proper communication between conspecifics.

224   During the experimental days, individuals lost 0.5%–8.4% of their body weight. The

percentage of the reduction in body weight significantly varied among experimental groups,

with LL suffering the greatest weight losses, followed by WL and WW, in that order (linear

mixed models [LMM], df = 2, F = 4.398, p = 0.025) (Fig. 4). Moreover, whether individuals

became dominant (or subordinate) during the experiment also had a marginal effect on body

weight reduction (df = 1, F = 3.955, p = 0.061). Sex (df = 1, F = 2.354, p = 0.145) and whether

individuals had been subjected to the light treatment (df = 1, F = 0.224, p = 0.641) had no

significant effects on the reduction. Social signals have been shown to confer benefits to both

conveyors and recipients [47]. The differential reduction in body weight observed across

experimental groups in this study, irrespective of social hierarchy, aligns with the tenets of

classical adaptive signaling theory and provides compelling evidence that facial coloration

yields socio-ecological benefits in *S. aequifasciatus*. This species forms social aggregates in the

Amazon River, an environment characterized by high predation pressure [36], and the act of

leaving the group, even to assume a subordinate position, carries fitness costs [48]. Here, the

individuals whose ability to modify facial coloration had been manipulated were possibly

unable to express the submissive signal toward dominant individuals and thus establish a

dominance hierarchy. Facial coloration may serve to mitigate superfluous competition within

social aggregates characterized by dominance hierarchies.

Peripheral innervation regulating facial coloration

The results of our behavioral experiments indicated that facial coloration in *S. aequifasciatus*

functions as a social signal in a similar manner to human facial expressions. One approach to

confirm whether the two phenomena are indeed analogous is to compare their underlying

regulatory mechanisms. Therefore, we conducted two experiments to elucidate the types of

peripheral nerves innervating the facial area subjected to changes in coloration in *S.*

aequifasciatus. Generally, discus fishes exhibit two primary mechanisms of color change:

physiological color change, mediated by hormonal and neural regulation of peripheral nerves,

and morphological color change, involving the accumulation or depletion of chromophores [49-51].

Previous behavioral experiments have shown that changes in facial coloration in discus fishes

occur rapidly, within a few seconds. Since morphological color changes usually require days to

weeks [51], we hypothesized that the above-mentioned rapid changes are physiological, which are

known to occur over shorter periods. In particular, they may be under neural control, which has

been shown to regulate very rapid color changes within seconds [44, 49]. In discus fishes, body

color changes, particularly those involving black coloration via the aggregation and dispersion

of melanosomes within melanophores, are generally regulated by the sympathetic nervous

system [44, 52, 53]. Sympathetic postganglionic neurons in discus fishes are known to possess

several neurotransmitters, including noradrenaline (NA), adenosine triphosphate (ATP),

neuropeptide Y, and acetylcholine [50, 54], the first two of which are the primary factors influencing

the motile activity of melanosomes [49, 55]. Therefore, we first examined whether NA and ATP

control facial coloration in discus fish by administering these neurotransmitters ex vivo. In this

experiment, scales (n = 20) with chromatophores were freshly collected from eye patches and

observed under a microscope. The scales were then immersed in neurotransmitters (1 µM NA or

10 mM ATP) or in PBS, which was used as a control. We quantified the rate of change in

melanosome dispersion (%) in the scales and observed significant differences between ATP

exposure and PBS exposure (two-way interaction between time and treatment, df = 1, F =

200.888, p < 0.001 by LMM), with melanosomes showing dispersion in the ATP exposure

group (Fig. 5a, c). Conversely, NA exposure resulted in significant melanosome aggregation

compared to PBS exposure (two-way interaction between time and treatment, df = 1, F =

89.651 p < 0.001) (Fig. 5b, d). These results suggested that changes in facial coloration in discus

fish also depended on sympathetic innervation. To further validate this hypothesis, we

investigated whether sympathetic postganglionic neurons (noradrenergic neurons) reached the

eye-patch region via tissue clearing. Tissue clearing is a methodology to visualize appropriately

labeled structures in chemically cleared, three-dimensional tissues, which is especially useful to

track neurons or blood vessels [56, 57]. We cleared the heads of X individuals using the slightly

modified DEEP-Clear method [42] and performed immunohistochemical staining using an anti-

tyrosine hydroxylase (TH) antibody (the key enzyme that converts tyrosine to L-DOPA, the

initial step in NA biosynthesis). Finally, we performed refractive index matching using CUBIC-

R reagent [58]. The results showed that TH-immunoreactive (TH-ir) fibers reached the eye-patch

region (Fig. 5e). Furthermore, morphological observations indicated that these fibers connected

to the scales containing the melanophores constituting the eye patch (Fig. 5f, g). These findings

strongly suggested that sympathetic nerves innervate the facial area subjected to changes in

coloration in discus fish.

Our results suggested that, unlike human facial expressions, which are regulated by

facial nerves [7] via the neurotransmitter acetylcholine [59], facial coloration in the discus fish

examined is regulated by the sympathetic nervous system. This has been similarly observed in

the Tanganyikan cichlid, *Astatotilapia burtoni* [40,44]. Facial coloration serves as a social signal

and plays a significant role in social interactions with other individuals [60, 61]. Therefore, its

functioning to convey facial expression and its peripheral neural regulation observed in this

study may not be limited to discus fish but could be prevalent among social fish species. One of

the interesting aspects of our findings is that, while facial expressions and coloration similarly

function as social signals in humans and fish respectively, they are controlled by different

nervous systems. In humans, they are regulated by voluntary nerves, whereas in fish they are

controlled by the autonomic nervous system. Regulation by voluntary nerves implies that

humans can deceive others by displaying facial expressions that do not reflect their authentic

emotions [62.63]. Recently, it has been reported that fish also possess the ability to deceive others

tactically [64]. However, considering that facial coloration in fish is regulated by the autonomic

nervous system, it is unlikely that they can use neurally controlled rapid variations in facial

color for intentional deception. Conversely, fish employ various mechanisms for controlling

body coloration beyond neural regulation, so deception using non-neural body color changes

may still be possible [65].

In humans, blushing may function as a supplementary emotional signal that enhances

facial expressions, though its impact is relatively less significant than that of facial expressions

[66]. This phenomenon occurs involuntarily, triggered by sympathetic activation [66]. While the

contexts and detailed mechanisms differ, blushing bears certain similarities to the coloration

mechanism observed in discus fishes. Consequently, these findings may point to a form of

convergent evolution with respect to the role of the sympathetic nervous system in mediating

emotional expression through facial features.

Evolution of facial expression in social fishes

Of the body coloration composed of melanophores, *S. aequifasciatus* intensively linked and

changed only facial coloration upon engaging in social behavior toward conspecifics. Our

findings further validated that facial coloration in this species serves to mitigate superfluous

aggressive competition among conspecifics, thereby benefiting both signalers and recipients

within the dominance hierarchy. This supports the hypothesis that facial coloration in discus

fishes exhibits functional parallels to mammalian facial expressions, predominantly those

observed in primates [19]. Conversely, the neural mechanisms underlying facial coloration in these

fishes markedly differ from those regulating facial expressions in mammals, strongly suggesting

a convergent evolution of facial expressions in each lineage.

In humans, facial expressions are hypothesized to have initially evolved as a

mechanism for modulating physiological states within an individual (e.g., widening of the eyes

in expressions of fear, facilitating enhanced peripheral vision) before assuming a secondary role

as social signals [67]. The vertical bars present on discus fishes, which are typically found among

driftwood, may serve as forms of cryptic or disruptive coloration [68]. Indeed, when confronted

with predators or subjected to prolonged intra-species aggression, discus fishes display vertical

bars, including eye-patch patterns, spanning their entire body [41]. Similarly to facial expression in

humans, facial coloration in discus fishes may have initially evolved as a cryptic form of

coloration mediated by melanophores, before subsequently being repurposed as a means to

convey submissive social signals localized solely to the facial region, undergoing

coevolutionary adaptations to facilitate their perception. Moreover, as an alternative prospective

evolutionary scenario other than cryptic coloration, the signals linked to long-term hierarchical

dominance cues within the facial region [40,43,44] may have undergone evolutionary divergence

toward immediate modification in response to social stimuli.

When a fish needs cryptic body coloration, for example when threatened by predators,

it experiences acute stress [69,70]. This stress activates the sympathetic nervous system, rapidly

increases systemic cortisol levels, and raises plasma glucose levels, creating a physiological

state conducive to fight-or-flight responses against the stressor [69-71]. A similar situation

assumably occurs when subordinate individuals try to avoid harassment from dominant

individuals [69,72-74]. Therefore, it should be plausible to hypothesize that the mechanism

regulating variations in body and facial coloration via the sympathetic nervous system to escape

from predator-related threats has secondarily adapted to cope with similar behavioral and

physiological contexts (for example to avoid harassment from dominant individuals), even

though the stressors are different.

However, facial coloration in discus fishes changes independently from body

coloration, suggesting the presence of distinct control mechanisms for these areas. It is

conceivable that discus fishes have developed neural mechanisms to gradually or alternatively

change the coloration of their face and body in response to different stimuli and social contexts.

Alternatively, differentiated neural pathways controlling facial and body coloration may have

evolved, as exemplified by the unique peripheral nerve branches found in these fish species,

such as the recurrent facial taste neurons in sea catfish, which have acquired taste buds on their

body surface and fins in response to socio-ecological demands over the course of evolution [75].

Although additional research is required, either scenario strongly supports the hypothesis that

social signals are centralized on the face in social animals, regardless of the mechanisms

controlling facial and body coloration.

In recent years, various studies have examined the idiosyncrasies of faces to individual

recognition [25,27-31], self-recognition [33], attention, and specialized cognitive functions in fish [28 32].

To the best of our knowledge, there have been no reports documenting instantaneous emotional

signals on the faces of fish, as observed in discus fishes. However, this may stem from a paucity

of research on facial expressions in this animal group rather than from an absence of such

expressions. Indeed, in numerous fish species, social signals indicative of long-term dominance

hierarchies are localized to the facial region [40,44,45], despite the face being an anatomically

limited part of the body.

A comparative study conducted on primates has revealed a correlation between the

efficacy of facial expression and social complexity [76]. Therefore, exploring the relationship

between social systems and facial coloration in fish represents the next step to comprehensively

elucidate the evolution of facial expression within this taxon. Our study demonstrates that

teleost fishes are valid animal models for probing the universality of facial expressions and their

underlying cognitive mechanisms in vertebrates.

Methods

Subjects and fish husbandry

Mature male and female *S. aequifasciatus* specimens were procured from ornamental fish

traders that imported them from the Brasilia Amazon and bred F1 individuals in Japan. These

specimens were housed in four separate water tanks. Uniform fish density and sex ratio were

meticulously maintained across all stacked tanks. Water temperature and pH were carefully

regulated and kept within the ranges of 27.0°C–30.0°C and 5.0–7.0, respectively. The fish were

nourished with an artificial diet formulated from beef heart and mosquito larvae (three times per

day). The sex of specimens was determined based on their distinctive body coloration and

morphological characteristics. In total, eight males and four females were used for *behavioral*

observation I, and 20 males for *behavioral observation II*. All observations and experiments

were conducted from 12.00 to 15.00 and recorded using a video camera (HDR-CX470 and

HDR-CX370, Sony Corporation, Tokyo, Japan) placed at ca. 30 cm in front of the water tanks.

Behavioral observation I: pigmentation patterns of S. aequifasciatus in different social contexts

The behavior of *S. aequifasciatus* in a water tank was observed to determine what social context

correlated with pigmentation patterns, which change instantly (Supplemental movie S1).

Driftwood and dry leaves were placed in the tank (360 L, 120 × 60 × 50 cm), and the water was

colored using humic acid to mimic the natural habitat [36,41]. Three mature individuals (two males

and one female) were held in this tank for 3 days to allow accustomization to the environment.

During behavioral observations, which lasted for 25 min, mosquito larvae were fed to the fish to

promote social interaction among them, and the pigmentation patterns of focal fish were

evaluated when they showed six specific social behaviors (i.e., aggressive behavior, counter

aggression, received aggression, feeding, and partner display) based on the ethogram described

in Satoh et al. (2016)[25]. Aggressive behavior was defined as a fish approaching the lateral side

of a non-partner while opening its mouth and spreading its gill covers and fins as well as a fish

aggressively biting and/or bumping into the non-partner. A focal fish receiving aggressive

behaviors from other experimental fish and then showing the same behaviors was defined as

counter-aggression, while the lack of any behaviors upon receiving aggression was defined as

received aggression. Partner display was defined as two fish approaching each other holding up

their heads and folding their pelvic fins toward their body. This type of behavior is observed

when fish form monogamous reproductive pairs [25]. Additionally, rest was defined as a state in

which the above behaviors were not exhibited; specifically, pigmentation patterns in this state

were recorded about 1 minute after the other behaviors were observed. Observations of the rest

state were conducted randomly five times in eight male and four female adults.

Three types of pigmentation patterns were detected in *S. aequifasciatus* (Fig. 1).

Typically, discus fishes have a total of eight vertical bars on their body, including the facial area

[41]. However, in this study, individuals did not show vertical bars on their body flank when such

bars were absent on their facial area.

Behavioral observation II: effects of facial coloration on aggressive behavior

The objective of these observations was to investigate the impact of facial coloration on the

frequency of harassment by dominant individuals toward subordinates. To this end, 10 pairs of

mature males (n = 20) were placed within an experimental water tank (162 L, 60 × 45 × 60 cm).

In these same-sex experimental pairs, the fish differed in body size by a minimum of 1.5 cm,

which ensured the establishment of a size-based hierarchy between them; smaller individuals

were designated as subordinates, while larger individuals assumed the role of dominants. The

experiment was focused on determining whether the presence of an eye-patch pattern in

subordinate individuals served as an emotional signal to mitigate the possibility of continuous

aggression (harassment) from dominant individuals, given that eye-patch patterns had been

frequently displayed by subordinates in response to aggressive behavior from conspecifics (Fig.

1b).

First of all, upon observing aggressive behavior by dominants against subordinates,

the facial coloration of the latter (i.e., absence or presence of the eye-patch pattern) was

recorded (hereafter called 'first aggression'). After the first aggression, whether the dominant

individuals continuously attacked the subordinates was also recorded. In this study, an

aggressive behavior that occurred within 5 seconds of the first aggression was defined as

continuous harassment. If this situation was produced, the facial coloration of subordinates was

recorded. In contrast, if dominant individuals did not continuously attack, the facial coloration

of subordinates was recorded 5 seconds after the first aggression. If dominant individuals

repeated the aggressive behavior three or more times within a single sequence of aggressive

behaviors, these were considered as part of the first aggression, and observations continued in

the same manner until the continuous harassment ended. During these observations, which

lasted for 20 min, a maximum of six consecutive aggressive behaviors against subordinates

occurred within a single behavioral sequence. Instances where subordinates did not exhibit the

eye-patch pattern upon first aggression by dominants and subsequently maintained their facial

coloration without alteration (*A to A*) were observed 97 times. Instances where subordinates did

not exhibit the eye-patch pattern upon first aggression but did so after it (*A to P*) were observed

69 times. Instances where subordinates exhibited the eye-patch pattern upon first aggression and

subsequently altered their facial coloration to a non-eye-patch pattern (*P to A*) were observed 9

times. Finally, instances where subordinates exhibited the eye-patch pattern upon first

aggression and subsequently maintained their facial coloration without alteration were observed

23 times. It should be noted that in 20 cases of aggression, facial coloration could not be

discerned; therefore, these data were removed from subsequent analyses

Experimental manipulation of social signals

To investigate the potential function of facial coloration as a social signal, body coloration in *S.*

aequifasciatus was experimentally manipulated using a light treatment known among discus

fish enthusiasts that targets the vertical bars, which are believed to serve as social signals.

Specifically, this treatment aimed to eliminate the vertical bars by exposing juvenile fish to

intense illumination. Initially, mature females and males exhibiting partner displays (Satoh et al.

2016) were transferred to separate water tanks (122 L, 60 × 45 × 45 cm or 91 L, 45 × 45 × 45

cm). Substrate bricks suitable for spawning were provided in these tanks, and humic acid was

periodically added to regulate water quality. Upon spawning, the offspring remained with their

parents for 10 days, as early free-swimming fry rely on parental mucus for sustenance (Satoh et

al. 2017). Subsequently, 50 fry from the same brood were placed in two smaller tanks. One tank

received illumination from a light source positioned approximately 30 cm above, thereby

constituting the light-treated group, while the other tank remained under ambient room lighting,

serving as the control for wild-type individuals. The juvenile fish were fed *Artemia* spp. nauplii

during the experiment. The lighting regime for the light-treated group was synchronized with

the room lighting, maintaining a 12-hour light / 12-hour dark cycle. After 3 months, the

juveniles were identified and moved to the same water tanks (162 L, 60 × 45 × 60 cm or 182 L,

90 × 45 × 45 cm) until they reached sexual maturity, which typically occurs over a period of

approximately 1.5–2.0 years. Facial coloration was evaluated in the light-treated individuals in

the same manner as in *behavioral observation I*, but none of them exhibited an eye-patch

pattern (Supplementary S2).

Light-treated and wild-type individuals were used to test whether facial coloration

functions analogously to emotional expression. Three types of experimental pairs were created:

a group consisting of wild-type individuals only (WW pair, n = 4 male pairs and 4 female pairs),

a group of wild-type and light-treated individuals (WL pair, n = 3 male pairs and 3 female

pairs), and a group of light-treated individuals only (LL pair, n = 3 male pairs and 4 female

pairs). These same-sex experimental pairs (42 individuals in total) were selected to ensure

homogeneity in kinship and were allowed to meet each other for the first time. After measuring

their body weight in 1 mg, they were placed in a water tank (122 L, 60 × 45 × 45 cm) for 9

days. During the experiment, the fish were individually fed with five mosquito larvae per day

using tweezers. This was to ensure that the dominant individuals did not monopolize the food.

The number of aggressive interactions between pair partners over a period of 20 minutes was

counted on days 1, 3, 5, 7, and 9. In this study, aggressive interaction was defined as total

aggressive display and direct aggression occurring between pair partners. After 9 days of

observations, the body weights of the experimental pairs were measured again. Among the

examined groups, individuals exhibiting more aggressive behaviors toward their experimental

partners during the experiment were defined as dominant, while those receiving more attacks as

submissive. The frequency of aggressive interaction was predicted to be higher in the LL group

than in the WW and WL groups if facial coloration has a function in emotional signaling.

Tissue clearing and immunohistochemistry

Three bred discus fish *S. aequifasciatus* individuals (22.5–33 mm SL) were used for tissue

clearing and immunohistochemistry, primarily following the methods described in Pende et al.

(2020)[42]. The fish were deeply anesthetized with benzocaine, and their head and body were

sectioned along a plane extending from the base of the pelvic fin to approximately the third to

fifth spine of the dorsal fin. After removing the internal organs and creating small holes in the

cornea, the heads were fixed in 4% (w/v) paraformaldehyde at 4°C overnight. The specimens

were washed four times (30 minutes each) with PBS (pH 7.4) at room temperature and

incubated in 0.2 M EDTA (pH 8.0) at 4°C for 48 hours, with the solution refreshed after 24

hours. After another four PBS washes at room temperature, they were transferred to pre-chilled

acetone and incubated at 4°C for 1 hour. The acetone was then refreshed, and the specimens

were incubated overnight at −20°C, followed by another refreshment and incubation overnight

at room temperature. After being washed four times with PBS at room temperature, the

specimens were treated with 3% (w/w) H_2O_2 in 0.8% (w/v) KOH for about 160 minutes at the

same temperature under intense light exposure. Subsequently, they were washed with 1% (v/v)

Triton X-100 in PBS (PBST) for two nights at 4°C and placed under vacuum for 60 minutes to

remove gas bubbles. They were then incubated in Solution-1.1 (pH 11.5) for two nights at 37°C,

with the solution being refreshed after the first night. This solution consisted of 10% (v/v)

THEED (Sigma-Aldrich, 87600-100 ml), 5% (v/v) Triton X-100, and 5% (w/v) urea in dH$_2$O,

and its pH was adjusted to 11.5 using 30% (w/w) ammonia solution. After five PBST washes

(10 minutes each), the specimens were treated with 1% (w/v) bovine serum albumin (BSA;

Proliant Biologicals, PRL68700-25G) in PBST overnight at 4°C. Then, they were incubated in

1:1,000 primary antibody (rabbit anti-TH; Merck Millipore, AB152) in PBST with 0.5% BSA

for 4 days at 4°C, followed by [*add number*] PBST washes for 2 days at 4°C. Next, the

specimens were incubated in 1:200 biotinylated goat anti-rabbit IgG secondary antibodies

(Vector Laboratories, BA-1000) in PBST with 0.5% BSA for 4 days at 4°C, washed in PBST for

2 days at 4°C, and incubated in a mixture of 1:100 avidin and 1:100 biotinylated horseradish

peroxidase (Vector Laboratories, PK-6100) in PBST for 4 days at 4°C, followed by PBST

washes for 2 days at 4°C. The specimens were stained in a reaction solution containing 0.05%

(w/v) 3,3'-diaminobenzidine, 0.01% H$_2$O$_2$, 0.04% (w/v) nickel ammonium sulfate, 0.5% (v/v)

Triton X-100, and 0.1 M phosphate buffer (pH 7.4) in ultra-pure water, followed by [*add*

number] PBST washes for 1 day at room temperature. For refractive index matching, the three

specimens were incubated in CUBIC-R+(N) (45% [w/w] antipyrine, 30% [w/w] nicotinamide,

0.5% [v/v] N-butyldiethanolamine in ultra-pure water) (Matsumoto et al. 2019) for 3 days at

room temperature. The eye-patch regions were then observed and photographed under a

microscope fitted with a digital camera (Olympus, SZX10 with DP25).

Ex vivo administration of NA and ATP to scale melanophores

Three bred discus fish *S. aequifasciatus* individuals (21–24 mm SL) were anesthetized with

benzocaine, and a total of 20 scales (ATP exposure group, n = 5; control for ATP exposure

group, n = 5; NA exposure group, n = 5; control for NA exposure group, n = 5) were collected

from the eye-patch region and kept in ice-cold PBS for a short period until the experiment was

started. The scales were placed on a concave slide glass filled with PBS, secured by pinching

the edges with a polypropylene sheet, and observed under a microscope (Olympus, BX53). No

melanophores were present in the pinched areas. It was difficult to completely standardize the

physiological state of each individual at the time of scale collection and the neural innervation

status across all scales. Therefore, the ATP exposure group was pre-treated with NA, and the NA

exposure group was pre-treated with ATP, and changes in melanosome movement due to

neurotransmitter exposure administration were specifically quantified. After removing the PBS

from the concave slide, 0.1 mM ATP in PBS was added to the NA exposure group, and 0.1 μM

NA in PBS was added to the ATP exposure group, followed by incubation at room temperature

for 2 minutes. Subsequently, the solutions were removed, and the scales were washed by adding

and quickly removing PBS. A total of 1 μM NA in PBS or PBS was added to the NA exposure

group, and a total of 10 mM ATP in PBS or PBS was added to the ATP exposure group.

Considering the moment each solution was applied as 0 seconds, photographs were taken using

a microscope fitted with a camera (Olympus, BX53 with DP74) at 1 second and every 10

seconds after that for 120 seconds. All images were adjusted for white balance using Adobe

Photoshop (version 25.9.1) and analyzed using ImageJ (version 2.14.0.). To exclude

xanthophores, the images were split into the three primary colors, and each monochrome one

was used in the analysis. Subsequently, one well-captured melanophore (e.g., focus alignment)

was randomly selected per scale, and its area was set as the ROI. Then, the area showing a

grayscale above the identical threshold for all images within the ROI was calculated. The rate of

variation in melanosome dispersion (%) was calculated using the following formula:

$$Rate\ of\ change\ in\ melanosome\ dispersion$$

$$= \frac{Final\ extent\ of\ melanosome\ dispersion}{Initial\ extent\ of\ melanosome\ dispersion} \times 100\ (\%)$$

Statistical analysis

All statistical analyses were performed in R (v. 4.3.3) [77] using LMMs or GLMMs in the lme4

package [78].

To evaluate whether facial coloration in subordinate individuals mitigated harassment

by dominant individuals, the probability of continued harassment was analyzed as a binary

variable within a binomial GLMM, with experimental pair ID incorporated as a random effect.

The variation in facial coloration, specifically whether the facial pattern transitioned before or

after the initial aggression by dominants (A to A, A to P, P to A, or P to P), was included as an

explanatory variable. Given that changes in facial coloration were observed to significantly

affect the likelihood of continuous harassment, post-hoc testing was conducted using separate

binomial GLMMs, again treating experimental pair ID as a random effect. For these post-hoc

GLMMs, the significance level (α) was adjusted via Bonferroni correction to $\alpha = 0.0083$.

With regard to the experimental manipulation of facial coloration, a negative binomial

GLMM was employed to analyze the data, with the experimental pair ID included as a random

effect. This model examined the two-way interaction between the experimental groups (WW,

WL, and LL pairs) and observation days (1, 3, 5, 7, and 9) as explanatory variables.

Furthermore, the reduction in body weight during the experiment was analyzed using an LMM,

again incorporating experimental pair ID as a random effect. The response variable was the

percentage reduction in body weight (calculated as post-experiment body weight divided by

pre-experiment body weight). The explanatory variables included the experimental group (WW,

WL, and LL pairs), the sex of focal individuals, their status (light-treated or wild-type), and

their social rank (dominant or subordinate).

Finally, the rate of variation in melanosome dispersion was examined upon exposure

to ATP and NA, which act as sympathetic neurotransmitters. Prior to analysis, the black area in

the selected melanophore (measured in pixels) was log-transformed (log10) to achieve

normalization. An LMM was employed in this analysis, with individual ID included as a

random effect. The two-way interaction between time and treatment (either ATP or PBS control)

was modeled [*using?*] as an explanatory variable. An analogous analysis was conducted for the

NA experiment. It should be noted that the results of these analyses are presented in Fig. 5 as

the percentage change in melanosome dispersion relative to the baseline measurements at the

start of the experiment for visualization.

References

1. Rhodes, G., Simmons, L., Barrett, L. & Dunbar, R. Oxford Handbook of Evolutionary

Psychology (2007).

2. McKone, E., Kanwisher, N. & Duchaine, B. C. Can generic expertise explain special

processing for faces ? Trends Cogn. Sci. 11, 8–15 (2007)

3. Schmidt, K. L. & Cohn, J. F. Human facial expressions as adaptations: evolutionary questions

in facial expression research. Am. J. Phys. Anthropol. 116, 3–24 (2001).

4. Kobayashi, H. & Kohshima, S. Unique morphology of the human eye. Nature 387, 767–768

(1997).

5. Tomasello, M., Hare, B., Lehmann, H. & Call, J. Reliance on head versus eyes in the gaze

following of great apes and human infants: the cooperative eye hypothesis. J. Hum. Evol. 52,

314–320 (2007).

6. Rinn, W. E. The neuropsychology of facial expression: a review of the neurological and

psychological mechanisms for producing facial expressions. Psychol. Bull. 95, 52–77 (1984).

7. Cattaneo, L. & Pavesi, G. The facial motor system. Neurosci. Biobehav. Rev. 38, 135–159

(2014).

8. Müri, R. M. Cortical control of facial expression. J. Comp. Neurol. 524, 1578–1585 (2016).

9. Allison, T., Puce, A. & McCarthy, G. Social perception from visual cues: role of the STS

region. Trends Cogn. Sci. 4, 267–278 (2000).

10. Engell, A. D. & Haxby, J. V. Facial expression and gaze-direction in human superior

temporal sulcus. Neuropsychologia 45, 3234–3241 (2007).

11. Xu, P., Peng, S., Luo, Y. J. & Gong, G. Facial expression recognition: a meta-analytic review

of theoretical models and neuroimaging evidence. Neurosci. Biobehav. Rev. 127, 820–836

(2021).

12. Hadders-Algra, M. Human face and gaze perception is highly context specific and involves

bottom-up and top-down neural processing. Neurosci. Biobehav. Rev. 132, 304–323 (2022).

13. Ekman, P. & Friesen, W. V. Constants across cultures in the face and emotion. J. Pers. Soc.

Psychol. 17, 124–129 (1971).

14. Boakes, R. From Darwin to Behaviourism: Psychology and the Minds of Animals

(Cambridge Univ. Press, New York, 1984).

15. Ekman, P. Darwin and Facial Expression: A Century of Research (2006).

16. Darwin, C. The Expression of the Emotions in Man and Animals (1872).

17. Parr, L. A., Waller, B. M. & Heintz, M. Facial expression categorization by chimpanzees

using standardized stimuli. Emotion 8, 216–231 (2008).

18. Vick, S. J., Waller, B. M., Parr, L. A., Smith Pasqualini, M. C. & Bard, K. A. A cross-species

comparison of facial morphology and movement in humans and chimpanzees using the facial

action coding system (FACS). J. Nonverbal Behav. 31, 1–20 (2007).

19. Leopold, D. A. & Rhodes, G. A comparative view of face perception. J. Comp. Psychol.

124, 233–251 (2010).

20. Langford, D. J. et al. Coding of facial expressions of pain in the laboratory mouse. Nat.

Methods 7, 447–449 (2010).

21. Bertin, A. et al. Captive Blue-and-yellow macaws (*Ara ararauna*) show facial indicators of

positive affect when reunited with their caregiver. Behav. Processes 206, 104833 (2023).

22. Tate, A. J., Fischer, H., Leigh, A. E. & Kendrick, K. M. Behavioural and neurophysiological

evidence for face identity and face emotion processing in animals. Philos. Trans. R. Soc. Lond.

B Biol. Sci. 361, 2155–2172 (2006).

23. Demaria, C. & Thierry, B. Responses to animal stimulus photographs in stumptailed

macaques (*Macaca arctoides*). Primates 29, 237–244 (1988).

24. Guo, K., Robertson, R. G., Mahmoodi, S., Tadmor, Y. & Young, M. P. How do monkeys

view faces?–A study of eye movements. Exp. Brain Res. 150, 363–374 (2003)

25. Satoh, S., Tanaka, H. & Kohda, M. Facial recognition in a discus fish (Cichlidae):

experimental approach using digital models. PLOS ONE 11, e0154543 (2016).

26. Siebeck, U. E., Parker, A. N., Sprenger, D., Mäthger, L. M. & Wallis, G. A species of reef

fish that uses ultraviolet patterns for covert face recognition. Curr. Biol. 20, 407–410 (2010).

27. Kohda, M. et al. Facial recognition in a group-living cichlid fish. PLOS ONE 10, e0142552

(2015).

28. Wang, M. Y. & Takeuchi, H. Individual recognition and the 'face inversion effect' in medaka

fish (Oryzias latipes). Elife, 6, e24728 (2017)

29. Saeki, T., Sogawa, S., Hotta, T. & Kohda, M. Territorial fish distinguish familiar neighbours

individually. Behaviour 155 , 279–293 (2018).

30. Hotta, T., Kawasaka, K., Satoh, S. & Kohda, M. Fish focus primarily on the faces of other

fish. Sci. Rep. 9, 8377 (2019).

31. Sogawa, S. et al. Male guppies recognize familiar conspecific males by their face. Zool. Sci.

40, 168–174 (2023).

32. Kawasaka, K., Hotta, T. & Kohda, M. Does a cichlid fish process face holistically?

Evidence of the face inversion effect. Anim. Cogn. 22, 153–162 (2019).

33. Kohda, M. et al. Cleaner fish recognize self in a mirror via self-face recognition like

humans. Proc. Natl Acad. Sci. U. S. A. 120, e2208420120 (2023).

34. Kano, F. & Tomonaga, M. How chimpanzees look at pictures: a comparative eye-tracking

study. Proc. Biol. Sci. 276, 1949–1955 (2009).

35. Farias, I. P. & Hrbek, T. Patterns of diversification in the discus fishes (*Symphysodon* spp.

Cichlidae) of the Amazon Basin. Mol. Phylogenet. Evol. 49, 32–43 (2008).

36. Crampton, W. G. Ecology and life history of an Amazon floodplain cichlid: the discus fish

Symphysodon (Perciformes: Cichlidae). Neotrop. Ichthyol. 6, 599–612 (2008).

37. Satoh, S., Tanoue, H., & Mohri, M. Costs and benefits of biparental mucus provisioning in

discus fish (*Symphysodon aequifasciatus*). Ichthyol Res, 65, 510-514 (2018)

38. Satoh, S., & Sowersby, W. Mucus provisioning behavior in teleost fishes: a novel model

system for the evolution of secretory provisioning in vertebrates. Ichthyol Res, 68, 1-10 (2021)

39. Satoh, S., Tanoue, H., Ruitton, S., Mohri, M., & Komatsu, T. Morphological and behavioral

ontogeny in larval and early juvenile discus fish *Symphysodon aequifasciatus*. Ichthyol Res, 64,

37-44 (2017)

40. Muske, L. E. & Fernald, R. D. Control of a teleost social signal. II. Anatomical and

physiological specializations of chromatophores. J. Comp. Physiol. A 160, 99–107 (1987).

41. Bleher, H. Bleher's Discus (Aquapress, Germany, 2006).

42. Pende, M. et al. A versatile depigmentation, clearing, and labeling method for exploring

nervous system diversity. Sci. Adv. 6, eaba0365 (2020).

43. Balzarini, V., Taborsky, M., Villa, F., & Frommen, J. G. Computer animations of color

markings reveal the function of visual threat signals in *Neolamprologus pulcher*. Curr Zool, 63(,

45-54 (2017)

44. Muske, L. E. & Fernald, R. D. Control of a teleost social signal. I. Neural basis for

differential expression of a color pattern. J. Comp. Physiol. A 160, 89–97 (1987).

45. Rodrigues, R. R., Carvalho, L. N., Zuanon, J., & Del-Claro, K. Color changing and

behavioral context in the Amazonian Dwarf Cichlid *Apistogramma hippolytae* (Perciformes).

Neot Ichthyo, 7, 641-646 (2009)

46. Seikai, T. Influences of fluorescent light irradiation, ocular side pigmentation, and source of

fishes on the blind side pigmentation in the young Japanese flounder, Paralichthys olivaceus.

Suisanzoshoku 39, 173–180 (1991).

47. Andersson, M. Sexual Selection (Princeton Univ. Press, 1994).

48. Krause, J. & Ruxton, G. D. Living in Groups (Oxford Univ. Press, Oxford, 2002).

49. Fujii, R. & Oshima, N. Control of chromatophore movements in teleost fishes. Zool. Sci. 3,

13–47 (1986).

50. Fujii, R. The regulation of motile activity in fish chromatophores. Pigment Cell Res. 13,

300–319 (2000).

51. Sugimoto, M. Morphological color changes in fish: regulation of pigment cell density and

morphology. Microsc. Res. Tech. 58, 496–503 (2002).

52. von Frisch, K. Beiträge zur physiologie der pigmentzellen in der fischhaut. Pflüger's Arch.

138, 319–387 (1911).

53. Fujii, R. & Novales, R. R. The nervous mechanism controlling pigment aggregation in

Fundulus melanophores. Comp. Biochem. Physiol 29, 109–124 (1969).

54. Funakoshi, K. & Nakano, M. The sympathetic nervous system of anamniotes. Brain Behav.

Evol. 69, 105–113 (2007).

55. Nilsson Sköld, H., Aspengren, S. & Wallin, M. Rapid color change in fish and amphibians –

function, regulation, and emerging applications. Pigment Cell Melanoma Res. 26, 29–38 (2013).

56. Konno, A., Matsumoto, N., Tomono, Y. & Okazaki, S. Pathological application of

carbocyanine dye-based multicolour imaging of vasculature and associated structures. Sci. Rep.

10, 12613 (2020).

57. Ueda, H. R. et al. Tissue clearing and its applications in neuroscience. Nat. Rev. Neurosci.

21, 61–79 (2020).

58. Matsumoto, K. et al. Advanced CUBIC tissue clearing for whole-organ cell profiling. Nat.

Protoc. 14, 3506–3537 (2019).

59. Yew, D. T., Webb, S. E. & Lam, E. T. Neurotransmitters and peptides in the developing

human facial nucleus. Neurosci. Lett. 206, 65–68 (1996)

60. Leong, C. Y. The quantitative effect of releasers on the attack readiness of the fish

Haplochromis burtoni (Cichlidae, Pisces). Z Vergl Physiol. 65, 29–50 (1969).

61. Heiligenberg, W., Kramer, U. & Schulz, V. The angular orientation of the black eye-bar in

Haplochromis burtoni (Cichlidae, Pisces) and its relevance to aggressivity. Z Vergl Physiol. 76,

168–176 (1972).

62. Schmidt, K. L. & Cohn, J. F. Human facial expressions as adaptations: evolutionary

questions in facial expression research. Am. J. Phys. Anthropol. 116, 3–24 (2001).

63. Straulino, E., Scarpazza, C. & Sartori, L. What is missing in the study of emotion

expression? Front. Psychol. 14, 1158136 (2023).

64. Bshary, R. Biting cleaner fish use altruism to deceive image–scoring client reef fish. Proc.

Biol. Sci. 269, 2087–2093 (2002).

65. Fernald, R. D. Decision-making in fish: genetics and social behaviour. In Dreher, J. C. &

Tremblay, L. (eds.), Decision Neuroscience: An Integrative Perspective 351–359 (Academic

Press, San Diego, 2017).

66. de Melo, C. M. & Gratch, J. Expression of emotions using wrinkles, blushing, sweating and

tears. In: Intelligent Virtual Agents: International Conference, IVA 2009 Amsterdam, The

Netherlands, September 14-16 (Springer Berlin Heidelberg, 2009).

67. Lee, D. H., Susskind, J. M. & Anderson, A. K. Social transmission of the sensory benefits of

eye widening in fear expressions. Psychol. Sci. 24, 957–965 (2013).

68. Ruxton, G. D., Allen, W. L., Sherratt, T. N. & Speed, M. P. Avoiding Attack: The

Evolutionary Ecology of Crypsis, Aposematism, and Mimicry (Oxford Univ. Press, 2019).

69. Wendelaar Bonga, S. E. The stress response in fish. Physiol. Rev. 77, 591–625 (1997).

70. Barton, B. A. Stress in fishes: a diversity of responses with particular reference to changes in

circulating corticosteroids. Integr. Comp. Biol. 42, 517–525 (2002).

71. Madaro, A. et al. How fish cope with stress?. In: Kristiansen, T., Fernö, A., Pavlidis, M. van

de Vis. H. (eds.), The Welfare if Fish (Springer Nature, Switzerland, 2020).

72. Fox, H. E., White, S. A., Kao, M. H. & Fernald, R. D. Stress and dominance in a social fish.

J. Neurosci. 17, 6463–6469 (1997).

73. Sloman, K. A., Metcalfe, N. B., Taylor, A. C. & Gilmour, K. M. Plasma cortisol

concentrations before and after social stress in rainbow trout and brown trout. Physiol.

Biochem. Zool. 74, 383–389 (2001).

74. Schjolden, J. & Winberg, S. Genetically determined variation in stress responsiveness in

rainbow trout: behavior and neurobiology. Brain Behav. Evol. 70, 227–238 (2007).

75. Denil, N. A., Yamashita, E., Kirino, M. & Kiyohara, S. Recurrent facial taste neurons of sea

catfish *Plotosus japonicus*: morphology and organization in the ganglion. J. Fish Biol. 82,

1773–1788 (2013).

76. Albohn, D. N. & Adams Jr, R. B. The social face hypothesis. Affect. Sci. 3, 539–545 (2022).

77. R Core Team. R: a language and environment for statistical computing. Version 4.3.3.

https://www.R-project.org/ (2023).

78. Bates, D., Maechler, M. & Bolker, B. lme4: linear mixed effects models using S4 classes.

http://CRAN.R-project.org/package=lme4 (2011).

79. Satoh, S., et al. Data from "Facial coloration and behaviors of discus fish". Figshare (___)

Acknowledgments

We thank the members of the Animal Sociology Laboratory of Osaka City University and the

Kutsukake Research Group of Graduate University for Advanced Studies for their helpful

comments. This study was supported financially by HAKUBI Project and KAKENHI (nos.

N19K23765, N19KK0189, 20J01170, and 23KK0131 to S.S; nos. 23H03868 to S.S. and K.F.).

Author contribution

S.S., K.F., A.K., S.T., M.K., and N.K. designed the study. S.S., K.F., S.M., T.B., and K.K.

collected the data. S.S. and K.F. conducted the analyses. S.S. and K.F. wrote the manuscript

with input from all the authors.

Competing interests

The authors declare no competing interests.

Ethics statement

All the experimental protocols were approved by the Animal Care and Use Committees at

Osaka City University and Kitazato University and adhered to the ASAB/ABS guidelines for

the treatment of animals in behavioral research.

Data availability

The data that supports the findings of this study are available in the figshare repository XXX [79].

Figure 1 Facial coloration in discus fish *Symphysodon aequifasciatus*. Typical chronological

changes of eye-patch pattern (a) and coloration associated with social behaviors (b). The bars

represent the possibility of coloration patterns emerging when individuals engage in specific

social behaviors. Three identifiable coloration patterns were detected: 1) absence of any vertical

black bar composed of melanophores; 2) presence of only one bar consisting of melanophores,

eye-patch pattern, and 3) presence of multiple bars throughout the body.

Figure 2 Effects of facial coloration in *S. aequifasciatus* subordinates on the continuous aggressive harassment by dominant individuals. The data are presented as mean values along with their corresponding 95% confidence intervals, and the sample sizes are indicated above the bars. Significant differences were determined using post-hoc binomial GLMMs, with p-values adjusted using Bonferroni correction (α level = 0.0083). The possibility of experiencing continuous harassment was notably higher in *A to A* than in *A to P* (df = 1, χ^2 = 54.186, p < 0.001) and *P to P* (df = 1, χ^2 = 10.05, p = 0.002), but similar to that of *P to A* (df = 1, χ^2 = 2.093, p = 0.148). Conversely, the probability of continuous harassment was significantly lower in *A to P* than in *P to A* (df = 1, χ^2 = 22.076, p < 0.001), but similar to that of *P to P* (df = 1, χ^2 = 3.3123, p = 0.0688). Additionally, there was a significant difference in the possibility of continuous harassment between *P to A* and *P to P* (df = 1, χ^2 = 9.690, p = 0.002).

Figure 3 Effects of the experimental manipulation of melanophores on aggressive social

interaction in *S. aequifasciatus*. The figure shows the relationship between the frequency of

aggressive interaction within experimental groups and days since the start of the experiment.

The experimental groups, consisting of wild-type × wild-type individuals (WW), wild-type ×

light-treated individuals (WL), and light-treated × light-treated individuals (LL), are indicated

by black, white, and gray bars, respectively. A significant interaction was detected between the

type of experimental group and the duration of the experiment (negative binomial GLMM, df =

2, χ^2 = 67.896, p < 0.001). Data are presented as mean values ± standard deviation (SD).

Figure 4 Reduction in body weight during the 9-day experiment in dominant and

subordinate *S. aequifasciatus*. The experimental groups, consisting of wild-type × wild-type

individuals (WW), wild-type × light-treated individuals (WL), and light-treated × light-treated

individuals (LL), are indicated by black, white, and gray bars, respectively. Significant

differences were detected in the percentage of reduction in body weight between experimental

groups (LMM, df = 2, F = 4.398, p = 0.025), while marginal differences were observed in

dominance rank (df = 1, F = 3.955, p = 0.061). Data are presented as mean values ± standard

deviation (SD).

Figure 5 Effect of the noradrenaline (NA) and adenosine triphosphate (ATP)

neurotransmitters used by the sympathetic postganglionic neurons on melanophores

constituting the eye patch and tyrosine hydroxylase-immunoreactive projection to the eye-

patch region. The peripheral nerves regulating the melanophores constituting the eye patch

were examined. **a–d** Temporal variations in the extent of melanosome dispersion upon the

exposure of melanophores constituting the eye patch to NA and ATP . The rate of variation in

melanosome dispersion upon exposure to ATP (a, c) and NA (b, d) was significantly different

from that observed in the PBS control (two-way interaction between time and treatment, both p

< 0.001). Data are presented as mean values ± standard error. **e–g** The tissues were cleared and

immunostained using an antibody against tyrosine hydroxylase (TH), revealing that TH-

immunoreactive (TH-ir) projections extended to the eye-patch region (white arrowheads). The

white dotted lines indicate the eye-patch region (**e**). The observations strongly suggested that

TH-ir fibers terminate at the scales possessing melanophores forming the eye patch: (**b**)

specimen whose scales had been removed to improve the visibility of the TH-ir fibers and **(e)**

specimen with intact scales.

 Supplementary

Supplemental movie S1 Typical example of facial coloration transition in the discus fish

Symphysodon aequifasciatus. To facilitate the viewing of changes in facial coloration patterns,

we presented a model of the reproductive partner on the monitor and observed its behavior

based on the method described in Satoh et al. (2016) [25].

Supplemental Figure S2 Coloration associated with social behaviors in light-treated *S.*

aequifasciatus. The bars represent the possibility of coloration patterns emerging when

individuals engage in specific social behaviors. Three identifiable coloration patterns were

detected: 1) absence of any vertical black bar composed of melanophores; 2) presence of only

one bar consisting of melanophores, eye-patch pattern, and 3) presence of multiple bars

throughout the body. These observations were made in the same manner as in *behavioral*

observation I (see Methods).