Development and validation of an open-source Hand Laterality Judgement Task for in-person and online studies

Marcos Moreno-Verdú[1]*, Siobhán M McAteer[1], Baptiste M Waltzing[1], Elise Van Caenegem[1], Robert M Hardwick[1]

[1] Brain, Action, and Skill Laboratory (BAS-Lab), Institute of Neuroscience (Cognition and Systems Division), UCLouvain, Belgium

ABSTRACT

The Hand Laterality Judgement Task (HLJT) is considered a measure of the ability to manipulate motor images. The 'biomechanical constraints' effect (longer reaction times for hand rotations towards anatomically difficult versus biomechanically easier movements) is considered the behavioural hallmark indicating motor imagery is being used. Previous work has used diverse HLJT paradigms, and there is no standardized procedure for the task. We developed an open-source, freely available version of the HLJT in PsychoPy2, which needs no programming skills and is highly customisable. Some studies suggest responding to the HLJT with the hands may interfere with performance, which would limit practical application of the task. We examined this potential issue using in-person and online versions. For the in-person version, 40 right-footed/handed individuals performed the HLJT with their feet or bimanually (N=20 each). For the online version, 60 right-handed individuals performed the task bimanually or unimanually (N=20 each). Bayesian mixed-effect analyses quantified the evidence for and against equivalence within and between the in-person and online versions. Both versions replicated previously described behavioural phenomena, including effects of angle, hand view, and the 'biomechanical constraints' effect. While responding with different effectors modified overall reaction times, it did not interact with other factors analysed, and did not affect accuracy or the 'biomechanical constraints' effect. There was also evidence for equivalence between in-person and online bimanual groups for all measures. We conclude that this open-source, standardized HLJT protocol (available at https://osf.io/8h7ec/) can reliably detect previously identified effects and works equally well in-person or online.

Key words: hand laterality; hand rotation; motor imagery; mental rotation; response mode; Bayesian inference.

Corresponding Author:
Dr. Marcos Moreno-Verdú
Brain, Action and Skill Laboratory (BAS-Lab), Institute of Neuroscience (Cognition and Systems Division), UCLouvain, Belgium
Address: Avenue Mounier 54, +2, 0430, 1200 Woluwe-Saint-Lambert, Belgium
Email: marcos.morenoverdu@uclouvain.be

INTRODUCTION

Motor imagery is a topic of increasingly growing scientific interest. It can be defined as the mental simulation of a movement without actual physical execution (Jeannerod, 2001). Motor imagery is relevant for a variety of purposes, from fundamental neuroscientific research to applied fields in sports and rehabilitation (Ladda et al., 2021; Zhao et al., 2023). Evidence indicates that performing motor imagery is subject to inter-individual variability (Floridou et al., 2022; Guillot et al., 2008) and depends on different imagery subprocesses (Collet et al., 2011), which has led to the development of 'motor imagery ability' measures (Suica et al., 2022). At the behavioural level, these assessments aim to evaluate the individual's ability to generate, maintain and manipulate motor imagery (Kraeutner et al., 2020).

The Hand Laterality Judgement Task (HLJT) has been extensively used as a measure of the ability to manipulate motor images (Parsons, 1987). In the task, individuals judge whether hand images rotated to different angles correspond to the left or right side of the body. Theoretically, most individuals will involuntarily use motor imagery to solve the task (Conson et al., 2020). That is, they will inadvertently simulate (i.e., imagine) rotating their own hand to decide the laterality. The fact that this behaviour appears unintentionally is one of the argued strengths of the HLJT over other imagery ability measures (e.g., questionnaires for imagery generation or mental chronometry for imagery maintenance), which rely on the individual's capacity of voluntarily performing motor imagery and subjectively reporting it afterwards (Suica et al., 2022; Williams et al., 2015). The HLJT assesses so-called *implicit* motor imagery ability (McAvinue & Robertson, 2008).

Research has generally suggested that one behavioural hallmark of the HLJT may indicate whether individuals are using motor imagery to solve it: the 'biomechanical constraints' effect (Parsons, 1994). According to this effect, the reaction time needed to judge laterally rotated stimuli would be greater than medially rotated stimuli considering a rotation in the frontal axis, a phenomenon explained by the inherent anatomical limitations of the actual movement. While the literature is not in complete agreement with this effect solely and uniquely reflecting motor imagery-based processing (Meng et al., 2016; Vannuscorps & Caramazza, 2016), it has been widely employed as an indication of motor imagery. This is partially because the effect only appears in the HLJT as opposed to other mental rotation tasks, such as letter rotation tasks (Bek et al., 2022, p. 20; Mibu et al., 2020), which suggest the task elicits a different cognitive strategy, probably based on egocentric processing.

While the HLJT has been widely used in psychological, neuroscientific, and clinical research, it has been substantially less used in applied contexts such as rehabilitation, sports science, or online studies. Moreover, there is no standardized version of the task, and previous studies have used a wide variety of experimental paradigms (e.g. different angles of rotation, views of the hand, etc.). Furthermore, researchers have generally implemented this task in software which are typically not prepared for running both in-person and online studies. New developments in open-source stimulus presentation software can overcome both issues while permitting the experimental setup to be flexible.

One of the peculiarities of the HLJT, which might explain its limited use in applied circumstances and online studies is that if participants respond using their own hands, a possible confound with task performance might arise, in the form of an interference or priming effect (Cocksworth & Punt, 2013). This may occur because the same effector is being used to process the stimulus and provide a response, potentially modifying how information is processed while the response is being prepared. Thus, verbal (Ionta et al., 2007) or foot (Brady et al., 2011) responses have been used to avoid this potential issue. We note, however, that studies which have used the HLJT with manual responses have broadly replicated the main behavioural phenomena of this paradigm, including the presence of the 'biomechanical constraints' effect (Bek et al., 2022; Hudson et al., 2006; Kraeutner et al., 2019; Saimpont et al., 2009). In fact, the assertion that hand response modes modify behaviour in the HLJT has only been directly tested in one previous study (Cocksworth & Punt, 2013). Further evidence in this regard is necessary to expand the applicability of the task to non-laboratory settings, where the individual could potentially need to respond with one or two hands depending on the specific situation (e.g., in case of unilateral motor disturbances or verbal communication disorders, or online studies).

The present study therefore had two central aims. Our first aim was to develop a standardized version of the HLJT as open-source software that is freely available. We publicly share both local and online running versions of the paradigm we used for this study (https://osf.io/8h7ec/). The task allows a high level of customizability and does not require extensive programming skills to be set up, with the ultimate goal of paving the way for its use in non-laboratory contexts. We 'validated' the in-person and online versions against each other using Bayesian analyses and equivalence tests, and investigated whether both versions could replicate the well-established behavioural phenomena in this task. The second goal was to test the hypothesis that manual responses interfere with performance in the HLJT, as this would facilitate future use in applied or online contexts. In the in-person version we compared a bimanual response mode against a foot response mode in terms of accuracy, reaction time and, critically, the 'biomechanical constraints' effect. We predicted that manual responses could modify task performance in terms of accuracy or reaction time, but not the biomechanical effect. Importantly, even in the presence of small discrepancies between response modes, we posited that meaningful differences would not emerge, and that responding with the hands would be practically equivalent than responding with the feet. In the online version, we compared the equivalence between bimanual and unimanual response modes, as the latter may be necessary for clinical applications.

METHODS

Study design and participants

This study used a mixed design, comparing measurements within- and between-subjects and in-person or online versions of the HLJT. All study procedures followed the Declaration of Helsinki (2013 revision) and were approved by the Ethics

Committee of the Institute for Research in the Psychological Sciences, UCLouvain
(N°: 2024-21). All participants were financially compensated for their time (€10/h).

Overall, 100 right-handed healthy individuals aged 18-35 years, with normal or
corrected-to-normal vision and no history of neurological damage participated.
Participants were identified as right-handed by self-assessment. All participants were
assessed with the Movement Imagery Questionnaire-3 (MIQ-3) to determine the
ability to generate motor imagery as total, kinesthetic, internal visual and external
visual imagery scores (Williams et al., 2012).

<u>Setting and participants in-person version:</u>

Forty right-handed individuals participated (see Table 1 for demographics). All
participants were also self-identified as right-footed by answering the question "If you
were to kick a ball would you do it with your right leg?", as previous work has
established this as an appropriate way to assess leg dominance (van Melick et al.,
2017). Participants were recruited via public advertisements, and they were assessed
at UCLouvain (Belgium). Participants were randomly allocated to 2 different groups
which only differed in the Response Mode. A computer random sequence generator
with 1:1 assignment was used. The "Foot" group (N = 20) responded to the HLJT
with their feet, whereas the "Bimanual" group (N = 20) responded with their hands
(see Response Modes for details).

<u>Setting and participants online version:</u>

Sixty right-handed individuals participated (see Table 1 for demographics). In order to
verify the task could be completed online, the experiment was conducted fully
remotely, and participants were recruited via the crowdsourcing platform Prolific
(https://www.prolific.com/). Participants were randomly allocated to 3 different groups
(N = 20 each) that completed the task in either a "Left Hand", "Bimanual" or "Right
Hand" condition (see Response Modes for details).

Table 1. Participants' characteristics for the in-person and online Hand Laterality Judgement Task versions according to their groups, including the Bayes Factors in favour of the null hypothesis of no differences between the groups (BF_{01}).

Characteristic	In-person version			Online version			
	Foot (N = 20)	Bimanual (N = 20)	Bayes Factor for null (BF_{01})	Left-Hand (N = 20)	Bimanual (N = 20)	Right-Hand (N = 20)	Bayes Factor for null (BF_{01})
Age, mean ± SD	24.80 ± 3.65	24.25 ± 3.21	2.92	26.20 ± 4.97	25.70 ± 3.63	26.05 ± 3.82 [1]	6.94
MIQ-3, mean ± SD							
Total score	68.40 ± 8.91	66.95 ± 7.44	2.86	61.30 ± 9.89	63.05 ± 11.75	61.79 ± 12.17	6.62
Kinesthetic	21.85 ± 3.67	22.05 ± 4.15	3.21	20.20 ± 4.26	20.90 ± 4.33	21.10 ± 3.93	6.15
Internal Visual	22.65 ± 3.80	22.30 ± 3.99	3.14	20.65 ± 3.88	21.05 ± 4.22	19.84 ± 4.68 [2]	5.46
External Visual	23.90 ± 2.94	22.60 ± 3.55	1.73	20.45 ± 3.98	21.26 ± 3.98 [3]	20.95 ± 4.25 [3]	6.26
Sex, n (%)			1.18				6.29
Female	12 (60%)	16 (80%)	-	9 (45%)	11 (55%)	9 (45%)	-
Male	8 (40%)	4 (20%)	-	11 (55%)	9 (45%)	11 (55%)	-

[1] Data showed from N=19 participants. [2] Data showed from N=19 participants. [3] Data showed from N=19 participants.

Note: BFs are interpreted following established benchmarks, considering 'no' evidence, 'anecdotal', 'moderate', 'strong', 'very strong' and 'extreme' evidence if the BF was ≈1, 1-3, 3-10, 10-30, 30-100 and >100, respectively (Andraszewicz et al., 2015; Jeffreys, 1998). Bayes factors are presented in relation to the null (BF_{01}) hypothesis.

General procedure, task, and stimuli

General procedure:

Experimental procedure is shown in Fig. 1A and Fig. 1B and was common for both versions. First, participants were assessed using an electronic version of the MIQ-3. The cross-culturally adapted English, Spanish or French versions were used depending on the native language of the participant (Robin et al., 2020; Trapero-Asenjo et al., 2021). Participants completed the questionnaire on their own. In the in-person version, the experimenter clarified any doubts or corrected the participants only if they asked or if they did not perform the movements as intended (e.g., if they performed a completely different movement or did the movement with the contralateral limb). In the online version, no clarifications were made.

Participants were assessed by the HLJT, whereby they judged whether images of hands that had been rotated depicted a right or left hand. Two experiments were created and run in PsychoPy2 version 2023.2.3 (Peirce et al., 2019). A 'local' experiment was used for the in-person version, whereas for the online version this was output to a PsychoJS experiment to be run online via PsychoPy2's associated webpage Pavlovia (https://pavlovia.org/) (Bridges et al., 2020). The files to run both experiments are freely accessible at https://osf.io/8h7ec/.

All participants were instructed to respond as quickly and accurately as possible through standardized on-screen instructions. The instructions were available in English, French or Spanish and had been reviewed by two native speakers in each case to ensure equivalence. In the in-person version, participants sat at a comfortable distance to the experiment computer (a 14-inch Dell laptop, 16gb RAM, 13th Gen i5-1345U, 1.6GHz, refresh rate = 60Hz) in a quiet room. In the online version, participants completed the experiment on their own computer, and were instructed to complete the task in a quiet room without distractions.

Task and experimental stimuli:

The task and experimental stimuli were identical in both versions. The experimental stimuli were real hand pictures as shown in Fig. 1C. Images of left hands were right-hand images that had been mirror reversed. Left or right images were presented in 8 rotational angles in the frontal axis (0°, 45°, 90°, 135°, 180°, 225°, 270°, 315° for right hands in the clockwise direction, and the opposite for left hands), and 2 possible views (palmar or dorsal). In total, 32 unique stimuli were used.

Each stimulus was preceded by a small fixation cross in the centre of the screen for 800ms. Stimuli were presented in the centre (PsychoPy dimensions: 0.45 x 0.45 units) and remained until a response was made. Visual feedback following each stimulus was provided for 300ms via two small boxes (PsychoPy dimensions: 0.07 x 0.07 units) placed at the bottom of the screen; the corresponding box turned green if a response was correct, and red if incorrect (Fig. 1B). Participants were allowed to familiarise themselves with the task and stimuli in a practice block with 1 repetition per unique stimulus (i.e., 32 trials) followed by 4 test blocks with 96 trials each (3 repetitions per unique stimulus). In each test block, stimuli were randomly presented in sub-blocks of 32 trials with 1 repetition per unique stimulus in each sub-block, to

minimize the likelihood of the same stimulus appearing more than twice in a row.
Only test blocks were used for the analyses, resulting in a total of N = 384 trials per
participant. Breaks between blocks were allowed, with a minimum duration of 10
seconds and no maximum limit.

Figure 1. Experiment overview. **Panel A** shows the overall structure of the experiment.
Participants started by completing the electronic version of the Movement Imagery
Questionnaire-3 (MIQ-3) before doing the Hand Laterality Judgement Task (HLJT), which
consisted in 1 practice block and 4 test blocks. **Panel B** shows the experimental stimuli used
Laterality (right/left) and rotational angle (0° to 315° in increments of 45°) were used to code
medial and lateral directions according to the anatomical position, which were later used for
the 'biomechanical constraints' effect analysis. **Panel C** shows an example of trials from the
HLJT, with visual feedback (green for correct responses, red for incorrect responses).
Panels D and E show all the possible Response Modes used in in-person (panel D) and
online (panel E) versions of the HLJT. The Foot and Bimanual groups used the keys 'S' and
'L' of the keyboard (left and right respectively). The Foot group responded on an external
keyboard placed on the floor from which all keys but 'S' and 'L' had been removed. The
Bimanual group used the index finger of their corresponding hand on the same external
keyboard placed on the desk. The Bimanual group in the online version used the same
paradigm as in-person, but on the computer's keyboard. The Left-hand and Right-hand
groups used the index and middle fingers placed over the 'G' and 'H' keys (left and right
respectively) on the computer's keyboard.

Response Modes across versions

Participants in both versions of the HLJT responded with one of the Response Modes depicted in Fig. 1D-E. All participants were given standardized, on-screen instructions which differed only according to the corresponding Response Mode.

In the in-person version (Fig. 1D), participant responses were recorded using an external QWERTY USB-keyboard (dell). All buttons were removed from this keyboard apart from the 'S' and 'L' keys, which were used to provide responses to left- and right-hand stimuli, respectively. This procedure allowed us to record both hand and foot responses with the same device. For the "Foot" Response Mode the keyboard was placed on the floor inside a custom-built wooden frame to help maintain its position. Participants sat at a desk and placed their left foot in contact with the 'S' key, and their right foot in contact with the 'L' key. They were instructed to maintain the position of the feet as stable as possible, with the heels in contact with the wooden frame, and did not wear shoes. Participants were allowed to adjust the positioning of their legs to optimise comfort, although their knees were always flexed at approximately 90°. They were asked to place their hands over their thighs and under the desk (i.e., without seeing them), and asked not to move them while performing the task. In the "Bimanual" mode, the keyboard was placed on the surface of the desk, and participants sat with their left index finger in contact with the 'S' key, and their right index finger in contact with the 'L' key. As such, participants could see their hands during the task.

In the online version (Fig. 1E), participants were instructed to complete the task either unimanually (with only the left or right hand), or bimanually. Participants in the "Bimanual" mode responded as in the in-person version, but on their own computer's keyboard. In the "Left-hand" and "Right-hand" Response Modes (only online), participants were instructed to respond with the index and middle fingers of the respective hand on their computer's keyboard. The keys in these two modes were the same and were centred on the keyboard ('G' for left images and 'H' for right images) to maintain relative spatial congruence in the required responses(Waltzing et al., 2024).

Statistical analysis

All analyses were performed in R version 4.3.3 (R Core Team 2024). Scripts and data are freely available at https://osf.io/8h7ec/. As a large part of the predictions for this study were focused on considering the evidence in favour of the null hypotheses or practically equivalent differences, a Bayesian framework was used. This allowed us to obtain evidence for and against the null/alternative hypothesis. Therefore, all primary results are reported from the posterior distribution as means and 95% Credible Intervals (95%CrI). Additionally, descriptive statistics from sample data (mean and SD) are presented in Supplementary Materials.

Accuracy and reaction time were analysed separately. First, trials where reaction time was <300ms or >3,000ms, likely reflecting anticipatory responses and no engagement with the task, respectively, were discarded. For accuracy, the proportion

of correct responses for each unique stimulus for each participant, after averaging
across repetitions, was obtained. For reaction time, the mean time for each unique
stimulus was considered only for the trials with correct responses. The measure of
the 'biomechanical constraints' effect was the difference between the mean reaction
time of medially vs. laterally rotated stimuli (Fig. 1C).

Separate models for the in-person and online datasets were first created and
analysed. Then, data from the two Bimanual groups (i.e., in-person and online) were
combined and analysed, to further validate the equivalence of online version of the
task with its in-person counterpart.

Statistical modelling was performed with multilevel models (i.e., mixed-effects
models), considering by-participant random intercepts using the 'brms' package
(Bürkner, 2017). These models were applied with a Gaussian identity likelihood for
reaction time and a zero-and-one inflated Beta likelihood for accuracy (as it allowed
the proportion of correct responses to be in the range 0-1 *including* both 0s and 1s in
the distribution) (Liu & Eugenio, 2018). The full model was always of type *'y ~*
Group*Angle*Laterality*View + (1|Participant)', where Group was a between-subject
factor representing different Response Modes (for comparisons within the same
version of the study) or setting (for comparisons between the in-person and online
versions).

For the 'biomechanical constraints' effect, a subset of the data considering only trials
where this effect could be present (i.e., rotations in the lateral (45°, 90°, 135°) and
medial (225°, 270°, 315°) directions) was used. The full model was *'Reaction Time ~*
Group*Direction*Laterality*View + (1|Participant)', where the effect of interest was the
Group * Direction interaction, or further higher-order interactions including these
factors. The biomechanical constraints effect was therefore analysed by collapsing
across the three rotation angles for each direction.

All models were run with weakly informative priors for all the beta parameters. For
reaction time, this was a normal distribution of mean = 0 and SD = 300ms, whereas
accuracy models were run with a normal distribution of mean = 0 and SD = 1 on the
logit scale. The same applied for the 'biomechanical constraints' models for reaction
time. Default (uniform) priors were used for the remaining parameters. All models
were run with 10 chains of 5,000 iterations each (1,000 warmup iterations per chain)
for an overall post-warmup of 40,000 iterations to inform the posterior distribution.
Model fit was assessed by visually inspecting posterior predictive checks and trace
plots, and R^2 values at convergence (R^2 < 1.01 was considered appropriate). More
details on the Bayesian models used can be found in Supplementary Materials.

For hypothesis testing, we used Bayesian Model Averaging to obtain the inclusion
Bayes Factor (BF_{inc}) for including each given effect or interaction, or against (BF_{01})
including it (Hinne et al., 2020). BF_{inc} were obtained by bridge sampling (Gronau
et al., 2017) using the 'bayestestR' package (Makowski et al., 2019). If evidence was
found for a given effect of Response Mode, or an interaction of Response Mode and
other factors, post-hoc comparisons were performed as equivalence tests
considering a Region of Practical Equivalence (ROPE) of 0.1*SD of the outcome
variable (Kruschke, 2018). Equivalence tests were run to examine evidence in favour

of the null hypothesis of equivalence (BF_{01}), or the alternative hypothesis of non-
equivalence (BF_{10}), always with equal-prior factor coding (Morey & Rouder, 2011).
For post-hoc comparisons not involving Response Mode, point-null hypothesis
testing was used. BFs for post-hoc comparisons were obtained via the Savage-
Dickey density ratio. All BFs were interpreted following established benchmarks,
considering 'no' evidence, 'anecdotal', 'moderate', 'strong', 'very strong' and 'extreme'
evidence if the BF was ≈1, 1-3, 3-10, 10-30, 30-100 and >100, respectively
(Andraszewicz et al., 2015; Jeffreys, 1998). Bayes factors are presented in relation to
the null (BF_{01}) or alternative (BF_{10}) hypothesis.

RESULTS

In-person version

Overall, 1.35 ± 0.27% (mean ± SD) of trials in the HLJT were rejected due to
extremely short (<300ms) or long (>3000ms) reaction time. The mean time to
complete the task was 17m 3s ± 1m 26.4s. There was evidence against including a
main effect of Response Mode or interactions between this factor and other terms for
reaction time, accuracy and the 'biomechanical constraints' effect. Therefore, we first
present results collapsing across groups, but we also report them separately (see
below for details).

<u>Reaction time:</u>

Overall, 13,109 valid trials were analysed. There was extreme evidence for including
the main effects of Angle, and View, and an interaction between them (BF_{inc} =
$3.21x10^{138}$, BF_{inc} = $3.17x10^{17}$ and BF_{inc} = $5.8x10^{17}$, respectively). Evidence against
including a main effect of Laterality, or two-way and three-way interactions was very
strong (BF_{01} > 35.71). There was evidence against including interactions of these
factors with Response Mode.

The main effect of Angle was explained by an increase in reaction time with stimulus
rotation (Fig. 2A, see Table 2 for pairwise comparisons), reaching the longest
reaction time at the maximum absolute rotation (180°). The main effect of View was
explained by palmar views (mean = 1038.69ms [979.97, 1096.95] 95%CrI) being
faster than dorsal views (mean = 1066.09ms [1007.54, 1124.87]), with inconclusive
evidence for the difference not being 0 (difference = -27.4ms [-57.8, 6.71], BF_{10} =
1.16).

The interaction between Angle and View suggested the presence of the
'biomechanical constraints' effect (Fig. 2B). Post hoc-analyses examined this
separately for palmar and dorsal views. Data for palmar views was consistent with a
'biomechanical constraints' effect, with the lateral rotations (45-135°) being slower
than their medial counterparts (225-315°). Pairwise comparisons showed extreme
evidence for a difference in each comparison (45° vs 315°: BF_{10} = 840.01; 90° vs
270°: BF_{10} = $1.48x10^5$; 135° vs 225°: BF_{10} = $1.91x10^{11}$). By contrast, there was no
evidence for a 'biomechanical constraints' effect for dorsal views, with moderate
evidence in favour of the null hypothesis for each comparison (45° vs 315°: BF_{01} =
4.17; 90° vs 270°: BF_{01} = 8.70; 135° vs 225°: BF_{01} = 7.04).

Table 2. Bayes Factors in favour of the null hypothesis of a zero difference (BF$_{01}$; in
bold) or the alternative (BF$_{10}$) hypothesis for the pairwise comparisons between
neighbouring Angles for Accuracy (%) and Reaction Time (milliseconds) in the Hand
Laterality Judgement Task.

Comparison	In-person version		Online version	
	Reaction Time	Accuracy	Reaction Time	Accuracy
0° vs 45°	**BF$_{01}$ = 2.48**	**BF$_{01}$ = 9.35**	**BF$_{01}$ = 1.35**	**BF$_{01}$ = 4.74**
45° vs 90°	BF$_{10}$ = 59.11	**BF$_{01}$ = 1.73**	BF$_{10}$ = 3.13x10^6	**BF$_{01}$ = 2.26**
90° vs 135°	BF$_{10}$ = 1.49x10^9	BF$_{10}$ = 1.39	BF$_{10}$ = 1.44x10^6	BF$_{10}$ = 111.46
135° vs 180°	BF$_{10}$ = 2.45x10^4	BF$_{10}$ = 21.60	BF$_{10}$ = 5.55x10^4	**BF$_{01}$ = 8.06**
180° vs 225°	BF$_{10}$ = 2.39x10^{18}	BF$_{10}$ = 5.96x10^4	BF$_{10}$ = 7.2x10^{15}	BF$_{10}$ = 1.96x10^5
225° vs 270°	BF$_{10}$ = 6.91x10^3	BF$_{10}$ = 227.97	BF$_{10}$ = 2.01x10^5	**BF$_{01}$ = 2.63**
270° vs 315°	BF$_{10}$ = 265.02	**BF$_{01}$ = 9.8**	BF$_{10}$ = 879.18	**BF$_{01}$ = 6.8**

<u>Accuracy:</u>

Overall, 15,153 valid trials were analysed. Only extreme evidence for including a
main effect of Angle was found (BF$_{inc}$ = 1.47x10^{24}; Fig. 2C), explained by a decrease
in accuracy with stimulus rotation, reaching its minimum value at the highest absolute
rotation (180°). Pairwise comparisons are shown in Table 2. Extreme evidence
against including all the remaining main effects and interactions was found (BF$_{01}$ >
6849.32).

<u>Biomechanical constraints effect:</u>

When considering only trials that could be affected by biomechanical constraints (i.e.
pooling data for 'lateral' and 'medial' stimuli while excluding 0° and 180° rotations), a
total of 10,053 valid trials were available for analysis. Extreme evidence for including
a main effect of Direction (BF$_{inc}$ = 5.1x10^{23}), along with a Direction x View interaction
(BF$_{inc}$ = 7.49x10^{10}) were found. Moderate evidence for including a main effect of
Laterality (BF$_{inc}$ = 6.39) and strong evidence for including a Direction x Laterality
interaction was found (BF$_{inc}$ = 11.72). Strong evidence for including a three-way
Direction x View x Laterality interaction (BF$_{inc}$ = 15.38) was also found.

The main effect of Direction was explained by lateral directions being slower than
medial directions, consistent with the 'biomechanical constraints' effect (lateral:
1079.38ms [1019.64, 1137.92]; medial: 968.66ms [910.30, 1027.83]; difference = -
111ms [-136, -86.8], BF$_{10}$ = 3.55x10^6). This effect was critically conditioned on View
(Fig. 2D), as for dorsal views, there was strong evidence in favour of the null
hypothesis of the difference being 0 (difference = -15.4ms [-50.7, 19.1], BF$_{01}$ =
12.99), whereas for palmar views there was extreme evidence for a difference not
being 0 (difference = -205.8ms [-240.6, -172.3], BF$_{10}$ = 1.55x10^{10}). Stimulus laterality
weakly conditioned the effect, as the difference was greater for right hands than for
left hands (right = -124ms [-158, -88.7], BF$_{10}$ = 4.62x10^6; left = -97.4ms [-132, -62.7],
BF$_{10}$ = 6.12x10^3). The three-way interaction revealed more marked 'biomechanical
constraints' for right palmar (difference = -218.56ms [-267.8, -170.3], BF$_{10}$ = 1.1x10^8)
than left palmar (difference = -193.08ms [-243.2, 147.0], BF$_{10}$ = 1.56x10^7), whereas

for dorsal views, neither side showed evidence for a biomechanical effect (right = -
29.54ms [-77.0, 19.9], BF_{01} = 7.52; left = -1.57ms [-50.4, 47.8], BF_{01} = 14.93).

Figure 2. Results of the in-person (N=40) version of the Hand Laterality Judgement Task.
Panels A and B show the Reaction Time and Accuracy measures by Rotation Angle and
View. **Panel C** shows the 'biomechanical constraints' effect (medial vs lateral difference in
milliseconds) for the corresponding pairs of angles. **Panel D** shows the 'biomechanical
constraints' effect collapsed across angles, split by View and Laterality.

Effects of Response Mode:

The main results are shown in Fig. 3. For reaction time, there was very strong
evidence against including a main effect of Response Mode (BF_{01} = 32.26), and very
strong to extreme evidence against including two-way, three-way, or four-way
interactions (BF_{01} > 83.33, BF_{01} > 1.46×10^{5} and BF_{01} > 1.32×10^{14}, respectively). The
Foot Group (mean = 979.47ms [898.86, 1061.71]) was faster than the Bimanual
Group (mean = 1125.32ms [1044.06, 1206.25]), with moderate evidence in favour of
non-equivalence (BF_{10} = 3.78; ROPE = 0 ± 31.27ms).

For accuracy, extreme evidence against including a main effect of Response Mode
was found (BF_{01} = 5.68×10^{8}), as well as interactions including this factor (BF_{01} =
3.64×10^{8}). Accuracy was comparable between the Foot Group (mean = 86.66%
[82.39, 90.33]) and the Bimanual Group (mean = 87.06% [82.6, 90.88]), with strong
evidence in favour of equivalence (BF_{01} = 20.83; ROPE = 0 ± 3.42%).

For the 'biomechanical constraints' effect model, evidence for including a Response
Mode x Direction interaction was inconclusive but favoured the null hypothesis (BF_{01}

= 1.46). The 'biomechanical constraints' effect was present in both groups (Foot = -89.5ms [-124, -55.5], BF_{10} = 3.58x10³; Bimanual = -132ms [-166, -96.9], BF_{10} = 6.83x10⁵), with moderate evidence for equivalence (BF_{01} = 4.9). Evidence against including interactions with View or Laterality was moderate (BF_{01} = 5.29 and BF_{01} = 4.52, respectively). Evidence against three-way interactions involving Response Mode, or a four-way interaction, was strong to extreme (BF_{01} > 52.63 and BF_{01} = 30581.04, respectively).

Figure 3. Comparison of the Foot and Bimanual Response Modes (N=20 each) of the in-person version of the Hand Laterality Judgement Task. **Panels A and B** show the Reaction Time and Accuracy measures by Rotation Angle and View. **Panel C** shows the 'biomechanical constraints effect' (medial vs lateral difference in milliseconds) for the corresponding pairs of angles. **Panel D** shows the 'biomechanical constraints' effect collapsing across angles but splitting by View and Laterality.

Online version

Overall, 4.36 ± 3.04% of trials were rejected due to extremely short (<300ms) or long (>3000ms) reaction time. The mean time to complete the task was 22m 22s ± 5m 40s. Evidence against including an effect of Response Mode was found in terms of accuracy and the 'biomechanical constraints' effect, as well as evidence against higher-order interactions in these models. For reaction time, evidence for including a main effect of Response Mode, and a Response Mode by View interaction was found (see below for details).

Reaction time:

Overall, 19,544 valid trials were analysed. Extreme evidence for including the main effects of Angle ($BF_{inc} = 2.76 \times 10^{187}$), View ($BF_{inc} = 9 \times 10^{32}$) and Laterality ($BF_{inc} = 551.26$), for their two-way interactions (Angle x View: $BF_{inc} = 1.69 \times 10^{33}$, Angle x Laterality: $BF_{inc} = 1030$, Laterality x View: $BF_{inc} = 1030$) and for a three-way interaction ($BF_{inc} = 1380$) was found.

The effect of Angle was explained by an increase in reaction time with stimulus rotation (Fig. 4A, see Table 2 for pairwise comparisons), reaching the longest reaction time at the maximum absolute rotation (180°). The effect of View was explained by dorsal stimuli showing longer reaction time than palmar stimuli (dorsal: 1223.21ms [1148.21, 1298.78]; palmar: 1178.04ms [1101.81, 1253.97]; $BF_{10} = 3200$). The effect of Laterality was explained by left stimuli showing longer reaction time compared to right stimuli (left: 1229.23ms [1154.24, 1306.47]; right: 1172.02ms [963.0, 1248.16]; $BF_{10} = 5.49 \times 10^{5}$). Critically, the interaction between Angle and View was consistent with the 'biomechanical constraints' effect (Fig. 4B). Post-hoc analysis for palmar views indicated that lateral rotations (45-135°) were slower than their medial counterparts (225-315°). Pairwise comparisons showed extreme evidence for the difference not being 0 at each rotation angle (45° vs 315°: $BF_{10} = 263.55$; 90° vs 270°: $BF_{10} = 9.67 \times 10^{8}$; 135° vs 225°: $BF_{10} = 3.46 \times 10^{14}$). Further post-hoc analysis identified that this effect did not appear for dorsal views, with anecdotal to strong evidence in favour of the null hypothesis (45° vs 315°: $BF_{01} = 2.58$; 90° vs 270°: $BF_{01} = 10$; 135° vs 225°: $BF_{01} = 13.51$).

Accuracy:

Overall, 22,036 valid trials were analysed. Extreme evidence for including the main effect of Angle ($BF_{inc} = 5.95 \times 10^{22}$), a main effect of View ($BF_{inc} = 2.4 \times 10^{4}$), and their interaction ($BF_{inc} = 4.4 \times 10^{4}$) was found. Extreme evidence against including a main effect of Laterality ($BF_{01} = 2.08 \times 10^{4}$) was found.

The main effect of Angle was driven by accuracy generally decreasing with stimulus rotation (Fig. 4C; see Table 2 for pairwise comparisons). The main effect of View was explained by strong evidence in favour of the palmar views being more accurate than the dorsal views (palmar: 89.78% [85.89, 93.1]; dorsal: 88.4% [84.63, 91.7]; $BF_{10} = 18.47$). The interaction between Angle and View was explained by the effect of Angle being stronger for dorsal views compared to palmar views at the comparisons 90° vs 135°, 135° vs 180° and 180° vs 225° (dorsal: $BF_{10} = 16.72$; $BF_{10} = 385.21$; $BF_{10} = 4.94 \times 10^{5}$; palmar: $BF_{01} = 1.49$; $BF_{10} = 2.05$; $BF_{10} = 5.03$, respectively).

Biomechanical constraints effect:

The model analysed 14,868 valid trials. There was extreme evidence for including a main effect of Direction ($BF_{inc} = 3.34 \times 10^{33}$), as shown by medial rotations (mean = 1126.22ms [1049.06, 1203.21]) being faster than lateral rotations (mean = 1233.66ms [1155.83, 1310.36]), consistent with the 'biomechanical constraints' effect (difference = -107ms [-128, -86.7], $BF_{10} = 8.03 \times 10^{8}$). This effect was critically conditioned on View, as shown by extreme evidence of including a Direction x View interaction ($BF_{inc} = 5.09 \times 10^{15}$), whereby the difference was only present for palmar views (difference = -197.6ms [-226.5, -168.0], $BF_{10} = 9.8 \times 10^{13}$) and not for dorsal

views (difference = -17.2ms [-46.7, 11.9], BF_{01} = 11.49). Evidence for including a
Direction x Laterality interaction was also extreme (BF_{inc} = 9.61x10^5), whereby the
medial vs lateral difference was larger for right hands (difference = -117ms [-146, -
88.6], BF_{10} = 4.79x10^6) than for left hands (difference = -98ms [-127, -68.9], BF_{10} =
5.29x10^4). Extreme evidence for including a three-way interaction (Fig. 4D) was also
found (BF_{inc} = 1.23x10^6), showing that for dorsal views, evidence against
biomechanical constraints was stronger for left hands (difference = 2.21ms [-38.9,
43.17), BF_{01} = 17.85) than right hands (difference = -36.74ms [-77.4, 3.99], BF_{01} =
3.76). In the palmar view, the effect was comparable between sides (right = -196.7ms
[-238.1, -156.1], BF_{10} = 6.12x10^8; left = -198.39ms [-239.6, -157.41], BF_{10} =
6.96x10^9).

Figure 4. Results of the online (N=60) version of the Hand Laterality Judgement Task.
Panels A and B show the Reaction Time and Accuracy measures by Rotation Angle and
View. **Panel C** shows the 'biomechanical constraints effect' (medial vs lateral difference in
milliseconds) for the corresponding pairs of angles. **Panel D** shows the 'biomechanical
constraints' effect collapsing across angles but splitting by View and Laterality.

Effects of Response Mode:

The main results are shown in Fig. 5. For reaction time, evidence for including a main
effect of Response Mode was extreme (BF_{inc} = 1.66x10^3). Evidence in favour of
equivalence was moderate comparing the Left-hand Group and the Bimanual Group
(BF_{01} = 4.51, ROPE = 0 ± 38.9ms). However, the Right-hand Group was slower than
the other two groups (Right-hand: 1364.03ms [1234.43, 1491.26]; Left-hand:
1115.60ms [988.68, 1245.60]; Bimanual: 1122.24ms [995.66, 1248.29]), and

evidence of non-equivalence was moderate comparing the Right-hand Group and the
Bimanual Group ($BF_{10} = 9.32$), and strong comparing the Right-hand Group and the
Left-hand Group ($BF_{10} = 11.74$). Response Mode only interacted with View (BF_{inc} =
3140), but not with Angle ($BF_{01} = 4.51$) and Laterality ($BF_{01} = 4.51$). Extreme
evidence against including three-way and four-way interactions was found ($BF_{01} >$
$3.57x10^6$). The interaction between Response Mode and View was driven by
evidence of non-equivalence between the groups being stronger (i.e. the Right-hand
group responding more slowly) for the dorsal views (Right-hand vs Bimanual: BF_{10} =
28.46; Right-hand vs Left-hand: $BF_{10} = 30.42$) than the palmar views (Right-hand vs
Bimanual: $BF_{10} = 2.71$; Right-hand vs Left-hand: $BF_{10} = 3.69$). Evidence for
equivalence between the Bimanual and Left-hand group did not change across views
(dorsal: $BF_{01} = 4.95$; palmar: $BF_{01} = 4.90$).

For accuracy, extreme evidence against including a main effect of Response Mode
($BF_{01} = 4.08x10^4$) was found. In fact, the overall accuracy was 89.31% [85.56, 92.54]
for the Bimanual Group, 89.2% [85.59, 92.34] for the Left-hand group and 88.77%
[84.62, 92.32] for the Right-hand Group. Evidence for equivalence was found
(Bimanual vs Right-hand: $BF_{01} = 15.38$; Bimanual vs Left-hand: $BF_{01} = 25.64$; Right-
hand vs Left-hand: $BF_{01} = 15.87$; ROPE = 0 ± 1.62%). Extreme evidence against
including all possible two-way or three-way interactions including Response Mode
($BF_{01} > 2.68x10^6$), and the four-way interaction ($BF_{01} = 4.59x10^{23}$) was found.

For the 'biomechanical constraints' model, very strong evidence against including a
Response Mode x Direction interaction ($BF_{01} = 90.91$), as well as a three-way
interaction with View ($BF_{01} = 50$) or Laterality ($BF_{01} = 1029.87$), or a four-way
interaction ($BF_{01} = 19493.18$) was found. The medial vs lateral difference was
present in all groups (Left-hand = -92.9ms [-129, -57.5], $BF_{10} = 2.9x10^3$; Bimanual = -
139.2ms [-175, -103.9], $BF_{10} = 4.69x10^6$; Right-hand = -90.0ms [-126, -54.8], BF_{10} =
$1.51x10^3$). The magnitude of the 'biomechanical constraints effect' was equivalent
between groups, with moderate to strong evidence in favour of the null hypotheses
(Bimanual vs Left-hand: $BF_{01} = 5.29$; Bimanual vs Right-hand: $BF_{01} = 4.37$; Left-hand
vs Right-hand: $BF_{01} = 55.56$).

Figure 5. Data from the online version of the Hand Laterality Judgement Task comparing Left-hand, Bimanual and Right-hand Response Modes (N=20 each). **Panels A and B** show the Reaction Time and Accuracy measures by Rotation Angle and View. **Panel C** shows the 'biomechanical constraints effect' (medial vs lateral difference in milliseconds) for the corresponding pairs of angles. **Panel D** shows the 'biomechanical constraints' effect collapsing across angles but splitting by View and Laterality.

Comparison of in-person and online versions (bimanual groups)

Socio-demographic characteristics were similar across the two Bimanual groups (age: $BF_{01} = 2.54$; sex: $BF_{10} = 1.38$), as well as MIQ-3 scores (Total: $BF_{01} = 1.17$; Kinesthetic: $BF_{01} = 2.59$; Internal Visual: $BF_{01} = 1.73$; External Visual: $BF_{10} = 2.6$). In the HLJT, rejection rates were similar for the online group compared to the in-person group (online = 2.03 ± 3.55%, in-person = 1.54 ± 1.54%, $BF_{01} = 2.84$). The overall time to complete the task was slightly longer for the online group than the in-person group (online = 21m 30s ± 5m 27.6s, in-person = 17m 42s ± 1m 17.4s; $BF_{10} = 9.17$).

For reaction time, 13,250 trials were analysed. Very strong evidence against including a main effect of Group (Fig. 6A), and higher-order interactions including this factor was found (all $BF_{01} > 5.37 \times 10^4$). Overall, the reaction time was equivalent across the groups (in-person = 1128.35ms [1018.9, 1235.87], online = 1114.05ms [1006.75, 1224.12], $BF_{01} = 5.16$; ROPE = 0 ± 34.77ms).

For accuracy (Fig. 6C), 15,086 trials were analysed. Extreme evidence against including a main effect of Group, and two-way, three-way or four-way interactions including this factor was found (all $BF_{01} > 6.06 \times 10^9$). In fact, overall accuracy showed

moderate evidence of equivalence (in-person = 87.77% [83.54, 91.37], online =
88.46% [84.5, 91.86], BF_{01} = 6.17; ROPE = 0 ± 1.66%).

For the 'biomechanical constraints' effect, 10,142 trials were analysed (Fig. 6D).
Extreme evidence against including a Group x Direction interaction was found (BF_{01}
= 47.62). In fact, the effect was similar across versions (in-person: -132ms [-166, -
96.9]; online = -139.2ms [-175, -103.9]), with very strong evidence for equivalence
(BF_{01} = 34.48; ROPE = 0 ± 32.54). Extreme evidence against including the
interactions of Group with View, Laterality, the three-way interaction and the four-way
interaction was found (all BF_{01} > 28.57).

Figure 6. Comparison of the in-person and online Bimanual groups (N=20 each) of the Hand
Laterality Judgement Task. **Panels A and B** show the Reaction Time and Accuracy
measures by Rotation Angle and View. **Panel C** shows the 'biomechanical constraints effect'
(medial vs lateral difference in milliseconds) for the corresponding pairs of angles. **Panel D**
shows the biomechanical constraints effect collapsing across angles but splitting by View
and Laterality.

DISCUSSION

This study developed an open-source version of the HLJT for both in-person and
online use. The study found that a bimanual Response Mode was largely equivalent
to responding with the feet in a lab setting. In addition, the bimanual Response Mode
in an online version was practically equivalent to its in-person counterpart. Both
versions reproduced the classical behavioural effects of this task, including the
'biomechanical constraints' effect. Finally, the comparison between bimanual and

unimanual Response Modes showed only a general increase in reaction time in the
group responding with the right hand relative to responding bimanually or with the left
hand (which were equivalent). This effect was slightly stronger for the dorsal view, did
not interact with any other factors of the task, and was not present for accuracy nor
affected the 'biomechanical constraints' effect.

Developing a standardized Hand Laterality Judgement Task

Standardisation of key parameters remains an unaddressed challenge in the field of
motor imagery, specifically in terms of assessment measures. This not only applies to
the HLJT, but a variety of methods (Suica et al., 2022). While we do not advocate for
the use of our specific paradigm in all future studies (as some decisions were made
based on feasibility or reliability considerations), we hope to partially address this
issue. By providing an open-source resource with reasonable default choices but
allowing flexibility for future studies (see below), we hope to contribute to creating a
common framework for researchers and clinicians interested in using the HLJT.

A wide variety of paradigms have been employed regarding the HLJT. This includes
different manipulations of the two main factors of rotation angle and hand view. Some
studies have used up to 12 different rotations (i.e. increments of 30°) (Cocksworth &
Punt, 2013) or as few as 4 (i.e. increments of 90°) (Conson et al., 2020; Saimpont
et al., 2009). Most studies use a number in-between these extremes, usually 6 (de
Vries et al., 2013; Ionta et al., 2007) or 8 angles (Brady et al., 2011; Mibu et al.,
2020). We decided to use 8 different rotations (i.e. increments of 45°) to maintain the
balance between using enough angles to identify a clear slope from 0° to 180°, while
reducing the overall number of trials required and therefore the overall time to
complete the task. With this paradigm, we could distinctly detect both the main
rotation effect for accuracy and reaction time, but also the 'biomechanical constraints'
effect at each specific rotation, in a task that lasted around 20 minutes overall.

The literature has been highly heterogeneous in the use of different hand views in the
HLJT. In the first well-known description of the task, Parsons used only hand
drawings of backs and palms of the hands (Parsons, 1987). Subsequently, other
studies incremented the variability in this parameter by introducing rotations in the
horizontal axis. For example, a 'thumb' view or a 'pinkie' view have been used, as
well as intermediate rotations in-between (Meng et al., 2016; Vannuscorps et al.,
2012). In fact, the effect of hand view is still controversial, as most studies suggest
that the 'biomechanical constraints' effect is only found in the palmar view and not in
the dorsal view (Conson et al., 2020; Meng et al., 2016; Mibu et al., 2020), which is
clearly consistent with our present results. However, some work suggests that
rotation in the horizontal axis is necessary to detect a 'biomechanical constraints'
effect, which was not observed with simple rotations in the frontal axis in either view
(Ter Horst et al., 2010). For the development of our proposed task, we decided to
maintain the 'traditional' palmar-dorsal paradigm to keep consistency with most of
previous studies. This paradigm clearly identified a 'biomechanical constraints' effect
for the palmar view and found strong evidence for its absence in the dorsal view. We
interpret this in line with a growing body of research suggesting that the palm and the
back of the hand may be processed using different cognitive strategies, and that only

the palm of the hand may trigger motor imagery-based processing (Conson et al.,
2021; Nagashima et al., 2019).

Several technical parameters of the HLJT paradigm used in this study were fixed or
chosen based on feasibility and reliability. For instance, we decided to include a
practice block with all the stimuli to allow the user to familiarise themselves with the
task. We consider this will be helpful for future applications, specifically for clinical
uses, as it will aid individuals to understand how the task works with minimal explicit
supervision by the researcher or clinician, which in our case was necessary for the
online version. Regarding the decision of providing feedback on accuracy on a trial-
by-trial basis, we chose to include this in the test blocks as we anticipated it would
enhance the engagement in the remote version of the task. To maintain consistency
across versions, we used the same procedure in the lab. We believe this may be a
key parameter to consider in future studies, as it provides participants with online
feedback during the task that allows them to adjust their performance if necessary. In
addition, we decided to keep the image on-screen until a response was provided, as
restricting the time that the participant has to process the stimulus would have limited
the generalisability of the paradigm to populations which are generally slower than
our sample of young, healthy individuals (e.g. older people or people with
neurological disorders). In addition, we decided to use 12 repetitions per unique
stimulus. A common rule of thumb in behavioural experiments is to include 8-10
repetitions (Matthews, 2011), but we decided to add more given the remote nature of
some of the comparisons, as we expected some participants would not be doing the
task as focused as if they were in the lab. Finally, we decided to allow participants to
have a break between test blocks and resume the task at their discretion (with a
minimum break of 10s). This, in turn, could explain why the online version took longer
than the in-person version, as participants may have taken longer breaks in the latter.
We made this decision to reduce potential effects of fatigue in future applications and
maintain engagement.

We are aware that the specific paradigm used in the present study, though sensible
for our purposes, may not meet all the requirements of future uses. For instance, in
the clinical field such a task could likely induce fatigue in some individuals and
therefore a shorter paradigm might be preferred, whereas a neuroscience study
might want to include more rotation angles or different hand views, or might prefer
not to provide feedback in the test trials, modify the number of repetitions, etc.
Therefore, the task that we publicly share has been modified to allow the user to
choose between all these possibilities. At the time of writing, the in-person task has
some predefined parameters that can be selected with a mouse click at the beginning
of the experiment, including technical considerations such as language (English,
Spanish and French are currently supported, and users could add their own
translations), whether to include a practice block or not, whether to provide feedback
in test blocks or not, the duration of this feedback (0.3s, 0.5s, 0.8s and 1s are
available) and the number of repetitions per unique stimulus (12, 8 and 4 are
available). Furthermore, the user can select specific parameters of the HLJT, such as
the number of rotational angles (4, 6, 8 and 12 are currently available) and the hand
views (palmar, dorsal or both). We have set sensible defaults for all these
parameters, but changing the defaults is as simple as changing the order of the

options in the experiment settings. Furthermore, the range of options for these and
other parameters could be easily extended without extensive programming
requirements, thanks to PsychoPy2's capabilities (Peirce et al., 2019). We hope all
the above options will allow a wide range of uses in future studies and applications.

The use of the HLJT in online studies and the use of the hands to respond to
the task

Recent advancements in software development have allowed us to leverage the use
of online platforms for collecting potentially more representative data, on a larger
scale and in a shorter period of time (Bridges et al., 2020; Helms et al., 2021;
Johnson et al., 2022). We aimed to translate this development into the field of motor
imagery research by testing an online version of the HLJT. Previous work had
suggested a confound with responding to the task with the hands would impede to
use this Response Mode in the task, as the use of the same effector could interfere
with information processing (Cocksworth & Punt, 2013). This would have limited the
development of an online HLJT, which requires participants to respond on their own
keyboard, and therefore was addressed in this study as part of the development
procedure.

Our findings, nonetheless, were consistent with the idea that a manual Response
Mode does not meaningfully interfere with this task. In the in-person version, we
found evidence for equivalence between feet and bimanual responses in terms of
accuracy and the 'biomechanical constraints' effect, and evidence against higher-
order interactions with all other factors. We only observed moderate evidence for
non-equivalence in terms of reaction time (the Bimanual group being approximately
50ms slower than the Foot group across all conditions, without interactions between
Response Mode and other factors). We consider this is the main 'effector confound'
of using the hands to respond, as reaction times are generally faster in manual
responses than foot responses in other tasks (Simonen et al., 1995). This effect is
consistent with previous work showing longer reaction times for manual responses
compared to responding verbally (Cocksworth & Punt, 2013), though they found
much larger differences (around 280ms). As they used unimanual Response Modes,
which arguably represent a more complex paradigm than responding bimanually, this
may explain the different magnitude of the effects. Both studies coincided in the
absence of interactions of Response Mode with the rotation angle and hand view,
and although Cocksworth & Punt observed lower accuracy for manual responses, the
effect was small (difference ≈ 2%). We did not find such an effect (between-group
difference < 1%), probably because we used equivalence tests instead of null-
hypothesis significance testing. Based on our results, we believe the 'effector
confound' arguably does not represent a meaningful difference which should prevent
to use bimanual responses in the task.

When comparing different bimanual and unimanual Response Modes, we found an
unexpected result. The group responding with their right-hand showed generally
longer reaction times than the other two groups. This partially contradicts previous
findings suggesting that both unimanual Response Modes were similar (Cocksworth
& Punt, 2013). Two (potentially interacting) reasons may explain this behaviour. First,

responding to this task with two fingers of one hand is arguably more complex than responding with one finger of each hand, which is a more intuitive method. Therefore, a relative slowness in the two unimanual groups was expected compared to the Bimanual group. However, both unimanual groups should have responded similarly, which was not the case. Our (speculative) interpretation relates to the second possible source of this effect; as all participants in our study were right-handed, they most likely were comparing their *dominant* hand to the observed stimuli while making the laterality judgements (i.e. they compared the orientation of the stimulus with their dominant hand, then judged if it could be congruent or not to make the right/left decision). There is previous evidence suggesting this effect may be present in the HLJT (Ní Choisdealbha et al., 2011). In the Left-hand group, this did not generate any conflicts between the processing of the stimulus and the response, as they were responding with their *non-dominant* hand. However, for the Right-hand group, a conflict might have arisen, as the hand used for the visual comparison was the same as the one used to provide the response. The fact that the effect was stronger for the dorsal stimuli further supports this idea, as the dorsal view of the hand is thought to trigger more visual than motor processing, perhaps because we are more familiar with seeing the back of our hands (Bläsing et al., 2013; Zapparoli et al., 2014). Notably, we did not find a corresponding effect for the Bimanual group, who also used their right-hand. We therefore speculate that a possible interaction between these effects may have been present; requiring participants to perform the more complex unimanual version of the task with their right hand may have interfered with the ability to compare their own right hand with the on screen stimulus.

Aside from the findings discussed above, all other comparisons across Response Modes provided strong evidence for equivalence, including all accuracy models and the 'biomechanical constraints' effect. While future studies should consider the above-mentioned particularities, we believe that for the purpose of applying the HLJT in applied and online contexts, these findings generally support the use of bimanual and unimanual responses in this task.

Strengths and Limitations

This study has some limitations. First, we did not investigate the differences between manual and verbal Response Modes, which have been also used in previous studies. We decided to compare a bimanual mode against a foot mode, as these are physically more similar than responding verbally, and this allows to rule out the possibility that slowness due to manual responses can be attributed to the response command simply reaching the effector more quickly. In addition, the feasibility of using hand and foot Response Modes in the task is higher as reaction times from button presses are unambiguous, whereas vocal responses require more training to prevent hesitations or pauses. Similarly, the accuracy of a response from a button press is immediately available allowing rapid feedback, whereas for vocal responses this is not currently possible. Future researchers could build upon our current open-source experiment to implement a vocal response version, if needed.

Second, as part of the study was run fully online, we were not able to assure participants in the unimanual groups were not using the contralateral hand to

facilitate performance, as we did not explicitly instruct them what they could or could
not do with it. However, given that most evidence indicated equivalence across
Response Modes in this version, as well as comparing online and in-person versions,
we believe the instructions given were specific enough to obtain reliable estimates in
this case. In the future, more explicit guidance could be incorporated.

Third, in the in-person version of the task, the sample was composed of mostly
females in the bimanual group, and this can influence HLJT performance (Conson
et al., 2020). Therefore, this introduces a potential limitation, as we did not include
sex as a covariate in any of the analyses because it was beyond the scope of this
paper and we had an unequal distribution between the groups.

Finally, we did not include a control task with non-biological stimuli, to further
establish whether the slowdown in processing that was observed is specific to the
HLJT or a general pattern also shown in other mental rotation tasks. As we were
mostly interested in the 'biomechanical constraints' effect, and it would not be present
in non-biological stimuli (Bek et al., 2022), we decided not to include such a control
task, as it would have doubled the time needed to complete the experiment,
potentially compromising the feasibility of the online version of the task.

CONCLUSIONS

An open-source, freely available Hand Laterality Judgement Task was developed for
in-person and online use. The task reproduced established phenomena of this
paradigm, both in-person and remotely, and across different Response Modes. For
the in-person version, evidence for equivalence between a foot and a bimanual
Response Mode was found in terms of accuracy and the 'biomechanical constraints'
effect. While reaction times were slightly longer in the bimanual group, we found
evidence against related higher-order interactions. For the online version, evidence
for equivalence between the bimanual and left-hand responses was found for all
measures, whereas longer reaction times were found for the Right-hand responses,
predominantly for the dorsal view of the hand. Evidence against all other higher-order
interactions was found. Evidence of equivalence between the two bimanual groups,
in-person and online, was also observed. We conclude that both in-person and online
versions reliably replicated key behavioural effects in the HLJT, providing a
standardized (but also highly customizable) version of the paradigm that can be
readily applied in future studies.

ACKNOWLEDGEMENTS

We would like to thank Andrés de los Santos Suárez for his assistance with the creation of the figures, and Alfredo Lerín Calvo, Celia López-Cañadillas and Alfredo Hernando Jorge for their help with participant recruitment.

CONFLICTS OF INTEREST

We have no conflicts of interest to disclose.

FUNDING

MMV, SMcA, BMW, and RMH are supported by an FNRS MIS Grant (FNRS F.4523.23). MMV is supported by an FNRS CR Fellowship (FNRS 1.B359.25). EVC is supported by an FNRS ASP fellowship (FNRS 1.AB19.24). GH is supported by an FNRS CDR award (FNRS J.0084.21).

AUTHOR ROLES (CRediT)

Author	Conceptualization	Methodology	Software	Validation	Formal Analysis	Investigation	Resources	Data Curation	Writing – original draft	Writing – reviewing & editing	Visualization	Supervision	Project Administration	Funding Acquisition
MMV	X	X	X	X	X	X		X	X	X	X		X	
SMcA		X	X		X					X	X			
BMW			X		X					X				
EVC										X				
RMH	X	X		X			X		X	X		X	X	X

REFERENCES

Andraszewicz, S., Scheibehenne, B., Rieskamp, J., Grasman, R., Verhagen, J., & Wagenmakers, E.-J. (2015). An Introduction to Bayesian Hypothesis Testing for Management Research. *Journal of Management*, *41*(2), 521-543. https://doi.org/10.1177/0149206314560412

Bek, J., Humphries, S., Poliakoff, E., & Brady, N. (2022). Mental rotation of hands and objects in ageing and Parkinson's disease: Differentiating motor imagery and visuospatial ability. *Experimental Brain Research*, *240*(7), 1991-2004. https://doi.org/10.1007/S00221-022-06389-5/TABLES/3

Bläsing, B., Brugger, P., Weigelt, M., & Schack, T. (2013). Does thumb posture influence the mental rotation of hands? *Neuroscience Letters*, *534*, 139-144. https://doi.org/10.1016/j.neulet.2012.11.034

Brady, N., Maguinness, C., & Ní Choisdealbha, A. (2011). My hand or yours? Markedly different sensitivity to egocentric and allocentric views in the hand laterality task. *PloS One*, *6*(8), e23316. https://doi.org/10.1371/journal.pone.0023316

Bridges, D., Pitiot, A., MacAskill, M. R., & Peirce, J. W. (2020). The timing mega-study: Comparing a range of experiment generators, both lab-based and online. *PeerJ*, *8*, e9414. https://doi.org/10.7717/peerj.9414

Bürkner, P.-C. (2017). brms: An R Package for Bayesian Multilevel Models Using Stan. *Journal of Statistical Software*, *80*, 1-28. https://doi.org/10.18637/jss.v080.i01

Cocksworth, R. L., & Punt, T. D. (2013). When the left hand does not know what the left hand is doing: Response mode affects mental rotation of hands. *Experimental Brain Research*, *228*(1), 87-95. https://doi.org/10.1007/s00221-013-3540-2

Collet, C., Guillot, A., Lebon, F., MacIntyre, T., & Moran, A. (2011). Measuring Motor Imagery Using Psychometric, Behavioral, and Psychophysiological Tools. *Exercise and Sport Sciences Reviews*, *39*(2), 85-92. https://doi.org/10.1097/JES.0b013e31820ac5e0

Conson, M., De Bellis, F., Baiano, C., Zappullo, I., Raimo, G., Finelli, C., Ruggiero, I., Positano, M., & Trojano, L. (2020). Sex differences in implicit motor imagery: Evidence from the hand laterality task. *Acta Psychologica*, *203*. https://doi.org/10.1016/j.actpsy.2020.103010

Conson, M., Di Rosa, A., Polito, F., Zappullo, I., Baiano, C., & Trojano, L. (2021). "Mind the thumb": Judging hand laterality is anchored on the thumb position. *Acta Psychologica*, *219*. https://doi.org/10.1016/j.actpsy.2021.103388

de Vries, S., Tepper, M., Feenstra, W., Oosterveld, H., Boonstra, A. M., & Otten, B. (2013). Motor imagery ability in stroke patients: The relationship between implicit and explicit motor imagery measures. *Frontiers in Human Neuroscience*, *7*, 790. https://doi.org/10.3389/fnhum.2013.00790

Floridou, G. A., Peerdeman, K. J., & Schaefer, R. S. (2022). Individual differences in mental imagery in different modalities and levels of intentionality. *Memory & Cognition*, *50*(1), 29-44. https://doi.org/10.3758/s13421-021-01209-7

Gronau, Q. F., Sarafoglou, A., Matzke, D., Ly, A., Boehm, U., Marsman, M., Leslie, D. S., Forster, J. J., Wagenmakers, E.-J., & Steingroever, H. (2017). A tutorial on bridge sampling. *Journal of Mathematical Psychology*, *81*, 80-97. https://doi.org/10.1016/j.jmp.2017.09.005

Guillot, A., Collet, C., Nguyen, V. A., Malouin, F., Richards, C., & Doyon, J. (2008). Functional neuroanatomical networks associated with expertise in motor imagery. *NeuroImage*, *41*(4), 1471-1483. https://doi.org/10.1016/j.neuroimage.2008.03.042

Helms, Y. B., Hamdiui, N., Kretzschmar, M. E. E., Rocha, L. E. C., van Steenbergen, J. E., Bengtsson, L., Thorson, A., Timen, A., & Stein, M. L. (2021). Applications and Recruitment Performance of Web-Based Respondent-Driven Sampling: Scoping Review. *Journal of Medical Internet Research*, *23*(1), e17564. https://doi.org/10.2196/17564

Hinne, M., Gronau, Q. F., van den Bergh, D., & Wagenmakers, E.-J. (2020). A Conceptual Introduction to Bayesian Model Averaging. *Advances in Methods and Practices in Psychological Science*, *3*(2), 200-215. https://doi.org/10.1177/2515245919898657

Hudson, M. L., McCormick, K., Zalucki, N., & Moseley, G. L. (2006). Expectation of pain replicates the effect of pain in a hand laterality recognition task: Bias in information processing toward the painful side? *European Journal of Pain (London, England)*, *10*(3), 219-224. https://doi.org/10.1016/j.ejpain.2005.03.009

Ionta, S., Fourkas, A. D., Fiorio, M., & Aglioti, S. M. (2007). The influence of hands posture on mental rotation of hands and feet. *Experimental Brain Research*, *183*(1), 1-7. https://doi.org/10.1007/s00221-007-1020-2

Jeannerod, M. (2001). *Neural simulation of action: A unifying mechanism for motor cognition*. *14*, 103-109. https://doi.org/10.1006/nimg.2001.0832

Jeffreys, H. (1998). *The Theory of Probability*. OUP Oxford.

Johnson, B. P., Dayan, E., Censor, N., & Cohen, L. G. (2022). Crowdsourcing in Cognitive and Systems Neuroscience. *The Neuroscientist: A Review Journal Bringing Neurobiology, Neurology and Psychiatry*, *28*(5), 425-437. https://doi.org/10.1177/10738584211017018

Kraeutner, S. N., El-Serafi, M., Lee, J. W., & Boe, S. G. (2019). Disruption of motor imagery
performance following inhibition of the left inferior parietal lobe. *Neuropsychologia*, *127*,
106-112. https://doi.org/10.1016/j.neuropsychologia.2019.02.016

Kraeutner, S. N., Eppler, S. N., Stratas, A., & Boe, S. G. (2020). Generate, maintain,
manipulate? Exploring the multidimensional nature of motor imagery. *Psychology of Sport*
and Exercise, *48*, 101673. https://doi.org/10.1016/j.psychsport.2020.101673

Kruschke, J. K. (2018). Rejecting or Accepting Parameter Values in Bayesian Estimation.
Advances in Methods and Practices in Psychological Science, *1*(2), 270-280.
https://doi.org/10.1177/2515245918771304

Ladda, A. M., Lebon, F., & Lotze, M. (2021). Using motor imagery practice for improving
motor performance – A review. *Brain and Cognition*, *150*, 105705.
https://doi.org/10.1016/j.bandc.2021.105705

Liu, F., & Eugenio, E. C. (2018). A review and comparison of Bayesian and likelihood-based
inferences in beta regression and zero-or-one-inflated beta regression. *Statistical Methods*
in Medical Research, *27*(4), 1024-1044. https://doi.org/10.1177/0962280216650699

Makowski, D., Ben-Shachar, M. S., & Lüdecke, D. (2019). bayestestR: Describing Effects
and their Uncertainty, Existence and Significance within the Bayesian Framework. *Journal*
of Open Source Software, *4*(40), 1541. https://doi.org/10.21105/joss.01541

Matthews, W. J. (2011). Stimulus Repetition and the Perception of Time: The Effects of Prior
Exposure on Temporal Discrimination, Judgment, and Production. *PLoS ONE*, *6*(5),
e19815. https://doi.org/10.1371/journal.pone.0019815

McAvinue, L. P., & Robertson, I. H. (2008). Measuring motor imagery ability: A review.
European Journal of Cognitive Psychology, *20*(2), 232-251.
https://doi.org/10.1080/09541440701394624

Meng, S., Oi, M., Sekiyama, K., & Saito, H. (2016). The neural mechanism of biomechanical
constraints in the hand laterality judgment task: A near-infrared spectroscopy study.
Neuroscience Letters, *627*, 211-215. https://doi.org/10.1016/j.neulet.2016.05.069

Mibu, A., Kan, S., Nishigami, T., Fujino, Y., & Shibata, M. (2020). Performing the hand
laterality judgement task does not necessarily require motor imagery. *Scientific Reports*,
10(1). https://doi.org/10.1038/s41598-020-61937-9

Morey, R. D., & Rouder, J. N. (2011). Bayes factor approaches for testing interval null
hypotheses. *Psychological Methods*, *16*(4), 406-419. https://doi.org/10.1037/a0024377

Nagashima, I., Takeda, K., Shimoda, N., Harada, Y., & Mochizuki, H. (2019). Variation in
performance strategies of a hand mental rotation task on elderly. *Frontiers in Human*
Neuroscience, *13*. https://doi.org/10.3389/fnhum.2019.00252

Ní Choisdealbha, Á., Brady, N., & Maguinness, C. (2011). Differing roles for the dominant
and non-dominant hands in the hand laterality task. *Experimental Brain Research*, *211*(1),
73-85. https://doi.org/10.1007/s00221-011-2652-9

Parsons, L. M. (1987). Imagined spatial transformations of one's hands and feet. *Cognitive*
Psychology, *19*(2), 178-241. https://doi.org/10.1016/0010-0285(87)90011-9

Parsons, L. M. (1994). Temporal and Kinematic Properties of Motor Behavior Reflected in
Mentally Simulated Action. *Journal of Experimental Psychology: Human Perception and*
Performance, *20*(4), 709-730. https://doi.org/10.1037/0096-1523.20.4.709

Peirce, J., Gray, J. R., Simpson, S., MacAskill, M., Höchenberger, R., Sogo, H., Kastman, E.,
& Lindeløv, J. K. (2019). PsychoPy2: Experiments in behavior made easy. *Behavior*
Research Methods, *51*(1), 195-203. https://doi.org/10.3758/s13428-018-01193-y

Robin, N., Coudevylle, G. R., Guillot, A., & Toussaint, L. (2020). French translation and
validation of the Movement Imagery Questionnaire-third version (MIQ-3f). *Movement &*
Sport Sciences - Science & Motricité, *108*, Article 108.
https://doi.org/10.1051/sm/2019035

Saimpont, A., Pozzo, T., & Papaxanthis, C. (2009). Aging affects the mental rotation of left
and right hands. *PloS One*, *4*(8), e6714. https://doi.org/10.1371/journal.pone.0006714

Simonen, R. L., Videman, T., Battié, M. C., & Gibbons, L. E. (1995). Comparison of foot and
hand reaction times among men: A methodologic study using simple and multiple-choice

repeated measurements. *Perceptual and Motor Skills*, *80*(3, Pt 2), 1243-1249. https://doi.org/10.2466/pms.1995.80.3c.1243

Suica, Z., Behrendt, F., Gäumann, S., Gerth, U., Schmidt-Trucksäss, A., Ettlin, T., & Schuster-Amft, C. (2022). Imagery ability assessments: A cross-disciplinary systematic review and quality evaluation of psychometric properties. *BMC Medicine*, *20*(1), 166. https://doi.org/10.1186/s12916-022-02295-3

Ter Horst, A. C., Van Lier, R., & Steenbergen, B. (2010). Mental rotation task of hands: Differential influence number of rotational axes. *Experimental Brain Research*, *203*(2), 347-354. https://doi.org/10.1007/s00221-010-2235-1

Trapero-Asenjo, S., Gallego-Izquierdo, T., Pecos-Martín, D., & Nunez-Nagy, S. (2021). Translation, cultural adaptation, and validation of the Spanish version of the Movement Imagery Questionnaire-3 (MIQ-3). *Musculoskeletal Science and Practice*, *51*, 102313. https://doi.org/10.1016/j.msksp.2020.102313

van Melick, N., Meddeler, B. M., Hoogeboom, T. J., Nijhuis-van der Sanden, M. W. G., & van Cingel, R. E. H. (2017). How to determine leg dominance: The agreement between self-reported and observed performance in healthy adults. *PloS One*, *12*(12), e0189876. https://doi.org/10.1371/journal.pone.0189876

Vannuscorps, G., & Caramazza, A. (2016). The origin of the biomechanical bias in apparent body movement perception. *Neuropsychologia*, *89*, 281-286. https://doi.org/10.1016/j.neuropsychologia.2016.05.029

Vannuscorps, G., Pillon, A., & Andres, M. (2012). Effect of biomechanical constraints in the hand laterality judgment task: Where does it come from? *Frontiers in Human Neuroscience*, *6*. https://doi.org/10.3389/fnhum.2012.00299

Waltzing, B., McAteer, S., Moreno-Verdú, M., Van Caenegem, E., Du, Y., & Hardwick, R. M. (2024). *Separate timescales for spatial and anatomical information processing during action observation* (p. 2024.10.07.616993). bioRxiv. https://doi.org/10.1101/2024.10.07.616993

Williams, S. E., Cumming, J., Ntoumanis, N., Nordin-Bates, S. M., Ramsey, R., & Hall, C. (2012). Further validation and development of the movement imagery questionnaire. *Journal of Sport and Exercise Psychology*, *34*(5), 621-646. https://doi.org/10.1123/jsep.34.5.621

Williams, S. E., Guillot, A., Di Rienzo, F., & Cumming, J. (2015). Comparing self-report and mental chronometry measures of motor imagery ability. *European Journal of Sport Science*, *15*(8), 703-711. https://doi.org/10.1080/17461391.2015.1051133

Zapparoli, L., Invernizzi, P., Gandola, M., Berlingeri, M., De Santis, A., Zerbi, A., Banfi, G., & Paulesu, E. (2014). Like the back of the (right) hand? A new fMRI look on the hand laterality task. *Experimental Brain Research*, *232*(12), 3873-3895. https://doi.org/10.1007/s00221-014-4065-z

Zhao, L. J., Jiang, L. H., Zhang, H., Li, Y., Sun, P., Liu, Y., & Qi, R. (2023). Effects of Motor Imagery Training for Lower Limb Dysfunction in Patients With Stroke: A Systematic Review and Meta-analysis of Randomized Controlled Trials. *American Journal of Physical Medicine & Rehabilitation*, *102*(5), 409-418. https://doi.org/10.1097/PHM.0000000000002107

                                     SUPPLEMENTARY MATERIALS

Details of Bayesian models

For accuracy, the zero-and-one inflated beta-regression models were modelled with
by-group (i.e. Response Mode) and by-Angle random intercepts and slopes for the
parameters 'phi', 'zoi' and 'coi', which estimated the measures of variability in the
beta distribution part of the model, the proportion of 0s and 1s in the distribution, and
the proportion of 1s within that proportion, respectively. This allowed the models to
converge better. These parameters were not used for any inferences, as the most
relevant measure was the beta distribution representing the continuous probability of
a correct response (i.e. Accuracy), which was modelled as explained in the Methods
section of this paper.

Customisability options in the Hand Laterality Judgement Task

Both local and online versions of the HLJT were developed in a modular manner, that
meaning that the level of flexibility is relatively high. The details about customisation
are fully described in a 'Readme' file specific to each version. As a general
description, in the current versions the experimenter can select the following
parameters:

• Language: English (default), Spanish and French. Language to display
instructions and messages throughout the task.
• Response mode: Both hands (default), Right hand or Left hand. Whether the
participant should respond bimanually or unimanually.
• Practice block: Yes (default) or No. Whether a practice block with all the
stimuli should be displayed before the actual test blocks.
• Number of angles: 4 angles with increments of 90° (default), 6 angles with
increments of 60°, 8 angles with increments of 45° or 12 angles with
increments of 30°.
• Hand views: Palmar and Dorsal (default), Palmar or Dorsal. Whether the
stimuli should appear in one hand view only or both.
• Number of repetitions: 12 (default), 8 or 4. Number of repetitions per unique
stimulus in the test blocks overall (i.e. in total excluding the practice block). As
of now the number of test blocks is fixed at 4, therefore, the total number of
repetitions must be a multiple of 4.
• Feedback: 0.3 seconds (default), 0.5 seconds, 0.8 seconds, 1 second or No
feedback. Time of feedback for accuracy (correct/incorrect) on a trial-by-trial
basis. This only applies to test blocks. If a practice block is included in the
experiment, feedback throughout this block is always provided.

In these versions, the inter-trial interval (ITI) for both the practice and test blocks, is
set at a random number between 0.6 and 1 seconds, with more probability in the
range 0.75 to 0.85 seconds, with the goal of having a random ITI throughout the
experiment with an average duration of ~0.8 seconds every 50 trials.

1008

Supplementary Tables with descriptives statistics in the in-person version.

Supplementary Table 1. Mean and SD for response time (milliseconds) for each Response Mode (group) according to Angle (0° to 315°), Laterality (left or right) and View (dorsal or palmar).

		Left				Right			
		Dorsal		Palmar		Dorsal		Palmar	
Angle	Group	Mean	SD	Mean	SD	Mean	SD	Mean	SD
0°	Foot	827.92	253.69	1017.97	423.06	793.29	278.66	904.85	347.17
	Bimanual	978.47	343.97	1126.10	481.00	924.41	414.39	1070.51	467.73
45°	Foot	910.84	349.62	1022.03	404.03	855.49	303.41	1001.50	385.67
	Bimanual	1081.33	470.16	1179.46	450.01	970.79	436.83	1152.25	444.00
90°	Foot	1079.28	407.96	1187.59	465.03	1005.50	387.16	1184.31	438.60
	Bimanual	1256.92	498.70	1410.35	532.55	1261.10	543.17	1369.96	552.18
135°	Foot	1328.12	538.27	1195.54	497.79	1318.98	505.93	1046.16	423.22
	Bimanual	1531.75	573.52	1492.67	617.45	1504.95	598.91	1311.45	523.80
180°	Foot	1084.28	415.45	977.93	409.21	1033.37	495.20	864.78	359.97
	Bimanual	1230.12	453.12	1116.48	433.89	1171.71	460.69	1010.36	380.81
225°	Foot	964.34	374.04	869.37	312.95	866.95	373.05	822.56	334.15
	Bimanual	1119.10	419.03	1000.62	370.66	1029.21	426.37	963.80	365.45
270°	Foot	829.88	286.18	871.26	322.64	773.60	301.17	799.48	320.38
	Bimanual	907.83	315.06	993.13	369.08	848.45	369.50	926.59	325.69
315°	Foot	827.92	253.69	1017.97	423.06	793.29	278.66	904.85	347.17
	Bimanual	978.47	343.97	1126.10	481.00	924.41	414.39	1070.51	467.73

Supplementary Table 2. Mean and SD for accuracy (%) for each Response Mode (group) according to Angle (0° to 315°), Laterality (left or right) and View (dorsal or palmar).

		Left				Right			
		Dorsal		Palmar		Dorsal		Palmar	
Angle	Group	Mean	SD	Mean	SD	Mean	SD	Mean	SD
0°	Foot	89.54	30.67	86.50	34.25	93.33	25.00	88.75	31.66
	Bimanual	86.25	34.51	91.67	27.70	88.75	31.66	90.76	29.03
45°	Foot	92.44	26.50	89.54	30.67	91.25	28.32	87.39	33.26
	Bimanual	90.25	29.72	87.87	32.72	87.45	33.20	86.55	34.19
90°	Foot	86.44	34.31	88.14	32.41	88.19	32.35	84.68	36.09
	Bimanual	89.41	30.84	90.72	29.08	83.97	36.77	81.20	39.16
135°	Foot	77.54	41.82	79.65	40.34	71.13	45.41	78.06	41.47
	Bimanual	90.68	29.14	85.71	35.07	75.95	42.83	80.95	39.35
180°	Foot	72.53	44.73	75.00	43.39	57.38	49.56	79.91	40.15
	Bimanual	76.96	42.20	74.01	43.96	69.91	45.97	75.11	43.33
225°	Foot	82.35	38.20	87.39	33.26	83.97	36.77	93.64	24.45
	Bimanual	88.84	31.55	90.21	29.78	84.52	36.25	89.54	30.67
270°	Foot	91.21	28.37	93.16	25.29	89.12	31.20	95.82	20.06
	Bimanual	93.31	25.05	94.58	22.68	90.00	30.06	95.00	21.84
315°	Foot	93.70	24.35	94.56	22.73	93.70	24.35	96.22	19.12
	Bimanual	92.47	26.45	92.08	27.06	93.72	24.30	92.44	26.50

1019 **Supplementary Tables with descriptives statistics in the online version.**

1020 **Supplementary Table 3**. Mean and SD for response time (milliseconds) for each
1021 Response Mode (group) according to Angle (0° to 315°), Laterality (left or right) and
1022 View (dorsal or palmar).

Angle	Group	Left				Right			
		Dorsal		Palmar		Dorsal		Palmar	
		Mean	SD	Mean	SD	Mean	SD	Mean	SD
0°	Bimanual	962.59	406.00	1032.30	414.57	858.16	337.01	969.95	395.07
	Left-hand	1028.31	387.03	1003.42	412.56	941.34	463.53	932.16	360.72
	Right-hand	1287.04	511.94	1188.90	500.35	1222.52	522.32	1225.30	499.89
45°	Bimanual	965.95	440.92	1133.96	457.16	902.99	394.37	1085.00	518.76
	Left-hand	1086.17	499.99	1100.83	431.18	955.33	426.05	962.00	386.29
	Right-hand	1293.51	552.02	1201.71	495.81	1245.36	533.03	1189.58	517.99
90°	Bimanual	1118.28	574.84	1180.72	469.77	1030.31	478.95	1152.42	550.65
	Left-hand	1133.55	462.22	1187.68	501.55	1030.35	434.90	1117.57	494.69
	Right-hand	1397.70	593.83	1361.04	588.06	1315.25	562.52	1387.97	620.70
135°	Bimanual	1219.58	500.47	1312.56	539.48	1162.24	496.69	1235.60	510.30
	Left-hand	1193.48	502.08	1355.50	539.56	1189.21	500.41	1297.34	565.08
	Right-hand	1441.74	551.71	1470.49	607.11	1452.80	574.97	1458.47	602.75
180°	Bimanual	1513.65	521.88	1249.11	531.03	1496.75	487.37	1195.21	531.28
	Left-hand	1330.55	505.66	1304.30	473.24	1412.99	518.96	1251.08	493.75
	Right-hand	1740.33	561.68	1547.38	626.33	1668.54	507.88	1406.76	625.81
225°	Bimanual	1272.35	506.11	1007.86	341.32	1120.06	452.76	993.53	421.90
	Left-hand	1282.56	552.32	1122.52	464.69	1137.60	466.82	1007.20	415.26
	Right-hand	1521.30	543.82	1251.39	585.67	1503.63	545.24	1194.16	522.04
270°	Bimanual	1073.38	402.84	971.10	386.32	1015.00	417.41	958.45	424.24
	Left-hand	1177.24	472.97	1051.99	406.00	990.64	414.94	905.69	339.61
	Right-hand	1394.91	518.85	1188.44	524.60	1409.68	544.31	1173.47	484.47
315°	Bimanual	954.32	401.12	991.58	380.06	872.23	382.46	973.24	433.82
	Left-hand	1082.04	507.01	970.98	384.74	874.04	352.48	926.03	384.57
	Right-hand	1235.62	531.45	1088.52	477.32	1245.80	520.03	1164.92	501.86

1023

1024

Supplementary Table 4. Mean and SD for accuracy (%) for each Response Mode (group) according to Angle (0° to 315°), Laterality (left or right) and View (dorsal or palmar).

Angle	Group	Left Dorsal Mean	SD	Left Palmar Mean	SD	Right Dorsal Mean	SD	Right Palmar Mean	SD
0°	Bimanual	94.47	22.91	94.98	21.88	97.06	16.93	94.09	23.63
	Left-hand	86.44	34.31	94.07	23.67	89.66	30.52	94.04	23.72
	Right-hand	89.74	30.40	95.67	20.40	86.16	34.61	94.40	23.05
45°	Bimanual	91.56	27.86	90.72	29.08	95.38	21.04	94.47	22.91
	Left-hand	89.79	30.35	92.41	26.55	90.21	29.78	91.38	28.13
	Right-hand	92.41	26.54	91.89	27.36	88.00	32.57	91.19	28.41
90°	Bimanual	89.79	30.35	90.04	30.01	89.92	30.18	84.55	36.22
	Left-hand	88.21	32.32	90.52	29.36	87.77	32.83	82.02	38.49
	Right-hand	91.52	27.92	92.89	25.76	88.94	31.44	90.37	29.57
135°	Bimanual	72.53	44.73	81.30	39.07	78.35	41.27	80.09	40.02
	Left-hand	88.60	31.86	84.55	36.22	77.92	41.57	76.44	42.53
	Right-hand	84.04	36.71	80.75	39.52	83.26	37.42	90.15	29.88
180°	Bimanual	65.78	47.55	83.48	37.22	60.34	49.02	83.98	36.76
	Left-hand	83.48	37.22	85.78	35.01	66.96	47.14	82.17	38.36
	Right-hand	77.39	41.94	85.71	35.07	71.36	45.32	89.20	31.11
225°	Bimanual	87.88	32.71	91.56	27.86	84.32	36.44	92.50	26.39
	Left-hand	91.38	28.13	91.91	27.32	85.04	35.74	92.77	25.96
	Right-hand	84.65	36.13	90.50	29.39	79.91	40.16	93.86	24.06
270°	Bimanual	94.92	22.02	94.56	22.73	94.14	23.53	94.96	21.93
	Left-hand	89.61	30.58	93.19	25.24	86.40	34.35	95.36	21.08
	Right-hand	90.54	29.33	95.09	21.66	89.47	30.76	95.24	21.34
315°	Bimanual	95.82	20.06	95.34	21.13	97.07	16.90	98.32	12.88
	Left-hand	91.42	28.07	95.76	20.19	91.49	27.96	95.38	21.04
	Right-hand	92.61	26.22	95.22	21.39	90.83	28.92	95.24	21.34