Phosphorylation Toggles the SARS-CoV-2 Nucleocapsid Protein Between Two Membrane-Associated Condensate States

Bruna Favetta[1], Huan Wang[2], Jasmine Cubuk[3], Arjun Singh[4], Mayur Barai[4], Cesar Ramirez[1], Adam J. Gormley[1], Sanjeeva Murthy[2], Gregory Dignon[4], Andrea Soranno[3], Zheng Shi[2], Benjamin S. Schuster[4]

1 Department of Biomedical Engineering, Rutgers, The State University of New Jersey, Piscataway, NJ 08854

2 Department of Chemistry and Chemical Biology, Rutgers, The State University of New Jersey, Piscataway, NJ 08854

3 Department of Biochemistry and Molecular Biophysics, Washington University in St Louis, St. Louis, MO 63110

4 Department of Chemical and Biochemical Engineering, Rutgers, The State University of New Jersey, Piscataway, NJ 08854

Corresponding author benjamin.schuster@rutgers.edu

Summary

The Nucleocapsid protein (N) of SARS-CoV-2 plays a critical role in the viral lifecycle by

regulating RNA replication and packaging the viral genome. N and RNA phase separate to form

condensates that may be important for these functions. Both functions occur at membrane

surfaces, but how N toggles between these two membrane-associated functional states is

unclear. Here, we reveal that phosphorylation switches how N condensates interact with

membranes, partly by modulating condensate material properties. Phosphorylated N forms liquid

condensates that wet membranes, reminiscent of N's role in RNA replication. In contrast,

unmodified N forms viscoelastic condensates that can be engulfed by membranes, evoking viral

genome packaging. These results suggest that phosphorylation serves as a regulatory switch

within the viral replication cycle by modulating N's association with membranes. We gained

mechanistic insight through structural analysis and molecular simulations, which suggest

phosphorylation induces a conformational change that softens condensate material properties.

Our studies also show that phosphorylation alters N's interaction with viral membrane proteins.

Together, our findings uncover a novel aspect of SARS-CoV-2 biology by identifying membrane

association as a key feature of N condensates and providing mechanistic insights into the

regulatory role of phosphorylation. Understanding this mechanism suggests potential therapeutic

targets for COVID infection.

Introduction

The COVID pandemic has focused attention on the mechanisms of SARS-CoV-2 viral

replication. SARS-CoV-2 is an enveloped virus with a non-segmented, positive-sense, single-

stranded, ~30 kb RNA genome[1]. In the virus core, genomic RNA (gRNA) is associated with

Nucleocapsid protein (N), forming a ribonucleoprotein (RNP) complex. N has several functions

during the viral life cycle but is primarily involved in protecting the viral RNA genome by binding,

condensing, and packaging it within the virion[2]. N has also been shown to be necessary for

efficient transcription and replication of viral RNA, and it additionally contributes to immune

evasion via sequestering stress granule proteins[3]. How N is regulated throughout the viral lifecycle

to perform its varied functions, and whether it is possible to therapeutically manipulate this

regulation to inhibit viral replication, remains unknown.

N is a multi-domain protein, consisting of two folded and three disordered domains (Fig.

1A), the diversity of which likely contributes to its ability to perform several functions throughout

the viral lifecycle[2]. The N-terminal folded domain (NTD) strongly binds with specific viral RNA

elements, stabilizing the RNP complex[4,5]. The C-terminal folded domain (CTD) mediates

dimerization of N, facilitating the formation of the helical nucleocapsid structure that protects the

viral RNA genome[6,7]. Both folded domains are flanked by disordered regions. The central

disordered region acts as a linker that is partially extended and thus retains some structure while

providing the protein conformational flexibility[8,9]. The serine/arginine (SR)-rich region (aa 175 –

206; Fig. 1B) within the linker is known to participate in both protein–protein and protein–RNA

interactions[10–12].

Importantly, this SR-rich region was identified as a site of phosphorylation of N (Fig. 1B)

that may regulate protein function, based on several lines of evidence. 1) At different stages of

the viral life cycle, N exists in two phosphorylation states. Abundant phosphorylated protein is

found inside infected cells, while unmodified protein is found within virions[13–18]. Regulation of

phosphorylation of N may act as a timer in the viral lifecycle, switching from replication,

transcription, and translation to assembly of new virions[19–21]. 2) Phosphorylation of the SR-rich

region of N has been shown to modify how the protein interacts with RNA, as well as affect the

transcription and translation of RNA[22,23]. 3) During viral replication, when N is likely

phosphorylated, it appears to form a thin layer around viral replication organelles (vROs), double

membrane vesicles filled with viral RNA commonly found in infected cells[16,24]. However, during

virion assembly, complexes of unmodified N and RNA remain as small, spherical structures while

the viral capsid is engulfed by the ER-Golgi intermediate complex (ERGIC) membrane[25]. Although

little attention has been given to N's interactions with membranes, these observations suggest

that phosphorylation of N and its membrane interactions are linked and may underlie N's multiple

functions. The goal of this paper is to elucidate this link between N phosphorylation and its

membrane interaction at the molecular level, providing insights critical for understanding COVID

infection.

We sought to understand the phosphorylation-dependent membrane interactions of N

through the lens of phase separation. N contains RNA-binding and disordered domains, which

enable N and RNA to interact in a dynamic manner. This results in the formation of droplets

enriched in protein and RNA, known as biomolecular condensates[3,8,26,27]. N undergoes phase

separation with RNA in vitro and in vivo[28,29]; experiments show that phosphorylation of N[19,26], type

of RNA[30,31], and temperature[30] modulate the condensate's propensity to phase separate and

affect its molecular dynamics. Biomolecular condensates can display a range of material

properties, from liquids to gels, which may be closely associated with their function[32,33]. The

material properties of a condensate need not be static, and cellular regulation can modulate

material properties and thus condensate function[34]. Carlson et al. showed that unmodified N and

RNA form gel-like condensates and discrete 15-nm particles, while phosphorylated N generates

a more liquid-like droplet. They hypothesized that this difference in material properties could be

the basis for a dual role of N during the viral lifecycle in both regulating RNA transcription and

facilitating nucleocapsid assembly[35,36]. However, how the phosphorylation-dependent material

properties of N condensates affect their interaction with membranes has not been studied. Recent

investigations on condensate-membrane interaction show that condensate material properties

are a key factor determining the mode of interaction between condensates and the membrane

surface[37–39]. Based on this, we hypothesized that phosphorylation tunes how N and RNA interact,

thus modulating the material properties of N and RNA condensates, consequently influencing

condensate-membrane interaction, and in turn allowing N to display the two distinct behaviors

observed throughout the viral life cycle (Fig. 1C-D).

Here, we tested this hypothesis using careful in vitro reconstitutions to map how

condensates formed by N and viral RNA fragments interact with membranes. We identified

membrane association as a feature of N condensates, and we found that this interaction is

regulated at least in part by N's phosphorylation state. Unmodified N condensates retain their

shape upon interaction with membranes, but phosphorylation reduces condensate viscoelasticity,

such that phosphorylated N (pN) condensates are softer and therefore wet membrane surfaces.

Phosphorylation also switches N's ability to interact with viral membrane proteins. These results

provide a mechanism by which N can toggle between distinct membrane-associated functions.

To understand these changes at a molecular level, we performed structural studies and molecular

simulations that revealed that phosphorylation strengthens interactions between pN monomers

within pN dimers and thus weakens protein-RNA binding. We propose a model where

phosphorylation acts as a functional switch during the viral lifecycle, shifting N's roles from RNA

replication to virion assembly by toggling its membrane interaction.

112
Results

We reconstituted N or pN and RNA condensates in vitro. We prepared recombinantly

expressed N that was either kept unmodified or was phosphorylated in vitro using GSK-3β and

SRPK1 kinases[17] (Fig. 1B). Up to 9 phosphorylation sites on N were confirmed using mass

spectrometry and Phos-Tag SDS-PAGE (Supplementary Fig. 1). We used in vitro transcription to

make fragments of viral RNA that are known to promote phase separation of N (Fig. 1E)[30]. First,

we tested a fragment containing the first 1000 base pairs from the 5' end of the viral genome that

contains important regulatory information[1]. Second, we tested the fragment of RNA that encodes

N (containing the first 75 nucleotides of the 5′ untranslated region recombined onto the N protein

coding sequence), given that it is highly produced during viral infection[1]. As expected, mixing N

or pN protein with either RNA at 37°C resulted in their condensation into protein- and RNA-rich

droplets (Fig. 1F). A change in morphology with regards to the degree of sphericity of droplets

already suggests that condensates are modulated by both type of RNA and phosphorylation

status of the protein (Supplementary Fig. 2), two factors that we investigate in depth.

Figure 1: Phosphorylation of the SARS-CoV-2 Nucleocapsid protein (N). A) Schematic of full-

length SARS-CoV-2 Nucleocapsid protein (N) used in experiments, showing N1, a disordered

domain; N2, RNA binding domain (PDB ID: 7VBD); N3, disordered linker domain containing the

Serine-Arginine-rich (SR) region; N4, dimerization domain (PDB ID: 6WZQ); and N5, a disordered

domain. B) Sites of phosphorylation in the SR region of the N3 domain. Our experiments used

two kinases – SRPK1 and GSK3β – resulting in a maximum of 10 phosphorylation sites. C)

Hypothesized form and roles of N and phosphorylated N (pN). pN may have dynamic functions,

forms liquid-like droplets with RNA and may localize to the surface of replication organelles.

Unmodified N forms solid-like, spherical assemblies with genomic RNA and facilitates new virus

encapsulation. N's form with RNA was characterized by Carlson et al., 2020, using light and

electron microscopy. D) A diagram of the SARS-CoV-2 lifecycle. Following viral entry (1), the

genome is uncoated (2) such that it can be read. Viral RNA is stored within viral replication

organelles (3) wherein RNA transcription and replication may be occurring. N binds to and

condenses genomic RNA exiting these organelles (4). In parallel, viral structural proteins are

produced (5), in preparation for viral encapsulation (6) and exit (7). E) Main RNA fragments used

in experiments are the first 1000 bases from the 5' end of the viral genome and the 1340 base

fragment encoding N. F) Droplets with varied morphologies form upon mixing 40 µM N (5% Alexa

Fluor 647 labeled) and 300 nM RNA fragments (5% Cy3 or Cy5 labeled) at 37˚C. Scale bar = 5

µm.

N condensate composition determines interaction with membranes

Prior work has shown that following viral infection, N accumulates in thin layers around

folded ER membranes that are likely vROs[40]. In contrast, later during viral budding events, N is

part of RNPs linked by the viral genome[25]. These RNPs do not fuse or grow during the budding

process, instead remaining as distinct complexes as the nucleocapsid is engulfed into new

virions[25,41]. We first asked whether we could reproduce these two behaviors of N in vitro.

We developed a system to study the interaction between N and RNA condensates and

membranes. We modeled membranes using giant unilamellar vesicles (GUVs). GUVs were made

with a lipid composition meant to approximate the human ER membrane[42,43], with 60% DOPC,

25% DOPE, 10% DOPS, and the addition of 5% Ni-NTA lipids. The Ni-NTA lipids allowed us to

tether membrane protein fragments to the GUV surface[44] to investigate whether the presence of

either of two viral membrane proteins would affect condensate-GUV interaction (Fig. 2A). The

viral non-structural protein 3 (Nsp3) localizes to vROs[45] and was suggested to both help form the

double layer of membranes as well as form pores that span the membranes[46–48]. Nsp3 is a known

interactor of N[49,50] and therefore is a logical candidate for modulating N's interaction with the

membranes enveloping vROs. We also studied the viral M protein that is present in the ERGIC

membrane where new virions form[51]. M and N are also known interactors, where M is thought to

anchor the RNP complex to the membrane during new virion formation[52,53]. Given the challenges

of producing and inserting transmembrane proteins into vesicles, we chose to study the domains

of the proteins that are known to interact with N. For M, we used a construct with the C-terminal

endodomain of the M protein, fused to a 6xHis tag and GFP[53]. For Nsp3, we took its ubiquitin-like

domain 1 (Ubl1)[49], and produced a construct with it fused to a 6xHis tag and GFP.

For this experiment, we used condensates composed of N or pN and the 1-1000 RNA

fragment. We added a small amount of GUVs and membrane protein fragments to the samples –

either Nsp3 or M fragments or a control solution (Fig. 2B). We observed the GFP signal from the

membrane proteins become localized to the surface of GUVs, with a similar degree of tethering

of Nsp3 and M fragments to the GUV surface (Supplementary Fig. 4). Next, we used optical

tweezers to control the position of condensates, moving them to and holding them at the GUV

surface. We then attempted to pull the condensates off the surface and observed whether they

remained bound or were mobile. For example, in a sample with pN, 1-1000 RNA, GUVs, and

Nsp3 fragments, we brought condensates composed of pN and 1-1000 RNA to the GUV and

observed as the condensates wet the surface (Fig. 2C).

We defined five types of interactions between condensates and membranes (Fig. 2D

legend): 1) With no interaction, the optical tweezer can move condensates away from a surface

it was in contact with. 2) With only binding, we observe the condensates remain attached to the

membrane surface even when the optical tweezer is attempting to dissociate the two, but no

additional condensate-membrane interactions occur. 3) During membrane wrapping, the

condensate binds to the membrane and the membrane surrounds the condensate over time. 4)

In partial wetting, the condensate not only binds to the membrane surface but also partially

deforms, expanding the area of contact between the condensate and membrane. 5) Finally, in

complete wetting, the condensates totally wet the membrane surface, forming a thin layer of

protein and RNA condensate.

We found that if no fragments of membrane proteins (Nsp3 or M) are present, no

interaction occurs between N and RNA condensates and the GUV surface, regardless of the

phosphorylation status of N (Fig. 2D and Movies S1-2). This suggests that N or pN and RNA

condensates have no intrinsic ability to interact with lipid membranes of the composition tested

(60% DOPC, 25% DOPE, 10% DOPS, 5% DGS-NTA(Ni), in 150 mM NaCl, 20 mM Tris-HCl, pH

7.5 buffer). In contrast, when the Nsp3 fragment is at the surface of GUVs, we observe

condensates composed of both N and pN binding to the membrane surface (Movies S3-4). With

unmodified N, condensates with 1-1000 RNA only partially wet the surface. Condensates with

phosphorylated N completely wet the surface, forming a thin layer of condensed material. This

behavior of pN and RNA condensates resembles the formation of a layer of protein on vROs that

is observed in infected cells[24]. Importantly, our results hint that by modulating the material

properties of N condensates, phosphorylation shifts the behavior observed from partial to

complete wetting. The thin layer of N that is observed surrounding vROs in infected cells may be

condensed pN wetting the surface due to pN-Nsp3 interactions.

When we repeated the experiment with the M fragment, we observed an important

difference in binding based on the status of N phosphorylation: while condensates with unmodified

N bound to the M-coated GUV surface, condensates with pN did not interact with the surface

(Movies S5-6). Therefore, phosphorylation of N may have unexpected effects beyond modulating

condensate material properties: it can act as a switch to prevent binding of pN to membrane

surfaces displaying the viral Membrane protein, where new virion formation typically occurs. (We

investigate how phosphorylation affects protein-protein interactions between N and M or Nsp3 in

Fig. 3). When condensates with N and 1-1000 RNA interacted with M-coated GUVs, we observed

binding (6/10 events) or membrane wrapping (4/10 events) (Fig. 2E), the difference between

either case likely driven by the membrane tension of the GUVs. In the cases where membrane

wrapping occurs, the protein and RNA condensates are partially engulfed into GUVs in the time

scale of minutes. The events of membrane wrapping are especially intriguing, given that they

resemble the direction of membrane bending required for encapsidation. These observations are

consistent with prior EM studies showing that RNPs may contribute to membrane bending after

recruitment to the membrane by M[54,55].

Given the important role that cholesterol plays in determining membrane fluidity, and

recent evidence pointing to its modulation of condensate-membrane interactions[56], we sought to

test the effect of adding cholesterol to our GUV membranes. Viral replication organelles originate

from ER membranes, which are known to be low in cholesterol content (~2 – 5%)[43]. Although the

SARS-CoV-2 virus buds from the nearby ERGIC[57], their membranes were found to include much

higher percentages of cholesterol (~25%)[58]. Therefore, we tested the effect of inclusion of 2, 5,

and 25% cholesterol on GUV membranes in place of DOPC (Supplementary Fig. 3). Addition of

cholesterol did not affect the binding interaction of condensates with membranes lacking viral

membrane fragments. Next, we observed that addition of cholesterol reduced the degree of

condensate wetting in cases where the GUV was coated with Nsp3 fragments, in a concentration-

dependent manner. However, as noted above, the ER-derived vRO membrane where N-Nsp3

interaction occurs is known to have low cholesterol content, and therefore we speculate that

cholesterol has a minor effect on N wetting to viral replication organelles. Finally, inclusion of

cholesterol did not affect the type of condensate-membrane interaction observed with M fragment

present. We conclude that although cholesterol can affect condensate-membrane interaction, in

the particular cases that are most physiologically relevant to SARS-CoV-2 infection, addition of

cholesterol did not change the interaction type we observed.

We quantified the affinity between condensates and GUVs by measuring the geometric

factor that considers the contact angles along the contact line between the GUV surface, the

condensate, and the external solution, as well as the intrinsic contact angle between condensate

and GUV (Fig. 2F-H)[39]. These are determined by the material properties of the condensate, the

strength of interaction between condensate and membrane surface, and the membrane tension

of the GUV. No interaction between condensates and the membrane results in the highest

geometric factors. An intermediate geometric factor is found with N condensates and Nsp3

fragments, where partial wetting is observed, and N condensates with M fragments, where either

binding or membrane wrapping are observed. A drastic reduction in geometric factor occurs with

pN and Nsp3, where complete wetting occurs. The different degrees of interaction also impact

the intrinsic contact angle between condensate and GUV. Partial wetting or membrane wrapping

cause an intermediate reduction in the angle, while complete wetting causes a drastic reduction

in the angle (Fig. 2H). Overall, with this panel of condensate compositions and membrane

surfaces, we were able to reproduce several behaviors that N displays during the viral lifecycle.

Figure 2: N condensate interaction with membranes. A) Fragments of the membrane proteins

used to test membrane interactions: C-terminal endodomain of the Membrane (M) protein and

the ubiquitin-like domain (Ubl1) of the Nsp3 protein. Both protein fragments are fused to a 6x

Histidine tag and GFP. B) Schematic of the experimental setup showing the M or Nsp3 protein

fragment tethered to the giant unilamellar vesicle (GUV) surface. Condensates composed of 40

µM N or pN plus 300 nM RNA are moved to the GUV surface using optical tweezers. C)

Representative widefield images showing that Nsp3 fragments preferentially localize to the GUV

surface (left) and condensates composed of pN and 1-1000 RNA wet the surface of GUVs after

being delivered to the surface using optical tweezers (OT) (right). White dashed line added to

denote the GUV surface. D) Representative widefield images showing the interaction between

condensates and membranes over time, for N vs. pN, and comparing GUVs with no membrane

protein, M fragment, and Nsp3 fragment. GUVs are labeled in green, and condensates (Cond.),

in blue. Interaction type is qualitatively classified. E) Timelapse imaging showing several

condensates (blue stars) being dragged by optical tweezers to the surface of a GUV, to which

they bind. Optical tweezers can be used to pull bound condensates which results in deformation

of the GUV surface (white arrows). Membrane wrapping occurs in one case. F) Sketch showing

the parameters that define the geometric factor (Φ) and intrinsic contact angle (θ_{in}). Interaction

leads to the formation of a contact line between the condensate interface and membrane. Contact

angles $\theta_c + \theta_e + \theta_i = 360°$ and we can calculate Φ to describe the degree of wetting. Wetting can

also be characterized by θ_{in} between condensate and membrane surfaces. G) Quantification of

the geometric factor from each combination of condensate composition and membrane surface.

Interactions that lead to membrane wrapping, partial wetting, or complete wetting result in reduced

geometric factors. H) Intrinsic contact angle from each combination of condensate composition

and membrane surface, calculated from the same image data set as G. n = 10 condensate-

membrane interactions from n = 3 independent trials. Individual data points are shown for each

interaction event. The lines indicate the median, lower quartile, and upper quartile. Scale bars =

5 µm.

Phosphorylated N cannot bind to the SARS-CoV-2 Membrane protein

Given our observation that pN condensates did not bind to GUVs with M fragments, we

examined whether phosphorylation affects protein-protein interactions between N and M or Nsp3.

Several groups have sought to understand the binding mechanism between coronavirus N and

M proteins[52,53], but no information is available on the effect of phosphorylation of N on its

interaction with M. A stretch of amino acids (168–208) within the linker region of N was identified

as critical for N-M interactions in the SARS-CoV-1 virus[52]. More recently, the linker domain of the

SARS-CoV-2 N was shown to be necessary for N to co-phase separate with M[53]. Together, these

studies point to a potential role of the SR-rich domain of N in interacting with M. Our results from

Fig. 2 suggest that phosphorylation of the SR region of N may inhibit binding between N and M.

We performed a partitioning experiment using confocal microscopy (Fig. 3A). We added

the fluorescently tagged M fragment or Nsp3 fragment to N or pN and observed the partitioning

of the membrane protein fragments into condensates (Fig. 3B-C). We confirmed that M has a

much stronger affinity to unmodified N than to pN. Unmodified N binds to M independent of the

presence of viral RNA fragments, though phase separation is promoted by the presence of 1-

1000 RNA. We observed contrasting results with phosphorylated N mixed with M. First, M does

not drive the condensation of pN without addition of RNA. Second, M does not partition

preferentially into condensates composed of pN and viral RNA (partition coefficient for M is 9.8 ±

0.3 in N + 1-1000 RNA condensates vs. 0.3 ± 0.1 in pN + 1-1000 RNA condensates; partition

coefficient is defined as the ratio of average fluorescence intensity inside vs. outside the

condensates). Given that M likely interacts with residues around N's SR domain[52], it is not

surprising that phosphorylation of the SR region affects binding between the two proteins. These

results are intriguing because they suggest a mechanism for timing of viral assembly. As noted

above, a majority of N within infected cells is phosphorylated, but N included within new virions is

unmodified[16]. pN may not be incorporated into newly formed virions because this protein is unable

to bind M and anchor the RNP to the membrane at the site of viral assembly.

As a comparison, we also investigated whether phosphorylation affects the partitioning of

Nsp3 protein's Ubl1 domain into N or pN condensates. Previous reports showed that Nsp3 Ubl1

domain partitions into condensates composed of both N and a phosphomimetic version of N[19].

Our results qualitatively agree with these findings, although we found pN has a lower affinity than

N for Nsp3, based on our measured partition coefficients (12.6 ± 0.1 for Nsp3 in N + 1-1000 RNA

condensates vs. 2.2 ± 0.1 in pN + 1-1000 RNA condensates). Given the hypothesized role of N

in delivering the viral genome to Nsp3-coated replication organelles following viral entry into cells,

and given that the films of N observed adhered to replication organelles[24] are likely composed of

phosphorylated N, it is not surprising that both unmodified and phosphorylated N bind to Nsp3.

Figure 3: N and pN interaction with membrane proteins, M and Nsp3. A) Schematic of the experimental setup where 6xHis-GFP-M$^{104-222}$ or 6xHis-GFP-Nsp3$^{1-111}$ is added to a condensate sample and partitioning of GFP-tagged protein is quantified using confocal microscopy. B) Representative confocal images from condensates composed of 40 μM N or pN plus either no RNA or 300 nM RNA; plus either 2 μM 6xHis-GFP-M$^{104-222}$ or 6xHis-GFP-Nsp3$^{1-111}$. 6xHis-GFP-M$^{104-222}$ binds to N but not pN condensates while 6xHis-GFP-Nsp3$^{1-111}$ partitions in regardless of

choice of N vs. pN. Normalized line profiles across condensates, representing averages of at least

n = 20 condensates from 3 independent trials. Scale bars = 5 µm. C) Quantification of partitioning

of GFP-tagged proteins into N or pN condensates by dividing average fluorescence inside

condensates by the average background fluorescence. Error bars represent one standard

deviation (±1 s.d.). p values were determined using two-way ANOVA followed by post hoc Tukey's

test. *p < 0.01.

Phosphorylation of N modulates the material properties of N condensates

Based on previous data on the liquid-like properties of pN condensates[19] and our

membrane binding experiments, we hypothesized that the dynamic nature of pN condensates is

what enables them to relax at the surface of GUVs. To explore the material properties of

condensates, we first used Fluorescence Recovery after Photobleaching (FRAP). We

photobleached either fluorescently labeled protein or RNA within a region of the condensate and

observed how the fluorescence recovered over time. The advantage of using FRAP here is that

it allows us to independently assess the dynamics of protein and RNA in our multi-component

condensates. N bound to 1-1000 RNA recovers more slowly than pN bound to the same RNA

(recovery half-life τ = 2.6 ± 0.1 min for N and 1.7 ± 0.1 min for pN) (Fig. 4A-B and Supplementary

Fig. 5). This is in agreement with previous data showing a more rapid FRAP recovery in

condensates with a phosphomimetic version of N compared to unphosphorylated N[19]. In addition,

we also observe that the total recovery of N is lower compared to pN (62% vs. 94%), suggesting

a pool of N is fixed to the RNA. Both metrics suggest that pN has a greater mobility than N when

bound to 1-1000 RNA. In contrast to the results obtained by photobleaching protein within

condensates, we observed almost no recovery of the fluorescence when RNA was bleached (Fig.

4A and C), independent of protein phosphorylation status. RNA appears to form a network onto

which protein can bind and dissociate[59]. Individual RNA molecules may move locally but do not

appear to diffuse long distances, likely due to their ability to form intermolecular base pairs.

Overall, our FRAP results support our hypothesis that differences in material properties contribute

in part to the type of condensate-membrane interaction observed. For example, condensates

composed of pN (which can wet surfaces) showed a greater protein recovery in FRAP

experiments, suggesting that the dynamic movement of pN allows the condensates to reorganize

at the surface of membranes over time.

To quantitatively assess how phosphorylation affects N condensate material properties,

we turned to passive microrheology. In passive microrheology, we embed 500 nm fluorescent

tracer beads into condensates and track their movement. The mean squared displacement (MSD)

of beads depends on their viscous and elastic environment[60]. Beads embedded within

condensates composed of pN + 1-1000 RNA displayed a greater MSD at all lag times when

compared to beads embedded in N + 1-1000 RNA condensates (Fig. 4D and Supplementary Fig.

6). The MSD of beads did not increase linearly with lag time for all samples, revealing that some

condensates behave as viscoelastic fluids under the experimental conditions. Using the

Generalized Stokes-Einstein Relation, we estimated viscous and elastic moduli. For N + 1-1000

RNA condensates, the elastic modulus dominates the viscous modulus at high frequencies (> 4

Hz). In contrast, for pN + 1-1000 RNA, the viscous modulus is dominant for all frequencies

measured (Fig. 4E – F). We then quantified the zero-shear viscosity of the samples, i.e., the

limiting value for viscosity at 0 Hz. For condensates composed of N + 1-1000 RNA, we found a

viscosity of 192 ± 3.6 Pa*s, while for condensates with pN, we measured a viscosity of 59 ± 3.4

Pa*s, representing a ~3x reduction in viscosity (Fig. 4G and Supplementary Fig. S7).

The change in viscosity upon phosphorylation of N may be due to modulation of protein-

protein and/or protein-RNA interactions, the latter via either specific or nonspecific binding to RNA.

We assessed each of these possibilities by testing the material properties of N or pN condensates

with unstructured RNA (polyrA) or with 5% PEG8000, which acts as a crowding agent to drive N

phase separation without RNA (Supplementary Fig. 2). Phosphorylation of N decreased the

viscosity of condensates with polyrA from 64 ± 4.2 Pa*s to 12 ± 0.4 Pa*s, suggesting that

nonspecific interactions between protein and RNA were disrupted. Furthermore, phosphorylation

decreased viscosity of condensates with PEG8000 from 11 ± 0.7 Pa*s to 4 ± 0.3 Pa*s, suggesting

that protein-protein interactions are also disrupted to some degree (Fig. 4G).

We also quantified the terminal relaxation time of the condensate network, which is the

inverse of the crossover frequency between the predominantly elastic and predominantly viscous

regimes. At timescales below the relaxation time, the condensates behave as elastic materials.

Elasticity is reduced by phosphorylation in both polyrA and 1-1000 RNA samples (Fig. 4H).

Condensates with N and polyrA or 1-1000 RNA have a relaxation time of 0.02 and 0.3 seconds,

respectively, while condensates with pN have no measurable relaxation timescale, suggesting

that phosphorylation disrupts the ability of N and RNA to crosslink (Supplementary Fig. 8). The

elasticity displayed by complexes formed between unmodified N and RNA may have a biological

function in protecting the RNA from mechanical stress[61].

We confirmed our material property measurements using micropipette aspiration

(MPA)[62,63]. We recorded how the length of the aspirated condensate in a micropipette changes

as a function of time in response to an applied pressure. We quantified the viscosity for the

different protein combinations and obtained the same trend and rank order that was observed

using microrheology (Fig. 4I and Supplementary Fig. 9). Viscosities quantified via MPA were

consistently lower than the zero-shear viscosities obtained from microrheology; when the two data

sets are plotted against each other, they have a linear fit with slope of 0.72 and R^2 of 0.99. Notably,

MPA measurements also showed that pN condensates have lower viscosity than the

corresponding N condensates. Together, our microrheology and micropipette aspiration results

point to N and RNA condensates being viscoelastic fluids whose properties depend on both RNA

type and phosphorylation status. Phosphorylation of N appears to loosen the network of

interactions between N and RNA, thus reducing the viscosity and elasticity of condensates.

Figure 4: Phosphorylation modulates material properties of N and RNA condensates. A)

Fluorescence recovery after photobleaching (FRAP) of N and RNA condensates, in which 5% of

N/pN is labeled with Alexa-647 and 5% of RNA is labeled with Cy-3. pN recovers to a greater

extent over time when compared to N, while in neither case does RNA signal recover. Scale bar

= 5 µm. B) Quantification of FRAP of protein in N vs. pN condensates (n = 3 independent trials).

C) Quantification of FRAP of RNA in N vs. pN condensates (n = 3 independent trials). D)

Ensemble MSD versus lag time (prior to noise correction) for the protein and RNA combinations

tested in this study. Inset: Representative trajectories from two-dimensional particle tracking

showing Brownian motion of beads in N vs. pN condensates with 1-1000 RNA. Each tick

represents 5 nm. E) Plot with the average viscous modulus (G", red) and the average elastic

modulus (G', blue) of N + 1-1000 RNA condensates as calculated from the MSDs (n ≥ 10 videos

from 3 independent samples) after noise correction. F) The average viscous and elastic moduli

after noise correction for pN + 1-1000 RNA condensates showing no crossover frequency in the

range studied. G) The zero-shear viscosity of the protein and RNA condensates studied,

calculated from the particle-tracking results after noise correction. Data from n ≥ 10 videos from 3

independent trials. H) Quantification of the timescales at which the elastic modulus dominates

(color) versus the viscous modulus dominates (grey) in protein and RNA condensates. I) Viscosity

of the protein and RNA condensates from micropipette aspiration. n = 3 independent trials. Error

bars represent one standard deviation (±1 s.d.).

RNA type also modulates condensate material properties

We repeated experiments with a second viral RNA fragment to assess whether our results

depend on RNA sequence and structure. RNA sequence and structure are known to affect the

material properties of other condensates, though most quantitative studies so far have focused

on shorter, artificial RNA sequences[64,65]. For N protein, evidence exists that the morphology of

condensates depends on the type of RNA included[66], however, the effect on condensate material

properties has not been investigated. Here, we compare our results with 1-1000 RNA with N RNA,

introduced previously, which encodes the N protein and is 1340 base pairs long. N RNA is an

important viral RNA to test because it is abundant in infected cells, and from a biophysical

perspective, it is (1) longer than 1-1000 RNA, and as such has a higher ensemble diversity (which

describes the diversity of conformations the RNA is predicted to fold into; it is predicted to be 198.5

for N RNA vs. 148.1 for 1-1000 RNA, based on ViennaRNA[67]), and (2) contains a different pattern

of preferred binding sites for N[66]. These factors suggest that N RNA forms a more entangled

network in condensates compared to 1-1000 RNA, resulting in the different condensate

morphologies observed (Fig. 1F). We found similar behavior between condensates with 1-1000

or N RNA (Supplementary Fig. 10), both in terms of binding to GUVs with different membrane

protein fragments, as well as their material response to phosphorylation (phosphorylation reduces

viscosity and elasticity). These results suggest that the change in N condensate material

properties and membrane interaction upon phosphorylation are not RNA structure dependent.

Across experiments, we did observe that condensates with the longer N RNA were less likely to

deform than their counterparts with 1-1000 RNA. This data hints that complexes of unmodified N

and longer fragments of RNA, such as genomic RNA, would appear solid-like and with an

important elastic response, which may play a role in mechanically protecting RNA within virions[61].

To further study these questions, additional analyses with varying RNA lengths and structures

would be required.

Phosphorylation weakens binding between N and RNA by promoting intra-dimer

interactions

We have shown that phosphorylation of N results in condensates that more readily deform

at membrane surfaces and that have lower viscosity and elasticity, both likely driven by a

loosening of the interaction network between protein and RNA. This change in interaction does

not appear to be dependent on specific RNA structures, as phosphorylation also drastically

softens condensates with unstructured polyrA (Fig. 4G-I). We therefore asked how

phosphorylation affects N, such that interaction between protein and RNA is diminished.

We leveraged fluorescence polarization to measure the binding affinity between N protein

and fluorescently labelled RNA (Fig. 5A-B). First, we assessed whether phosphorylation affects

binding of N to a stem-loop structure (SL4) present in the 1-1000 RNA fragment. SL4 was

identified as a preferred binding site for N's RNA binding domain[30,31]. Phosphorylation of N

reduces the binding affinity to SL4 RNA, increasing the dissociation constant (K_D) from 10.9 ± 1.4

nM to 59.0 ± 0.9 nM. Although this is a six-fold reduction in binding affinity, the concentrations at

which our membrane interaction and material property experiments were conducted are well

above the nanomolar K_{DS} measured, and therefore this change in binding cannot explain the

material property changes observed. Next, to test the hypothesis that nonspecific binding between

protein and RNA is disrupted by phosphorylation, we measured the binding affinity between N or

pN and a 30-base polyrA. Phosphorylation causes a twenty-fold change in K_D from 71.7 ± 0.8 nM

to 1571.0 ± 2.1 nM, larger than that observed for the SL4 RNA. We conclude that phosphorylation

reduces N's interaction with unstructured RNA more so than structured RNA. Importantly, for both

polyrA and SL4, phosphorylation of N also reduced the maximum polarization. This suggests a

greater rotational freedom in the protein-RNA complexes formed with pN that could be driven

either by the formation of a more compact complex or a conformation in which the protein-bound

RNA retains more rotational mobility[68]. We anticipated that phosphorylation would weaken binding

between N and RNA due to electrostatic repulsion. However, our fluorescence polarization results

suggest that phosphorylation may also be inducing a conformational change in N that affects its

RNA binding properties.

Protein + RNA	K_D (nM)	Δ Polarization (mP)
N + SL4 RNA	10.9 ± 1.4	252.2 ± 6.2
pN + SL4 RNA	59.0 ± 0.9	152.6 ± 3.5

Protein + RNA	K_D (nM)	Δ Polarization (mP)
N + polyrA	71.7 ± 0.8	217.1 ± 5.8
pN + polyrA	1571.0 ± 2.1	106.2 ± 2.3

Figure 5: N protein phosphorylation weakens RNA binding affinity. A) Quantification of

binding affinity between N vs. pN and the viral stem loop 4 RNA (SL4) based on a change in

normalized fluorescence polarization (minimum polarization set to 0). n = 3 independent trials. B)

Quantification of binding affinity between N vs. pN and unstructured 30 base polyrA from

normalized fluorescence polarization. n = 3 independent trials.

To assess the conformational change of the N protein dimer following phosphorylation, we

used Small Angle X-ray Scattering (SAXS), a technique compatible with the size and largely

disordered nature of N protein. Our SAXS results indeed show that N and pN are partially

disordered dimers in solution. This is in agreement with previously published data on unmodified

N protein;[9] SAXS of pN had not been previously reported. With phosphorylation, we found a

significant decrease in radius of gyration (R_g), from 60.2 ± 1.3 Å for N to 54.4 ± 2.0 Å for pN, a

finding supported by both the pair distance distribution and the Guinier approximation (the

decrease in R_g was consistent across concentrations) (Fig. 6A-B and Supplementary Fig. 11-12).

The shape of the pair distance distribution function also changes with phosphorylation. The

presence of multiple peaks in the unmodified N data highlights the multi-domain nature of the N

protein dimer[69]. The lack of these peaks in the phosphorylated N data suggests the domains can

no longer be resolved, indicating there may be new interactions bringing the different parts of the

protein together. Next, analyzing normalized Kratky plots confirms both N and pN display behavior

typical of a multidomain protein with flexible linkers (Fig. 6C)[70]. The elevated tail for N indicates

greater extension of the protein compared to pN. Using the scattering data to construct bead

models for N and pN allows us to visualize these changes (Fig. 6D-E and Supplementary Fig.

13). Our bead models show a new region of electron density between monomers within pN

dimers, suggesting that indeed new interactions within pN dimers take place that are not present

in N dimers.

To better understand how the conformation of N changes upon phosphorylation, we used

single-molecule Förster Resonance Energy Transfer (smFRET). First, for full-length N, we

confirmed that phosphorylation does not affect the conformation of the dimerization domain[71] or

the tendency of N to dimerize (Supplementary Fig. 14). Next, we used smFRET to study the

conformation of the linker region. We probed two constructs of N with fluorescent labels flanking

the linker region (at residues 172 and 245): full-length N and truncated N[1-246], which lacks the

dimerization and C-terminal disordered domains. We performed experiments at two protein

concentrations: low concentration (100 pM labeled protein), at which N is in its monomeric form,

and high concentration (100 pM labeled protein + 1 μM unlabeled protein for full-length N or 4 μM

unlabeled protein for N[1-246]), at which dimers form if the dimerization domain is present. We

measured the distribution of transfer efficiencies for each protein construct at each concentration

(Supplementary Fig. 15). The distributions represent a dynamic ensemble of conformations, as

supported by the corresponding analysis of donor lifetime vs. transfer efficiency (Supplementary

Fig. 15). Therefore, we used the mean transfer efficiencies to calculate the root mean square

distance (RMSD) between labelled residues, which can be compared across samples to

understand the degree of expansion of the linker (Supplementary Fig. 15). Consistent with

previous measurements of the same constructs[8], we observed a mean transfer efficiency across

the linker of approximately 0.6 for full-length N and 0.75 for N[1-246]. We found that phosphorylation

causes an expansion of the linker region, which was unexpected given that we previously

measured by SAXS a reduction in radius of gyration of the N dimer with phosphorylation.

However, analysis of the sequence charge decoration (SCD) of the linker supports our smFRET

findings and suggests that addition of several negative charges to the domain causes local

repulsion that results in expansion[72] (Supplementary Fig. 15). Importantly, our smFRET results

show that the degree of expansion due to phosphorylation is greatly reduced in the full-length

construct under dimer conditions. To reconcile the SAXS and smFRET results, one possibility is

that phosphorylation of the N dimer drives formation of new interactions across linker domains of

the dimer that counteract the expansion caused by phosphorylation.

To explore the new interactions that arise upon N protein phosphorylation, we conducted

coarse-grained molecular simulations of the protein. We combined the HPS-Urry model for

simulation of disordered proteins[73,74] with an elastic network model and Center of Mass (COM)

bead placement scheme for modeling folded domains[75]. To mimic the effects of phosphorylation,

we used phosphomimetic mutations (S to D and T to E). We found reasonable agreement

between the radii of gyration measured in SAXS and those predicted in the simulations, which

were 55.5 Å for N and 52.4 Å for phosphomimetic N (SAXS results were 60.2 ± 1.3 Å for N to 54.4

± 2.0 Å for pN). Fig. 6F depicts three conformations of N protein that were captured by the

simulation, with a range of radii of gyration. Next, we compared the pair distance distributions

from our simulations of N and phosphomimetic N protein with SAXS data (Fig. 6G-H). Although

the simulation captured the overall collapse of N upon phosphorylation, it did not fully capture the

change in conformation of the protein that we observed through SAXS, which is reflected in the

shape of the pair distance distribution function[69]. This indicates that while the simulation model

may capture accurate conformations, it lacks the level of detail needed to consistently match

experimental data across the full ensemble.

To refine our simulation, we incorporated information from our SAXS experiments. We

used a Bayesian Maximum Entropy ensemble reweighting scheme to adjust the statistical weights

of the simulated conformations so that their pair distance distribution better matches the

experimental SAXS data[76] (Supplementary Fig. 16). This helps identify subsets of conformations

that explain the experimental data, which may have been undersampled due to shortcomings in

the simulation or missing environmental effects. For example, the phosphomimetic mutations

used in simulations lack the additional steric bulk of phosphate groups and simulations may not

fully capture how the ionic conditions in experiments modulate electrostatic interactions involving

the phosphate groups[77]. Upon reweighting, the radii of gyration obtained for N and

phosphomimetic N more closely matched the values obtained from SAXS, with 58.5 Å for N and

55.0 Å for phosphomimetic N (Supplementary Fig. 17A-B). The pair distance distribution function

from the reweighted simulation also more closely resembled the data obtained from SAXS (Fig.

6F-G). For example, with reweighting, the discrepancy between the pair distance distribution for

N reduces from a χ^2 of 7.5 to 2.2. Finally, with reweighting, we also observe agreement between

simulations and smFRET results regarding the expansion of the linker domain (Supplementary

Fig. 17).

We then investigated the change in intra- and inter-domain interactions predicted by the

simulation with phosphomimetic mutations (Fig. 6I-J). With the simulation, we can quantify the

change in number of contacts between residues both within each monomer and across monomers

in a dimer. We found that most intramonomer interactions decrease in the phosphomimetic protein

compared to unmodified N, with the exception of N2-N2 and N3-N4 interactions. While the N2

domain was almost entirely held folded by the harmonic constraints of the simulation, the last 13

residues of the domain were left flexible due to their high b-factor in the reference crystal structure.

Following the phosphomimetic mutations to the nearby linker (N3), these 13 residues interact with

the folded part of the N2 domain at a higher frequency (Supplementary Fig. 18A). Separately, the

increase in N3-N4 interactions observed is likely driven by the negative charges added to the N3

domain interacting with a positive patch of residues in the beginning of the N4 folded domain

(Supplementary Fig. 18B). Interestingly, we observe a significant decrease in intramonomer

interaction within the N3 domain, which matches our smFRET observations. The main driver of

this loss of interaction is a decrease in contact between nearby residues of the SR-rich region

following the addition of several negative charges due to the phosphomimetic mutations

(Supplementary Fig. 18C). In terms of intermonomer interactions, we found a gain in interaction

for almost all combinations of domains of N. In particular, we observe that the linker (N3) gains

contacts with the other monomer's N2, N3, and N4 domains. The N3 domain gains interactions

with an arginine-rich patch on the N2 domain and it interacts directly with the SR-rich region of

the opposite N3 domain, which is likely also driven by interactions with arginine residues

(Supplementary Fig. 18D-E). Finally, the N3 domains interacts with a positively charged patch in

the N4 domain, which is rich in both arginine and lysine residues (Supplementary Fig. 18F).

Therefore, binding between the negatively charged phosphomimetic residues and positively

charged arginine and lysine residues are the main source of new interactions within the

phosphomimetic N dimers.

Together, our SAXS, smFRET, and simulation results support that a conformational

change in N occurs as a result of phosphorylation. Previous molecular simulation of a fragment

of N containing the linker domain (N3) had found a phosphorylation-induced increase in intra- and

inter-molecular contacts due to the formation of salt bridges between phosphate groups and

arginine side chains[26]. Our results partly agree with these findings: while intramonomer N3

interactions decrease, binding between arginine residues and phosphate groups across members

of a pN dimer appears to be the basis for the new inter-monomer interactions observed. Overall,

we speculate that new interactions within and between monomers occupy the linker and thus

interfere with N's ability to bind to RNA. The more compact conformation of pN dimers and

reduced affinity between protein and RNA[10] explain our earlier observations from the binding

affinity assay and material property measurements. Therefore, this new model ties together the

conformational change of N upon phosphorylation with its reduced ability to interact with RNA,

which in turn affects its condensate material properties and its form when interacting with

membranes.

Figure 6: N protein phosphorylation weakens RNA binding affinity due to change in protein conformation. A) Representative pairwise interatomic distance distribution P(r) derived from

SAXS for N and pN. N or pN concentration = 3 mg/mL. B) The maximum dimension (D_{max}) and

radius of gyration (R_g) for N and pN derived from the pair distance distributions. Error bars

represent one standard deviation (±1 s.d.). p values were determined using two-sided student's

t-test; asterisk indicates $p < 0.05$. C) Normalized Kratky plot comparing the scattering of N vs. pN,

indicating a structural change has occurred due to phosphorylation. Concentrations shown for N

or pN are 1, 1.5, and 3 mg/mL from lighter to darker. D) Bead model representation for the N

dimer developed from SAXS results (left) and hypothesized conformation of N (right). E) Bead

model representation for the pN dimer developed from SAXS results (left) and hypothesized

conformation of pN highlighting new intermolecular interactions (right). F) Snapshots from N

simulation depicting conformations that demonstrate the protein's representative collapsed and

extended states. The radius of gyration of each selected conformation is noted. G) Comparison

of the pairwise interatomic distance distribution P(r) derived from SAXS and molecular dynamics

simulation before and after reweighting with SAXS data for N. H) Comparison of the pairwise

interatomic distance distribution P(r) derived from SAXS and molecular dynamics simulation

before and after reweighting with SAXS data for pN. I) Change in contact probability between all

residues within a dimer, compared between N and phosphomimetic N simulations. Lower left and

upper right quadrants represent intramonomer interactions while the upper left and lower right

quadrants represent intermonomer interactions. Blue coloring indicates more frequent

interactions occurred in phosphomimetic N compared to unmodified. J) Analysis of the difference

in intra-monomer (left) and inter-monomer (right) interactions between N and phosphomimetic N,

per domain of the protein (as defined by the schematic of N above). Positive numbers / blue

coloring indicates more frequent interactions occurred in phosphomimetic N compared to

unmodified. Below, sum of change in interactions across each domain for both intra- and inter-

domain interactions.

Discussion

Despite significant interest, there is still limited knowledge about how the SARS-CoV-2 N

protein is regulated during viral replication. Our work was motivated by three open questions.

First, N is thought to have several functions during viral replication, including regulating viral RNA

transcription and translation, and facilitating encapsulation of genomic RNA[2,3]. How can one

protein perform such distinct functions? Second, N's regulatory functions likely occur near viral

replication organelles[40], while its role in RNA encapsulation is known to take place at the surface

of the ERGIC membrane, where viral budding occurs[25]. How does N associate with these different

membranes? Third, N is known to phase separate in vitro and when expressed in cells[3,30,53].

However, is phase separation important for the function of N during viral replication? Building

upon prior research, we reasoned that phosphorylation holds a clue to answering these three

questions. Phosphorylation was previously hypothesized to be the key mechanism of regulation

of N during viral replication[19]. Two populations of N exist inside infected cells, with the majority of

N being phosphorylated, but with N that is incorporated into new virions remaining unmodified[13–

17]. Phosphorylation was also shown to alter the behavior of N condensates[19]. We combined these

observations to formulate a new hypothesis: that phosphorylation may be responsible for toggling

N between its functions by modulating the material properties of N condensates and thus

switching between its two membrane-associated states.

In this work, we reveal membrane association as a key feature of N phase separation. We

argue that the phosphorylation state of N determines its two membrane-associated behaviors in

infected cells: 1) being adhered to membrane surfaces and 2) being tightly bound to RNA when

engulfed into new virions. Our experiments show that phosphorylation modulates the material

properties of N and RNA condensates, and thus the type of interaction between condensates and

membranes, which is important for the function of N. We next studied N at a molecular level and

identified that phosphorylation results in new interactions within and between the monomers of a

dimer, thus interfering with the protein's ability to interact with RNA. We propose a model that ties

together an understanding of how phosphorylation acts as a switch controlling the conformation

of N and its behavior during viral replication (Fig. 7). In its phosphorylated form, N prefers

intermolecular interactions between N monomers within a dimer over interaction with RNA, thus

resulting in a looser network formation, condensates that can remodel at the membrane surface,

and RNA that is more accessible during replication steps. In its unmodified form, N binds tightly

to RNA, and facilitates its encapsulation after being anchored to the membrane surface by N-M

interactions. Through this model, we suggest that phase separation is critical to SARS-CoV-2

replication and that phosphorylation is a key mechanism through which phase separation is

regulated to allow N to complete its functions at membrane surfaces.

Figure 7: Model of N protein form and membrane-associated state during the viral lifecycle.

Inside the cell (1), the vast majority of N is found in its phosphorylated form, localized to the

surface of viral replication organelles (2). Phosphorylation promotes linker-linker interactions

across N dimers that weaken interaction between N and RNA (inset, left). This looser protein-RNA

interaction network within condensates results in a relatively low viscosity and elasticity that may

facilitate the molecular diffusion needed during RNA transcription and replication. A second

population of N that is destined to form new virions binds to new viral genomic RNA (gRNA) and

condenses RNA into small spherical complexes (3). This unmodified protein binds tightly to gRNA

through both specific and nonspecific interactions (inset, right). These assemblies have high

viscosity and elasticity that may support a protective function of N towards gRNA. The gRNA and

N capsid is engulfed by the ERGIC membrane, facilitated by N and M interactions (4). New virions

exit the infected cells (5).

Our results are distinct from recently published data on the effect of phosphorylation on

the molecular conformation of N determined using NMR. Botova et al. found a rigidification of the

SR region upon phosphorylation that their data suggests is a result of new interactions with the

RNA binding domain[78]. Their conclusions are due to the similarity in the chemical shift

perturbations observed when N is phosphorylated vs. when RNA binding occurs. However, the

rigidification of the SR region could also be due to the presence of bulky phosphate groups.

Though new interactions between N's RNA binding domain and the linker region could be

overlooked by our SAXS-derived reconstruction given its low resolution, our results point to a

larger conformational shift of the dimer. Stuwe et al. also studied the effect of phosphorylation on

N using NMR, though the construct studied lacked the N-terminal RNA binding domain, thus

potentially skewing the analysis of intra- and inter- molecular interactions[79]. They found that

phosphorylation changes the behavior of a leucine-rich helix (LRH) found in N's linker: in

unmodified N, higher order oligomers form through LRH self-interaction that is significantly

weakened by phosphorylation. In contrast with these results, our data presents an alternative

model in which phosphorylation of N promotes intermolecular interactions between the SR

regions of monomers within a dimer and thus disrupts non-specific interaction between pN and

unstructured RNA. We do not observe a significant change in the oligomerization behavior of N

following phosphorylation (Supplementary Fig. 14).

Our findings suggest several pathways towards development of novel COVID

therapeutics. One is to target the relevant kinases. A recent study suggested that N in cells is

sequentially phosphorylated by host kinases SRPK1, GSK-3, and CK1[17]. Indeed, inhibiting

SRPK1 was found to reduce viral replication, while blocking GSK-3 activity decreased replication

in cells and reduced infection in patients[17,80]. Second, our results suggest that a population of

unmodified N protein is necessary for viral encapsulation and formation of new virions. Although

the source of this population of unmodified N is less understood, a phosphatase was identified to

act on the murine coronavirus N protein[81]. Similar phosphatases should be identified for the

SARS-CoV-2 N protein and explored as treatment targets. Third, our results also highlight that

phase separation may be critical for the functions of N protein, and particularly that the material

properties of condensates can support N's functions. Therefore, condensate-modifying

molecules, which may not directly interact with N but do partition into condensates and affect their

properties, may also interrupt viral replication and should explored as treatments. Fourth, an

alternative approach to inhibiting viral replication is to target RNA structure. We identified that

RNA structure is important in determining the material properties of N and RNA condensates, and

as such, molecules that disrupt viral RNA structure may be potential treatments for viral infection.

For example, Vögele et al. identified a molecule that disrupts the double-stranded structure of

stem loop 4 (SL4) in viral RNA[82]. Further investigation is needed to assess whether this type of

disruption modulates the material properties of N and RNA condensates in a manner and extent

that impairs N's functions in viral replication.

Although significant progress has been made in understanding the mechanisms of SARS-

CoV-2 replication in infected cells, here we turned to an in vitro system to investigate the

biophysical mechanisms driving the behaviors observed in cells. Where possible, we opted to

include components in our system that mimic physiological conditions, such as in vitro

phosphorylated N instead of a phosphomimetic version, and fragments of viral RNA instead of

unstructured RNA. Using an in vitro system has allowed us to investigate mechanisms that would

be challenging to measure in cells, such as quantifying how specific membrane protein domains

affect condensate-membrane interactions, how specific RNA structures affect strength of N

protein binding, and how phosphorylation affects the conformation of N protein. However, we

recognize that within the crowded environment of the cell, and with many other interaction

partners present during viral infection, the behaviors of N may be different than those we studied

in vitro. For example, phosphorylation of N protein has been shown to promote its interaction with

14-3-3 proteins, which are abundant in cells[83,84]. In our system, we did not consider how binding

to 14-3-3 proteins may change how pN interacts with other viral membrane proteins and

membranes. Therefore, further studies in infected cell models are necessary to verify our findings.

For example, in Scherer et al., the location of pN was identified as the surface of replication

organelles by studying infected cells that were fixed and stained. This experiment should be

repeated with cells that are treated with SRPK1/2 inhibitors to verify how inhibition of

phosphorylation affects the localization of N.

In conclusion, we have investigated the connection between N protein's molecular

conformation, its material properties when phase separated with RNA, and its membrane-

associated behavior. We show that phosphorylation may act as a switch to toggle N between its

two membrane-associated states during viral replication. Given the complexity of the system

studied, further investigations are needed to better understand how our results apply to other N

protein interactions that occur in infected cells. This includes considering factors such as the

complex composition of lipid membranes, as well as the presence of both human mRNA and the

much longer viral genomic RNA.

Acknowledgements

We thank Christine Roden and Amy Gladfelter for helpful discussion, for sharing plasmids

for making RNA fragments and fluorescently labeled viral RNA; Mihai Solotchi and Smita Patel

for assistance with fluorescence polarization experiments and helpful discussions; Gabriela

Tirado-Mansilla for assistance preparing SAXS samples; Bineet Sharma for help with GUV

production; Srinivas Chakravartula and Haiyan Zheng for assistance with mass spectrometry;

Kevin Corbett and Qiaozhen Ye for sharing the GFP-Mfragment plasmid; and Jean Baum and

Vikas Nanda for helpful discussion. This work was supported by NIH grants R35GM142903 (to

B.S.), R35GM147027 (to Z.S), R35GM138296 (to A.G.), R35GM150589 (to G.D.) and National

Science Foundation grant DMREF-2118860 (to S.M. and A.G.). X-ray scattering measurements

were done at the NSLS-II beamline 16-ID (LiX) at Brookhaven National Laboratory. The LiX

beamline is supported by an NIGMS P30 Grant (P30GM133893), NIH Grant S10 OD012331, the

DOE Office of Biological and Environmental Research (KP1605010), and the DOE Office of

Science, Office of Basic Energy Sciences Program under contract number DE-SC0012704 to

CBMS. Figure 1C-D: Created in BioRender. Favetta, B. (2024) BioRender.com/o21g999. Figure

2A-B: BioRender.com/s97t568. Figure 7: Created in BioRender. Favetta, B. (2022)

BioRender.com/382j936.

756   Author Contributions

BF: conceptualization, experimental design, data collection and analysis, writing—original draft,

writing—review and editing; HW: data collection for MPA experiments; JC and AS: experimental

design, data collection and analysis for smFRET experiments; AS: data collection for simulations;

MB: data analysis for microrheology experiments; CR: data collection and analysis for SAXS

experiments; GD: experimental design, data collection and analysis for simulations; SM:

experimental design, data collection, analysis and supervision for SAXS experiments; AJG:

supervision of SAXS data analysis and funding acquisition; ZS: supervision, experimental design

for MPA and GUV experiments and funding acquisition; BSS: conceptualization, supervision,

experimental design, data analysis, writing—original draft, writing—review and editing, funding

acquisition.

768   Declaration of Interests

The authors declare no competing interests.

Resource Availability

Lead contact: Requests for further information and resources should be directed to and will be

fulfilled by the lead contact, Benjamin S Schuster (benjamin.schuster@rutgers.edu).

Materials availability: Plasmids generated in this study will be deposited to Addgene (links will

be made available upon publication).

Data and code availability: All the quantitative analyses discussed in this paper were generated

based on data and computer codes that will be made publicly available (links will be made

available upon publication).

Supporting information

SI Appendix: Table S1, Supplementary Fig. 1–18, SI References

Movies S1-6, related to Figure 2.

Materials and methods

Cloning

All genes of interest were cloned into pET vectors in frame with N-terminal 6x-His tags. A

TEV protease site was inserted between the 6x-His tag and the Nucleocapsid protein coding

sequence. Constructs were cloned by DNA assembly (NEBuilder HiFi DNA Assembly Master Mix;

New England Biolabs). Gene sequences were verified by Sanger sequencing (Genewiz).

Protein expression and purification

For bacterial expression, plasmids were transformed into BL21(DE3) competent E. coli

(New England BioLabs). Colonies picked from fresh plates were grown for 12 hours at 37 °C in 5

mL LB while shaking at 250 rpm. This starter culture was then used to inoculate 0.5 L cultures.

Cultures were grown overnight in 2 L baffled flasks in Terrific Broth medium (Fisher Scientific)

supplemented with 4 g/L glycerol at 18°C while shaking at 250 rpm. Once the OD_{600} reached

approximately 1, expression was induced with 500 µM isopropyl β-D-1-thiogalactopyranoside

(IPTG). The pET vectors used contained a kanamycin resistance gene; kanamycin was used at

concentrations of 50 µg/mL in cultures[85]. After overnight expression at 18 ˚C, bacterial cells were

pelleted by centrifugation at 4100 x g at 10 ˚C. Pellets were resuspended in lysis buffer (1 M NaCl,

20 mM Tris, 20 mM imidazole, EDTA-free protease inhibitor, pH 7.5) and lysed by sonication while

on ice. Lysate was clarified by centrifugation at 25000 x g for 30 minutes at 10 ˚C. The clarified

lysate was then filtered with a 0.22 µm filter.

Proteins were purified using an AKTA Pure FPLC with 1 mL nickel-charged HisTrap

columns (Cytiva) for affinity chromatography of the His-tagged proteins. For N protein, after

injecting proteins onto the column, the column was washed with 15 column volumes of 3 M NaCl,

20 mM Tris, 20 mM imidazole, pH 7.5. For all other proteins, after injecting proteins onto the

column, the column was washed with 5 column volumes of 500 mM NaCl, 20 mM Tris, 20 mM

imidazole, pH 7.5. Proteins were eluted with a linear gradient up to 500 mM NaCl, 20 mM Tris,

500 mM imidazole, pH 7.5. Histidine tags were cleaved from the N protein using TEV protease

during dialysis. TEV protease was added to proteins and dialyzed overnight using 10 kDa MWCO

membranes (Slide-A-Lyzer G2, Thermo Fisher) into 300 mM NaCl, 20 mM Tris, 20 mM imidazole,

5mM DTT, pH 7.5 buffer at 4°C. The reaction mixture was purified using a nickel resin gravity

column (HisPur Ni-NTA Resin, Thermo Fisher) and the flow through was collected. Flow through

aliquots were concentrated and buffer exchanged into storage buffer (300mM NaCl, 20 mM Tris-

HCl, pH 7.5) using a 10kDa MWCO centrifugal filter (Amicon Ultra, Sigma). Proteins were either

reserved for the phosphorylation protocol or snap frozen in liquid N_2 in single-use aliquots and

stored at -80 °C.

SDS-PAGE

For chromatographically purified proteins, SDS-PAGE was run using NuPAGE 4-12% Bis-

Tris gels (Invitrogen) and stained using a Coomassie stain (GelCode Blue Safe Protein, Thermo

Scientific).

Fluorescent labeling of N protein

Purified N protein was dialyzed into PBS buffer with 0.1 mL sodium bicarbonate per 1 mL

protein solution. Protein was then labeled by adding a 3:1 molar ratio of Alexa 647 NHS ester

(Fisher Scientific) stored in DMSO. The mixture was incubated at 4°C for 1 h with rocking.

Unbound dye was removed by size exclusion chromatography (Superdex 200 Increase 10/300

GL, Cytiva). For phase separation assays, percent of dyed protein was adjusted to 5% of total by

dilution with undyed protein.

Phosphorylation protocol

~80 μM N protein was prepared in a buffer containing 300 mM NaCl, 20 mM Tris, 1mM

dithiothreitol (DTT), 10 mM $MgCl_2$, and 2 mM ATP in a 200 μL reaction mixture. 5 μL GSK-3β and

5 μL SRPK (SignalChem Biotech) were added to the mixture. After incubation at 37°C for 120 min,

phosphorylation was confirmed using a SuperSep Phos-tag acrylamide gel (FujiFilm Wako

Chemicals). Enzymes were removed from the reaction mixture using GST-based affinity

chromatography. Briefly, 1 mL of gluthathione resin (Glutathione Sepharose 4B, Cytiva) was

packed in a gravity column. The column was washed with 300mM NaCl, 20mM Tris, pH 7.5 buffer.

The 200 uL of reaction mixture was diluted to a total of 2 mL using wash buffer and poured into

the column. The column flow through was collected. The column was washed with an additional

1 mL of wash buffer, which were collected. The flow through and wash fractions were combined

and the 3 mL of solution containing phosphorylated N protein was concentrated and buffer

exchanged using an Amicon Ultra 0.5 mL 10K MWCO centrifugal filter, according to manufacturer

instructions.

RNA In vitro transcription protocol

RNA production was carried out according to established protocols[86]. Templates were

gifted from the Gladfelter lab where they were synthesized (IDT) and cloned into pJet

(ThermoFisher Scientific K1231) plasmids using blunt end cloning. Directionality and sequence

were confirmed using Sanger sequencing (Azenta). Plasmids were linearized and then amplified

using PCR. 10 ng of plasmid was used as starting material and primers are noted in the

Supplemental Materials. Melting temperatures for the 1-1000 and N fragments were 69˚C. 5 µL

of PCR product was loaded onto an agarose gel to determine size and purity using SYBR™ Gold

Nucleic Acid Gel Stain (Invitrogen) and 1 kb Plus DNA Ladder (New England Biolabs). If the PCR

product was pure then the sample was purified (NEB DNA Clean-up kit). 100 ng of purified DNA

was used as a template for *in vitro* transcription (New England Biolabs) carried out according to

the manufacturer's instructions. Following incubation at 37°C for 18 h, *in vitro* transcription

reactions were treated with DNAse (New England Biolabs) according to the manufacturer's

instructions. Following DNAse treatment, reactions were purified with an RNA purification kit (NEB

T2040L). Purified RNA was verified for purity and size using an agarose gel and RNA Gel Loading

Dye (NEB B0363S) and RiboRuler High Range RNA Ladder (Thermo Scientific SM183).

Concentration was measured using a Nanodrop One spectrophotometer (ThermoFisher

Scientific).

Fluorescently labeled RNA was gifted by the Gladfelter lab. Cy3 RNA was transcribed from

the same template used above and using the same protocol described above, but with the addition

of 0.1 µL of Cy3 labeled UTP to each reaction (Sigma PA53026).

Microscopy

For microscopy experiments, protein samples were prepared as follows: N or pN protein

aliquots were thawed at room temperature. Proteins were then diluted and mixed with RNA (if

any) to obtain a final solution with 40 µM protein and 300 nM viral RNA or 1 mg/mL polyrA (unless

otherwise noted) and buffer conditions of 20 mM Tris-HCl, 150 mM NaCl, pH 7.5. Protein

concentrations were measured based on their absorbance at 280 nm using a Nanodrop

spectrophotometer (ThermoFisher).

Protein samples were plated on 1.5 thickness slides that were coated with 5% Pluronic F-

127 (Sigma-Aldrich) for a minimum of 10 minutes. The slides were washed with buffer solution

prior to plating the protein samples. A silicone spacer (0.5 mm) and a microfluidic temperature

controller (Cherry Biotech) were attached to the slide and the temperature of the sample was set

to 37˚C during observation.

Confocal imaging was performed on a Zeiss Axio Observer 7 inverted microscope

equipped with an LSM900 laser scanning confocal module and employing a 63x/1.4 NA plan-

apochromatic, oil-immersion objective. GFP was excited to fluoresce with a 488 nm laser, Cy3

and Alexa-567 with a 561 nm laser and Cy5 with a 630 nm laser. Confocal fluorescence images

were captured using GaAsP detectors. Transmitted light images were collected with either the

ESID module or an Axiocam 702 sCMOS camera (Zeiss), in both cases using a 0.55 NA

condenser.

Droplet image analysis

Image analysis and data processing for Figures 2, 3 and 4A-C and Supplementary Figures

2 and 8B-C were performed in MATLAB R2023a. The fluorescence intensity profile of the

condensates with fluorescently labeled proteins was measured by using the Hough Transform to

identify droplet locations and drawing a line that spanned the droplet diameter plus 1/4th of a

radius length in each direction across the droplets. Line-scan graphs were generated in MATLAB.

Total intensity and partitioning graphs were generated in MATLAB. Condensate perimeter and

area were calculated using MATLAB's inbuilt regionprops function.

Giant unilamellar vesicle electroformation

Giant unilamellar vesicles were prepared by electroformation[87]. Briefly, 20 uL of a lipid

solution containing 60% DOPC, 25% DOPE, 10% DOPS, 5% DGS-NTA(Ni) in chloroform (Avanti

Lipids) was prepared. ~10 µL lipid solution was spread onto indium tin oxide (ITO)-coated glasses

and dried under vacuum for 12 h. The plates were assembled into a chamber with a Teflon spacer

and the swelling solution (~1 mL of 300mM sucrose) was introduced. For electroformation, a

sinusoidal electric field of 2.0 Vpp and 10 Hz was applied using a function generator for 2.5 h,

after which the frequency was reduced to 5 Hz for 30 min. Osmolarity of the sucrose solution (301

± 1 mOsM) was lower than the osmolarity of the condensate NaCl solutions (324 ± 1 mOsM) to

ensure vesicles were not too tense. The solution osmolarities were confirmed using a freezing-

point osmometer (Advanced Instruments model 3D3).

Contact angles measurement and geometric factor calculation

A detailed explanation of the contact angles used in this work has been published

elsewhere[39]. Briefly, we measured the three contact angles θ_c, θ_e, θ_i from fluorescent images of

GUVs and condensates (for samples with GFP) or from brightfield images (for samples without

GFP). From these angles, we calculated the geometric factor, Φ. From Mangiarotti et al., 2024,

when $\Phi = -1$ there is complete wetting of the membrane by the condensate phase, while $\Phi = +1$

corresponds to dewetting of the membrane by the condensate phase. The geometric factor, Φ, is

negative if the membrane prefers the condensate over the exterior buffer and positive otherwise[56].

We followed the same procedure to measure θ_{in}, the intrinsic contact angle.

Fluorescence Recovery After Photobleaching

Circular bleach regions of approximate radius R = 1 µm were drawn in the center of protein

droplets. Alexa-647 was imaged and bleached with a 640 nm laser. Cy3 was imaged and

bleached with a 561 nm laser. Recovery curves were fit to a single exponential recovery model

$f(t) = A * (1 - e^{-t/\tau})$ to calculate the recovery timescale, т.

Passive Microrheology

Yellow–green carboxylate-modified polystyrene beads (0.5 µm diameter; FluoSpheres,

Invitrogen) were used for video particle-tracking (VPT) microrheology measurements. Each

sample was prepared by mixing protein and RNA to a final concentration of 40 µM protein and

300 nM viral RNA or 1 mg/mL polyrA in 150 mM NaCl, 20 mM Tris-HCl buffer, pH 7.5. Next, 50 µL

of the protein and RNA sample was mixed with 1 uL of a 1:100 dilution of the fluorescent tracer

bead solution and the sample was plated on a 1.5 thickness slide pre-treated with Pluronic F127.

The sample was covered with a CherryTemp microfluidic temperature controller chip and the

sample was set to 37 ˚C and incubated for 1 hour.

VPT measurements were conducted on a Zeiss Axio Observer 7 inverted microscope

equipped with an Axiocam 702 monochrome sCMOS camera (Zeiss), employing a 63x, 1.4-

numerical aperture plan-apochromatic oil-immersion objective. The microscope focus was

adjusted to the midsection of the protein droplets for VPT acquisition. Epifluorescence video

imaging was initiated at the 1-hour timepoint, with fluorescence excitation using a 475-nm light-

emitting diode (Colibri 7, Zeiss). Videos of the tracer beads diffusing within the condensate were

collected at 100 frames per second for 2,000 frames. Imaging was conducted at 37 ˚C. For each

sample, three independent samples were made on different days, and ~20 videos were collected

from each sample, with each video containing ~5–50 tracer beads. Viscosity data presented in

Fig. 4D - G and Supplementary Fig 8D - E are the average of these independent trials.

Data analysis was conducted using the open-source particle tracking package TrackPy

(v0.5.0)[88] in Python and customized as needed. The TrackPy particle tracking code was used to

analyze the collected videos, starting with extracting particle trajectories. The MSD was calculated

from the trajectories of individual beads, followed by calculating the ensemble-average MSD. To

remove the static error from the MSD curves for calculating viscosities, we corrected the

ensemble-average MSD by subtracting the noise floor from the MSD curves[89]. In general, the

ensemble-average MSD often scales as a power law with lag time τ, as given by

$$< r^2(\tau) > = 2dD\tau^{\alpha}$$

where d is the number of dimensions (here d = 2, because data collection and analysis were

conducted in the x–y plane), D is the diffusion coefficient, and α is the diffusivity exponent. For a

purely viscous fluid, the diffusivity exponent α is unity, and the Stokes-Einstein relation can be

used to calculate the viscosity[90]. The α values for all the condensates tested were in the range of

0.3–1.1. We therefore used the Generalized Stokes-Einstein Relation (GSER) to measure the

viscoelastic properties of the condensates[90]. The frequency dependent GSER in the Fourier

domain is represented by the following equation:

$$950 \qquad G^*(\omega) = G'(\omega) + iG''(\omega) = \frac{d\, k_B\, T}{3\pi a(i\omega) <\Delta r^2(\omega)>}$$

where G*(ω) is complex shear modulus, k_B is Boltzmann's constant, T is the temperature, a is the

bead radius, G'(ω) and G" (ω) are the frequency-dependent storage (elastic) and loss (viscous)

moduli, respectively , and $<\Delta r^2(\omega)>$ is the unilateral Fourier transform of the MSD. We use an

algebraic approach proposed by Mason et al. to estimate the Fourier transform of the MSD, which

approximates the local MSD as a power-law function[60]. The algebraic expression is given by the

following equation:

$$957 \qquad |G^*(\omega)| = \frac{d\, k_B\, T}{3\pi a <\Delta r^2(\tau = 1/\omega)> \Gamma[1 + \alpha(\tau = 1/\omega)]}$$

$$958 \qquad G'(\omega) = |G^*(\omega)|\cos\left(\frac{\pi\alpha(\omega)}{2}\right)$$

$$959 \qquad G''(\omega) = |G^*(\omega)|\sin\left(\frac{\pi\alpha(\omega)}{2}\right)$$

where $\alpha(\tau) = d\ln(\Delta r^2(\tau))/d\ln(\tau)$ is the local power law exponent describing the logarithmic

slope of $\Delta r^2(\tau)$ at $\tau = 1/\omega$ and Γ is the Gamma function. Next, cubic spline interpolation fitting is

used on the calculated G' and G" data to reduce measurement noise generated in the algebraic

conversion[91]. We plot the fitted G' and G" to identify the cross-over frequency, or the timescale at

which the material transitions from an elastic-dominant to a viscous dominant regime. To extract

the zero-shear viscosity from the viscoelastic moduli, we use the following equation:

$$966 \qquad \eta(\omega) = \frac{G''(\omega)}{\omega}$$

and obtain the limit of η(ω) at low frequencies[60]. The presented viscoelastic moduli and zero-

shear viscosities are the average from multiple videos (at least n = 10) taken from 3 independent

samples.

The noise floor of the 500 nm beads was measured by allowing a solution of beads in

water to dry on the glass surface of a slide, resulting in beads adhered to the glass surface. We

acquired the trajectories of the beads adhered to the glass surface using the same parameters

and experimental setup as used for VPT studies of the samples.

Micropipette aspiration

The micropipette aspiration experiments were carried out on a Ti2-A inverted fluorescence

microscope (Nikon, Japan) equipped with a motorized stage and two motorized 4-axes

micromanipulators (PatchPro-5000, Scientifica) and a multi-trap optical tweezers (Tweez305,

Aresis, Slovenia) according to the protocol we reported previously[62,63]. An oil immersion

objective (100X; NA 1.30; Nikon) was integrated with an objective heating collar (OKOlab) and

temperature controller (OKOlab) for 37 °C measurements. Micropipettes were made by pulling

glass capillaries using a pipette puller (PUL-1000, World Precision Instruments). The pipette tip

was then cut to achieve an opening diameter ~5 µm. Subsequently, the pipette was bent to an

angle of approximately 40° using a microforge (DMF1000, World Precision Instruments).

MPA experiments were carried out in glass-bottom dishes (ES56291, Azer Scientific, US)

that were pre-treated with 5% Pluronic F-127 (P2443-250G, Sigma) for 10 minutes to prevent

adhesion of condensates to the glass. The micropipette was filled with the same buffer used in

microscopy experiments (150 mM NaCl, 20 mM Tris-HCl, pH 7.5) using a MICROFIL needle

(World Precision Instruments). The filled micropipette was then mounted onto a micromanipulator.

The rear end of the pipette was connected to an automatic pressure controller (Flow-EZ, Fluigent;

Pressure resolution 1 Pa).

Optical tweezers were used to contact and merge droplets to achieve a large (> 5 µm)

condensate for easier MPA measurements and analysis. A secondary micropipette was used to

hold the condensate during MPA. To minimize sample evaporation, 1.5 mL Milli-Q water was

added to the edge of the dish, and the dishes were covered with a thin plastic wrap with a ~2 mm

hole for pipette insertion. We observed that in vitro condensates always wet the inner wall of

uncoated micropipettes. Therefore, the analysis of the MPA data follows the protocol described in

Roggeveen et al[63]. Briefly, normalized aspiration length (aspiration length, Lp, over the pipette

radius, Rp) was segmented according to the pressure steps. For each segment, the slope of a

linear fitting of (Lp/Rp)^2 vs. time is equal to the effective shear rate. Then, the slope of aspiration

pressure vs. shear rate graph gives 4η.

Fluorescence Polarization

Fluorescence polarization measurements were performed on a Tecan Spark microplate

reader in a 384-well black plate at 20°C. A monochromator set the excitation wavelength at 485

nM and the emission wavelength at 535 nm, with a 20 nm bandwidth. Purified N or pN protein

was serially titrated in 150 mM NaCl, 20 mM Tris-HCl, pH 7.5 buffer and incubated with a constant

4 nM FAM-labeled RNA (Azenta) for 10 min at room temperature (20˚C) prior to measurement.

The data was analyzed using Matlab V2023a, with a one-site binding curve (hyperbola)

fitted to the data. K$_D$ values from three experiments were averaged and the standard deviation

calculated. Data was normalized by subtracting the initial polarization value from each dataset.

The one-site binding equation utilized is $mP = \frac{mP_{Max} *[N]}{K_D+[N]}$, in which mP is the observed

millipolarization, mP$_{MAX}$ is the maximum polarization, [N] is the concentration of protein

(unmodified or phosphorylated).

smFRET

Single-molecule fluorescence measurements were performed as described in Cubuk et

al. 2021[8] with a Picoquant MT200 instrument (Picoquant, Germany). Briefly, FRET experiments

were performed by exciting the donor dye with a laser power of ~100 µW (measured at the back

aperture of the objective) at 488 nm wavelength. For pulsed interleaved excitation of donor and

acceptor, we used a repetition rate of 20 MHz for the donor excitation and a delay of approximately

25 ns for acceptor excitation. Acceptor excitation was achieved by using a white laser (SuperK

Extreme, NKT Photonics, Denmark), filtered by a z582/15 band pass filter (Chroma). Acceptor

excitation power was adjusted to match the acceptor emission intensity to that of the donor

(between 50 and 70 µW). Single-molecule FRET efficiency histograms were acquired from

samples with protein concentrations of 100 pM labeled protein and the population with

stoichiometry corresponding to 1:1 donor:acceptor labeling was selected. Photon detection

events were stored with 16 ps resolution.

Proteins were designed and prepared as described in Cubuk et al. 2021[8]. Following

preparation, protein for smFRET experiments was phosphorylated using the protocol described

above, with the exception of the use of 200 mM β-mercaptoethanol instead of 5 mM DTT. All

samples were prepared in 50 mM HEPES pH 7.4, 150mM KCl, 200 mM β-mercaptoethanol (for

photoprotection), 0.001% Tween 20 (for limiting surface adhesion) unless otherwise stated. All

measurements were performed in custom-made glass cuvettes coated with PEG. Each sample

was measured for at least 30 min at room temperature (295 ± 0.5 K). Protein concentrations were

(1) 100 pM labeled protein for the low concentrations or (2) 100 pM labeled protein + 1 µM

unlabeled protein for high concentration for the full-length construct or (3) 100 pM labeled protein

+ 4 µM unlabeled protein for high concentration for N^{1-246}.

Determination of root mean square inter-dye distances from mean FRET transfer

efficiencies was conducted as described in Cubuk et al. 2024[92]. Briefly, the Gaussian chain model

was employed in the conversion, which relies on a single parameter, the root-mean-squared inter-

dye distance $r=<R^2>^{1/2}$. Estimates for this parameter were obtained by numerically solving: $< E >$

$= \int_0^\infty E(R)P(R)dR$ where <E> is the mean transfer efficiency, R is the inter-dye distance, $P(R)$

represents the Gaussian chain distribution, and $E(R)$ is the Förster equation for the dependence

of transfer efficiency on distance R and Förster radius R_0.

smFRET lifetime analysis

Analysis of fluorescence lifetime from mean FRET transfer efficiencies was conducted

as described in Cubuk et al. 2024[71].

smFRET dimerization constant

Modeling of dimerization was conducted as described in Cubuk et al. 2024[71].

Small-Angle X-ray Scattering

Small-angle X-ray scattering (SAXS) measurements were made at 16-ID-C LIXS beamline

(National Synchrotron Light Source II (NSLS-II), Brookhaven National Laboratory; 15.14 keV X-

rays (λ= 0.8189 Å) and two Pilatus 1 M detectors). Samples were prepared by diluting protein into

150mM NaCl, 20mM Tris-HCl buffer to the concentrations desired (0.5 mg/mL – 4 mg/mL). Data

over a q range of 0.005–0.25 Å^{-1} was analyzed. Background subtraction was done using the

scattering from the storage buffer (150mM NaCl, 20mM Tris-HCl, pH 7.5) and by scaling the buffer

and sample intensities at the $q \sim 2$ Å^{-1} water peak.

The data were analyzed in BioXTAS RAW 2.1 with ATSAS 3.0.4 to determine the radius

of gyration (R_g) by Guinier analysis, the compactness of the particle by Kratky plots and pair-

distance distribution functions, $P(r)$. R_g from $P(r)$ was compared with the Guinier R_g to ensure

internal consistency in data analyses. D_{max} was calculated from $P(r)$. Bead model reconstructions

using a dummy atom model were obtained from the $P(r)$ functions generated and stored as .out

files using GNOM in BioXTAS RAW 2.1 using the SAXS data as input. Bead models were

generated using the DAMMIN program in the ATSAS 3.0.4 software package assuming single-

phase objects.

LC-MS

Whole Protein: A sample with protein at a concentration of 10 µM was dissolved in 50%

ammonium acetate solution and diluted 1:10 fold. The sample was then transferred into an LC

vial. Reverse phase liquid chromatography (RPLC) was performed on an Acquity column BEH -

C4 -300A, 1.7 micron (300 µm x 100 mm, Waters Corporation). Column temperature was set at

40C. The following mobile phases were used A: Water with 0.1% formic acid; B: acetonitrile with

0.1% formic acid. The M-Class micro flow LC (Waters Instruments) was operated at a flow rate

of 8 µL/min, and 5 µL of sample was injected with an autosampler after equilibration of the

column. For the separation of proteins, the following concentration gradients were used: starting

with 5 – 90% B for 55 minutes and then back to 5% B for minutes 55 - 62. Prior to injection on

the RPLC column, a blank run was performed. ESI-MS experiments were performed using a

quadrupole-TWIMS-TOF hybrid mass spectrometer (Synapt G2 HDMS; Waters Corporation) in

positive ionization mode with m/z scan range 400 - 4000. The applied experimental parameters

were capillary voltage, 3.0 kV; sampling cone voltage, 40 V; source offset 80C; source

temperature, 100C; desolvation temperature 250C; cone gas 50 L/Hr and desolvation gas 600

L/Hr. Data analysis was performed using MassLynx 4.1 and protein deconvolution was

performed using Mas Ent 1.

Pepsin Digest: For pepsin digestion, protein samples were prepared with 2 µL of protein

and 38 µL PBS and 60 µL of denaturation buffer (2 M urea, 20 mM TCEP, 0.8% formic acid).

Samples were injected into a Vanquish LC system coupled to a Q Exactive HF mass

spectrometer and digested online using a protease type XVIII/pepsin column (NovaBioassays).

Peptides were first trapped on a Waters BEH C18 trap column (2.1 mm × 2 cm), then separated

on a Hypersil Gold C18 analytical column (2.1 mm × 5 cm, 1.9 µm particle size) using a multi-

step gradient at 5 °C: 0–4 min at 2% B, ramping to 10% B in 0.3 min, followed by 10% to 40% B

over 7.6 min. MS data were acquired at 120,000 resolution over a 300–2000 m/z range. The top

20 most intense precursor ions were selected for fragmentation using HCD (collision energy

27), with MS/MS scans acquired at 15,000 resolution and a dynamic exclusion of 20 seconds.

Data were analyzed in Proteome Discoverer 3.0 using Sequest HT against a custom database

containing target proteins and kinases. Precursor and fragment mass tolerances were set to

±10 ppm and ±0.02 Da, respectively. N-terminal acetylation was set as a dynamic protein

terminus modification, while methionine oxidation and phosphorylation at serine, threonine, and

tyrosine were set as dynamic modifications. A non-specific protease setting was used. Peptide-

spectrum matches were validated using the Fixed Value PSM Validator (maximum delta CN

< 0.05), and phosphosite confidence was assessed with PhosphoRS.

Trypsin Digest: Tryptic digests were prepared using the Single-Pot Solid-Phase-

enhanced Sample Preparation (SP3) method with 10 µg of protein per sample. Following protein

binding and washing, digestion was initiated, and after 30 minutes at 37 °C, 20 µL was removed,

acidified with 1 µL of 10% formic acid, and stored at −80 °C. The remaining sample was

incubated overnight at 37 °C to complete digestion. Peptides from each timepoint were prepared

for LC-MS/MS by mixing 10 µL of digest with 10 µL of 0.1% TFA (1:1), resulting in a final

concentration of 0.1 µg/µL. From this, 5 µL (0.5 µg) was injected into a Q Exactive HF mass

spectrometer using a 70-minute data-dependent acquisition (DDA) method. MS/MS data were

searched against a protein database using MSFragger.

BS3 Cross-linking Experiment

1108   Protein (2-10 µM in PBS at pH 7.5) was chemically cross-linked with

1109   bis(sulfosuccinimidyl) suberate (BS3) (Thermo Fisher) (50 µM) at room temperature for 30 min

1110   and quenched by adding Tris (pH 7.5) to a 20 mM final concentration. Reaction mixtures were

1111   then analyzed by SDS-PAGE.

1112   **Molecular dynamics simulations**

1113   All molecular dynamics (MD) simulations were performed using OpenMM 7.6.0[93] in the

NVT ensemble. We simulated single truncated monomers, or full dimers of the N-protein using

Langevin dynamics at a constant temperature of 310 K, with a friction coefficient of 0.01 ps^{-1} and

an integration time step of 10 fs. Each system was enclosed in a cubic periodic box with periodic

boundary conditions. Each chain was simulated for 260 ns, including a 10 ns equilibration phase

followed by 250 ns of production dynamics. From the resulting trajectories, the first 50 ns of

simulation were discarded as additional equilibration time before analysis.

Force field and interaction potentials:

1121   We employed the HPS-Urry coarse-grained energy model[74], which includes bonded

1122   interactions, non-bonded interactions, and an elastic network model (ENM) for structured regions.

1123   The total interaction energy U_{total} is given by:

1124
$$U_{total} = \sum_{i>j} \phi_{ij}^{elec} + \sum_{i>j} \phi_{ij}^{HPS} + \sum_{i=1}^{N-1} k_b (r_{i,i+1} - r_0)^2 + \sum_{(i,j)\in\, pairs} \frac{1}{2} k_d (r_{ij} - r_0^{(ij)})^2$$

1125 where ϕ_{ij}^{elec} represents electrostatic interactions, ϕ_{ij}^{HPS} denotes hydrophobic interactions, and the

1126 third term models bonded interactions between adjacent residues using a spring constant $k_b =$

1127 $20\,kj/\text{Å}^2$ and an equilibrium bond length r_0 = 3.82 Å. The last term corresponds to the elastic

1128 network model (ENM) which restrains the folded domains and is defined below.

1129 Hydrophobic interactions are modeled using the Ashbaugh-Hatch potential[94]:

1130
$$\phi_{ij}(r) = \begin{cases} \phi_{ij}^{LJ}(r) + \left(1 - \lambda_{ij}\right)\epsilon, & for\ r \leq 2^{1/6}\sigma_{ij}, \\ \lambda_{ij}\phi_{ij}^{LJ}(r), & for\ > 2^{1/6}\sigma_{ij} \end{cases}$$

1131
1132 with the Lennard-Jones potential defined as:

1133
$$\phi_{ij}^{LJ}(r) = 4\epsilon\left[\left(\frac{\sigma_{ij}}{r}\right)^{12} - \left(\frac{\sigma_{ij}}{r}\right)^6\right]$$

1134
1135 In this, ϵ = 0.2 kcal/mol. λ_{ij} and σ_{ij} can be calculated by the arithmetic average of

1136 hydrophobicity values (($\lambda_{ij} = \lambda_i + \lambda_j)/2$) and size (($\sigma_{ij} = \sigma_i + \sigma_j)/2$), where σ_i and λ_i are the

1137 size and stickiness respectively of each amino acid type. Values of λ come from the Urry

1138 hydropathy scale[74,95], and σ from the predicted volume of amino acid residues[96].

1139 Electrostatics are modeled using a Debye–Hückel potential[97]:

1140
$$\phi_{ij}^{el}(r) = \frac{q_i q_j}{4\pi Dr}e^{-\kappa r}$$

where q_i represents the charge of residue 'i' located at the center of the corresponding bead. D

is the dielectric constant of the solvent, and κ is the inverse Debye screening length. For our

calculations, we set D = 80, the dielectric constant of water, and κ = 1 nm^{-1}, which corresponds to

salt concentrations of approximately 100 mM.

Elastic Network Model (ENM) for Folded domains:

To maintain structural integrity of folded domains, we applied an elastic network model

(ENM) using a harmonic potential between non-bonded residue pairs within a cutoff distance

based on the reference PDB structure[98]. The ENM potential is defined as:

1149
$$u_{ENM}(r) = \sum_{(i,j) \in \, pairs} \frac{1}{2} k_d \left(r_{ij} - r_0^{(ij)} \right)^2$$

where r_{ij} is the current distance between two beads, r_0^{ij} is the reference distance between residues

i and j from the crystal structure, and k_d is the force constant. For structured domains, each bead

was placed at the center of mass calculated from all atoms in a residue (COM representation),

while for intrinsically disordered domains, each bead was placed at the C_α atom position as has

been used in similar models[75]. This model ensures that secondary and tertiary structures within

folded domains remain stable throughout the simulation. Non-bonded interactions, including

Ashbaugh-Hatch and Debye- Hückel potentials, are excluded for pairs restrained by the ENM.

The Pair Distance Distribution Function (PDDF) P(r) for each model was calculated by

finding the cartesian distance between all residue pairs i,j. Contact between residues was defined

as any 2 residues that were less than 1 nm apart, while ignoring interactions with itself, and 2

subsequent residues in sequence. Finally, a Bayesian Maximum Entropy ensemble reweighting

scheme was implemented to adjust the statistical weights of the simulated conformations

according to Lima Costa and Fushman et al[76].

References

1. Kim, D. *et al.* The Architecture of SARS-CoV-2 Transcriptome. *Cell* **181**, 914-921.e10 (2020).

2. McBride, R., van Zyl, M. & Fielding, B. C. The coronavirus nucleocapsid is a multifunctional protein. *Viruses* vol. 6 2991–3018 (2014).

3. Perdikari, T. M. *et al.* SARS-CoV-2 nucleocapsid protein phase-separates with RNA and with human hnRNPs. *EMBO J.* **39**, 1–15 (2020).

4. Korn, S. M., Dhamotharan, K., Jeffries, C. M. & Schlundt, A. The preference signature of the SARS-CoV-2 Nucleocapsid NTD for its 5'-genomic RNA elements. *Nat. Commun.* **14**, 1–17 (2023).

5. Roden, C. A. *et al.* Double-stranded RNA drives SARS-CoV-2 nucleocapsid protein to undergo phase separation at specific temperatures. *Nucleic Acids Res.* **50**, 8168–8192 (2022).

6. Ye, Q., West, A. M. V., Silletti, S. & Corbett, K. D. Architecture and self-assembly of the SARS-CoV-2 nucleocapsid protein. *Protein Sci.* **29**, 1890–1901 (2020).

7. Morse, M., Sefcikova, J., Rouzina, I., Beuning, P. J. & Williams, M. C. Structural domains of SARS-CoV-2 nucleocapsid protein coordinate to compact long nucleic acid substrates. *Nucleic Acids Res.* **51**, 290–303 (2023).

8. Cubuk, J. *et al.* The SARS-CoV-2 nucleocapsid protein is dynamic, disordered, and phase separates with RNA. *Nat. Commun.* **12**, 1–17 (2021).

9. Zeng, W. *et al.* Biochemical characterization of SARS-CoV-2 nucleocapsid protein. *Biochem. Biophys. Res. Commun.* **527**, 618–623 (2020).

10. Ribeiro-Filho, H. V. *et al.* Structural dynamics of SARS-CoV-2 nucleocapsid protein induced by RNA binding. *PLoS Comput. Biol.* **18**, 1–30 (2022).

11. Tung, H. Y. L. & Limtung, P. Mutations in the phosphorylation sites of SARS-CoV-2 encoded nucleocapsid protein and structure model of sequestration by protein 14-3-3. *Biochem. Biophys. Res. Commun.* **532**, 134–138 (2020).

12. Wu, C. *et al.* Characterization of SARS-CoV-2 nucleocapsid protein reveals multiple functional consequences of the C-terminal domain. *iScience* **24**, 102681 (2021).

13. Bouhaddou, M. *et al.* The Global Phosphorylation Landscape of SARS-CoV-2 Infection. *Cell* **182**, 685-712.e19 (2020).

14. Wu, C. H. *et al.* Glycogen synthase kinase-3 regulates the phosphorylation of severe acute respiratory syndrome coronavirus mucleocapsid protein and viral replication. *J. Biol. Chem.* **284**, 5229–5239 (2009).

15. Fung, T. S. & Liu, D. X. Post-translational modifications of coronavirus proteins: roles and function. **13**, 405–430 (2018).

16. Patton, D., Stohlman, S. A., Fleming, J. & Lai, A. M. C. Synthesis and Subcellular Localization of the Murine Coronavirus Nucleocapsid Protein. *Virology* **130**, 527–532 (1983).

17. Yaron, T. M. *et al.* Host protein kinases required for SARS-CoV-2 nucleocapsid phosphorylation and viral replication. *Sci. Signal.* **15**, 1–16 (2022).

18. Ino, Y. *et al.* Phosphopeptide enrichment using Phos-tag technology reveals functional phosphorylation of the nucleocapsid protein of SARS-CoV-2. *J. Proteomics* **255**, 104501 (2022).

19. Carlson, C. R. *et al.* Phosphoregulation of Phase Separation by the SARS-CoV-2 N

Protein Suggests a Biophysical Basis for its Dual Functions. *Mol. Cell* **80**, 1092-1103.e4
(2020).

20.   Ju, X. *et al.* A novel cell culture system modeling the SARS-CoV-2 life cycle. *PLoS*
Pathog. **17**, 1–23 (2021).

21.   Syed, A. M. *et al.* SARS-CoV-2 evolution balances conflicting roles of N protein
phosphorylation. *SSRN-Unpublished* 7–11 (2023) doi:10.1371/journal.ppat.1012741.

22.   Wu, C. H., Chen, P. J. & Yeh, S. H. Nucleocapsid phosphorylation and RNA helicase
DDX1 recruitment enables coronavirus transition from discontinuous to continuous
transcription. *Cell Host Microbe* **16**, 462–472 (2014).

23.   Peng, T. Y., Lee, K. R. & Tarn, W. Y. Phosphorylation of the arginine/serine dipeptide-rich
motif of the severe acute respiratory syndrome coronavirus nucleocapsid protein
modulates its multimerization, translation inhibitory activity and cellular localization. *FEBS*
J. **275**, 4152–4163 (2008).

24.   Scherer, K. M. *et al.* SARS-CoV-2 nucleocapsid protein adheres to replication organelles
before viral assembly at the Golgi/ERGIC and lysosome-mediated egress. *Sci. Adv.* **8**,
(2022).

25.   Wolff, G. *et al.* A molecular pore spans the double membrane of the coronavirus
replication organelle. *Science (80-.).* **369**, 1395–1398 (2020).

26.   Savastano, A., Ibáñez de Opakua, A., Rankovic, M. & Zweckstetter, M. Nucleocapsid
protein of SARS-CoV-2 phase separates into RNA-rich polymerase-containing
condensates. *Nat. Commun.* **11**, (2020).

27.   Chen, H. *et al.* Liquid–liquid phase separation by SARS-CoV-2 nucleocapsid protein and
RNA. *Cell Res.* **30**, 1143–1145 (2020).

28.   Wang, S. *et al.* Targeting liquid–liquid phase separation of SARS-CoV-2 nucleocapsid
protein promotes innate antiviral immunity by elevating MAVS activity. *Nat. Cell Biol.* **23**,
718–732 (2021).

29.   Zhao, M. *et al.* GCG inhibits SARS-CoV-2 replication by disrupting the liquid phase
condensation of its nucleocapsid protein. *Nat. Commun.* **12**, 1–14 (2021).

30.   Iserman, C. *et al.* Genomic RNA Elements Drive Phase Separation of the SARS-CoV-2
Nucleocapsid. *Mol. Cell* **80**, 1078-1091.e6 (2020).

31.   Roden, C. A. *et al.* Double-stranded RNA drives SARS-CoV-2 nucleocapsid protein to
undergo phase separation at specific temperatures. *Nucleic Acids Res.* **50**, 8168–8192
(2022).

32.   Lasker, K. *et al.* The material properties of a bacterial-derived biomolecular condensate
tune biological function in natural and synthetic systems. *Nat. Commun.* **13**, 5643 (2022).

33.   Yamasaki, A. *et al.* Liquidity Is a Critical Determinant for Selective Autophagy of Protein
Condensates. *Mol. Cell* **77**, 1163-1175.e9 (2020).

34.   Kota, D. & Zhou, H.-X. Macromolecular Regulation of the Material Properties of
Biomolecular Condensates. *J. Phys. Chem. Lett.* 5285–5290 (2022)
doi:10.1021/acs.jpclett.2c00824.

35.   Carlson, C. R. *et al.* Phosphorylation modulates liquid-liquid phase separation of the
SARS-CoV-2 N protein. *bioRxiv* 1–31 (2020) doi:10.1101/2020.06.28.176248.

36.   Carlson, C. R. *et al.* Reconstitution of the SARS-CoV-2 ribonucleosome provides insights
into genomic RNA packaging and regulation by phosphorylation. *J. Biol. Chem.* **298**,
102560 (2022).

37. Yuan, F. *et al.* Membrane bending by protein phase separation. *Proc. Natl. Acad. Sci. U.*
S. A. **118**, (2021).

38. Agudo-Canalejo, J. *et al.* Wetting regulates autophagy of phase-separated compartments
and the cytosol. *Nature* **591**, 142–146 (2021).

39. Mangiarotti, A., Chen, N., Zhao, Z., Lipowsky, R. & Dimova, R. Wetting and complex
remodeling of membranes by biomolecular condensates. *Nat. Commun.* **14**, 1–15 (2023).

40. Scherer, K. M. *et al. SARS-CoV-2 nucleocapsid protein adheres to replication organelles*
before viral assembly at the Golgi/ERGIC and lysosome-mediated egress. *Sci. Adv* vol. 8
https://www.science.org (2022).

41. Klein, S. *et al.* SARS-CoV-2 structure and replication characterized by in situ cryo-
electron tomography. *Nat. Commun.* **11**, (2020).

42. Collins, L. T. *et al.* Elucidation of SARS-Cov-2 Budding Mechanisms through Molecular
Dynamics Simulations of M and E Protein Complexes. *J. Phys. Chem. Lett.* **12**, 12249–
12255 (2021).

43. van Meer, G. Membrane lipids, where they are and how they behave: Sphingolipids on
the move. *FASEB J.* **24**, 112–124 (2010).

44. Pramanik, S., Steinkühler, J., Dimova, R., Spatz, J. & Lipowsky, R. Binding of His-tagged
fluorophores to lipid bilayers of giant vesicles. *Soft Matter* **18**, 6372–6383 (2022).

45. Verheije, M. H. *et al.* The Coronavirus Nucleocapsid Protein Is Dynamically Associated
with the Replication-Transcription Complexes. *J. Virol.* **84**, 11575–11579 (2010).

46. Zimmermann, L. *et al.* SARS-CoV-2 nsp3 and nsp4 are minimal constituents of a pore
spanning replication organelle. *Nat. Commun.* **14**, (2023).

47. Klatte, N., Shields, D. C. & Agoni, C. Modelling the Transitioning of SARS-CoV-2 nsp3
and nsp4 Lumenal Regions towards a More Stable State on Complex Formation. *Int. J.*
Mol. Sci. **24**, (2023).

48. Hurst, K. R., Koetzner, C. A. & Masters, P. S. Characterization of a Critical Interaction
between the Coronavirus Nucleocapsid Protein and Nonstructural Protein 3 of the Viral
Replicase-Transcriptase Complex. *J. Virol.* **87**, 9159–9172 (2013).

49. Hurst, K. R., Ye, R., Goebel, S. J., Jayaraman, P. & Masters, P. S. An Interaction
between the Nucleocapsid Protein and a Component of the Replicase-Transcriptase
Complex Is Crucial for the Infectivity of Coronavirus Genomic RNA. *J. Virol.* **84**, 10276–
10288 (2010).

50. Ni, X., Han, Y., Zhou, R., Zhou, Y. & Lei, J. Structural insights into ribonucleoprotein
dissociation by nucleocapsid protein interacting with non-structural protein 3 in SARS-
CoV-2. *Commun. Biol.* **6**, (2023).

51. Zhang, Z. *et al.* Structure of SARS-CoV-2 membrane protein essential for virus assembly.
Nat. Commun. **13**, (2022).

52. He, R. *et al.* Characterization of protein-protein interactions between the nucleocapsid
protein and membrane protein of the SARS coronavirus. *Virus Res.* **105**, 121–125 (2004).

53. Lu, S. *et al.* The SARS-CoV-2 nucleocapsid phosphoprotein forms mutually exclusive
condensates with RNA and the membrane-associated M protein. *Nat. Commun.* **12**,
(2021).

54. Klein, S. *et al.* SARS-CoV-2 structure and replication characterized by in situ cryo-
electron tomography. *Nat. Commun.* **11**, 1–10 (2020).

55. Bergner, T. *et al.* Near-Native Visualization of SARS-CoV-2 Induced Membrane

Remodeling and Virion Morphogenesis. *Viruses* **14**, 1–17 (2022).

56. Mangiarotti, A., Schmidt, K. V., Lipowsky, R. & Dimova, R. Lipid packing and cholesterol
content regulate membrane wetting by biomolecular condensates. *bioRxiv* 1–31 (2024).

57. Pinto, A. L. *et al.* Ultrastructural insight into SARS-CoV-2 entry and budding in human
airway epithelium. *Nat. Commun.* **13**, 1–14 (2022).

58. Saud, Z. *et al.* The SARS-CoV2 envelope differs from host cells, exposes procoagulant
lipids, and is disrupted in vivo by oral rinses. *J. Lipid Res.* **63**, 100208 (2022).

59. Ma, W., Zheng, G., Xie, W. & Mayr, C. In vivo reconstitution finds multivalent RNA–RNA
interactions as drivers of mesh-like condensates. *Elife* **10**, 1–32 (2021).

60. Mason, T. G. Estimating the viscoelastic moduli of complex fluids using the generalized
Stokes-Einstein equation. *Rheol. Acta* **39**, 371–378 (2000).

61. Wang, Z., Lou, J. & Zhang, H. Essence determines phenomenon: Assaying the material
properties of biological condensates. *Journal of Biological Chemistry* vol. 298 (2022).

62. Wang, H., Kelley, F. M., Milovanovic, D., Schuster, B. S. & Shi, Z. Surface tension and
viscosity of protein condensates quantified by micropipette aspiration. *Biophysical*
Reports vol. 1 (2021).

63. Roggeveen, J. V., Wang, H., Shi, Z. & Stone, H. A. A calibration-free model of
micropipette aspiration for measuring properties of protein condensates. *Biophys. J.* **123**,
1393–1403 (2024).

64. Alshareedah, I., Moosa, M. M., Pham, M., Potoyan, D. A. & Banerjee, P. R.
Programmable viscoelasticity in protein-RNA condensates with disordered sticker-spacer
polypeptides. *Nat. Commun.* **12**, 1–14 (2021).

65. Laghmach, R. *et al.* RNA chain length and stoichiometry govern surface tension and
stability of protein-RNA condensates. *iScience* **25**, (2022).

66. Iserman, C. *et al.* Genomic RNA Elements Drive Phase Separation of the SARS-CoV-2
Nucleocapsid. *Mol. Cell* **80**, 1078-1091.e6 (2020).

67. Lorenz, R. *et al.* ViennaRNA Package 2.0. *Algorithms Mol. Biol.* **6**, 122–128 (2011).

68. Demchenko, A. . Fluorescence Polarization and Rotational Mobility. in *Ultraviolet*
Spectroscopy of Proteins 198–207 (Springer, 1986).

69. Svergun, D. I. & Koch, M. H. J. Small-angle scattering studies of biological
macromolecules in solution. *Reports Prog. Phys.* **66**, 1735–1782 (2003).

70. Kikhney, A. G. & Svergun, D. I. A practical guide to small angle X-ray scattering (SAXS)
of flexible and intrinsically disordered proteins. *FEBS Lett.* **589**, 2570–2577 (2015).

71. Cubuk, J. *et al.* The dimerization domain of SARS CoV 2 Nucleocapsid protein is partially
disordered as a monomer and forms a high affinity dynamic complex. *bioRxiv* (2024).

72. Ghosh, L. S. K. A theoretical method to compute sequence dependent configurational
properties in charged polymers and proteins. *J. Chem. Phys* **143**, (2015).

73. Dignon, G. L., Zheng, W., Kim, Y. C., Best, R. B. & Mittal, J. Sequence determinants of
protein phase behavior from a coarse-grained model. *PLoS Comput. Biol.* **14**, 1–23
(2018).

74. Regy, R. M., Thompson, J., Kim, Y. C. & Mittal, J. Improved coarse-grained model for
studying sequence dependent phase separation of disordered proteins. *Protein Sci.* **30**,
1371–1379 (2021).

75. Cao, F., Bülow, S. von, Tesei, G. & Lindorff-Larsen, K. A coarse-grained model for

disordered and multi-domain proteins. *bioRxiv* 2024--02 (2024) doi:10.1002/pro.5172.

76. Gama Lima Costa, R. & Fushman, D. Reweighting methods for elucidation of
conformation ensembles of proteins. *Curr. Opin. Struct. Biol.* **77**, 102470 (2022).

77. Bernetti, M., Hall, K. B. & Bussi, G. Reweighting of molecular simulations with explicit-
solvent SAXS restraints elucidates ion-dependent RNA ensembles. *Nucleic Acids Res.*
49, (2021).

78. Botova, M. *et al.* A specific phosphorylation-dependent conformational switch of SARS-
CoV-2 nucleoprotein inhibits RNA binding. *Sci. Adv.* (2024) doi:10.1126/sciadv.aax2323.

79. Stuwe, H. *et al.* Phosphorylation in the Ser/Arg-rich region of the nucleocapsid of SARS-
CoV-2 regulates phase separation by inhibiting self-association of a distant helix. *J. Biol.*
Chem. **300**, 107354 (2024).

80. Liu, X. *et al.* Targeting the Coronavirus Nucleocapsid Protein through GSK-3 Inhibition.
medRxiv 2021.02.17.21251933 (2021).

81. Mohandas, D. V. & Dales, S. Endosomal association of a protein phosphatase with high
dephosphorylating activity against a coronavirus nucleocapsid protein. *FEBS Lett.* **282**,
419–424 (1991).

82. Vögele, J. *et al.* High-resolution structure of stem-loop 4 from the 5′-UTR of SARS-CoV-2
solved by solution state NMR. *Nucleic Acids Res.* **51**, 11318–11331 (2023).

83. Tugaeva, K. V. *et al.* The Mechanism of SARS-CoV-2 Nucleocapsid Protein Recognition
by the Human 14-3-3 Proteins :SARS-CoV-2 N association with host 14-3-3 proteins. *J.*
Mol. Biol. **433**, 166875 (2021).

84. Tugaeva, K. V. *et al.* Human 14-3-3 Proteins Site-selectively Bind the Mutational Hotspot
Region of SARS-CoV-2 Nucleoprotein Modulating its Phosphoregulation. *J. Mol. Biol.*
435, 167891 (2023).

85. Studier, F. W. Stable Expression Clones and Auto-Induction for Protein Production in E.
coli. in *Methods in molecular biology (Clifton, N.J.)* vol. 1091 17–32 (2014).

86. Langdon, E. M. *et al.* mRNA structure determines specificity of a polyQ-driven phase
separation. **1**, 922–927 (2018).

87. Angelova, M. I. & Dimitrov, D. S. Liposome Electro formation. 303–311 (1986).

88. Allan, D. B., Thomas, C., Keim, N. C., van der Wel, C. M. & Verweij, R. W. soft-
matter/trackpy: Trackpy. (2021).

89. McGlynn, J. A., Wu, N. & Schultz, K. M. Multiple particle tracking microrheological
characterization: Fundamentals, emerging techniques and applications. *J. Appl. Phys.*
127, (2020).

90. Furst EM, S. T. *Microrheology*. (Oxford University Press, 2017).
doi:10.1093/oso/9780199655205.001.0001.

91. Tassieri, M., Evans, R. M. L., Warren, R. L., Bailey, N. J. & Cooper, J. M. Microrheology
with optical tweezers: Data analysis. *New J. Phys.* **14**, (2012).

92. Cubuk, J. *et al.* The disordered N-terminal tail of SARS-CoV-2 Nucleocapsid protein
forms a dynamic complex with RNA. *Nucleic Acids Res.* **52**, 2609–2624 (2024).

93. Eastman, P. *et al.* OpenMM 7: Rapid development of high performance algorithms for
molecular dynamics. *PLoS Comput. Biol.* **13**, 1–17 (2017).

94. Ashbaugh, H. S. & Hatch, H. W. Natively Unfolded Protein Stability as a Coil-to-Globule
Transition in Charge/Hydropathy Space. *JACS* 9536–9542 (2008).

95.   Urry, D. W. *et al.* Hydrophobicity scale for proteins based on inverse temperature
1387   transitions. *Biopolymers* **32**, 1243–1250 (1992).

1388   96.   Kim, Y. C. & Hummer, G. Coarse-grained Models for Simulations of Multiprotein
1389   Complexes: Application to Ubiquitin Binding. *J. Mol. Biol.* **375**, 1416–1433 (2008).

1390   97.   Debye, P. & Huckel, E. De la theorie des electrolytes. i. abaissement du point de
1391   congelation et phenomenes associes. *Phys. Zeitschrift* **24**, 185–206 (1923).

1392   98.   Bahar, I., Atilgan, A. R. & Erman, B. Direct evaluation of thermal fluctuations in proteins
1393   using a single-parameter harmonic potential. *Fold. Des.* **2**, 173–181 (1997).

1394