S100a4[+] alveolar macrophages accelerate the progression of precancerous atypical adenomatous hyperplasia by promoting the angiogenic function regulated by fatty acid metabolism

Hong Huang[1,2*], Ying Yang[2*], Qiuju Zhang[3], Yongfeng Yang[2,4], Zhenqi Xiong[3], Shengqiang Mao[2], Tingting Song[2], Yilong Wang[2], Zhiqiang Liu[2], Hong Bu[1,5#], Li Zhang[2,4,6,7#], and Le Zhang[3,8#]

[1]Institute of Clinical Pathology, Key Laboratory of Transplantation Engineering and Immunology, Ministry of Health, West China Hospital, Sichuan University, Chengdu, Sichuan, China

[2]Institute of Respiratory Health, Frontiers Science Center for Disease-related Molecular Network, West China Hospital, Sichuan University, Chengdu, Sichuan, China

[3]College of Computer Science, Sichuan University, Chengdu, Sichuan, China

[4]State Key Laboratory of Respiratory Health and Multimorbidity, West China Hospital, Sichuan University, Chengdu, Sichuan, China

[5]Department of Pathology, West China Hospital, Sichuan University, Chengdu, Sichuan, China

[6]Department of Pulmonary and Critical Care Medicine, West China Hospital, Sichuan University, Chengdu, Sichuan, China

[7]Precision Medicine Key Laboratory of Sichuan Province, West China Hospital, Sichuan University, Chengdu, Sichuan, China

[8]Key Laboratory of Systems Health Science of Zhejiang Province, Hangzhou Institute for Advanced Study, University of Chinese Academy of Sciences, Hangzhou, Zhejiang, China.

[*]These authors contributed equally to this work.

[#]**Corresponding author:**

Hong Bu, MD, Department of Pathology, Institute of Clinical Pathology, Key

Laboratory of Transplantation Engineering and Immunology, Ministry of Health, West

China Hospital, Sichuan University, Chengdu, 610041, Sichuan Province, China.

Email: hongbu@scu.edu.cn

Tel.: +86-18980601026

Li Zhang, PhD, Department of Pulmonary and Critical Care Medicine, Institute of

Respiratory Health, State Key Laboratory of Respiratory Health and Multimorbidity,

Frontiers Science Center for Disease-related Molecular Network, Precision Medicine

Key Laboratory of Sichuan Province, West China Hospital, Sichuan University,

Chengdu 610041, Sichuan Province, China.

Email: zhangli2809@wchscu.cn

Tel.: +86-18980602214

Le Zhang, PhD, College of Computer Science, Sichuan University, Chengdu 610065,

Sichuan Province, China; Key Laboratory of Systems Health Science of Zhejiang

Province, Hangzhou Institute for Advanced Study, University of Chinese Academy of

Sciences, Hangzhou 310024, Zhejiang Province, China.

Email: zhangle06@scu.edu.cn

Tel.: +86-13883147240

Abstract

Lung cancer is preceded by premalignant lesions, and what factors drive this transformation and the potential regulatory mode in the context of tumor initiation remain to be elucidated. In the course of precancerous lesions, we found a phasic shift in metabolic patterns. Macrophages are a heterogeneous cell population with high plasticity in the tumor microenvironment. Single-cell interaction and metabolic analyses highlighted a cellular state, S100a4$^+$ alveolar macrophages, which exhibited distinct fatty acid metabolic activity, such as palmitic acid metabolism, at the atypical adenomatous hyperplasia stage, accompanied by an angiogenic-promoting function in a pre-neoplastic setting. These findings were reproducible in human single-cell transcriptomes and had been confirmed by histopathological staining and in vitro cell coculture assays. Taken together, the results from this study demonstrated that the S100a4$^+$ alveolar macrophage subset contributes to tumorigenesis by altering its metabolic state, suggesting that metabolic interventions targeting this cell state in the early stage of disease may delay neoplastic transformation of the lung epithelium.

Keywords: precancerous lesion, atypical adenomatous hyperplasia, alveolar macrophage, fatty acid metabolism, angiogenesis, lung cancer

Introduction

The ontology of lung cancer is a multi-gene-involved, multi-stage, long-term, and extremely complex pathological process (Greaves & Maley, 2012). Emerging evidence indicates that multistep tumorigenesis develops through a series of progressive pathologic changes known as preneoplastic or precursor lesions, which present prior to the onset of cancer (Pich et al., 2022). With the development of imaging technology and the application of molecular biomarkers for lung cancer screening in recent years, the detection rate of early lung cancer, especially precancerous lesions, has increased significantly. Some of the abnormalities in precancerous lesions are reversible; early diagnosis and guiding precancerous patients away from carcinogens or early intervention and blocking may reverse their further development to minimize the risk of cancer (Pennycuick et al., 2020; Teixeira et al., 2019; Yatabe, Borczuk, & Powell, 2011).

As the most common histologic subtype of lung cancer, lung adenocarcinoma (LUAD) predominantly arises from alveolar type 2 (AT2) cells (Zacharias et al., 2018). For the classification of LUAD and its precursors, atypical adenomatous hyperplasia (AAH) is considered to be the first step in a continuum of histomorphologic changes in malignant adenocarcinomas and is in continuity with preinvasive adenocarcinoma in situ (AIS) in terms of morphological alterations, then microinvasive lesions termed minimally invasive adenocarcinoma (MIA), and eventually invasive adenocarcinoma (IA) (S. Sivakumar et al., 2017; Travis et al., 2011). However, the scarcity and difficult accessibility of adequate samples limit the investigation of the cellular and molecular landscape of LUAD precursors. Alternatively, studies also found AAH-like lesions in mouse models, as well as adenomas and AIS, prior to the appearance of LUAD (Nemanja Despot Marjanovic et al., 2020; W. Weichert & A. Warth, 2014). These models accurately mimic human lung precancerous lesions at the molecular and histopathological levels, providing well-suited materials for our study. Gradually shifting the target of lung cancer

treatment from middle and advanced patients with clinical symptoms to

asymptomatic patients with early or precancerous lesions is precisely the new

concept of oncotherapy in the future (Reynolds et al., 2023; Umar, Dunn, &

Greenwald, 2012).

The tumor microenvironment (TME) homeostasis is determined by close crosstalk

within and across all cellular components, including malignant, immune, endothelial,

and stromal cells (Vitale, Manic, Coussens, Kroemer, & Galluzzi, 2019). Unveiling

potential communications between malignant cells and TME underlying oncogenesis

is critical for the future development of mechanism-informed, subset-targeted tumor

immunotherapy strategies (Feng & Gao, 2021; Wen, Lambrecht, Ju, & Tacke, 2021).

Tumor-associated macrophages (TAMs) are a cell population with plasticity and

heterogeneity in the TME. As an important player in tumor pathobiology, on the one

hand, macrophages may be educated by neoplastic cells to provide a favorable

microenvironment, supporting the malignant transformation, disease progression,

metastasis, and resistance to treatment; and on the other, they may play a role in

anti-tumor immunity (Mantovani, Marchesi, Malesci, Laghi, & Allavena, 2017;

Ruffell, Affara, & Coussens, 2012; Sica & Mantovani, 2012). Part of such

heterogeneity may be attributed to the diversity of phenotypic characteristics,

metabolic patterns, and functional profiles of TAMs in response to environmental

perturbations (Cassetta & Pollard, 2018).

Tumorigenesis relies on reprogramming of cellular metabolism, and metabolic

mining of immune cells in the TME has expanded our understanding of

tumor-associated metabolic alterations at different stages of tumor development

(Bian et al., 2020; W. Li et al., 2018). Metabolic changes of TAMs in TME have

been reported to be accompanied by phenotypic and functional changes (Vitale et al.,

2019; Xia et al., 2020). The metabolic landscape of TAMs in the context of tumor

initiation and the interdependent relationship of development, metabolism, and

functional plasticity in TAMs remain largely unraveled. Exploring the major

metabolic circuits by which TAMs remodel the TME and digging metabolic clues

that affect the functional polarization of TAMs will contribute to the proposal of

immunometabolic strategies that utilize TAMs for tumor prevention and therapy.

In order to simulate the process of human lung adenocarcinogenesis, we established

a spontaneous LUAD mouse model and used single-cell RNA sequencing

(scRNA-seq) to parse the metabolic changes of malignant epithelial cells and

macrophages during the precancerous period, explore the metabolic heterogeneity

and pro-tumor mechanism of alveolar macrophages, and probe the key links of

metabolic pattern shifts whereby the alveolar macrophage subset conditions

epithelial cell transformation. In-depth understanding of the molecular and cellular

mechanisms underlying the acquisition of aberrant phenotypes at precancerous and

further carcinogenesis stages may accelerate the identification of precancerous

metabolic intervention targets. Therefore, the diagnosis and prevention of lung

cancer can be advanced to the stage of precancerous lesions, and measures can be

taken to prevent or reverse the further development of precancerous lesions.

Results

Histopathology profiling and scRNA-seq of precancerous lesions in mice

A/J mice have the highest incidence of spontaneous lung tumors among various

mouse strains, and this probability significantly increased with age (Landau, Wang,

Yang, Ding, & Yang, 1998). To more comprehensively mirror the tumor initiation

and progression process of human lung cancer, A/J mice were maintained for 12-16

months for spontaneous lesions, which resulted in three recognizable precancerous

lesions in the lung. A total of 19 tissues were histologically categorized by two

pathologists into four subtypes, including eight normal tissues, three AAHs, three

adenomas, and five AISs, which were confirmed by Ki67 staining, spanning the

cascade from normal to precancerosis (Figure 1B and 1C). To characterize cell

diversity along tumorigenesis, we conducted a time-ordered single-cell

transcriptomic profiling starting with normal tissue and ending with AIS (Figure 1A).

For each sample, we isolated single cells without prior selection for cell types and

used the 10x Chromium platform to generate RNA-seq data. After removing

low-quality cells, a total of 119,698 cells that passed quality control were retained

for subsequent analysis, which yielded a median of 910 detected genes per cell. The

cell count and genes expressed in each sample were provided in Figure S1A and

S1B.

To identify distinct cell populations based on gene expression patterns, we

performed dimensionality reduction and unsupervised cell clustering using methods

implemented in Seurat, followed by removing batch effects among multiple samples.

As shown using Uniform Manifold Approximation and Projection (UMAP), profiles

along the cascade were derived, and a total of 13 main cell clusters were finally

identified, which we defined as the single-cell transcriptome atlas of mouse lung in

premalignant lesions (Figure 1D). Based on the expression of canonical markers, we

classified immune cells into T cells (Cd3e, Cd3d, Cd3g), B cells (Cd79a, Ms4a1,

Igkc), neutrophils (Retnlg, S100a8, S100a9), macrophages (Mrc1, Cd68, C1qa),

monocytes (F13a1, Ms4a6c), dendritic cells (H2-Eb1, H2-Aa, H2-Ab1), and mast

cells (Csf1, Il6) and identified five clusters of non-immune cells, including epithelial

cells (Sftpc, Scgb1a1, Sftpb), fibroblasts (Col1a1, Dcn, Gsn), endothelial cells (Cdh5,

Ramp2, Cldn5), mesothelials (Gpm6a, Upk3b, Msln), and smooth muscle cells

(Acta2, Tagln, Myh11). In addition, a cluster expressing proliferating markers

(Top2a, Mki67) was identified, named cell cycle cells (Figure 1E). Each of these

populations was captured from different pathological stages of different mice (Figure

1F). The proportion of cell types across the four stages showed irregular changes,

while transcriptional homogeneity was reduced with precancerous progression,

illustrating the importance of heterogeneity in tumorigenesis and also proving the

reliability of the sampling in this study (Figure S1C, S1D, and 1G).

Identification of initiation-associated epithelial marker and resolution of

metabolic pattern in malignant epithelial cells

Tumorigenesis relies on reprogramming of cellular metabolism. To obtain

comprehensive cell type-specific metabolic profiles, metabolism-related genes were

extracted from the Kyoto Encyclopedia of Genes and Genomes (KEGG) database,

and metabolic clustering was performed on scRNA-seq data. The metabolic subsets

and cell types corresponded well, indicating metabolic heterogeneity among cell

types (Figure S2A and S2B). Through Gene Set Enrichment Analysis (GSEA)

against metabolic pathways, 16 subsets manifested distinct metabolic activities, with

neutrophils, epithelial cells, and macrophages showing relatively higher metabolic

activities (Figure S2C).

Generally, malignant transformation of epithelial cells is the starting point for the

development of precancerous lesions. AT2 cells, ciliated cells, and AT1 cells were

identified in our epithelial transcriptome (Figure 2A). At the evolutionary stages of

precancerous LUAD, despite possible sample heterogeneity and other interference,

we observed increased interactions between epithelial cells and surrounding stromal

and immune cells in the microenvironment, indicating gradually frequent cell-cell

communication during this process (Figure 2B). It was worth noting that the

Spp1-Cd44 ligand-receptor complex showed more interactions in the AAH stage

than in the AIS stage, between epithelial cells and other cell types, especially

myeloid cells (macrophages, monocytes, and neutrophils). The SPP1-CD44 axis was

reported to be involved in tumor initiation and growth, and CD44 levels were found

to be higher in AAH compared to AIS and IA (Kerr et al., 2004; Pietras et al., 2014).

In addition, the pattern of the App-Cd74 signaling suggested incremental

communications between epithelial cells and immune cells during the occurrence of

precancerous lesions, and it was reported to be implicated in tumor progression and

metastasis (Anderson et al., 2023). Then we sought to identify initiation-associated

epithelial markers underlying LUAD precancerosis. Ldha was gradually enriched in

the precancerous occurrence of lung cancer, and the increased expression trend along

with the tumor initiation process was verified by immunofluorescent staining (Figure

S2D and 2C). Moreover, analysis of The Cancer Genome Atlas (TCGA) database

indicated that the initiation-associated marker LDHA was highly expressed in LUAD

and showed an increasing trend in the early stages of the disease, and its high

expression was correlated with poor prognosis of LUAD patients (Figure 2D-F).

Thus, it deserves to be further evaluated as a potentially promising biomarker during

lung tumorigenesis.

To distinguish malignant cells from non-malignant cells based on karyotypes,

chromosomal copy number variations (CNVs) from each epithelial cell's profile

were inferred (Figure 2G and S2E). There were some CNV variations in the normal

group, which may be related to the nature of the algorithm, the sample collection

operation, or the late processing of the cells; these cells were excluded from

subsequent analyses. To explore the phasic metabolic changes in malignant epithelial

cells, metabolic activity was quantified using the scMetabolism package. The results

indicated that malignant epithelial cells in the adenoma stage were enriched in lipid

metabolism-related pathways, while in the AIS stage, significant enrichment of

glycometabolism-related pathways was observed (Figure 2H). The Gene Set

Variation Analysis (GSVA) enrichment analysis showed a concordant variation trend

(Figure 2I). These data suggested that epithelial cells manifest diverse metabolic

patterns during malignant transition.

S100a4^{+} alv-macro exhibited active fatty acid metabolism in the AAH phase

With the development of tumor metabolism research, it has been found that the

metabolism of other cell types in the microenvironment besides epithelial cells can

also regulate tumorigenesis. Macrophage was the most prominent cell type that

interacted with malignant epithelial cells, as shown by Cell Chat analysis (Figure

S3A). Macrophages were composed of alveolar macrophages (Itgax and Siglecf) and

interstitial macrophages (Itgam and Cx3cr1) in subclustering (Figure S3B and S3C).

Further analysis revealed a higher correlation between alveolar macrophages and

malignant epithelial cells (Figure 3A and S3D). To resolve the metabolic pattern of

alveolar macrophages at a higher resolution, we subclassified them into four

subpopulations based on gene expression (Figure S3E and S3F). Enrichment

analysis was performed for each of the four subpopulations, and scMetabolism

analysis revealed that one of the alveolar macrophage subpopulations already

showed a metabolically active state at the AAH stage, especially lipid metabolism,

which we hereafter termed as S100a4^{+} alv-macro (Figure 3B and C). As a

representative, the overall abundance trends of glycerolipid metabolism,

glycerophospholipid metabolism, and biosynthesis of unsaturated fatty acids were

displayed in Figure 3D. Next, we used the Compass algorithm to model the

metabolic flux of cells for S100a4^{+} alv-macro, quantitatively analyzed the metabolic

activity of cells, and then focused on metabolic enzymes and genes related to lipid

metabolism for in-depth analysis. Fatty acid metabolism was found to be relatively

more active in the AAH phase, which was represented by changes in palmitoyl-CoA

desaturase, palmitoyl-CoA hydrolase, and carnitine palmitoyl-transferase (Figure

3E). Consistently, the expression levels of related genes Cpt1a and Acot2 were also

higher in precancerous lesions relative to normal circumstance (Figure 3F). In our preliminary validation, the presence of F4/80[+]/CD11c[+]/S100A4[+] alv-macro was identified in the corresponding tissues by multiplexed immunohistochemistry staining, and their number and proportion were observed to be significantly greater in AAH samples than in normal samples (Figure S3G, S3H, and 3G). CPT1A is responsible for the fatty acid β-oxidation in mitochondria (Nakamura, Yudell, & Loor, 2014), and its positivity was found more frequently in the S100a4[+] alv-macro of the AAH group (Figure 3G), suggesting fatty acid metabolism in this subpopulation was activated at the AAH stage.

Pro-angiogenic function of S100a4[+] alv-macro was associated with fatty acid metabolism

Metabolic profile is coupled with phenotype and functional program of macrophages. Among all pathological stages, although there was no significant difference in the cell proportion of S100a4[+] alv-macro in macrophages, its proportion in alveolar macrophages was obviously the highest in the AAH stage (Figure S4A and 4A). To evaluate changes in the functional programs of this macrophage subpopulation in the context of precancerous lesions, we performed enrichment analysis of the well-known macrophage phenotypes. It was revealed that the capacities for angiogenesis, M2-like polarization, and immunosuppression were found to be stronger in AAH or other precancerous stages relative to the normal stage (Figure 4B). The pro-angiogenic function predominated in the AAH stage, while M2-like and immunosuppressive functions were more prominent in the subsequent precancerous progression. Correlation analysis was used to measure the relationship between these functional programs and metabolic activity, where pro-angiogenic and M2-like phenotypes of S100a4[+] alv-macro were proved to be positively correlated with lipid metabolism (Figure S4B). Further investigation showed that both of them were positively correlated with palmitic acid-related metabolic reactions (DESAT16_2_pos, C160CPT1_pos, and RE0577C_pos, Figure 4C). Next, we

queried the angiogenesis-related gene signature in S100a4$^+$ alv-macro, and the
expression levels of Anxa2 and Ramp1 were detected to be higher in the AAH group
than the normal group (Figure 4D). ANXA2 supports angiogenesis in specific
tumor-related settings (Huang et al., 2022), and its expression level was positively
correlated with that of Cpt1a in S100a4$^+$ alv-macro (Figure 4E). What's more, the
correlation between CPT1A and ANXA2 was verified in the immunostaining of
AAH tissues, revealing more F4/80$^+$/CD11c$^+$/S100A4$^+$/CPT1A$^+$/ANXA2$^+$ alv-macro
compared to normal tissues (Figure 4F). These results suggested that the
pro-angiogenic function of S100a4$^+$ alv-macro might be related to its fatty acid
metabolic status.

S100a4$^+$ alv-macro displayed the potential to promote the early malignant transformation of lung epithelial cells in vitro

To further validate our findings from scRNA-seq and histological staining,
S100a4-overexpressed (OE) alveolar macrophages were established by transfection
of the mS100a4 vector into the murine MH-S cell line, and empty vector was
transfected as negative control (NC) cells, which were verified by quantitative
real-time polymerase chain reaction (qPCR) and western blotting (Figure 5A and
5B). Then coculture assays were conducted to functionally prove the protumorigenic
characteristics of S100a4$^+$ alv-macro in vitro. After culture with conditioned medium
(CM) of S100a4-OE MH-S, the proliferation and migration ability of mouse lung
epithelial MLE12 cells were significantly promoted, and the cell cycle was more
arrested in the G2/M phase (Figure 5C-F). Reactive oxygen species (ROS) produced
within cells can lead to genetic mutations that trigger malignant transformation of
cells (Santibáñez-Andrade et al., 2023), and we found markedly elevated ROS levels
in OE-CM cocultured epithelial cells (Figure 5G). Mitochondrial morphological
change is one of the organelle indications of malignant transformation of cells
(Missiroli, Perrone, Genovese, Pinton, & Giorgi, 2020). As shown in the electron
microscope images, compared with the cocultured epithelial cells in the NC group,

we observed the acquisition of irregular shapes, obvious vacuoles, and disorganized cristae in mitochondria of the OE group, which might be associated with energy metabolism in the process of cell transformation (Figure 5H). In addition, we performed reverse verification with siRNA and demonstrated that the functional phenotype of epithelial cells did not change after S100a4-knockdown MH-S coculture (Figure S5).

During the in vitro carcinogenesis process, a battery of assays allows for the identification of the cells in the initiation, promotion, or aggressive stages of tumorigenesis (Barguilla et al., 2023). We found increased expression of DNA damage marker p-γH2AX and decreased expression of adhesion protein E-Cadherin, but no marked changes in N-Cadherin, Vimentin, and stem-like markers (CD44 and CD133) (Figure 5I). The expression of tumorigenesis-associated proteins (c-MYC, p-ERK, SPA, VEGF, and HIF-1α) was elevated in OE-CM-cultured MLE12 (Figure 5J). Combined with the above observations and the absence of difference in invasion potential, we speculated that the coculture partially activated the epithelial-mesenchymal transition (EMT) program, and the transformation of lung epithelial cells was still in the early or intermediate stage and had not yet reached the advanced stage, which validated our findings to a certain extent and might correspond to the AAH stage in histology. In addition, the expression of pro-tumor indicators in macrophages (VEGF, MMP9, TGF-β, and HIF-1α) was also higher in S100a4-OE MH-S (Figure 5K). Next, we detected tumor-inducing factors secreted by these unique macrophages using Cytokine Antibody Array. We noted the production of macrophage inflammatory protein (MIP)-2, hepatocyte growth factor (HGF), tumor necrosis factor α (TNF-α), IL-1α, MMP9, and CD40, and these cytokine-related biological processes were mainly involved in the regulation of angiogenesis and immune response (Figure 5L and 5M). In summary, we demonstrated in vitro that S100a4[+] alv-macro promote the early malignant transformation of lung epithelial cells by secreting tumor-promoting cytokines.

S100a4[+] alv-macro drove angiogenesis by promoting Cpt1a-mediated fatty acid metabolism

Next, we sought to identify the metabolic and functional changes found in the AAH phase mentioned above. It was demonstrated that the pro-tumor polarization of macrophages is dependent on the accumulation of intracellular lipid droplets as a stable source of fatty acid metabolism (H. Wu et al., 2019). As expected, in S100a4-OE MH-S, we observed an increase in lipid droplet content, as well as enhanced function of mitochondria where fatty acid β-oxidation takes place (Figure 6A). The pro-angiogenic effect of this subset was confirmed by tube formation assay of human umbilical vein endothelial cells (HUVECs) (Figure 6B). Moreover, fatty acid metabolism-related CPT1A and angiogenesis-related ANXA2 were also up-regulated in S100a4-OE MH-S (Figure 6C and 6D). ANXA2 is considered to be a cellular receptor for plasminogen (PLG), often expressed on the surface of endothelial cells, macrophages, and tumor cells, which activates a cascade of pro-angiogenic factors by promoting the conversion of PLG to plasmin, thereby promoting angiogenesis and tumor progression (Semov et al., 2005; Sharma, 2019). We found synergistic upregulation of ANXA2 and PLG expression in S100a4-OE MH-S and cocultured HUVEC and MLE12, which may help explain how ANXA2 induction was involved in angiogenesis and malignant transformation (Figure 6D).

Then we aimed to figure out the contribution of CPT1A to angiogenesis and pro-tumor function in S100a4[+] alv-macro. Following the addition of the CPT1A inhibitor etomoxir (ETO) to S100A4-OE MH-S, the cocultured MLE12 showed restoration of cell proliferation, migration, and ROS production, and the cocultured HUVEC also exhibited decreased tube-forming capacity (Figure 6E-H). At the protein expression level, we also identified the recovery of angiogenesis-related proteins in MH-S and cocultured MLE12, as well as a regression in the levels of pro-tumorigenic and M2 markers in MH-S (Figure 6I and 6J). It was also found that the effect of CPT1A may be dependent on the activation of PPAR-γ signaling.

Therefore, S100a4$^+$ alv-macro may facilitate angiogenesis through

CPT1A-PPAR-γ-mediated fatty acid metabolism, thereby triggering tumorigenesis.

S100A4$^+$ alv-macro in similar patterns was present in human precancerous

AAH lesions

Finally, we were dedicated to corroborating the correspondence between the results

analyzed in human samples and our observations in the mouse model. A total of 23

human tissues were included in our study to proceed with the aforementioned

analysis, including nine normal lung tissues, three AAHs, four AISs, three MIAs,

and four IAs, all of which have been pathologically confirmed by hematoxylin and

eosin (H&E) staining and pathologist assessment (Figure 7A). The annotation of

scRNA-seq profiles yielded 15 cell types, in which macrophages and AT2 cells

similarly exhibited significant enrichment of metabolic activities (Figure S6A-C).

Malignant cells were extracted from epithelial cells by inferring chromosomal copy

number alterations, and GSVA enrichment analysis of malignant epithelial cells was

performed across the four pathological stages (Figure S6D and S6E). Along with

scMetabolism analysis, the results collectively suggested that human malignant

epithelial cells showed relatively active lipid metabolism during the AAH phase, as

for the IA phase, glycometabolism and oxidative phosphorylation pathways were

obviously enriched (Figure S6F). Besides, the enrichment of macrophage migration

and chemotaxis indicated the vital role of macrophages throughout the whole

malignant transformation process of LUAD.

In order to figure out whether phenotypically and functionally similar subset of

alveolar macrophages also exist in human LUAD precancerous lesions, human

alveolar macrophages were further subdivided into four cell subsets, which led us to

find that the same was true for human S100A4$^+$ alv-macro (Figure 7B and 7C). In

addition, the activities of palmitoyl-CoA hydrolase and carnitine

palmitoyl-transferase were also verified to be elevated during the AAH phase,

corresponding to metabolic reactions RE0577C_pos and C160CPT1_pos in the

391 Compass analysis (Figure 7D). For the macrophage phenotype, several functional

392 programs with higher abundance in the AAH phase in the mouse scRNA-seq

393 followed a consistent trend in human data, including angiogenesis (Figure 7E). In

394 terms of gene expression, the expression of fatty acid metabolism-related genes

395 CPT1A and ACOT2 and angiogenesis-related gene ANXA2 in the AAH stage was

396 also in line with expectations (Figure 7F). Furthermore, we also noticed greater

397 amounts of S100A4$^+$ alv-macro in human AAH tissues and verified the correlation

398 between CPT1A and ANXA2 in this subset (Figure 7G). Taken together, these

399 results confirmed the presence of S100A4$^+$ alv-macro with fatty acid metabolic

400 properties and pro-angiogenic function in the development of human precancerous

401 AAH.

Discussion

Here, our scRNA-seq data provided a comprehensive resource to investigate cell state changes alongside tumor initiation in a mouse model of spontaneous tumors simulating the oncogenic transformation in clinical LUAD. We extensively mapped the transcriptional landscape of cell subpopulations and switching cell interactions, some of which were essential in reshaping the microenvironment favoring tumor evolution. In the course of precancerous lesions, the transcriptional heterogeneity was gradually enriched, indicating underlying progressive molecular and cellular changes during carcinogenesis. We identified an initiation-associated marker, LDHA; it was suggested that this glycolysis-related gene was significantly upregulated in LUAD and could serve as an independent prognostic indicator of unfavorable overall survival and recurrence-free survival (Yu et al., 2018), illustrating the potentially vital role of metabolic alternations in tumorigenesis.

Based on the understanding that TME can affect the state of epithelial cells, metabolic fluctuations in this context are closely linked to cell phenotype and function (Buck, Sowell, Kaech, & Pearce, 2017; de Visser & Joyce, 2023; J. Wang et al., 2021). Macrophages are no exception to this rule; their metabolic profile can be shaped by specific histological TME, and the cellular state is thus activated and exhibits remarkable plasticity (Kumar et al., 2019; Mazzone, Menga, & Castegna, 2018). Besides, it has been demonstrated that macrophages contribute to the induction of early lung cancer lesions (Casanova-Acebes et al., 2021), and purine metabolism has been confirmed to determine the pro-tumor phenotype of the macrophage population (S. Li et al., 2022). In our exploration of malignant epithelial cells in the initiation of LUAD, alveolar macrophages were identified as the cell population with which they interacted most strongly. The scRNA-seq approaches enable the resolution of macrophage heterogeneity, assisting us in understanding how these transcriptionally defined subsets connect to macrophage metabolic status and functional phenotypes. Individual macrophage subsets possess distinct metabolic

profiles; our analysis highlighted a specific cell subset, which persisted throughout

lung carcinogenesis and was observably defined by the expression of S100a4 that

had previously not been implicated in tumor-associated alveolar macrophages. The

most intriguing finding about this cellular state was that the fatty acid metabolism

became active during the AAH phase, accompanied by angiogenetic-like properties.

S100A4, which was regarded as an angiogenesis regulator, has been stated to

enhance macrophage protumor phenotype by control of fatty acid oxidation

(Kazakova et al., 2023; S. Liu et al., 2021). What is more convincing is that

S100A4$^+$ alv-macro with similar characteristics was extended to scRNA-seq profiles

of human precancerous AAH lesions. CPT1A was thought to play a crucial role in

this metabolic paradigm shift, and CPT1A-mediated fatty acid oxidation has been

proved to be conducive to tumor metastasis, radiation resistance, and immune escape

(Z. Liu et al., 2023; Tan et al., 2018; Y.-N. Wang et al., 2018). We demonstrated the

regulation of fatty acid metabolism by CPT1A in S100a4$^+$ alv-macro as well as the

involvement of PPAR-γ. Nevertheless, the molecular mechanism that drives the

acquisition of metabolic and functional switching properties specific to this cell state

still requires further characterization in the context of precancerous lesions. It has

been reported that CD36 is the main effector of the S100A4/PPAR-γ pathway, and its

mediated fatty acid uptake plays an important role in the tumor-promoting function

of macrophages (S. Liu et al., 2021).

In terms of subset functional phenotype, S100a4$^+$ alv-macro enriched for fatty acid

metabolism held restricted M1-like and phagocytosis capacities at the AAH stage but

exhibited M2-like, angiogenesis, and immunosuppressive features, creating a

permissive environment for neoplasia. The pro-angiogenic function was dominant

and confirmed by tube formation assay of HUVEC, and the regulation of CPT1A on

angiogenesis was also revealed. The role of ANXA2 in angiogenesis has been

widely recognized, and it may facilitate plasmin production by binding to S100A4

and then trigger angiogenesis and malignant cell transformation (Grindheim, Saraste,

& Vedeler, 2017; Y. Liu, Myrvang, & Dekker, 2015). Precancerous lesions of LUAD

are angiogenic, and pro-angiogenic factors secreted by cells, including S100a4[+]

alv-macro, induce endothelial cell sprouting and chemotaxis, leaving the angiogenic

switch activated, prompting the formation of new blood vessels on the basis of the

original ones to supply oxygen and nutrients to sustain tumor initiation (Chen et al.,

2024; Kayser et al., 2003; van Hinsbergh & Koolwijk, 2008). Under hypoxic

conditions, HIF-1α activates numerous factors that contribute to the angiogenic

process, including VEGF, which promotes vascular permeability, and MMP9, which

breaks down the ECM, promotes endothelial cell migration, and recruits pericytes to

provide structural support (Raza, Franklin, & Dudek, 2010; Sakurai & Kudo, 2011).

Cytokines secreted into the microenvironment activate macrophages, which

subsequently produce angiogenic factors, further promoting angiogenesis (Sica,

Schioppa, Mantovani, & Allavena, 2006). Furthermore, TGF-β and HGF activate

vascular endothelial cells and promote proliferation and migration, as well as induce

the expression of pro-angiogenic factors such as VEGF (Vimalraj, 2022; Watabe,

Takahashi, Pietras, & Yoshimatsu, 2023). Macrophage-derived TNF-α and IL-1α

lead tumor cells to produce potent angiogenic factors IL-8 and VEGF, which affect

angiogenesis and tumor growth (Torisu et al., 2000). MIP2 and CD40 were also

identified as pro-tumor factors associated with angiogenesis (Kollmar, Scheuer,

Menger, & Schilling, 2006; Murugaiyan, Martin, & Saha, 2007). It is worth noting

that our mining of S100a4[+] alv-macro remains at the precancerous initiation stage,

and further experimental designs are needed to verify its specific contribution at

more aggressive stages. For example, FACS sorting of the subpopulation at different

stages of disease progression, respectively, for precise functional characterization;

continuous monitoring of the fluctuation of the above factors in bronchoalveolar

lavage fluid at corresponding periods; and intratracheal instillation of primary

S100a4[+] alv-macro to observe the pathological progression of precancerous lesions.

Besides, as our previous in vitro results were obtained based on cell lines, we will

optimize the experimental conditions to achieve coculture of primary macrophage

subset and epithelial cells and establish transgenic mouse models for in vivo

validation.

With the elaborate resolution of TME, macrophage-related therapy is considered to

be promising. So far, macrophage-targeted therapy has demonstrated clinical

efficacy in Gaucher's disease and advanced hematological malignancies (Barton et

al., 1991; Ossenkoppele et al., 2013). In lung cancer, an attempt to enhance

anti-PD-1 therapy in NSCLC by depleting myeloid-derived suppressor cells with

gemcitabine was prematurely terminated because of insufficient data collected;

another clinical trial of TQB2928 monoclonal antibody promoting macrophage

phagocytosis of tumor cells in combination with a third-generation EGFR TKI for

advanced NSCLC is now recruiting. Moreover, preclinical studies on

macrophage-targeted therapy combined with immune checkpoint inhibitors are being

extensively conducted in NSCLC, and it was suggested that blockade of purine

metabolism can reverse macrophage immunosuppression, and a synergetic effect can

be achieved when combined with anti-PD-L1 therapy, which inspired the direction

of our early intervention strategies (H. Wang, Arulraj, Anbari, & Popel, 2024; Yang

et al., 2025).

In this study, our results provided the unique patterns of cellular components during

the initiation process of LUAD and gave evidence that S100a4$^+$ alv-macro enhances

the function of pro-angiogenesis by promoting fatty acid metabolism, thereby

accelerating the process of precancerous lesions. Early intervention in solid cancers

at the precancer stage may effectively prevent their progression (Prieto et al., 2023).

In the cascade of tumor initiation, AAH is represented as a probable forerunner of

LUAD, serving as a potential hub in carcinogenesis (Mori, Rao, Popper, Cagle, &

Fraire, 2001). On the basis of the newly identified pro-tumor subpopulation and

relevant metabolic and functional elements, the development of novel

immunotherapies targeting the immunometabolism will provide alternative choices

for interventions at the AAH stage, paving the way for potential benefits for patients.

Materials and methods

A/J mouse model and human samples

A/J mice were obtained from Shanghai Model Organisms Center (Shanghai, China).

The 200 mice were divided into groups and housed in an air-conditioned room with

a 12-h light/dark cycle and standard chow diet. Mice were killed by decapitation at

different months of age. A gross examination of all organs was conducted, the lungs

were removed and bisected for tissue fixation and the preparation of single cell

suspensions. All the animal surgical and experimental procedures were approved by

the Ethics Board of Experimental Animals, West China Hospital, Sichuan University

(Approval No. 20230208003).

Human clinical specimens from ten patients were enrolled with informed consent

from West China Hospital. All samples were obtained during surgery and prior to

any other treatment. Histological classification of precancerous lesions and tumors

was determined according to the WHO guidelines (Travis et al., 2015). Fresh

specimens were instantly divided into two pieces for single cell suspension

preparation and histopathological staining. The utilization of human samples was

approved by the Medical Ethics Committee and Institutional Review Board of West

China Hospital, with all participants providing written informed consents.

H&E staining

Lung tissue was further analyzed by histopathology to confirm pathological stage.

H&E staining was performed on 4 μm paraffin sections according to routine staining

protocol. Microscopic H&E-stained slides were independently reviewed by clinical

or veterinary pathologists from the Department of Pathology, West China Hospital.

Histopathological classification of mouse (AAH, adenoma, and AIS) and human

(AAH, AIS, MIA, and IA) lesions was performed as previously described (Nemanja

Despot Marjanovic et al., 2020; Smruthy Sivakumar et al., 2017; Travis et al., 2011;

Wilko Weichert & Arne Warth, 2014).

Single cell suspension preparation

Resected samples were washed in Hanks' balanced salt solution (HBSS, GIBCO,

Grand Island, USA) on ice immediately, minced into small pieces with a sterile

surgical scalpel, and then transferred into a conical tube. The mechanical

dissociation was followed by an enzymatic digestion. The tissue fragments were

digested in HBSS with 2 mg/mL collagenase I and 1 mg/mL collagenase IV

(Washington, Lakewood, USA) for 30-45 min at 37 °C. The suspension was then

filtered using 70 μm and 30 μm cell strainers (Corning, NY, USA) successively, and

the cell pellet was resuspended with red blood cell (RBC) lysis buffer (Invitrogen,

Carlsbad, USA) to discard RBCs. Following cell sorting, dead cells were removed

from the cell suspension based on exclusion of 7-aminoactinomycin D (7-AAD, Life

Technologies Corporation, Gaithersburg, USA), and live cells were sorted into

phosphate buffer solution (PBS) with 0.04% bovine serum albumin (BSA) for

subsequent scRNA-seq.

Single-cell RNA library construction and sequencing

Single cell capturing and cDNA library generation were performed using the Single

Cell 3' Chip Kit and 10x Chromium 3' Library & Gel Bead Kit v3 (10x Genomics,

Pleasanton, USA) following the manufacturer's instructions. The barcoded cDNA

was purified with Dynabeads MyOne SILANE beads (Thermo Fischer Scientific,

Waltham, USA) before amplification by PCR. Amplified cDNA product was

size-selected using SPRIselect Reagent (Thermo Fisher Scientific). After adaptor

ligation and sample index PCR, libraries were sequenced on an Illumina NovaSeq

6000.

Single-cell RNA sequencing analysis and identification of marker genes

Raw gene expression matrices were generated per sample using CellRanger (version

4.0.0). These matrices were then combined by R (version 4.0.5) and converted into a

Seurat data object using the Seurat R package (version 4.3.0.1). We excluded cells

based on any one of the following criteria: (1) more than 10,000 or fewer than 500

unique molecular identifiers (UMIs); (2) more than 5,000 or fewer than 300

expressed genes; (3) over 10% UMIs that were derived from the mitochondrial

genome; (4) genes that were expressed in fewer than 10 cells. Subsequently, we

carried out downstream analyses by the Seurat workflow (Hao et al., 2021) as

described below:

Firstly, we normalized the counts on the filtered matrix for each gene to the total

library size by the 'NormalizeData' function. Secondly, we employed the

'FindVariableGenes' function to identify the most highly variable genes for

unsupervised clustering. Thirdly, these highly variable genes were centered to a

mean of zero and were scaled by the standard deviation by the 'ScaleData' function

of Seurat. Fourthly, principal component analysis (PCA) was performed by the

function 'RunPCA'. The 'RunHarmony' function was used to integrate samples and

remove batch effects. The 'FindClusters' function was used to calculate the clusters

of the cells. UMAP plots were used to visualize clusters of cells localized in the

graph-based clusters by the 'RunUMAP' function. Markers for each cluster were

identified by finding differentially expressed genes between cells from the individual

cluster versus cells from all other clusters by the 'FindAllMarkers' function.

Estimating heterogeneity within stages

Heterogeneity of single-cell profiles within a stage was quantified by examining the

average pairwise Normalized Mutual Information (NMI) (N. D. Marjanovic et al.,

2020) between the profiles for each stage. Firstly, we selected 100 differentially

expressed genes for each of the 13 cell type clusters and the top 100 genes for each

of the top 20 components in harmony reduction. Next, we discretized expression per

gene into 10 bins. Considering the differences in the number of cells across samples,

we subsampled 100 cells from each stage 500 times and calculated the median NMI

across each within-stage sampled pair. NMI was calculated between each pair of

cells x and y by first calculating the mutual information (Eq. 1) and then normalizing

it by the entropy of each cell (Eq. 2).

$$I(x,y) = \sum_x \sum_y p(x,y) log \frac{p(x,y)}{p(x)p(y)} \qquad (1)$$

601
$$NMI(x, y) = \frac{I(x,y)}{\sqrt{H(x)H(y)}}. \qquad\qquad (2)$$

where $p(x)$ and $p(y)$ represent the probability distributions of cell x and cell y

over all discretized gene expression values, respectively. $p(x, y)$ denotes the joint

probability distribution of cell x and cell y over all discretized gene expression

values. $H(x)$ and $H(y)$ correspondingly indicate the entropy of cell x and cell y

across all discretized gene expression values.

Single-cell metabolic state analysis

According to the Molecular Signatures Database (Subramanian et al., 2005), we

extracted 2,710 genes associated with the metabolic pathways for dimensional

reduction and clustering analysis. We employed PCA to process the metabolic genes

and used UMAP to do dimensional reduction for the top 20 components. After that,

we employ the 'FindClusters' function of Seurat for clustering analysis. Next, we

used the 'FindAllMarkers' function of Seurat to detect the specific expression of

metabolic genes for each metabolic state and used Single-sample GSEA to explore

the dynamic pattern of metabolic pathways by SingleSeqGset (Cillo, 2021) (version

0.1.2.9) of R.

Pathway activity analysis

GSVA (Hänzelmann, Castelo, & Guinney, 2013) is a widely used approach to assess

pathway enrichment of samples for gene expression data. We used the GSVA

package (version 1.51.0) to carry out enrichment analysis for the annotated KEGG

and Gene Ontology_Biological Process (GOBP) gene sets in the msigdbr package

(Subramanian et al., 2005) (version 7.5.1), followed by differential pathway analysis

using the limma package (Ritchie et al., 2015) (version 3.54.2). We not only

visualized such pathways that have significant differences (adj. p.val < 0.05), but

also employed the scMetabolism package (Y. Wu et al., 2022) (version 0.2.1) to do

enrichment analysis and visualization for KEGG and Reactome metabolic pathways.

Cell–cell communication analysis

We employed the CellChat (version 1.6.1) algorithm (Jin et al., 2021) to analyze and

quantify cell-cell communication between all major cell types. CellChat identifies

differentially over-expressed ligands and receptors for each cell group and models

the probability of communication between cell groups by the law of mass action.

Significant interactions are identified based on a permutation test (Good, 2013)

which can randomly permute the group labels of cells. We then used chord plots and

circle plots to describe the number of cell-cell interactions and communication

strength and visualized significant interactions (ligand-receptor pairs) from some cell

groups to other cell groups by the 'netVisual_bubble' function of CellChat.

InferCNV analysis

Malignant epithelial cells were identified using inferCNV (Tickle et al., 2019)

(version 1.14.2). We scored each cell for the extent of CNV signal and plotted the

cells on a dendrogram, and then we cut the dendrogram into 6 clusters. We labeled

all cells clustered with normal cells in the control group as "non-mag" and the

remaining clusters as "mag".

Correlation analysis between cell groups

We calculated the correlation between different cell groups based on the expression

of each cell on the genes. First, the mean value of gene expression was taken as the

representative value for each group of cells on each gene. Then, the top 1000 genes

with the greatest standard deviation were extracted to calculate the correlation

between groups by Spearman's (Spearman, 1987) method, and finally we used the

corrplot (Wei et al., 2013) (version 0.92) function to visualize the correlation matrix.

Cellular metabolic activity modeling

We modeled the metabolic state of individual cells by the Compass algorithm

(Wagner et al., 2021). We extracted the normalized gene expression matrix as input

and then ran Compass with default settings. The output of Compass is a reactions.tsv

file, where the numbers correspond to penalties for each reaction per cell. We

employ a negative log to have scores. If the score is great, the reaction is more

active.

Gene set module score

The AddModuleScore function of Seurat can be used to score any gene set. Its workflow is to firstly calculate the average value of all genes in the target gene set on a single-cell level, secondly divide the expression matrix into several bins based on the average value, and thirdly randomly select the control genes (genes outside the target gene set, default 100 genes) as the background value from each bin. Finally, we count all target and background genes as an average value, respectively. The score for the gene set is determined by subtracting the average background value from the average target gene value. Here we used this function to score gene sets of interest at the single-cell level.

Correlation analysis between gene sets

For correlation analysis between gene sets, the score of each gene set was defined as above. Here, we used the Pearson (Pearson, magazine, & science, 1901) method for correlation analysis and employed the ggpubr package (Kassambara, 2020) (version 0.6.0) to plot scatter plots.

Macrophage function analysis

Gene sets for well-known macrophage functions (M1-like polarization, M2-like polarization, Angiogenesis, Phagocytosis, Antigen presentation, and Immunosuppression) were collected from databases and relevant literature (Table S1). The GSVA package was used to analyze the functional enrichment of these gene sets.

Immunofluorescence staining

The sections of mouse lung tissue were first blocked with normal goat serum (10%). Primary antibodies were diluted and incubated with the sections at 4 °C overnight: anti-Ki67 (Abcam, Cambridge, USA) and anti-LDHA (Proteintech, Wuhan, China). The slides were then washed three times with PBS and incubated with Alexa-conjugated secondary antibodies (Jackson ImmunoResearch Laboratories, West Grove, USA) for 1 h at room temperature. The nuclei were finally

685 counterstained with DAPI (4', 6-diamidino-2-phenylindole, Sigma-Aldrich, St. Louis,

686 USA), and the sections were mounted. Images were captured by a fluorescence

687 microscope (Olympus, Tokyo, Japan). The pictures presented represent the results of

688 three sections, each of which was collected for nine fields.

Multiplex immunohistochemistry staining

Multiplex immunohistochemistry staining is a tyramine amplification-based method

for in situ labeling of tissue proteins. Combined with spectroscopic imaging

technology, the rich protein information contained in the tissue can be obtained. The

New Opal Polaris 7-color Manual IHC Kit (Akoya Biosciences, Massachusetts, USA)

was used for multiple staining. The Opal multi-labeling method used indirect

fluorescent labeling in which tissues were incubated with primary antibody and then

labeled with horseradish peroxidase (HRP) secondary antibody and fluorescence

amplification working solution. The non-covalently bound antibody was then

washed by thermal repair, then replaced with another primary antibody and

fluorescein substrate, and so on to achieve multiple labeling, followed by DAPI

staining and anti-fluorescence quencher sealing. Detailed experimental procedures

were described in the kit instructions. The multispectral tissue imaging system

(PerkinElmer Vectra®) and analysis software (Phenochart 1.0) were used for image

processing. The following antibodies were used in the staining: F4/80 (Cell

Signaling Technology, Danvers, USA), CD11c (Abcam), S100A4 (Abcam), CPT1A

(Abcam), ANXA2 (Abcam), and CD68 (Abcam).

Transient transfection in MH-S cells

The murine alveolar macrophage MH-S cell line was purchased from Procell Life

Science & Technology (Wuhan, China) and cultured in RPMI-1640 medium

(GIBCO) supplemented with 10% fetal bovine serum (FBS, GIBCO) and 0.05 mM

2-mercaptoethanol (Sigma-Aldrich). The plasmids of mS100a4 and NC were

commercially obtained from Hanbio (Shanghai, China) and transiently transfected

into MH-S cells with Lipofectamine 3000 (Invitrogen, Carlsbad, CA), respectively,

according to the manufacturer's instructions. S100a4 siRNAs (OBiO, Shanghai, China) were also transfected in the manner described above. ETO (MedChem Express) was used for targeted inhibition of CPT1A in MH-S.

Coculture assay in vitro

The mouse alveolar epithelial MLE12 cell line was kindly provided by Dr. Yi Li and cultured in DMEM/F12 medium (GIBCO) with 10% FBS. For the coculture experiment, the transfected MH-S cells were incubated with serum-free medium overnight. The CM was centrifuged at 200 g for 5 min and filtered through a 0.22 μm syringe filter. The collected CM was then applied to MLE12 cells.

qPCR

Total RNA was extracted by the SevenFast Total RNA Extraction Kit (Seven, Beijing, China). Reverse transcription for cDNA conversion was conducted with PrimeScript RT Master Mix (TAKARA, Tokyo, Japan). qPCR analysis of mRNA expression was measured by TB Green Premix Ex Taq (TAKARA) and normalized to β-actin. All reactions were performed in triplicate. The primers for target genes are listed below:

S100a4-F: TCCACAAATACTCAGGCAAAGAG

S100a4-R: GCAGCTCCCTGGTCAGTAG

β-actin-F: GGCTGTATTCCCCTCCATCG

β-actin-R: CCAGTTGGTAACAATGCCATGT

Western blotting

Total protein was extracted using lysis buffer containing protease inhibitor cocktail (MedChem Express, Monmouth Junction, USA) and quantitated by BCA protein assay kit (KeyGEN BioTECH, Beijing, China). Proteins were separated by sodium dodecyl sulfate–polyacrylamide gel electrophoresis (SDS-PAGE) and then transferred to a polyvinylidene fluoride (PVDF) membrane. Immunoblotting was performed with the following primary antibodies: S100A4 (Abcam), β-Actin (Cell Signaling Technology), SPA (Boster, Wuhan, China), E-Cadherin (Proteintech),

N-Cadherin (ABclonal, Wuhan, China), Vimentin (ABclonal), CD44 (Affinity,

Jiangsu, China), CD133 (Proteintech), p-ERK (Santa Cruz Biotechnology, Santa

Cruz, USA), MMP9 (Cell Signaling Technology), p-γH2AX (Abcam), VEGF (Santa

Cruz Biotechnology), TGF-β (Cell Signaling Technology), HIF-1α (Abcam),

c-MYC (Abcam), CPT1A (Abcam), ANXA2 (Abcam), PLG (Abcam), PPAR-γ

(Proteintech), CD206 (Abcam), and ARG1 (Proteintech).

Cell proliferation and migration assay

Cell viability was assessed by using Cell Counting Kit-8 (CCK8, 4A Biotech Co.,

Beijing, China). Cell proliferation was measured by colony formation assay. Cell

migration was determined by a wound-healing assay.

Cell cycle assay

PI/RNase Staining Solution (Sungene Biotech Co., Tianjin, China) was used for cell

cycle distribution measurement.

Intracellular ROS measurement

The intracellular ROS level was detected using the Reactive Oxygen Species Assay

Kit (Beyotime, Shanghai, China). After coculture, MLE12 cells were loaded with

DCFH-DA probe in situ and incubated in a 37 °C incubator for 30 min. Cells were

harvested, and the fluorescence intensity was measured by flow cytometry.

Transmission electron microscopy

Cell samples were prefixed with 2.5% glutaraldehyde, refixed with 1% osmium

tetroxide, dehydrated step by step with acetone, and then subjected to osmosis and

embedded with Epon-812 embedding agent. 60-90 nm ultrathin sections were made

by an ultrathin microtome and transferred to copper mesh, then stained with uranyl

acetate for 10-15 min, followed by lead citrate for 1-2 min at room temperature.

Image acquisition was carried out with a JEM-1400FLASH transmission electron

microscope (JEOL, Japan).

Mouse cytokine antibody array

CM from S100a4-OE and NC MH-S were collected for cytokine detection by

RayBio® G-Series Arrays (Guangzhou, China). Like a traditional sandwich-based

ELISA, this array uses a matched pair of cytokine-specific antibodies for detection.

After incubation with the sample, the target cytokines are captured by the antibodies

printed on the solid surface. A second biotin-labeled detection antibody is then added,

which recognizes a different epitope of the target cytokine. The

cytokine-antibody-biotin complex can then be visualized through the addition of the

streptavidin-conjugated Cy3 equivalent dye. After capturing the spot densities with a

laser scanner, array data analysis and functional enrichment analysis were

conducted.

Measurement of lipid droplet content and mitochondrial function

BODIPY 493/503 (GLPBIO, Montclair, USA) was used to detect intracellular lipid

droplet accumulation. TMRM and MitoSOX Green (Thermo Fisher Scientific) were

used to assess mitochondrial membrane potential and mitochondrial ROS production.

The fluorescence intensity was measured by flow cytometry.

Tube formation assay

HUVECs (2×10^4 cells per well) were seeded in 96-well plates coated with Matrigel

(Corning) and incubated in the CM from MH-S. Tube formation was examined

under an inverted microscope after 6 h. The total tube length and the number of

nodes were measured using ImageJ software.

Statistics

Statistical significance was determined using two-tailed unpaired Student's t-test,

and p-value <0.05 was considered significant (*p <0.05, **p <0.01, ***p <0.001,

****p <0.0001). The correlation coefficient (R) was calculated using the Pearson

method with 95% confidence intervals (CI), and corresponding p-values were added

to the plots. Statistical analyses were performed using R, Python, and GraphPad

Prism.

Data availability

The scRNA-seq data of mouse and human in this study have been deposited to codeocean (https://codeocean.com/capsule/4741227/tree).

Reference

Anderson, A. N., Conley, P., Klocke, C. D., Sengupta, S. K., Robinson, T. L., Fan,
Y., . . . Wong, M. H. (2023). Analysis of uveal melanoma scRNA sequencing
data identifies neoplastic-immune hybrid cells that exhibit metastatic potential.
BioRxiv : the Preprint Server For Biology. doi:10.1101/2023.10.24.563815

Barguilla, I., Maguer-Satta, V., Guyot, B., Pastor, S., Marcos, R., & Hernández, A.
(2023). In Vitro Approaches to Determine the Potential Carcinogenic Risk of
Environmental Pollutants. *International Journal of Molecular Sciences, 24*(9).
doi:10.3390/ijms24097851

Barton, N. W., Brady, R. O., Dambrosia, J. M., Di Bisceglie, A. M., Doppelt, S. H.,
Hill, S. C., . . . Argoff, C. E. (1991). Replacement therapy for inherited
enzyme deficiency--macrophage-targeted glucocerebrosidase for Gaucher's
disease. *The New England Journal of Medicine, 324*(21), 1464-1470.

Bian, Y., Li, W., Kremer, D. M., Sajjakulnukit, P., Li, S., Crespo, J., . . . Zou, W.
(2020). Cancer SLC43A2 alters T cell methionine metabolism and histone
methylation. *Nature, 585*(7824), 277-282. doi:10.1038/s41586-020-2682-1

Buck, M. D., Sowell, R. T., Kaech, S. M., & Pearce, E. L. (2017). Metabolic
Instruction of Immunity. *Cell, 169*(4), 570-586. doi:10.1016/j.cell.2017.04.004

Casanova-Acebes, M., Dalla, E., Leader, A. M., LeBerichel, J., Nikolic, J., Morales, B.
M., . . . Merad, M. (2021). Tissue-resident macrophages provide a
pro-tumorigenic niche to early NSCLC cells. *Nature, 595*(7868), 578-584.
doi:10.1038/s41586-021-03651-8

Cassetta, L., & Pollard, J. W. (2018). Targeting macrophages: therapeutic approaches
in cancer. *Nature Reviews. Drug Discovery, 17*(12), 887-904.
doi:10.1038/nrd.2018.169

Chen, Y.-C., Hsu, C.-L., Wang, H.-M., Wu, S.-G., Chang, Y.-L., Chen, J.-S., . . . Yang,
P.-C. (2024). Multi-omics Analysis Reveals Molecular Changes During Early
Progression of Precancerous Lesions to Lung Adenocarcinoma in
Never-Smokers. *Cancer Research*. doi:10.1158/0008-5472.CAN-24-0821

Cillo, A. (2021). Singleseqgset: Single-Cell Gene Set Enrichment Analysis.

de Visser, K. E., & Joyce, J. A. (2023). The evolving tumor microenvironment: From
cancer initiation to metastatic outgrowth. *Cancer Cell, 41*(3), 374-403.
doi:10.1016/j.ccell.2023.02.016

Feng, D., & Gao, B. (2021). From basic liver immunology to therapeutic
opportunities for liver diseases. *Cellular & Molecular Immunology, 18*(1), 1-3.
doi:10.1038/s41423-020-00607-2

Good, P. (2013). *Permutation tests: a practical guide to resampling methods for
testing hypotheses*: Springer Science & Business Media.

Greaves, M., & Maley, C. C. (2012). Clonal evolution in cancer. *Nature, 481*(7381),
306-313. doi:10.1038/nature10762

Grindheim, A. K., Saraste, J., & Vedeler, A. (2017). Protein phosphorylation and its
role in the regulation of Annexin A2 function. *Biochimica Et Biophysica Acta.*

General Subjects, 1861(11 Pt A), 2515-2529.
doi:10.1016/j.bbagen.2017.08.024

Hänzelmann, S., Castelo, R., & Guinney, J. (2013). GSVA: gene set variation analysis
for microarray and RNA-Seq data. *BMC Bioinformatics, 14*(1), 7.
doi:10.1186/1471-2105-14-7

Hao, Y., Hao, S., Andersen-Nissen, E., Mauck, W. M., 3rd, Zheng, S., Butler, A., . . .
Satija, R. (2021). Integrated analysis of multimodal single-cell data. *Cell,
184*(13), 3573-3587 e3529. doi:10.1016/j.cell.2021.04.048

Huang, Y., Jia, M., Yang, X., Han, H., Hou, G., Bi, L., . . . Ouyang, X. (2022).
Annexin A2: The diversity of pathological effects in tumorigenesis and
immune response. *International Journal of Cancer, 151*(4), 497-509.
doi:10.1002/ijc.34048

Jin, S., Guerrero-Juarez, C. F., Zhang, L., Chang, I., Ramos, R., Kuan, C.-H., . . . Nie,
Q. (2021). Inference and analysis of cell-cell communication using CellChat.
Nature Communications, 12(1), 1088. doi:10.1038/s41467-021-21246-9

Kassambara, A. J. R. p. v. (2020). Ggpubr:"ggplot2" Based Publication Ready Plots.
2020.

Kayser, K., Nwoye, J. O., Kosjerina, Z., Goldmann, T., Vollmer, E., Kaltner, H., . . .
Gabius, H.-J. (2003). Atypical adenomatous hyperplasia of lung: its incidence
and analysis of clinical, glycohistochemical and structural features including
newly defined growth regulators and vascularization. *Lung Cancer
(Amsterdam, Netherlands), 42*(2), 171-182.

Kazakova, E., Rakina, M., Sudarskikh, T., Iamshchikov, P., Tarasova, A., Tashireva,
L., . . . Larionova, I. (2023). Angiogenesis regulators S100A4, SPARC and
SPP1 correlate with macrophage infiltration and are prognostic biomarkers in
colon and rectal cancers. *Frontiers In Oncology, 13*, 1058337.
doi:10.3389/fonc.2023.1058337

Kerr, K. M., MacKenzie, S. J., Ramasami, S., Murray, G. I., Fyfe, N., Chapman, A.
D., . . . King, G. (2004). Expression of Fhit, cell adhesion molecules and
matrix metalloproteinases in atypical adenomatous hyperplasia and pulmonary
adenocarcinoma. *The Journal of Pathology, 203*(2), 638-644.

Kollmar, O., Scheuer, C., Menger, M. D., & Schilling, M. K. (2006). Macrophage
inflammatory protein-2 promotes angiogenesis, cell migration, and tumor
growth in hepatic metastasis. *Annals of Surgical Oncology, 13*(2), 263-275.

Kumar, S., Sharife, H., Kreisel, T., Mogilevsky, M., Bar-Lev, L., Grunewald, M., . . .
Keshet, E. (2019). Intra-Tumoral Metabolic Zonation and Resultant
Phenotypic Diversification Are Dictated by Blood Vessel Proximity. *Cell
Metabolism, 30*(1). doi:10.1016/j.cmet.2019.04.003

Landau, J. M., Wang, Z. Y., Yang, G. Y., Ding, W., & Yang, C. S. (1998). Inhibition of
spontaneous formation of lung tumors and rhabdomyosarcomas in A/J mice by
black and green tea. *Carcinogenesis, 19*(3), 501-507.

Li, S., Yu, J., Huber, A., Kryczek, I., Wang, Z., Jiang, L., . . . Zou, W. (2022).

Metabolism drives macrophage heterogeneity in the tumor microenvironment. *Cell Reports, 39*(1), 110609. doi:10.1016/j.celrep.2022.110609

Li, W., Tanikawa, T., Kryczek, I., Xia, H., Li, G., Wu, K., . . . Zou, W. (2018). Aerobic Glycolysis Controls Myeloid-Derived Suppressor Cells and Tumor Immunity via a Specific CEBPB Isoform in Triple-Negative Breast Cancer. *Cell Metabolism, 28*(1). doi:10.1016/j.cmet.2018.04.022

Liu, S., Zhang, H., Li, Y., Zhang, Y., Bian, Y., Zeng, Y., . . . Qin, Z. (2021). S100A4 enhances protumor macrophage polarization by control of PPAR-γ-dependent induction of fatty acid oxidation. *Journal For Immunotherapy of Cancer, 9*(6). doi:10.1136/jitc-2021-002548

Liu, Y., Myrvang, H. K., & Dekker, L. V. (2015). Annexin A2 complexes with S100 proteins: structure, function and pharmacological manipulation. *British Journal of Pharmacology, 172*(7), 1664-1676. doi:10.1111/bph.12978

Liu, Z., Liu, W., Wang, W., Ma, Y., Wang, Y., Drum, D. L., . . . Wang, X.-Y. (2023). CPT1A-mediated fatty acid oxidation confers cancer cell resistance to immune-mediated cytolytic killing. *Proceedings of the National Academy of Sciences of the United States of America, 120*(39), e2302878120. doi:10.1073/pnas.2302878120

Mantovani, A., Marchesi, F., Malesci, A., Laghi, L., & Allavena, P. (2017). Tumour-associated macrophages as treatment targets in oncology. *Nature Reviews. Clinical Oncology, 14*(7), 399-416. doi:10.1038/nrclinonc.2016.217

Marjanovic, N. D., Hofree, M., Chan, J. E., Canner, D., Wu, K., Trakala, M., . . . Tammela, T. (2020). Emergence of a High-Plasticity Cell State during Lung Cancer Evolution. *Cancer Cell, 38*(2). doi:10.1016/j.ccell.2020.06.012

Marjanovic, N. D., Hofree, M., Chan, J. E., Canner, D., Wu, K., Trakala, M., . . . Tammela, T. (2020). Emergence of a High-Plasticity Cell State during Lung Cancer Evolution. *Cancer Cell, 38*(2), 229-246 e213. doi:10.1016/j.ccell.2020.06.012

Mazzone, M., Menga, A., & Castegna, A. (2018). Metabolism and TAM functions-it takes two to tango. *The FEBS Journal, 285*(4), 700-716. doi:10.1111/febs.14295

Missiroli, S., Perrone, M., Genovese, I., Pinton, P., & Giorgi, C. (2020). Cancer metabolism and mitochondria: Finding novel mechanisms to fight tumours. *EBioMedicine, 59*, 102943. doi:10.1016/j.ebiom.2020.102943

Mori, M., Rao, S. K., Popper, H. H., Cagle, P. T., & Fraire, A. E. (2001). Atypical adenomatous hyperplasia of the lung: a probable forerunner in the development of adenocarcinoma of the lung. *Modern Pathology : an Official Journal of the United States and Canadian Academy of Pathology, Inc, 14*(2), 72-84.

Murugaiyan, G., Martin, S., & Saha, B. (2007). CD40-induced countercurrent conduits for tumor escape or elimination? *Trends In Immunology, 28*(11), 467-473.

Nakamura, M. T., Yudell, B. E., & Loor, J. J. (2014). Regulation of energy
metabolism by long-chain fatty acids. *Progress In Lipid Research, 53*, 124-144.
doi:10.1016/j.plipres.2013.12.001
Ossenkoppele, G. J., Lowenberg, B., Zachee, P., Vey, N., Breems, D., Van de
Loosdrecht, A. A., . . . Debnam, P. M. (2013). A phase I first-in-human study
with tefinostat - a monocyte/macrophage targeted histone deacetylase inhibitor
- in patients with advanced haematological malignancies. *British Journal of
Haematology, 162*(2), 191-201. doi:10.1111/bjh.12359
Pearson, K. J. T. L., Edinburgh,, magazine, D. p., & science, j. o. (1901). LIII. On
lines and planes of closest fit to systems of points in space. *2*(11), 559-572.
Pennycuick, A., Teixeira, V. H., AbdulJabbar, K., Raza, S. E. A., Lund, T., Akarca, A.
U., . . . Janes, S. M. (2020). Immune Surveillance in Clinical Regression of
Preinvasive Squamous Cell Lung Cancer. *Cancer Discovery, 10*(10),
1489-1499. doi:10.1158/2159-8290.CD-19-1366
Pich, O., Bailey, C., Watkins, T. B. K., Zaccaria, S., Jamal-Hanjani, M., & Swanton, C.
(2022). The translational challenges of precision oncology. *Cancer Cell, 40*(5),
458-478. doi:10.1016/j.ccell.2022.04.002
Pietras, A., Katz, A. M., Ekström, E. J., Wee, B., Halliday, J. J., Pitter, K. L., . . .
Holland, E. C. (2014). Osteopontin-CD44 signaling in the glioma perivascular
niche enhances cancer stem cell phenotypes and promotes aggressive tumor
growth. *Cell Stem Cell, 14*(3), 357-369. doi:10.1016/j.stem.2014.01.005
Prieto, L. I., Sturmlechner, I., Graves, S. I., Zhang, C., Goplen, N. P., Yi, E. S., . . .
Baker, D. J. (2023). Senescent alveolar macrophages promote early-stage lung
tumorigenesis. *Cancer Cell, 41*(7). doi:10.1016/j.ccell.2023.05.006
Raza, A., Franklin, M. J., & Dudek, A. Z. (2010). Pericytes and vessel maturation
during tumor angiogenesis and metastasis. *American Journal of Hematology,
85*(8), 593-598. doi:10.1002/ajh.21745
Reynolds, A. R., Moschetta, M., Yohannes, A. R., Walcott, F., Ashford, M., Szucs,
Z., . . . Galbraith, S. (2023). A View on Drug Development for Cancer
Prevention. *Cancer Discovery, 13*(5), 1058-1083.
doi:10.1158/2159-8290.CD-22-0776
Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., & Smyth, G. K.
(2015). limma powers differential expression analyses for RNA-sequencing
and microarray studies. *Nucleic Acids Research, 43*(7), e47-e47.
doi:10.1093/nar/gkv007 %J Nucleic Acids Research
Ruffell, B., Affara, N. I., & Coussens, L. M. (2012). Differential macrophage
programming in the tumor microenvironment. *Trends In Immunology, 33*(3),
119-126. doi:10.1016/j.it.2011.12.001
Sakurai, T., & Kudo, M. (2011). Signaling pathways governing tumor angiogenesis.
Oncology, 81 Suppl 1, 24-29. doi:10.1159/000333256
Santibáñez-Andrade, M., Quezada-Maldonado, E. M., Rivera-Pineda, A., Chirino, Y.
I., García-Cuellar, C. M., & Sánchez-Pérez, Y. (2023). The Road to Malignant

Cell Transformation after Particulate Matter Exposure: From Oxidative Stress
to Genotoxicity. *International Journal of Molecular Sciences, 24*(2).
doi:10.3390/ijms24021782

Semov, A., Moreno, M. J., Onichtchenko, A., Abulrob, A., Ball, M., Ekiel, I., . . .
Alakhov, V. (2005). Metastasis-associated protein S100A4 induces
angiogenesis through interaction with Annexin II and accelerated plasmin
formation. *The Journal of Biological Chemistry, 280*(21), 20833-20841.

Sharma, M. C. (2019). Annexin A2 (ANX A2): An emerging biomarker and potential
therapeutic target for aggressive cancers. *International Journal of Cancer,*
144(9), 2074-2081. doi:10.1002/ijc.31817

Sica, A., & Mantovani, A. (2012). Macrophage plasticity and polarization: in vivo
veritas. *The Journal of Clinical Investigation, 122*(3), 787-795.
doi:10.1172/JCI59643

Sica, A., Schioppa, T., Mantovani, A., & Allavena, P. (2006). Tumour-associated
macrophages are a distinct M2 polarised population promoting tumour
progression: potential targets of anti-cancer therapy. *European Journal of*
Cancer (Oxford, England : 1990), 42(6), 717-727.

Sivakumar, S., Lucas, F. A. S., McDowell, T. L., Lang, W., Xu, L., Fujimoto, J., . . .
Kadara, H. (2017). Genomic Landscape of Atypical Adenomatous Hyperplasia
Reveals Divergent Modes to Lung Adenocarcinoma. *Cancer Res, 77*(22),
6119-6130. doi:10.1158/0008-5472.CAN-17-1605

Sivakumar, S., Lucas, F. A. S., McDowell, T. L., Lang, W., Xu, L., Fujimoto, J., . . .
Kadara, H. (2017). Genomic Landscape of Atypical Adenomatous Hyperplasia
Reveals Divergent Modes to Lung Adenocarcinoma. *Cancer Research, 77*(22),
6119-6130. doi:10.1158/0008-5472.CAN-17-1605

Spearman, C. (1987). The proof and measurement of association between two things.
By C. Spearman, 1904. *Am J Psychol, 100*(3-4), 441-471.

Subramanian, A., Tamayo, P., Mootha, V. K., Mukherjee, S., Ebert, B. L., Gillette, M.
A., . . . Mesirov, J. P. (2005). Gene set enrichment analysis: a
knowledge-based approach for interpreting genome-wide expression profiles.
Proc Natl Acad Sci U S A, 102(43), 15545-15550.
doi:10.1073/pnas.0506580102

Tan, Z., Xiao, L., Tang, M., Bai, F., Li, J., Li, L., . . . Cao, Y. (2018). Targeting
CPT1A-mediated fatty acid oxidation sensitizes nasopharyngeal carcinoma to
radiation therapy. *Theranostics, 8*(9), 2329-2347. doi:10.7150/thno.21451

Teixeira, V. H., Pipinikas, C. P., Pennycuick, A., Lee-Six, H., Chandrasekharan, D.,
Beane, J., . . . Janes, S. M. (2019). Deciphering the genomic, epigenomic, and
transcriptomic landscapes of pre-invasive lung cancer lesions. *Nature*
Medicine, 25(3), 517-525. doi:10.1038/s41591-018-0323-0

Tickle, T., Tirosh, I., Georgescu, C., Brown, M., Haas, B. J. K. C. O., Broad Institute
of MIT, & Harvard. (2019). inferCNV of the Trinity CTAT Project.

Torisu, H., Ono, M., Kiryu, H., Furue, M., Ohmoto, Y., Nakayama, J., . . . Kuwano, M.

(2000). Macrophage infiltration correlates with tumor stage and angiogenesis in human malignant melanoma: possible involvement of TNFalpha and IL-1alpha. *International Journal of Cancer, 85*(2), 182-188.

Travis, W. D., Brambilla, E., Nicholson, A. G., Yatabe, Y., Austin, J. H. M., Beasley, M. B., . . . Wistuba, I. (2015). The 2015 World Health Organization Classification of Lung Tumors: Impact of Genetic, Clinical and Radiologic Advances Since the 2004 Classification. *Journal of Thoracic Oncology : Official Publication of the International Association For the Study of Lung Cancer, 10*(9), 1243-1260. doi:10.1097/JTO.0000000000000630

Travis, W. D., Brambilla, E., Noguchi, M., Nicholson, A. G., Geisinger, K. R., Yatabe, Y., . . . Yankelewitz, D. (2011). International association for the study of lung cancer/american thoracic society/european respiratory society international multidisciplinary classification of lung adenocarcinoma. *Journal of Thoracic Oncology : Official Publication of the International Association For the Study of Lung Cancer, 6*(2), 244-285. doi:10.1097/JTO.0b013e318206a221

Umar, A., Dunn, B. K., & Greenwald, P. (2012). Future directions in cancer prevention. *Nature Reviews. Cancer, 12*(12), 835-848. doi:10.1038/nrc3397

van Hinsbergh, V. W. M., & Koolwijk, P. (2008). Endothelial sprouting and angiogenesis: matrix metalloproteinases in the lead. *Cardiovascular Research, 78*(2), 203-212.

Vimalraj, S. (2022). A concise review of VEGF, PDGF, FGF, Notch, angiopoietin, and HGF signalling in tumor angiogenesis with a focus on alternative approaches and future directions. *International Journal of Biological Macromolecules, 221*, 1428-1438. doi:10.1016/j.ijbiomac.2022.09.129

Vitale, I., Manic, G., Coussens, L. M., Kroemer, G., & Galluzzi, L. (2019). Macrophages and Metabolism in the Tumor Microenvironment. *Cell Metabolism, 30*(1), 36-50. doi:10.1016/j.cmet.2019.06.001

Wagner, A., Wang, C., Fessler, J., DeTomaso, D., Avila-Pacheco, J., Kaminski, J., . . . Yosef, N. (2021). Metabolic modeling of single Th17 cells reveals regulators of autoimmunity. *Cell, 184*(16), 4168-4185.e4121. doi:10.1016/j.cell.2021.05.045

Wang, H., Arulraj, T., Anbari, S., & Popel, A. S. (2024). Quantitative systems pharmacology modeling of macrophage-targeted therapy combined with PD-L1 inhibition in advanced NSCLC. *Clinical and Translational Science, 17*(6), e13811. doi:10.1111/cts.13811

Wang, J., Wang, Y., Chu, Y., Li, Z., Yu, X., Huang, Z., . . . Zheng, L. (2021). Tumor-derived adenosine promotes macrophage proliferation in human hepatocellular carcinoma. *Journal of Hepatology, 74*(3), 627-637. doi:10.1016/j.jhep.2020.10.021

Wang, T., Wang, Z., Niu, R., & Wang, L. (2019). Crucial role of Anxa2 in cancer progression: highlights on its novel regulatory mechanism. *Cancer Biology & Medicine, 16*(4), 671-687. doi:10.20892/j.issn.2095-3941.2019.0228

Wang, Y.-N., Zeng, Z.-L., Lu, J., Wang, Y., Liu, Z.-X., He, M.-M., . . . Xu, R.-H. (2018). CPT1A-mediated fatty acid oxidation promotes colorectal cancer cell metastasis by inhibiting anoikis. *Oncogene, 37*(46), 6025-6040. doi:10.1038/s41388-018-0384-z

Watabe, T., Takahashi, K., Pietras, K., & Yoshimatsu, Y. (2023). Roles of TGF-β signals in tumor microenvironment via regulation of the formation and plasticity of vascular system. *Seminars In Cancer Biology, 92*, 130-138. doi:10.1016/j.semcancer.2023.04.007

Wei, T., Simko, V., Levy, M., Xie, Y., Jin, Y., & Zemla, J. J. R. p. v. (2013). corrplot: Visualization of a correlation matrix. *230*(11).

Weichert, W., & Warth, A. (2014). Early lung cancer with lepidic pattern: adenocarcinoma in situ, minimally invasive adenocarcinoma, and lepidic predominant adenocarcinoma. *Current Opinion In Pulmonary Medicine, 20*(4), 309-316. doi:10.1097/MCP.0000000000000065

Weichert, W., & Warth, A. (2014). Early lung cancer with lepidic pattern: adenocarcinoma in situ, minimally invasive adenocarcinoma, and lepidic predominant adenocarcinoma. *Curr Opin Pulm Med, 20*(4), 309-316. doi:10.1097/MCP.0000000000000065

Wen, Y., Lambrecht, J., Ju, C., & Tacke, F. (2021). Hepatic macrophages in liver homeostasis and diseases-diversity, plasticity and therapeutic opportunities. *Cellular & Molecular Immunology, 18*(1), 45-56. doi:10.1038/s41423-020-00558-8

Wu, H., Han, Y., Rodriguez Sillke, Y., Deng, H., Siddiqui, S., Treese, C., . . . Glauben, R. (2019). Lipid droplet-dependent fatty acid metabolism controls the immune suppressive phenotype of tumor-associated macrophages. *EMBO Molecular Medicine, 11*(11), e10698. doi:10.15252/emmm.201910698

Wu, Y., Yang, S., Ma, J., Chen, Z., Song, G., Rao, D., . . . Gao, Q. (2022). Spatiotemporal Immune Landscape of Colorectal Cancer Liver Metastasis at Single-Cell Level. *Cancer Discov, 12*(1), 134-153. doi:10.1158/2159-8290.Cd-21-0316

Xia, H., Li, S., Li, X., Wang, W., Bian, Y., Wei, S., . . . Zou, W. (2020). Autophagic adaptation to oxidative stress alters peritoneal residential macrophage survival and ovarian cancer metastasis. *JCI Insight, 5*(18). doi:10.1172/jci.insight.141115

Yang, L., Li, A., Yu, W., Wang, H., Zhang, L., Wang, D., . . . Zhang, Y. (2025). Blockade of purine metabolism reverses macrophage immunosuppression and enhances anti-tumor immunity in non-small cell lung cancer. *Drug Resistance Updates : Reviews and Commentaries In Antimicrobial and Anticancer Chemotherapy, 78*, 101175. doi:10.1016/j.drup.2024.101175

Yatabe, Y., Borczuk, A. C., & Powell, C. A. (2011). Do all lung adenocarcinomas follow a stepwise progression? *Lung Cancer (Amsterdam, Netherlands), 74*(1). doi:10.1016/j.lungcan.2011.05.021

1093   Yu, C., Hou, L., Cui, H., Zhang, L., Tan, X., Leng, X., & Li, Y. (2018). LDHA
1094   upregulation independently predicts poor survival in lung adenocarcinoma, but
1095   not in lung squamous cell carcinoma. *Future Oncology (London, England),*
1096   *14*(24), 2483-2492. doi:10.2217/fon-2018-0177

1097   Zacharias, W. J., Frank, D. B., Zepp, J. A., Morley, M. P., Alkhaleel, F. A., Kong,
1098   J., . . . Morrisey, E. E. (2018). Regeneration of the lung alveolus by an
1099   evolutionarily conserved epithelial progenitor. *Nature, 555*(7695), 251-255.
1100   doi:10.1038/nature25786

1101

1102

Author contribution

Hong Huang: Conceptualization; Methodology; Validation; Investigation; Resources;

Writing – original draft preparation; Writing – review & editing; Visualisation

Ying Yang: Methodology; Validation; Investigation; Resources

Qiuju Zhang: Software; Formal analysis; Data curation; Visualisation

Yongfeng Yang: Conceptualization; Methodology; Writing – review & editing

Zhenqi Xiong: Software; Formal analysis; Data curation

Shengqiang Mao: Software; Formal analysis

Tingting Song: Software; Formal analysis

Yilong Wang: Validation; Investigation

Zhiqiang Liu: Validation; Resources

Hong Bu: Conceptualization; Writing – review & editing; Supervision; Project

administration

Li Zhang: Conceptualization; Resources; Writing – review & editing; Supervision;

Project administration; Funding acquisition

Le Zhang: Software; Writing – review & editing; Supervision; Project administration;

Funding acquisition

1120

1121 **Competing interests**

1122 The authors declare no conflict of interest.

1123

Funding

This work was supported by the National Natural Science Foundation of China [92159302 and 62372316], National Science and Technology Major Project [2021YFF1201200], Sichuan Science and Technology Program Key Project [2024YFHZ0091], Sichuan Science and Technology Project [2022ZDZX0018], and 1.3.5 Project for Disciplines of Excellence, West China Hospital, Sichuan University [ZYGD2200].

Figure legends

Figure 1. Histopathological grading and single-cell transcriptome profiles of mouse LUAD precancerous lesions.

A. Analysis and experimental flow chart of this study.

B. H&E images of normal tissue and three stages of precancerous lesions (AAH, adenoma, and AIS) in mice. Scale bar: 100 µm.

C. Immunofluorescence staining of Ki67 at the four histopathological stages. Scale bar: 100 µm.

D. UMAP plot of 13 cell types from mouse scRNA-seq data.

E. Dotplot of marker genes in all cell types.

F. Cell distribution at the four stages.

G. Reduced transcriptional homogeneity with progression of precancerous lesions. Transcriptional heterogeneity between cells is inversely proportional to normalized mutual information (NMI).

AAH: atypical adenomatous hyperplasia; AIS: adenocarcinoma in situ; MIA: minimally invasive adenocarcinoma; IA: invasive adenocarcinoma; CM: conditioned medium; DC: dendritic cell; SMC: smooth muscle cell; ***$p < 0.001$.

Figure 2. Identification of initiation-associated epithelial marker LDHA and analysis of malignant epithelial cells.

A. UMAP plot of epithelial cell subtypes.

B. Dotplot showing the significance (P-value) and strength (communication probability) of specific interactions between epithelial cells and other cell types at the four stages. Data was obtained by Cell Chat analysis.

C. Immunofluorescence staining of LDHA at the four stages. Scale bar: 50 µm.

D. LDHA expression in LUAD and normal tissues from the TCGA database.

E. LDHA expression in stage I-IV LUAD and normal tissues from the TCGA database.

F. Correlation of LDHA expression with overall survival of LUAD patients in the TCGA database.

G. Copy number variations (CNVs) (red, amplifications; blue, deletions) across the chromosomes (columns) inferred from the scRNA-seq of each epithelial cell (rows).

H. scMetabolism analysis of malignant epithelial cells at precancerous stages.

I. GSVA enrichment analysis of malignant epithelial cells at precancerous stages.

AT1/2: type I/II alveolar epithelial cells; LUAD: lung adenocarcinoma; *$p < 0.05$.

Figure 3. S100a4[+] alv-macro was active in lipid metabolism at the AAH stage.

A. Malignant epithelial cells showed the strongest communication weight with alveolar macrophages, as shown by Cell Chat analysis.

B. UMAP plot of S100a4 expression in alveolar macrophages.

C. scMetabolism analysis of S100a4[+] alv-macro at the four stages.

D. Changes in scores of representative lipid metabolism-related gene sets across the four stages.

E. Compass analysis showing the reaction activities of fatty acid metabolism across the four stages.

F. Density plots of Cpt1a and Acot2 at the four stages. The x-axis represents the gene expression level, and the y axis represents the density of numbers of cells.

G. Multiplex immunofluorescence validation of F4/80[+]/CD11c[+]/S100A4[+] alv-macro in mouse normal

1174   and AAH tissues and comparison of tissue expression of CPT1A in this subpopulation. Scale bar:

1175   10 µm.

1176   EC: endothelial cell.

1177

Figure 4. The pro-angiogenic function of S100a4$^+$ alv-macro was related to fatty acid metabolism.

A. Cell proportion comparison of S100a4$^+$ alv-macro in alveolar macrophages.

B. Macrophage functional program analysis of S100a4$^+$ alv-macro across the four stages.

C. Pearson correlation analysis of angiogenesis and M2-like function with fatty acid metabolic

reactions in S100a4$^+$ alv-macro.

D. Density plots of Anxa2 and Ramp1 at the four stages.

E. Correlation analysis of Cpt1a and Anxa2 expression in S100a4$^+$ alv-macro.

F. Multiplex immunofluorescence validation of the correlation between CPT1A and ANXA2

expression in F4/80$^+$/CD11c$^+$/S100A4$^+$ alv-macro of mouse normal and AAH tissues. Scale bar: 10

µm.

1188

Figure 5. S100a4-OE MH-S promoted malignant transformation of MLE12 epithelial cells in vitro.

A. S100a4 mRNA expression level in MH-S after transfection.

B. S100A4 protein expression level in MH-S after transfection.

C. CCK8 assay showing cell proliferation of MLE12 after coculture with S100a4-OE MH-S.

D. Colony-forming ability of MLE12 after coculture, as shown by the colony formation assay.

E. Cell cycle distribution of MLE12 after coculture, as shown by cell cycle analysis.

F. Wound healing assay showing cell migration of MLE12 after coculture.

G. Intracellular ROS level of MLE12 after coculture.

H. Transmission electron microscopy showing the morphological changes of mitochondria in MLE12

after coculture.

I. Western blotting of DNA damage marker p-γH2AX, EMT markers (E-Cadherin, N-Cadherin, and

Vimentin), and stem-like markers (CD44 and CD133) in MLE12 after coculture.

J. Western blotting of tumorigenesis-associated proteins (c-MYC, p-ERK, SPA, VEGF, and HIF-1α)

in MLE12 after coculture.

K. Western blotting of macrophage pro-tumor indicators (VEGF, MMP9, TGF-β, and HIF-1α) in

S100a4-OE MH-S.

L. Heatmap of cytokine assay for differential factors secreted after MH-S transfection.

M.   Enrichment analysis of GO biological processes for differential factors.

OE: overexpression; NC: negative control; ROS: reactive oxygen species; *p < 0.05, **p < 0.01, ***p

< 0.001, ****p < 0.0001.

1209

Figure 6. S100a4-OE MH-S controlled pro-angiogenic and pro-tumorigenic functions through CPT1A

induction

A. Determination of intracellular lipid droplet accumulation, mitochondrial membrane potential, and

mitochondrial ROS by flow cytometry.

B. Tube formation of HUVECs after coculture with S100a4-OE MH-S, quantified by total tube length

and number of nodes.

C. Western blotting of fatty acid metabolism-related CPT1A in S100a4-OE MH-S.

D. Western blotting of ANXA2 and PLG in S100a4-OE MH-S and cocultured HUVEC and MLE12.

E. CCK8 assay of cocultured MLE12 after treatment of ETO to S100a4-OE MH-S.

F. Wound healing assay of cocultured MLE12 after treatment of ETO to S100a4-OE MH-S.

G. Intracellular ROS production of cocultured MLE12 after treatment of ETO to S100a4-OE MH-S.

H. Tube formation of cocultured HUVECs after treatment of ETO to S100a4-OE MH-S.

I. Western blotting of fatty acid metabolism-related proteins (CPT1A and PPAR-γ),
angiogenesis-related proteins (ANXA2, VEGF, TGF-β, HIF-1α, and MMP9), and M2 polarization
markers (CD206 and ARG1) in S100a4-OE MH-S after treatment of ETO.

J. Western blotting of angiogenesis-related proteins (ANXA2, VEGF, and HIF-1α) in cocultured
MLE12 after treatment of ETO to S100a4-OE MH-S.

MFI: mean fluorescence intensity; ETO: etomoxir; *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p <$
0.0001.

1229

Figure 7. S100A4[+] alv-macro with a similar pattern in the human AAH stage

A. H&E images of five stages of human LUAD development (normal, AAH, AIS, MIA, and IA).
Scale bar: 100 µm.

B. UMAP plot of subclusters of alveolar macrophages.

C. UMAP plot of S100A4 expression in alveolar macrophages.

D. Compass analysis of the reaction activities of fatty acid metabolism in S100A4[+] alv-macro across
the five stages.

E. Macrophage functional analysis of S100A4[+] alv-macro across the five stages.

F. Dotplot of expression of CPT1A, ACOT2, and ANXA2 in the five stages.

G. Multiplex immunofluorescence staining of CPT1A and ANXA2 expression in
CD68[+]/CD11c[+]/S100A4[+] alv-macro of human normal and AAH tissues. Scale bar: 20 µm.