Ketogenesis protects against MASLD-MASH progression through fat oxidation-

independent mechanisms

Eric D. Queathem[1,2,3], David B. Stagg[1], Alisa B. Nelson[1], Alec B. Chaves[4], Scott B. Crown[4], Kyle

Fulghum[1], D. Andre d'Avignon[1], Justin R. Ryder[5], Patrick J. Bolan[6], Abdirahman Hayir[1], Jacob

R. Gillingham[1,2], Shannon Jannatpour[7], Ferrol I. Rome[1], Ashley S. Williams[4], Deborah M.

Muoio[4,8,9], Sayeed Ikramuddin[7], Curtis Hughey[1], Patrycja Puchalska[1*], Peter A. Crawford[1,2*]

[1]Division of Molecular Medicine, Department of Medicine, University of Minnesota Medical

School, Minneapolis, MN, 55455 USA

[2]Department of Biochemistry, Molecular Biology and Biophysics, University of Minnesota

Medical School, Minneapolis, MN, 55455 USA

[3]Department of Integrative Biology and Physiology, University of Minnesota Medical School,

Minneapolis, MN, 55455 USA

[4]Duke Molecular Physiology Institute and Sarah W. Stedman Nutrition and Metabolism Center,

Duke University Medical Center, Durham, NC 27701, USA

[5]Department of Surgery, Lurie Children's Hospital, Chicago, IL 60611 USA

[6]Department of Radiology, University of Minnesota Medical School, Minneapolis, MN, 55455

USA

[7]Department of Surgery, University of Minnesota, Minneapolis, MN, USA

[8]Division of Endocrinology, Metabolism, and Nutrition, Department of Medicine, Duke University

Medical Center, Durham, NC 27710, USA

[9]Department of Pharmacology and Cancer Biology, Duke University Medical Center, Durham,

NC 27710, USA

*Contact for correspondence, crawforp@umn.edu or ppuchals@umn.edu

Abstract.

The progression of metabolic-dysfunction-associated steatotic liver disease (MASLD) to

metabolic-dysfunction-associated steatohepatitis (MASH) involves complex alterations in both

liver-autonomous and systemic metabolism that influence the liver's balance of fat accretion and

disposal. Here, we quantify the relative contribution of hepatic oxidative pathways to liver injury

in MASLD-MASH. Using NMR spectroscopy, UHPLC-MS, and GC-MS, we performed stable-

isotope tracing and formal flux modeling to quantify hepatic oxidative fluxes in humans across

the spectrum of MASLD-MASH, and in mouse models of impaired ketogenesis. We found in

humans with MASH, that liver injury correlated positively with ketogenesis and total fat

oxidation, but not with turnover of the tricarboxylic acid cycle. The use of loss-of-function mouse

models demonstrated that disruption of mitochondrial HMG-CoA synthase (HMGCS2), the rate-

limiting step of ketogenesis, impairs overall hepatic fat oxidation and induces a MASLD-MASH-

like phenotype. Furthermore, disruption of mitochondrial β-hydroxybutyrate dehydrogenase

(BDH1), the terminal step of ketogenesis, also impaired fat oxidation, but surprisingly did not

exacerbate steatotic liver injury. Taken together, these findings suggest that quantifiable

variations in overall hepatic fat oxidation may not be a primary determinant of MASLD-to-MASH

progression, but rather, that maintenance of hepatic ketogenesis could serve a protective role

through alternative fat oxidation-independent mechanisms.

Keywords

MASLD-MASH, obesity, ketogenesis, fat oxidation, TCA cycle turnover, stable-isotope tracing,

metabolic flux analysis

Introduction

The global prevalence of metabolic-dysfunction-associated steatotic liver disease (MASLD) is

surging and is associated with mortality and multiple comorbidities.[1,2] The natural history of

MASLD is tightly linked to obesity and insulin resistance, and involves both liver-autonomous

and systemic metabolic abnormalities that drive the ectopic storage of triacylglycerol (TAG)

species in hepatocytes (*i.e.,* hepatic steatosis), which is strongly associated with cellular injury,

inflammation, and the development of metabolic-dysfunction-associated steatohepatitis

(MASH), a progressive fibrotic liver disease that markedly increases the risks of cirrhosis and

hepatocellular carcinoma.[3–5] While the severity of hepatic steatosis and the underlying metabolic

disturbances precipitating TAG accretion may predict histological evidence of liver injury,

steatosis alone does not account for MASLD-MASH pathogenesis. For any degree of steatosis,

the development of MASH is marked by the appearance of oxidative and inflammatory stress,

coupled to hepatocellular injury (*i.e.,* hepatocyte ballooning), which collectively can be quantified

using the NAFLD (corresponding to the prior term referencing MASLD, nonalcoholic fatty liver

disease) activity score (NAS), a histological scoring system ranging 0-8, which encompasses

steatosis (0-3), inflammation (0-3) and hepatocyte ballooning (0-2). MASH is defined as a NAS

score ≥ 4, and typically coincides with fibrosis, though fibrosis is not included when calculating

NAS.[6]

Currently there is only one FDA approved drug available for the treatment of MASLD-

MASH, while several others have failed due to poor efficacy and/or toxicity, illustrating a need to

better understand the underlying drivers of hepatic fat accumulation, and the mechanistic link

this has to the development of liver injury and advanced liver disease.[7,8] Though sodium-

glucose cotransporter 2 (SGLT2) inhibitors and incretin mimetics show great promise for

MASLD, approaches that (i) are liver-specific, indicated for patients without diabetes or body

mass index (BMI) > 30, (ii) limit the requirement for lifelong high-cost pharmacotherapy, and/or

(iii) limit toxicities, remain an unmet need.[9,10] Prior studies have correlated the accumulation of

liver TAGs in obesity and/or MASLD to a rise in hepatic *de novo* lipogenesis (DNL)[11–16], a

diminution in polyunsaturated fatty acids (PUFAs)[17–21], and an acceleration in hepatic oxidative

fluxes[22–25]. Other studies have shown no changes in hepatic fat oxidation[26–29], or have reported

impairments in oxidative metabolism.[30–34] Therefore, the lack of consensus regarding

pathological alterations in hepatic oxidative fluxes throughout the natural history of MASLD-

MASH progression illustrates one of the key obstacles in developing effective liver-targeted

therapeutics.

Oxidative metabolism in the liver yields reducing equivalents (RE's) in the forms of

reduced nicotinamide adenine dinucleotide (NADH) and flavin adenine dinucleotide ($FADH_2$),

which are primarily sourced from hepatic fat oxidation via (1) the β-oxidation pathway, which

generates acetyl-CoA, and (2) terminal oxidation of acetyl-CoA in the tricarboxylic acid (TCA)

cycle.[35] RE production is also directly coupled to the production of ATP, which fuels

phosphoenolpyruvate (PEP)-derived gluconeogenesis (GNG) via the non-oxidative entry

(anaplerosis) and exit (cataplerosis) of intermediates through the TCA cycle[36]. In both human

and rodent models of MASLD, hepatic TCA cycle turnover and PEP-derived GNG are increased

relative to controls,[25,36,37] and very low density lipoprotein (VLDL) secretion increases with

worsening steatosis[38,39]. The stimulation of fat disposal pathways may offload excess TAGs,

however, the rise in lipid disposal is not sufficient to compensate for the rate of lipid appearance,

leading to ectopic accumulation of fat, which predisposes the liver to injury. In addition to VLDL

secretion and terminal oxidation in the TCA cycle, ketogenesis, which produces D-beta-

hydroxybutyrate (D-βOHB) and acetoacetate (AcAc), is a major conduit supporting fat disposal

in the liver. Congenital deficiency of the fate-committing enzyme of ketogenesis, 3-hydroxy-3-

methylglutaryl-CoA (HMG-CoA) synthase 2 (HMGCS2), is linked to hepatomegaly and fatty

liver.[40,41] Moreover, programmed ketogenic insufficiency in mice is associated with more

aggressive MASLD progression.[42–46] However, the relative rate of ketogenesis in MASLD,

compared to controls, has varied among studies.[22,27,34,37,47] As such, whether alterations in

ketogenesis in human MASLD-MASH plays driver, bystander, or compensating roles is

debated, as are the mechanisms that link ketogenesis to oxidative metabolism and liver injury.

To date, *in vivo* hepatic oxidative fluxes have only been quantified in humans with

uncomplicated MASLD but have not been measured in humans with histologically confirmed

MASH. To understand the metabolic drivers and/or predictors of MASLD-MASH development,

progression, and resolution, we quantified hepatic oxidative fluxes in histologically confirmed

patients with MASH using nuclear magnetic resonance (NMR) spectroscopy, ultra-high

performance liquid chromatography coupled to mass spectrometry (UHPLC-MS), and

sophisticated $^2H/^{13}C$ stable isotope tracing. Loss of ketogenesis mouse models were used to

determine the relationships among ketogenesis, the rate of overall fat oxidation, and liver injury.

Together, our observations are consistent with the notion that while ketogenesis and fat

oxidation are key inputs into the liver's balance of lipid accretion and disposal, the influence of

ketogenesis in overall hepatic fat oxidation may not be a primary determinant of MASLD-to-

MASH progression. Therefore, maintenance of hepatic ketogenesis could serve as a critical

protective role through alternative fat oxidation-independent mechanisms.

Results

Hepatic ketogenesis correlates with NAS in humans. During the progression of

uncomplicated MASLD, reports of hepatic oxidative fluxes have been varied, with studies

reporting (1) accelerations[22–25], (2) no changes[26–29], or (3) impairments[30–34] in oxidative

metabolism. However, oxidative fluxes have never been quantified *in vivo* in humans with

histologically confirmed MASH **(Figure 1A).** To determine rates of hepatic fat oxidation and

GNG in humans with MASH, participants were recruited from two clinical trials, NCT03997422

and NCT03587831. Recruited participants remained weight stable during the screening process

and during the conduction of metabolic assessments. Participants with a BMI ≥ 35 were

screened with magnetic resonance imaging (MRI) to identify patients with a liver proton density

fat fraction (PDFF) > 5%, indicative of hepatic steatosis, then a total of 14 participants were

recruited to the study outlined in **Figure 1B.** After liver biopsy collection, glucose homeostasis

and body composition were quantified using a frequently sampled intravenous glucose

tolerance test (FSIVGTT), and dual-energy X-ray absorptiometry (DXA), respectively.

Thereafter, during a 20hr fast, oral ^2H_2O and [U-$^{13}C_3$] propionate tracers were delivered to enrich

the circulating glucose pool across hydrogen and carbon atoms, whose positional isotopomeric

labeling distribution supports modeling of *relative* reaction velocities (i.e., flux) through hepatic

intermediary metabolic pathways *in vivo.* Simultaneous to the quantification of plasma glucose

^{13}C- and ^2H-positional enrichment, the absolute rates of endogenous glucose production (EGP)

(V_{EGP}) and ketogenesis ($V_{Ra\beta OHB}$) were measured at metabolic and isotopic steady state, after a

2hr infusion of [3,4-$^{13}C_2$]glucose and [U-$^{13}C_4$]βOHB, respectively, thereby allowing the absolute

rates of hepatic intermediary metabolic pathways to be quantified **(Supplemental Figure**

1).[35,48–50] During the screening phase, and prior to metabolic flux assessments, a liver biopsy

was collected and used to histologically grade liver health, which was collectively summarized

by the NAS score, which ranged from 1-8 across all participants. All 14 patients exhibited

histological signs of liver steatosis, and 12 out of 14 participants displayed signs of either

hepatocyte cell injury (*i.e.,* ballooning) and/or lobular inflammation **(Figure 1C).** Nine of the 14

participants exhibited histopathological fibrosis. As expected, NAS strongly correlated with

PDFF (r = 0.61, P value < 0.05) and liver fibrosis (r = 0.79, P value < 0.001) **(Figure 1D)**, but did

not correlate with BMI **(Figure 1E).** NAS also correlated positively with the acute insulin

response to glucose (AIRg) (r = 0.74, P value = 0.004) during the FSIVGTT, suggestive of a

correlation between insulin resistance and NAS **(Figure 1F)**, however, NAS did not correlate

with the homeostatic model assessment for insulin resistance (HOMA-IR) **(Figure 1G).**

We then examined the metabolic flux modeling data acquired in this cohort **(Figure 2A-**

B). Across all 14 participants, 30 ± 8% of V_{EGP} arose from glycogenolysis ($V_{Glycogen}$), 19 ± 7%

from glycerol ($V_{Glycerol}$) and 51 ± 7% from PEP (V_{PEP}) **(Figure 2C).** Next, by simultaneously

quantifying the rates of (1) TCA cycle flux, V_{CS}, (2) total ketone body (TKB) production, V_{RaTKB},

and (3) GNG sourcing pathways, $V_{Glycerol}$ and V_{PEP}, the majority of hepatic fat oxidation and RE

production could be quantified. The majority of RE's produced in the fasted liver originated from

β-oxidation, accounting for 40.0 ± 14.5 µmol RE/min/kg-lean body mass (LBM), while RE's

derived from the TCA cycle accounted for 15.8 ± 4.8 µmol RE /min/kg-LBM, and GNG

accounted for 2.3 ± 1.4 µmol RE /min/kg-LBM **(Figure 2D)**. To determine relationships among

modeled hepatic fluxes, we performed regression analyses and obtained Pearson correlation

coefficients **(Supplemental Figure 2A)**. As expected, V_{EGP} correlated individually with $V_{Glycogen}$,

$V_{PEP,}$ and total anaplerosis, V_{PEPCK}. Moreover, expected correlations among V_{PEPCK} and pyruvate

cycling V_{PK+ME} (r = 0.99, P value < 0.0001), V_{CS} and V_{PK+ME} (r = 0.64, P value= 0.01), and V_{CS}

and V_{PEPCK} (r = 0.64, P value = 0.01) were observed **(Supplemental Figure 2B)**. Additional

correlations were observed between V_{EGP} and V_{PK+ME} (r = 0.60, P value = 0.02), between V_{EGP}

and V_{PEPCK} (r = 0.66, P value = 0.01),, and between V_{EGP} and V_{CS} (r = 0.52, P value = 0.06,

trend), **(Supplemental Figure 2C)**, as well as between V_{PEP} and V_{PK+ME} (r = 0.71, P value <

0.01), between V_{PEP} and V_{PEPCK} (r = 0.80, P value < 0.001), and between V_{PEP} and V_{CS} (r = 0.51,

P value = 0.06, *trend*) **(Supplemental Figure 2D)**. Inverse correlations were observed between

the percentage of V_{EGP} attributable to glycerol and V_{PK+ME}, $V_{PEPCK,}$ and V_{CS} **(Supplemental**

Figure 2E). No statistically significant correlations were observed between the ketogenic flux

$V_{RaβOHB}$ and any other measured flux **(Supplemental Figure 2A)**. However, as expected, total

hepatic ketogenesis (V_{RaTKB}) was highly correlated with total fat oxidation (r = 0.99, P value <

0.0001) **(Supplemental Figure 2F)**. Interestingly, total fat oxidation was not correlated with V_{CS}

(r = 0.21, P value > 0.05) **(Supplemental Figure 2G)**. Taken together, these data reflect

internal consistency in the modeling data, demonstrating expected relationships between TCA

cycle flux and GNG, as well as among TCA cycle flux, ketogenesis, and total fat oxidation in this

cohort of participants.

Given the validity of the flux modeling data collected, we next sought to determine the

relationships among these measured hepatic fluxes and MASLD-MASH progression.

Surprisingly, despite the known relationships between insulin resistance and V_{EGP} in many

populations, and the relationship between insulin resistance and NAS, V_{EGP} did not correlate

with NAS in this population (r = 0.20, P value = 0.49) **(Figure 2E).** Moreover, V_{CS}, $V_{Glycogen}$,

$V_{Glycerol}$, V_{PEP}, V_{PK+ME}, and V_{PEPCK} all showed no correlation with NAS in this population **(Figures**

2F, Supplemental Figure 3A-B). However, surrogates for the rate of ketogenesis correlated

directly with NAS [$V_{Ra\beta OHB}$, **Figure 2G** (r = 0.64, P value = 0.01); and V_{RaAcAc} (r = 0.63, P value =

0.01) and V_{RaTKB} (r = 0.65, P value = 0.01), **Supplemental Figure 3C**]**.** Thus, consistent with the

strong correlation between ketogenesis and total fat oxidation **(Supplemental Figure 2F),** both

total fat oxidation (r = 0.63, P value = 0.02) and total RE production (r = 0.62, P value = 0.02)

rates correlated directly with NAS (**Figure 2H-I**). The mitochondrial redox ratio of βOHB:AcAc

did not correlate with NAS **(Supplemental Figure 3D).** $V_{Ra\beta OHB}$ trended positive with liver PDFF

(r = 0.46, p value = 0.09) and with AIRg (r = 0.53, p value = 0.06) **(Supplemental Figure 3E-F),**

as well as with circulating transaminase levels **(Supplemental Figure 3G).** Though not included

in NAS, fibrosis is also a strong prognostic indicator of MASLD progression. However, total fat

oxidation and RE production rate only trended towards a positive correlation with fibrosis

(Figure 2J-K). Interestingly, fibrosis was inversely correlated with the rate of anaplerosis

relative to the TCA cycle (V_{PEPCK} / V_{CS}) (r = -0.66, P value = 0.01), and positively correlated with

the rate of the TCA cycle relative to EGP (V_{CS} / V_{EGP}) (r = 0.57, P value = 0.03) **(Figure 2L-M).**

These ratios provide insight into the distribution of RE's harvested from fat oxidation, and

suggest that with MASLD progression, that the sourcing of RE's to glucose production may

wane, consistent with the pathological course of the condition through MASH. Collectively,

these data demonstrate that ketogenesis and total fat oxidation increased in this population of

participants during the natural history of MASLD progression, whereas TCA cycle flux remained

relatively stable. Given (i) the relationship between ketogenesis and MASLD and (ii) the fact that

ketogenesis is a strong predictor of total fat oxidation in the liver, we next sought to determine

the mechanistic connections among these indices using mouse models.

Loss of HMGCS2 predisposes the liver to steatosis and dysregulated hepatic energy

metabolism. Our previous findings in *ex vivo* perfused livers suggested that GNG was impaired

in the setting of HMGCS2 knockdown-driven ketogenesis insufficiency.[42,49] Because hepatic

GNG is bioenergetically coupled to fat oxidation, this suggested that overall hepatic fat oxidation

might be impaired in the absence of ketogenesis. Nonetheless, ketogenesis insufficient mice

maintained euglycemia when fasted, which possibly is due to compensation from

glycogenolysis.[42,49,51] To test if sustained fat oxidation was impaired in the setting of ketogenic

insufficiency *in vivo,* we fed control and hepatocyte-specific HMGCS2-null mice a high-fat,

carbohydrate-restricted (HFCR) diet, and hypothesized that in the absence of dietary

carbohydrates, hepatic GNG would fail to support glycemia **(Figure 3A)**. We first confirmed loss

of HMGCS2 function in hepatocyte-specific HMGCS2 knock out (KO) mice[45,52] by demonstrating

that chow-fed HMGCS2-Liver-KO mice failed to mount a ketogenic response to fasting, but

maintained a normal ratio of βOHB:AcAc **(Figure 3B-C).**[53] When littermate control mice were

placed on a HFCR diet for 1 week, 4 hr fasted total ketone bodies were elevated 3.3 ± 0.2-fold,

as expected (Student's t-test, P value < 0.001); however, only βOHB was increased, and not

circulating AcAc, which was lower in the HFCR diet-fed mice **(Figure 3D).** As a result, the ratio

of βOHB:AcAc was increased 10.7 ± 0.2-fold, indicative of an elevated mitochondrial

NADH:NAD$^+$ ratio **(Figure 3E).** Moreover, unexpectedly, while circulating total ketones in

HFCR-fed HMGCS2-Liver-KO mice (662 ± 152 μM) trended lower than littermate controls (1071

± 349 μM) (Student's t-test, P value = 0.08), they were markedly elevated compared to chow-

fed HMGCS2-Liver-KO animals (66 ±5 μM) **(Figure 3B, 3F).** Compared to littermate controls

maintained on HFCR diet, loss of HMGCS2 markedly decreased the ratio of βOHB:AcAc in

HFCR diet fed mice, but increased the total amount of circulating L-βOHB by 3.8 ± 0.1-fold,

which does not contribute to mitochondrial $NADH:NAD^+$ ratio **(Figure 3G-I).** To address whether

the ketonemia in HFCR-diet fed HMGCS2-Liver-KO mice was attributable to ketogenesis or to

possible impairments in peripheral ketone body disposal, we quantified static abundances of

ketone bodies in liver tissue and found that HFCR-fed HMGCS2-Liver-KO livers showed a 69

±14% diminution in total ketone bodies (Student's t-test, P value < 0.01), and increased liver L-

βOHB, supportive of marked impairment of conventional ketogenesis in livers of HFCR-fed

HMGCS2-Liver-KO mice **(Figure 3J-K).**

To quantify the systemic physiological response to acute HFCR-feeding of HMGCS2-

Liver-KO mice, we quantified gravimetric indices and metabolite concentrations one week after

transitioning from chow. HMGCS2-Liver-KO mice lost more weight than littermate controls when

fed a HFCR diet but developed severe hepatomegaly **(Figure 4A-B).** This was accompanied by

a 3.5 ± 0.1-fold accumulation of liver TAGs (**Figure 4C-D),** which could be reversed by

refeeding mice a standard chow-diet for one week **(Figure 4E).** In both the random-fed state,

and 4 hour fasted state, HMGCS2-Liver-KO mice fed a HFCR diet for one week had elevated

blood non esterified fatty acids (NEFAs), no differences in circulating TAGs, decreased blood

glucose concentrations, and depleted hepatic glycogen stores **(Figure 4F-G).**

To test the hypothesis that the impairment of hepatic fat oxidation could be linked to

hypoglycemia in HFCR-fed HMGCS2-Liver-KO mice, we quantified hepatic glucose and

associated oxidative fluxes *in vivo* in conscious, unrestrained mice, using a previously

established GCMS-based approach[54–56]. To accomplish this, vascular catheters were placed in

the carotid artery and jugular vein of 16-week-old mice fed a standard-chow diet. Five days into

post-surgical recovery, mice were switched to HFCR diet for 2 days. Seven days post-surgery,

^2H_2O, [U-$^{13}C_3$]propionate and [6,6-$^2H_2$]glucose stable isotope tracers were intravenously infused,

allowing quantification of *in vivo* fluxes via $^2H/^{13}C$ metabolic flux analysis **(Figure 5A).**[48,55,56]

When fasted, HFCR-fed HMGCS2-Liver-KO mice had lower rates of whole-body glucose

production (V_{EGP}) (59.6 ± 8.2 μmol/min/kg body weight (BW)), compared to littermate control

mice (90.1 ± 12 μmol/min/kg BW) (Student's T-test, P value = 0.003) **(Figure 5B).** Body weight

was not different between groups (data not shown). Both littermate controls and HMGCS2-

Liver-KO mice showed very low rates of glycogenolysis (V_{PYGL}), however, as expected, HFCR-

fed HMGCS2-Liver-KO mice showed a 31 ± 8% (Student's t-test, P value < 0.01) diminution in

total gluconeogenesis (V_{Aldo}), which was linked to a reduction in sourcing of glucose from both

glycerol (V_{GK}) and PEP (V_{Enol}) **(Figure 5C).** Gluconeogenic fluxes are tightly coupled to the entry

(anaplerosis) and exit (cataplerosis) of nutrients through the TCA cycle, thus we next examined

glucose linked hepatic oxidative fluxes. While anaplerosis from propionyl-CoA (V_{PCC}) was

decreased in HMGCS2-Liver-KO mice, most anaplerotic and associated fluxes were not

different between genotypes (V_{PC}, V_{LDH}, and V_{PK+ME}). Cataplerosis (V_{PCK}) was also similar

between HMGCS2-Liver-KO and WT littermates (Student's t-test, P value > 0.05) **(Figure 5D)**.

However, HMGCS2-Liver-KO mice showed increased TCA cycle fluxes as evidenced by higher

V_{CS} (+72 ± 6%, Student's t-test, P value < 0.001) and V_{SDH} (+42 ± 6%, Student's t-test, P value

< 0.01) **(Figure 5D).** Thus, the V_{PCK}/V_{CS} ratio was decreased 50 ± 10% (Student's t-test, P value

< 0.001) in HMGCS2-Liver-KO mice, and V_{CS}/V_{EGP} ratio was increased 158 ± 8% (Student's t-

test, P value < 0.001) **(Figure 5E).** Moreover, while V_{PC} was unchanged, static concentrations of

acetyl-CoA (quantified by LC-MS/MS) were increased 77 ± 17% (Student's t-test, P value <

0.01) in livers of HFCR-fed HMGCS2-Liver-KO mice, indicating an absence of allosteric

activation of PC activity due to increased acetyl-CoA in HFCR-fed HMGCS2-Liver-KO mice

(Figure 5F). Moreover, static levels of liver succinyl-CoA and propionyl-CoA were decreased,

also suggestive of diminished relative anaplerosis **(Figure 5E-F).** The total pool of energy

adenylates (ATP, ADP, and AMP) was diminished 39 ± 14% in livers of HFCR-fed HMGCS2-

Liver-KO mice, with no impairment in calculated energy charge **(Supplemental Figure 4A-C)**.

Static liver concentrations of redox metabolites NAD^+ and NADH were also both diminished 45

$\pm$ 13% and 38 $\pm$ 14%, respectively, but no difference in static NADH:NAD+ ratio was observed

in livers of HFCR-fed HMGCS2-Liver-KO mice **(Supplemental Figure 4D-F).** Together, these

findings underscore interconnections among ketogenesis, overall fat oxidation,

anaplerosis/cataplerosis, and total EGP. Relative responses of modeled fluxes in HFCR-fed

HMGCS2-Liver-KO mice, quantified *in vivo* via intravenous infusions, are consistent with our

prior observations performed using *ex vivo* portal vein liver perfusions in chow-fed HMGCS2

knockdown mice, which revealed: (1) increased mitochondrial acetyl-CoA content, (2) elevated

V_{CS} absolute flux rate, and (3) diminished V_{PEPCK}/V_{CS} ratio.[42,49] We next sought to quantify the

contribution of ketogenesis to global hepatic fat oxidation, and how this influences the course of

MASLD pathology.

Ketogenic insufficiency impairs total hepatic fat oxidation. To determine if ketogenic

insufficiency impaired global rates of hepatic fat oxidation **(Figure 6A)**, including those

components accounted for by ketogenesis and V_{CS}, we returned to ketogenesis insufficient

models in *ex vivo* perfused livers in carbohydrate-replete settings to avoid counterregulatory

responses to hypoglycemia. Our group has previously characterized the HMGCS2 knockdown

model of ketogenic insufficiency via antisense oligonucleotide (ASO) extensively, and showed

(1) nearly complete abrogation of HMGCS2 protein in the liver, but its preservation in non-

hepatic tissues; (2) abrogation of fasting-induced ketogenesis; (3) elevation of V_{CS} and increase

in V_{CS}/V_{GNG} ratio; and (4) increased predisposition to steatotic liver injury compared to

scrambled ASO-treated controls.[42,43,49,53] Compared to livers of scrambled ASO-treated controls,

the livers of chow-fed ketogenesis insufficient mice exhibited an 80 $\pm$ 24% diminution in total fat

oxidation (Student's t-test, P value < 0.001), a 76 $\pm$ 24% impairment in RE's generated from β-

oxidation, and a strong trend towards a 39 $\pm$ 23% decrease in total RE production rate

(Student's t-test, P value = 0.056), despite a trend toward a 70 ± 23% increase in TCA cycle-

sourced RE's (Student's t-test, P value = 0.1108) **(Figure 6B-C).** There were no differences in

RE's generated through GNG. Concordant observations were observed in livers of HMGCS2

knockdown mice fed a 42% kcal from fat western diet for 8 weeks, in which V_{CS} was increased

155 ± 30% (Student's t-test, P value = 0.03), and as such RE's produced by the TCA cycle

strongly trended towards a >2-fold increase in the HMGCS2 knockdown liver (Student's t-test, P

value = 0.05), suggesting that loss of ketogenesis stimulates terminal oxidation on western diet

(Figure 6D-F). However, the data support the hypothesis that the rate of total fat oxidation is

impaired by ketogenic insufficiency.

To address underlying mechanisms for impaired total fat oxidation in the setting of

ketogenic impairment, we isolated mitochondria from control C57BL/6NJ mouse livers and

performed mitochondrial stress testing using our standard creatine kinase clamp protocol,

allowing quantification of respiratory flux (JO_2) across variations in energy demand (ΔG_{ATP})

(Figure 6G).[57] Treatment of mitochondria with hymeglusin (HG), a small molecule inhibitor of

both HMGCS enzymes (HMGCS1 is cytoplasmic enzyme and is thus not included in our

mitochondrial preparations) showed the expected diminution in the pool size of HMG-CoA (the

product of the HMGCS2 reaction), as well as AcAc and βOHB within mitochondria, coinciding

with abrogated βOHB production rates in mitochondria stimulated with octanoylcarnitine + α-

ketoglutarate **(Figure 6H).**[58,59] With increasing energy demand (decreasing absolute value of

ΔG_{ATP}), JO_2 rates increased with a diminished slope in HG-treated mitochondria, indicating an

overall respiratory impairment in the setting of ketogenic insufficiency **(Figure 6I)**. This

coincided with an accumulation of acetyl-CoA, AcAc-CoA, as well as C4-C8 acyl-CoA species,

and a depletion in free CoA **(Figure 6J-K)**. Finally, mitochondrial redox was more reduced

[NAD(P)H, % reduced] across levels of energy demand in HG-treated mitochondria **(Figure 6L).**

While this result may appear to contradict the normal redox ratio observed in livers of HFCR-fed

HMGCS2-Liver-KO mice **(Figure S4F)**, static concentrations of total redox nucleotides may not

reflect variations in cycling enzymatically-bound pools.[50,60] Taken together, these findings show

that programmed ketogenic insufficiency causes impaired total fat oxidation *in vivo*, *ex vivo*, and

in isolated mitochondria. A potential contributor to this impairment includes CoA trapping, which

impedes procession though β-oxidation and the TCA cycle, which is supported by our prior

findings in the setting of ketogenic insufficiency.[42] A second contributor to diminished fat

oxidation rates is reduced mitochondrial matrix redox potential in the setting of ketogenic

insufficiency, captured in this flux-based assay that quantifies redox nucleotide turnover among

actively cycling enzymatic pools.

Loss of mitochondrial βOHB Dehydrogenase (BDH1) impairs hepatic fat oxidation but

does not provoke liver injury in HFD-fed mice. Ketogenic insufficiency via HMGCS2 loss of

function causes steatotic liver injury.[42–46] Our novel findings here show that HMGCS2 loss of

function diminishes rates of global liver fat oxidation in a manner correlated with more reduced

mitochondrial redox potential. To directly interrogate the role of the ketogenic reaction directly

regulating mitochondrial redox potential, catalyzed by mitochondrial BDH1 (which converts

AcAc and NADH to βOHB, recovering NAD$^+$, in a near-equilibrium reaction) in (1) total hepatic

fat oxidation and (2) the response to steatotic liver injury, we studied mice in which BDH1 had

been deleted specifically in hepatocytes (BDH1-Liver-KO mice) **(Figure 7A)**. Our prior studies

demonstrated impairments of oxidative metabolism in the livers of BDH1-Liver-KO mice

maintained on a standard chow diet.[50] Here, we fed mice a 40% high-fat western diet for 18

weeks. As expected, and consistent with observations in BDH1-Liver-KO mice maintained on a

standard chow diet, circulating βOHB concentrations and total ketone bodies were modestly

diminished after an 18h fast **(Supplemental Figure 5A)**. Body weights, liver weights, circulating

glucose concentrations were all normal **(Supplemental Figure 5B-D).** Livers were then

perfused *ex vivo* via the portal vein with a physiologically relevant mixture of long chain fatty

acids (LCFAs) conjugated to bovine serum albumin (BSA). While there was no statistically

significant diminution in total ketogenesis in livers of BDH1-Liver-KO mice, all ketone bodies

produced by the liver emerged as AcAc, with no βOHB production **(Figure 7B).** Total glucose

production was diminished 26 ± 13% (Student's t-test, P value = 0.0390) in livers of BDH1-Liver-

KO mice, which coincided with a 48 ± 12% and 30 ± 13% decrease in sourcing of both PEP

(Student's t-test, P value < 0.001) and glycogen (Student's t-test, P value = 0.04) to glucose,

respectively **(Figure 7C-D).** Consistent with our previous findings in livers of BDH1-Liver-KO

mice maintained on a chow diet, the rates of pyruvate cycling (V_{PK+ME}), anaplerosis and

cataplerosis (V_{PEPCK}), and TCA cycle turnover (V_{CS}) were all suppressed 42 ± 16% (Student's t-

test, P value = 0.0043), 15 ± 13% (Student's t-test, P value = 0.0005), and 39 ± 17% (Student's

t-test, P value = 0.0160), respectively, in livers of BDH1-Liver-KO mice of HFD **(Figure 7E).**

Furthermore, total fat oxidation was decreased 26 ± 12% (Student's t-test, P value = 0.0170)

and RE production rate was decreased 20 ± 10% (Student's t-test, P value = 0.0259) in the

BDH1 KO liver **(Figure 7F-G),** demonstrating that loss of the NAD^+ regenerating step of

ketogenesis alone is sufficient to impair overall fat oxidation in the liver.

To determine whether the diminution in hepatic fatty acid (FA) oxidation in livers of

BDH1-Liver-KO mice was linked to increased hepatic steatosis or injury, we performed

histological and molecular analyses. Hematoxylin and eosin staining showed normal lipid

droplet storage, and no evidence of increased fibrosis via picrosirius red staining compared to

littermate controls **(Figure 7H).** Gene expression biomarkers of fibrosis (*Col1a1, Col3a1,*

Col4a1, Acta2) were normal, and quantities of tissue lipid peroxides (hydroxyalkenals, indices of

oxidative stress) were not increased in livers of BDH1-Liver-KO mice **(Figure 7I-J).** Maintaining

BDH1-Liver-KO mice and littermate controls on a 60% HFD, rather than a western diet, also

showed no evidence of increased steatosis, inflammation, or fibrosis – with a trend towards a 56

± 30% diminution in fibrosis in livers of BDH1-Liver-KO mice (Student's t-test, P value = 0.06)

(Supplemental Figure 6B-C). Moreover, gene expression biomarkers of fibrosis and

inflammation (*Nox, Emr1, and Tnfa*) were all diminished in livers of BDH1-Liver-KO mice

(Supplemental Figure 6C). Collectively, these findings demonstrate (1) that homeostatic

ketogenesis is required to maintain fat oxidation and its coupling to hepatic glucose production,

with loss of either HMGCS2 or BDH1 showing impairments in hepatic fat oxidation, and (2) that

the impairment in fat oxidation does not fully explain why loss of HMGCS2 predisposes the liver

to HFD-induced liver injury.

Discussion.

The natural history of MASLD-MASH proceeds through parallel derangements of both liver-

specific and systemic metabolic pathways, as well as canonical signaling cascades among

many cell types. Characteristic signatures include imbalanced lipid synthesis (DNL, and TAG

synthesis from DNL-sourced FAs and free FAs (FFAs) derived from adipose lipolysis) and

disposal (mitochondrial FA oxidation, and TAG secretion as VLDL particles). While most studies

concur that DNL is augmented during the progression of MASLD-MASH, and VLDL secretion,

while increased, does not match the rate of lipid accumulation, controversy surrounds variations

in the rate of mitochondrial fat oxidation and whether variations in this rate have a causal role.

This significantly impacts the selection of therapeutic approaches and molecular targets worthy

of pursuit. In this study, we quantified hepatic oxidative fluxes in humans with histologically

confirmed MASLD-MASH and demonstrated that the histopathological degree of liver injury (via

NAS) was positively correlated with ketogenesis and total fat oxidation, but was not correlated

with TCA cycle turnover. Use of mouse models suggested a driver role of ketogenesis in overall

hepatic fat oxidation and a causal role in preventing liver injury, as elimination of HMGCS2

showed an impairment in overall hepatic fat oxidation. This impairment of hepatic fat oxidation

correlates with increased steatotic liver injury as shown in numerous studies performed by our

group and others in the setting of ketogenic insufficiency due to deletion of HMGCS2.[42–46] Thus,

the increase in ketogenesis observed in patients with worsening MASH could be posed to serve

as a compensatory mechanism, increasing fat disposal through oxidation, in the face of

stagnating TCA cycle flux. This position is challenged, however, by the results observed in mice

lacking BDH1 in hepatocytes, whose deficiency also quantitatively diminishes overall hepatic FA

oxidation rates, but is not associated with worsened steatotic liver disease. In fact, signatures of

a modestly adaptive response to diminished hepatic FA oxidation is evident in the livers of

BDH1-Liver-KO mice. Taken together, these results suggest that while maintenance of

ketogenic sufficiency may be important to slow the progression of MASLD, the salutary

mechanism conferred by ketogenesis may not directly relate to maximizing the overall rate of

hepatic FA oxidation. Therefore, approaches aimed primarily at stimulating further hepatic fat

oxidation may not ameliorate the course of commonly-observed MASLD, and could exacerbate

liver injury.

Quantified rates of hepatic fat oxidation in humans with MASLD, relative to steatosis-

free, but weight-matched controls, have been variously reported in the literature.[22–25,30–37,47,61,62]

Much of this may relate to the populations studied, including (1) the stage of progression; (2) the

degree of adiposity and insulin resistance; (3) whether or not the participant was weight-neutral

at the time of study; (4) the genotype (*e.g.,* PNPLA3); (5) whether or not the participant was fed,

fasting, or undergoing a clamp study; and lastly, (6) the specific methodology employed. While

in vivo hepatic oxidative fluxes have been quantified in humans with uncomplicated MASLD,

and isolated mitochondrial function has been measured in liver biospecimens harvested from

participants with MASH, *in vivo* hepatic oxidative fluxes had not been quantified in humans with

histologically confirmed MASH. In well-performed studies by the Burgess and Browning group,

formally modeled ketogenesis negatively correlated with worsening hepatic steatosis (quantified

by PDFF) in overweight and obese participants with BMI ≤ 35.[22,37] Moreover, circulating ketones

and expression levels of ketogenic enzymatic machinery decreased within worsening NAS in a

bariatric surgery cohort studied by the Rector group, but no study has tested whether

endogenous ketogenesis correlates with histopathologically-confirmed liver injury.[33,34]

Consistent with our findings, a recent study in non-obese participants with MASLD by the

Petersen and Shulman group showed that participants with compounding cardiometabolic

disturbances have increased rates of βOHB turnover in comparison to participants with simple

steatosis alone, or with healthy livers, but liver biopsies were not available in that study.[27] The

cohort studied in our experiments reflects a patient population with advanced obesity at

elevated risk of progression from MASH to cirrhosis, and among the metabolic fluxes modeled,

the only direct correlate of NAS was endogenous ketogenesis ($V_{Ra\beta OHB}$). We propose that the

escalation of TCA cycle flux early in the progression of MASLD may become attenuated with

further progression, and indeed, the two participants we studied with the greatest V_{CS} magnitude

(>5 µmol/min/kg-LBM) had NAS 4, and those exhibiting a NAS > 4 did not exceed 5

µmol/min/kg-LBM. Thus with respect to variation of TCA cycle flux, our data are generally

consistent with those of the Burgess and Browning group.[22,25] Our data in both humans and

mouse models suggest that ketogenesis may serve an adaptive counterregulatory role,

especially with further MASH progression.

Our findings suggest that impairments in fat oxidation do not play a casual role in linking

impaired ketogenesis to liver injury, implying that the link between ketogenesis and MASLD-

MASH progression involves attributes of ketogenesis beyond fat oxidation. Concordance

between our observations in modeled hepatic fluxes of the human participants with MASH and

mouse models of ketogenic insufficiency strongly substantiate this claim. In addition to the direct

association between ketogenesis and human NAS, increasing histopathological evidence of

fibrosis was inversely correlated with V_{PEPCK}/V_{CS} ratio and directly with V_{CS}/V_{EGP} ratio **(Figure**

2L-M), suggestive of decoupling between TCA cycle flux and sourcing of its derived RE's to

GNG with increasing liver injury. Similarly, V_{PCK}/V_{CS} ratio was decreased in steatosis and injury-

prone HMGCS2-Liver-KO mice, and V_{CS}/V_{EGP} ratio was increased **(Figure 5E)**. Conversely, this

imbalance was not observed in livers of BDH1-Liver-KO mice, whose V_{PEPCK}, V_{EGP}, and V_{CS}

were all coordinately diminished **(Figure 7E).** Alternatively, and not mutually exclusively,

mechanisms linked to the increase in AcAc production in livers of BDH1-Liver-KO mice could

explain why the loss of BDH1 protects the liver from injury.[43]

Recently, we reported a novel role for ketogenesis in supporting FA elongation, PUFA

biosynthesis and phospholipid metabolism in the liver.[45] We demonstrated that (1) ketogenesis

furnishes hepatocytes with recycled acetyl-CoA that can be used for FA biosynthesis, via both

the DNL and FA elongation pathways, and (2) that incorporation of ketone body-derived carbon

into FA elongation proceeds solely via the cytosolic enzyme acetoacetyl-CoA synthetase

(AACS), through a metabolic pathway independent of ketone body sourcing of DNL.

Furthermore, we demonstrated that loss of HMGCS2 impairs FA elongation and diminishes

hepatic PUFAs, specifically those esterified in phosphatidylethanolamine (PE) pools, suggesting

the ketogenesis could influence mitochondrial function, and VLDL assembly and secretion

through PUFA-dependent mechanisms, though these possibilities are untested. While it is

tempting to speculate that this pathway linking ketogenesis to PUFA biosynthesis explains the

protective nature of ketogenesis in MASLD-MASH progression, the observation that AcAc

production is increased in the absence of BDH1 suggests that FA elongation and PUFA

biosynthesis could be stimulated in a substrate-driven manner. PUFAs inhibit DNL in the liver

via inhibiting SREBP1c mediated gene transcription,[63–68] and therefore the rise in ketone body-

sourced PUFAs in the BDH1 KO could drive both acute and chronic inhibition of DNL, while

simultaneously stimulating VLDL secretion. This hypothesis is further supported by the

observation that loss of HMGCS2, which abrogates production of all ketones and exhibits

diminished liver PUFAs, induces a DNL gene expression profile.[42,45]

Collectively, these findings in humans with MASH and in loss-of-ketogenesis function

mouse models suggest that ketogenesis may compensate in a fat-overloaded, progressively

injured liver, thereby maintaining rates of overall fat oxidation, but perhaps more importantly, by

burning fat through a metabolic pathway that confers additional benefits that terminal oxidation

in the TCA cycle may not bestow during MASLD progression. The specific regulatory

mechanisms, whether deleterious or salutary, that coordinate the fate of β-oxidation-derived

acetyl-CoA through the TCA cycle or ketogenesis, remain incompletely understood. Classical

mechanisms include oxaloacetate availability, allosteric regulation of citrate synthase, NADH

redox potential, and ΔG_{ATP}. Physiological variation in the abundance of HMGCS2 protein could

serve as a regulatory mechanism, but this has not been definitively proven. Irrespective, the

myriad pathways through which endogenous ketogenesis and exogenous ketone bodies could

influence DNL, PUFA synthesis, VLDL secretion, fat oxidation, and signaling through

inflammatory and immune pathways, opens many opportunities to understand the range of

therapeutic opportunities for metabolic and signaling roles to ameliorate MASLD.

Limitations. The human studies are limited by a small sample size (14 participants), and the

inability to study sex as a biological variable (most recruited participants were female). However,

this is the first population in which isotope tracers have been used to model oxidative fluxes in

participants with a BMI ≥ 35 and histopathologically graded NAS. Furthermore, while all

participants were insulin resistant, only one participant had diabetes, so it is not possible to

make conclusions relevant in the scope of MASLD within diabetes. Two independent

experimental approaches were leveraged to delete HMGCS2 protein in mice, but the

concordance in phenotypes across these models supports the rigor of our findings. Using

metabolic flux modeling, we demonstrated that disruption of ketogenesis impaired overall

hepatic fat oxidation, which was quantified primarily via an indirect flux modeling-based assay

predicated on the assumption that mitochondrial acetyl-CoA turnover at steady-state is

mathematically equivalent to total ketogenesis (V_{RaTKB}) + TCA cycle turnover (V_{CS}) +

Acetogenesis ($V_{Acetate}$). A limitation of this approach is the indirect measurement of fat oxidation,

the assumption that fat oxidation is primarily sourced from exogenous fat sources during portal

vein perfusions, and the assumption that octanoate sources little acetate. The findings

generated using this methodology, however, are reinforced by the concordance among all the

fluxes modeled across all orthogonal experimental methodologies employed here. Collectively,

an impairment in fat oxidation was consistency observed in the setting of ketogenic

insufficiency, as shown across various models including: (1) *in vivo* fluxes in HFCR-diet-fed

HMGCS2-Liver-KO mice; (2) *ex vivo* portal vein octanoate-perfused livers from HMGCS2 ASO-

knockdown mice maintained on either chow or western diet; (3) *ex vivo* portal vein octanoate-

perfused livers from BDH1-Liver-KO mice maintained on chow diet; (4) *ex vivo* portal vein

LCFA-perfused livers from BDH1-Liver-KO maintained on Western diet; and (5) *ex vivo* isolated

mitochondria treated with HG, which inhibits HMGCS2 and simultaneously impaired JO$_2$ linked

to fat oxidation. Collectively, these data strongly support the hypothesis that ketogenesis plays

an integral role in regulating fat oxidation. A modest limitation of *in vivo* ketogenesis

measurements in humans was that ketogenesis was modeled as a single pool, with V_{RaAcAc}

calculated from $V_{Ra\beta OHB}$ and the ratio of βOHB:AcAc. Future experiments will infuse independent

[^{13}C]AcAc and [^{13}C]βOHB tracers to directly measure V_{RaAcAc}, and interconversion of AcAc and

βOHB through BDH1 via a two-pool model. Finally, though male and female mice were used in

these studies, as delineated in the figure legends, sex as a biological variable was not

specifically considered in these studies.

Materials and Methods

Human MASLD-MASH recruitment and study design **(Figure 1 and Supplemental Figure 1).**

Patients with abnormal liver function, prediabetes or type 2 diabetes were screened to identify

participants who meet eligibility requirements for two clinical studies (ClinicalTrials.gov,

NCT03997422 and NCT03587831). The full list of initial inclusion and exclusion criteria are

provided in **Supplemental Table 1.** Fourteen participants with a BMI ≥ 35 and a PDFF > 5%,

quantified by MRI were enrolled and were not given instruction or counseling to lose weight.

Insulin sensitivity and body composition were assessed via a FSIVGTT and DXA imaging,

respectively, followed by *in vivo* hepatic oxidative flux measurements. After enrollment, and prior

to flux measurements, a liver biopsy was obtained by an interventional radiologist under direct

image guidance with ultrasound-guided approach with percutaneous puncture using a 16-gauge

Tru-cut needle with standard protocols in place to detect complications. Histological severity

was scored independently by two blinded pathologists based on hematoxylin and eosin-stained

sections, assessing for macrovesicular steatosis (0-3), lobular inflammation (0-3), and acinar

zone 3 hepatocellular injury, or ballooning degeneration (0-2), which were summed to express

the NAFLD activity score, NAS. In addition to NAS, fibrosis (0-4) was also quantified using

picrosirius red staining.

PDFF Measurements. Each participant underwent a 20-minute MRI scan to measure liver fat

content non-invasively. MRI scans were performed on a 3T Prisma MR system (Siemens

Medical, Erlangen, Germany) at the UMN Center for Clinical Imaging Research. The PDFF

image (MRI-PDFF) was acquired using a 3D GRE with 6 TE values (TR=9ms,flip angle=4,

TE=1.15, 2.46, 3.69, 4.92, 6.15, 7.38 ms, 2x2x3.5mm resolution, one 17s breath-hold), and

reconstructed using the vendor's software to produce a PDFF map.

Assessment of Glucose Homeostasis via FSIVGTT. For three days prior to admission to the

clinical research unit (CRU), participants consumed a weight-maintaining diet formulated to

approximate the previous 3-day dietary record. An evening meal consisting of 40% daily caloric

intake and 40-50% carbohydrate and 30-40% fat was consumed before the 0700 CRU

admission for FSIVGTT. Two intravenous catheters were placed, one for sampling and the other

for administration of dextrose and insulin. Glucose, insulin and C-peptide were quantified at

baseline and multiple times over the course of 180 minutes. HOMA-IR was calculated using the

equation, HOMA-IR = fasting insulin (µU/ml) × fasting glucose (mmol/l) / 22.5 levels. At time 0,

dextrose 0.3 grams/kg BW was given intravenously. Insulin, 0.03 units/kg BW was then injected

at 20 minutes to augment glucose disposal. Insulin and C-peptide levels were measured for 10

minutes and the area under the curve was calculated as a measure of insulin secretory function,

termed AIRg. A minimal model assessment was done to calculate total body insulin resistance

(Si), then the Disposition Index (DI) was calculated as the product of AIRg x Si. For FSIVGTT

modeling, glucose was measured via UV test enzymatic reference method with hexokinase,

insulin was measured via electrochemiluminescence immunoassay (ECLIA), and C-peptide was

measured via chemiluminescent immunoassay.

Body Composition Measurements via DXA Imaging. Body composition was measured by a DXA

scanner, the GE Healthcare Lunar i (GE Healthcare Lunar with enCORE software version 16.2).

Stable isotope delivery in humans. After FSIVGTT and DXA scan, participants remained in the

CRU and received a low-carbohydrate lunch meal. Participants then began fasting and did not

receive an evening meal. Stable isotope intake began at 2200 hours. Participants were provided

with three equal doses of 70% deuterated water (^2H_2O, D_2O, 5 g/kg body water, calculated as

60% of BW in men and 50% of BW in women) every 4 hours between 2200 and 0600. These

doses, along with 0.5% ^2H_2O available for ad libitum consumption, sufficiently labeled body

water for measuring flux parameters. From 0800 to 0900 on day 2, three doses of [U-

$^{13}C_3$]propionate (200 mg/dose) were administered orally to label TCA cycle intermediates for

profiling of hepatic mitochondrial fluxes. At 0900, intravenous stable isotope infusion began with

[3,4-$^{13}C_2$]glucose [0.563 mg/kg BW bolus, immediately followed by a two-hour infusion (0.0563

mg/min/kg BW)] and [U-$^{13}C_4$]βOHB [0.1 mg/kg BW bolus, immediately followed by two-hour

infusion (0.01 mg/min/kg BW)]. At the end of the 2h infusion, ~50 mL of blood was drawn,

centrifuged to isolate serum, then aliquots were frozen at -80°C until further analysis.

Hepatic Flux Modeling Sample Preparation and ^{2}H/^{13}C NMR Data Acquisition. The positional

^{13}C and ^{2}H isotopomers of plasma glucose encodes quantitative information about hepatic

biochemical pathways, thus allowing for noninvasive modeling of hepatic metabolism *in vivo.*

Direct quantification of the positional isotopomer populations of plasma glucose was achieved

using ^{13}C and ^{2}H NMR, post derivatization into 1,2-isopropylidene glucofuranose (monoacetone

glucose [MAG]) as previously described.[25,36,48,49] MAG was reconstituted in 510 µL acetonitrile

spiked with 5% ^{2}H-depleted H_2O, then was transferred to a 5 mm 528-pp Wilmad NMR tube

(Vineland, NJ) and analyzed on a Bruker Avance III 600 NMR instrument equipped with a

CryoProbe. During ^{2}H NMR data collection, the magnet was not maintained on a field-frequency

lock, as the lock channel is used to acquire ^{2}H signal. Data was collected at 48°C over 1380 Hz

using a 68 µs (90°) pulse, 1.3s delay between transients and Waltz16 decoupling on the proton

channel during a 1s acquisition period. Depending on signal-to-noise (S/N), 4,000-40,000

transients were collected. Magnet stability was sufficient to result in deuterium line-widths of 2-3

Hz, allowing for integration to establish relative ^{2}H enrichments at the number 2, 5 and 6

positions of ^{2}H-labeled MAG, reflecting the relative ^{2}H enrichments of plasma glucose. For ^{13}C

data collection, 40 µL of d_3-acetonitrile was spiked into each sample used to collect the ^{2}H NMR

spectra, then a field-frequency lock was established. For data acquisition, a 12 µs excitation

pulse with 22,320 Hz collection window and waltz16 decoupling for ^{1}H was used. Depending on

S/N, 4,000 to 40,000 transients were collected, then signal magnitudes were established by

integration. The ^{13}C isotopomers that appear in glucose provide a readout on oxidative

metabolism in the liver, which are encoded in the multiplet pattern arising at the C2 resonance

of MAG, centered at ~85 ppm. The relative populations of the D12 doublet, D23 doublet and

Q123 quartet that arise when carbons 1 and 2, carbons 2 and 3, or carbons 1, 2 and 3 are

labeled, respectively, then are used to mathematically model TCA cycle turnover, anaplerosis,

cataplerosis and pyruvate cycle as described below.

Quantifying Whole-body βOHB Turnover. βOHB turnover was calculated at metabolic and

isotopic steady-state from the dilution of intravenously infused D-[U-$^{13}C_4$] βOHB. Briefly,

unlabeled and ^{13}C βOHB were quantified using UHPLC-MS/MS, then the fractional ^{13}C

enrichment of plasma βOHB and tracer infusion rate were used to calculate whole-body

turnover rates of βOHB using the equation:

$$\textit{Whole Body βOHB Turnover (μmol/min/kg BW)} = \frac{F}{\left(\frac{TTR}{TTR+1}\right)} - F$$

Where F is the tracer infusion rate and TTR is the Tracer : Tracee ratio ([$^{13}C_4$] βOHB / [^{12}C]

βOHB). After flux measurements, turnover rates are corrected for LBM from DXA imaging, and

expressed as μmol βOHB turned over/min/kg-LBM. Because the liver is the primarily site of

systemic ketone body production, whole-body turnover rates of ketone bodies at steady-state

are mathematically equivalent to hepatic endogenous βOHB production rates **($V_{RaβOHB}$).**

Quantifying Whole-body Glucose Turnover. Glucose turnover was calculated at metabolic and

isotopic-steady state from the dilution of intravenously infused [3,4-$^{13}C_2$] glucose. The fractional

enrichment of plasma glucose was determined from the ^{13}C NMR spectra of MAG, by

measuring the ratio (R) of enrichment at C3 and C4, relative to the natural abundance of the

methyl groups from the acetone moiety of the MAG derivative. Fractional ^{13}C enrichment (L_p)

was then calculated using linear regression, from a standard curve of known glucose

enrichments and the equation $L_p = 1.33$ (R) + 0.031. Assuming that the fractional ^{13}C

enrichment of infused [3,4-$^{13}C_2$] glucose (L_i) was 99%, and the known tracer infusion rate (F),

then whole body glucose turnover at metabolic and isotopic steady state could be calculated as:

$$\textit{Whole Body Glucose Turnover (mg glucose/min/kg BW)} = F \ x \ \left(\frac{Li-Lp}{Lp}\right)$$

Using the DXA scan information, final flux units were expressed as µmoles glucose turned

over/min/kg-LBM. In the fasted state the majority (>90%) of whole-body glucose production is

derived from the liver, therefore total glucose turnover rates are mathematically equivalent to

total hepatic EGP rates (V_{EGP}).

Quantifying Hepatic Glucose Sourcing Fluxes. Using the ^2H enrichment patterns encoded in

plasma glucose, the fractional contribution of various glucose sourcing pathways can be

calculated based on the following equations:

Fraction of EGP from Glycogen $= \frac{H2-H5}{H2}$

Fraction of EGP from Glycerol $= \frac{H5-H6s}{H2}$

Fraction of EGP from PEP $= \frac{H6s}{H2}$

Where H2, H5 and H6s correspond to the relative ^2H enrichment at the 2, 5 and 6s carbons of

MAG. By multiplying each fraction by total EGP (V_{EGP}), flux through each hepatic glucose

sourcing pathway can be calculated as follows:

Glycogenolysis ($V_{Glycogen}$) (µmol glucose/min kg LBM) $= V_{EGP} \times (\frac{H2-H5}{H2})$

Gluconeogenesis from Glycerol ($V_{Glycerol}$) (µmol glucose/min/kg LBM) $= V_{EGP} \times \left(\frac{H5-H6s}{H2}\right)$

Gluconeogenesis from PEP (V_{PEP}) (µmol glucose/min/kg LBM) $= V_{EGP} \times (\frac{H6s}{H2})$

Quantifying Hepatic Oxidative Fluxes. The rate of hepatic TCA cycle turnover, anaplerosis,

cataplerosis and pyruvate cycling, can be modeled in the liver using the ^{13}C isotopomers that

arise in plasma glucose from orally administered [U-$^{13}C_3$] propionate. Briefly, at metabolic

steady-state, glucose-linked rates of anaplerosis must equal cataplerosis, which are modeled

relative to the turnover rate of the hepatic TCA cycle, assumed to be equal to the rate of CS

(V_{CS}). Because the non-oxidative entry (anaplerosis) and exit (cataplerosis) of carbon through

the TCA cycle supports glucose anabolism via PEP-sourced GNG, changes in oxidative fluxes

distort the ^{13}C enrichment patterns of PEP which are transmitted and encoded in the multiplet

resonances of plasma glucose. Using the known rate of PEP-derived GNG (V_{PEP}), quantified as

described above, and the ^{13}C isotopomer populations quantified via ^{13}C NMR, the absolute rate

of the TCA cycle turnover (V_{CS}), as well as anaplerotic and cataplerotic fluxes (V_{PEPCK}), and label

recycling rates (V_{PK+ME}) can be calculated as follows:

Pyruvate Cycling (V_{PK+ME}) (μmol/min/kg LBM) = V_{PEP} x $\left(\frac{D12-Q}{Q-D23}\right)$

Anaplerosis (V_{PC}) and Cataplerosis (V_{PEPCK}) (μmol/min/kg LBM) = V_{PEP} x $\left(\frac{D12-D23}{Q-D23}\right)$

TCA Cycle Turnover (V_{TCA}) (μmol/mi /kg LBM) = V_{PEP} x $\left(\frac{D23}{Q-D23}\right)$

where D12 represents the doublet arising when C1 and C2 of glucose labeled, D23 represents

the doublet arising when C2 and C3 are labeled, and Q is the quartet arising when C1, C2 and

C3 are all labeled. Pyruvate cycling (V_{PK+ME}) reflects the summed rate of pyruvate kinase (PK)

and malic enzyme (ME). Total anaplerosis is modeled as flux of PC (V_{PC}), while cataplerosis is

modeled as flux through PEP carboxykinase (PEPCK) (V_{PEPCK}), which at steady-state are equal

to one another ($V_{PC} = V_{PEPCK}$).

Quantifying Hepatic Fat Oxidation Rates. By assuming that the liver is in a metabolic steady-

state, the rate of acetyl-CoA production via β-oxidation can be assumed to be equal to the rate

of acetyl-CoA disposal, thereby allowing total fat oxidation rates to be indirectly quantified as the

sum of all disposal routes for hepatic acetyl-CoA. In the fasted liver, β-oxidation derived acetyl-

CoA is (1) principally converted to ketone bodies which are largely released from the liver, (2)

terminally oxidized in the TCA cycle, or (3) secreted as acetate. By quantifying each disposal

pathway for acetyl-CoA, the majority of acetyl-CoA disposal can be accounted for, and used as

a surrogate proxy of total acetyl-CoA turnover and fat oxidation at steady-state by the equation:

$$\textit{Total Fat Oxidation (µmol/min/kg LBM)} = 2 \times (V_{Ra\beta OHB} + V_{RaAcAc}) + V_{TCA} + V_{Acetate}$$

Fat oxidation is expressed in µmoles of acetyl-CoA turned over/min/kg-LBM. $V_{Ra\beta OHB}$ was

measured via a tracer dilution of intravenously infused $[U-^{13}C_4]\beta OHB$. V_{RaAcAc} was calculated

from $V_{Ra\beta OHB}$ turnover rates and the ratio of βOHB:AcAc. Acetate production rates ($V_{Acetate}$) was

assumed to be negligible in humans.

Quantifying Hepatic RE Production. The production of RE's, principally NADH and $FADH_2$, can

be estimated from the above fluxes as previously described.[35,69] This approach assumes that all

acetyl-CoA originates from palmitate, and that the sourcing of V_{PEP} reflects 90% lactate and

10% pyruvate/alanine, in accordance with the typical hepatocellular redox state. The model

assumes that complete β-oxidation of palmitate, a 16-carbon saturated FA which yields a total

of 8 acetyl-CoA units, requiring 7 rounds of the β-oxidation cascade which collectively yields 14

RE's, or an average of 1.75 RE's per molecule of acetyl-CoA produced (14 RE / 8 Ac-CoA =

1.75 RE / Ac-CoA). By quantifying the rate for each disposal pathway for acetyl-CoA, the

turnover of RE's generated from fat oxidation can be calculated from the stoichiometry of each

pathway as follows. For every mole of AcAc produced, two moles of acetyl-CoA turned over,

and thus total RE's generated from β-oxidation that are accounted for by AcAc **(RE_{AcAc})** = 3.5 x

V_{RaAcAc}. For every mole of βOHB produced, two moles of acetyl-CoA turned over, but conversion

of AcAc to βOHB consumes 1 RE, and thus net RE's generated from β-oxidation that are

accounted for by βOHB **($RE_{\beta OHB}$)** = 2.5 x $V_{Ra\beta OHB}$. For every turn of the TCA cycle, one molecule

of acetyl-CoA is consumed by CS (V_{CS}), accounting for 1.75 RE's, but each turn also generates

an additional 4 RE's, and thus total RE's accounted for by the TCA cycle **($RE_{TCA\text{-}total}$)** = 5.75 x

V_{CS}. $RE_{TCA\text{-}total}$ can be broken down into (1) total RE's generated from β-oxidation **($RE_{\beta ox\text{-}CS}$)** =

1.75 x V_{CS}, and (2) total RE's generated exclusively from the dehydrogenases of the TCA cycle

(not β-oxidation) **(RE$_{TCA\text{-}only}$)** = 4 x V$_{CS.}$ Total RE's generated from β-oxidation can then be

estimated as the sum of RE's accounted for by AcAc, βOHB and the TCA cycle

dehydrogenases as follows:

RE's generated from β-oxidation **(RE$_{βox}$)** = RE$_{AcAc}$ + RE$_{βOHB}$ + RE$_{βox\text{-}CS}$

Total RE balance in the liver requires further accounting for GNG and three enzymes in

particular: glycerol-3-phosphate dehydrogenase (G3PDH), glyceraldehyde-3-phosphate

dehydrogenase (GAPDH), and lactate dehydrogenase (LDH). G3PDH generates 1 RE during

GNG from glycerol. GAPDH consumes 1 RE during GNG from PEP, however, if the substrate

used to source PEP is lactate, then LDH will generate 1 RE to offset this. Assuming that GNG is

sourced 90% from lactate, and 10% from pyruvate + alanine, then total RE's generated from

GNG **(RE$_{GNG}$)** = V$_{Glycerol}$ − (0.1 x V$_{PEP}$). From this, total RE's turner over in the liver can then be

calculated at steady-state as the sum of RE's generated from β-oxidation, TCA cycle flux and

GNG:

Total RE production rate (μmol RE/min/kg-LBM) = RE$_{βOX}$ + RE$_{TCA\text{-}only}$ + RE$_{GNG}$

Animal Models and Diets. All animal experiments were approved by the Institutional Animal

Care and Use Committee at the University of Minnesota. All mice were bred on a C57BL/6NJ

background, and received autoclaved water ad libitum. Mice were housed on corncob bedding

in groups of 4 to 5 with lights off between 2000 and 0600 in a room maintained at 22°C. Sex of

mice studied are indicated within each figure legend. Hepatocyte-specific HMGCS2 null mice

were generated as described previously by crossing homozygous Hmgcs2$^{flox/flox}$ mice, with

heterozygous Cre recombinase expressing mice, driven by the albumin promoter[70]. Littermate

wild-type (WT) Alb-Cre-negative (*Hmgcs2$^{flox/flox}$*, Alb-Cre$^{-/-}$) animals were used as controls for

Cre-positive (*Hmgcs2$^{flox/flox}$*, Alb-Cre$^{+/-}$) HMGCS2 KO mice, which are maintained on a

C57BL6/NJ substrain hybrid background. Hmgcs2 ASO-mediated loss of function was initiated

in 6-week-old mice by i.p. injection (25 mg/kg) with murine *Hmgcs2*-targeted ASOs or

scrambled sequence control ASOs (Ionis Pharmaceuticals) biweekly as described previously[42].

Hepatocyte-specific BDH1 null mice were generated as previously described[50], by crossing

homozygous Bdh1$^{flox/flox}$ mice[71] to heterozygous Cre recombinase expressing mice, driven by

the albumin promoter[70]. Littermate WT Cre-negative (Bdh1$^{flox/flox}$, Cre$^{-/-}$) animals were used as

controls for Cre-positive (Bdh1$^{flox/flox}$, Cre$^{+/-}$) BDH1 KO mice. Food and water were provided ad

libitum to mice in temperature- and humidity-controlled conditions maintained on a 14:10 h

light/dark cycle. Mice were fed either a standard chow diet, or a high-fat diet. Standard chow

diet (Teklad Global, 2016) consisted of 12% kcal from fat, 66% kcal from carbohydrates and

22% kcal from protein. Western diet (Envigo, TD.88137) consisted of 42% kcal from fat, 43%

kcal from carbohydrates and 15% kcal from protein. 60% high-fat diet (Research Diets, D12492)

consisted of 60% kcal from fat, 20% kcal from carbohydrates, and 20% kcal from protein. HFCR

diet (Envigo, TD.160153.PWD) consisted of 90.5% kcal from fat, 0% kcal from carbohydrates

and 9.5% kcal from protein.

Quantifying Liver TAGs. 20 mg frozen liver tissue was homogenized in 1 mL of 0.1x PBS, then

total protein was quantified via BCA assay. Total lipids were extracted from 0.5 mg of protein via

a Bligh-dyer lipid extraction, then total lipids were quantified colorimetrically using Infinity TAG

reagent (Thermo, TR22421).

Quantifying Liver Glycogen Content. Total glycogen was assayed as described previously.[49]

Briefly, 100 mg of frozen liver tissue was digested in 300 μL of 30% KOH (w/v), then glycogen

was precipitated followed by enzymatic digestion using 0.3 mg/mL amyloglucosidase in 0.2 M

sodium acetate buffer at pH 4.8 (Sigma 10115, 70 U/mg). Total glucose was quantified using ^{1}H

NMR then total glycogen was expressed in glucose units per mg of liver tissue.

Serum NEFA, TAG and Blood Glucose Measurements. Serum NEFAs (Wako, 633-52001) and

TAGs (Thermo, TR22421) were measure colorimetrically using commercial kits. Blood glucose

was measured using glucose meters (Metene).

Quantification of Total Ketone Bodies via UHPLC-MS/MS. Ketone bodies were formally

quantified using UHPLC-MS/MS as described previously[50,72]. Briefly, [U-^{13}C$_4$]AcAc and [3,4,4,4-

D$_4$]βOHB internal standards were spiked into ice cold MeOH:ACN (1:1), then ketones were

extracted from serum, perfusate, plasma, or sample extracts, followed by separation via reverse-

phase UHPLC, and detected via parallel reaction monitoring (PRM) on a QExactive Plus hybrid

quadrupole-orbitrap mass spectrometer. Quantification of ^{13}C labeled ketone bodies was

achieved post reduction with NaBD$_4$ as previously described.[45,72]

Quantification of Glucose and βOHB via ^{1}H NMR. 1.8 mL of liver effluent were collected, snap

frozen and stored at -80°C till analysis. To prepare samples, the solvent was removed at 45°C

under vacuum, then samples were reconstituted in 550 µl of ^{2}H$_2$O (99.9%) spiked with 0.3 mM

d8-trimethyl-silyl propionate (TSP). ^{1}H-NMR signals were acquired using a Bruker Avance III 600

NMR instrument equipped with a CryoProbe, then the integrated intensities of the α-anomeric

proton on glucose carbon-1, the methyl signal for βOHB, and the tri-methyl signal from TSP, were

used to calculate molar concentrations of the respective substrates. For all ^{1}H-NMR collections,

spectra were collected by conventional pulse-and-collect measurements under quantitative

conditions (10-ppm spectral range using ~15 µs [90°] excitation pulse and 22-second delay

between each of 20 transients).

Stable Isotope Infusions in Mice. Jugular vein and carotid artery catheters were implanted into

mice for isotope infusion and sampling as previously described.[56,73,74] The exteriorized ends of

the implanted catheters were flushed with 200 U·mL^{-1} heparinized saline and sealed with

stainless-steel plugs. After surgery, mice were housed individually for ~7 days of postoperative

recovery before stable isotope infusions studies. All mice were within 10% of presurgical weight

before stable isotope infusions. Mice were fasted during the duration of the infusions, and mice

were fasted a total of 7 hours when samples were collected. Two hours into the fast, the

exteriorized catheters were connection to infusion syringes. After a 1-hour acclimation period,

an arterial blood sample (80 µL) was collected to determine glucose natural abundance

enrichment, followed by a stable isotope infusion protocol to measure glucose-linked hepatic

oxidative fluxes as previously described.[55,56] Briefly, a ^2H_2O (99.9%)-saline bolus containing

[6,6-$^2H_2$] glucose (99%) was infused over 25 min to enrich body water and prime (440

µmol·kg^{-1}) the glucose pool. This was followed by a continuous infusion of [6,6-$^2H_2$] glucose (4.4

µmol·kg^{-1}·min^{-1}) till the end of the experiment. A primed (1.1 mmol·kg^{-1}), continuous (0.055

mmol·kg^{-1}·min^{-1}) intravenous infusion of [U-^{13}C] propionate was started 2 h after the ^2H_2O

bolus. 100 µL arterial samples were collected at 90, 100, 110, and 120 mins after the [U-$^{13}C_3$]

propionate bolus to determine arterial glucose concentration and enrichment. Donor

erythrocytes were provided by constant rate infusion throughout the experiment to prevent a

decline in hematocrit. Mice were euthanized by cervical dislocation following the final sample

collection, then tissues were rapidly excised, freeze-clamped in liquid nitrogen. All samples were

stored at −80°C.

Glucose Derivatization, GC-MS Analysis and in vivo $^2H/^{13}C$ Metabolic Flux Analysis in conscious

unrestrained Mice. Approximately 40 µL of plasma at each time point was used to synthesize di-

O-isopropylidene propionate, aldonitrile pentapropionate, and methyloxime pentapropionate

derivatives of glucose, then GC-MS analysis was utilized to collect uncorrected mass

isotopomer distributions (MIDs) for six fragment ions as previously described.[54,55,75] The

metabolic flux analysis methodology used in these studies followed that previously described

with minor modifications.[54] To summarize, a metabolic reaction network was constructed using

Isotopomer Network Compartmental Analysis (INCA) software.[76] The reaction network defined

the carbon and hydrogen atom transitions for hepatic glucose and associated oxidative

reactions. The reaction velocities (i.e., flux) through each network reaction was determined

relative to CS flux (V_{CS}) by minimizing the sum of squared residuals between simulated and

experimentally determined MIDs of the six fragment ions described above. Flux estimates were

repeated 50 times from random initial values. Goodness of fit was assessed by a Chi-square

test (P = 0.05) and confidence intervals of 95% were determined as previously described.[56]

Mouse body weights and the [6,6-^{2}H$_2$] glucose infusion rate were used to determine absolute

values. In the GCMS-based model, total EGP (V_{EGP}) is equal to the sum of glycogenolysis

(V_{PYGL}) and GNG (V_{Aldo}), which are modeled as the rates of glycogen phosphorylase (PYGL)

and aldolase (Aldo), respectively. Sourcing of GNG from glycerol (V_{GK}) and PEP (V_{Enol}) are

modeled as the rates of glycerol kinase (GK) and enolase (Enol), respectively. Pyruvate cycling

(V_{PK+ME}) is equal to the summed rate of PK and ME and total cataplerosis is still equal to flux

through PEPCK (V_{PCK}). Anaplerosis is equal to the sum of PC (V_{PC}) and propionyl-CoA

carbobxylase (PCC) (V_{PCC}), and accounts for entry of unlabeled substrates through LDH (V_{LDH}).

Turnover of the 1st span of the TCA cycle turnover is modeled as the rate of CS (V_{CS}), whereas

2nd span TCA turnover, modeled as the rate of succinate dehydrogenase (SDH) (V_{SDH}) reflects

the sum of V_{CS} and V_{PCC}.

Mouse Liver Histology. Following euthanasia, mouse liver sections were fixed in 10% neutral

buffered formalin, then tissue embedding, sectioning, and staining with hematoxylin and eosin

was performed by the University of Minnesota Clinical and Translational Science Institute

histology core.

Quantifying Energy Charge, Redox State and Short-chain Acyl-CoAs. Energy nucleotides (ATP,

ADP and AMP), redox nucleotides (NAD$^+$ and NADH) and high-energy thioester containing acyl-

CoAs (acetyl-CoA, propionyl-CoA and succinyl-CoA) were measured as previously

described[50,77], with modifications. Briefly, metabolites were extracted, separated using ion-

pairing reverse-phase UHPLC-MS/MS on a C$_{18}$ column (Waters Xbridge, 150 x 2.1mm, 3μm),

then detected on a Thermo QExactive Plus mass spectrometer, operated in positive ionization

mode, using PRM transitions as previously described[50,77]. Relative pool size (ion counts)

normalized to biomass (mg total protein) was quantified for all analytes. Nucleotides were

detected as adducts of dibutylamine acetate. Energy charge and Redox State can be estimated

in the liver using the pool size of nucleotides and the equations:

$$Energy\ Charge\ = \frac{([ATP]+0.5[ADP])}{Total\ Energy\ Nucletodies}$$

$$Total\ Energy\ Nucleotides = [ATP] + [ADP] + [AMP]$$

$$Redox\ State\ = \frac{[NAD+]}{[NADH]}$$

$$Total\ Redox\ Nucleotides = [NAD+]\ +\ [NADH]$$

Ex vivo Mouse Liver Perfusions. Liver perfusions were performed as described previously[49].

Briefly, mice were anesthetized with i.p. sodium pentobarbital injection, externally sterilized with

70% ethanol, then mice were dissected to expose the hepatic portal vein, followed by

cannulation with a 24-gauge catheter needle. The cannula was secured, then the abdominal

aorta, inferior vena cava and cardiac right atrium were severed to isolate the liver from

peripheral tissues and prevent recirculation. Livers were perfused with oxygenated Krebs-

Henseleit buffer (118 mM NaCl, 4.7 mM KCl, 2.5 mM CaCl$_2$, 1.2 mM KH$_2$PO$_4$, 1.22 mM MgSO$_4$,

25 mM NaHCO$_3$, pH 7.4), warmed to ~45^{o}C and supplemented with 1.5 mM sodium lactate,

0.15 mM sodium pyruvate, 0.25 mM glycerol, 0.1 mM [U-^{13}C$_3$] propionate and 3% ^{2}H$_2$O.

Perfusion buffers were also supplemented with exogenous FAs. Either 0.2 mM sodium

octanoate was added directly to buffers, or 0.8 mM of a LCFA mixture (30% palmitate, 35%

oleate, 8% stearate, and 27% linoleate) was bound to FA-free BSA at a BSA:FA molar ratio of

1:3, then dialyzed using a capillary dialysis filtration system and a Reveclear-300 dialysis

cartridge (Baxter, 114745L) prior to addition to buffer. Specifics of buffer composition can be

found in each figure legend. Perfusion buffer was oxygenated by continuous bubbling of a 95%

O_2, 5% CO_2 gas mixture through the buffer reservoir using a fritted glass tube, and was

delivered using a peristaltic pump at a flow rate of ~8 mL/min for 60 minutes. Buffer and effluent

samples were collected every 10 minutes. At the completion of the perfusion, the liver was

freeze-clamped in liquid nitrogen, and subsequently weighed to obtain total liver weight for flux

normalizations. Perfusion buffer (input), effluent (output) and tissues were stored at -80°C until

analysis.

Flux Calculations in Perfused Livers. Stable-isotope based flux modeling used *in vivo* in humans

and rodents as described above was used to study metabolism of the *ex vivo* perfused liver

within minor modifications as described previously.[49] Briefly, total EGP rates **(V_{EGP})** was

calculated from the rate of appearance of unlabeled glucose in the perfusate. βOHB production

($V_{RaβOHB}$) and AcAc production **(V_{RaAcAc})** was calculated from the rate of appearance of

unlabeled ketones in the perfusate. Livers were perfused with ^2H_2O and [U-$^{13}C_3$] propionate,

which generate positional isotopomer populations that allow intrahepatic fluxes to be quantified

as described during *in vivo* experiments in humans and mice. The ^2H-enrichment of glucose at

H2, H5 and H6s reports on the fractional contribution of glycogen, glycerol and PEP to glucose

production, as described above. V_{EGP} measured from total glucose production in the perfusate,

therein allows $V_{Glycogen}$, $V_{Glycerol}$ and V_{PEP} to be quantified as described above. The ^{13}C-

enrichment of glucose at carbons C1, C2 and C3 from [U-$^{13}C_3$] propionate distort the multiplet

nuclear magnetic resonances of the C2 carbon in a manner that quantitatively encodes the ratio

of TCA cycle turnover relative to anaplerosis, cataplerosis and pyruvate cycling. Because at

steady-state the oxidative metabolic network herein described is coupled to PEP-linked GNG,

by normalizing fluxes to V_{PEP} (measured as described above), absolute fluxes through the TCA

cycle, anaplerosis, cataplerosis and pyruvate cycling can be calculated.

Hepatic Fat Oxidation Calculations in ex vivo perfused livers. In perfusions, total fat oxidation

was calculated as:

Total Fat Oxidation (µmol/min/kg LBM) = 2 ($V_{Ra\beta OHB}$ + V_{RaAcAc}) + V_{CS} + $V_{Acetate}$

and expressed as µmol acetyl-CoA turned over/min/g liver. $V_{Ra\beta OHB}$, V_{RaAcAc} and V_{CS} are

calculated as described above. Here, $V_{Acetate}$ is quantified as the appearance of unlabeled

acetate in the perfusate, and included in the *ex vivo LCFA* oxidation measurements. For

octanoate perfused livers, it was assumed that very little acetate arose from octanoate, and

therefore was not included in these calculations.

RE Production Calculations in Liver Perfusions. The same theory and assumption used to

model RE production rate *in vivo* in humans was used to model RE production rate in *ex vivo*

perfused liver, with minor modifications. For LCFA perfusions, RE's from each pathway were

calculated using the same equations as above. For octanoate perfused livers, the following

modifications were made. Because octanoate only generates 4 acetyl-CoA molecules, via three

rounds of β-oxidation which generate 6 RE, the ratio of RE's generated per acetyl-CoA molecule

is 1.5 (6 RE / 4 Ac-CoA = 1.5), and not 1.75 as when palmitate is metabolized. Based on this

the RE's generated from each pathway can be calculated as follows:

$RE_{AcAc} = 3 \times V_{RaAcAc}$

$RE_{\beta OHB} = 2 \times V_{Ra\beta OHB}$

$RE_{TCA\text{-}total} = 5.5 \times V_{CS}$

$RE_{TCA\text{-}only} = 4 \times V_{CS}$

$RE_{\beta ox\text{-}TCA} = 1.5 \times V_{CS}$

$RE_{\beta OX} = RE_{Acetate} + RE_{AcAc} + RE_{\beta OHB} + RE_{\beta ox}$

Total RE production rate can then be calculated based on $V_{Glycerol}$ and V_{PEP} based on the

assumptions that V_{PEP} is sourced 90% from lactate and 10% for pyruvate/alanine.

RE_{GNG} = V3 – (0.1 x V4)

Total RE produced (µmol/min/g liver) = $RE_{\beta OX}$ + $RE_{TCA\text{-}only}$ + RE_{GNG}

Liver mitochondrial isolation. Mice were fasted for 18h with access to water. Livers were excised

and placed in ice-cold mannitol-sucrose isolation media (MSIM; 220mM Mannitol, 70mM sucrose,

10mM Tris, 1mM EDTA; pH=7.2) supplemented with 5g/L fatty-acid free bovine serum albumin

(BSA) (MSIM+BSA). Livers were minced in 5ml MSIM and liver pieces were rinsed three times in

MSIM+BSA then transferred to an ice-chilled borosilicate glass vessel for homogenization.

Homogenized liver tissue was centrifuged at 700 x g for 10 min at 4°C and the supernatant was

filtered through two layers of gauze and centrifuged at 9,000 x g for 10 min at 4°C. The

supernatant was discarded, and pellets were gently resuspended in 10 ml MSIM+BSA and

centrifuged at 7,000 x g for 5 min at 4°C. The supernatant was discarded, and pellets were

resuspended in 1.2 ml MSIM+BSA and centrifuged at 9,000 x g at 4°C. MSIM+BSA was aspirated

from each tube and mitochondrial pellets were resuspended in 1.2 ml of MSIM without BSA and

centrifuged at 9,000 x g at 4°C. MSIM without BSA was aspirated from each tube and

mitochondrial pellets were resuspended in MSIM without BSA. Protein content was determined

via a BCA protein assay and mitochondria were resuspended at a final concentration of 20 mg/ml

for all functional assays.

Mitochondrial respiratory control. Liver mitochondria were isolated as described above. Steady-

state oxygen consumption rates (JO_2) were determined using a modified version of the creatine

kinase (CK) energetic clamp as described previously.[78] Briefly, isolated mitochondria (0.150

mg/ml) were added to Buffer Z (105 mM MES potassium salt, 30 mM KCl, 10 mM KH_2PO_4, 5 mM

$MgCl_2$) supplemented with Creatine (5 mM), PCr (1.5 mM), CK (20 U/mL), fatty acid free BSA

(2.5 mg/mL), and EGTA (1 mM) with either 10uM Hymeglusin (HG) or the vehicle control (DMSO),

followed by the addition of respiratory substrates (Pc/AKG; 50µM Palmitoyl-L-Carnitine, 5mM

alpha-ketoglutarate), and ATP (5mM). Next, sequential additions of PCr was used to adjust the

free energy of ATP hydrolysis (ΔG_{ATP}). Following the respiration assays, 1.6mL of the

mitochondrial buffer slurry was immediately collected and chilled on ice-water prior to differential

centrifugation (9,0000g for 5 minutes at 4 °C) to separate the mitochondrial supernatant.

Supernatants were flash frozen on dry-ice and stored at -80 °C until further analysis.

Measuring mitochondrial NAD(P)H/NAD(P)+ redox. Fluorescent measurements of mitochondrial

NAD(P)H/NAD(P)$^+$ redox were determined using a QuantaMaster Spectrofluorometer (QM-400;

Horiba Scientific) as described previously.[78] All assays were conducted at 37°C in a 0.2ml reaction

buffer. The buffer for all assays was Buffer Z supplemented with creatine (Cr; 5 mM),

phosphocreatine (PCr; 1.5 mM), creatine kinase (CK; 20 U/ml), and either 10uM Hymeglusin (HG)

or DMSO (VCTRL). At the start of each assay, isolated liver mitochondria (0.4 mg/ml) were added

to the assay buffer, followed by the addition of respiratory substrates (Pc/AKG) alone to capture

100% reduction within the NAD(P)H/NAD(P)$^+$ couple, and then sequential additions of PCr to

obtain the following final concentrations: 3, 6, 9, 12, 15mM. After the final PCr addition,

alamethicin (12.5ug/ml) was added to permeabilize the mitochondria. NAD(P)H/NAD(P)$^+$ during

the experiment was expressed as a percent reduced according to the formula: % Reduction = (F-

F0%)/(F100%-F0%), where the 0% reduction state is represented as the fluorescent signal

(Ex/Em, 340/450nm) recorded in the presence of alamethicin.

Measuring Mitochondrial βOHB Production. Following the mitochondrial respiration control

assays described above, 1.6mL of the mitochondria-buffer slurry was immediately collected and

chilled on ice-water prior to differential centrifugation (9,000xg for 5 minutes at 4°C) to isolate the

mitochondrial supernatant. βOHB levels in the supernatant were determined via a commercial kit

(Wako Autokit 3-HB, Fujifilm Healthcare Solutions), according to manufacturer specifications, with

modifications. Briefly, the principle of the assay is to colorimetrically monitor the reduction of

NAD+→NADH coupled to the oxidization of βOHB present in the assay samples. The reaction is

catalyzed by the addition of βOHB dehydrogenase present in the R2 working reagent. To ensure

AcAc does not interfere with the absorbance readings, AcAc in the samples is converted to

acetone following the addition of acetoacetone decarboxylase present in the R1 working reagent.

Prior to running the assay, mitochondrial supernatants were diluted 3-fold in Buffer Z. 20µL of

mitochondrial supernatant samples and βOHB standard curve samples (0-12.5µM) were loaded

onto a 96-well flat-bottom plate in duplicate. 180µL of R1 working reagent was added to the

standard and sample wells to monitor background absorbance at 405 nm for 5 minutes 37°C.

Following, 180µL/well of R2 working reagent was added, and absorbance was kinetically

monitored 405nm was monitored for 24 minutes at 37°C. For each of the βOHB standards,

absorbance slope values (Δ absorbance/Δ seconds) were used to derive a linear standard curve

to derive βOHB concentration present in each unknown sample. Unknown values were corrected

for protein and total assay volume and expressed as nmol βOHB/mg protein.

Mitochondrial incubations for metabolite profiling. Isolated liver mitochondria (0.02mg/mL) were

incubated in 2mL Buffer Z supplemented with creatine (Cr; 5 mM), phosphocreatine (PCr; 1.5

mM), creatine kinase (CK; 20 U/ml), respiratory substrates (Pc/AKG), and either 10uM

Hymeglusin (HG) or the vehicle control (VCTRL; DMSO), for 5 minutes at 37 °C with constant

stirring. Subsequently, 5mM of ATP was added to reach respiratory state 3, followed by a 5-

minute incubation at a fixed energy demand (ΔG_{ATP} = -14.26 kcals/mol) imposed by the addition

of PCr (final concentration = 6mM). 1.6mL of the mitochondrial buffer slurry was immediately

collected and chilled on ice-water prior to differential centrifugation (9,0000g for 5 minutes at 4

°C) to separate the mitochondrial pellet and supernatant. Both fractions were flash frozen on dry-

ice and stored at -80 °C until further analysis.

Mitochondrial acyl-CoA analysis by LC-MS/MS. Acyl-CoA analysis was performed as described

in previous reports.[78–80] Briefly, frozen mitochondrial pellets (0.4mg) were sonicated for ~5

minutes in 1.1 ml extraction buffer (1:1 methanol/water containing 5% acetic acid) containing 20

µl 0.01 mM D9 pentanoyl-CoA as internal standard. Homogenates were centrifuged at 4°C for 15-

minute at 18,000g and 900 µl of supernatant was used for solid phase extraction (SPE). For SPE,

the cartridge (1-ml ion exchange cartridge packed with 100 mg of 2-2(pyridyl)ethyl silica gel

(Sigma)) was pre-activated with 1 ml of methanol and then with 1 ml of extraction buffer (1:1

methanol/water containing 5% acetic acid). The acyl-CoAs trapped on the silica gel cartridge were

released with (i) 1 ml of a 1:1 mixture of 50 mm ammonium formate, pH 6.3, and methanol and

then (ii) 1 ml of a 1:3 mixture of 50 mm ammonium formate, pH 6.3 and (iii) 1 ml of methanol. The

combined effluent was dried with nitrogen gas and stored at −80 °C until LC-MS analysis. A 40 µl

sample was injected on a Pursuit XRs 5 C18 column (150 × 2.0 mm, 5µm), protected by a guard

column (Pursuit XRs 5 C18 5 µm, 10 × 2.0 mm, 5µm) with temperature controlled at 25 °C, in an

ExionLC AD liquid chromatograph system. The Analyst software (version 1.6) was used for data

record and processing. Relative pool size estimates were determined by comparing analyte peak

areas to D9 pentanoyl-CoA peak area, then normalized to the VCTRL (set to 1) to represent fold-

change in the pool size of CoA in the presence of HG. The detailed LC-MS/MS conditions can

refer to our previous reports.[78–80]

Mitochondrial organic acid profiling by LC-MS/MS. Organic acids in the incubation buffer were

methoximated and silylated to TBDMS-derivatives and profiled by a gas chromatography mass

spectrometry (GCMS) method. Briefly, 200 µl of incubation buffer was spiked with 20 µl 0.05 mM

norvaline as internal standard and then vortexed vigorously with 200 µl methanol and 1mL

acetonitrile. Homogenates were then centrifuged at 4C for 15-minute at 18,000g and 675 µl of the

resulting supernatant was dried under nitrogen gas at 37C. TBDMS-derivatives were prepared

for GC-MS with MOX and TBDMS as described previously.[81,82] GC-MS analyses were performed

on an Agilent 7890B GC system equipped with a HP-5MS capillary column (30 m, 0.25 mm i.d.,

0.25 µm-phase thickness; Agilent J&W Scientific), connected to an Agilent 5977A Mass

Spectrometer operating under ionization by electron impact (EI) at 70 eV. Relative pool size

estimates were determined by comparing analyte peak areas to norvaline peak area (m/z 288).

Quantification of hydroxyalkenal species in liver tissue. Frozen liver tissue was homogenized in

ice-cold water using 2.8- and 1.4-mm zirconium oxide beads at 4°C, then an appropriate amount

of internal standard was spike in based on total protein determined using the BCA assay, followed

by lipid extraction using a modified Bligh-dyer as previously described.[45,49] Lipid extracts were

then derivatized with carnosine and quantified using shotgun lipidomics as previously described.[83]

Gene Expression. For mRNA expression analysis, samples were harvested and homogenized in

Qiazol, then total RNA was isolated using Qiagen's RNeasy protocol. First-strand cDNA was

synthesized using iScript (BioRad, 170-8891), then quantitative real-time PCR was performed on

a BioRad CFX384 Real-Time thermocycler using SsoAdvanced Universal SYBR Green supermix

(BioRad, 170-5274). mRNA expression was calculated using the $2^{-\Delta\Delta Ct}$ method, where $\Delta Ct =$

housekeeping gene Ct – gene of interest CT, then expressed as fold-change relative to the intact

liver. mRNA levels were normalized to the intact liver, which was set to 1. Housekeeping gene

was *Rpl32* for all targets. Primer sequences are given in **Supplemental Table 2.**

Statistics and Data Analysis. All analyses were performed using GraphPad Prism version 9.

Unpaired 2-tailed Student's t-tests or 1-way ANOVA, was used to determine statistically

significant differences. P-value < 0.05 was accepted as significant for all tests. Information

about statistical analysis is included in each figure legend. Data are presented as mean ± SD,

unless otherwise indicated.

1000

1001

Acknowledgements

The authors acknowledge support from the National Institutes of Health (grants DK091538,

DK122832, HL166142, DK136772, and AG069781). The authors are also incredibly grateful to

the human participants who contributed to this study. The authors thank Rebecca Hollister,

Nicholas G. Evanoff and the Center for Pediatric Obesity Medicine at the University of

Minnesota Medical School for performing DXA scans. The authors also thank Oyedele Adeyi

and Khalid Amin for liver histopathology reports.

1009

<u>Acronyms</u>

3-hydroxy-3-methylglutaryl-CoA (HMG-CoA)

3-hydroxy-3-methylglutaryl-CoA synthase 2 (HMGCS2),

acetoacetate (AcAc)

acute insulin response to glucose (AIRg)

antisense oligonucleotide (ASO)

body mass index (BMI)

body weight (BW)

bovine serum albumin (BSA)

citrate synthase (CS)

clinical research unit (CRU)

D-beta-hydroxybutyrate (D-βOHB)

de novo lipogenesis (DNL)

dual-energy X-ray absorptiometry (DXA)

endogenous glucose production (EGP)

fatty acid (FA)

flavin adenine dinucleotide ($FADH_2$)

free fatty acids (FFAs)

frequently sampled intravenous glucose tolerance test (FSIVGTT)

gluconeogenesis (GNG)

glyceraldehyde-3-phosphate dehydrogenase (GAPDH)

glycerol-3-phosphate dehydrogenase (G3PDH),

high-fat carbohydrate-restricted (HFCR) diet

homeostatic model assessment for insulin resistance (HOMA-IR)

hymeglusin (HG)

Isotopomer Network Compartmental Analysis (INCA)

knock out (KO)

lactate dehydrogenase (LDH)

lean body mass (LBM)

long chain fatty acids (LCFAs)

magnetic resonance imaging (MRI)

malic enzyme (ME).

mass isotopomer distributions (MIDs)

mass spectrometry (MS)

metabolic-dysfunction-associated steatohepatitis (MASH)

metabolic-dysfunction-associated steatotic liver disease (MASLD)

monoacetone glucose (MAG)

NAFLD activity score (NAS)

nicotinamide adenine dinucleotide (NADH)

non esterified fatty acids (NEFAs)

nonalcoholic fatty liver disease (NAFLD)

nuclear magnetic resonance (NMR) spectroscopy

parallel reaction monitoring (PRM)

phosphoenolpyruvate (PEP)

polyunsaturated fatty acids (PUFAs)

proton density fat fraction (PDFF)

pyruvate kinase (PK)

reducing equivalents (RE's)

respiratory flux (JO2)

signal-to-noise (S/N)

sodium-glucose cotransporter 2 (SGLT2)

tandem mass spectrometry (MS/MS)

total ketone bodies (TKBs)

triacylglycerol (TAG)

tricarboxylic acid (TCA) cycle

U.S. Food and Drug Administration (FDA)

ultra-high performance liquid chromatography (UHPLC)

very low-density lipoprotein (VLDL)

References.

(1) Younossi, Z. M.; Golabi, P.; Paik, J. M.; Henry, A.; Van Dongen, C.; Henry, L. The Global Epidemiology of Nonalcoholic Fatty Liver Disease (NAFLD) and Nonalcoholic Steatohepatitis (NASH): A Systematic Review. *Hepatology* **2023**, *77* (4), 1335. https://doi.org/10.1097/HEP.0000000000000004.

(2) Riazi, K.; Azhari, H.; Charette, J. H.; Underwood, F. E.; King, J. A.; Afshar, E. E.; Swain, M. G.; Congly, S. E.; Kaplan, G. G.; Shaheen, A.-A. The Prevalence and Incidence of NAFLD Worldwide: A Systematic Review and Meta-Analysis. *Lancet Gastroenterol. Hepatol.* **2022**, *7* (9), 851–861. https://doi.org/10.1016/S2468-1253(22)00165-0.

(3) Ipsen, D. H.; Lykkesfeldt, J.; Tveden-Nyborg, P. Molecular Mechanisms of Hepatic Lipid Accumulation in Non-Alcoholic Fatty Liver Disease. *Cell. Mol. Life Sci.* **2018**, *75* (18), 3313–3327. https://doi.org/10.1007/s00018-018-2860-6.

(4) Postic, C.; Girard, J. Contribution of de Novo Fatty Acid Synthesis to Hepatic Steatosis and Insulin Resistance: Lessons from Genetically Engineered Mice. *J. Clin. Invest.* **2008**, *118* (3), 829–838. https://doi.org/10.1172/JCI34275.

(5) Pierantonelli, I.; Svegliati-Baroni, G. Nonalcoholic Fatty Liver Disease: Basic Pathogenetic Mechanisms in the Progression From NAFLD to NASH. *Transplantation* **2019**, *103* (1), e1. https://doi.org/10.1097/TP.0000000000002480.

(6) Sheka, A. C.; Adeyi, O.; Thompson, J.; Hameed, B.; Crawford, P. A.; Ikramuddin, S. Nonalcoholic Steatohepatitis: A Review. *JAMA* **2020**, *323* (12), 1175–1183. https://doi.org/10.1001/jama.2020.2298.

(7) Harrison, S. A.; Bedossa, P.; Guy, C. D.; Schattenberg, J. M.; Loomba, R.; Taub, R.; Labriola, D.; Moussa, S. E.; Neff, G. W.; Rinella, M. E.; Anstee, Q. M.; Abdelmalek, M. F.; Younossi, Z.; Baum, S. J.; Francque, S.; Charlton, M. R.; Newsome, P. N.; Lanthier, N.; Schiefke, I.; Mangia, A.; Pericàs, J. M.; Patil, R.; Sanyal, A. J.; Noureddin, M.; Bansal, M. B.; Alkhouri, N.; Castera, L.; Rudraraju, M.; Ratziu, V. A Phase 3, Randomized, Controlled Trial of Resmetirom in NASH with Liver Fibrosis. *N. Engl. J. Med.* **2024**, *390* (6), 497–509. https://doi.org/10.1056/NEJMoa2309000.

(8) Ferguson, D.; Finck, B. N. Emerging Therapeutic Approaches for the Treatment of NAFLD and Type 2 Diabetes Mellitus. *Nat. Rev. Endocrinol.* **2021**, *17* (8), 484–495. https://doi.org/10.1038/s41574-021-00507-z.

(9) Jang, H.; Kim, Y.; Lee, D. H.; Joo, S. K.; Koo, B. K.; Lim, S.; Lee, W.; Kim, W. Outcomes of Various Classes of Oral Antidiabetic Drugs on Nonalcoholic Fatty Liver Disease. *JAMA Intern. Med.* **2024**, *184* (4), 375–383. https://doi.org/10.1001/jamainternmed.2023.8029.

(10) Mak, S. K.; Wong, P. N.; Lee, K. F.; Fung, L. H.; Wong, A. K. Intracardiac Thrombus in an Adult Patient with Nephrotic Syndrome. *Nephrol. Dial. Transplant. Off. Publ. Eur. Dial. Transpl. Assoc. - Eur. Ren. Assoc.* **1996**, *11* (8), 1627–1630.

(11) Syed-Abdul, M. M.; Moore, M. P.; Wheeler, A. A.; Ganga, R. R.; Diaz-Arias, A.; Petroski, G. F.; Rector, R. S.; Ibdah, J. A.; Parks, E. J. Isotope Labeling and Biochemical Assessment of Liver-Triacylglycerol in Patients with Different Levels of Histologically-Graded Liver Disease. *J. Nutr.* **2023**, *153* (12), 3418–3429. https://doi.org/10.1016/j.tjnut.2023.09.018.

(12) Lambert, J. E.; Ramos–Roman, M. A.; Browning, J. D.; Parks, E. J. Increased De Novo Lipogenesis Is a Distinct Characteristic of Individuals With Nonalcoholic Fatty Liver Disease. *Gastroenterology* **2014**, *146* (3), 726–735. https://doi.org/10.1053/j.gastro.2013.11.049.

(13) Flannery, C.; Dufour, S.; Rabøl, R.; Shulman, G. I.; Petersen, K. F. Skeletal Muscle Insulin Resistance Promotes Increased Hepatic de Novo Lipogenesis, Hyperlipidemia, and

Hepatic Steatosis in the Elderly. *Diabetes* **2012**, *61* (11), 2711–2717. https://doi.org/10.2337/db12-0206.

(14) Donnelly, K. L.; Smith, C. I.; Schwarzenberg, S. J.; Jessurun, J.; Boldt, M. D.; Parks, E. J. Sources of Fatty Acids Stored in Liver and Secreted via Lipoproteins in Patients with Nonalcoholic Fatty Liver Disease. *J. Clin. Invest.* **2005**, *115* (5), 1343–1351. https://doi.org/10.1172/JCI23621.

(15) Schwarz, J.-M.; Linfoot, P.; Dare, D.; Aghajanian, K. Hepatic de Novo Lipogenesis in Normoinsulinemic and Hyperinsulinemic Subjects Consuming High-Fat, Low-Carbohydrate and Low-Fat, High-Carbohydrate Isoenergetic Diets. *Am. J. Clin. Nutr.* **2003**, *77* (1), 43–50. https://doi.org/10.1093/ajcn/77.1.43.

(16) Diraison, F.; Moulin, P.; Beylot, M. Contribution of Hepatic de Novo Lipogenesis and Reesterification of Plasma Non Esterified Fatty Acids to Plasma Triglyceride Synthesis during Non-Alcoholic Fatty Liver Disease. *Diabetes Metab.* **2003**, *29* (5), 478–485. https://doi.org/10.1016/s1262-3636(07)70061-7.

(17) Araya, J.; Rodrigo, R.; Videla, L. A.; Thielemann, L.; Orellana, M.; Pettinelli, P.; Poniachik, J. Increase in Long-Chain Polyunsaturated Fatty Acid n - 6/n - 3 Ratio in Relation to Hepatic Steatosis in Patients with Non-Alcoholic Fatty Liver Disease. *Clin. Sci. Lond. Engl. 1979* **2004**, *106* (6), 635–643. https://doi.org/10.1042/CS20030326.

(18) Allard, J. P.; Aghdassi, E.; Mohammed, S.; Raman, M.; Avand, G.; Arendt, B. M.; Jalali, P.; Kandasamy, T.; Prayitno, N.; Sherman, M.; Guindi, M.; Ma, D. W. L.; Heathcote, J. E. Nutritional Assessment and Hepatic Fatty Acid Composition in Non-Alcoholic Fatty Liver Disease (NAFLD): A Cross-Sectional Study. *J. Hepatol.* **2008**, *48* (2), 300–307. https://doi.org/10.1016/j.jhep.2007.09.009.

(19) Arendt, B. M.; Ma, D. W. L.; Simons, B.; Noureldin, S. A.; Therapondos, G.; Guindi, M.; Sherman, M.; Allard, J. P. Nonalcoholic Fatty Liver Disease Is Associated with Lower Hepatic and Erythrocyte Ratios of Phosphatidylcholine to Phosphatidylethanolamine. *Appl. Physiol. Nutr. Metab. Physiol. Appl. Nutr. Metab.* **2013**, *38* (3), 334–340. https://doi.org/10.1139/apnm-2012-0261.

(20) Spooner, M. H.; Jump, D. B. Nonalcoholic Fatty Liver Disease and Omega-3 Fatty Acids: Mechanisms and Clinical Use. *Annu. Rev. Nutr.* **2023**, *43* (1), 199–223. https://doi.org/10.1146/annurev-nutr-061021-030223.

(21) Spooner, M. H.; Garcia-Jaramillo, M.; Apperson, K. D.; Löhr, C. V.; Jump, D. B. Time Course of Western Diet (WD) Induced Nonalcoholic Steatohepatitis (NASH) in Female and Male Ldlr-/- Mice. *PloS One* **2023**, *18* (10), e0292432. https://doi.org/10.1371/journal.pone.0292432.

(22) Fletcher, J. A.; Deja, S.; Satapati, S.; Fu, X.; Burgess, S. C.; Browning, J. D. Impaired Ketogenesis and Increased Acetyl-CoA Oxidation Promote Hyperglycemia in Human Fatty Liver. *JCI Insight* **2019**, *4* (11). https://doi.org/10.1172/jci.insight.127737.

(23) Koliaki, C.; Szendroedi, J.; Kaul, K.; Jelenik, T.; Nowotny, P.; Jankowiak, F.; Herder, C.; Carstensen, M.; Krausch, M.; Knoefel, W. T.; Schlensak, M.; Roden, M. Adaptation of Hepatic Mitochondrial Function in Humans with Non-Alcoholic Fatty Liver Is Lost in Steatohepatitis. *Cell Metab.* **2015**, *21* (5), 739–746. https://doi.org/10.1016/j.cmet.2015.04.004.

(24) Satapati, S.; Sunny, N. E.; Kucejova, B.; Fu, X.; He, T. T.; Méndez-Lucas, A.; Shelton, J. M.; Perales, J. C.; Browning, J. D.; Burgess, S. C. Elevated TCA Cycle Function in the Pathology of Diet-Induced Hepatic Insulin Resistance and Fatty Liver[S]. *J. Lipid Res.* **2012**, *53* (6), 1080–1092. https://doi.org/10.1194/jlr.M023382.

(25) Sunny, N. E.; Parks, E. J.; Browning, J. D.; Burgess, S. C. Excessive Hepatic Mitochondrial TCA Cycle and Gluconeogenesis in Humans with Nonalcoholic Fatty Liver Disease. *Cell Metab.* **2011**, *14* (6), 804–810. https://doi.org/10.1016/j.cmet.2011.11.004.

(26) Petersen, K. F.; Befroy, D. E.; Dufour, S.; Rothman, D. L.; Shulman, G. I. Assessment of Hepatic Mitochondrial Oxidation and Pyruvate Cycling in NAFLD by 13C Magnetic Resonance Spectroscopy. *Cell Metab.* **2016**, *24* (1), 167–171. https://doi.org/10.1016/j.cmet.2016.06.005.

(27) Petersen, K. F.; Dufour, S.; Mehal, W. Z.; Shulman, G. I. Glucagon Promotes Increased Hepatic Mitochondrial Oxidation and Pyruvate Carboxylase Flux in Humans with Fatty Liver Disease. *Cell Metab.* **2024**, *0* (0). https://doi.org/10.1016/j.cmet.2024.07.023.

(28) Lund, M. T.; Kristensen, M.; Hansen, M.; Tveskov, L.; Floyd, A. K.; Støckel, M.; Vainer, B.; Poulsen, S. S.; Helge, J. W.; Prats, C.; Dela, F. Hepatic Mitochondrial Oxidative Phosphorylation Is Normal in Obese Patients with and without Type 2 Diabetes. *J. Physiol.* **2016**, *594* (15), 4351–4358. https://doi.org/10.1113/JP272105.

(29) Kotronen, A.; Seppälä-Lindroos, A.; Vehkavaara, S.; Bergholm, R.; Frayn, K. N.; Fielding, B. A.; Yki-Järvinen, H. Liver Fat and Lipid Oxidation in Humans. *Liver Int.* **2009**, *29* (9), 1439–1446. https://doi.org/10.1111/j.1478-3231.2009.02076.x.

(30) Cortez-Pinto, H.; Chatham, J.; Chacko, V. P.; Arnold, C.; Rashid, A.; Diehl, A. M. Alterations in Liver ATP Homeostasis in Human Nonalcoholic SteatohepatitisA Pilot Study. *JAMA* **1999**, *282* (17), 1659–1664. https://doi.org/10.1001/jama.282.17.1659.

(31) Rector, R. S.; Thyfault, J. P.; Uptergrove, G. M.; Morris, E. M.; Naples, S. P.; Borengasser, S. J.; Mikus, C. R.; Laye, M. J.; Laughlin, M. H.; Booth, F. W.; Ibdah, J. A. Mitochondrial Dysfunction Precedes Insulin Resistance and Hepatic Steatosis and Contributes to the Natural History of Non-Alcoholic Fatty Liver Disease in an Obese Rodent Model. *J. Hepatol.* **2010**, *52* (5), 727–736. https://doi.org/10.1016/j.jhep.2009.11.030.

(32) Croci, I.; Byrne, N. M.; Choquette, S.; Hills, A. P.; Chachay, V. S.; Clouston, A. D.; O'Moore-Sullivan, T. M.; Macdonald, G. A.; Prins, J. B.; Hickman, I. J. Whole-Body Substrate Metabolism Is Associated with Disease Severity in Patients with Non-Alcoholic Fatty Liver Disease. *Gut* **2013**, *62* (11), 1625–1633. https://doi.org/10.1136/gutjnl-2012-302789.

(33) Moore, M. P.; Cunningham, R. P.; Meers, G. M.; Johnson, S. A.; Wheeler, A. A.; Ganga, R. R.; Spencer, N. M.; Pitt, J. B.; Diaz-Arias, A.; Swi, A. I. A.; Hammoud, G. M.; Ibdah, J. A.; Parks, E. J.; Rector, R. S. Compromised Hepatic Mitochondrial Fatty Acid Oxidation and Reduced Markers of Mitochondrial Turnover in Human NAFLD. *Hepatol. Baltim. Md* **2022**, *76* (5), 1452–1465. https://doi.org/10.1002/hep.32324.

(34) Moore, M. P.; Shryack, G.; Alessi, I.; Wieschhaus, N.; Meers, G. M.; Johnson, S. A.; Wheeler, A. A.; Ibdah, J. A.; Parks, E. J.; Rector, R. S. Relationship Between Serum β-Hydroxybutyrate and Hepatic Fatty Acid Oxidation in Individuals with Obesity and NAFLD. *Am. J. Physiol.-Endocrinol. Metab.* **2024**, ajpendo.00336.2023. https://doi.org/10.1152/ajpendo.00336.2023.

(35) Satapati, S.; Kucejova, B.; Duarte, J. A. G.; Fletcher, J. A.; Reynolds, L.; Sunny, N. E.; He, T.; Nair, L. A.; Livingston, K. A.; Fu, X.; Merritt, M. E.; Sherry, A. D.; Malloy, C. R.; Shelton, J. M.; Lambert, J.; Parks, E. J.; Corbin, I.; Magnuson, M. A.; Browning, J. D.; Burgess, S. C. Mitochondrial Metabolism Mediates Oxidative Stress and Inflammation in Fatty Liver. *J. Clin. Invest.* **2015**, *125* (12), 4447–4462. https://doi.org/10.1172/JCI82204.

(36) Burgess, S. C.; Hausler, N.; Merritt, M.; Jeffrey, F. M. H.; Storey, C.; Milde, A.; Koshy, S.; Lindner, J.; Magnuson, M. A.; Malloy, C. R.; Sherry, A. D. Impaired Tricarboxylic Acid Cycle Activity in Mouse Livers Lacking Cytosolic Phosphoenolpyruvate Carboxykinase. *J. Biol. Chem.* **2004**, *279* (47), 48941–48949. https://doi.org/10.1074/jbc.M407120200.

(37) Fu, X.; Fletcher, J. A.; Deja, S.; Inigo-Vollmer, M.; Burgess, S. C.; Browning, J. D. Persistent Fasting Lipogenesis Links Impaired Ketogenesis with Citrate Synthesis in Humans with Nonalcoholic Fatty Liver. *J. Clin. Invest. 133* (9), e167442. https://doi.org/10.1172/JCI167442.

(38) Adiels, M.; Taskinen, M.-R.; Packard, C.; Caslake, M. J.; Soro-Paavonen, A.; Westerbacka, J.; Vehkavaara, S.; Häkkinen, A.; Olofsson, S.-O.; Yki-Järvinen, H.; Borén, J. Overproduction of Large VLDL Particles Is Driven by Increased Liver Fat Content in Man. *Diabetologia* **2006**, *49* (4), 755–765. https://doi.org/10.1007/s00125-005-0125-z.

(39) Choi, S. H.; Ginsberg, H. N. Increased Very Low Density Lipoprotein (VLDL) Secretion, Hepatic Steatosis, and Insulin Resistance. *Trends Endocrinol. Metab. TEM* **2011**, *22* (9), 353–363. https://doi.org/10.1016/j.tem.2011.04.007.

(40) Conboy, E.; Vairo, F.; Schultz, M.; Agre, K.; Ridsdale, R.; Deyle, D.; Oglesbee, D.; Gavrilov, D.; Klee, E. W.; Lanpher, B. Mitochondrial 3-Hydroxy-3-Methylglutaryl-CoA Synthase Deficiency: Unique Presenting Laboratory Values and a Review of Biochemical and Clinical Features. *JIMD Rep.* **2018**, *40*, 63–69. https://doi.org/10.1007/8904_2017_59.

(41) Bouchard, L.; Robert, M. F.; Vinarov, D.; Stanley, C. A.; Thompson, G. N.; Morris, A.; Leonard, J. V.; Quant, P.; Hsu, B. Y.; Boneh, A.; Boukaftane, Y.; Ashmarina, L.; Wang, S.; Miziorko, H.; Mitchell, G. A. Mitochondrial 3-Hydroxy-3-Methylglutaryl-CoA Synthase Deficiency: Clinical Course and Description of Causal Mutations in Two Patients. *Pediatr. Res.* **2001**, *49* (3), 326–331. https://doi.org/10.1203/00006450-200103000-00005.

(42) Cotter, D. G.; Ercal, B.; Huang, X.; Leid, J. M.; d'Avignon, D. A.; Graham, M. J.; Dietzen, D. J.; Brunt, E. M.; Patti, G. J.; Crawford, P. A. Ketogenesis Prevents Diet-Induced Fatty Liver Injury and Hyperglycemia. *J. Clin. Invest.* **2014**, *124* (12), 5175–5190. https://doi.org/10.1172/JCI76388.

(43) Puchalska, P.; Martin, S. E.; Huang, X.; Lengfeld, J. E.; Daniel, B.; Graham, M. J.; Han, X.; Nagy, L.; Patti, G. J.; Crawford, P. A. Hepatocyte-Macrophage Acetoacetate Shuttle Protects against Tissue Fibrosis. *Cell Metab.* **2019**, *29* (2), 383-398.e7. https://doi.org/10.1016/j.cmet.2018.10.015.

(44) Asif, S.; Kim, R. Y.; Fatica, T.; Sim, J.; Zhao, X.; Oh, Y.; Denoncourt, A.; Cheung, A. C.; Downey, M.; Mulvihill, E. E.; Kim, K.-H. Hmgcs2-Mediated Ketogenesis Modulates High-Fat Diet-Induced Hepatosteatosis. *Mol. Metab.* **2022**, *61*, 101494. https://doi.org/10.1016/j.molmet.2022.101494.

(45) Queathem, E. D.; Moazzami, Z.; Stagg, D. B.; Nelson, A. B.; Fulghum, K.; Hayir, A.; Seay, A.; Gillingham, J. R.; d'Avignon, D. A.; Han, X.; Ruan, H.-B.; Crawford, P. A.; Puchalska, P. Ketogenesis Supports Hepatic Polyunsaturated Fatty Acid Homeostasis via Fatty Acid Elongation. *BioRxiv Prepr. Serv. Biol.* **2024**, 2024.07.09.602593. https://doi.org/10.1101/2024.07.09.602593.

(46) Ramakrishnan, S.; Mooli, R. G. R.; Han, Y.; Fiorenza, E.; Kumar, S.; Bello, F.; Nallanagulagari, A.; Karra, S.; Teng, L.; Jurczak, M. Hepatic Ketogenesis Regulates Lipid Homeostasis via ACSL1-Mediated Fatty Acid Partitioning. July 18, 2023. https://doi.org/10.21203/rs.3.rs-3147009/v1.

(47) Sanyal, A. J.; Campbell-Sargent, C.; Mirshahi, F.; Rizzo, W. B.; Contos, M. J.; Sterling, R. K.; Luketic, V. A.; Shiffman, M. L.; Clore, J. N. Nonalcoholic Steatohepatitis: Association of Insulin Resistance and Mitochondrial Abnormalities. *Gastroenterology* **2001**, *120* (5), 1183–1192. https://doi.org/10.1053/gast.2001.23256.

(48) Deja, S.; Fu, X.; Fletcher, J. A.; Kucejova, B.; Browning, J. D.; Young, J. D.; Burgess, S. C. Simultaneous Tracers and a Unified Model of Positional and Mass Isotopomers for Quantification of Metabolic Flux in Liver. *Metab. Eng.* **2020**, *59*, 1–14. https://doi.org/10.1016/j.ymben.2019.12.005.

(49) d'Avignon, D. A.; Puchalska, P.; Ercal, B.; Chang, Y.; Martin, S. E.; Graham, M. J.; Patti, G. J.; Han, X.; Crawford, P. A. Hepatic Ketogenic Insufficiency Reprograms Hepatic Glycogen Metabolism and the Lipidome. *JCI Insight* **2018**, *3* (12), e99762, 99762. https://doi.org/10.1172/jci.insight.99762.

(50) Stagg, D. B.; Gillingham, J. R.; Nelson, A. B.; Lengfeld, J. E.; d'Avignon, D. A.; Puchalska, P.; Crawford, P. A. Diminished Ketone Interconversion, Hepatic TCA Cycle Flux, and

Glucose Production in D-β-Hydroxybutyrate Dehydrogenase Hepatocyte-Deficient Mice. *Mol. Metab.* **2021**, *53*, 101269. https://doi.org/10.1016/j.molmet.2021.101269.

(51) Feola, K.; Venable, A. H.; Broomfield, T.; Villegas, M.; Fu, X.; Burgess, S.; Huen, S. C. Hepatic Ketogenesis Is Not Required for Starvation Adaptation in Mice. *Mol. Metab.* **2024**, *86*, 101967. https://doi.org/10.1016/j.molmet.2024.101967.

(52) Alexander, M.; Upadhyay, V.; Rock, R.; Ramirez, L.; Trepka, K.; Puchalska, P.; Orellana, D.; Ang, Q. Y.; Whitty, C.; Turnbaugh, J. A.; Tian, Y.; Dumlao, D.; Nayak, R.; Patterson, A.; Newman, J. C.; Crawford, P. A.; Turnbaugh, P. J. A Diet-Dependent Host Metabolite Shapes the Gut Microbiota to Protect from Autoimmunity. bioRxiv May 13, 2024, p 2023.11.02.565382. https://doi.org/10.1101/2023.11.02.565382.

(53) Puchalska, P.; Nelson, A. B.; Stagg, D. B.; Crawford, P. A. Determination of Ketone Bodies in Biological Samples via Rapid UPLC-MS/MS. *Talanta* **2021**, *225*, 122048. https://doi.org/10.1016/j.talanta.2020.122048.

(54) Hasenour, C. M.; Wall, M. L.; Ridley, D. E.; Hughey, C. C.; James, F. D.; Wasserman, D. H.; Young, J. D. Mass Spectrometry-Based Microassay of (2)H and (13)C Plasma Glucose Labeling to Quantify Liver Metabolic Fluxes in Vivo. *Am. J. Physiol. Endocrinol. Metab.* **2015**, *309* (2), E191-203. https://doi.org/10.1152/ajpendo.00003.2015.

(55) Hughey, C. C.; James, F. D.; Bracy, D. P.; Donahue, E. P.; Young, J. D.; Viollet, B.; Foretz, M.; Wasserman, D. H. Loss of Hepatic AMP-Activated Protein Kinase Impedes the Rate of Glycogenolysis but Not Gluconeogenic Fluxes in Exercising Mice. *J. Biol. Chem.* **2017**, *292* (49), 20125–20140. https://doi.org/10.1074/jbc.M117.811547.

(56) Rome, F. I.; Shobert, G. L.; Voigt, W. C.; Stagg, D. B.; Puchalska, P.; Burgess, S. C.; Crawford, P. A.; Hughey, C. C. Loss of Hepatic Phosphoenolpyruvate Carboxykinase 1 Dysregulates Metabolic Responses to Acute Exercise but Enhances Adaptations to Exercise Training in Mice. *Am. J. Physiol. Endocrinol. Metab.* **2023**, *324* (1), E9–E23. https://doi.org/10.1152/ajpendo.00222.2022.

(57) Fisher-Wellman, K. H.; Davidson, M. T.; Narowski, T. M.; Lin, C.-T.; Koves, T. R.; Muoio, D. M. Mitochondrial Diagnostics: A Multiplexed Assay Platform for Comprehensive Assessment of Mitochondrial Energy Fluxes. *Cell Rep.* **2018**, *24* (13), 3593-3606.e10. https://doi.org/10.1016/j.celrep.2018.08.091.

(58) Tomoda, H.; Ohbayashi, N.; Morikawa, Y.; Kumagai, H.; Omura, S. Binding Site for Fungal Beta-Lactone Hymeglusin on Cytosolic 3-Hydroxy-3-Methylglutaryl Coenzyme A Synthase. *Biochim. Biophys. Acta* **2004**, *1636* (1), 22–28. https://doi.org/10.1016/j.bbalip.2003.11.005.

(59) Le Foll, C.; Dunn-Meynell, A. A.; Miziorko, H. M.; Levin, B. E. Regulation of Hypothalamic Neuronal Sensing and Food Intake by Ketone Bodies and Fatty Acids. *Diabetes* **2014**, *63* (4), 1259–1269. https://doi.org/10.2337/db13-1090.

(60) Veech, R. L.; Todd King, M.; Pawlosky, R.; Kashiwaya, Y.; Bradshaw, P. C.; Curtis, W. The "Great" Controlling Nucleotide Coenzymes. *IUBMB Life* **2019**, *71* (5), 565–579. https://doi.org/10.1002/iub.1997.

(61) Schmid, A. I.; Szendroedi, J.; Chmelik, M.; Krššák, M.; Moser, E.; Roden, M. Liver ATP Synthesis Is Lower and Relates to Insulin Sensitivity in Patients With Type 2 Diabetes. *Diabetes Care* **2011**, *34* (2), 448–453. https://doi.org/10.2337/dc10-1076.

(62) Felig, P.; Wahren, J.; Hendler, R.; Brundin, T. Splanchnic Glucose and Amino Acid Metabolism in Obesity. *J. Clin. Invest.* **1974**, *53* (2), 582–590. https://doi.org/10.1172/JCI107593.

(63) Xu, J.; Nakamura, M. T.; Cho, H. P.; Clarke, S. D. Sterol Regulatory Element Binding Protein-1 Expression Is Suppressed by Dietary Polyunsaturated Fatty Acids. A Mechanism for the Coordinate Suppression of Lipogenic Genes by Polyunsaturated Fats. *J. Biol. Chem.* **1999**, *274* (33), 23577–23583. https://doi.org/10.1074/jbc.274.33.23577.

(64) Yahagi, N.; Shimano, H.; Hasty, A. H.; Amemiya-Kudo, M.; Okazaki, H.; Tamura, Y.; Iizuka, Y.; Shionoiri, F.; Ohashi, K.; Osuga, J.; Harada, K.; Gotoda, T.; Nagai, R.; Ishibashi, S.; Yamada, N. A Crucial Role of Sterol Regulatory Element-Binding Protein-1 in the Regulation of Lipogenic Gene Expression by Polyunsaturated Fatty Acids. *J. Biol. Chem.* **1999**, *274* (50), 35840–35844. https://doi.org/10.1074/jbc.274.50.35840.

(65) Botolin, D.; Wang, Y.; Christian, B.; Jump, D. B. Docosahexaneoic Acid (22:6,n-3) Regulates Rat Hepatocyte SREBP-1 Nuclear Abundance by Erk- and 26S Proteasome-Dependent Pathways. *J. Lipid Res.* **2006**, *47* (1), 181–192. https://doi.org/10.1194/jlr.M500365-JLR200.

(66) Takeuchi, Y.; Yahagi, N.; Izumida, Y.; Nishi, M.; Kubota, M.; Teraoka, Y.; Yamamoto, T.; Matsuzaka, T.; Nakagawa, Y.; Sekiya, M.; Iizuka, Y.; Ohashi, K.; Osuga, J.; Gotoda, T.; Ishibashi, S.; Itaka, K.; Kataoka, K.; Nagai, R.; Yamada, N.; Kadowaki, T.; Shimano, H. Polyunsaturated Fatty Acids Selectively Suppress Sterol Regulatory Element-Binding Protein-1 through Proteolytic Processing and Autoloop Regulatory Circuit. *J. Biol. Chem.* **2010**, *285* (15), 11681–11691. https://doi.org/10.1074/jbc.M109.096107.

(67) Deng, X.; Dong, Q.; Bridges, D.; Raghow, R.; Park, E. A.; Elam, M. B. Docosahexaenoic Acid Inhibits Proteolytic Processing of Sterol Regulatory Element-Binding Protein-1c (SREBP-1c) via Activation of AMP-Activated Kinase. *Biochim. Biophys. Acta BBA - Mol. Cell Biol. Lipids* **2015**, *1851* (12), 1521–1529. https://doi.org/10.1016/j.bbalip.2015.08.007.

(68) Green, C. J.; Pramfalk, C.; Charlton, C. A.; Gunn, P. J.; Cornfield, T.; Pavlides, M.; Karpe, F.; Hodson, L. Hepatic de Novo Lipogenesis Is Suppressed and Fat Oxidation Is Increased by Omega-3 Fatty Acids at the Expense of Glucose Metabolism. *BMJ Open Diabetes Res. Care* **2020**, *8* (1), e000871. https://doi.org/10.1136/bmjdrc-2019-000871.

(69) Malloy, C. R.; Jones, J. G.; Jeffrey, F. M.; Jessen, M. E.; Sherry, A. D. Contribution of Various Substrates to Total Citric Acid Cycle Flux and Anaplerosis as Determined by 13C Isotopomer Analysis and O2 Consumption in the Heart. *Magma N. Y. N* **1996**, *4* (1), 35–46. https://doi.org/10.1007/BF01759778.

(70) Postic, C.; Shiota, M.; Niswender, K. D.; Jetton, T. L.; Chen, Y.; Moates, J. M.; Shelton, K. D.; Lindner, J.; Cherrington, A. D.; Magnuson, M. A. Dual Roles for Glucokinase in Glucose Homeostasis as Determined by Liver and Pancreatic β Cell-Specific Gene Knock-Outs Using Cre Recombinase*. *J. Biol. Chem.* **1999**, *274* (1), 305–315. https://doi.org/10.1074/jbc.274.1.305.

(71) Horton, J. L.; Davidson, M. T.; Kurishima, C.; Vega, R. B.; Powers, J. C.; Matsuura, T. R.; Petucci, C.; Lewandowski, E. D.; Crawford, P. A.; Muoio, D. M.; Recchia, F. A.; Kelly, D. P. The Failing Heart Utilizes 3-Hydroxybutyrate as a Metabolic Stress Defense. *JCI Insight* **2019**, *4* (4), e124079, 124079. https://doi.org/10.1172/jci.insight.124079.

(72) Puchalska, P.; Nelson, A. B.; Stagg, D. B.; Crawford, P. A. Determination of Ketone Bodies in Biological Samples via Rapid UPLC-MS/MS. *Talanta* **2021**, *225*, 122048. https://doi.org/10.1016/j.talanta.2020.122048.

(73) Ayala, J. E.; Bracy, D. P.; Malabanan, C.; James, F. D.; Ansari, T.; Fueger, P. T.; McGuinness, O. P.; Wasserman, D. H. Hyperinsulinemic-Euglycemic Clamps in Conscious, Unrestrained Mice. *J. Vis. Exp. JoVE* **2011**, No. 57, 3188. https://doi.org/10.3791/3188.

(74) Rome, F. I.; Hughey, C. C. Disrupted Liver Oxidative Metabolism in Glycine N-Methyltransferase-Deficient Mice Is Mitigated by Dietary Methionine Restriction. *Mol. Metab.* **2022**, *58*, 101452. https://doi.org/10.1016/j.molmet.2022.101452.

(75) Antoniewicz, M. R.; Kelleher, J. K.; Stephanopoulos, G. Measuring Deuterium Enrichment of Glucose Hydrogen Atoms by Gas Chromatography/Mass Spectrometry. *Anal. Chem.* **2011**, *83* (8), 3211–3216. https://doi.org/10.1021/ac200012p.

(76) Young, J. D. INCA: A Computational Platform for Isotopically Non-Stationary Metabolic Flux Analysis. *Bioinformatics* **2014**, *30* (9), 1333–1335. https://doi.org/10.1093/bioinformatics/btu015.

(77) Fu, X.; Deja, S.; Kucejova, B.; Duarte, J. A. G.; McDonald, J. G.; Burgess, S. C. Targeted Determination of Tissue Energy Status by LC-MS/MS. *Anal. Chem.* **2019**, *91* (9), 5881–5887. https://doi.org/10.1021/acs.analchem.9b00217.

(78) Tr, K.; Gf, Z.; Mt, D.; Ab, C.; Sb, C.; Jm, J.; Dh, S.; Pa, G.; Dm, M. Pyruvate-Supported Flux through Medium-Chain Ketothiolase Promotes Mitochondrial Lipid Tolerance in Cardiac and Skeletal Muscles. *Cell Metab.* **2023**, *35* (6). https://doi.org/10.1016/j.cmet.2023.03.016.

(79) Li, Q.; Zhang, S.; Berthiaume, J. M.; Simons, B.; Zhang, G.-F. Novel Approach in LC-MS/MS Using MRM to Generate a Full Profile of Acyl-CoAs: Discovery of Acyl-Dephospho-CoAs. *J. Lipid Res.* **2014**, *55* (3), 592–602. https://doi.org/10.1194/jlr.D045112.

(80) Zhang, G.-F.; Kombu, R. S.; Kasumov, T.; Han, Y.; Sadhukhan, S.; Zhang, J.; Sayre, L. M.; Ray, D.; Gibson, K. M.; Anderson, V. A.; Tochtrop, G. P.; Brunengraber, H. Catabolism of 4-Hydroxyacids and 4-Hydroxynonenal via 4-Hydroxy-4-Phosphoacyl-CoAs. *J. Biol. Chem.* **2009**, *284* (48), 33521–33534. https://doi.org/10.1074/jbc.M109.055665.

(81) Williams, A. S.; Crown, S. B.; Lyons, S. P.; Koves, T. R.; Wilson, R. J.; Johnson, J. M.; Slentz, D. H.; Kelly, D. P.; Grimsrud, P. A.; Zhang, G.-F.; Muoio, D. M. Ketone Flux through BDH1 Supports Metabolic Remodeling of Skeletal and Cardiac Muscles in Response to Intermittent Time-Restricted Feeding. *Cell Metab.* **2024**, *36* (2), 422-437.e8. https://doi.org/10.1016/j.cmet.2024.01.007.

(82) Williams, A. S.; Koves, T. R.; Davidson, M. T.; Crown, S. B.; Fisher-Wellman, K. H.; Torres, M. J.; Draper, J. A.; Narowski, T. M.; Slentz, D. H.; Lantier, L.; Wasserman, D. H.; Grimsrud, P. A.; Muoio, D. M. Disruption of Acetyl-Lysine Turnover in Muscle Mitochondria Promotes Insulin Resistance and Redox Stress without Overt Respiratory Dysfunction. *Cell Metab.* **2020**, *31* (1), 131-147.e11. https://doi.org/10.1016/j.cmet.2019.11.003.

(83) Wang, M.; Fang, H.; Han, X. Shotgun Lipidomics Analysis of 4-Hydroxyalkenal Species Directly from Lipid Extracts after One-Step in Situ Derivatization. *Anal. Chem.* **2012**, *84* (10), 4580–4586. https://doi.org/10.1021/ac300695p.

Figure Legends

Figure 1. Characteristics of liver-specific, and systemic metabolic health in bariatric humans with MASLD-MASH. (A) The initial stages of metabolic-dysfunction associated steatotic liver disease (MASLD) begins with the accumulation of fat in the liver (i.e., hepatic steatosis), which has been linked to an acceleration in: de novo lipogenesis (DNL), the turnover of the tricarboxylic acid (TCA) cycle, and phosphoenolpyruvate (PEP)-derived gluconeogenesis (GNG). Metabolic shifts during the progression of MASLD to metabolic-dysfunction associated steatohepatitis (MASH) remain poorly understood. **(B)** Study design. MRI-based screening of bariatric participants (BMI ≥ 35) identified patients with a liver proton density fat fraction (PDFF) greater than 5%. Each participant underwent a liver biopsy, FSIVGTT and DXA. After an overnight fast, a panel of eight hepatic intermediary metabolic fluxes were quantified at steady-state using a multi-stable-isotope tracing approach. Metabolic fluxes were correlated with clinical markers of liver and systemic health. **(C)** The breakdown of the NAFLD activity score (NAS) in all 14 participants. **(D)** Correlation matrix of NAS with individual histological scores for steatosis, ballooning, inflammation, and fibrosis, as well as PDFF. Correlation of NAS with **(E)** body mass index (BMI), **(F)** acute insulin response to glucose (AIRg), and **(G)** the homeostatic model assessment for insulin resistance (HOMA-IR). Pearson correlations coefficients (r) are given on each graph along with a line of best fit calculated using linear regression. For the correlation matrix, Pearson correlations coefficients (r) are shown in heat map format with the magnitude of the correlation given by the right-hand legend and displayed in each square. Correlations were accepted as significant if $p < 0.05$ and are shown on each graph, or in each box on the correlation matrix an indicated, * $p < 0.05$, ** $p < 0.01$, ***$p < 0.001$.

Figure 2. Liver injury in MASLD-MASH correlates with endogenous ketogenesis and hepatic fat oxidation. Flux through hepatic intermediary metabolic pathways is quantified *in vivo* non-invasively following administration of stable isotope tracers. Participants orally ingested

heavy water (^2H_2O) and [U-$^{13}C_3$] propionate. [3,4-$^{13}C_2$] glucose and [U-$^{13}C_4$]βOHB were intravenously infused allowing whole body glucose and βOHB turnover to be quantified at metabolic and isotopic steady state. **(A)** Fractional sourcing of glucose production can be quantified from the ^2H enrichment pattern of plasma glucose using ^2H NMR. The product of total glucose production and fractional sourcing thereby allowing absolute reaction velocities for glucose sourcing pathways to be quantified in the liver. **(B)** Administration of [U-$^{13}C_3$] propionate ^{13}C-enriches TCA cycle intermediates, which are in exchange with phosphoenolpyruvate (PEP). Using ^{13}C NMR, the multiplet arising from the C2 resonance of plasma glucose, the resulting metabolic network models oxidative and anaplerotic nodes of the TCA cycle in parallel to glucose production. Normalizing fluxes to the rate of PEP conversion to glucose (panel A), the absolute reaction velocities of the TCA cycle, anaplerosis, cataplerosis and pyruvate cycling can be quantified. **(C)** Average percent of endogenous glucose production (V_{EGP}) derived from glycogen, glycerol and PEP across all 14 participants, highlighting that TCA cycle-sourced PEP is the major contributor to V_{EGP} in the fasted state. **(D)** Average RE's derived from GNG, β-oxidation and the TCA cycle. The correlation of NAS score with **(E)** V_{EGP}, **(F)** TCA cycle turnover (V_{CS}), **(G)** endogenous ketogenesis ($V_{RaβOHB}$), **(H)** total fat oxidation, and **(I)** reducing equivalent (RE) turnover. The correlation of liver fibrosis score with **(J)** total fat oxidation, **(K)** RE production rate, **(L)** the ratio of anaplerosis relative to TCA cycle turnover (V_{PEPCK} / V_{CS}), and **(M)** the ratio of TCA cycle turnover relative to total glucose production (V_{CS} / V_{EGP}). Data are expressed as either mean ± standard deviation or shown as correlations. Pearson correlations coefficients (r) are shown on each group along with a line of best fit calculated using linear regression. Correlations were accepted as significant if $p<0.05$. P values are shown on each graph.

Figure 3. Ketogenic insufficiency induced by loss of hepatocyte HMGCS2 impairs fasting ketosis and provokes a non-physiological rise in L-βOHB in HFCR-diet-fed mice. (A)

Ketogenesis-null mice were generated by deleting 3-hydroxymethylglutaryl-CoA synthase 2 (HMGCS2) in hepatocytes, the rate limiting step of ketone body production, by crossing homozygous *Hmgcs2*^{flox/flox} mice with transgenic heterozygous mice expressing Cre recombinase driven under the albumin promoter. Mice were maintained on standard chow diet then age-matched littermate control male wild-type (WT) and HMGCS2 knock out (HMGCS2-Liver-KO) mice were switched to a high-fat carbohydrate-restricted (HFCR) diet for ~1 week, and studied either in the random-fed or fasted state. **(B)** Validation of functional loss of HMGCS2 was confirmed *in vivo* in standard-chow fed mice by demonstrating HMGCS2-Liver-KO mice failed to increase total ketone bodies (TKB) during a 24 hour fast (left), marked by a decrease in both AcAc and βOHB (right, 4hr fasted) (n=4-8/group). **(C)** Ratio of βOHB:AcAc in 4hr fasted WT and HMGCS2-Liver-KO mice fed a standard chow diet (n=4-8/group). Wild-type mice were fed a high-fat carbohydrate-restricted (HFCR) diet for one week, then fasted for 4 hours. **(D)** AcAc, βOHB and TKBs in chow or HFCR-fed mice after 4 hours fasting (n=6-8/group). **(E)** The ratio of βOHB:AcAc in chow or HFCR-fed mice after 4 hours fasting (n=6-8/group). **(F)** Serum TKB analysis in littermate control and HMGCS2-Liver-KO mice fed a HFCR-diet (n=4-6/group). **(G)** The ratio of βOHB:AcAc in littermate control and HMGCS2-Liver-KO mice fed a HFCR-diet (n=4-6/group). **(H)** Absolute and **(I)** relative enantiomeric distribution of the serum βOHB pool in littermate control and HMGCS2-Liver-KO mice fed a HFCR-diet (n=2-3/group). **(J)** Absolute and **(K)** relative TKB analysis of intrahepatic ketones from littermate control and HMGCS2-Liver-KO mice fed a HFCR-diet and fasted for 4 hours (n=4-8/group). Data are expressed as mean ± standard deviation. Statistical differences were determined by Student's t-tests and accepted as significant is p<0.05. * p<0.05, ** p<0.01, ***p<0.001, as indicated. NS = not statistically significant.

Figure 4. Loss of hepatocyte HMGCS2 induces hepatomegaly, liver steatosis and relative hypoglycemia in HFCR diet-fed mice. Littermate control (WT) and HMGCS2-Liver-KO mice

maintained on standard chow diet were switched to high-fat carbohydrate-restricted (HFCR) diet for ~1 week. **(A)** Net change in body weight after switching to HFCR diet (n=7/group). **(B)** Total liver weight (grams), and relative liver weight (% of body weight) (n=3-8/group). Total liver triacylglycerols (TAG) in random-fed livers quantified **(C)** colorimetrically, and shown **(D)** histologically. **(E)** Total liver TAGs in random-fed livers of HFCR diet fed mice, after 1 week of refeeding standard chow (n=4/group). Systemic physiological markers of whole-body metabolism in **(F)** random-fed and **(G)** 4-hour fasted mice, including: blood non-esterified fatty acids (NEFAs), TAGs, and glucose. Also included is total liver glycogen content. (n=4-11/group). Data are expressed as mean ± standard deviation. Statistical differences were determined by Student's t-tests and accepted as significant is $p<0.05$. * $p<0.05$, ** $p<0.01$, ***$p<0.001$, as indicated. NS = not statistically significant.

Figure 5. Ketogenic insufficiency diminishes gluconeogenesis in HFCR diet-fed mice, and disrupts hepatic energy metabolism. (A) Overview of *in vivo* hepatic flux modeling study design. *(Top)* Fluxes were measured in ~16-week-old female littermate control (WT) and HMGCS2-Liver-KO mice. Five days after surgical implantation of indwelling cannulas into the carotid artery and jugular vein, mice were switched from standard chow to a HFCR diet, then fluxes were measured after a two-day acclimation period. *(Middle)* Multiple stable isotope tracers were administered including [6,6-^{2}H$_2$] glucose, [U-^{13}C$_3$] propionate, and heavy water (^{2}H$_2$O). Red blood cells from donor mice were also infused to maintain blood volume. Samples collected for flux analysis are during the final 30 minutes after metabolic and isotopic steady-state has been reached. *(Bottom)* Diagram of reaction velocities (V) (i.e., *fluxes*) modeled using a mass isotopomer distribution analysis of derivatized plasma glucose acquired via GC-MS. **(B)** Rates of whole-body endogenous glucose production (V_{EGP}) at metabolic- and isotopic-steady state measured via a tracer dilution of [6,6-^{2}H$_2$] glucose. >90% of whole-body EGP comes from the liver, thus V_{EGP} ~ hepatic glucose production. (n=4-6/group). **(C)** Absolute rates of glucose

sourcing pathways including: glycogenolysis (V_{PYGL}), total gluconeogenesis (V_{Aldo}), and rates of glycerol flux to glucose (V_{GK}) and phosphoenolpyruvate (PEP) flux to glucose (V_{Enol}). (n=4-6/group). **(D)** Absolute rates for key hepatic reactions in the oxidative flux network shown in panel A including: lactate dehydrogenase (V_{LDH}), PEP carboxykinase (PCK) (V_{PCK}) (i.e., total TCA cycle cataplerosis), anaplerosis into the TCA cycle via pyruvate carboxylase (PC) (V_{PC}), anaplerosis into the TCA cycle via propionyl-CoA carboxylase (PCC) (V_{PCC}), pyruvate cycling as the sum of pyruvate kinase (PK) and malic enzyme (ME) (V_{PK+ME}), citrate synthase flux (V_{CS}) (i.e., TCA cycle turnover) and succinate dehydrogenase flux (V_{SDH}) (n=4-6/group). **(E)** The ratio of total anaplerosis relative to TCA cycle turnover (V_{PCK}/ V_{CS}), and the ratio of TCA cycle turnover relative to total glucose production (V_{CS} / V_{EGP}) (n=4-6/group). The pool size of various molecules that are central to hepatic energy metabolism were quantified in livers of HFCR fed fasted mice including **(F)** acetyl-CoA, succinyl-CoA and propionyl-CoA. Data are expressed as mean ± standard deviation. Statistical differences were determined by Student's t-tests and accepted as significant is p<0.05. * p<0.05, ** p<0.01, ***p<0.001, as indicated. NS = not statistically significant.

Figure 6. Ketogenic insufficiency impairs hepatic fat oxidation. (A-C) Prior studies have demonstrated that ketogenesis-insufficiency induced by loss of HMGCS2 predisposes the liver to high fat diet-induced injury, coupled with an accumulation of mitochondrial acetyl-CoA and acceleration of the TCA cycle. Livers of scrambled control antisense oligonucleotide (ASO) or mouse Hmgcs2 ASO were perfused *ex vivo* with Krebs-Henseleit buffer supplemented with 0.2 mM octanoate, 1.5 mM sodium lactate, 0.15 mM sodium pyruvate, 0.25 mM glycerol, 0.2 mM [U-$^{13}C_3$] propionate and 3% ^2H_2O, then quantified oxidative fluxes using stable isotope tracing and formal flux modeling. First fat oxidation was studied in livers of chow fed mice. **(B)** Total fat oxidation quantified as the summation of (2 x ketogenesis) + TCA cycle turnover in perfused livers from control and Hmgcs2 ASO mice maintained on a standard chow diet (n=10-11/group).

(C) Reducing equivalent (RE) turnover (NADH + FADH$_2$) broken down into RE's generated from gluconeogenesis (GNG), β-oxidation, and the TCA cycle, in perfused livers from control and Hmgcs2 ASO mice maintained on a standard chow diet (n=10-11/group). **(D-F)** Fat oxidation was studied in ex vivo perfused livers of ASO-treated mice maintained on a 40% high-fat high-sucrose western diet (WD) for 8 weeks. **(D)** TCA cycle turnover in perfused livers from control ASO and Hmgcs2 ASO mice maintained on WD (n=5-6/group). **(E)** Total fat oxidation quantified as the summation of (2 x ketogenesis) + TCA cycle turnover in perfused livers from control and Hmgcs2 ASO mice maintained WD (n=5-6/group). **(F)** RE production rate (NADH + FADH$_2$) broken down into RE's generated from GNG, β-oxidation, and the TCA cycle, in perfused livers from control and Hmgcs2 ASO mice maintained on WD (n=5-6/group). **(G)** Mitochondrial Diagnostics Workflow. Liver mitochondria isolated by differential centrifugation and fueled by 50 uM palmitoylcarnitine+5 mM α-ketoglutarate, were used to assess the effects of Hymeglusin (HG) versus its vehicle control (VCTRL) on βOHB production, respiratory kinetics, and redox potential during a creatine kinase (CK) energetic clamp. Parallel experiments were performed to obtain post-CK clamp mitochondrial pellets and supernatant for LC/MS metabolite profiling. **(H)** Relative pool size of 3-hydroxymethylglutaryl-CoA (HMG-CoA), AcAc and βOHB (BHB) in mitochondria, and βOHB production measured in supernatant collected after the CK clamp bioenergetics assays (n=4/group). **(I)** Respiration (JO$_2$) plotted as a function of energy demand (ΔG$_{ATP}$ (kcals/mol)) (n=4/group). Respiratory sensitivity, or JO$_2$ conductance, was calculated by measuring the change in JO$_2$ against the change in energy demand (ΔG$_{ATP}$). Results of Student's t-test of slopes between VCTRL and HG is shown on the graph. **(J)** Relative pool size of short chain acyl-CoA species, and **(K)** free coenzyme A in mitochondria (n=4/group). **(L)** Redox potential ((NAD(P)H % reduction) plotted as a function of energy demand (ΔG$_{ATP}$ (kcals/mol)) (n=4/group). Relative pool sizes shown are from mitochondria after incubation at a fixed energy demand (ΔG$_{ATP}$ = -14.26 kcals/mol). Data are expressed as mean ± standard deviation. Statistical differences were determined by Student's t-tests or two-way ANOVA and

accepted as significant if p<0.05. * p<0.05, ** p<0.01, ***p<0.001, as indicated. # = significant

by 2-way ANOVA (p <0.05). NS = not statistically significant.

Figure 7. Loss of hepatocyte BDH1 impairs total hepatic fat oxidation but does not exacerbate HFD-induced liver fibrosis. (A) Hepatocyte-specific BDH1-null mice were generated by crossing homozygous *Bdh1*[flox/flox] mice with transgenic heterozygous mice expressing Cre recombinase driven under the albumin promoter. Mice were maintained on standard chow diet then age-matched male littermate control (WT) and BDH-Liver-KO mice were switched to a 40% high-fat high-sucrose WD for ~16 weeks, after which oxidation fluxes were studied in *ex vivo* perfused livers using a mixture of long-chain fatty acids (LCFAs) bound to BSA. **(B)** Total ketone body production *(left)* is normal in livers of BDH-Liver-KO mice, however, only AcAc and not βOHB is produced *(right)* (n=10-11/group). **(C)** Total hepatic glucose production (V_{HGP}) and **(D)** the sourcing of glucose in LCFA perfused livers from littermate control and BDH-Liver-KO mice (n=10-11/group). **(E)** Absolute rates of oxidative pathways including: pyruvate cycling (V_{PK+ME}) *(left),* total anaplerosis and cataplerosis (V_{PEPCK}) *(middle)*, and TCA cycle turnover (V_{CS}) *(right)* in LCFA perfused livers from WT and BDH1 KO mice (n=10-11/group). **(F)** Total fat oxidation calculated at (2 x ketogenesis) + acetogenesis + TCA cycle turnover (n=10-11/group). **(G)** Reducing equivalent (RE) turnover (NADH + $FADH_2$) broken down into RE's generated from gluconeogenesis (GNG) *(purple bar)*, β-oxidation *(white bar)*, and the TCA cycle *(blue bar)*, in LCFA perfused livers from WT and BDH1 KO mice. **(H)** Liver H and E and picrosirius histological staining of livers of littermate control and BDH-Liver-KO mice maintained on 40% WD. **(I)** Relative gene expression for fibrotic gene markers measured via RT-qPCR in livers of WT and BDH1 KO mice fed a 40% WD (n=11-12/group). **(J)** Absolute levels of lipid peroxide 4-hydroxyalkenal species in livers of littermate control and BDH-Liver-KO mice fed a 40% WD (n=10-11/group). Data are expressed as mean ± standard deviation. Statistical differences were determined by Student's t-tests and accepted as

significant is p<0.05. * p<0.05, ** p<0.01, ***p<0.001, as indicated. NS = not statistically

significant.

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

Fat Oxidation in Chow-fed HMGCS2-null perfused livers

Fat Oxidation in Western Diet-fed HMGCS2-null perfused livers

Inhibiting HMGCS2 Impairs Fat Oxidation in isolated mitochondria

Figure 7.