Single-step in vitro ribosome reconstitution mediated by two GTPase factors, EngA

and ObgE

Aya Sato[1,3], Weng Yu Lai[1,2,3], Yusuke Sakai[1,4,5], *Yoshihiro Shimizu[1]

[1]Laboratory for Cell-Free Protein Synthesis, RIKEN Center for Biosystems Dynamics

research (BDR), Kobe, Hyogo 650-0047, Japan.

[2]Graduate School of Frontier Biosciences, Osaka University, Osaka 565-0871, Japan

[3]These authors contributed equally to this study.

[4]Laboratory of Nucleic Acid Nanotechnology, Institute for Quantitative Biosciences, The

University of Tokyo, Tokyo, Japan

[5]Department of Medical Biochemistry and Biophysics, Karolinska Institutet, Stockholm,

Sweden

*Correspondence should be addressed to:

Yoshihiro Shimizu, Ph.D.

Laboratory for Cell-free Protein Synthesis, RIKEN Center for Biosystems Dynamics

(BDR), IIB Bldg. S304, 6-7-1, Minatojima-minamimachi, Chuo-ku, Kobe, Hyogo 650-

0047, Japan

Tel: +81-78-569-8878

Email: yshimizu@riken.jp

Abstract

When bacterial ribosomes are assembled in vitro, manipulation of incubation temperature and magnesium ion concentration have been an essential procedure, which is a crucial step for the assembly of active large subunits. The present study tackles with this issue to develop a single-step procedure, which can be performed in a physiological condition. We found that GTPase factors EngA and ObgE can complement the changes in temperature and magnesium ion concentrations. In the presence of these factors, ribosome assembly can proceed under physiological conditions, with magnesium ion concentrations below 10 mM, potassium concentrations around 100 mM, and temperatures of 37 °C. Both the ribosome assembly and translation processes were successfully integrated in the reconstituted cell-free protein synthesis system. Furthermore, we found that these GTPase factors can reassemble the ribosomes to an active state, whose structure was disrupted by EDTA chelation of magnesium ions, indicating that these two factors can reversibly induce the ribosome structure to an intact state. The findings are essential for the bottom-up construction of synthetic cells.

Introduction

Ribosomes are large intracellular complexes that play a central role in a protein translation system. They consist of two subunits, each composed of one or a few rRNAs and dozens of ribosomal proteins. Maturation processes of these complex assemblies have been studied for decades. For the bacterial ribosomes, the construction of assembly maps of both subunits from *Escherichia coli*, describing the order of ribosomal proteins bind to rRNA, is well known as milestone achievements[1,2]. However, many aspects have yet to be elucidated, including differences between the comparatively slow in vitro reconstitution processes and the more efficient and refined cellular assembly processes, accompanied inclusion of various assembly factors, and effects of a variety of rRNA modifications on these maturation processes. Various in vivo and in vitro studies are still being performed, aiming for a deeper understanding of these processes, by incorporating advanced technologies, including quantitative mass spectrometry (qMS), cryo-electron microscopy (cryo-EM), and single molecule imaging[3-8].

Recently, with the increasing focus on constructing synthetic cells from the bottom-up viewpoint[9,10], the importance of the studies on assembling ribosomes from individual parts in vitro has been highlighted. The integrated synthesis, assembly, and translation (iSAT) method developed by Jewett *et al.* demonstrated that in the presence of cell extracts, the assembly of both subunits proceeds using transcribed rRNAs and ribosomal proteins purified from native ribosomes, termed total proteins of the 30S subunit (TP30) and total proteins of the 50S subunit (TP50)[11]. Cell extracts-free ribosome assembly methods have been developed by the integration with the reconstituted cell-free protein synthesis system (PURE system)[12], enabling the in vitro evolution of 16S rRNA[13]. Also, TP-free methods have been constructed by preparing individually purified recombinant ribosomal proteins for both subunits[14-17]. The use of cell-free expressed ribosomal proteins is also addressed[8,16,18-20], aiming for the development of self-replicating synthetic cells in the future[21].

These cell-free synthetic biology studies can help us understand the minimally required building blocks for producing functional assemblies. For the small subunit, functionally active subunits have been shown to be assembled by merely mixing synthetic parts, transcribed rRNA and recombinant ribosomal proteins with minimal post-translational modifications[16]. The functional large subunit has also been shown to be assembled with recombinant proteins with minimal post-translational modifications[17]. However, two major issues exist with the large subunit reconstitution method.

The first issue is the requirement of posttranscriptional modifications in 23S rRNA. According to Green and Noller, modified nucleotides in certain regions of rRNA are

necessary for the peptidyl transferase activity of the reconstituted large subunit from *E.*
coli[22]. For *Bacillus stearothermophilus*, it has been shown that the subunit with the
peptidyl transferase activity can be reconstituted even with the transcribed 23S rRNA[23].
The addition of some chemical substances into the assembly mixtures for *E. coli*
ribosomes has also been reported to be effective for the peptidyl transferase activity of
the reconstituted subunit using transcribed 23S rRNA[24]. All of these experiments have
focused on the peptidyl transferase activity of the reconstituted subunit. Whether or not
the assembled subunits are competent for the whole translation processes, including
initiation, elongation, and termination steps, remains unclear. By contrast, protein
synthesis activity of the reconstituted large subunit was observed if the iSAT method is
applied[11]. It can be considered that some factors in the cell extract, including rRNA
modification enzymes, are involved in the functional assembly of the large subunit.

The second issue is the need for a two-step procedure that involves temperature and
ion concentration changes in the reaction solution. Nierhaus and Dohme revealed that two
significant steps are required for the successful assembly of the large subunit[25,26]. In the
first step, 5S rRNA, 23S rRNA, and ribosomal proteins are incubated at 44 °C to form
41S or 48S intermediates. Then, by raising the magnesium ion concentration from 4 mM
to 20 mM and the temperature to 50 °C, active subunits formation is completed by further
incubation (**Fig. 1a**). The need to raise the temperature to a non-physiological value and
to manipulate the ion concentrations can be a major barrier when attempting the bottom-
up construction of synthetic cells. It is necessary to develop the assembly methods under
physiological conditions without temperature changes or ion concentration manipulations.
It is noteworthy that the second issue also does not apply to the iSAT method. In the
method, the ribosome assembly and the translation processes are seamlessly linked in the
same reaction solution, without manipulation of the temperature and ion concentrations[11].
It is highly likely that specific factors in the cell extracts complement this issue.

The present study aimed to tackle with the second issue by using ribosome biogenesis
factors. It was found that changes in temperatures are effective and changes in magnesium
ion concentration are essential for the assembly of translation competent ribosomes. We
found that the presence of GTPase factors EngA[27] and ObgE[28,29] is necessary to
complement these changes. In the presence of these factors, ribosome assembly can
proceed under physiological conditions, with magnesium ion concentrations below 10
mM, potassium concentrations around 100 mM, and temperatures of 37 °C. Both the
ribosome assembly and translation processes were successfully linked together in the
PURE system, demonstrating that the ribosome assembly proceeds in the same solution
conditions as the translation reaction, and the assembled ribosomes are competent for the

whole translation processes. Furthermore, it was revealed that unfolded ribosomes, whose structure was disrupted by EDTA chelation of magnesium ions[30], could be restored to an active state with both the supplementation of magnesium ions and the addition of EngA and ObgE, indicating that these two factors can reversibly induce the ribosome structure to an intact state.

Results

Effect of changes in temperature and ion concentration on the ribosome assembly

In our previous study on small subunits, we have found that the assembly proceeds autonomously without the support of any factors[16], whereas temperature and ion concentration changes are necessary for the assembly of large subunits[25]. To evaluate the impact of these changes on the reconstitution of entire ribosomes, we performed in vitro ribosome assembly experiment using total ribosomal RNAs (5S, 16S, and 23S rRNA) and total proteins of the 70S ribosome (TP70) prepared from the purified native ribosomes[31]. Four patterns of temperature and magnesium ion concentration changes were applied to clarify each effect (**Fig. 1b**). The reconstituted ribosomes were purified from the assembly mixtures and then they were added into the PURE system without the ribosome to test superfolder green fluorescent protein (sfGFP) synthesis. When no magnesium ion concentration changes were applied, sfGFP synthesis by the assembled ribosomes was not detected, and when no temperature changes were applied, sfGFP synthesis decreased by about one-third (**Fig. 1b**). The results showed that the changes in magnesium ions are particularly essential, indicating that the two-step procedure established for the reconstitution of large subunits[25] (**Fig. 1a**) is also an essential key process for the assembly of entire ribosomes with translation activity, consistent with the previous statement[31].

Single-step ribosome assembly using ribosome biogenesis factors

Based on the differences between the iSAT assembly experiments[11] and those using purified parts, including this study (**Fig. 1b**), we hypothesized that some factors in the cell extracts used in iSAT, particularly ribosome biogenesis factors, may complement the changes in temperature and divalent ion concentration required in the two-step procedure, and facilitate a single-step assembly in the cell-free reaction mixtures (**Fig. 2a**). Furthermore, in previous experiments examining the effects of ribosome biogenesis factors on small subunit reconstitution, we found that the factors Era and RsgA (YjeQ) were effective in promoting subunit assembly, both being GTPase factors[15,16]. Therefore, we focused on GTPase factors, which have been suggested to be involved in large subunit

biogenesis. Based on literature surveys, we selected BipA[32], EngA (Der)[27], EngB
(YihA/YsxC)[33], HflX[34], LepA[35], and ObgE[28,29], which were prepared as recombinant
proteins. Surprisingly, when these six factors were mixed with total ribosomal RNAs (5S,
16S, and 23S rRNA) and TP70, along with an energy regeneration system to maintain
GTPase activity, we found that it was possible to assemble the ribosomes with sfGFP
synthesis activity at a constant but relatively low magnesium concentration of 8 mM at
37 °C (**Fig. 2b**). The sfGFP synthesis activity significantly dropped at a magnesium
concentration of 17 mM in the presence of these six factors. Almost no sfGFP synthesis
activity was observed in the absence of six factors, even with 8 mM magnesium,
suggesting the selected factors exhibit some activity that compensates for the magnesium
ion concentration changes in the two-step procedure. The optimal magnesium
concentration for this reaction was 8 mM (**Fig. 3a**), and the optimal monovalent salt
concentration for ribosome assembly was several hundred millimolar, consistent with
previous assembly experiments[25] (**Fig. 3b**).
162
Coupling of the assembly and translation reactions
The results with GTPase factors suggested that the factors were able to complement the
temperature and ion concentration changes in the two-step procedure, realizing a single-
step ribosome assembly. Because the required monovalent and divalent ion
concentrations are within the range for the optimum conditions of the translation reactions
(**Fig. 3**), it appeared probable to integrate the ribosome assembly and translation processes
in a single reaction mixture. To this end, we attempted to construct a coupled reaction, by
adding 3 rRNAs, total proteins of the 70S ribosome (TP70), and selected GTPases into
the ribosome-omitted PURE system (**Fig. 4a**). The result showed that after a lag period
that likely corresponds to the time required for the ribosome assembly, increase of sfGFP
fluorescence was observed (**Fig. 4b**). An optimal magnesium concentration was 8 mM
(**Fig. 4b, c**), consistent with the large subunit assembly experiment (**Fig. 3a**). The
synthesis of sfGFP was unaffected by a monovalent salt concentration above 100 mM
(**Fig. 4d**). While high salt concentrations were effective for large subunit assembly (**Fig.
3b**), they may have negatively impacted translation reactions. An empirically translation-
competent salt concentration of 100-250 mM[36-38] was found to be optimal. The results
demonstrated that under physiological conditions suitable for translation reactions, a
single-step ribosome assembly has proceeded in the solution, where the assembled
ribosomes are competent for the whole translation processes, resulting in sfGFP synthesis.
182
EngA and ObgE are responsible for the single-step ribosome assembly process

We examined which of the selected six factors plays a role in facilitating the single-step
ribosome assembly. Since more than one factor could be involved in the process, we
addressed the activity of ribosomes assembled using GTPase mixtures with each factor
removed one by one. When tested in the coupled reaction system, sfGFP synthesis activity
significantly decreased in the absence of ObgE and was not detected in the absence of
EngA (**Fig. 5a**). Similar results were obtained in an uncoupled system (**Fig. 5b**),
suggesting that EngA and ObgE are the responsible factors which may complement the
changes in temperature and ion concentration in the two-step protocol for large subunit
assembly. Furthermore, when only ObgE was added to the coupled system, sfGFP
synthesis was not detected, similar to the case without any factors, and when only EngA
was added, comparatively slow sfGFP synthesis was observed (**Fig. 5c**). When both
factors were added, sfGFP synthesis was highly active, indicating that ribosome assembly
was achieved with the presence of EngA and ObgE.
197
EngA and ObgE are required for the reassembly of unfolded ribosomes
It has been shown that adding EDTA to ribosomes collapses their structure to a level
detectable by electron microscopy[30]. The rRNA used in this study was extracted from
native ribosomes with phenol, and it is possible that the partial structure, formed
depending on magnesium ion, was still retained, potentially reducing ObgE dependency.
We added EDTA to native ribosomes to collapse their structure and then restored the
environment by adding magnesium ions. We tested whether ribosome activity would be
restored in the presence of ObgE and EngA. The results showed that active ribosomes
were formed only in the presence of both factors (**Fig. 6**), suggesting that both ObgE and
EngA are equally necessary for reassembling ribosomes.
208
Discussion
In the present study, the change in divalent ion concentration was crucial for the
translationally active ribosome assembly (**Fig. 1**). In contrast, in the presence of GTPase
factors EngA and ObgE, ribosome assembly and translation reactions could be coupled
under constant magnesium ion concentrations (8 mM), indicating the significant role of
these two factors under physiological conditions (**Fig. 2-5**). Although the dependency on
ObgE was slightly mitigated in these experiments, both factors were indispensable for the
reversible unfolding and regeneration of ribosomes by manipulation of EDTA and
magnesium ion concentrations (**Fig. 6**), suggesting that both of two factors are essential
in ribosome biogenesis, compensating for the divalent ion concentration and temperature
changes in the conventional in vitro assembly conditions. It is possible that some tertiary

structures completed by ObgE catalysis might have been retained in phenol-extracted
rRNA prepared from native ribosomes.

The two GTPases identified in this study belong to the 13 highly conserved GTPases
in bacteria, implying their significant importance for cell viability[39]. GTPases EngB, HflX,
and LepA, examined in this study, also belong to the 13 members and have been suggested
to be involved in ribosome assembly[33-35], whereas no significant activity was detected in
the present study. This does not deny their involvement in ribosome assembly under
different assembly conditions, and further verification is necessary. We note that an
important finding regarding HflX function, which is different from ribosome assembly
pathway, has been found very recently[40]. BipA is not included in the 13 members but is
reported as a homolog of EF-G and LepA and reported to be involved in the ribosome
assembly process[32], and so it was also tested in this study.

The finding that the two GTPases compensate for the changes in temperature and
divalent ion concentration provides new insights into the 50S subunit assembly process.
These changes are known to be required for late-stage assembly processes[26] and therefore,
it is suggested that the two factors catalyze the late-stage assembly processes. Indeed,
cryo-EM based studies have suggested the involvement of these two factors in the late-
stage assembly processes[5]. The structures of *E. coli* 50S subunits with EngA[27] and ObgE[29]
have been individually elucidated, and structural analysis of 50S subunits in the assembly
process obtained by affinity purification with tagged ObgE has been performed[28].
Additionally, in the cryo-EM study on the assembly process of the 50S subunit of *Bacillus*
subtilis, a deep relationship with YphC, a homolog of EngA, was demonstrated[33]. These
reports suggest the involvement of the two factors in the maturation of critical structures
in the late-stage process, including functional core (FC) and peptidyl transferase center
(PTC), and the binding of ribosomal proteins involved in the late-stage processes like
uL16 and bL36. The present study clarifies the deep relationship between these critical
maturation steps in the late-stage processes and the essential operations in the in vitro
assembly process identified by Nierhaus and Dohme[25].

In addition to the structural analyses, genetic studies have suggested the involvement
of both proteins in the 50S assembly process[41,42]. Interestingly, both proteins have been
reported as suppressor proteins of cold-sensitive RlmE (RrmJ) knockout mutants[43]. RlmE
is an enzyme that methylates U2552 in domain V of 23S rRNA, where PTC is located.
The 50S subunits obtained from the RlmE-knockout mutants show some translational
defects and structural differences, particularly in the FC and PTC regions, compared to
the wild-type subunits[44]. Since the cell growth of RlmE knockout mutants restores with
increased expression of EngA and ObgE[43], the structural differences in ribosomes may

also be recovered by these factors. This suggests that one of the roles of these two factors
includes the maturation of these regions, and their roles might overlap.

However, the present analysis shows that active ribosomes are not assembled with
ObgE alone, and the activity of ribosomes assembled with EngA alone is significantly
lower than when both factors are used (**Fig. 5**). Furthermore, the regeneration experiment
of ribosome structure by EDTA chelation of magnesium ions and subsequent restoration
with magnesium ions revealed that the presence of both factors is essential (**Fig. 6**). This
suggests that these two factors individually play different roles in the late-stage assembly
process. There has been no structural study focusing on both EngA and ObgE at the same
time. According to the individual cryo-EM studies, a C-terminal GTPase domain of ObgE
binds to the GTPase-associated center (GAC), and its N-terminal domain binds to the
interface region towards the PTC from the A site space[28,29]. In contrast, an N-terminal
GTPase domain (GD1) of EngA binds near the L1 stalk, its C-terminal GTPase domain
(GD2) binds to the P site space, and its C-terminal KH domain binds to the interface
region towards the PTC[27]. The N-terminal domain of ObgE and the C-terminal region of
EngA, which is not fully observed with electron density maps, are very close, suggesting
that they might not bind to the subunit simultaneously. Moreover, the complex obtained
by affinity purification with tagged ObgE shows additional electron density maps
corresponding to YjgA and RluD, which may prevent EngA binding[28]. Therefore, ObgE
and EngA might independently support the late-stage assembly processes in different
steps, suggesting that at least two critical maturation steps in the late-stage assembly
processes exist. Further structural analysis will elucidate the details of these processes.

Unlike gram-negative bacteria, represented by *E. coli*, gram-positive bacteria,
represented by *B. subtilis,* might require a different set of GTPases. Previous reports
showed that intermediate 45S or 44.5S subunits accumulate in cells lacking YphC (a
homolog of EngA), RbgA, and YsxC (a homolog of EngB)[33,45]. These intermediates lack
ribosomal proteins binding in the late-stage processes, such as uL16 and bL36, suggesting
stalling before critical maturation steps. Although *B. subtilis* contains a gene annotated as
obgE, this protein does not appear in a series of studies, while RbgA, which has no
corresponding protein in *E. coli*, and YsxC, a homolog of EngB, are essential. This
suggests that diversity in support systems for critical maturation steps has arisen during
evolution.

The cryo-EM structure of 50S subunits during the assembly process, which are
affinity-purified with tagged ObgE, suggested the presence of three factors that may work
cooperatively with ObgE, including RsfS, YjgA, and RluD[28]. By contrast, the assembly
process progresses well only with ObgE and EngA in the present study. Several scenarios

can explain these differences. First, it can be considered that the factors identified in the
cryo-EM analysis might be supportive but not essential, consistent with the fact that the
genes for these factors are not essential for the cell viability. As another possibility, their
involvement in rRNA folding can be considered. In the present study, we use rRNA
extracted from native ribosomes (**Fig. 1-5**) or unfolded ribosomes with EDTA (**Fig. 6**).
The requirement of ObgE is slightly different between these experiments, suggesting
different degrees of rRNA folding in both preparations, which might dependent on the
presence of magnesium ions. Moreover, it can be considered that secondary structures
formed by simple complementary duplex formation are still preserved in both
preparations. In contrast, newly transcribed rRNA is produced without any secondary and
tertiary structures. We have attempted the 50S subunit assembly with transcribed rRNAs
but failed to achieve active subunits with the two factors (data not shown), suggesting
further multifaceted mechanisms might be required for the formation of the 50S subunits.
They might include rRNA modifications and associated rRNA folding, and the protein
group identified by the cryo-EM study may function in such processes. Further in-depth
analysis based on the developed assay system will clarify the overall assembly process of
the 50S subunits.

The present study reveals that the process corresponding to the changes in temperature
and magnesium concentration, identified by Nierhaus and Dohme[25], is catalyzed by two
GTPases. In reactions using cell extracts like iSAT[11], the presence of these two factors in
the extracts may have facilitated the ribosome assembly without any changes in
temperature and ion concentration. While improvement of the iSAT is extensively
performed[46-48], the addition of two factors identified in this study may contribute to such
studies. The two GTPases may also be one of the key factors for the coupling systems
using cell-free systems for expressing ribosomal proteins[18,20], as well as the future
developments towards self-replicating synthetic cells[9,10]. At the same time, investigation
of the mechanisms of actions of the two GTPases, through structural analysis using cryo-
EM, protein composition analysis using qMS, and dynamic analysis using imaging
techniques will lead to a deeper understanding of ribosome assembly mechanisms.

Methods
Sample preparation for in vitro experiments. Native ribosomes and the PURE system
components were prepared as described previously[49]. Preparation of TP70 was performed
according to the conventional methods[31]. Total ribosomal RNAs containing 5S, 16S, and
23S rRNAs Ribosomal RNAs were prepared from the native ribosomes via water-
saturated phenol extraction followed by 2-propanol precipitation, according to the same

literature[31]. Ribosome biogenesis factors used in this study were prepared as follows. *E.*
coli genes for BipA, EngB, HflX, and LepA were cloned into pET15b (Merck Millipore,
USA) as His-tag fusion protein. The genes for EngA and ObgE were cloned into pET15b
(Merck Millipore, USA) as small ubiquitin-like modifier (SUMO) protein-fusion proteins
where His-tag, SUMO protein, and EngA or ObgE were tandemly arranged. Detailed
sequences are described in the supplementary data (**Supplementary Data 1**). BipA,
EngB, HflX, and LepA proteins were purified in a same way as His-tagged translation
factors[49]. EngA and ObgE were purified in a same way as uS2 according to our previous
study[15]. DNA templates for the sfGFP synthesis was prepared as previously described[16].
337

Ribosome assembly with a two-step procedure. Reaction mixtures (25 µL) contained
20 mM Tris-HCl, pH 7.4, 4 mM $Mg(OAc)_2$, 400 mM NH_4Cl, 7 mM 2-mercaptoethanol,
22.5 pmol total rRNAs, and 27 pmol TP70. Reactions were carried out at 44 °C for 20
min and then 1 µL of 400 mM $Mg(OAc)_2$ solution or water was added to change the
magnesium ion concentration. The mixtures were further incubated at 44 °C or 50 °C for
90 min. The mixtures were centrifugated at 5,000 *g* for 5 min at 25 °C and then
supernatants were recovered. The assembled ribosomes were concentrated and buffer
exchanged using 0.5 mL centrifugal filter units MWCO 30 kDa with ribosome buffer (10
mM Hepes-KOH, pH 7.6, 10 mM $Mg(OAc)_2$, 30 mM K(OAc), and 1 mM DTT). The
activity of ribosomes was monitored in the home-made PURE system[49], where reaction
mixtures (20 µL) containing the assembled ribosomes from 25 µL mixtures and 5 nM
DNA template encoding sfGFP were incubated at 37 °C in Mx3005P (Agilent
Technologies, USA), and sfGFP fluorescence was monitored during incubation.
351

Ribosome assembly with ribosome biogenesis factors. The assembly reactions were
performed based on the PURE system reaction mixtures[49], in anticipation of future
integration with a cell-free translation system. The reaction mixtures (25 µL) contained
50 mM Hepes-KOH, pH7.6, 2 mM spermidine, 1 mM DTT, 2 mM ATP, 2 mM GTP, 1
mM UTP, 1 mM CTP, 10 mM creatine phosphate, 0.1 mM each 20 amino acids, 0.25 µg
each of creatine kinase, nucleotide diphosphate kinase, BipA, EngA, EngB, HflX, LepA,
and ObgE, varying concentrations of $Mg(OAc)_2$ and potassium glutamate, 22.5 pmol total
rRNAs, and 27 pmol TP70. The mixtures were incubated at 37 °C for 110 min and they
were centrifugated at 5,000 *g* for 5 min at 25 °C, followed by the recovery of supernatants.
The assembled ribosomes were concentrated and buffer exchanged using 0.5 mL
centrifugal filter units MWCO 30 kDa with ribosome buffer. The activity of ribosomes
was monitored in the home-made PURE system[49], where reaction mixtures (20 µL)

containing the assembled ribosomes from 25 µL mixtures and 5 nM DNA template encoding sfGFP were incubated at 37 °C in Mx3005P (Agilent Technologies, USA), and sfGFP fluorescence was monitored during incubation.

Integration of ribosome assembly and translation systems. Reaction mixtures (20 µL) were based on the home-made PURE system[49] without ribosomes. In addition to the standard components, 0.25 µg each of BipA, EngA, EngB, HflX, LepA, and ObgE, 22.5 pmol total rRNAs, and 27 pmol TP70 were added in to the mixtures. Concentrations of concentrations of $Mg(OAc)_2$ and potassium glutamate are varied according to the experiments and designated in the figure legends. After the addition of 5 nM DNA template encoding sfGFP, the mixtures were incubated at 37 °C in Mx3005P (Agilent Technologies, USA), and sfGFP fluorescence was monitored during incubation.

Reassembly of unfolded ribosomes. First, 0.5 µL of 5 µM ribosome (0.5 pmol) was mixed with 1 µL of 100 mM EDTA (final concentration was 67 mM) and incubated on ice for 30 min. Then, PURE translation mixtures were prepared. In this experiment, we used solution II (Enzyme Mix) of PUREfrex 2.0 (GeneFrontier Corporation) instead of using home-made translation factors and enzymes, whereas other components were based on the home-made PURE system[49]. In addition to the standard components, the mixtures (10 µL) contained 1.5 µL ribosome solution (0.5 pmol in 67 mM EDTA), 0.5 µL of solution II of PUREfrex 2.0, 0.2 µg each of EngA and ObgE. The concentration of potassium glutamate was 150 mM and the concentration of $Mg(OAc)_2$ was 32 mM to restore the actual magnesium ion concentration. After the addition of 5 nM DNA template encoding sfGFP, the mixtures were incubated at 37 °C in Mx3005P (Agilent Technologies, USA), and sfGFP fluorescence was monitored during incubation.

Acknowledgments
This work was supported by CREST (JPMJCR20S4 to YShimizu) from Japan Science and Technology Agency (JST), a Grant-in-Aid (20H05701 to YShimizu) from the Japan Society for the Promotion of Science (JSPS), the Human Frontier Science Program (RGP0043/2017 to YShimizu), and an intramural Grant-in-Aid from the RIKEN Center for Biosystems Dynamics Research (to YShimizu).

Author Contributions
Y.Shimizu. designed the study. A.S., W.Y.L., Y.Sakai. and Y.Shimizu. wrote the manuscript. A.S., W.Y.L., and Y.Sakai performed the experiments. All authors discussed

400 the results and commented on the manuscript.

Competing interests

The authors declare no competing interests.

References

1. Mizushima, S. & Nomura, M. Assembly mapping of 30S ribosomal proteins from *E. coli*. *Nature* **226**, 1214-1218 (1970).

2. Röhl, R. & Nierhaus, K. H. Assembly map of the large subunit (50S) of *Escherichia coli* ribosomes. *Proc Natl Acad Sci U S A* **79**, 729–733 (1982).

3. Shajani, Z., Sykes, M. T. & Williamson, J. R. Assembly of bacterial ribosomes. *Annu Rev Biochem* **80**, 501–526 (2011).

4. Davis, J. H. & Williamson, J. R. Structure and dynamics of bacterial ribosome biogenesis. *Philos Trans R Soc Lond B Biol Sci* **372**, 20160181 (2017).

5. Seffouh, A., Nikolay, R. & Ortega, J. Critical steps in the assembly process of the bacterial 50S ribosomal subunit. *Nucleic Acids Res* **52**, 4111–4123 (2024).

6. Duss, O., Stepanyuk, G. A., Puglisi, J. D. & Williamson, J. R. Transient Protein-RNA Interactions Guide Nascent Ribosomal RNA Folding. *Cell* **179**, 1357-1369.e16 (2019).

7. Rodgers, M. L. & Woodson, S. A. Transcription Increases the Cooperativity of Ribonucleoprotein Assembly. *Cell* **179**, 1370-1381.e12 (2019).

8. Levy, M., Falkovich, R., Daube, S. S. & Bar-Ziv, R. H. Autonomous synthesis and assembly of a ribosomal subunit on a chip. *Sci Adv* **6**, eaaz6020 (2020).

9. Dogterom, M., Kamat, N. P., Jewett, M. C. & Adamala, K. P. Progress in Engineering Synthetic Cells and Cell-Free Systems. *ACS Synth Biol* **13**, 695–696 (2024).

10. De Capitani, J. & Mutschler, H. The Long Road to a Synthetic Self-Replicating Central Dogma. *Biochemistry* **62**, 1221–1232 (2023).

11. Jewett, M. C., Fritz, B. R., Timmerman, L. E. & Church, G. M. In vitro integration of ribosomal RNA synthesis, ribosome assembly, and translation. *Mol Syst Biol* **9**, 678 (2013).

12. Shimizu, Y. *et al.* Cell-free translation reconstituted with purified components. *Nat Biotechnol* **19**, 751–755 (2001).

13. Murase, Y., Nakanishi, H., Tsuji, G., Sunami, T. & Ichihashi, N. In Vitro Evolution of Unmodified 16S rRNA for Simple Ribosome Reconstitution. *ACS Synth Biol* **7**, 576–583 (2018).

436   14.   Culver, G. M. & Noller, H. F. Efficient reconstitution of functional *Escherichia*
437   *coli* 30S ribosomal subunits from a complete set of recombinant small subunit
438   ribosomal proteins. *RNA* **5**, 832–843 (1999).

439   15.   Tamaru, D., Amikura, K., Shimizu, Y., Nierhaus, K. H. & Ueda, T. Reconstitution
440   of 30S ribosomal subunits in vitro using ribosome biogenesis factors. *RNA* **24**,
441   1512–1519 (2018).

442   16.   Shimojo, M. *et al.* In vitro reconstitution of functional small ribosomal subunit
443   assembly for comprehensive analysis of ribosomal elements in *E. coli*. *Commun*
444   *Biol* **3**, 142 (2020).

445   17.   Aoyama, R. *et al.* In vitro reconstitution of the *Escherichia coli* 70S ribosome
446   with a full set of recombinant ribosomal proteins. *J Biochem* **171**, 227–237 (2022).

447   18.   Li, J. *et al.* Cogenerating Synthetic Parts toward a Self-Replicating System. *ACS*
448   *Synth Biol* **6**, 1327–1336 (2017).

449   19.   Levy, M. *et al.* Boundary-Free Ribosome Compartmentalization by Gene
450   Expression on a Surface. *ACS Synth Biol* **10**, 609–619 (2021).

451   20.   Kosaka, Y. *et al.* Reconstitution of ribosome self-replication outside a living cell.
452   *biorxiv* doi: 10.1101/2022.08.29.5056922 (2022).

453   21.   Laohakunakorn, N. *et al.* Bottom-Up Construction of Complex Biomolecular
454   Systems With Cell-Free Synthetic Biology. *Front Bioeng Biotechnol* **8**, 213
455   (2020).

456   22.   Green, R. & Noller, H. F. In vitro complementation analysis localizes 23S rRNA
457   posttranscriptional modifications that are required for *Escherichia coli* 50S
458   ribosomal subunit assembly and function. *RNA* **2**, 1011–1021 (1996).

459   23.   Green, R. & Noller, H. F. Reconstitution of functional 50S ribosomes from in vitro
460   transcripts of *Bacillus stearothermophilus* 23S rRNA. *Biochemistry* **38**, 1772–
461   1779 (1999).

462   24.   Semrad, K. & Green, R. Osmolytes stimulate the reconstitution of functional 50S
463   ribosomes from in vitro transcripts of *Escherichia coli* 23S rRNA. *RNA* **8**, 401–
464   411 (2002).

465   25.   Nierhaus, K. H. & Dohme, F. Total reconstitution of functionally active 50S
466   ribosomal subunits from *Escherichia coli*. *Proc Natl Acad Sci U S A* **71**, 4713–
467   4717 (1974).

468   26.   Nikolay, R. *et al.* Structural Visualization of the Formation and Activation of the
469   50S Ribosomal Subunit during In Vitro Reconstitution. *Mol Cell* **70**, 881-893.e3
470   (2018).

27.   Zhang, X. *et al.* Structural insights into the function of a unique tandem GTPase

EngA in bacterial ribosome assembly. *Nucleic Acids Res* **42**, 13430–13439 (2014).

28.   Nikolay, R. *et al.* Snapshots of native pre-50S ribosomes reveal a biogenesis factor

network and evolutionary specialization. *Mol Cell* **81**, 1200-1215.e9 (2021).

29.   Feng, B. *et al.* Structural and functional insights into the mode of action of a

universally conserved Obg GTPase. *PLoS Biol* **12**, e1001866 (2014).

30.   King, T. C., Rucinsky, T., Schlessinger, D. & Milanovich, F. *Escherichia coli*

ribosome unfolding in low Mg^{2+} solutions observed by laser Raman spectroscopy

and electron microscopy. *Nucleic Acids Res* **9**, 647–661 (1981).

31.   Nierhaus, K. H. Reconstitution of ribosomes in Ribosomes and protein synthesis:

A Practical Approach (Spedding G. eds.) 161-189, IRL Press at Oxford University

Press, New York (1990).

32.   Goh, K. J. *et al.* Translational GTPase BipA Is Involved in the Maturation of a

Large Subunit of Bacterial Ribosome at Suboptimal Temperature. *Front Microbiol*

12, 686049 (2021).

33.   Ni, X. *et al.* YphC and YsxC GTPases assist the maturation of the central

protuberance, GTPase associated region and functional core of the 50S ribosomal

subunit. *Nucleic Acids Res* **44**, 8442–8455 (2016).

34.   Polkinghorne, A. *et al.* Chlamydophila pneumoniae HflX belongs to an

uncharacterized family of conserved GTPases and associates with the *Escherichia*

coli 50S large ribosomal subunit. *Microbiology (Reading)* **154**, 3537–3546 (2008).

35.   Gibbs, M. R. *et al.* Conserved GTPase LepA (Elongation Factor 4) functions in

biogenesis of the 30S subunit of the 70S ribosome. *Proc Natl Acad Sci U S A* **114**,

980–985 (2017).

36.   Kim, D. M., Kigawa, T., Choi, C. Y. & Yokoyama, S. A highly efficient cell-free

protein synthesis system from *Escherichia coli*. *Eur J Biochem* **239**, 881–886

(1996).

37.   Kigawa, T. *et al.* Cell-free production and stable-isotope labeling of milligram

quantities of proteins. *FEBS Lett* **442**, 15–19 (1999).

38.   Sun, Z. Z. *et al.* Protocols for implementing an *Escherichia coli* based TX-TL cell-

free expression system for synthetic biology. *J Vis Exp* e50762 (2013).

39.   Verstraeten, N., Fauvart, M., Versées, W. & Michiels, J. The universally conserved

prokaryotic GTPases. *Microbiol Mol Biol Rev* **75**, 507–542 (2011).

40.   Seely, S. M., Basu, R. S. & Gagnon, M. G. Mechanistic insights into the

alternative ribosome recycling by HflXr. *Nucleic Acids Res* **52**, 4053–4066 (2024).

41.   Sato, A. *et al.* The GTP binding protein Obg homolog ObgE is involved in

ribosome maturation. *Genes Cells* **10**, 393–408 (2005).

42. Jiang, M. *et al.* The Escherichia coli GTPase CgtAE is involved in late steps of
large ribosome assembly. *J Bacteriol* **188**, 6757–6770 (2006).
43. Tan, J., Jakob, U. & Bardwell, J. C. A. Overexpression of two different GTPases
rescues a null mutation in a heat-induced rRNA methyltransferase. *J Bacteriol*
184, 2692–2698 (2002).
44. Wang, W. *et al.* Loss of a single methylation in 23S rRNA delays 50S assembly at
multiple late stages and impairs translation initiation and elongation. *Proc Natl*
Acad Sci U S A **117**, 15609–15619 (2020).
45. Seffouh, A. *et al.* Structural consequences of the interaction of RbgA with a 50S
ribosomal subunit assembly intermediate. *Nucleic Acids Res* **47**, 10414–10425
(2019).
46. Liu, Y., Fritz, B. R., Anderson, M. J., Schoborg, J. A. & Jewett, M. C.
Characterizing and alleviating substrate limitations for improved in vitro
ribosome construction. *ACS Synth Biol* **4**, 454–462 (2015).
47. Fritz, B. R., Jamil, O. K. & Jewett, M. C. Implications of macromolecular
crowding and reducing conditions for in vitro ribosome construction. *Nucleic*
Acids Res **43**, 4774–4784 (2015).
48. Caschera, F. *et al.* High-Throughput Optimization Cycle of a Cell-Free Ribosome
Assembly and Protein Synthesis System. *ACS Synth Biol* **7**, 2841–2853 (2018).
49. Shimizu, Y. & Ueda, T. PURE Technology. *Methods Mol Biol* **607**, 11–21 (2010).
528
529

Figures

Figure 1. Ribosome assembly with two-step procedure. (a) Schematic of the two-step
procedure identified by Nierhaus and Dohme. In the step 1, comparatively low
magnesium ion concentration and low incubation temperature are required to assemble
the intermediate state of the 50S subunit (41S/48S particle), whereas in the step 2,
increasing the incubation temperature and magnesium ion concentration are essential for
the assembly of translationally active 50S subunit. (b) Individual effect of temperature
and magnesium concentration changes on the activity of assembled ribosomes.

Figure 2. Ribosome assembly with ribosome biogenesis factors. (a) Working
hypothesis of this study. We hypothesized that some of the ribosome biogenesis factors
take over the role of temperature and magnesium concentration changes in the two-step
procedure. We selected six GTPase factors as candidates of these factors. (b) Single-step
ribosome assembly by selected ribosome biogenesis factors. Activity of assembled
ribosomes in the presence or absence of six factors at 8 or 17 mM magnesium ion
concentration at 37 °C. Concentration of potassium glutamate for the ribosome assembly
experiment was fixed at 200 mM.

Figure 3. Effect of magnesium and potassium concentrations on the single-step ribosome assembly using six GTPase factors. (a) Effect of magnesium concentration on the assembled ribosome activity. (b) Effect of potassium concentration on the assembled ribosome activity. Relative endpoint sfGFP fluorescence at 4 h timepoint, compared to 8 mM magnesium or 300 mM potassium, respectively, is shown. Potassium concentration was fixed at 200 mM (a) and magnesium concentration was fixed at 8 mM (b).

567
568

Figure 4. Coupling of the assembly and translation reactions. (a) Schematic of the
coupled reaction. Three rRNAs (5S, 16S, and 23S) and TP70 are added into the ribosome-
free PURE system. Ribosomes are assembled in the PURE system with a help of ribosome
biogenesis factors and then, assembled ribosomes exhibit the translation activity, which
can be detected as sfGFP fluorescence. (b) Time course of sfGFP fluorescence in the
coupled reaction system. Effect of magnesium concentration is shown. (c) Effect of
magnesium concentration on the assembled ribosome activity. (d) Effect of potassium
concentration on the assembled ribosome activity. Relative endpoint sfGFP fluorescence
at 8 h timepoint, compared to 8 mM magnesium or 150 mM potassium, respectively, is
shown. Potassium concentration was fixed at 150 mM (b, c) and magnesium
concentration was fixed at 8 mM (d).

Figure 5. Identification of the responsible factors. (a) Ribosome activities assembled
in the absence of each factor in the coupled system. (b) Ribosome activities assembled in
the absence of each factor in the uncoupled system. (c) Ribosome activities assembled
with only EngA and ObgE. When coupled system is applied, magnesium concentration
was fixed at 8 mM and potassium concentration was fixed at 150 mM (a, c). When
uncoupled system was used, magnesium concentration was fixed at 8 mM and potassium
concentration was fixed at 250 mM for the ribosome assembly reaction (b). Relative
endpoint sfGFP fluorescence at 8 h timepoint, compared to that using all six factors (a, b)
or time course of sfGFP fluorescence (c) are shown.

597 **Figure 6. Reassembly of unfolded ribosomes using EngA and ObgE.** Ribosomes were
598 unfolded by the addition of high concentration of EDTA and then restored the
599 environment by the addition of magnesium ions. Followingly, reassembly of the ribosome
600 was addressed by using EngA and ObgE. Time course of sfGFP fluorescence are shown
601 in the presence or absence of EngA and ObgE..

22