A generalist microbial predator shows innate attraction to more profitable prey, but does not learn from experience

P. M. Shreenidhi[1,2], Rachel I. McCabe[1], Heng Liang[1], Margaret I. Steele[1,3], David C. Queller[1], Joan E. Strassmann[1]

[1]Department of Biology, Washington University in St. Louis, St. Louis, Missouri, USA
[2]Department of Molecular Genetics, University of Toronto, Toronto, Ontario, Canada
[3]Department of Biological Sciences, Texas Tech University, Lubbock, Texas, USA

Corresponding author: P. M. Shreenidhi

Corresponding author email: shreenidhi.perukkaranai@utoronto.ca

Author contributions: P.M.S., D.C.Q., and J.E.S. designed research. P.M.S., R.I.M., H.L., and M.I.S., performed research. P.M.S., R.I.M., H.L., M.I.S., D.C.Q., and J.E.S. analyzed the data. P.M.S., D.C.Q., and J.E.S. wrote the paper with inputs from R.I.M., H.L., and M.I.S.

Competing Interest Statement: The authors declare no competing interests.

Acknowledgements

This material is based upon work supported by the National Science Foundation (DEB 1753743 and DEB 2237266). We thank Debra Brock for inputs during the development of the project. Robert Insall provided insight on short versus long-range chemoattraction. We also thank members of the Queller-Strassmann lab for useful discussions.

1 Abstract

How do generalists adjust to variation in prey abundance and profitability and seek out their

preferred prey? We investigated this question in the soil protist *Dictyostelium discoideum*, a

generalist predator of many species of bacteria. Despite their generalist diet, amoebas

proliferate more quickly on some bacteria than on others. We tested amoeba chemoattraction

towards 23 bacterial species and found that they are generally more attracted to the more

profitable prey bacteria. Naïve amoebas were also preferentially more attracted to an edible

mutant rather than the inedible wild type of a soil *Pseudomonas* isolate. These results suggest

that *D. discoideum* amoebas have an innate prey preference that is adaptive. We also tested

how experience with different prey bacteria affects chemoattraction in amoebas. Given the

huge number of bacterial species in soil, learning from experience should be advantageous.

However, we found no evidence that experience with prey bacteria affects preference. Our

results suggest that generalist amoebas are innately attracted to the more profitable prey

bacteria and this innate attraction cannot be overridden by recent experience.

17 Introduction

Predation, where one individual kills and consumes another, is a fundamental interaction

between organisms that can profoundly shape organismal evolution. One key component of a

predator's fitness is the ability to sense its prey. Specialist predators are often superior at

detecting their prey as compared to generalists (Bernays 2001). Since specialist predators

feed on few prey species, they need to track fewer features in their environment to quickly

and accurately detect their preferred prey. But generalist predators feed on a diversity of prey

species. This very abundance of choice may hinder their ability to detect relevant information

about specific prey.

Generalist predators, or more broadly, generalist consumers, can overcome this problem

either by innate or learned mechanisms. Generalists might possess innate preferences for

certain environmental features, such as shape or colour, that can drive foraging decisions. For

example, though honey bees, *Apis mellifera,* harvest pollen and nectar from many flower

species, they exhibit an innate preference for some colours (Giurfa et al. 1995). Similarly, the

generalist hoverfly, *Eristalinus aeneus,* which feeds on plant pollen and nectar is attracted to

artificial floral objects that are bright, radially symmetrical, and possess plant odours (Mishra

et al. 2025). However, plasticity in behaviour and the ability to learn from experience are also

critical components of foraging behaviour in generalists (Bernays 2001; Mery and Burns

2010). Learning can increase the efficiency of decision making by generalists and allow them

to track the most profitable or abundant resources. Honey bees, despite their preference for

some colours, can quickly be trained to prefer alternative colours if they are associated with

greater rewards (Giurfa et al. 1995). Also, many avian predators learn to avoid chemically

defended insects based on their warning coloration (Guilford 1990).

Prey preference has been extensively studied in macroscopic predators such as birds or

insects that rely on visual and olfactory cues to track their prey (Guilford 1990; Dukas 2008).

These preference behaviours are encoded in the nervous system. But the ability to track prey

is crucial even to microscopic organisms that lack a nervous system. Soil protists are often

generalist predators that feed on diverse prey bacteria (Johnke et al. 2014; Amacker et al.

2022). Protist predation is a major cause of bacterial mortality in the environment (Jousset

2012). Predation by protists can shape bacterial communities, facilitate nutrient cycling and

drive the evolution of bacterial defences (Sherr and Sherr 2002; Jousset 2012; Smith et al.

2023). However, protists use fundamentally different methods for prey detection compared to

macroscopic predators.

Prey detection can be challenging for soil protists because of the complexity of soil

environments. Soil is a heterogenous environment that varies spatially and temporally in the

abundance and diversity of prey bacteria. Even a gram of soil contains extremely diverse

micro-organisms and may contain anywhere between two thousand and eight million

bacterial species (Gans et al. 2005; Schloss and Handelsman 2006). Soil is composed of

minute particles of clay, sand, and decayed plant material separated by water and air-filled

pores (Vos et al. 2013).The particles and pores lead to a patchy distribution of resources

resulting in fragmented bacterial communities (Raynaud and Nunan 2014; Negassa et al.

2015). Protists must navigate this maze-like soil habitat to seek out their preferred prey

bacteria.

Dictyostelium discoideum, also known as the social amoeba, is a microbial predator well-

known for its fascinating lifecycle which includes both unicellular and multicellular stages

(Strassmann and Queller 2011). *D. discoideum* in its simplest form persists as unicellular

amoebas which are generalist predators of soil bacteria in temperate forest soils. Prey bacteria

that have been co-isolated with *D. discoideum* are diverse and range across at least four

bacterial phyla: Proteobacteria, Actinobacteria, Firmicutes and Bacteroidetes (Brock et al.

2018). Despite being generalist predators, these amoebas divide faster when feeding on some

prey bacteria than on others (Shreenidhi et al. 2024). Under laboratory conditions, they can

double in number in three hours on some of the most edible species of bacteria but take as

long as ten hours to double on the less edible species. Some species of bacteria are plainly

inedible, while some others are pathogenic (Brock et al. 2018; Steele et al. 2023). Bacteria

possess an arsenal of defences to fend off protist predation, including secretion of toxins and

chemorepellents, modification of surface properties, and biofilm formation (Jousset 2012;

Smith et al. 2023). The edibility of a bacterium could therefore be considered a joint

phenotype (Queller 2014) of the amoeba's offense and the bacteria's defence. Thus, given the

variation in bacterial profitability and defensive responses, it would be advantageous if the

amoebas could either innately sense the suitability of a bacterium for predation from a

distance or modulate their preference based on their prior experience with a prey bacterium.

D. discoideum amoebas use chemotaxis to locate their prey bacteria (King and Insall 2009).

Amoebas are attracted to bacterially secreted cyclic adenosine monophosphate (cAMP)

(Meena and Kimmel 2017; Rashidi and Ostrowski 2019) and folic acid (Pan et al. 1972;

Lima et al. 2014). These two chemoattractants are detected using two different G-protein

coupled receptors, cAR1 and fAR1 respectively. There may of course be other, unknown

chemoattractants. *D. discoideum* amoebas are also known to be more attracted to Gram-

negative bacteria than to Gram-positive bacteria (Rashidi and Ostrowski 2019). However,

there is wide-variation in edibility among Gram-negative bacteria (Brock et al. 2018; Steele

et al. 2023; Shreenidhi et al. 2024).

In addition to innate preferences, we hypothesized that experience with prey bacteria may

play an important role in modulating prey preference in amoebas. The question of whether

unicellular organisms can learn from experience and change their behaviour through

associative conditioning was a highly disputed topic in the 20th century (Gershman et al.

2021), but there is growing evidence that single-celled protists are sometimes capable of

associative learning (Carrasco-Pujante et al. 2021). There are multiple reasons why learning

would be beneficial for *D. discoideum* as a generalist predator. Innate preferences may fail to

recognize rare species or differentiate between closely related prey that differ in edibility. To

account for this, amoebas might learn from experience to be more attracted to profitable

bacteria and to avoid unprofitable or harmful bacteria. Prey experience might also direct

amoebas towards familiar prey because of the cost of prey-switching. Amoebas experience a

reduction in population growth rate when switched from one species of prey bacteria to

another (Shreenidhi et al. 2024). These costs are likely physiological and related to changing

transcriptional responses to match the new prey bacterium. Thus, it is possible that

experience with a prey bacterium primes amoebas to be more attracted towards one

bacterium compared to others.

In this study, we measured the chemotaxis of *D. discoideum* amoebas towards 23

phylogenetically diverse species of prey bacteria that were previously isolated from close

association with *D. discoideum* from soil environments. We then tested if the chemoattraction

of amoebas was explained by the edibility of the prey bacteria. At a finer level, we also tested

whether amoebas could differentiate between an inedible *Pseudomonas* strain and an edible

transposon mutant of the same species. Next, we tested if amoebas displayed any evidence of

learning such that they are differentially attracted to a bacterium when they have prior

experience of it being profitable prey.

Methods

Population growth rate data

To test for correlation between innate attraction and population growth rate of amoebas on

different bacteria, we used previously published population growth rate data of *D. discoideum*

for 20 of the 23 bacterial species (Shreenidhi et al. 2024). We used similar assay protocols to

estimate population growth rate of *D. discoideum* on our other four bacterial species. We

estimated the Malthusian parameter **r** over 20 h on plates.

Innate prey preference experiments

We performed the innate preference experiments on AX4 amoebas that had been cultured

axenically for at least 4 days on HL5 medium including glucose (Formedium) with

antibiotics (300 µg/ml Streptomycin, 100 µg/ml Carbenicillin), free of any prey bacteria. By

culturing the amoebae axenically we avoided any effects of learned preference for recently

experienced bacteria. Amoebas were inoculated at a starting concentration between 5×10^5 –

2×10^6 cells/ml in 25 ml of HL5. The flasks were placed in a shaking incubator at 21ºC, 100

rpm. The amoebas were transferred into fresh HL5 media every 3-4 days to maintain them at

a concentration below 2×10^7 cells/ml. We collected amoebas for the experiment by

centrifuging the cultures at 300*g*, 10ºC for 3 minutes. We discarded the spent media and

washed the amoeba pellet once more in KK2 buffer (2.25 g KH_2PO_4 (Sigma-Aldrich) and

0.67 g K_2HPO_4 (Fisher Scientific) per litre). The amoeba pellet was suspended in 2 ml KK2

buffer. We estimated amoeba concentration using a haemocytometer and diluted them to a

final concentration of 10^7 cells/ml for the experiment.

The bacteria for the experiments were cultured on SM/5 (2 g glucose (Fisher Scientific), 2 g

BactoPeptone (Oxoid), 2 g yeast extract (Oxoid), 0.2 g $MgSO_4$ (Fisher Scientific), 1.9 g

KH_2PO_4 (Sigma-Aldrich), 1 g K_2HPO_4 (Fisher Scientific), 15g agar (Sigma-Aldrich) per litre

of deionized water) or LB plates (Fisher Scientific), and resuspended to a concentration of

OD_{600} 5 in KK2 buffer.

We used assay protocols similar to Rashidi and Ostrowski (2019) for the chemotaxis

experiments. The experiments were performed in 6-well plates containing 2 ml of 2% high

purity agar in KK2 buffer per well. In the centre of each well, we pipetted a 2µl droplet of the

amoeba suspension (10^7 cells/ml). A 2 µl spot of the bacterial suspension was placed at 0.65

cm on one side of the amoeba spot and a 2 µl droplet of KK2 buffer at the same distance on

the other side (Figure 1A). The spots were placed using a template to keep them identically

spaced. We measured the chemotaxis of AX4 amoebas towards 23 species of soil bacteria

that had been collected in close association with *D. discoideum,* with three technical

replicates per bacterium (Table 1). The experiments were performed over four different days.

To account for any day-specific effects, we also performed chemotaxis experiments towards

Klebsiella pneumoniae, the standard laboratory food given to *D. discoideum*, on all four days.

We imaged (phase contrast, 10x magnification) the plates at 6 h from set-up using the Leica

Thunder Live Cell Microscope with an automated stage.

Innate preference for an edible mutant

We studied the innate attraction of amoebas towards the wild-type *Pseudomonas* strain 20P

3.2.4 (20P324) and its mariner transposon mutant (20P324 6-C11) (Steele et al. 2024). This

mutant strain is susceptible to predation by *D. discoideum* amoebas, whereas the wild-type

bacterium is inedible (Steele et al. 2024). Using the same assay conditions as described

above, we performed experiments where the central spot of AX4 amoebas was flanked on

opposite sides by a spot of bacterial suspension (wild type or mutant) and KK2 buffer. We

also performed experiments where the wild type and the transposon mutant were paired

directly against each other, with the amoeba spot was flanked by both bacterial suspensions.

We repeated these experiments on three days with three technical replicates per day. We

imaged (phase contrast, 10x magnification) the plates at 6 h from set-up using the Leica

Thunder Live Cell Microscope with an automated stage.

Whole genome sequencing of edible *Pseudomonas* mutant 20P324 6-C11

We extracted Genomic DNA from 20P324 6-C11 cells using the Qiagen DNeasy Blood and

Tissue Kit (Qiagen). Illumina sequencing was performed by SeqCenter. Illumina sequencing

libraries were prepared using the Illumina DNA Prep kit with a target insert size of 280 bp

and sequenced using the Illumina NovaSeq X Plus (2x151bp paired end reads). Bcl-convert

(v4.2.4) was used for demultiplexing and adapter trimming. Geneious Prime was used to map

reads to the wild type reference genome (NCBI RefSeq assembly GCF_025466075.1), call

variants, and exclude variants in low coverage regions. The 20P324 6-C11 genome was

assembled using Unicycler (v0.4.9) (Wick et al. 2017), and annotated using the Rapid

Annotation using Subsystem Technology (RAST) server (Aziz et al. 2008; Overbeek et al.

2014; Brettin et al. 2015). The mutant was determined to have three mariner transposon

insertions not present in the wild type (previously sequenced), which were located in the

gacA gene, *cyoE* gene, and an in intergenic region between tRNA-dihydrouridine synthase

and a gene encoding a hypothetical protein. The edible phenotype is likely caused by the

disruption of the *gacA* gene, which positively regulates secondary metabolism (Stallforth et

al. 2013).

Learned prey preference experiment

We studied learned prey preference in a wild strain of *D. discoideum,* QS9, isolated from

Mountain Lake Biological Station, Virginia in 2000 and then stored in a -80 °C freezer

(Fortunato et al. 2003). To test how experience with a prey bacterium affects prey preference,

we first "familiarized" the amoebas to that bacterium. We chose eight bacteria on which *D.*

discoideum has a high population growth rate so that the familiar bacterium provides a

positive stimulus. To culture amoebas, we spread 200 µl of 1.5 OD_{600} bacterial suspension on

SM/5 plates and pipetted 50 µl of *D. discoideum* spore solution containing 200,000 spores.

The amoebas were collected from the resulting vegetative front after 3 - 4 days with a sterile

loop. For some bacterial species where the front was hard to see, we collected amoebas from

the entire plate after ~ 36 - 40 hours. We suspended the amoebas in KK2 buffer and

centrifuged them at 300*g*, 10 ºC for 3 minutes. We washed the amoeba pellet three more

times to remove any bacteria. We used the same chemotaxis set-up as described in the

"Innate prey preference experiments", with a central spot of amoebas flanked by two spots of

potential attractants. But here the amoebas were given a choice between the bacterium they

were cultured on ("familiar bacterium") and a new bacterium ("novel bacterium") (Figure

1B). We did this experiment with eight bacterial species in a full factorial design. Amoebas

cultured on each of eight familiar bacteria were tested against seven novel bacteria (Table 1,

bacterial species highlighted in green). All experiments for amoebas conditioned to a given

familiar bacterium were done on the same day, with three technical replicates for every

choice between familiar and novel bacterium. We imaged the plates at both 4 h and 6 h from

set-up.

Image Analysis

For all chemotaxis experiments, we used ImageJ version 1.53 (Schneider et al. 2012) to

estimate the number of amoebas that moved out of the amoeba spot towards each of the two

potential attractants. We drew a line tangential to the amoeba spot at a point that was closest

to each potential attractant to define a boundary and counted only those amoebas that moved

past this boundary (Figure 1A), as was previously done (Rashidi and Ostrowski 2019). We

did the following image processing steps: *Convert to 8-bit > Set Threshold > Make Binary >*

Fill holes > Apply Watershed. We manually counted the cells for the innate preference

experiments because AX4 amoebas are irregularly shaped and harder to count using

automation. We used the *Analyze particles* feature on the relevant parts of the image to count

the number of cells for the learned preference experiments that was performed on QS9.

Amoebas had not reached the bacteria in any of these experiments, so attraction could not

have been caused by feeding amoebas.

Data analysis & Statistics

Innate prey preference experiments

We performed all statistical analysis in R (version 4.2.1) (R Core Team 2022). To estimate

net amoeba movement towards a given bacterium, we subtracted amoeba movement towards

KK2 buffer from amoeba movement towards the bacterial spot. To account for day-specific

effects (Figure S1), we scaled all amoeba movement estimated on a given day by dividing it

by amoeba movement towards *Klebsiella pneumoniae* (averaged across 3 replicates) on that

day. We refer to these values corrected for both buffer and day effects as normalized amoeba

movement.

To investigate the relationship between innate preference of amoebas and edibility of the

bacterium, we needed to control for the phylogenetic non-independence of data points. We

used a phylogeny including all 23 species of soil bacteria used in our experiments as a

random factor in our models to control for this. These bacteria are a subset of a larger

collection of bacterial strains isolated from close association with *D. discoideum* (Brock et al.

2018). We reconstructed the phylogeny in Brock et al. 2018 by using the previously

published 16s sequences of the bacteria (deposited by Brock et al. on GenBank) and 26

species of anchor bacteria (full length 16s sequences downloaded from SILVA rRNA

database). A multiple sequence alignment of all sequences was generated using MUSCLE on

Geneious Prime. IQ-TREE version 2.3.6 was used to construct a maximum-likelihood tree

with model optimization and 100 bootstraps to assess branch support (Minh et al. 2020). We

pruned the tree to the 23 species of bacteria used in chemotaxis experiments using iTOL

(Letunic and Bork 2021) and used this tree in newick format for all downstream analysis. We

forced the phylogenetic tree to be ultrametric using *chronos* function (lambda = 1) in *ape*

package in R (Paradis and Schliep 2019) and used *inverseA* function in *MCMCglmm* package

(Hadfield 2010) to convert the phylogeny into a covariance matrix.

We then used the R package *MCMCglmm* (Generalized linear-mixed model models using

Markov Chain Monte Carlo approach) to examine the relationship between normalized

amoeba movement (response variable, henceforth referred to as innate preference) and mean

population growth rate of amoebas (fixed effect), controlling for phylogenetic relationships

among the bacteria by modelling the bacterial phylogeny as a random effect. We used the

default prior for the fixed effect and weakly informative priors for residual variance and

random effects (V=1, nu = 0.002) to not overly influence posterior distributions. We report

the pMCMC values and the R^2 for the fixed effect calculated as described using Nakagawa

and Schielzeth (2013).

To test for chemokinesis (stimulation of undirected movement), we looked for correlations

between average movement towards buffer (response variable) and average movement

towards bacterium (fixed effect), scaled as before by average movement towards *Klebsiella*

pneumoniae on a given day. We used a MCMCglmm to model this with Bacterial phylogeny

as the random factor.

Innate preference to *Pseudomonas* edibility mutants

To estimate net amoeba movement towards either the wildtype or mutant *Pseudomonas* in

buffer experiments, we subtracted amoeba movement towards KK2 buffer from amoeba

movement towards the bacterial spot. To test whether net amoeba movement significantly

differed based on bacterial identity, we used a Linear Mixed Model (LMM) with bacterial

identity (Wild type or Mutant) as a fixed effect, Day as a random effect (Day1, Day2, Day3)

and net amoeba movement as the response variable using the R package *lmerTest*

(Kuznetsova et al. 2017). To test if amoeba were differentially attracted to either wild-type or

mutant bacterium when presented with both, we first calculated proportional amoeba

movement to a given bacterium. Next, we tested if arcsine square root transformed

proportions were significantly different from 0.5 using a linear mixed model with Day as a

random effect.

Learned preference experiments

We counted the number of amoebas moving towards the familiar bacterium or novel

bacterium in each replicate at 4 h and 6 h from set-up. We used the average from these two

time points for each replicate for downstream analysis. Replicates with zero amoeba

movement were removed from the analysis (2 reps), as they interfere with calculation of

proportions. We define $f_{i,j}$ as the proportional movement of amoebas towards familiar

bacterium i, when paired with novel bacterium j, averaged over all three replicates.

We tested if the overall proportional movement (collection of $f_{i,j}$'s) was significantly

different from 0.5 using a linear mixed effects model with familiar bacterium identity

modelled as random effect. We also tested if proportional movement towards familiar

bacterium (collection of $f_{i,j}$'s for a given i) was dependent on familiar bacterium identity by

using a linear model with familiar bacterium identity as fixed effect and arcsine square root

transformed proportional movement data as response variable. We used the *emmeans*

package to calculate the estimated marginal means and confidence intervals (Lenth et al.

2022).

For a given bacterium *i*, we have two proportional amoeba movement measures, one when

the bacterium is familiar (F_i), and another when the bacterium is novel (N_i). F_i, the estimated

proportional movement towards bacterium *i* when familiar, averaged over all seven novel

bacteria, is calculated as:

318
$$F_i = \frac{\sum_{j \neq i} f_{i,j}}{7}$$

319   N_i, the estimated proportional movement towards bacterium *i* when novel, averaged over all

320   seven familiar bacteria, is calculated as:

$$N_i = \frac{\sum_{j \neq i}(1 - f_{j,i})}{7}$$

To test if learned preference of amoebas is correlated with innate preference, we built a MCMCglmm model with innate preference towards prey bacterium (data from the innate preference assay) as fixed effect and average proportional movement towards bacterium when familiar (F_i) as the response variable.

To account for innate attractiveness of bacteria when testing if experience with a prey affects chemoattraction of amoebas, we calculated F_i (average movement when familiar) - N_i (average movement when novel) for all 8 species of bacteria and tested if this was significantly different from zero using MCMCglmm. Since these data are also collected across different species of bacteria, we added in the phylogeny of the 8 bacterial species assayed as a random effect. We followed the same steps as described previously to construct the phylogeny, except we pruned the phylogenetic tree down to the 8 species of bacteria used in the learned preference experiments. We used the default priors for the fixed effect and weakly informative prior for residual variance and random effects (V=1, nu = 0.002) to not overly influence posterior distributions. We report the pMCMC values and the R^2 for the fixed effect calculated as described using Nakagawa & Schielzeth (2013).

Results

Amoebas preferentially detect and direct movement towards better prey.

To estimate innate prey preference in *D. discoideum* amoebas, we measured the movement of naïve amoebas towards 23 species of soil bacteria or towards KK2 buffer (Figure 1A, Table 1). Amoeba movement was generally directed towards the bacteria and rarely towards buffer

(Figure S2). As our measure of innate preference, we subtracted movement towards KK2

from movement towards bacteria and scaled all amoeba movement by movement towards

Klebsiella penumoniae on the same day to get normalized net amoeba movement towards

different species of bacteria (Figure S3). To test if innate response was adaptive, we tested

the relationship between normalized movement and population growth rate of amoebas on

these bacteria, correcting for the bacterial phylogeny. We found that normalized movement

increases with increasing population growth rate (Figure 2; MCMCglmm, Fixed effect

posterior mean = 4.58, 95% CI (1.00, 8.11), pMCMC = 0.011, fixed effect R^2 = 0.21). Thus,

amoebas possess an adaptive innate response where they tend to be innately attracted to more

profitable bacteria. We found no evidence that this effect dependent on Gram status of the

bacteria (Figure S4).

While chemotaxis is the directed movement of amoebas in response to a chemical gradient,

chemokinesis refers to random undirected movement in response to chemical cues. If some

bacteria release chemokines that increase amoeba movement overall, then we might observe

an increase in amoeba movement in all directions (though presumably weaker at the far end

of the amoeba spot). To test if chemokinesis can explain the observed movement patterns, we

tested if movement towards KK2 buffer was correlated with movement towards bacteria. We

found no significant relationship between these two measures suggesting that these patterns

are not driven by chemokinesis (Figure S5: MCMCglmm, Fixed effect posterior mean = 0.06,

95% CI (-0.01, 0.13), pMCMC = 0.12).

Naïve amoebas are more attracted to an edible mutant over the inedible wild type of a

***Pseudomonas* strain**

To further test the idea that innate preference of amoebas correlates with bacterial edibility,

we measured the chemotaxis of axenically cultured AX4 amoebas towards an inedible

Pseudomonas soil isolate (20P 3.2.4) and its edible transposon mutant strain (20P324 6-C11).

We found that net amoeba movement towards the edible mutant was greater than net amoeba

movement (corrected for movement towards buffer) towards the inedible wild type (Figure

3A: Estimate LMM: Estimated net movement (Mutant – WT) = 16.1), t_{14} = 2.38, p-value =

0.03). When amoebas were presented with both bacterial strains simultaneously, we found

that the proportion of amoebas that moved towards the edible mutant was significantly

greater than 0.5 (Figure 3B: Estimated proportional movement = 0.79, 95 % CI (0.53,0.94), t_8

= 3.373, p-value = 0.009).

Disruption of the *gacA* gene might be the cause of differential attraction of the amoebas

towards *Pseudomonas* bacteria

Whole genome sequencing of *Pseudomonas* 20P 3.2.4 and the mutant revealed transposon

insertions in two genes, *gacA* and *cyoE*, and in an intergenic region. The *cyoE* gene is

involved in aerobic respiration, while the *gacA* gene is a member of the two-component

signal transduction system, GacA/GacS. GacA regulates the transcription of hundreds of

genes, including genes involved in the production secondary metabolites that are important

for defence against bacterial competitors, plant pathogens and protist predators (Jousset et al.

2009; Hassan et al. 2010; Stallforth et al. 2013). The disruption of the *gacA* gene likely

affects the production of many secondary metabolites and is likely to be the cause of

increased attraction of amoebas towards the mutant *Pseudomonas*.

Amoebas do not show a preference for familiar prey

Because the innate effect explains 21% of the variance, there is plenty of scope for improvement via learned effects. To test if prior experience with prey bacteria affects preference in amoebas, amoebas were familiarized to different bacteria and then given a choice between the bacterium they were cultured on (familiar bacterium) or a new bacterium (novel bacterium) (Figure 1B). We chose eight bacteria on which *D. discoideum* has a high population growth rate so that the familiar bacterium provides a positive stimulus, and tested whether amoebas would learn and be more attracted to this good prey. There was no overall trend of preference for the familiar bacterium (Fig 4A: p = 0.476).

However, we did find wide variation in proportional attraction of amoebas towards the bacterium that depended on bacterial identity (Figure 4A: Familiar bacterium identity: $F_{8,48}$ = 2.81, p-value = 0.012). For example, when amoebas were familiarized to *Pseudomonas sp.* 14P 8.1.1(*Pk*), they seemingly prefer the familiar bacterium (*Pk*) over the novel bacterium (Estimated marginal means = 0.13, 95% CI (-0.006, 0.27), p = 0.06). However, when amoebas were familiarized to *Variovorax sp.* 20P 10.2.4 (*Vb*), they preferred the novel bacterium on average over the familiar bacterium (*Vb*) (Estimated marginal means = -0.18, 95% CI (-0.32, -0.04), p = 0.01).

We found that this pattern of variation in attraction towards the familiar bacterium may be explained by the innate attraction of the amoebas towards the bacterium (Figure 4B; MCMCglmm, Fixed effect posterior mean = 0.16, 95% CI (-0.02, 0.36), pMCMC = 0.07, R^2 = 0.28). The pattern is not quite significant but, to be conservative, we tested if removing the estimated effects of innate preference affects our results. Because our experiments employ a

fully factorial design, each prey bacterium appears in both familiar and novel treatments.

Therefore, we accounted for innate preference for the bacterium by comparing the average

movement towards bacterium when it is familiar to average movement towards the bacterium

when it is novel ($F_{i} - N_{j}$) (because innate attraction would be part of both F_i and N_j and cancel

out). The difference between the average familiar and novel proportions was not different

from zero and the trend is in the direction opposite to learning (Figure 4C; MCMCglmm,

Fixed effect posterior mean = -0.08, 95% CI (-0.19, 0.04), pMCMC = 0.151).

Discussion

All predators must find prey, but generalists that have many suitable prey may have the most

difficulty evolving rules to identify them and distinguish degrees of suitability. This may be

especially true for microbial predators that encounter many potential bacterial prey species

and lack nervous systems to process the information. In this paper, we tested for both innate

and learned prey preferences of *D. discoideum* amoebas towards their prey bacteria, as

measured by chemoattraction. We found that naïve amoebas are more attracted, on average,

to prey bacteria that are more profitable. Thus, innate prey preferences of *D. discoideum*

amoebas are generally adaptive.

We also found that such innate preferences can depend on the presence or loss of a single

gene in the bacterium. Naive amoebas were more attracted to an edible mutant of the 20P

3.2.4 *Pseudomonas* strain compared to the inedible wild type. The mutant differed at three

sites but the likely causal mutation was a transposon insertion in the *gacA* gene. This gene

positively regulates secondary metabolism, and is important for predation resistance in many

species of *Pseudomonas* including the *Pseudomonas* strain 20P 3.2.4 (Stallforth et al. 2013;

Inglis et al. 2018; Steele et al. 2024). 20P 3.2.4 is predicted to have 13 secondary metabolite

biosynthetic gene clusters based on analysis with antiSMASH (Steele et al., 2023). Targeted

deletion of the *gacA* gene in 20P 3.2.4 is known to increase susceptibility to predation by

amoebas similar to the increase seen in the transposon mutant (Steele et al. 2024). It is

possible that secondary metabolites that confer predation resistance also act as

chemorepellents. Thus, bacteria may be signalling their unsuitability for predation using

some secondary metabolites much like the way many insects signal their unsuitability as prey

using warning colouration. It is also possible that *gacA* downregulates production of the

normal chemoattractants. Either way, the result suggests modulating chemoattraction is a

novel function of *gacA*.

The positive relationship between innate preference and prey profitability hints at the long

evolutionary history of amoeba-bacteria interactions and how these innate responses have

been selected to enhance fitness over evolutionary time. But as with any correlation we

cannot establish the direction of causation between innate preference and prey profitability. It

seems likely that variation in amoeba performance on different bacteria shaped prey

preferences. But it is also possible that amoebas improved their performance on bacteria that

were already attractive.

Our data shows that only 21 % of variance in amoeba attraction is explained by prey

profitability suggesting that there are other constraints on this response. For example, *D.*

discoideum amoebas use chemotaxis not only to locate their bacterial prey, but also to locate

other conspecific amoebas during starvation to initiate the multicellular cycle. Thus, innate

chemoattraction towards bacterial prey cannot be changed to the extent that it hinders the

ability of the amoebas to find other conspecifics. Indeed, cAMP is a common chemoattractant

signal used by *D. discoideum* to detect both bacterial prey and conspecifics using the

chemoreceptor cAR1 (Klein et al. 1988; Meena and Kimmel 2017; Rashidi and Ostrowski

2019). A greater constraint on optimal amoeba preferences is evolution of the prey species.

Because prey bacteria are constantly evolving, it seems impossible to evolve innate

preferences that remain adaptive on all encountered bacterial species. Prey would presumably

be under selection to become "invisible" by reducing secretion of attractants, and might also

evolve mimicry of secretions of unattractive bacteria. Thus, whatever the reasons for the

imperfection of innate attraction, there would likely be considerable advantage to adjust

preferences via learning.

However, we found that prior experience with a prey species did not generally seem to

influence subsequent attraction of *D. discoideum* amoebas to that bacterium. Because these

were all profitable prey species, we expected amoebas might learn to be more attracted after

experiencing them, but they did not. The idea that amoebas can change their prey preference

based on experience rests on the assumption that the prey bacteria secrete different

chemoattractants, or combinations of chemoattractants, and that amoebas can distinguish

among them, probably by using different chemoreceptors or combinations of chemoreceptors.

Experience with a prey bacterium could then lead to upregulation of those specific

chemoreceptors, or to give greater weight to information from those chemoreceptors, thus

reinforcing preference. The only prey chemoattractants of *D. discoideum* that are well

understood are folate and cAMP (Artemenko et al. 2014). The response to these

chemoattractants is mediated by two highly-ligand specific G-protein coupled receptors

(GPCRs): the cAR1 receptor for cAMP and the fAR1(GrlL) receptor for folate (Pan et al.

2016; Meena and Kimmel 2017). But amoebas possess 61 putative receptors in the GPCR

family (Heidel et al. 2011). Many of these potential GPCRs are of unknown function and

some could be undiscovered chemoreceptors that play a role in modulating responses based

on experience. All that would be needed is a GPCR with a different ligand binding domain to

activate the signalling machinery that stimulates actin remodelling to trigger chemotaxis.

Though we observed no consistent change in amoeba response based on experience with

prey, variation in amoeba movement in learning experiments was partly explained by their

innate preference for a specific bacterium. It is possible that the bacteria assayed in our

experiments do not secrete sufficiently different chemoattractants, or that amoebas only use

folate and cAMP to detect their prey bacteria. We know that amoeba attraction towards these

two chemoattractants is dose-dependent where higher doses of the attractant lead to more

amoeba movement (Meena and Kimmel 2017; Rashidi and Ostrowski 2019). Thus, it is

possible that the prey bacteria we tested vary in the levels of cAMP or folate production and

that amoebas simply prefer prey bacteria that produce more cAMP or folate, irrespective of

prey experience. However, the lack of observed learning may have explanations other than an

absence of multiple chemoattractants.

The levels of environmental heterogeneity experienced by an organism play an important role

in the evolution of innate and learned responses (Mery and Burns 2010). Innate behavioural

responses evolve when the environment is relatively stable over several generations, while

plastic/learned responses evolve when the environment changes rapidly over the organism's

lifetime (Stephens 1991). The presence of an innate adaptive prey preferences and the lack of

evidence for learning in our experiments suggests that perhaps amoebas do not experience

very much environmental heterogeneity in prey species. This seems surprising given the

diversity of bacteria in the soil but the micro-distribution of prey species is worth further

investigation.

The importance of self-generated gradients in long-range movement may be a more likely

explanation for why we observed no learning. Chemotactic gradients established solely by

the diffusion of attractants from a local source are only effective at guiding directional

movement of cells over short distances (500 μm) (Tweedy et al. 2016). The gradient might be

too shallow to establish any directional movement over large distances. At long distances (5

mm) self-generated gradients play an important role (Tweedy et al. 2016). Self-generated

gradients occur when cells produce enzymes that break down the chemoattractant locally.

This can steepen the attractant gradient and allow directional movement towards the source.

D. discoideum amoebas secrete deaminases to break down folate (Bernstein et al. 1981) and

cAMP phosphodiesterase to breakdown cAMP (Shaulsky et al. 1998). Our experiments

involve distances well above 500 μm, so attractant breakdown is likely to be quite important.

Amoeba movement in this context might be limited to chemoattractants that can be degraded

such as folate and cAMP. In experiments using much shorter distances, amoebas may be able

to respond to, and perhaps learn from, chemoattractants that they cannot degrade.

In conclusion, our study shows that innate preferences are a key component of foraging

behaviour in a microbial predator, *Dictyostelium discoideum*. We found no evidence for

learned prey attraction behaviours. This is a puzzling result that merits further exploration.

Future work could focus on the microstructure of bacterial soil communities experienced by

amoebas and if that is conducive for learning. Further characterization of GPCRs could reveal

whether or not the amoebas possess a flexible mechanism for in sensing different

chemoattractants. Finally, experiments involving attraction over shorter distances might

uncover learned foraging behaviours at smaller scales that we tested.

References

Amacker, N., Z. Gao, J. Hu, A. L. C. Jousset, G. A. Kowalchuk, and S. Geisen. 2022. Protist

feeding patterns and growth rate are related to their predatory impacts on soil

bacterial communities. FEMS Microbiology Ecology 98:fiac057.

Artemenko, Y., T. J. Lampert, and P. N. Devreotes. 2014. Moving towards a paradigm:

common mechanisms of chemotactic signaling in Dictyostelium and mammalian

leukocytes. Cell. Mol. Life Sci. 71:3711–3747.

Aziz, R. K., D. Bartels, A. A. Best, M. DeJongh, T. Disz, R. A. Edwards, K. Formsma, S. Gerdes,

E. M. Glass, M. Kubal, F. Meyer, G. J. Olsen, R. Olson, A. L. Osterman, R. A. Overbeek,

L. K. McNeil, D. Paarmann, T. Paczian, B. Parrello, G. D. Pusch, C. Reich, R. Stevens, O.

Vassieva, V. Vonstein, A. Wilke, and O. Zagnitko. 2008. The RAST Server: rapid

annotations using subsystems technology. BMC Genomics 9:75.

Bernays, E. A. 2001. Neural limitations in phytophagous insects: Implications for diet

breadth and evolution of host affiliation. Annual Review of Entomology 46:703–727.

Bernstein, R. L., M. Tabler, D. Vestweber, and R. Van Driel. 1981. Extracellular folate

deaminase of *Dictyostelium discoideum*. Biochimica et Biophysica Acta (BBA) -

General Subjects 677:295–302.

Brettin, T., J. J. Davis, T. Disz, R. A. Edwards, S. Gerdes, G. J. Olsen, R. Olson, R. Overbeek, B.

Parrello, G. D. Pusch, M. Shukla, J. A. Thomason, R. Stevens, V. Vonstein, A. R.

Wattam, and F. Xia. 2015. RASTtk: a modular and extensible implementation of the

RAST algorithm for building custom annotation pipelines and annotating batches of

genomes. Sci Rep 5:8365.

Brock, D. A., T. S. Haselkorn, J. R. Garcia, U. Bashir, T. E. Douglas, J. Galloway, F. Brodie, D. C.

Queller, and J. E. Strassmann. 2018. Diversity of free-living environmental bacteria

and their interactions with a bactivorous amoeba. Front. Cell. Infect. Microbiol.

8:411.

Carrasco-Pujante, J., C. Bringas, I. Malaina, M. Fedetz, L. Martínez, G. Pérez-Yarza, M.

Dolores Boyano, M. Berdieva, A. Goodkov, J. I. López, S. Knafo, and I. M. De la

Fuente. 2021. Associative Conditioning Is a Robust Systemic Behavior in Unicellular

Organisms: An Interspecies Comparison. Front Microbiol 12:707086.

De Souza, J. T., M. Mazzola, and J. M. Raaijmakers. 2003. Conservation of the response

regulator gene gacA in Pseudomonas species. Environmental Microbiology 5:1328–

1340.

Dukas, R. 2008. Evolutionary Biology of Insect Learning. Annual Review of Entomology

53:145–160. Annual Reviews.

Fortunato, A., J. E. Strassmann, L. Santorelli, and D. C. Queller. 2003. Co-occurrence in

nature of different clones of the social amoeba, Dictyostelium discoideum.

Molecular Ecology 12:1031–1038.

Gans, J., M. Wolinsky, and J. Dunbar. 2005. Computational Improvements Reveal Great

Bacterial Diversity and High Metal Toxicity in Soil. Science 309:1387–1390.

Gershman, S. J., P. E. Balbi, C. R. Gallistel, and J. Gunawardena. 2021. Reconsidering the

evidence for learning in single cells. eLife 10:e61907. eLife Sciences Publications, Ltd.

Giurfa, M., J. Núñez, L. Chittka, and R. Menzel. 1995. Colour preferences of flower-naive

honeybees. J Comp Physiol A 177:247–259.

Guilford, T. 1990. The evolution of aposematism. P. *in* Insect defences: adaptive

mechanisms and strategies of prey and predators. Suny Press.

Hadfield, J. D. 2010. MCMC Methods for Multi-Response Generalized Linear Mixed Models:

The MCMCglmm R Package. Journal of Statistical Software 33:1–22.

Hassan, K. A., A. Johnson, B. T. Shaffer, Q. Ren, T. A. Kidarsa, L. D. H. Elbourne, S. Hartney, R.

Duboy, N. C. Goebel, T. M. Zabriskie, I. T. Paulsen, and J. E. Loper. 2010. Inactivation

of the GacA response regulator in Pseudomonas fluorescens Pf-5 has far-reaching

transcriptomic consequences. Environmental Microbiology 12:899–915.

Heidel, A. J., H. M. Lawal, M. Felder, C. Schilde, N. R. Helps, B. Tunggal, F. Rivero, U. John, M.

Schleicher, L. Eichinger, M. Platzer, A. A. Noegel, P. Schaap, and G. Glockner. 2011.

Phylogeny-wide analysis of social amoeba genomes highlights ancient origins for

complex intercellular communication. Genome Research 21:1882–1891.

Inglis, R. F., O. Asikhia, E. Ryu, D. C. Queller, and J. E. Strassmann. 2018. Predator-by-

Environment Interactions Mediate Bacterial Competition in the Dictyostelium

discoideum Microbiome. Front. Microbiol. 9.

Johnke, J., Y. Cohen, M. de Leeuw, A. Kushmaro, E. Jurkevitch, and A. Chatzinotas. 2014.

Multiple micro-predators controlling bacterial communities in the environment.

Current Opinion in Biotechnology 27:185–190.

Jousset, A. 2012. Ecological and evolutive implications of bacterial defences against

predators: antipredator defences and bacterial ecology. Environmental Microbiology

14:1830–1843.

Jousset, A., L. Rochat, M. Péchy-Tarr, C. Keel, S. Scheu, and M. Bonkowski. 2009. Predators

promote defence of rhizosphere bacterial populations by selective feeding on non-

toxic cheaters. The ISME Journal 3:666–674.

King, J. S., and R. H. Insall. 2009. Chemotaxis: finding the way forward with Dictyostelium.

Trends in Cell Biology 19:523–530.

Klein, P. S., T. J. Sun, C. L. Saxe, A. R. Kimmel, R. L. Johnson, and P. N. Devreotes. 1988. A

Chemoattractant Receptor Controls Development in Dictyostelium discoideum.

Science 241:1467–1472. American Association for the Advancement of Science.

Kuznetsova, A., P. B. Brockhoff, and R. H. B. Christensen. 2017. lmerTest Package: Tests in

Linear Mixed Effects Models. Journal of Statistical Software 82:1–26.

Lenth, R. V., P. Buerkner, M. Herve, M. Jung, J. Love, and F. Miguez. 2022. Emmeans:

Estimated Marginal Means, aka Least-Squares Means, Version 1.8. 1-1.

Letunic, I., and P. Bork. 2021. Interactive Tree Of Life (iTOL) v5: an online tool for

phylogenetic tree display and annotation. Nucleic Acids Research 49:W293–W296.

Lima, W. C., D. Balestrino, C. Forestier, and P. Cosson. 2014. Two distinct sensing pathways

allow recognition of Klebsiella pneumoniae by Dictyostelium amoebae. Cellular

Microbiology 16:311–323.

Meena, N. P., and A. R. Kimmel. 2017. Chemotactic network responses to live bacteria show

independence of phagocytosis from chemoreceptor sensing. eLife 6:e24627.

Mery, F., and J. G. Burns. 2010. Behavioural plasticity: an interaction between evolution and

experience. Evol Ecol 24:571–583.

Minh, B. Q., H. A. Schmidt, O. Chernomor, D. Schrempf, M. D. Woodhams, A. von Haeseler,

and R. Lanfear. 2020. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic

Inference in the Genomic Era. Molecular Biology and Evolution 37:1530–1534.

Mishra, A., A. Jain, P. S. Iyer, A. Suryanarayanan, K. Nordström, and S. B. Olsson. 2025.

Innate floral object identification in a solitary pollinator employs a combination of

both visual and olfactory cues. Sci Nat 112:18.

Nakagawa, S., and H. Schielzeth. 2013. A general and simple method for obtaining R^2 from

generalized linear mixed-effects models. Methods Ecol Evol 4:133–142.

Negassa, W. C., A. K. Guber, A. N. Kravchenko, T. L. Marsh, B. Hildebrandt, and M. L. Rivers.

2015. Properties of Soil Pore Space Regulate Pathways of Plant Residue

Decomposition and Community Structure of Associated Bacteria. PLOS ONE

10:e0123999.

Overbeek, R., R. Olson, G. D. Pusch, G. J. Olsen, J. J. Davis, T. Disz, R. A. Edwards, S. Gerdes,

B. Parrello, M. Shukla, V. Vonstein, A. R. Wattam, F. Xia, and R. Stevens. 2014. The

SEED and the Rapid Annotation of microbial genomes using Subsystems Technology

(RAST). Nucleic Acids Res 42:D206-214.

Pan, M., X. Xu, Y. Chen, and T. Jin. 2016. Identification of a Chemoattractant G-Protein-

Coupled Receptor for Folic Acid that Controls Both Chemotaxis and Phagocytosis.

Developmental Cell 36:428–439.

Pan, P., E. M. Hall, and J. T. Bonner. 1972. Folic Acid as Second Chemotactic Substance in the

Cellular Slime Moulds. Nature New Biology 237:181–182.

Paradis, E., and K. Schliep. 2019. ape 5.0: an environment for modern phylogenetics and

evolutionary analyses in R. Bioinformatics 35:526–528.

Queller, D. C. 2014. Joint phenotypes, evolutionary conflict and the fundamental theorem of

natural selection. Philosophical Transactions of the Royal Society B: Biological

Sciences 369:20130423.

R Core Team, A. 2022. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria. 2012.

Rashidi, G., and E. A. Ostrowski. 2019. Phagocyte chase behaviours: discrimination between

Gram-negative and Gram-positive bacteria by amoebae. Biol. Lett. 15:20180607.

Raynaud, X., and N. Nunan. 2014. Spatial Ecology of Bacteria at the Microscale in Soil. PLOS

ONE 9:e87217.

Schloss, P. D., and J. Handelsman. 2006. Toward a Census of Bacteria in Soil. PLOS

Computational Biology 2:e92.

Schneider, C. A., W. S. Rasband, and K. W. Eliceiri. 2012. NIH Image to ImageJ: 25 years of

image analysis. Nat Methods 9:671–675.

Shaulsky, G., D. Fuller, and W. F. Loomis. 1998. A cAMP-phosphodiesterase controls PKA-

dependent differentiation. Development 125:691–699.

Sherr, E. B., and B. F. Sherr. 2002. Significance of predation by protists in aquatic microbial

food webs. Antonie van Leeuwenhoek 81:16.

Shreenidhi, P. M., D. A. Brock, R. I. McCabe, J. E. Strassmann, and D. C. Queller. 2024. Costs

of being a diet generalist for the protist predator *Dictyostelium discoideum*. Proc.

Natl. Acad. Sci. U.S.A. 121:e2313203121.

Smith, W. P. J., B. R. Wucher, C. D. Nadell, and K. R. Foster. 2023. Bacterial defences:

mechanisms, evolution and antimicrobial resistance. Nat Rev Microbiol 21:519–534.

Stallforth, P., D. A. Brock, A. M. Cantley, X. Tian, D. C. Queller, J. E. Strassmann, and J. Clardy.

2013. A bacterial symbiont is converted from an inedible producer of beneficial

molecules into food by a single mutation in the *gacA* gene. Proc. Natl. Acad. Sci.

U.S.A. 110:14528–14533.

Steele, M. I., J. M. Peiser, S. P. Dawson, D. C. Queller, and J. E. Strassmann. 2024. The genetic

basis of predation resistance in Pseudomonas species associated with the

bactivorous soil amoeba Dictyostelium discoideum. bioRxiv 2024–08.

Steele, M. I., J. M. Peiser, P. M. Shreenidhi, J. E. Strassmann, and D. C. Queller. 2023.

Predation-resistant *Pseudomonas* bacteria engage in symbiont-like behavior with the

social amoeba *Dictyostelium discoideum*. The ISME Journal 17:2352–2361.

Stephens, D. W. 1991. Change, regularity, and value in the evolution of animal learning.

Behavioral Ecology 2:77–89.

Strassmann, J. E., and D. C. Queller. 2011. Evolution of cooperation and control of cheating

in a social microbe. Proceedings of the National Academy of Sciences 108:10855–

10862.

Tweedy, L., D. A. Knecht, G. M. Mackay, and R. H. Insall. 2016. Self-Generated

Chemoattractant Gradients: Attractant Depletion Extends the Range and Robustness

of Chemotaxis. PLOS Biology 14:e1002404.

Vos, M., A. B. Wolf, S. J. Jennings, and G. A. Kowalchuk. 2013. Micro-scale determinants of

bacterial diversity in soil. FEMS Microbiol Rev 37:936–954.

Wick, R. R., L. M. Judd, C. L. Gorrie, and K. E. Holt. 2017. Unicycler: Resolving bacterial

genome assemblies from short and long sequencing reads. PLOS Computational

Biology 13:e1005595.

Figures

A

Innate prey preference

0.65 cm

2µl spot of $OD_{600} = 5$ bacteria

2µl spot of 20,000 naive amoebas

2µl spot of KK2 buffer

Count number of amoebas in the defined boundaries

B

Learned prey preference

Familiar bacterium   Amoebas   Novel bacterium   Familiar bacterium   Amoebas   Novel bacterium

Amoebas familiarized to green bacterium are given a choice between green bacterium or novel bacterium

Green bacterium also appears as novel bacterium in other experiments

Figure 1.Experimental set-up. A) Innate prey preference in *D. discoideum* amoebas: A

spot of AX4 amoebas was placed between spots of bacteria and KK2 buffer. The amoeba

spot was imaged after 6 hours to estimate the number of amoebas that move towards the

bacteria and buffer. **B) Learned prey preference in *D. discoideum* amoebas**: QS9 amoebas

cultured on bacterium A (referred to as familiar bacterium) were given a choice between a

familiar and a novel bacterium. The number of amoebas that move towards the familiar and

novel bacterium were counted at 4 h and 6 h. Each bacterium appears as both familiar and

novel in these experiments. Though we represented only four bacterial species in this figure,

the full design included fully factorial tests with eight species.

Figure 2. Amoebas are innately attracted to more profitable prey bacteria. Normalized movement of naïve amoebas towards different species of bacteria increases with increasing population growth rate of amoebas on these bacteria (MCMCglmm, Fixed effect posterior mean = 4.58, 95% CI (1.00, 8.11), pMCMC = 0.011, R^2 = 0.21). Each point is an average of three biological replicates of amoeba population growth rate and three technical replicates of amoeba movement on 23 species of soil bacteria. The solid blue line represents the mean value of slope and intercept from the MCMC glmm model. The fine blue lines are a sampling of 100 slopes and intercepts that lie within the 95% CI distribution of these values from the model.

Figure 3. Amoebas prefer the edible transposon mutant over the inedible wild type of

Pseudomonas **sp. 20P 3.2.4.** A). In experiments with a choice between a bacterium and

buffer, net amoeba movement towards the edible mutant is significantly higher than

movement towards the wild type *Pseudomonas* bacterium (LMM: p = 0.03). B) In

experiments with a choice between the two bacterial strains, proportional amoeba

movement towards the edible mutant is significantly different from 0.5 (p = 0.009).

Figure 4. A) Amoebas varied in the level of attraction towards the familiar bacterium.

Proportional movement of amoebas towards familiar bacterium averaged across 7 species of

novel bacteria. The points represent $f_{i,j}$ (movement towards familiar bacterium i when paired

against novel bacterium j). For each familiar bacterium, the black squares and error bars mark

the mean $f_{i,j}$ across all seven novel bacteria (= F_i) and its 95% confidence interval. **B) The**

variation in preference for the familiar bacterium was positively correlated with the

innate preference for the bacterium (MCMCglmm, Fixed effect posterior mean = 0.16,

95% CI (-0.02, 0.36), pMCMC = 0.07, R^2 = 0.28). Here the colored squares represent F_i. **C)**

Prey experience does not modulate prey preference in *D. discoideum* amoebas.

Difference in proportional movement of amoebas when they are familiar compared to when

they are novel ($F_i - N_j$), averaged across all partner bacterial species. There was no significant

difference in average movement of amoebas when bacteria are familiar compared to when

bacteria are novel. The white circle with the error bars marks the mean plus 95% confidence

intervals.

754 Table 1: List of all bacteria used in innate preference experiments. The bacteria marked

755 in green were also used in learned preference experiments.

Strain number	Closest species by 16s (Brock2018)	Abbreviation	Phylum/Class	Gram status
18P 8.1.1	*Achromobacter aegrifaciens*	Aa	Betaproteobacteria	Negative
20P 9.1.2	*Agrobacterium rubi*	Ar	Alphaproteobacteria	Negative
18P 2.2.1	*Agrobacterium tumefaciens*	At	Alphaproteobacteria	Negative
20P 7.2.1	*Bacillus aryabhattai*	Ba	Firmicutes	Positive
20P 6.1.2	*Brucella papionis*	Bp	Alphaproteobacteria	Negative
17D 9.2.1	*Bacillus toyonensis*	Bt	Firmicutes	Positive
18P 8.2.2	*Buttiauxella warmboldiae*	Bw	Gammaproteobacteria	Negative
14P 4.3.2	*Chryseobacterium rhizosphaerae*	Cr	Bacteroidetes	Negative
18P 6.2.3	*Comamonas testosteroni*	Ct	Betaproteobacteria	Negative
20P 3.2.2	*Flavobacterium ginsengisoli*	Fg	Bacteroidetes	Negative
Lab clone	*Klebsiella pneumoniae*	Kp	Gammaproteobacteria	Negative
11P 7.1.1	*Mycobacterium arabiense*	Ma	Actinobacteria	Positive
14P 6.2.1	*Microbacterium maritypicum*	Mm	Actinobacteria	Positive
18P 5.1.3	*Nocardia coeliaca*	Nc	Actinobacteria	Positive
14P 8.1.1	*Pseudomonas kuykendallii*	Pk	Gammaproteobacteria	Negative
14P 5.3.2	*Pseudomonas migulae*	Pm	Gammaproteobacteria	Negative
20P 2.1.2	*Pandoraea oxalativorans*	Po	Betaproteobacteria	Negative
20P 2.1.1	*Paenibacillus pabuli*	Pp	Firmicutes	Positive
17P 9.1.1	*Rhodococcus agglutinans*	Ra	Actinobacteria	Positive
20P 10.2.1	*Sphingobacterium ginsensidimutans*	Sg	Bacteroidetes	Negative
14P 6.2.3	*Stenotrophomonas maltophilia*	Sm	Gammaproteobacteria	Negative
5S 2.1.1	*Staphlycoccus saprophyticus*	Ss	Firmicutes	Positive
20P 9.1.1	*Shinella zoogloeoides*	Sz	Alphaproteobacteria	Negative
20P 10.2.4	*Variovorax boronicumulans*	Vb	Betaproteobacteria	Negative

757   **Supplementary figures:**

760   **Figure S1**: A) Movement of AX4 amoebas towards *Klebsiella penumoniae* on different days.

761   B) Movement towards other bacteria on different days. Since chemotaxis experiments

762   towards *Klebisella penumoniae* were repeated on all days, we can see that there are day-

763   specific effects that need to be corrected for.

Figure S2: Data from innate preference experiments. The movement towards the bacterial

spot is represented by green bars, and movement towards KK2 buffer is represented by grey

bars. Amoebas generally move towards the bacteria and rarely to buffer. The bar marks the

means across the 3 technical replicates points (more for the *Kp* control).

Figure S3: Normalized movement of AX4 amoebas towards different species of bacteria. We

first calculated net amoeba movement towards a bacterium in each replicate by subtracting

the number of amoebas that moved towards KK2. To make amoeba movement comparable

across different days, we divided net movement towards bacterium by average net movement

towards *Klebsiella pneumoniae* on that day. Bar marks the mean across 3 technical replicates

(points).

Figure S4: Normalized movement of amoebas versus their growth rate on various bacteria.

We found no significant effect interaction effect between Gram status and mean growth rate

on normalized amoeba movement (p = 0.37). We used a linear model with normalized

amoeba movement as the response variable, and included mean amoeba population growth

rate and bacterial Gram status as fixed effects, along with their interaction.

788 **Figure S5:** Normalized movement towards KK2 buffer vs. Normalized movement towards

789 bacteria. Here normalization means correction for any day effect, where we divided all

790 movement on a given day by mean movement towards *Klebsiella pneumoniae* on that day.

791 We found no significant relation between movement towards buffer and movement towards

792 bacteria (Fixed effect posterior mean = 0.06, 95% CI (-0.01, 0.13), pMCMC = 0.12).

794