Multi-timescale neural adaptation underlying long-term musculoskeletal reorganization.

(Multi-timescale neural adaptation**)**

*Roland Philipp[1,2], Yuki Hara[3], Naohito Ohta[1,2], Naoki Uchida[1,2], Tomomichi Oya[1,4], Tetsuro Funato[2], and *Kazuhiko Seki[1]

[1]National Center of Neurology and Psychiatry, Department of Neurophysiology, 4-1-1 Ogawa-Higashi, Kodaira, Tokyo 1878502, Japan

[2]University of ElectroCommunications, Graduate School of Informatics and Engineering, Dept. of Mechanical and Intelligent Systems Engineering. W2-217, 1-5-1 Chofugaoka, Chofu, Tokyo, Japan

[3]National Center of Neurology and Psychiatry, Department of Orthopaedic Surgery, 4-1-1 Ogawa-Higashi, Kodaira, Tokyo 1878502, Japan

[4]Western Institute for Neuroscience, University of Western Ontario, London, N6A 3K7, Canada

*Corresponding authors:
Kazuhiko Seki and Roland Philipp
Email: seki@ncnp.go.jp and roland@ncnp.go.jp

Author Contributions: R.P. and K.S. designed all experiments and wrote the paper. R.P., Y.H., N.U., and N.O. performed the experiments and analyzed the data. R.P., Y.H., T.O., and K.S. performed the surgeries. T.F. provided conceptual advice and wrote the paper. All authors reviewed the results and approved the final version of the manuscript.

Conflict of Interest Statement: The authors declare no competing interests.

Number of pages: 83 (92, including supplementary information)

Number of figures: 13 (18, including supplementary figures)

Number of multimedia: 7

Number of words for abstract: 173

Number of words in introduction: 706

Number of words in discussion: 1851

ABSTRACT

The central nervous system (CNS) can effectively control body movements despite environmental changes. While much is known about adaptation to external environmental changes, less is known about responses to internal bodily changes. This study investigates how the CNS adapts to long-term alterations in the musculoskeletal system using a tendon transfer model in non-human primates. We surgically relocated finger flexor and extensor muscles to examine how the CNS adapts its strategy for finger movement control by measuring muscle activities during grasping tasks. Two months post-surgery, the monkeys demonstrated significant recovery of grasping function despite the initial disruption. Our findings suggest a two-phase CNS adaptation process: an initial phase enabling function with the transferred muscles, followed by a later phase abolishing this enabled function and restoring a control strategy that, while potentially less conflicted than the maladaptive state, resembled the original pattern, possibly representing a 'good enough' solution. These results highlight a multi-phase CNS adaptation process with distinct time constants in response to sudden bodily changes, offering potential insights into understanding and treating movement disorders.

SIGNIFICANCE STATEMENT

After major changes to the body's mechanics, the nervous system adapts using strategies on multiple timescales. Our primate tendon transfer study shows that core muscle synergy groupings remain stable, reflecting a default to modular control. However, the activation of these synergies changes dramatically; an initial, rapid 'swap' of their timing proves to be **maladaptive**, impairing motor function. This conflict is only resolved through the gradual development of **slower, compensatory strategies** over several weeks. This process highlights the fundamental tension the CNS faces when its reliance on stable motor modules conflicts with the need for flexible control, offering insights into neural plasticity and staged rehabilitation.

Introduction

The central nervous system (CNS) continuously adapts bodily functions in response to both external and internal challenges. Experimental models based on external perturbations—such as altered gravitational fields or distorted sensory feedback—have illuminated mechanisms of sensorimotor adaptation (Sugita, 1996; Davidson and Wolpert, 2003; Luauté et al., 2009; Fleury et al., 2019). Because the changes in the external environment can be controlled by the experimental design, these models provide an opportunity to assess how the CNS adapts to them. However, the transient and predictable nature of these changes may not fully capture the demands posed by internal, long-lasting, and unpredictable alterations to the body's internal environment

In contrast, the internal changes such as developmental growth (Power and Schlaggar, 2017), fatigue (Green, 1997), postural sway (Zemková, 2022), and sensory disorientation (Schärli et al., 2024) impose distinct challenges. Because identifying the source, extent, and time-constant of changes in the internal environment is usually difficult, assessing the corresponding CNS adaptation is also challenging. Particularly, structural alterations to the musculoskeletal system—whether due to injury, disease, or surgery—fundamentally change the

body's biomechanics and sensorimotor associations, but the quantification of

these changes is usually difficult. Accordingly, the way CNS remaps its motor

control strategies corresponding to the changes is not yet well understood

(Walker et al., 2004), although understanding such adaptations is crucial for

elucidating the pathophysiology of motor impairments observed in chronic

musculoskeletal conditions such as (Hunter and Eckstein, 2009) and muscular

dystrophy (Mercuri and Muntoni, 2013).

To address this question, in this study, we employed a tendon transfer (TT)

surgery model (Sperry, 1940), which introduces a controlled, sustained change

to the musculoskeletal structure. TT surgery is a clinically well-established

procedure (Gardenier et al., 2020) that surgically re-attaches the tendon of a

specific muscle to that of a surrounding one. This procedure relocates a specific

muscle, so that its contraction generates a new mechanical action and,

consequently, novel somatosensory feedback. Because the internal change is

controlled by the experimenter, the TT model provides a powerful platform to

investigate how the CNS adapts to a new internal state. Another unique feature

of TT model is that it places permanent changes of the internal environment while

leaving the CNS anatomically intact. Unlike CNS lesion models (Hoffmann et al.,

2009) —where the injury itself disrupts neural circuits and thereby complicates

the assessment of adaptive capacity of the CNS—TT offers a distinct advantage:

it allows investigation of CNS-driven adaptation without the confounding effects

of direct neural damage. Tendon transfer has been used in various species, with

studies suggesting different adaptive capacities, ranging from limited adaptation

in adult rodents (Sperry, 1940; Sperry, 1942; Slawinska and Kasicki, 2002;

Bowlus et al., 2003) to more substantial adjustments in cats (Yumiya et al., 1979;

Loeb, 1999) and primates, including humans (Lee and Seung, 1999; Wester et

al., 2018; Gaetz et al., 2023).   Therefore, adaptability to TT appears enhanced in

primates and humans. Altogether, this approach provides a controlled platform

to examine how the CNS adapts to musculoskeletal changes in primates.

According to earlier reports, there are two motor control strategies that could be

involved when the CNS needs to face the sensorimotor remapping posed by

structural alterations to the musculoskeletal system. First, the CNS may employ

modular building blocks such as muscle synergies—coordinated activations of

muscle groups that reduce the dimensionality of motor control (d'Avella et al.,

2003; Bizzi and Cheung, 2013). Second, skilled behaviors like fine finger

movements often require fractionation—the capacity to activate muscles

independently. When confronted with structural changes to the musculoskeletal

system, does the CNS adapt by modulating existing synergies, or by shifting

toward more fractionated control strategies?

This study aimed to identify long-term adaptive mechanisms of the primate CNS

following structural changes to the musculoskeletal system. We investigated

whether the CNS adapts by modulating existing muscle synergies or by altering

the fractionated control of muscles specifically affected by the surgical alteration.

We surgically altered the limb structure by performing crossed tendon transfers

of finger extensor and flexor muscle tendons. Using a trained finger-grasping task

as our behavioral readout, we examined how the CNS recalibrates muscle activity

to regain skilled motor function. Our findings provide new insights into the

dynamic reorganization of motor control following structural changes of the bodily.

RESULTS

In this study, we developed a novel crossed tendon transfer animal model. This

involved surgically swapping the tendons of two antagonistic finger muscles in a

macaque monkeys' forearm (Fig. 1). The effectiveness and consistency of the

surgery was confirmed by measuring: (i) the distance traveled by muscle fibers

and their intramuscular tendons in the transferred muscle (Fig. 2B, C); and (ii) the

amount of fingertip displacement (Fig. 2D, E). Both measurements were induced

by percutaneous electrical stimulation over the transferred muscles (Fig. 2A). For

a detailed description of the results of these measurements see the Methods

section. The two monkeys were assigned slightly different tasks, allowing us to

examine controlled grasping in Monkey A and a more natural grasp in Monkey B

(refer to the Methods section, and Figs. 3 and Fig. 4 for further details). The two

monkeys ultimately performed related but distinct grasping tasks, a

methodological divergence that provided a valuable opportunity to test the

generality of the core adaptive mechanisms. Monkey A performed a controlled

grasping task requiring a fine precision grip, designed to study adaptation of fine

motor control (Fig. 3). While the same task was initially planned for both animals,

Monkey B performed this task inconsistently, frequently varying its grip strategy,

and we could not reinforce this monkey to perform in single strategy consistently.

Therefore, to ensure reliable task engagement and in accordance with the ethical

principle of Reduction and Refinement (the '3Rs'; (Tannenbaum and Bennett,

2015)), the task for Monkey B was modified to a more naturalistic food retrieval

grasp that the animal performed consistently (Fig. 4).

Once the monkeys had mastered their respective tasks, we recorded control

sessions and then performed the crossed TT. Post-surgery, and once the

monkeys had fully recovered and were able to perform the task independently

(~3–4 weeks), EMG recordings and behavior were resumed.

Functional Recovery Follows a Period of Maladaptive Behavior

Despite the now reversed roles of the transferred muscles, the monkeys were

able to recover their grasping performance within two months, assessed by both

the return of key behavioral metrics to pre-surgical levels (Fig. 5A) and their

consistent ability to perform the task independently (Supplementary Videos S3,

S4, S7). However, this recovery was not immediate. The initial post-surgical

period was instead defined by a phase of significant *maladaptation*, primarily

characterized by aberrant reaching. These aberrant movements included

inefficient, 'explorative' trajectories in Monkey A and target overshoots in Monkey

B (Supplementary Videos S2, S3, S6). This impairment was quantified across

two key behavioral metrics: grip formation times (Fig. 5A) and aberrant reaching

times (Fig. 5B). First, task-related grip formation times were significantly longer

immediately post-TT (Monkey A: 197.7 ± 92.2 vs. 660.6 ± 221 ms, $p = 0.014$;

Monkey B: 169.7 ± 14 vs. 316.3 ± 31.9 ms, $p = 0.016$); it took approximately 10

days for this metric to return to pre-TT levels and 20 to 32 days, respectively, to

stabilize. Second, the duration of the aberrant reaching component itself was

substantially elevated post-surgery. Monkey A exhibited these movements from

day 29 (1978 ± 715 ms) to 73 (93.4 ± 3.7 ms), while Monkey B showed them from

day 20 (103.4 ms) to 62 (107.6 ms), with the duration peaking at days 38-46 (167

± 1.4 ms). Ultimately, overcoming these maladaptive behaviors restored

movement efficiency and precision, leading to the stabilization of grasping

performance by the end of the experimental period.

A two-phase adaptation is observed in the EMG activity of individual muscles

Initially, we investigated whether EMG activity exhibited any changes post-TT. Given that we interchanged the tendons of an antagonistic muscle pair, functionally rendering the FDS an extensor and the EDC a flexor (as confirmed mechanically, Fig. 2), a biomechanically sound adaptation would be to use the former finger extensor (EDC) for finger flexion, and the former finger flexor (FDS) for finger extension during a grasping task. We therefore analyzed EMG activity from both transferred and non-transferred muscles to determine if these expected functional changes occurred. Figure 6 shows a comparison of EMG activity profiles for the two transferred muscles, EDC and FDS, in Monkey A (Fig. 6A–D). Prior to surgery, the control showed distinct and contrasting EMG profiles in both muscles. For instance, peak activity in EDC (▼) occurred long after completion of the grasping action (indicated by vertical dashed lines at 0% task range) at 15% task time (Fig. 6A), coinciding with the animal pre-shaping its hand to prepare for the subsequent grasp. Whereas peak activity in FDS (▽) was observed immediately after completion of the grasping action (0.49%; Fig. 6C). The question we posed was whether the post-surgery activity of EDC resembled

its original profile (without adaptation to surgery) or the activity of FDS (the

anticipated profile following surgical adaptation), and vice versa. Our findings

suggested that EMG activity patterns largely shifted in a manner consistent with

the new mechanical function imposed by the transfer. Early after TT, peak activity

of EDC occurred at -0.58% task time (Fig. 6B; days 29 [red line] and 64 [orange]),

which aligns with peak activity of FDS prior to TT (Fig. 6C; black line). Also, for

FDS, peak activity occurred at 11% task time (Fig. 6D; ▼), which aligned with

peak activity of EDC (Fig. 6A).

This was corroborated by the results in Monkey B (Fig. 6, panels K-M). Although

the EMG patterns of both muscles varied significantly as the two monkeys

performed different types of grasping actions, we found that the EMG of EDC

peaked at -6.12% before (Fig. 6K; ▼) and 0.35% and 1.05% after TT (Fig. 6L; ▽,

days 22 and 36, respectively), closely matching the timing for FDS control data

(Fig. 6M). The corresponding analysis for FDS could not be performed in Monkey

B, since the EMG signal of FDS was lost early after TT (see Methods for details).

We were able to extend our recordings to 122- and 64-days post-TT for monkeys

A and B, respectively. This allowed us to examine whether the adaptations

observed in the early period were consistent throughout the experimental period.

These additional days (Fig. 6B, D; 69 [cyan], 79 [blue], and 99 [black] post-TT;

see Fig. S1 for recordings from all days) showed that the EMG activity profiles for

both muscles unexpectedly returned to their pre-surgery state. For example, as

already described, the EMG signal of EDC at days 29 and 64 shared the

characteristics of FDS, but by the following day (only 5 days later, day 69 [light

blue] post-TT) already exhibited pre-TT characteristics of EDC, continuing as

such until the end of the recording period (day 79 [blue] and 99 [black]). This was

confirmed for the EDC muscle of Monkey B (Fig. 6K, L; Fig. S1U-AO for

recordings from all days).

Figure 6G illustrates cross-correlation coefficients between the EDC EMG profile

post-TT (EDC-post) and the original EDC control data (EDC-pre) in Monkey A.

Figure 6I shows coefficients for EDC-post and FDS-pre correlations in the same

monkey. The coefficient corresponding to the original profile decreased to

between -0.3 and 0.2 over approximately 65 days (Fig. 6G). In contrast, the

coefficient corresponding to the expected EMG profile exhibited an inverse

pattern, with low coefficients pre-TT and a high coefficient (of 0.9) post-TT (Fig.

6I). Cross-correlation analysis for FDS (Fig. 6H, J) revealed a similar pattern.

Cross-correlation coefficients for FDS-pre- vs. FDS-post-analysis declined from

1 to negative values, only to rebound to their original values within a month (day

79; Fig. 6H). This outcome aligns with our previous observations (Fig. 6B, D), and

were corroborated by the results from Monkey B (Fig. 6K-Q). Here, cross-

correlation coefficients dropped from +0.95 to +0.1 for EDC-pre vs. EDC-post

comparison, and increased from -0.5 to 0.9 for the FDS-pre- vs. EDC-post

comparison (Fig. 6P, Q). After 42 days post-TT, cross-correlation coefficients

started to gradually increase or decrease to +0.9 and -0.7, respectively. This

indicates that the anticipated initial changes in EMG activity profiles are not

permanent. Instead, they revert to their original patterns two months post-TT.

Interestingly, changes in EMG pattern were not isolated to the transferred

muscles; non-transferred muscles exhibited a variety of complex adaptations

(Figs. 6, S1-S3). Many agonists adapted in concert with their transferred synergist,

following the same two-phase "swap-and-revert" pattern seen in Figure 6. In

Monkey A, these included extensors ED23 and ECU, while in Monkey B, they

included ED23, ED45, and ECU (see Fig. S1 for details). In contrast, other

muscles showed patterns that were incompatible with a simple swap. For

example, the non-transferred flexor carpi radialis (FCR) in Monkey A showed a

distinct adaptive profile characterized by a drastic initial decrease in one activity

peak and a consistent increase in a later peak (Fig. 6E, F). A similar incompatible

pattern was seen in palmaris longus (PL) (Fig. S1H, I, R, S). Finally, some

muscles, like the extensor carpi radialis (ECR) in Monkey B, remained relatively

stable post-surgery (Fig. 6N, O).

Adaptation occurs through modulating the activation of stable muscle

synergies

Alternation of EMG activity in non-transferred muscle suggests one possibility:

TT might alter the control strategy of coordinated muscle activity for hand

movement by modifying the transferred muscles and their agonists as a cohesive

unit (i.e., muscle synergy). To test this, we conducted a similar analysis on

temporal activation coefficients of muscle synergies. We computed synergy

weights (Fig. 7A, B and Fig. 8C, G for Monkey A; and Fig. 7E, F and 8 D, H for

Monkey B) and temporal activation coefficients (Fig. 7 C, D and Fig. 8 E, I for

Monkey A; and Fig. 7 G, H and Fig. 8 F, J for Monkey B) from EMGs using non-

negative matrix factorization (NMF) (Lee and Seung, 1999). We extracted four

muscle synergies that accounted for more than 80% (Cheung et al., 2012) of the

'variance accounted for' (VAF) (Figs. 8A, B, 85% for both monkeys; the results of

the original and shuffled data sets are shown). Among these synergies, one was

a finger flexor synergy and another was a finger extensor synergy, identified

before the crossed tendon transfer in each monkey (Fig. 7A, B, E, F and Fig. S4).

These were referred to as synergy A and synergy B, respectively.

We examined if the composition and activation coefficient of each synergy

changed, corresponding to the changes observed in the transferred and non-

transferred muscles. First, we analyzed changes in the composition of these

synergies. The cosine distance for the same synergy pairs (e.g., synergy A of

pre-TT and synergy A of post-TT in all recording days (Fig. S4A, E; monkey A

and B, respectively) was low (mean [± SD]: 0.03 [± 0.03] in Monkey A and 0.09

[± 0.11] in Monkey B) and different pairs (e.g., synergy A of pre-TT and synergy

B of post-TT) was high (mean [± SD]: 0.67 [± 0.08] in Monkey A and 0.61 [± 0.14]

in Monkey B). There was a one-to-one correspondence between synergy pairs

(p < 0.0001, two-way analysis of variance [ANOVA] due to synergy factor; p <

0.01, Bonferroni post-hoc test between synergy of the same pair and different

pair). This was demonstrated in both monkeys (S4A-D and E-H for monkey A and

B, respectively). These results indicate that the composition of these four

synergies and the contribution of each muscle do not change drastically before

and after TT surgery.

Second, we analyzed changes in the activation coefficients of these synergies.

Prior to TT, the activation coefficients for the flexor synergy (synergy A) reached

a peak at +0.97% (Fig. 7C). Post-TT, these coefficients shifted towards later times

shortly after surgery (+10.7%), but eventually returned to their original timing.

Meanwhile, the extensor synergy (synergy B) peaked at 12.62% pre-TT and

moved to -0.97% shortly after surgery. Like the flexor synergy, the extensor

synergy also reverted to its original timing in subsequent sessions (Fig. 7D).

These characteristics corresponded well to the ones found in the transferred

muscles (Fig. 6), supporting our hypothesis that the transferred muscle and its

agonist changes as a cohesive unit.

Figure 8E shows the temporal changes for synergy C, a wrist flexor synergy (Fig.

8C and S4C). While two peaks were observed prior to surgery (+1.46% and

+15%), the early post-surgery sessions did not exhibit any significant peaks.

However, later sessions displayed a single peak at +14.6% task time, with an

increasing magnitude as a function of recovery days. These characteristic

changes corresponded well to a patten that was incompatible to the transferred

muscles found in FCR and PL, as described above. Indeed, both muscles

showed highest synergy weights for this synergy (Fig. 7C and S4C). Synergy D

was a wrist extensor synergy (Fig. 8G and S4D), and underwent a single notable

change on day 29, but quickly reverted to its original pattern (Fig. 8I). These

pattens of changes in synergy 3 and 4, both contributed minimally to the EMG of

transferred muscles. This suggests that some non-transferred muscles change

their EMG activity following the mode, which is independent to the one observed

in the transferred muscle and may reinforce the functional adaptation made by

the transferred muscle and its synergists.

Several of these results were also confirmed in Monkey B. Activation coefficients

of extensor synergy B peaked at -8.75% before TT, increasing to +3.5% after TT

(Fig. 7G). This peak activity persisted until the last day (day 69 post-TT), when it

shifted back to an earlier time than -10% of the task range. This pattern

corresponds well to that found in the transferred muscles (EDC; Fig. 6K, L).

Again, the characteristics of these changes in activation coefficients of muscle

synergies were also quantitatively confirmed by cross-correlation analysis of all

four muscle synergies in both monkeys. The coefficients were plotted over the

course of experimental days in relation to tendon surgery (Fig. 9). In this analysis,

the activation coefficient of each synergy was cross-correlated with either the one

from synergy A (finger flexor synergy; Fig. 9C, D, G, H) or synergy B (extensor

synergy; Fig. 9A, B, E, F) before TT. As previously demonstrated for the FDS and

EDC muscles, the temporal activation coefficients of muscle synergies A and B

(flexor and extensor) displayed a distinct pattern. After cross-correlation with their

own control data, both the flexor and extensor synergies showed coefficients of

1 prior to surgery. These became negative for 66 days post-surgery before

reverting close to their original values (Fig. 9B, C). However, after cross-

correlation with control data of the antagonistic muscle synergy (i.e., extensor for

flexor synergy and vice versa, Fig. 9B, D), the pattern was reversed. Coefficients

started with negative values prior to TT and shifted to positive coefficients near 1

shortly after surgery. Around 66 days later, the correlation coefficients returned

to negative values. In short, these results mirror our findings between the

transferred muscles (Fig. 6G, H). For synergies C and D, the changes in cross-

correlation coefficients over time were markedly different. For synergy C,

coefficients begin to steadily increase when cross-correlated with control data of

the extensor synergy (synergy B, Fig. 10E), and decreased with the flexor

synergy (synergy A, Fig. 10G). This suggests an increasing contribution to finger

extension. In contrast, synergy D does not exhibit any specific pattern following

TT.

In Monkey B, this cross-correlation analysis revealed a more varied, or differential,

pattern of adaptation across the four synergies (Figs. 9 and 10). The primary

extensor, Synergy B, mirrored the results in Monkey A, showing a clear swap-

and-revert pattern (Fig. 9F, H). In contrast, Synergy A did not show a clear

reversal; its correlation coefficients gradually converged towards zero, likely due

to the absence of the FDS EMG signal in the analysis (Fig. 9E, G). Of the

secondary synergies, Synergy C showed no discernible trend (Fig. 10E, G), while

Synergy D exhibited a complex pattern of an immediate change, a quick reversion,

and a second gradual change late in the adaptation period (Fig. 10F, H).

In summary, both monkeys exhibited a distinct two-phase adaptation following

TT. We define the 'early phase' as the period from initial post-surgical recovery

up to the reversal of the swapped activation patterns (approximately days 20/29

to ~65 post-TT, see Fig. 9), characterized by the transferred muscles/synergies

adopting antagonistic temporal profiles. The 'late phase' encompasses the period

following this switch-back (after ~day 66 post-TT), where original activation

timings were largely restored.

The Early Adaptation Phase is Characterized by a Maladaptive Neural and

Behavioral Profile

The early adaptation phase was defined by a distinct neural strategy that

correlated with significant behavioral deficits. First, the period dominated by

aberrant reaching movements and prolonged grip times (Fig. 5B) was precisely

when the swapped activation of Synergy A and B was most prominent (Fig. 9).

This temporal link provides strong evidence that this initial 'swap' strategy was, in

fact, maladaptive, as the flawed neural control directly underpinned the

impairments in movement efficiency and precision.

Second, we found that the net activity of muscles representing certain muscle

synergies (aggregated average EMG; aaEMG) showed distinct, synergy-specific

changes over time (Fig. 11). A key difference was observed in the finger extensor

synergy (Synergy B). In Monkey A, this synergy demonstrated what we term a

neuromuscular 'arms race': a steady and significant increase in aaEMG post-TT

until day 64 ($p < 0.0001$) as the CNS appeared to escalate activation to overcome

the internal conflict between the transferred EDC (now a flexor) and its non-

transferred synergists (which remained extensors), before abruptly reverting to

its pre-TT level (Fig. 11B, F). In contrast, while Monkey B also showed a

significant initial surge in Synergy B activity (day 22: $p < 0.0001$), this was

followed by a gradual return to baseline and below (day 64: $p < 0.001$), lacking a

sustained 'arms race' (Fig. 11J, N). This discrepancy is likely attributable to the

different force requirements of the tasks. The 'arms race' in Monkey A may have

been driven by the need to generate high grip force against the synergy's internal

conflict, whereas Monkey B's delicate food-retrieval task did not require such high

forces, thus lacking the same drive for escalating, energetically costly activation.

For the finger flexor synergy (Synergy A), Monkey A showed a general decrease

in activity ($p = 0.0004$ at day 79; Fig. 11A, E), while Monkey B showed a

consistent and significant increase throughout the experiment (day 22: $p = 0.0004$; day 64: $p = 0.0008$). This was expected, as Monkey B's Synergy A relied

on FDP and PL to compensate for the loss of FDS function (Fig. 11I, M). Synergy

C exhibited a general increase in activity during the post-TT phase in both

monkeys (Fig. 11G, O), while Synergy D showed no significant differences in

Monkey B and only minor changes in Monkey A (Fig. 11H, P). All statistical

comparisons were made against the pre-surgery control period using a two-

sample t-test with Bonferroni correction ($\alpha = 0.01$).

Taken together, these distinct patterns of aggregated EMG activity—especially

the escalating, conflicted activation in Synergy B—further illustrate that the early

adaptive phase was characterized by an unstable and inefficient neural control

strategy.

Recruitment of a Compensatory Tenodesis Strategy Enables Late

Adaptation

The development of a compensatory strategy, primarily involving the wrist flexor

synergy (Synergy C), appears crucial for enabling the final phase of adaptation.

The increasing contribution of this synergy is supported by cross-correlation

analysis, which shows that the correlation between Synergy C and the extensor

synergy (Synergy B) steadily increased over time after an initial drop (Fig. 10A; p

< 0.05, two-sample t-test). Increased reliance on synergy C following TT suggests

a compensatory strategy to overcome the difficulty in finger extension, crucial for

grasping. This is despite the recruitment of the finger extensor and wrist flexor

synergies, which seemed inadequate to achieve effective flexion. The

interactions between the cumulative proximal-to-distal joint angles and musculo-

tendon length can be complex. In our experiments, as in human patients who

receive tendon transfers (Zajac, 1992), those interactions can affect the ranges

of finger flexion-extension that can be achieved for a given wrist angle (i.e.,

passive and active tenodesis). Briefly, biomechanically, wrist flexion can enable

finger extension (i.e., a dynamic tenodesis where extensor tendons of the fingers

cross both the wrist and interphalangeal joints): Finger extension can cancel out

the increase in total musculo-tendon length produced by wrist flexion. This

compensatory interaction across joints allows for some degree of functional hand

use, even in the absence of active finger movement (Cash and Jones, 2011) (see

the Discussion for a complete description).

Further kinematic analysis of Monkey B (Fig. 12; no analysis on Monkey A due

to low-resolution video images) indicated that this monkey indeed exhibited a

similar active tenodesis as human patients. The angle at the

metacarpophalangeal (MCP) joint, responsible for finger extension, steadily

increased (Fig. 12A) after surgery, indicating finger extension. Concurrently, the

wrist joint angle decreased (Fig. 12B), indicating wrist flexion. This coordinated

movement suggests a compensatory strategy to achieve finger extension during

grasping. Statistical analysis confirmed that MCP and wrist joint angles differed

significantly between pre- and post-TT ($p < 0.0001$, ANOVA; $\alpha = 0.01$). Therefore,

we suggest that deteriorated finger extension after TT was assisted by a

compensatory wrist flexion synergy.

Discussion

The central nervous system's (CNS) response to profound musculoskeletal

change is a fundamental problem in motor control. This study sought to determine

whether the primate CNS adapts to such change by flexibly modulating stable

muscle synergies or by developing more fractionated, independent muscle

control. We found that the CNS initially defaulted to a modular strategy,

repurposing entire synergies by swapping their activation timings. This simple

solution, however, proved to be maladaptive, creating a mechanical conflict that

impaired motor function. This early maladaptive phase was ultimately resolved

through the gradual development of compensatory movements, leading to a

"good enough" functional recovery. This multi-stage process, operating on

different timescales, highlights the intricate balance between modularity and

flexibility in neural adaptation.

The CNS Defaults to a Modular Strategy, Leading to a Maladaptive Conflict

Our primary finding provides a direct answer to the question posed in the

introduction: the CNS adapted to the tendon transfer not by developing fine-

grained, fractionated control, but by implementing a modular strategy by

repurposing entire co-activation modules (Overduin et al., 2008; Berger et al.,

2013; Bizzi and Cheung, 2013; Takei et al., 2017). The initial and most immediate

neural change was a wholesale swap of the temporal activation patterns of the

primary flexor and extensor synergies (Synergy A and B). The fact that the

muscle weight compositions of all four synergies remained remarkably stable

throughout the months-long experiment (Fig. S4), in the face of drastically altered

biomechanics, strongly supports the hypothesis that these synergies represent

stable, neurally constrained building blocks (d'Avella et al., 2003; Bizzi and

Cheung, 2013; Takei et al., 2017). Our result indicated that this modular approach

is likely the default for adaptation to altered musculoskeletal systems because it

is computationally "easy" and metabolically "cheap" (Flash and Sejnowski, 2001).

It relies on the simple retiming of pre-existing, stable neural circuits—thought to

be encoded in the spinal cord and brainstem—which is a far less demanding

operation than the "difficult" alternative of fractionated control (Bizzi and Cheung,

2013). The latter would require extensive synaptic plasticity and cortical

remapping to selectively uncouple muscles that were previously co-activated, a

process associated with the slower, more deliberate learning of an entirely new

motor skill (Kitago and Krakauer, 2013). Thus, the simple and economical

strategy is prioritized in the CNS to deal with an acute alteration of the musculo-

skeletal system.

However, this adherence to established synergy structures created a

fundamental conflict. Activating the original extensor synergy (Synergy B) after

the transfer now inevitably co-activated the surgically transferred EDC, which

functioned mechanically as a flexor, alongside non-transferred muscles like ED23

and ECU, which remained anatomical extensors. This internal mechanical

antagonism appears to be the root cause of the early maladaptive phase. This

interpretation is supported by two key lines of evidence from our results. First, the

period of severe behavioral impairment, characterized by aberrant reaching and

inefficient grasping (Fig. 5B), occurred precisely when this flawed "swap" strategy

was active. Second, the aggregated EMG activity revealed a sustained and

significant increase in the total activation of the conflicted Synergy B in Monkey

A (Fig. 11B, F), likely reflecting an energetically costly effort to overcome the

internal mechanical antagonism. This scenario can be viewed through the lens of

optimal control and cost-benefit analysis, where a cost function is minimized

(Wolpert, 1997). The initial "swap" strategy, while computationally cheap to select,

incurred an unacceptably high operational cost in terms of both poor task

performance (high error) and excessive energy expenditure. It was this

profoundly unfavorable cost-benefit ratio that likely served as the critical error

signal that drove the CNS to abandon this initial strategy.

Resolution Through Slower Compensatory Adaptations

The CNS did not persist in the failed swap-based strategy. The high metabolic

and computational cost of activating a mechanically conflicted synergy likely

triggered the second rapid adaptation: the "switch-back" of synergy activation

timings toward their original patterns (Fig. 7, 9). This rapid reversion, occurring

over just a few days, is characteristic of an error-based learning mechanism, a

form of adaptation that is profoundly impaired by cerebellar damage across a

range of tasks, including adaptation to force fields (Smith and Shadmehr, 2005),

prismatic shifts (Martin et al., 1996), and split-belt treadmills (Morton and Bastian,

2006). It is plausible that the CNS operates with an implicit threshold for an

acceptable cost/performance ratio; once the persistent task failure and high

muscular co-contraction of the swap strategy exceeded this threshold, a swift

recalibration was initiated. This reversion, despite the remaining mechanical

antagonism from the transferred tendons, represents a "good enough" solution

where functional success is prioritized over perfect efficiency (Mussa-Ivaldi and

Bizzi, 1995; Ranganathan et al., 2013; Gijsberts et al., 2014). The rapid timescale

of this change is highly consistent with the cerebellum's proposed role as a

forward model, predicting the sensory consequences of motor commands and

driving rapid learning in response to sensory prediction errors (Popa et al., 2016).

Crucially, this switch-back to a less conflicted state, presumably representing the

involvement of the error-based learning process, was likely enabled only by the

concurrent development of slower, compensatory strategies that provided an

alternative means to achieve the task goal. The primary compensation was a

learned use of the tenodesis effect. Over weeks, the monkeys gradually

increased the activation of the wrist flexor synergy (Synergy C) during hand

opening. This wrist flexion biomechanically generates passive finger extension

(Zajac, 1992; Cash and Jones, 2011), providing a viable new method for hand

pre-shaping. This learned, compensatory behavior, confirmed by kinematic

analysis (Fig. 12), ultimately allowed the CNS to abandon the maladaptive

synergy swap. This gradual, exploratory process is distinct from the following

rapid adaptation and represents a form of motor skill acquisition, a process which

may be associated with plasticity in cortical structures like the motor cortex and

basal ganglia. (Kitago and Krakauer, 2013).

This multi-timescale process, conceptually summarized in Figure 13, aligns well

with influential two-state models of motor learning, which propose parallel "fast"

and "slow" learning systems (Smith et al., 2006a). These models posit that a fast

system responds strongly to error but has poor retention, while a slow system

responds weakly to error but retains information well, enabling robust long-term

memory. The initial synergy swap and subsequent "switch-back" appear to be

governed by the fast system (likely cerebellar), which responds strongly to error

but has poor retention. The gradual learning of the tenodesis effect is the

signature of the slow system (likely cortical and/or basal ganglia-dependent),

which integrates experience over a longer period to build robust, lasting skills

(Trewartha et al., 2014). The interaction between these systems is key: as the

slow system gradually develops the effective tenodesis compensation, it reduces

the overall task error. This reduction in error removes the primary drive for the

fast, error-correcting system to continue its search for a solution. With its error

signal quieted, the fast system allows the control policy to stabilize, and the CNS

"settles" into the "good enough" compensatory state. This suggests a control

architecture where the successful output of a slow, deliberate skill-learning

system can gate and ultimately supersede the output of a fast, reactive adaptation

system, leading to a stable and functional long-term motor plan.

Broader Implications and Limitations

Our findings reveal a complex interaction between adaptive processes operating

on different timescales. The initial, rapid synergy swap may be linked to a

'sensory surprise' from altered proprioceptive feedback (Kitazawa et al., 1995;

Martin et al., 1996; Smith et al., 2006b; Edwards et al., 2012; Petitet et al., 2018),

driving a fast but flawed response. This multi-level network reorganization

presents a more complex challenge than the gain modulation within single or

specific CNS pathways seen in studies of spinal reflex conditioning (Thompson

and Wolpaw, 2014). Further understanding the interaction between the global

synergy reorganization observed in this study and the underlying local

sensorimotor loops will require considering factors not assessed here, such as

the roles of spinal reflex pathways (Smith et al., 2006c) and fusimotor drive

(Vallbo et al., 1979; Hospod et al., 2007). The altered biomechanics would have

immediately changed proprioceptive feedback, triggering rapid modulation of

spinal reflex circuits and requiring recalibration of the fusimotor system, which

adjusts muscle spindle sensitivity via alpha-gamma co-activation. A

comprehensive model must ultimately connect the global synergy reorganization

with these local, interacting feedback and control loops.

While our primary analysis focused on the grasping, the adaptation encompassed

the entire reach-to-grasp sequence, as evidenced by the initial deficits in reaching

movements (Supplementary Videos S2, S3, S6; Fig. 5B). The observation that

both monkeys, engaged in both a highly controlled learned task and a more

naturalistic grasping task, showed similar adaptive patterns strengthens the

generalizability and ecological validity of our findings (Berniker et al., 2013).

Furthermore, the qualitative reports of proficient real-world hand use indicate that

the learned compensatory strategies were not narrow "tricks" but robust solutions

that transferred to unconstrained activities, demonstrating their "portability" to the

complexities of natural behavior.

However, a limitation is that this generalization was not tested systematically.

Another critical point is that our paradigm cannot distinguish between the

unlearning of a maladaptive strategy and the relearning of a functional one. This

distinction is crucial for both motor memory theory and clinical practice (Kitago et

al., 2013). The disappearance of the initial "swap" strategy could be due to

passive decay (forgetting), active inhibition of the memory (extinction), or the

formation of a new, stronger memory that outcompetes it (interference) (Villalta

et al., 2013). The literature increasingly suggests that unlearning is an active

process, and that the new, successful compensatory strategy likely supplants the

maladaptive one through interference. This question is vital for rehabilitation:

should therapy focus on "erasing" bad habits, or on intensively training better

ones to overwrite them? Our results lend weight to the latter, suggesting

functional recovery is a process of competitive memory formation.

Implications for Sensorimotor Rehabilitation

These findings may hold implications for sensorimotor rehabilitation. The

observation of a distinct maladaptive phase followed by compensation and

recovery suggests a staged approach could be beneficial (Hatem et al., 2016;

Kakavas et al., 2025). An early phase of therapy might focus on minimizing

sensory prediction errors caused by the altered proprioceptive feedback, and

reducing maladaptive responses through techniques aimed at integrating the

altered body schema (e.g., using visual feedback or task-oriented practice) (Nudo

et al., 1996; Takeuchi and Izumi, 2012; Zeiler and Krakauer, 2013; Foell et al.,

2014; Levin et al., 2015; De Nunzio et al., 2018; Sattin et al., 2023; van Vliet et

al., 2023). This stage would guide the fast, error-driven learning system. A later

phase could then employ task-specific training to promote effective

compensatory strategies, facilitating the abandonment of maladaptive patterns

and the consolidation of a stable motor strategy (Murata et al., 2008; Maier et al.,

2019). This second stage engages the slow, skill-acquisition system by creating

a rich, problem-solving environment. Finally, a third stage would focus on

intensive, high-repetition practice to automate the new functional motor plan,

making it robust for real-world use through dosage and generalization

(Winterbottom and Nilsen, 2024). This neurobiologically-informed staging—

guiding fast adaptation, then facilitating slow skill learning, then driving

consolidation—may offer a more logical and effective path to functional recovery.

616
617
618

METHODS

Animals

Data were collected using two male macaque monkeys (*Macaca fuscata*; Monkey A: 7.8 kg and Monkey B: 9.9 kg) purpose-bred at the National Bioresource Project (NBRP). They were kept in custom-made primate cages, allowing for potential pair housing. Both monkeys were trained to perform a simple grasping task. During experimental recordings, the monkeys were seated in a primate chair without further restraint. Their head movement remained unrestricted at all times.

Procedures

All procedures were designed to minimize discomfort and pain and approved by the institutional Animal Care and Use Committees at the National Center of Neurology and Psychiatry (NCNP), Tokyo, Japan. Details of the surgical operations, experimental setup, and procedures for electromyography (EMG) recordings have been previously described (Takei and Seki, 2010).

EMG implant surgery

Both animals were familiarized with the experimental setup and trained to perform

the behavioral task prior to surgery. After an initial training period, EMG

electrodes were chronically implanted subcutaneously into several muscles of the

left forearm including the FDS, FDP, and EDC (Figs. 1, 3C, and 4C). Individual

muscles were localized and confirmed using electrical microstimulation before

implanting EMG electrodes chronically by two methods. For muscles involved in

the crossed tendon transfer procedure, muscle fascia was cut and EMG wires

with looped ends were placed on top of the muscle belly. These were then

secured by reclosing the fascia with absorbable suture threaded through the loop

(AS-361, stainless-steel Cooner wire, Conner Wire Co., Chatsworth, CA, USA).

All other muscles were directly implanted into the muscle belly using a 22-gauge

hypodermic needle. Each wire was threaded into the needle tip, folded back along

the shaft, and inserted into the muscle before carefully retracting the needle.

Electrode separation for bipolar recordings was approximately 5–10 mm. Note

that some EMG recording sites were lost over time. Surgical procedures were

carried out under deep general anesthesia (sevoflurane 1.5 to 2.5% in 2:1

O_2/N_2O) and with full aseptic precautions. Heart rate, blood pressure, body

temperature, and blood oxygen saturation were monitored throughout surgery.

Analgesics and antibiotics were administered intramuscularly for at least one

week postoperatively.

Tendon-Transfer Surgery

After a minimum four-week recovery period from EMG implant surgery, TT

surgery was performed. The FDS and EDC tendons were cut as distally as

possible (immediately below and above the wrist for FDS and EDC, respectively)

to avoid damaging the Golgi tendon organ located near the junction between the

muscle fibers and tendon (Schoultz and Swett, 1972; Jami, 1992). Tendons were

then guided either through the gap between the radius and ulnar bone (Monkey

A), or around the wrist and then re-attached to the tendon of the antagonist

muscle using a tendon graft harvested from the plantaris tendon of the lower limb

(Monkey B; EDC → FDS: direct connection; and FDS → EDC: tendon grafting).

This was only necessary in Monkey B since the tendons used for the cross-

transfer were too short. This surgical procedure aimed to reverse the primary

mechanical actions of the manipulated muscles, making the FDS tendon an

effective finger extensor and the EDC tendon an effective finger flexor. The

success and nature of this mechanical rearrangement were verified post-

operatively via direct muscle stimulation (see Methods subsection 'Tendon

transfer confirmation' and Fig. 2). The general organization and function of the

major forearm muscles involved in finger flexion and extension in *Macaca fuscata*

are broadly comparable to those in humans (Vanhoof et al., 2021; Yan et al.,

2022).

Surgical procedures were carried out under deep general anesthesia

(sevoflurane 1.5 to 2.5% in 2:1 O_2/N_2O) and with full aseptic precautions. Heart

rate, blood pressure, body temperature, and blood oxygen saturation were

monitored throughout surgery. Analgesics and antibiotics were administered

intramuscularly for at least one week postoperatively. Both monkeys wore a

plaster cast post-surgery, but its effectiveness was limited (lasting approximately

one week).

Tendon transfer confirmation

To verify the success and long-term stability of TT surgery, two procedures (Fig.

2) were implemented. During the first procedure, the monkey was sedated and

its arm was secured to a metal frame (Fig.2A). Electrical stimulation was then

applied to either the FDS or EDC muscle (50 mA, DS8R, Digitimer, Welwyn

Garden City, UK), while ultrasound scans of tendon movements were

concurrently captured (SONIMAGE MX1, Konica Minolta, Inc., Tokyo, Japan

(Nordez et al., 2009; Dieterich et al., 2017).

Figure 2B shows a sonogram of the FDS muscle and its intramuscular tendons.

The left side displays a static image of the FDS muscle at a specific moment, with

white arrows indicating the FDS tendon used for measurement. The right side

shows staggered images (top white box) of the FDS tendon displacement

triggered by muscle stimulation. The lower right inset illustrates the measurement

area. The displacement wave area represents the intensity of muscle contraction,

and was calculated by measuring the average duration (a, in seconds) and

amplitude (b, in cm) of three successive waves. The area (a*b/2) and regression

line for FDS (red) and EDC (blue) (R^2 > 0.5 for FDS) for days 0, 7, and 105 post-

TT are shown (Fig. 2C). The results suggest that the muscle contractions induced

by direct electrical stimulation remained nearly constant.

For the second procedure, high-speed videos (37U Series Color Industrial Camera, The Imaging Source, Charlotte, NC, USA) of the monkey's finger movements were recorded while electrically stimulating the FDS or EDC. Markers were placed on the nails of the index, middle, and ring fingers (Fig. 2A) to measure finger displacement in xyz-dimensions using DeepLabCut software (Mathis et al., 2018; Nath et al., 2019). The sum of Euclidean distances of each marker from the origin of the three-dimensional (3D) coordinate system was computed as a scalar quantity. Post-surgery, movement along the z-axis reversed, indicating a shift from finger flexion to extension due to TT (Fig. 2D; blue: pre-TT on surgery day, dark brown: post-TT on surgery day, light brown: 1 wk post-TT, and red: 3 wk post-TT). The scalar quantities of each finger did not significantly change during muscle stimulation on days 0-, 7-, and 105-days post-TT (Fig. 2E), suggesting there was no postoperative tendon rupture or slackening.

Data recordings

EMG signals were recorded using the AlphaLab SnR system (Alpha Omega Engineering, Hamerkava St. 6, Ziporit Industrial Zone, P.O.Box 810, Nof HaGalil

(Nazareth Illit) 1789062, Israel), displayed online and then stored on a hard drive

for later off-line analysis using MATLAB (MathWorks, Natick, MA, USA). Data

were recorded at 11 kHz and processed for analysis as follows. 1. Down-sampled

to 5 kHz; 2. 50 Hz high-pass filtered (6th order Butterworth filter); 3. rectified; 4.

20 Hz low-pass filtered (6th order Butterworth filter); and 5. down-sampled to 100

Hz. Behavioral task events were recorded as transistor–transistor logic (TTL)

signals using the AlphaLab SnR system. Monkey behavior was further recorded

by two cameras (Sanyo VPC-WH1, 60 fps [Sanyo, Osaka, Japan]; and Casio EX-

100F, 240 fps [Casio, Tokyo, Japan] for monkeys A and B, respectively) from two

different angles (top and side view). The images were later used to detect

additional behavioral events (contact time with the object and pull onset for

Monkey A; and food touch and food lift onset for Monkey B), and detection of

possible changes in the animal's general movement pattern. Kinovea software

was   used   for   this   video   analysis   (free   and   open-source   software;

https://www.kinovea.org/).

Behavioral task

Before and after EMG surgery, the monkeys were trained on a simple grasping

task that involved a small object attached to a rod (Fig. 3B for Monkey A; and Fig.

4B for Monkey B). For Monkey A, object 1 was a small rod placed between two

side walls encouraging the monkey to grasp the rod with a precision grip using

the tips of the index finger and thumb (controlled grasp). The force used to

compress the spring while pulling was low and the distance moved short. Object

2 had to be grasped in the same way. However, the force to compress the spring

was higher and the distance moved longer (object 1: 300 cN, 4 mm; and object

2: 800 cN, 30 mm). The monkey was asked to grasp and hold the objects for

approximately 300–500 ms (Fig. 3A–D). After completing the trial, the monkey

was rewarded with a piece of fruit that had to be taken from the experimenter's

palm. The task sequence for Monkey A ranged from object 1 hold onset to object

2 hold offset (Fig. 3C, D; and 'Data Analysis' below). For Monkey B, object 1 was

a small rod that was grasped and pulled (300 cN, 4 mm) using a lateral

prehension grip (like using a door key), and resembling a power grip rather than

a natural grasp. However, instead of grasping object 2, Monkey B picked up a

piece of fruit from an allocated location. In doing so, the monkey had to cross the

path of a photo cell which was detected by the recording system (Fig. 4A–D). The

task sequence to be analyzed for Monkey B ranged from object 1 hold onset to

food touch, which was indicated by a LED (Fig. 4C, D; and 'Data Analysis' below).

For Monkey A, the task sequence was as follows (Fig. 3A; and Supplementary

Video S1). From a starting position in front of the monkey, it had to lift its arm and

move towards object 1, grasp, pull, and then hold the object for 500 ms.

Immediately after releasing object 1, Monkey A proceeded to move its hand

towards object 2, which was located to the right of object 1. Again, the monkey

had to grasp, pull, and then hold the object for another 500 ms. Each hold period

was accompanied by an audio signal which stopped once the hold duration was

sufficient. After completing the task, which was signaled by another acoustic

signal, the monkey was rewarded with a piece of fruit presented by the

experimenter and taken by the monkey.

For Monkey B, the first part of the sequence was identical. However, after

releasing object 1, Monkey B was required to pick up a piece of fruit from an

allocated location (Fig. 4A, B; and Supplementary Video S5). In doing so, the

monkey passed a photocell in front of the food well. This event was detected by

the recording system and a video camera (a red LED triggered by activation of

the photocell). For the analysis, only object 1 and the food grasp were considered

in monkeys A and B, respectively.

Data analysis

<u>*EMG analysis*</u>

EMG data were normalized to the average time for each monkey to complete a

trial (object 1 hold onset ➜ object 2 hold offset/LED offset = 100%). Data were

cropped and aligned according to the time stamps (Obj1 hold offset ± 15%; and

LED onset [food touch] ± 15% for monkeys A and B, respectively). Recorded

EMGs from the pre- and post-tendon surgery period were averaged for each

recording and compared over experimental sessions. For both monkeys, the

muscles used for EMG analysis were: FDS and EDC for pre- (control) and EDC

for post-TT (experimental).

Both monkeys required time to recover from surgery. Once they were able to

perform the task on their own, data collection started (29 and 20 days post-

surgery for monkeys A and B, respectively). After tendon surgery, EMG signals

for FDS, FCU, and FCR deteriorated in Monkey B. Therefore, there is no

experimental data for these muscles in this monkey and they were excluded from

the muscle synergy analysis.

Synergy analysis

The EMG envelopes obtained from pre-processing raw EMG data were divided

by the mean value to normalize activity. Muscle synergies were then extracted

for each session using NMF (d'Avella et al., 2003). NMF decomposes the EMG

data matrix M, as a product of two matrices C and W:

$M = CW^T + re = \Sigma c_i w_i^T + re$

where the vector wi and column of matrix W are muscle synergies; the vector ci,

and column of matrix C are their temporal coefficients; and re represents the

residuals. Multiplicative update rules were used for decomposition (Lee and

Seung, 1999). Updating matrices for decomposition from 20 different random

initial matrices was started, and factorization results with least residuals among

the 20 results were used. To improve robustness of the muscle synergies, one

set of muscle synergies was extracted from EMG data of multiple trials of each

day. A k-fold cross validation was used, with k = 4 i.e., EMG data set of multiple

trials during the day was split into four data sets, in which three data sets were

for training and one data set was for testing. Synergy matrix W was calculated

from three training data sets using NMF, and then a coefficient matrix C was

calculated by decomposing the EMG matrix M of the test data set with a fixed W

matrix of the training data set. This operation was repeated by changing the test

data set for each four data sets. Finally, the extracted four C matrices and four W

matrices were averaged to obtain daily synergies.

The number of synergies was determined as the number where VAF exceeded

a threshold. This threshold was set as 0.8. To clarify the effect of TT, the number

of synergies for each monkey was determined, assuming that trial-by-trial

variation in synergies within the same monkey was small. MATLAB 'nnmf'

function was used for NMF.

To determine whether the weights of muscle synergies changed after TT, the

following procedure was used. First, the average of each pre-TT spatial synergy

was calculated and used as control data to calculate the cosine distance for all

post-TT spatial synergies. This generated four distance relationships for each

pre-TT synergy. If the change in spatial synergy before and after TT was small,

then only one of the four cosine distances (e.g., synergy A before TT and synergy

A after TT) should be significantly smaller. Two-way ANOVA was conducted

using the type of synergy and session as factors for the cosine distance. The

significance of synergy pairs was identified using the Bonferroni post-hoc test.

Cross-correlation analysis

Cross-correlation analysis was performed to examine pre- and post-surgery EMG

signals as well as temporal activation coefficients of extracted muscle synergies.

Cross-correlation coefficients were calculated using MATLAB (MathWorks,

corrcoef.m function). These were plotted for different muscles and muscle

synergy activation coefficients over experimental sessions for comparison.

Based on cross-correlation analysis, five 'landmark days' were identified for

further analysis (Fig. 6P). These days represent distinct stages in the recovery

process and excluded pre-surgery control data. The first landmark day was

chosen from one of the initial recording sessions after TT surgery (days 29 and

22 in monkeys A and B, respectively) when the cross-correlation coefficients had

changed significantly compared with control data. The second day was then

chosen from a time period just before the switch-back, when the cross-correlation

coefficients had started to return to their original values (days 64 and 36). Another

day was then picked from the period when the coefficients were still changing

significantly (days 69 and 44) and before starting to saturate. At this point the next

day was defined (days 79 and 48). Finally, the final landmark day was chosen

from one of the last recording sessions when the behavior had fully recovered

(days 99 and day 64 for monkeys A and B, respectively).

Behavioral analysis

To examine behavioral recovery, the duration, onset, and offset of object and

food grasps were analyzed. Event times extracted from the video analysis were

used for alignment of EMG data and subsequent cross-correlation and synergy

analyses, along with recorded event TTL signals. For each experimental session,

the first 20 trials of each video recording were analyzed. In Monkey A, the

following events were detected and in-between times stored in ms: touch onset,

pull onset, and hold onset (resulting in grasp duration and pull time) for object 1

and object 2. In Monkey B, the following events were detected and in-between

times stored: touch onset, pull onset, and hold onset (resulting in grasp duration

and pull time) for object 1; and for the food grasp component of the task, LED

onset, food-touch onset, food-lift onset, LED offset, and movement end (contact

of food with mouth).

Contact times (touch onset ➔ pull onset) for Monkey A were plotted in ms (mean

± SD) over number of days from tendon surgery for object 1. For Monkey B,

contact times with food (food-touch ➔ food-lift onset) were plotted in ms (mean

± SD).

ANOVA was performed to compare control data recorded before and after tendon

surgery. In total, 5 and 3 control sessions vs. 35 and 22 experimental sessions

were used for analyses in monkeys A and B, respectively.

To quantify the observed aberrant reaching behavior, video footage was

analyzed and the means of 10 consecutive trials calculated for each session as

follows. For Monkey A, time spent within a specific spatial window covering the

area behind and between the two objects was measured; the monkey passed

through this space almost exclusively while exhibiting the impairment (see Fig.

3E, crossing of the yellow dotted line). In Monkey B, the time spent in contact with

the rear plate of the object was measured (see Fig. 4G).

Joint kinematics analysis

To determine whether compensatory movements occurred, steady state changes

in joint angles were examined before and after recovery. Using DeepLabCut

(Mathis et al., 2018; Nath et al., 2019), key points on the fingers and wrists were

tracked from experimental videos. From the obtained key points, the vector v_1

connecting the wrist and fingers and vector v_2 connecting the arm and wrist were

estimated. The joint angle θ was calculated as follows:

888
$$cos(\theta) = \frac{v_2 \cdot v_3}{|v_2|\,|v_3|} \rightarrow \theta = cos^{-1}\left(\frac{v_2 \cdot v_3}{|v_2|\,|v_3|}\right)$$

889
890

891   **DATA, MATERIALS, AND SOFTWARE AVAILABILITY**

893   Matlab scripts and raw data will be made available online. All study data are

894   included in the article and/or SI Appendix.

ACKNOWLEDGEMENTS:

We would like to express our sincere gratitude to Drs. Francisco Valero-Cuevas and Andrea d'Avella for their invaluable comments and constructive suggestions on the earlier version of the manuscript, and Drs Nicholas Schweighofer and Gerald E. Loeb for stimulating discussion. We would also like to thank Dr. Naomichi Ogihara and Prof. Toshiyasu Nakamura for their valuable insights and technical advice on tendon surgery in the hand and forearm of primates. We are grateful to Drs. Kumiko Oida and Chika Sasaki for their medical care of the experimental animals, vital help during surgery, and training of the animals. Lastly, we would like to thank Masahi Koizumi for designing parts of the experimental setup and Drs. Joachim Confais, Shinji Kubota, Saeka Tomatsu, Tatsuya Umeda, and Natsumi Uchida for their support and assistance during the proceeding pilot study. We thank Rachel James, PhD, from Edanz (https://jp.edanz.com/ac) for editing a draft of this manuscript.

This work was supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan (26120003, 26250013, 15K21754, 19H05724, 19H01092, 23H05488, and 24K21313), the Japan Agency for Medical Research and Development

REFERENCES

Berger DJ, Gentner R, Edmunds T, Pai DK, d'Avella A (2013) Differences in

Adaptation Rates after Virtual Surgeries Provide Direct Evidence for

Modularity. The Journal of Neuroscience 33:12384-12394.

Berniker M, O'Brien MK, Kording KP, Ahmed AA (2013) An Examination of the

Generalizability of Motor Costs. PLOS ONE 8:e53759.

Bizzi E, Cheung VC (2013) The neural origin of muscle synergies. Frontiers in

Computational Neuroscience Volume 7 - 2013.

Bowlus TH, Lane RD, Stojic AS, Johnston M, Pluto CP, Chan M, Chiaia NL,

Rhoades RW (2003) Comparison of reorganization of the somatosensory

system in rats that sustained forelimb removal as neonates and as

adults. J Comp Neurol 465:335-348.

Cash DJW, Jones JWM (2011) The role of tenodesis in surgery of the upper

limb. The Journal of Bone & Joint Surgery British Volume 93-B:285-292.

Cheung VC, Turolla A, Agostini M, Silvoni S, Bennis C, Kasi P, Paganoni S,

Bonato P, Bizzi E (2012) Muscle synergy patterns as physiological

markers of motor cortical damage. Proc Natl Acad Sci U S A 109:14652-

14656.

d'Avella A, Saltiel P, Bizzi E (2003) Combinations of muscle synergies in the

construction of a natural motor behavior. Nature Neuroscience 6:300-

308.

Davidson PR, Wolpert DM (2003) Motor learning and prediction in a variable

environment. Curr Opin Neurobiol 13:232-237.

De Nunzio AM, Schweisfurth MA, Ge N, Falla D, Hahne J, Gödecke K, Petzke

F, Siebertz M, Dechent P, Weiss T, Flor H, Graimann B, Aszmann OC,

Farina D (2018) Relieving phantom limb pain with multimodal sensory-

motor training. J Neural Eng 15:066022.

Dieterich AV, Botter A, Vieira TM, Peolsson A, Petzke F, Davey P, Falla D (2017)

Spatial variation and inconsistency between estimates of onset of muscle

activation from EMG and ultrasound. Sci Rep 7:42011.

Edwards MJ, Adams RA, Brown H, Pareés I, Friston KJ (2012) A Bayesian

account of 'hysteria'. Brain 135:3495-3512.

Flash T, Sejnowski TJ (2001) Computational approaches to motor control. Curr

Opin Neurobiol 11:655-662.

Fleury L, Prablanc C, Priot AE (2019) Do prism and other adaptation paradigms

really measure the same processes? Cortex 119:480-496.

Foell J, Bekrater-Bodmann R, Diers M, Flor H (2014) Mirror therapy for

phantom limb pain: brain changes and the role of body representation.

Eur J Pain 18:729-739.

Gaetz W, Dockstader C, Furlong PL, Amaral S, Vossough A, Schwartz ES,

Roberts TPL, Scott Levin L (2023) Somatosensory and motor

representations following bilateral transplants of the hands: A 6-year

longitudinal case report on the first pediatric bilateral hand transplant

patient. Brain Res 1804:148262.

Gardenier J, Garg R, Mudgal C (2020) Upper Extremity Tendon Transfers: A

Brief Review of History, Common Applications, and Technical Tips. Indian

J Plast Surg 53:177-190.

Gijsberts A, Bohra R, Sierra González D, Werner A, Nowak M, Caputo B, Roa

MA, Castellini C (2014) Stable myoelectric control of a hand prosthesis

using non-linear incremental learning. Front Neurorobot 8:8.

Green HJ (1997) Mechanisms of muscle fatigue in intense exercise. J Sports

Sci 15:247-256.

Hatem SM, Saussez G, Della Faille M, Prist V, Zhang X, Dispa D, Bleyenheuft Y

(2016) Rehabilitation of Motor Function after Stroke: A Multiple

Systematic Review Focused on Techniques to Stimulate Upper Extremity

Recovery. Front Hum Neurosci 10:442.

Hoffmann G, Kamper DG, Kahn JH, Rymer WZ, Schmit BD (2009) Modulation

of stretch reflexes of the finger flexors by sensory feedback from the

proximal upper limb poststroke. J Neurophysiol 102:1420-1429.

Hospod V, Aimonetti J-M, Roll J-P, Ribot-Ciscar E (2007) Changes in Human

Muscle Spindle Sensitivity during a Proprioceptive Attention Task. The

Journal of Neuroscience 27:5172-5178.

Hunter DJ, Eckstein F (2009) Exercise and osteoarthritis. J Anat 214:197-207.

Jami L (1992) Golgi tendon organs in mammalian skeletal muscle: functional

properties and central actions. Physiol Rev 72:623-666.

Kakavas G, Brilakis E, Papatzikou M, Malliaropoulos N, Mazeas J, Forelli F

(2025) Reverse Linear Neuro Periodization Model for Rehabilitation After

Arthroscopic Rotator Cuff Repair: A Narrative Review. Clinics and

Practice 15:105.

Kitago T, Krakauer JW (2013) Motor learning principles for neurorehabilitation.

Handb Clin Neurol 110:93-103.

Kitago T, Ryan SL, Mazzoni P, Krakauer JW, Haith AM (2013) Unlearning

versus savings in visuomotor adaptation: comparing effects of washout,

passage of time, and removal of errors on motor memory. Front Hum

Neurosci 7:307.

Kitazawa S, Kohno T, Uka T (1995) Effects of delayed visual information on the

rate and amount of prism adaptation in the human. J Neurosci 15:7644-

7652.

Lee DD, Seung HS (1999) Learning the parts of objects by non-negative matrix

factorization. Nature 401:788-791.

Levin MF, Weiss PL, Keshner EA (2015) Emergence of virtual reality as a tool

for upper limb rehabilitation: incorporation of motor control and motor

learning principles. Phys Ther 95:415-425.

Loeb GE (1999) Asymmetry of hindlimb muscle activity and cutaneous reflexes

after tendon transfers in kittens. J Neurophysiol 82:3392-3405.

Luauté J, Schwartz S, Rossetti Y, Spiridon M, Rode G, Boisson D, Vuilleumier P

(2009) Dynamic changes in brain activity during prism adaptation. J

Neurosci 29:169-178.

Maier M, Ballester BR, Verschure PFMJ (2019) Principles of Neurorehabilitation

After Stroke Based on Motor Learning and Brain Plasticity Mechanisms.

Frontiers in Systems Neuroscience Volume 13 - 2019.

Martin TA, Keating JG, Goodkin HP, Bastian AJ, Thach WT (1996) Throwing

while looking through prisms. II. Specificity and storage of multiple gaze-

throw calibrations. Brain 119 (Pt 4):1199-1211.

Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M

(2018) DeepLabCut: markerless pose estimation of user-defined body

parts with deep learning. Nature Neuroscience 21:1281-1289.

Mercuri E, Muntoni F (2013) Muscular dystrophies. Lancet 381:845-860.

Morton SM, Bastian AJ (2006) Cerebellar contributions to locomotor adaptations

during splitbelt treadmill walking. J Neurosci 26:9107-9116.

Murata Y, Higo N, Oishi T, Yamashita A, Matsuda K, Hayashi M, Yamane S

(2008) Effects of motor training on the recovery of manual dexterity after

primary motor cortex lesion in macaque monkeys. J Neurophysiol

99:773-786.

Mussa-Ivaldi FA, Bizzi E (1995) The Modular Organization of Motor Control:

What Frogs Can Teach Us About Adaptive Learning. IFAC Proceedings

Volumes 28:413-418.

Nath T, Mathis A, Chen AC, Patel A, Bethge M, Mathis MW (2019) Using

1033   DeepLabCut for 3D markerless pose estimation across species and

1034   behaviors. Nature Protocols 14:2152-2176.

1035 Nordez A, Gallot T, Catheline S, Guével A, Cornu C, Hug F (2009)

1036   Electromechanical delay revisited using very high frame rate ultrasound.

1037   J Appl Physiol (1985) 106:1970-1975.

1038 Nudo RJ, Wise BM, SiFuentes F, Milliken GW (1996) Neural substrates for the

1039   effects of rehabilitative training on motor recovery after ischemic infarct.

1040   Science 272:1791-1794.

1041 Overduin SA, d'Avella A, Roh J, Bizzi E (2008) Modulation of Muscle Synergy

1042   Recruitment in Primate Grasping. The Journal of Neuroscience 28:880-

1043   892.

1044 Petitet P, O'Reilly JX, O'Shea J (2018) Towards a neuro-computational account

1045   of prism adaptation. Neuropsychologia 115:188-203.

1046 Popa LS, Streng ML, Hewitt AL, Ebner TJ (2016) The Errors of Our Ways:

1047   Understanding Error Representations in Cerebellar-Dependent Motor

1048   Learning. Cerebellum 15:93-103.

1049 Power JD, Schlaggar BL (2017) Neural plasticity across the lifespan. Wiley

1050   Interdiscip Rev Dev Biol 6.

Ranganathan R, Adewuyi A, Mussa-Ivaldi FA (2013) Learning to be lazy:

exploiting redundancy in a novel task to minimize movement-related

effort. J Neurosci 33:2754-2760.

Sattin D, Parma C, Lunetta C, Zulueta A, Lanzone J, Giani L, Vassallo M,

Picozzi M, Parati EA (2023) An Overview of the Body Schema and Body

Image: Theoretical Models, Methodological Settings and Pitfalls for

Rehabilitation of Persons with Neurological Disorders. Brain Sci 13.

Schärli A, Hecht H, Mast FW, Hossner EJ (2024) How spotting technique affects

dizziness and postural stability after full-body rotations in dancers. Hum

Mov Sci 95:103211.

Schoultz TW, Swett JE (1972) The fine structure of the Golgi tendon organ. J

Neurocytol 1:1-26.

Slawinska U, Kasicki S (2002) Altered Electromyographic Activity Pattern of Rat

Soleus Muscle Transposed into the Bed of Antagonist Muscle. The

Journal of Neuroscience 22:5808-5812.

Smith MA, Shadmehr R (2005) Intact ability to learn internal models of arm

dynamics in Huntington's disease but not cerebellar degeneration. J

Neurophysiol 93:2809-2821.

Smith MA, Ghazizadeh A, Shadmehr R (2006a) Interacting Adaptive Processes

with Different Timescales Underlie Short-Term Motor Learning. PLOS

Biology 4:e179.

Smith MA, Ghazizadeh A, Shadmehr R (2006b) Interacting adaptive processes

with different timescales underlie short-term motor learning. PLoS Biol

4:e179.

Smith SA, Mitchell JH, Garry MG (2006c) The mammalian exercise pressor

reflex in health and disease. Exp Physiol 91:89-102.

Sperry RW (1940) The functional results of muscle transposition in the hind limb

of the rat. Journal of Comparative Neurology 73:379-404.

Sperry RW (1942) Transplantation of motor nerves and muscles in the forelimb

of the rat. Journal of Comparative Neurology 76.

Sugita Y (1996) Global plasticity in adult visual cortex following reversal of

visual input. Nature 380:523-526.

Takei T, Seki K (2010) Spinal interneurons facilitate coactivation of hand

muscles during a precision grip task in monkeys. J Neurosci 30:17041-

17050.

Takei T, Confais J, Tomatsu S, Oya T, Seki K (2017) Neural basis for hand

muscle synergies in the primate spinal cord. Proceedings of the National

Academy of Sciences 114:8643-8648.

Takeuchi N, Izumi S (2012) Maladaptive plasticity for motor recovery after

stroke: mechanisms and approaches. Neural Plast 2012:359728.

Tannenbaum J, Bennett BT (2015) Russell and Burch's 3Rs then and now: the

need for clarity in definition and purpose. J Am Assoc Lab Anim Sci

54:120-132.

Thompson AK, Wolpaw JR (2014) Operant conditioning of spinal reflexes: from

basic science to clinical therapy. Front Integr Neurosci 8:25.

Trewartha KM, Garcia A, Wolpert DM, Flanagan JR (2014) Fast but fleeting:

adaptive motor learning processes associated with aging and cognitive

decline. J Neurosci 34:13411-13421.

Vallbo AB, Hagbarth KE, Torebjörk HE, Wallin BG (1979) Somatosensory,

proprioceptive, and sympathetic activity in human peripheral nerves.

Physiol Rev 59:919-957.

van Vliet P, Carey LM, Turton A, Kwakkel G, Palazzi K, Oldmeadow C, Searles

A, Lavis H, Middleton S, Galloway M, Dimech-Betancourt B, O'Keefe S,

Tavener M (2023) Task-specific training versus usual care to improve

upper limb function after stroke: the "Task-AT Home" randomised

controlled trial protocol. Frontiers in Neurology 14.

Vanhoof MJM, van Leeuwen T, Galletta L, Vereecke EE (2021) The forearm and

hand musculature of semi-terrestrial rhesus macaques (Macaca mulatta)

and arboreal gibbons (fam.Hylobatidae). Part II. Quantitative analysis.

Journal of Anatomy 238:321-337.

Villalta JI, Landi SM, Fló A, Della-Maggiore V (2013) Extinction Interferes with

the Retrieval of Visuomotor Memories Through a Mechanism Involving

the Sensorimotor Cortex. Cerebral Cortex 25:1535-1543.

Walker JG, Jackson HJ, Littlejohn GO (2004) Models of adjustment to chronic

illness: using the example of rheumatoid arthritis. Clin Psychol Rev

24:461-488.

Wester K, Hove LM, Barndon R, Craven AR, Hugdahl K (2018) Cortical

Plasticity After Surgical Tendon Transfer in Tetraplegics. Front Hum

Neurosci 12:234.

Winterbottom L, Nilsen DM (2024) Motor Learning Following Stroke:

Mechanisms of Learning and Techniques to Augment Neuroplasticity.

Phys Med Rehabil Clin N Am 35:277-291.

Wolpert DM (1997) Computational approaches to motor control. Trends Cogn

Sci 1:209-216.

Yan Y, Sobinov AR, Bensmaia SJ (2022) Prehension kinematics in humans and

macaques. J Neurophysiol 127:1669-1678.

Yumiya H, Larsen KD, Asanuma H (1979) Motor readjustment and input-output

relationship of motor cortex following cross-connection of forearm

muscles in cats. Brain Research 177:566-570.

Zajac FE (1992) How musculotendon architecture and joint geometry affect the

capacity of muscles to move and exert force on objects: a review with

application to arm and forearm tendon transfer design. J Hand Surg Am

17:799-804.

Zeiler SR, Krakauer JW (2013) The interaction between training and plasticity in

the poststroke brain. Curr Opin Neurol 26:609-616.

Zemková E (2022) Physiological Mechanisms of Exercise and Its Effects on

Postural Sway: Does Sport Make a Difference? Front Physiol 13:792875.

1138

1139

1140

1141

Figure 1: Muscle anatomy of the macaque forearm and the tendon transfer procedure.

A schematic of the primary forearm muscles involved in the study, showing both the dorsal and volar views. The diagram illustrates the surgical crossed tendon transfer of the extensor digitorum communis (EDC) and flexor digitorum superficialis (FDS) tendons. All labeled muscles were implanted with EMG electrodes. Muscle Abbreviations: BRD: brachioradialis, ECR: extensor carpi radialis, ECU: extensor carpi ulnaris, ED2,3: extensor digitorum-2,3, ED4,5: extensor digitorum-4,5, EPL: extensor pollicis longus, FCR: flexor carpi radialis, FCU: flexor carpi ulnaris, FDP: flexor digitorum profundus, PL: palmaris longus, PT: pronator teres, *(The deltoid (DEL) muscle was also implanted in Monkey B but is not shown as it is a shoulder muscle.)*

1154

1155

1156

Figure 2: *Long term confirmation of tendon surgery effectiveness to alter mechanical properties.*

(A) Set-up for the ultrasound measurement and video recordings of the stimulation induced movements of the EDC and FDS tendons. (B) Sonogram of the FDS muscle and its intramuscular tendons. Left side (B-mode, i.e., brightness mode) shows the still image of the monkeys forearm at a given point in time. Right side (M-mode, i.e., motion mode) shows the stagged images of the FDS tendon displacement induced by muscle stimulation (50mA). White arrows demarcate the FDS tendon which was used for the measurement. Grayscale gradations correspond to tissue densities: hyperechoic regions (white) denote denser structures like surface of bones and tendons, while hypoechoic areas (black) signify less dense tissues such as adipose tissue and musculature. Inset demonstrates the area measurement. The area of the displacement waves was measured in the M-mode, representing the strength of muscle contraction. We measured the duration (a, sec) and amplitude (b, cm) of three waves and calculated the average. Area = a*b/2(cm/sec) for days 0, 7 and 105 after tendon transfer. (C) Areas under the wave measured in the M-mode for 3 experimental days (0, 7 and 105 days post-TT) and regression lines in red and blue for FDS

and EDC, respectively. R2>0.5 for FDS. The data suggested that muscle contractions induced by direct electrical stimulation were nearly constant. (D) Markers placed on the index, middle and ring finger nails (A) were used to measure finger displacement in xyz-dimensions. We calculated the sum of the Euclidean distances of each marker from the origin of the 3D coordinate system as a scalar quantity. Observing the movement along the Z-axis, it became reversed post-surgery indicating a reversal from finger flexion to extension due to tendon transfer (D, blue = pre-TT at surgery day; dark brown = post-TT at surgery day; light brown = 1wk post-TT; red = 3wks post-TT). The scalar quantity of the fingers during muscle stimulation did not change much at day 0, 7 and 105 days (E), suggesting that there was no tendon rupture or slackening of the tendons postoperatively (EDC stimulation, left; FDS stimulation, right).

1191

Figure 3: *Experimental set-up, task sequence and typical EMG (monkey A).*

(A) Schematic of the behavioral sequence (hook → grasp → release). (B) Schematic of the task object using a rod requiring monkey A to perform a controlled grasp. (C) Typical EMG traces of a control session (high-pass filtered) for all recorded muscles. Gray boxes represent the task sequence. Obj 1 ON: start of the hold period of object 1. Obj 1 OFF: end of the hold period of object 1, i.e., object release. Obj 2 ON: start of the hold period of object 2. Obj 2 OFF: end of the hold period of object 2, i.e., object release. Tendons of the muscles marked with * were cross-transferred. (D) Rectified and smoothed EMG for all recorded

1202 muscles (average for one recording session; amplitude [μV] over task sequence
1203 [%]). Horizontal bars illustrate the corresponding behavioral periods, red vertical
1204 lines indicate peak amplitude for each muscle). (E) The time the monkey spent
1205 on the left side of the yellow dotted line while moving from object 1 to object 2
1206 was measured and used to quantify the mal-adaptive behavior.
1207
1208

1209

Figure 4: Experimental set-up, task sequence and typical EMG (monkey B).

(A) Schematic of the task sequence (picking up food). (B) Schematic of the task requiring monkey B to pick up food from a groove allowing for a more natural grasp. (C) Typical EMG traces of a control session (high-pass filtered) for all recorded muscles. Gray boxes represent the task. Obj 1 ON: start of the hold period of object 1. Obj 1 OFF: end of object 1's hold period, i.e. object release. LED ON: approximate start of the food touch. LED OFF: approximate time of food retrieval. Tendons of the muscles marked with * were cross-transferred. (D) rectified and smoothed EMG for all recorded muscles (average for one recording session; amplitude [μV] over task sequence [%]). Horizontal bars illustrate the corresponding behavioral periods, red vertical lines indicate peak amplitude for each muscle). (E) Development of the metacarpophalangeal joint (MCP) and wrist joint angle [deg] over task range [%]. Decreasing and increasing angles indicate extension or flexion of the MCP or wrist angle, respectively. Data are aligned on LED onset (food retrieval) and indicate consistency between

1225   behavioral observation (A), EMG (C-D) and kinematics (E). (F) Illustration of joint

1226   angle measurement using a still picture of the monkey while performing the task.

1227   (G) Example for mal-adaptive behavior in monkey B. The time the monkey spent

1228   in contact with or behind the object plate was measured and used to quantify the

1229   mal-adaptive behavior.

1230

1231

Figure 5: *Behavioral recovery is preceded by maladaptive motor control in both monkeys.*

(A) Grip formation times. Plotted is the average time (mean ± SD, n=20 trials) from initial touch to action onset (pull or lift) for Monkey A and Monkey B. Grip formation times were significantly elevated immediately post-surgery before returning to baseline. (B) Quantification of aberrant reaching. Plotted is the average time (mean ± SD, n=10 trials) spent executing aberrant movements. For Monkey A, this was the duration spent moving behind the target (see Fig. 3E); for Monkey B, it was the duration in contact with the object's rear plate (see Fig. 4G). These behaviors were prominent in the early post-surgical phase and diminished over time. Filled squares indicate a significant difference from pre-TT baseline ($p < 0.05$, two-sample t-test). All data are plotted over days relative to the tendon transfer (TT) surgery (day 0).

1245

1246

Figure 6: A *Two-Phase Adaptation of Muscle Activity is Observed in Both Monkeys Following Tendon Transfer.*

The figure compares EMG activity profiles and cross-correlation analyses for Monkey A (left) and Monkey B (right), demonstrating a consistent pattern of motor adaptation consisting of an initial functional swap followed by a later reversion. A representative non-transferred muscle is included for each monkey. Monkey A (A-J): Profiles are aligned on object release. Monkey B (K-Q): Profiles are aligned on food touch. Filled triangles (▼) indicate peak activity occurring during finger extension, while open triangles (▽) indicate peak activity occurring during finger flexion. Small triangles on correlation plots denote landmark days.

1257

1258

Figure 7: Muscle Synergy Compositions Remain Stable While Their Activation Timings Show a Two-Phase Adaptation.

The figure presents the spatial muscle weights and temporal activation coefficients for the primary finger flexor (Synergy A) and extensor (Synergy B) for Monkey A (A-D) and Monkey B (E-H). (A, B, E, F) Spatial Synergy Weights: The muscle weight contributions for each synergy are shown for the pre-surgery control period and for five post-surgery landmark days, demonstrating that synergy structure was preserved. (C, D, G, H) Temporal Activation Coefficients: The activation profiles show the same two-phase adaptation seen in individual muscles: an initial swap followed by a later reversion. Symbols and alignment are as described in Fig. 6.

1271

Figure 8: Analysis of Secondary Muscle Synergies and Variance Accounted For (VAF).

This figure provides supporting data for the synergy analysis, including the VAF plots justifying the use of four synergies, and the detailed analysis of the secondary Synergies C and D for both Monkey A and Monkey B. (A-B) Variance Accounted For (VAF). The plots show the cumulative variance in the EMG data explained by an increasing number of synergies for Monkey A (left) and Monkey B (right). In both original datasets (blue lines), four synergies were sufficient to account for more than 80% of the variance (demarcated by the black horizontal line), justifying the dimensionality of the synergy model. VAF for corresponding

shuffled data is shown for comparison (red lines). (C-F) Analysis of Synergy C (Wrist Flexor Synergy). This panel group shows the spatial weights (C, D) and temporal activation coefficients (E, F) for Synergy C in Monkey A and Monkey B, respectively. Spatial Weights (C, D): Pre-surgery control profiles (left sub-panels) are compared to post-surgery landmark days (right sub-panels), demonstrating that the muscle contributions to this synergy remained largely stable. Temporal Activation (E, F): In Monkey A, the temporal profile (C) shows a notable increase in activation during the late adaptation phase, consistent with its recruitment for a compensatory strategy. The adaptive pattern for Monkey B (E) was less distinct. (G-J) Analysis of Synergy D. This panel group shows the spatial weights (G, H) and temporal activation coefficients (I, J) for Synergy D in Monkey A and Monkey B. This synergy, which contributes to wrist extension, exhibited relatively minor and inconsistent changes following the tendon transfer in both animals. For all temporal plots, profiles are aligned on object release (Monkey A) or food touch (Monkey B) at 0% task range. Post-surgery plots show data from selected landmark days as indicated by the color legend.

Figure 9: Cross-correlation analysis reveals the two-phase adaptation of primary synergy activation.

Cross-correlation coefficients for the primary flexor (Synergy A) and extensor (Synergy B) synergies are plotted over post-surgery days for Monkey A (A-D) and Monkey B (E-H). For each synergy, the activation pattern was cross-correlated with its own pre-surgery profile (A, B, E, F) and with that of its antagonist (C, D, G, H). The results quantitatively demonstrate the swap-and-revert pattern in both animals. The // represents the recovery period.

1308

1309

Figure 10: Cross-correlation analysis of secondary synergy activation.

Cross-correlation coefficients for the secondary synergies (C and D) are plotted
over post-surgery days for Monkey A (A-D) and Monkey B (E-H). For each
synergy, the activation pattern was cross-correlated with the pre-surgery profiles
of Synergy A (red lines) and Synergy B (blue lines) to reveal their changing
relationships to the primary flexor and extensor commands.

1316

1317

Figure 11: Aggregated and averaged EMG (aaEMG) reveals synergy-specific adaptive patterns.

Aggregated and averaged electromyography (EMG) activities for the main contributing muscles of each of the four synergies for Monkey A (A-H) and Monkey B (I-P). For each recording session, EMG activity within a ±15% task range window was summed for the contributing muscles and then averaged. The left columns for each monkey (A-D, I-L) show the aaEMG over all post-surgery days. The right columns (E-H, M-P) show the mean (± SEM) aaEMG for the pre-surgery period ("pre") and the five selected landmark days. Monkey A (A-H) The contributing muscles for each synergy were: Synergy A: flexor digitorum superficialis (FDS) and flexor carpi ulnaris (FCU). Synergy B: extensor digitorum communis (EDC), extensor digitorum-2,3 (ED23), and extensor carpi ulnaris (ECU). Synergy C: palmaris longus (PL) and flexor carpi radialis (FCR). Synergy D: flexor digitorum profundus (FDP), brachioradialis (BRD), and extensor carpi radialis (ECR). Monkey B (I-P) The contributing muscles for each synergy were: Synergy A: flexor digitorum profundus (FDP) and palmaris longus (PL). Synergy B: extensor digitorum communis (EDC), extensor digitorum-2,3 (ED23) and

extensor digitorum-4,5 (ED45). Synergy C: extensor carpi radialis (ECR), brachioradialis (BRD), and deltoid (DEL). Synergy D: extensor carpi ulnaris (ECU). Vertical colored bars on the time-series plots indicate the landmark days shown in the bar plots. The // on the time axis represents the recovery period during which no recordings were taken. Asterisks on bar plots indicate a significant difference from the pre-surgery control period ($p < 0.01$, *$p < 0.001$, **$p < 0.0001$; two-sample t-test with Bonferroni correction, $\alpha = 0.01$).

1344

Figure 12: Kinematic analysis reveals a compensatory tenodesis strategy in Monkey B.

Changes in joint angles at the wrist and metacarpophalangeal (MCP) joint for Monkey B (mean of 20 trials ±SD for each landmark day, taken 83ms before food touch). (A) The MCP joint angle increased post-surgery, indicating greater finger extension. (B) Concurrently, the wrist joint angle decreased, indicating wrist flexion. This coordinated movement pattern is characteristic of an active tenodesis effect, suggesting it was used as a compensatory strategy to achieve finger extension. Statistical analysis confirmed these changes were significant ($p < 0.0001$, $\alpha = 0.01$, ANOVA).

1355

Figure 13: A Proposed Model of Multi-Timescale Adaptation Following Tendon Transfer.

This schematic illustrates the hypothesized interaction between fast and slow adaptive processes driving recovery. The initial Tendon Transfer triggers a rapid but maladaptive 'swap' of motor commands (Fast Adaptation 1), leading to a maladaptive state. During this phase, two slower processes are hypothesized to occur in parallel: a costly 'arms race' within the conflicted synergy (Slow Process A, red curve) and the gradual development of a functional compensatory strategy (Slow Process B, green curve). When the 'arms race' reaches a threshold of unsustainable cost (dashed blue line), a second Fast Adaptation ('Switch-Back', 2) is triggered. This allows for the abandonment of the flawed strategy and the adoption of a stable, 'good enough' solution, which is now supported by the newly learned compensatory strategy. The gray line represents the observed neural data (e.g., cross-correlation coefficients of electromyography and temporal activation coefficients of muscle synergies), which reflects this two-phase process.