Multi-timescale neural adaptation underlying long-term musculoskeletal reorganization.

(Multi-timescale neural adaptation**)**

*Roland Philipp[1,2], Yuki Hara[3], Naohito Ohta[1,2], Naoki Uchida[1,2], Tomomichi Oya[1,4], Tetsuro Funato[2], and *Kazuhiko Seki[1]

[1]National Center of Neurology and Psychiatry, Department of Neurophysiology, 4-1-1 Ogawa-Higashi, Kodaira, Tokyo 1878502, Japan
[2]University of ElectroCommunications, Graduate School of Informatics and Engineering, Dept. of Mechanical and Intelligent Systems Engineering. W2-217, 1-5-1 Chofugaoka, Chofu, Tokyo, Japan
[3]National Center of Neurology and Psychiatry, Department of Orthopaedic Surgery, 4-1-1 Ogawa-Higashi, Kodaira, Tokyo 1878502, Japan
[4]Western Institute for Neuroscience, University of Western Ontario, London, N6A 3K7, Canada

*Corresponding authors:
Kazuhiko Seki and Roland Philipp
Email: seki@ncnp.go.jp and roland@ncnp.go.jp

Author Contributions: R.P. and K.S. designed all experiments and wrote the paper. R.P., Y.H., N.U., and N.O. performed the experiments and analyzed the data. R.P., Y.H., T.O., and K.S. performed the surgeries. T.F. provided conceptual advice and wrote the paper. All authors reviewed the results and approved the final version of the manuscript.

Conflict of Interest Statement: The authors declare no competing interests.

Number of pages: 74 (96, including supplementary text)
Number of figures: 10 (20, including supplementary figures)
Number of multimedia: 7
Number of words for abstract: 173
Number of words in introduction: 557
Number of words in discussion: 1436

ABSTRACT

The central nervous system (CNS) can effectively control body movements despite environmental changes. While much is known about adaptation to external environmental changes, less is known about responses to internal bodily changes. This study investigates how the CNS adapts to long-term alterations in the musculoskeletal system using a tendon transfer model in non-human primates. We surgically relocated finger flexor and extensor muscles to examine how the CNS adapts its strategy for finger movement control by measuring muscle activities during grasping tasks. Two months post-surgery, the monkeys demonstrated significant recovery of grasping function despite the initial disruption. Our findings suggest a two-phase CNS adaptation process: an initial phase enabling function with the transferred muscles, followed by a later phase abolishing this enabled function and restoring a control strategy that, while potentially less conflicted than the maladaptive state, resembled the original pattern, possibly representing a 'good enough' solution. These results highlight a multi-phase CNS adaptation process with distinct time constants in response to sudden bodily changes, offering potential insights into understanding and treating movement disorders.

SIGNIFICANCE STATEMENT

After major changes to the body's mechanics, the nervous system adapts motor

control using strategies operating on multiple timescales. Our primate tendon

transfer study shows that core muscle synergy groupings remain stable,

reflecting modular control principles. However, the activation of these synergies

changes substantially, with rapid initial shifts followed by slower adjustments over

weeks. This process highlights the fundamental tension the CNS faces in

balancing reliance on stable muscle synergies with the need for flexible control

to manage new mechanics. Understanding this adaptive process offers insights

into neural plasticity and informs potential rehabilitation strategies.

INTRODUCTION:

The central nervous system (CNS) adapts bodily functions in response to various challenges, including external changes like altered gravitational fields or altered sensory feedback (Sugita, 1996; Davidson and Wolpert, 2003; Luauté et al., 2009; Fleury et al., 2019). While such paradigms have provided models for motor adaptation to external perturbations, less is understood about adaptations to internal changes, such as growth (Power and Schlaggar, 2017), muscle fatigue (Green, 1997), postural sway (Zemková, 2022), disorientation from bodily rotations (Schärli et al., 2024), and structural alterations to the musculoskeletal system (Walker et al., 2004). Unlike external changes, which are often transient and predictable, internal changes, particularly structural alterations, require the CNS to remap motor control within the context of the body's altered biomechanics and sensory feedback. Although external tools can alter effective limb dynamics (Imamizu et al., 2000), our focus is on the CNS response to changes in the body's own physical structure. Understanding adaptation to such internal changes is crucial for understanding the pathophysiology of chronic conditions such as osteoarthritis or muscular dystrophy, where progressive musculoskeletal changes lead to significant motor dysfunction (Walker et al., 2004).

To manage the complexity of the motor system, the CNS may utilize modular

building blocks such as muscle synergies, which coordinate groups of muscles

into functional units (d'Avella et al., 2003; Bizzi and Cheung, 2013). Efficient

motor control often relies on muscle synergies, reducing the degrees of freedom

the CNS must control. However, skilled behavior, like dexterous hand movements,

often relies on fractionation – the ability to selectively activate muscles

independently. For instance, precise finger movements like typing or playing a

musical instrument require a high degree of fractionation. This creates a tension:

how does the CNS balance the stability and efficiency of modular synergistic

control with the flexibility needed when the musculoskeletal system undergoes

change? Does adaptation rely on modulating existing synergies or developing

more fractionated control?

Here, we use a tendon transfer (TT) surgery model to explore how animals adjust

limb control following long-term musculoskeletal changes. The TT model offers a

unique perspective on CNS plasticity in response to altered internal physical

environments (Sperry, 1940), contrasting with CNS lesion models where the

injury itself complicates assessing adaptive capabilities. Tendon transfer provides

a unique model to study CNS adaptation to altered musculoskeletal mechanics

because it creates a relatively well-defined and localized change, allowing

researchers to isolate the effects of altered biomechanics from other confounding

factors. Tendon transfer is used in various species, with studies suggesting

different adaptive capacities, ranging from limited adaptation in adult rodents

(Sperry, 1940; Sperry, 1942; Slawinska and Kasicki, 2002; Bowlus et al., 2003)

to more substantial adjustments in cats (Yumiya et al., 1979; Loeb, 1999) and

primates, including humans (Lee and Seung, 1999; Wester et al., 2018; Gaetz et

al., 2023). Adaptability to TT appears enhanced in primates and humans.

However, the organization and temporal progression of CNS adaptations post-

TT remain unclear. This study aimed to identify long-term adaptive mechanisms

of the primate CNS to musculoskeletal changes. We surgically altered the

musculoskeletal system via crossed tendon transfer of finger extensor and flexor

muscle tendons. We focused on determining how these modifications impact

CNS regulation of limb muscle activity during recovery, with finger-grasping task

control serving as the primary metric for assessing functional adaptation. Through

this analysis, we sought to better understand long-term CNS adaptations to

substantial musculoskeletal changes.

RESULTS

In this study, we developed a novel crossed tendon transfer animal model. This involved surgically swapping the tendons of two antagonistic finger muscles in a macaque monkeys' forearm (Fig. 1). The effectiveness and consistency of the surgery was confirmed by measuring: (i) the distance traveled by muscle fibers and their intramuscular tendons in the transferred muscle (Fig. 2B, C); and (ii) the amount of fingertip displacement (Fig. 2D, E). Both measurements were induced by percutaneous electrical stimulation over the transferred muscles (Fig. 2A). For a detailed description of the results of these measurements see the Methods section. Here, we will primarily discuss the adaptations observed by electromyography (EMG) post-tendon transfer (TT), as well as behavioral recovery in the monkey. The two monkeys were assigned slightly different tasks, allowing us to examine controlled grasping in Monkey A and a more natural grasp in Monkey B (refer to the Methods section, and Figs. 2, 3 and S1 for further details). Once the monkeys had mastered their respective tasks, we recorded control sessions and then performed the crossed tendon transfer. Post-surgery, and once the monkeys had fully recovered and were able to perform the task independently (~3–4 weeks), EMG recordings and behavior were examined.

Despite the now reversed roles of the transferred muscles, the monkeys were

able to recover their grasping performance within two months (Fig. 4). Functional

recovery was primarily assessed based on the return of key behavioral metrics,

such as object/food contact time (Fig. 4), to pre-surgical baseline levels, as well

as the animal's consistent ability to perform the task independently without

assistance (see Methods and Supplementary Video legends). However, none of

the parameters measured returned promptly to pre-TT levels. For example,

contact time with the object (Monkey A, Fig. 4A) or food (Monkey B, Fig. 4C) were

much longer immediately after TT than in the first 6 (or 5) sessions post-TT (197.7

± 92.2 vs. 660.6 ± 221 ms; p = 0.014, and 169.7 ± 14 vs. 316.3 ± 31.9 ms; p =

0.016, respectively. Two-sample t-test; mean ± SD; filled squares in Fig. 4). It

took around 10 days for both monkeys to recover back to pre-TT levels (p > 0.05,

two-sample t-test) and fully stabilize (after 20 and 32 days, respectively). All

measured indices eventually returned to pre-TT levels by the end of the

experiment.

Transferred muscles exhibited antagonistic EMG activity post-TT

Initially, we investigated whether EMG activity exhibited any changes post-TT. Given that we interchanged the tendons of an antagonistic muscle pair, functionally rendering the FDS an extensor and the EDC a flexor (as confirmed mechanically, Fig. 2), a biomechanically sound adaptation would be to use the former finger extensor (EDC) for finger flexion, and the former finger flexor (FDS) for finger extension during a grasping task. Our focus was to determine if these expected changes did indeed occur by EMG of both the transferred and non-transferred muscles.

Figure 5 shows a comparison of EMG activity profiles for the two transferred muscles, EDC and FDS, in Monkey A (Fig. 5C–F). Prior to surgery, the control showed distinct and contrasting EMG profiles in both muscles. For instance, peak activity in EDC (▼) occurred long after completion of the grasping action (indicated by vertical dashed lines at 0% task range) at 15% task time (Fig. 5C), coinciding with the animal pre-shaping its hand to prepare for the subsequent grasp. Whereas peak activity in FDS (▽) was observed immediately after completion of the grasping action (0.49%; Fig. 5E). The question we posed was

whether the post-surgery activity of EDC resembled its original profile (without

adaptation to surgery) or the activity of FDS (the anticipated profile following

surgical adaptation), and vice versa. Our findings suggested that EMG activity

patterns largely shifted in a manner consistent with the new mechanical function

imposed by the transfer. Early after TT, peak activity of EDC occurred at -0.58%

task time (Fig. 5D; days 29 [red line] and 64 [orange]), which aligns with peak

activity of FDS prior to TT (Fig. 5E; black line). Also, for FDS, peak activity

occurred at 11% task time (Fig. 5F; ▼), which aligned with peak activity of EDC

(Fig. 5C).

This was corroborated by the results in Monkey B (Fig. S2). Although the EMG

patterns of both muscles varied significantly as the two monkeys performed

different types of grasping actions, we found that the EMG of EDC peaked at -

6.12% before (Fig. S2A; ▼) and 0.35% and 1.05% after TT (Fig. S2B; ▽, days

22 and 36, respectively), closely matching the timing for FDS control data (Fig.

S2C). Note that the corresponding analysis using FDS could not be performed in

Monkey B, since the EMG signal of FDS was lost early after TT (see Methods for

details).

We were able to extend our recordings to 122- and 64-days post-TT for monkeys

A and B, respectively. This allowed us to examine whether the adaptations

observed in the early period were consistent throughout the experimental period.

These additional days (Fig. 5D, F; 69 [cyan], 79 [blue], and 99 [black] post-TT;

see Fig. S3 for recordings from all days) showed that the EMG activity profiles for

both muscles unexpectedly returned to their pre-surgery state. For example, as

already described, the EMG signal of EDC at days 29 and 64 shared the

characteristics of FDS, but by the following day (only 5 days later, day 69 [light

blue] post-TT) already exhibited pre-TT characteristics of EDC, continuing as

such until the end of the recording period (day 79 [blue] and 99 [black]). This was

confirmed for the EDC muscle of Monkey B (Fig. S2A, B; Fig. S3U-AO for

recordings from all days).

Figure 5G illustrates cross-correlation coefficients between the EDC EMG profile

post-TT (EDC-post) and the original EDC control data (EDC-pre) in Monkey A.

Figure 5I shows coefficients for EDC-post and FDS-pre correlations in the same

monkey. The coefficient corresponding to the original profile decreased to

between -0.3 and 0.2 over approximately 65 days (Fig. 5G). In contrast, the

coefficient corresponding to the expected EMG profile exhibited an inverse

pattern, with low coefficients pre-TT and a high coefficient (of 0.9) post-TT (Fig.

5I). Cross-correlation analysis for FDS (Fig. 5H–J) revealed a similar pattern.

Cross-correlation coefficients for FDS-pre- vs. FDS-post-analysis declined from

1 to negative values, only to rebound to their original values within a month (day

79; Fig. 5H). This outcome aligns with our previous observations (Fig. 5D, F), and

were corroborated by the results from Monkey B (Fig. S2). Here, cross-correlation

coefficients dropped from +0.95 to +0.1 for EDC-pre vs. EDC-post comparison,

and increased from -0.5 to 0.9 for the FDS-pre- vs. EDC-post comparison (Fig.

S2D, E). After 42 days post-TT, cross-correlation coefficients started to gradually

increase or decrease to +0.9 and -0.7, respectively. This indicates that the

anticipated initial changes in EMG activity profiles are not permanent. Instead,

they revert to their original patterns two months post-TT.

In summary, both monkeys exhibited a distinct two-phase adaptation following

TT. We define the 'early phase' as the period from initial post-surgical recovery

up to the reversal of the swapped activation patterns (approximately days 20/29

to ~65 post-TT, see Fig. 7), characterized by the transferred muscles/synergies

adopting antagonistic temporal profiles. The 'late phase' encompasses the period

following this switch-back (after ~day 66 post-TT), where original activation

timings were largely restored.

Non-transferred muscles exhibited characteristic changes in EMG activity

post-TT

Interestingly, changes in EMG pattern were not isolated to the transferred

muscles. When the analysis was repeated on non-transferred muscles in both

monkeys (Figs. S3–5), we also found comparable changes in some muscles. This

quantitative analysis is described in the next section; here, we briefly describe

activity characteristics. In Monkey A, similar changes as in the EDC (Fig. 5C, D)

were also found in extensor digitorum-2,3 (ED23), extensor carpi ulnaris (ECU),

and extensor carpi radialis (ECR) (Fig. S4B–D and L–N), i.e., an initial shift of

activity towards pre-surgery FDS that was followed by a later reversion to the

original pattern. The non-transferred flexor carpi ulnaris (FCU) (Fig. S4G, Q)

exhibited changes that mimicked those observed in FDS (Fig. 5E, F). We found

that the non-transferred muscles in Monkey B (Fig. S4U-AO) also exhibited the

changes observed in the transferred muscles. For example, ED23, ED45, ECU,

and flexor digitorum profundus (FDP) (e.g., ED23; Fig. S4AH) exhibited similar

changes as in EDC (Fig. S4AG), initially adopting flexor properties and later

reverting to their original pattern.

In addition, some muscles also exhibited a patten that was incompatible with the

transferred muscles. For example, before TT, PL, and flexor carpi radialis (FCR)

exhibited peak activity at event alignment in monkey A (e.g., PL at object

release; Fig. S4H, I), which again increased gradually over time towards the end

of the shown task range. After TT, however, the amplitude of both peaks initially

decreased drastically, followed by a consistent increase in the amplitude of the

later peak. There was no two-step change observed in the transferred muscles

in any of the subsequent recording days (Fig. S4R, S).

Muscle synergies exhibited antagonistic properties post-TT

Alternation of EMG activity in non-transferred muscle suggests an interesting

possibility: TT might alter the control strategy of coordinated muscle activity for

hand movement by modifying the transferred muscles and their agonists as a

cohesive unit (i.e., muscle synergy). To test this, we conducted a similar analysis

on temporal activation coefficients of muscle synergies. We computed synergy

weights (Fig. 6A–D and S7A-D for Monkey A; and Fig. S6A-D and S7E-H for

Monkey B) and temporal activation coefficients (Fig. 6E–H for Monkey A; and Fig.

S6F–I for Monkey B) from EMGs using non-negative matrix factorization (NMF)

(Lee and Seung, 1999). We extracted four muscle synergies that accounted for

more than 80% (Cheung et al., 2012) of the 'variance accounted for' (VAF) (Figs.

6E and S6E, 85% for both monkeys; the results of the original and shuffled data

sets are shown). Among these synergies, one was a finger flexor synergy and

another was a finger extensor synergy, identified before the crossed tendon

transfer in each monkey (Fig. 6A, B and S7A, B). These were referred to as

synergy A and synergy B, respectively.

We examined if the composition and activation coefficient of each synergy

changed, corresponding to the changes observed in the transferred and non-

transferred muscles. First, we analyzed changes in the composition of these

synergies. The cosine distance for the same synergy pairs (e.g., synergy A of

pre-TT and synergy A of post-TT in all recording days (Fig.S7A, E; monkey A and

B, respectively) was low (mean [± SD]: 0.03 [± 0.03] in Monkey A and 0.09 [±

0.11] in Monkey B) and different pairs (e.g., synergy A of pre-TT and synergy B

of post-TT) was high (mean [± SD]: 0.67 [± 0.08] in Monkey A and 0.61 [± 0.14]

in Monkey B). There was a one-to-one correspondence between synergy pairs

($p < 0.0001$, two-way analysis of variance [ANOVA] due to synergy factor; $p <$

0.01, Bonferroni post-hoc test between synergy of the same pair and different

pair). This was demonstrated in both monkeys (S7A-D and E-H for monkey A and

B, respectively). These results indicate that the composition of these four

synergies and the contribution of each muscle do not change drastically before

and after TT surgery.

Figure 6F–I presents a comparison of the activation coefficients for all four muscle

synergies, before and after the TT procedure (for Monkey B, refer to Fig. S6F–I;

Fig. S8 for all experimental days). Prior to TT, the activation coefficients for the

flexor synergy (synergy A) reached a peak at +0.97% (Fig. 6F). Post-TT, these

coefficients shifted towards later times shortly after surgery (+10.7%), but

eventually returned to their original timing. Meanwhile, the extensor synergy

(synergy B) peaked at 12.62% pre-TT and moved to -0.97% shortly after surgery.

Like the flexor synergy, the extensor synergy also reverted to its original timing in

subsequent sessions (Fig. 6G). These characteristics corresponded well to the

ones found in the transferred muscles (Fig. 5), supporting our hypothesis that the

transferred muscle and its agonist changes as a cohesive unit.

Figure 6H shows the temporal changes for synergy C, a wrist flexor synergy (Fig.

6C and S7C). While two peaks were observed prior to surgery (+1.46% and

+15%), the early post-surgery sessions did not exhibit any significant peaks.

However, later sessions displayed a single peak at +14.6% task time, with an

increasing magnitude as a function of recovery days. These characteristic

changes corresponded well to a patten that was incompatible to the transferred

muscles found in FCR and PL, as described above. Indeed, both muscles

showed highest synergy weights for this synergy (Fig. 6C and S7C). Synergy D

was a wrist extensor synergy (Fig. 6D and S7D), and underwent a single notable

change on day 29, but quickly reverted to its original pattern (Fig. 6H). These

pattens of changes in synergy 3 and 4, both contributed minimally to the EMG of

transferred muscles. This suggests that some non-transferred muscles change

their EMG activity following the mode, which is independent to the one observed

in the transferred muscle and may reinforce the functional adaptation made by

the transferred muscle and its synergists.

Several of these results were also confirmed in Monkey B. Activation coefficients

of extensor synergy B peaked at -8.75% before TT, increasing to +3.5% after TT

(Fig. S6G). This peak activity persisted until the last day (day 69 post-TT), when

it shifted back to an earlier time than -10% of the task range. This pattern

corresponds well to that found in the transferred muscles (EDC; Fig. S2).

Again, the characteristics of these changes in activation coefficients of muscle

synergies were also quantitatively confirmed by cross-correlation analysis of all

four muscle synergies. The coefficients were plotted over the course of

experimental days in relation to tendon surgery (Fig. 7). In this analysis, the

activation coefficient of each synergy was cross-correlated with either the one

from synergy A (finger flexor synergy; Fig. 7C, D, G, H) or synergy B (extensor

synergy; Fig. 7A, B, E, F) before TT. As previously demonstrated for the FDS and

EDC muscles, the temporal activation coefficients of muscle synergies A and B

(flexor and extensor) displayed a distinct pattern. After cross-correlation with their

own control data, both the flexor and extensor synergies showed coefficients of

1 prior to surgery. These became negative for 66 days post-surgery before

reverting close to their original values (Fig. 7B, C). However, after cross-

correlation with control data of the antagonistic muscle synergy (i.e., extensor for

flexor synergy and vice versa, Fig.4A, D), the pattern was reversed. Coefficients

started with negative values prior to TT and shifted to positive coefficients near 1

shortly after surgery. Around 66 days later, the correlation coefficients returned

to negative values. In short, these results mirror our findings between the

transferred muscles (Fig. 5G, H). For synergies C and D, the changes in cross-

correlation coefficients over time were markedly different. For synergy C,

coefficients begin to steadily increase when cross-correlated with control data of

the extensor synergy (synergy B, Fig.4E), and decreased with the flexor synergy

(synergy A, Fig.4G). This suggests an increasing contribution to finger extension.

In contrast, synergy D does not exhibit any specific pattern following TT. We also

found a differential pattern of adaptation between transferred and non-transferred

muscles in Monkey B (Fig. S9F-H).

Overall, we observed early and late changes between two synergy pairs (synergy

A and B) in both monkeys, performing controlled and natural grasping. Another

characteristic change that reinforced finger extension in other synergies was also

observed (i.e., synergy C in Monkey A).

Rationale for the early reverse and late switch-back changes

So far, our results suggest a two-step adaptation following TT: an early and a late

phase. In the initial (early adaptation) phase, the temporal activation patterns of

the transferred muscles and their host synergies largely shifted consistent with

their new mechanical roles (Figs. 2-4). However, the later phase restored pre-TT

EMG profiles and muscle synergies.

This two-step adaptation raises a key question: why? The early changes (where

extensor muscles exhibited flexor properties and vice versa) seem reasonable

and might be unique to CNS adaptation of non-human primates to TT. However,

the later reversal seems puzzling. The monkeys were already performing the task

relatively well (e.g., Fig. 4). Why did the early adaptation need to switch back to

the original pattern?

Here, we hypothesize that the TT forced the monkeys to explore new control

strategies of both transferred and non-transferred muscles (early adaptation),

and the reversing activity was a consequence of the monkey's exploitation of the

previous strategy. This hypothesis assumes that the early adaptation was a

"maladaptation" resulting from a now conflicting mechanical outcome of the

agonistic muscles (i.e., transferred vs. non-transferred muscles). For example, in

monkey A, the finger extensor synergy (synergy B) is composed of EDC, ED23,

and ED45, and their combination remained unchanged after TT and recovery (Fig.

6A–D and S7A–D). However, the transferred EDC became an anatomical finger

flexor, while ED23 and ED45 continued to be anatomical extensors. This led to

paradoxical activation of synergistic muscles with the finger extension synergy

having similar activity to the finger flexion synergy. Therefore, after early

adaptation, monkeys needed to establish a novel strategy to control the altered

musculoskeletal systems. Early maladaptation must have triggered this exploring

process.

This hypothesis appears to be supported by two additional observations. First,

we found unusual behavioral characteristics during target reaching and grasping

after TT (Supplementary Videos S2, S3, and S6). We compared the kinematic

performance of grasping during early adaptation, late adaptation, and pre-TT

periods. Following TT, monkeys exhibited clear functional impairments during the

early adaptation phase before eventually recovering proficiency (Figs. 4).

Behavioral observations revealed specific difficulties impacting fundamental

reach control: Monkey A often required external assistance or support from its

contralateral limb to lift its forearm and displayed prolonged 'explorative' finger

movements before achieving a grasp (Supplementary Video S2); Monkey B

frequently overshot the target and collided with parts of the apparatus

(Supplementary Video S6). Quantitative analysis confirmed these periods of

impaired motor control, showing significantly increased durations for key

behavioral events, such as contact times (Figs. 4A, C), and increased time spent

in non-task-related workspace areas (Figs. 4B, D). These specific maladaptive

reaching behaviors gradually diminished over several weeks, largely resolving by

approximately day 70-75 post-surgery in Monkey A and day 60-65 in Monkey B

(Figs. 4; Supplementary video S4 and S7). This early post-surgical behavior is

considered maladaptive as it reflects substantially reduced movement efficiency

and control precision (e.g., needing assistance, longer task/contact times,

overshooting, ineffective finger movements), likely stemming from the profound

changes in sensorimotor feedback and consequently inaccurate internal models

resulting from the surgically altered musculoskeletal system. These observations

were further supported by measuring the time spent for this maladaptive

behavioral component in both monkeys (Figs. 4B, D). Monkey A exhibited

malcoordinated reaching from day 29 (1978 ± 715 ms) to 73 (93.4 ± 3.7 ms) (see

Methods for additional details). The duration decreased progressively over time.

Monkey B showed malcoordinated reaching from day 20 (103.4 ms) to 62 (107.6

ms). The duration peaked at days 38–46 (167 ±1.4 ms) and then declined.

Notably, the periods dominated by malcoordinated movements coincided with the

observed maladaptive swapping of synergy A and B activity, occurring between

the early and late adaptation phases. This precise temporal coincidence between

the flawed synergy activation strategy and the behavioral impairments provides

strong support for the hypothesis that the maladaptive neural control directly

contributed to the observed impairments in reach control.

Second, we found that the net activity of muscles representing certain muscle

synergies (aggregated average EMG; aaEMG) were unstable during the period

between early (reversal) and late (switch-back) adaptation in their temporal

activation patterns (Fig. 5). In Monkey A (Fig. 8; see Fig. S10 and Extended result

in SI Appendix for Monkey B), muscles of the finger flexor synergy (synergy A,

Fig. 8A) tended to show a decrease in aaEMG post-TT, maintaining this lower

activity level throughout the experiment (p = 0.0004 at day 79, two-sample t-test,

Bonferroni-corrected; α = 0.01; panel E). Because the monkeys performed

successful grasping after TT, this suggests that finger flexion is performed by

muscles other than synergy A. The finger extensor synergy (synergy B, Fig. 8B)

demonstrated a steady increase in aaEMG post-TT until day 64 ($p < 0.0001$, two-

sample t-test, Bonferroni-corrected; $\alpha = 0.01$, panel F), when the activity abruptly

reverted to near its pre-TT level. Increased activity of synergy B suggests a

maladaptive attempt by the monkey to achieve finger flexion. This strategy

involved enhancing activity of synergy B (which includes the EDC muscle, an

atomical flexor) after TT. However, as previously established, this approach

cannot effectively promote finger flexion, leading to eventual abandonment by the

monkey. This 'arms race' dynamic, evidenced by the escalating aaEMG, directly

reflects the mechanical conflict hypothesized to arise from activating the EDC

(now acting as a flexor due to the transfer) within the extensor synergy, further

supporting the idea that this maladaptive state drove the search for alternative

strategies. While directly proving the causal chain is difficult, this hypothesis

provides a plausible framework linking the observed mechanical conflicts,

behavioral deficits, and multi-stage neural adaptation.

Recruitment of novel, compensatory movements aiding finger extension

Muscles of synergy C (wrist flexion synergy) initially showed significantly decreased aaEMG post-TT (days 69 and 79: $p < 0.0001$ and p = 0.001, respectively; two-sample *t*-test, Bonferroni-corrected; α = 0.01, panel G) but showed a gradual increase starting from day 64. This is consistent with the cross-correlation analysis (Fig. 4E), where cross-correlation coefficients between synergy C (wrist flexor) and synergy B (finger extensor) initially dropped post-TT but then first rapidly and then steadily increased over time. Increased reliance on synergy C following TT suggests a compensatory strategy to overcome the difficulty in finger extension, crucial for grasping. This is despite the recruitment of the finger extensor and wrist flexor synergies, which seemed inadequate to achieve effective flexion. The interactions between the cumulative proximal-to-distal joint angles and musculo-tendon length can be complex. In our experiments, as in human patients who receive tendon transfers (Zajac, 1992), those interactions can affect the ranges of finger flexion-extension that can be achieved for a given wrist angle (i.e., *passive and active tenodesis*). Briefly, biomechanically, wrist flexion can enable finger extension (i.e., a dynamic tenodesis where extensor tendons of the fingers cross both the wrist and

interphalangeal joints): Finger extension can cancel out the increase in total

musculo-tendon length produced by wrist flexion. This compensatory interaction

across joints allows for some degree of functional hand use, even in the absence

of active finger movement (Cash and Jones, 2011) (see the Discussion for a

complete description).

Further kinematic analysis of Monkey B (Fig. 9; no analysis on Monkey A due to

low-resolution video images) indicated that this monkey indeed exhibited a similar

active tenodesis as human patients. The angle at the metacarpophalangeal

(MCP) joint, responsible for finger extension, steadily increased (Fig. 9A) after

surgery, indicating finger extension. Concurrently, the wrist joint angle decreased

(Fig. 9B), indicating wrist flexion. This coordinated movement suggests a

compensatory strategy to achieve finger extension during grasping. Statistical

analysis confirmed that MCP and wrist joint angles differed significantly between

pre- and post-TT ($p < 0.0001$, ANOVA; $\alpha = 0.01$). Therefore, we suggest that

deteriorated finger extension after TT was assisted by a compensatory wrist

flexion synergy.

What triggered the late adaptation?

Assuming that TT caused maladaptation and forced the monkeys to explore new

control strategies of both transferred and non-transferred muscles, what triggered

the late adaptation and why did the grasping performance increase after this

adaptation? To address this question, we summarized our experimental results

in a simple model (Fig. 10) consisting of multiple adaptation processes to TT with

different time constants: a slow dynamics adaptation (developing over

weeks/months, e.g., compensatory changes) and a fast dynamics adaptation (i.e.,

processes occurring rapidly over days, such as the switch-back event). In this

model, tendon cross-transfer surgery (black arrow) evoked changes in

coordination of finger muscle activity (i.e., swapping temporal activity of flexor

and extensor synergies; left blue arrow). These changes were sufficient to reach

a threshold (blue dotted line) and trigger (arrow 1) adaptation with slow dynamics

(A and B). This slow process may have different time constants. For example,

synergy B (Monkey A), which comprises EDC and ED23/ED45, consistently

elevated its activity to achieve successful flexion after TT (red line "A"). This 'arms

race' is instigated by conflicting activation of the 'new flexor' (EDC) and its

previous agonists (ED23/45), which still mechanically function as finger extensors.

In contrast, synergy A can reduce its activity as does synergy D (containing FDP),

and accomplish the necessary finger flexion by simply adjusting its timing (Figs.

3, S3, and S4), thereby circumventing the 'arms race'. However, this 'arms race'

observed in synergy B can only persist until a second threshold (red dotted line)

is reached, whereby a further increase in muscle activity becomes unsustainable.

At this point, the second fast adaptation process (blue arrow i.e., switch-back) is

recruited, allowing the monkeys to cease flexing their fingers by excessive

activation of synergy B, thus concluding the 'arms race". Meanwhile, an

alternative, sustainable strategy to perform finger extension has been developed

(Fig. 9). Specifically, the monkey increasingly recruited synergy C (wrist flexors,

PL and FCR) to aid finger extension using the tenodesis effect (wrist flexion),

which was more effective after the 'arms race' was concluded. These implications

can also be applicable to Monkey B (see Extended result in SI Appendix).

DISCUSSION

In our crossed tendon transfer (TT) model, we identified multiple distinct modes of CNS adaptation, each distinguished by their characteristic time course. Notably, changes involving the relocated muscles (FDS and EDC) and their synergists (synergy A and B) exhibited rapid alterations within a few days, during both the early and later phases of adaptation. We propose that the early changes represent a maladaptive phase, while the later restoration of original patterns, coupled with compensation, might represent the achievement of a 'good enough' functional state. In addition, changes involving certain non-relocated muscles were slower and more gradual throughout the adaptation period, suggesting a compensatory role. These slower adaptations appear to mitigate maladaptive responses and contribute to improved performance after achieving a "good enough" adaptation, possibly through exploitation of an active tenodesis effect. We will first explore the potential causes of this distinct adaptation in relocated versus non-relocated muscles, followed by a discussion on the potential interactions between these different adaptation modes.

Relocated muscles and their synergists: early and late adaptation

Given the stability of muscle synergy composition (Fig. S7), the CNS adapted to

relocated muscles (FDS and EDC) by swapping the temporal activation patterns

of the original flexor and extensor synergies, and also their host synergies

(synergy A and B). This suggests that selectively altering single muscles within

co-activated groups is challenging for the CNS, with much daily muscle activity

potentially generated through co-activation modules (Overduin et al., 2008;

Berger et al., 2013; Takei et al., 2017; Madarshahian and Latash, 2022). The

stable muscle synergy composition after surgery supports modular control via

synergies (d'Avella et al., 2003; Bizzi and Cheung, 2013).

However, this adherence to established synergy structures created a

fundamental conflict following crossed tendon transfer. Activating the original

extensor synergy (Synergy B), for instance, now inevitably co-activated the

surgically transferred EDC, which functioned mechanically as a flexor (Fig. 6A-D,

S7). This highlights a tension between maintaining the integrity of these neural

modules and the need for fractionated control – the ability to selectively drive

muscles independently – when faced with altered biomechanics. Consequently,

the CNS opted to swap temporal activation of the muscle synergy, leaving the

synergy itself unchanged.

The observed initial adaptation, where the temporal activation patterns of the

entire flexor and extensor synergies were swapped (Fig. 6E-F, Fig. 7), can be

interpreted in light of this tension. Rather than attempting the potentially complex

task of fractionating control within the extensor synergy (e.g., activating

ED23/ECU without EDC), the CNS appeared to adopt this simpler strategy of

repurposing the entire synergy module based on the new primary function of the

transferred muscle (EDC as flexor, FDS as extensor). However, this initial

adaptation turned out to be maladaptive because this strategy failed functionally

due to the unresolved mechanical conflict within the activated synergy (Figs. 4B,

8B). It involved muscles unaffected by the TT, leading to a steady increase of

antagonistic muscle activity and thus antagonistic proprioceptive feedback in the

non-transferred muscles over time.

Notably, the maladaptive swapped-activation strategy observed early on was

abandoned during the late adaptation phase, with synergy activation patterns

reverting towards their original state (Fig. 6, 7). This reversion occurred despite

remaining mechanical antagonism, suggesting the CNS did not achieve perfect

fine-grained fractionation but instead adopted a different strategy, perhaps driven

by the high cost of the internal conflict (Fig. 8B, 'arms race'). This outcome aligns

with principles of "good enough" motor control (Mussa-Ivaldi and Bizzi, 1995;

Ranganathan et al., 2013; Gijsberts et al., 2014), where functional success and

potentially reduced effort are prioritized over perfect efficiency, especially when

faced with constraints. Indeed, functional recovery was largely achieved (Fig. 4A)

and maladaptive behaviors reduced (Fig. 4B) during this late phase. Crucially,

this reversion to the less conflicted original timing was likely enabled only by the

concurrent development of slower compensatory strategies, like the tenodesis

effect (Fig. 9, discussed below), which provided an alternative means to achieve

task goals.

The rapid nature of both the initial synergy swap and the later switch-back—

typically stabilizing within days—suggests distinct fast adaptive processes,

potentially triggered by strong error signals. The initial maladaptation could

plausibly be linked to the 'sensory surprise' from altered proprioceptive feedback

(Kitazawa et al., 1995; Martin et al., 1996; Smith et al., 2006; Edwards et al.,

2012; Petitet et al., 2018) driving a rapid but flawed response. While lingering

discomfort cannot be fully excluded after the initial recovery period, the specific,

evolving nature of the synergy changes over weeks seems primarily attributable

to the fundamental sensorimotor disruption itself rather than ongoing pain.

Contextualizing these findings requires considering different forms of plasticity.

For instance, while operant conditioning demonstrates that even spinal reflexes

can be substantially modified over weeks, primarily through gain modulation

within specific pathways (Thompson and Wolpaw, 2014), adapting to the tendon

transfer studied here presented a distinct challenge. The drastically altered

sensorimotor mapping and mechanics likely necessitated the broader, multi-level

network reorganization reflected in our complex multi-timescale findings, rather

than primarily tuning an existing response.

It is important to note that while our primary analysis focused on muscle synergies

during grasping epochs, the tendon transfer also induced significant deficits in

the preceding reaching movements (Supplementary Videos S2, S3, S6; Figs. 4B,

D), highlighting that adaptation encompassed the entire reach-to-grasp sequence.

Although the reach component was not quantitatively analyzed here, its recovery

likely interacted strongly with the adaptation of grasp control, given the need to

coordinate the full limb trajectory under new mechanical constraints. Furthermore,

the development of compensatory strategies, such as the tenodesis effect aiding

hand pre-shaping, likely influenced both reach and grasp components.

Slow and gradual adaptation of several non-relocated muscles

The initial maladaptive phase, characterized by swapped synergy activation

patterns, coincided with significant functional deficits, including malcoordination

and increased task times (Fig. 4; Supplementary Videos S2, S3, S6). In contrast,

slower adaptive processes emerged over weeks to months in non-transferred

muscles (e.g., FDP, ECR, PL, and FCR in Monkey A, Fig. S4), suggesting

compensatory mechanisms.

A key compensation was the gradual development of wrist flexion to substitute

for reduced finger extension. Specifically, the increased activation of Synergy C

(wrist flexors FCR and PL, Fig. 6C) during hand opening and pre-shaping (Figs.

S3Q, S and S8C; S3AL and S8E) indicates the monkeys' learned use of the

tenodesis effect. This biomechanical coupling, where wrist flexion generates

passive finger extension (Zajac, 1992; Cash and Jones, 2011), provided a viable

alternative for hand opening, ultimately enabling the 'switch-back' of Synergies A

and B and the recovery of task performance (Fig. 4).

Additionally, the non-transferred finger flexor FDP showed a slow, gradual

increase in activity, independent of its original synergy (Synergy D) (Fig. S4F, P;

Fig. S3M). This suggests complex, slow-timescale adjustments, potentially

involving altered synergistic relationships or novel independent muscle control.

Neural adaptation from different adaptive modules

A key question concerns the interaction between rapid and gradual adaptation

processes. Our model (Fig. 6) proposes that the large initial sensory prediction

error (SPE) from the tendon transfer triggered a fast adaptive process – the

observed synergy swap – which, despite being maladaptive, rapidly altered the

sensorimotor state. From this altered state, slower processes were subsequently

recruited, including both a continued maladaptive strategy (Process A, e.g., the

Synergy B 'arms race') and the development of compensatory movements

(Process B, e.g., tenodesis via Synergy C). The cumulative cost of the

maladaptive process likely triggered a second fast adaptation – the switch-back

to the original synergy timing. This final shift was enabled by the compensatory

strategy (Process B), resulting in a stable 'good enough' functional outcome

despite residual antagonism (Ranganathan et al., 2013). Further understanding

requires considering factors not assessed here, such as the roles of spinal reflex

pathways (Smith et al., 2006) and fusimotor drive (Vallbo et al., 1979; Hospod et

al., 2007).

Implication for sensorimotor rehabilitation

Our findings, revealing distinct phases including initial maladaptation followed by

slower compensation and functional recovery, hold potential implications for

rehabilitation. A staged approach mirroring these phases could be beneficial: an

early phase might focus on minimizing sensory prediction errors and reducing

maladaptive responses through techniques aimed at integrating the altered body

schema (e.g., using visual feedback or task-oriented practice with managed

error) (Nudo et al., 1996; Takeuchi and Izumi, 2012; Zeiler and Krakauer, 2013;

Foell et al., 2014; Levin et al., 2015; De Nunzio et al., 2018; Sattin et al., 2023;

van Vliet et al., 2023).

Subsequently, a later phase could employ task-specific training to promote

effective compensatory strategies, similar to the slow adaptations observed here.

Encouraging functional compensation might facilitate the eventual abandonment

of maladaptive patterns and the consolidation of a stable, functional motor

strategy, consistent with the eventual recovery observed (Murata et al., 2008).

METHODS

Animals

Data were collected using two male macaque monkeys (*Macaca fuscata*; Monkey
A: 7.8 kg and Monkey B: 9.9 kg) purpose-bred at the National Bioresource Project
(NBRP). They were kept in custom-made primate cages, allowing for potential
pair housing. Both monkeys were trained to perform a simple grasping task.
During experimental recordings, the monkeys were seated in a primate chair
without further restraint. Their head movement remained unrestricted at all times.

Procedures

All procedures were designed to minimize discomfort and pain and approved by
the institutional Animal Care and Use Committees at the National Center of
Neurology and Psychiatry (NCNP), Tokyo, Japan. Details of the surgical
operations, experimental setup, and procedures for electromyography (EMG)
recordings have been previously described (Takei and Seki, 2010).

EMG implant surgery

Both animals were familiarized with the experimental setup and trained to perform the behavioral task prior to surgery. After an initial training period, EMG electrodes were chronically implanted subcutaneously into several muscles of the left forearm including the FDS, FDP, and EDC (Figs. 1, 3C, and S1C). Individual muscles were localized and confirmed using electrical microstimulation before implanting EMG electrodes chronically by two methods. For muscles involved in the crossed tendon transfer procedure, muscle fascia was cut and EMG wires with looped ends were placed on top of the muscle belly. These were then secured by reclosing the fascia with absorbable suture threaded through the loop (AS-361, stainless-steel Cooner wire, Conner Wire Co., Chatsworth, CA, USA). All other muscles were directly implanted into the muscle belly using a 22-gauge hypodermic needle. Each wire was threaded into the needle tip, folded back along the shaft, and inserted into the muscle before carefully retracting the needle. Electrode separation for bipolar recordings was approximately 5–10 mm. Note that some EMG recording sites were lost over time. Surgical procedures were carried out under deep general anesthesia (sevoflurane 1.5 to 2.5% in 2:1 O2/N2O) and with full aseptic precautions. Heart rate, blood pressure, body

temperature, and blood oxygen saturation were monitored throughout surgery.

Analgesics and antibiotics were administered intramuscularly for at least one

week postoperatively.

Tendon-Transfer Surgery

After a minimum four-week recovery period from EMG implant surgery, TT

surgery was performed. The FDS and EDC tendons were cut as distally as

possible (immediately below and above the wrist for FDS and EDC, respectively)

to avoid damaging the Golgi tendon organ located near the junction between the

muscle fibers and tendon (Schoultz and Swett, 1972; Jami, 1992). Tendons were

then guided either through the gap between the radius and ulnar bone (Monkey

A), or around the wrist and then re-attached to the tendon of the antagonist

muscle using a tendon graft harvested from the plantaris tendon of the lower limb

(Monkey B; EDC → FDS: direct connection; and FDS → EDC: tendon grafting).

This was only necessary in Monkey B since the tendons used for the cross-

transfer were too short. This surgical procedure aimed to reverse the primary

mechanical actions of the manipulated muscles, making the FDS tendon an

effective finger extensor and the EDC tendon an effective finger flexor. The

success and nature of this mechanical rearrangement were verified post-

operatively via direct muscle stimulation (see Methods subsection 'Tendon

transfer confirmation' and Fig. 2). The general organization and function of the

major forearm muscles involved in finger flexion and extension in *Macaca fuscata*

are broadly comparable to those in humans (Vanhoof et al., 2021; Yan et al.,

2022).

Surgical procedures were carried out under deep general anesthesia

(sevoflurane 1.5 to 2.5% in 2:1 O_2/N_2O) and with full aseptic precautions. Heart

rate, blood pressure, body temperature, and blood oxygen saturation were

monitored throughout surgery. Analgesics and antibiotics were administered

intramuscularly for at least one week postoperatively. Both monkeys wore a

plaster cast post-surgery, but its effectiveness was limited (lasting approximately

one week).

Tendon transfer confirmation

To verify the success and long-term stability of TT surgery, two procedures (Fig.

2) were implemented. During the first procedure, the monkey was sedated and

its arm was secured to a metal frame (Fig.2A). Electrical stimulation was then

applied to either the FDS or EDC muscle (50 mA, DS8R, Digitimer, Welwyn

Garden City, UK), while ultrasound scans of tendon movements were

concurrently captured (SONIMAGE MX1, Konica Minolta, Inc., Tokyo, Japan

(Nordez et al., 2009; Dieterich et al., 2017).

Figure 2B shows a sonogram of the FDS muscle and its intramuscular tendons.

The left side displays a static image of the FDS muscle at a specific moment, with

white arrows indicating the FDS tendon used for measurement. The right side

shows staggered images (top white box) of the FDS tendon displacement

triggered by muscle stimulation. The lower right inset illustrates the measurement

area. The displacement wave area represents the intensity of muscle contraction,

and was calculated by measuring the average duration (a, in seconds) and

amplitude (b, in cm) of three successive waves. The area (a*b/2) and regression

line for FDS (red) and EDC (blue) (R^2 > 0.5 for FDS) for days 0, 7, and 105 post-

TT are shown (Fig. 2C). The results suggest that the muscle contractions induced

by direct electrical stimulation remained nearly constant.

For the second procedure, high-speed videos (37U Series Color Industrial

Camera, The Imaging Source, Charlotte, NC, USA) of the monkey's finger

movements were recorded while electrically stimulating the FDS or EDC. Markers

were placed on the nails of the index, middle, and ring fingers (Fig. 2A) to

measure finger displacement in xyz-dimensions using DeepLabCut software

(Mathis et al., 2018; Nath et al., 2019). The sum of Euclidean distances of each

marker from the origin of the three-dimensional (3D) coordinate system was

computed as a scalar quantity. Post-surgery, movement along the z-axis

reversed, indicating a shift from finger flexion to extension due to TT (Fig. 2D;

blue: pre-TT on surgery day, dark brown: post-TT on surgery day, light brown: 1

wk post-TT, and red: 3 wk post-TT). The scalar quantities of each finger did not

significantly change during muscle stimulation on days 0-, 7-, and 105-days post-

TT (Fig. 2E), suggesting there was no postoperative tendon rupture or slackening.

Data recordings

EMG signals were recorded using the AlphaLab SnR system (Alpha Omega

Engineering, Hamerkava St. 6, Ziporit Industrial Zone, P.O.Box 810, Nof HaGalil

(Nazareth Illit) 1789062, Israel), displayed online and then stored on a hard drive

for later off-line analysis using MATLAB (MathWorks, Natick, MA, USA). Data

were recorded at 11 kHz and processed for analysis as follows. 1. Down-sampled

to 5 kHz; 2. 50 Hz high-pass filtered (6th order Butterworth filter); 3. rectified; 4.

20 Hz low-pass filtered (6th order Butterworth filter); and 5. down-sampled to 100

Hz. Behavioral task events were recorded as transistor–transistor logic (TTL)

signals using the AlphaLab SnR system. Monkey behavior was further recorded

by two cameras (Sanyo VPC-WH1, 60 fps [Sanyo, Osaka, Japan]; and Casio EX-

100F, 240 fps [Casio, Tokyo, Japan] for monkeys A and B, respectively) from two

different angles (top and side view). The images were later used to detect

additional behavioral events (contact time with the object and pull onset for

Monkey A; and food touch and food lift onset for Monkey B), and detection of

possible changes in the animal's general movement pattern. Kinovea software

was used for this video analysis (free and open-source software;

https://www.kinovea.org/).

 Behavioral task

Before and after EMG surgery, the monkeys were trained on a simple grasping

task that involved a small object attached to a rod (Fig. 3B for Monkey A; and Fig.

S1B for Monkey B). For Monkey A, object 1 was a small rod placed between two

side walls encouraging the monkey to grasp the rod with a precision grip using

the tips of the index finger and thumb (controlled grasp). The force used to

compress the spring while pulling was low and the distance moved short. Object

2 had to be grasped in the same way. However, the force to compress the spring

was higher and the distance moved longer (object 1: 300 cN, 4 mm; and object

2: 800 cN, 30 mm). The monkey was asked to grasp and hold the objects for

approximately 300–500 ms (Fig. 3A–D). After completing the trial, the monkey

was rewarded with a piece of fruit that had to be taken from the experimenter's

palm. The task sequence for Monkey A ranged from object 1 hold onset to object

2 hold offset (Fig. 3C, D; and 'Data Analysis' below). For Monkey B, object 1 was

a small rod that was grasped and pulled (300 cN, 4 mm) using a lateral

prehension grip (like using a door key), and resembling a power grip rather than

a natural grasp. However, instead of grasping object 2, Monkey B picked up a

piece of fruit from an allocated location. In doing so, the monkey had to cross the

path of a photo cell which was detected by the recording system (Fig. S1A–D).

The task sequence to be analyzed for Monkey B ranged from object 1 hold onset

to food touch, which was indicated by a LED (Fig. S1C, D; and 'Data Analysis'

below).

For Monkey A, the task sequence was as follows (Fig. 3A; and Supplementary

Video S1). From a starting position in front of the monkey, it had to lift its arm and

move towards object 1, grasp, pull, and then hold the object for 500 ms.

Immediately after releasing object 1, Monkey A proceeded to move its hand

towards object 2, which was located to the right of object 1. Again, the monkey

had to grasp, pull, and then hold the object for another 500 ms. Each hold period

was accompanied by an audio signal which stopped once the hold duration was

sufficient. After completing the task, which was signaled by another acoustic

signal, the monkey was rewarded with a piece of fruit presented by the

experimenter and taken by the monkey.

For Monkey B, the first part of the sequence was identical. However, after

releasing object 1, Monkey B was required to pick up a piece of fruit from an

allocated location (Fig. S1A, B; and Supplementary Video S5). In doing so, the

monkey passed a photocell in front of the food well. This event was detected by

the recording system and a video camera (a red LED triggered by activation of

the photocell). For the analysis, only object 1 and the food grasp were considered

in monkeys A and B, respectively.

Data analysis

<u>*EMG analysis*</u>

EMG data were normalized to the average time for each monkey to complete a

trial (object 1 hold onset ➔ object 2 hold offset/LED offset = 100%). Data were

cropped and aligned according to the time stamps (Obj1 hold offset ± 15%; and

LED onset [food touch] ± 15% for monkeys A and B, respectively). Recorded

EMGs from the pre- and post-tendon surgery period were averaged for each

recording and compared over experimental sessions. For both monkeys, the

muscles used for EMG analysis were: FDS and EDC for pre- (control) and EDC

for post-TT (experimental).

Both monkeys required time to recover from surgery. Once they were able to

perform the task on their own, data collection started (29 and 20 days post-

surgery for monkeys A and B, respectively). After tendon surgery, EMG signals

for FDS, FCU, and FCR deteriorated in Monkey B. Therefore, there is no

experimental data for these muscles in this monkey and they were excluded from

the muscle synergy analysis.

Synergy analysis

The EMG envelopes obtained from pre-processing raw EMG data were divided

by the mean value to normalize activity. Muscle synergies were then extracted

for each session using NMF (d'Avella et al., 2003). NMF decomposes the EMG

data matrix M, as a product of two matrices C and W:

$M = CW^T + re = \Sigma c_i w_i^T + re$

where the vector wi and column of matrix W are muscle synergies; the vector ci,

and column of matrix C are their temporal coefficients; and re represents the

residuals. Multiplicative update rules were used for decomposition (Lee and

Seung, 1999). Updating matrices for decomposition from 20 different random

initial matrices was started, and factorization results with least residuals among

the 20 results were used. To improve robustness of the muscle synergies, one

set of muscle synergies was extracted from EMG data of multiple trials of each

day. A k-fold cross validation was used, with k = 4 i.e., EMG data set of multiple

trials during the day was split into four data sets, in which three data sets were

for training and one data set was for testing. Synergy matrix W was calculated

from three training data sets using NMF, and then a coefficient matrix C was

calculated by decomposing the EMG matrix M of the test data set with a fixed W

matrix of the training data set. This operation was repeated by changing the test

data set for each four data sets. Finally, the extracted four C matrices and four W

matrices were averaged to obtain daily synergies.

The number of synergies was determined as the number where VAF exceeded

a threshold. This threshold was set as 0.8. To clarify the effect of TT, the number

of synergies for each monkey was determined, assuming that trial-by-trial

variation in synergies within the same monkey was small. MATLAB 'nnmf'

function was used for NMF.

To determine whether the weights of muscle synergies changed after TT, the

following procedure was used. First, the average of each pre-TT spatial synergy

was calculated and used as control data to calculate the cosine distance for all

post-TT spatial synergies. This generated four distance relationships for each

pre-TT synergy. If the change in spatial synergy before and after TT was small,

then only one of the four cosine distances (e.g., synergy A before TT and synergy

A after TT) should be significantly smaller. Two-way ANOVA was conducted

using the type of synergy and session as factors for the cosine distance. The

significance of synergy pairs was identified using the Bonferroni post-hoc test.

Cross-correlation analysis

Cross-correlation analysis was performed to examine pre- and post-surgery EMG

signals as well as temporal activation coefficients of extracted muscle synergies.

Cross-correlation coefficients were calculated using MATLAB (MathWorks,

corrcoef.m function). These were plotted for different muscles and muscle

synergy activation coefficients over experimental sessions for comparison.

Based on cross-correlation analysis, five 'landmark days' were identified for

further analysis (Figs. 2G, H, and S2D). These days represent distinct stages in

the recovery process and excluded pre-surgery control data. The first landmark

day was chosen from one of the initial recording sessions after TT surgery (days

29 and 22 in monkeys A and B, respectively) when the cross-correlation

coefficients had changed significantly compared with control data. The second

day was then chosen from a time period just before the switch-back, when the

cross-correlation coefficients had started to return to their original values (days

64 and 36). Another day was then picked from the period when the coefficients

were still changing significantly (days 69 and 44) and before starting to saturate.

At this point the next day was defined (days 79 and 48). Finally, the final landmark

day was chosen from one of the last recording sessions when the behavior had

fully recovered (days 99 and day 64 for monkeys A and B, respectively).

Behavioral analysis

To examine behavioral recovery, the duration, onset, and offset of object and

food grasps were analyzed. Event times extracted from the video analysis were

used for alignment of EMG data and subsequent cross-correlation and synergy

analyses, along with recorded event TTL signals. For each experimental session,

the first 20 trials of each video recording were analyzed. In Monkey A, the

following events were detected and in-between times stored in ms: touch onset,

pull onset, and hold onset (resulting in grasp duration and pull time) for object 1

and object 2. In Monkey B, the following events were detected and in-between

times stored: touch onset, pull onset, and hold onset (resulting in grasp duration

and pull time) for object 1; and for the food grasp component of the task, LED

onset, food-touch onset, food-lift onset, LED offset, and movement end (contact

of food with mouth).

Contact times (touch onset ➜ pull onset) for Monkey A were plotted in ms (mean

± SD) over number of days from tendon surgery for object 1. For Monkey B,

contact times with food (food-touch ➜ food-lift onset) were plotted in ms (mean

± SD).

ANOVA was performed to compare control data recorded before and after tendon

surgery. In total, 5 and 3 control sessions vs. 35 and 22 experimental sessions

were used for analyses in monkeys A and B, respectively.

To quantify the observed malcoordinated reaching behavior, video footage was

analyzed and the means of 10 consecutive trials calculated for each session as

follows. For Monkey A, time spent within a specific spatial window covering the

area behind and between the two objects was measured; the monkey passed

through this space almost exclusively while exhibiting the impairment (see Fig.

3E, crossing of the yellow dotted line). In Monkey B, the time spent in contact with

the rear plate of the object was measured (see Fig. S1G).

Joint kinematics analysis

To determine whether compensatory movements occurred, steady state changes

in joint angles were examined before and after recovery. Using DeepLabCut

(Mathis et al., 2018; Nath et al., 2019), key points on the fingers and wrists were

tracked from experimental videos. From the obtained key points, the vector v_1

connecting the wrist and fingers and vector v_2 connecting the arm and wrist were

estimated. The joint angle θ was calculated as follows:

920
$$cos(\theta) = \frac{v_2 \cdot v_3}{|v_2| \, |v_3|} \rightarrow \theta = cos^{-1}\left(\frac{v_2 \cdot v_3}{|v_2| \, |v_3|}\right)$$

923 **DATA, MATERIALS, AND SOFTWARE AVAILABILITY**

924

925 Matlab scripts and raw data will be made available online. All study data are

926 included in the article and/or SI Appendix.

927

ACKNOWLEDGEMENTS:

We would like to express our sincere gratitude to Drs. Francisco Valero-Cuevas

and Andrea d'Avella for their invaluable comments and constructive suggestions

on the earlier version of the manuscript, and Drs Nicholas Schweighofer and

Gerald E. Loeb for stimulating discussion. We would also like to thank Dr.

Naomichi Ogihara and Prof. Toshiyasu Nakamura for their valuable insights and

technical advice on tendon surgery in the hand and forearm of primates. We are

grateful to Drs. Kumiko Oida and Chika Sasaki for their medical care of the

experimental animals, vital help during surgery, and training of the animals. Lastly,

we would like to thank Masahi Koizumi for designing parts of the experimental

setup and Drs. Joachim Confais, Shinji Kubota, Saeka Tomatsu, Tatsuya Umeda,

and Natsumi Uchida for their support and assistance during the proceeding pilot

study. We thank Rachel James, PhD, from Edanz (https://jp.edanz.com/ac) for

editing a draft of this manuscript.

This work was supported by Grants-in-Aid for Scientific Research from the

Ministry of Education, Culture, Sports, Science, and Technology of Japan

(26120003, 26250013, 15K21754, 19H05724, 19H01092, 23H05488, and

24K21313), the Japan Agency for Medical Research and Development

REFERENCES

958 Berger DJ, Gentner R, Edmunds T, Pai DK, d'Avella A (2013) Differences in Adaptation Rates
959   after Virtual Surgeries Provide Direct Evidence for Modularity. The Journal of
960   Neuroscience 33:12384-12394.

961 Bizzi E, Cheung VC (2013) The neural origin of muscle synergies. Frontiers in Computational
962   Neuroscience Volume 7 - 2013.

963 Bowlus TH, Lane RD, Stojic AS, Johnston M, Pluto CP, Chan M, Chiaia NL, Rhoades RW
964   (2003) Comparison of reorganization of the somatosensory system in rats that sustained
965   forelimb removal as neonates and as adults. J Comp Neurol 465:335-348.

966 Cash DJW, Jones JWM (2011) The role of tenodesis in surgery of the upper limb. The Journal of
967   Bone & Joint Surgery British Volume 93-B:285-292.

968 Cheung VC, Turolla A, Agostini M, Silvoni S, Bennis C, Kasi P, Paganoni S, Bonato P, Bizzi E
969   (2012) Muscle synergy patterns as physiological markers of motor cortical damage. Proc
970   Natl Acad Sci U S A 109:14652-14656.

971 d'Avella A, Saltiel P, Bizzi E (2003) Combinations of muscle synergies in the construction of a
972   natural motor behavior. Nature Neuroscience 6:300-308.

973 Davidson PR, Wolpert DM (2003) Motor learning and prediction in a variable environment.
974   Curr Opin Neurobiol 13:232-237.

975 De Nunzio AM, Schweisfurth MA, Ge N, Falla D, Hahne J, Gödecke K, Petzke F, Siebertz M,
976   Dechent P, Weiss T, Flor H, Graimann B, Aszmann OC, Farina D (2018) Relieving
977   phantom limb pain with multimodal sensory-motor training. J Neural Eng 15:066022.

978 Dieterich AV, Botter A, Vieira TM, Peolsson A, Petzke F, Davey P, Falla D (2017) Spatial
979   variation and inconsistency between estimates of onset of muscle activation from EMG
980   and ultrasound. Sci Rep 7:42011.

981 Edwards MJ, Adams RA, Brown H, Pareés I, Friston KJ (2012) A Bayesian account of 'hysteria'.
982   Brain 135:3495-3512.

983 Fleury L, Prablanc C, Priot AE (2019) Do prism and other adaptation paradigms really measure
984   the same processes? Cortex 119:480-496.

985 Foell J, Bekrater-Bodmann R, Diers M, Flor H (2014) Mirror therapy for phantom limb pain:
986   brain changes and the role of body representation. Eur J Pain 18:729-739.

987 Gaetz W, Dockstader C, Furlong PL, Amaral S, Vossough A, Schwartz ES, Roberts TPL, Scott
988   Levin L (2023) Somatosensory and motor representations following bilateral transplants
989   of the hands: A 6-year longitudinal case report on the first pediatric bilateral hand
990   transplant patient. Brain Res 1804:148262.

Gijsberts A, Bohra R, Sierra González D, Werner A, Nowak M, Caputo B, Roa MA, Castellini C

(2014) Stable myoelectric control of a hand prosthesis using non-linear incremental

learning. Front Neurorobot 8:8.

Green HJ (1997) Mechanisms of muscle fatigue in intense exercise. J Sports Sci 15:247-256.

Hospod V, Aimonetti J-M, Roll J-P, Ribot-Ciscar E (2007) Changes in Human Muscle Spindle

Sensitivity during a Proprioceptive Attention Task. The Journal of Neuroscience

27:5172-5178.

Imamizu H, Miyauchi S, Tamada T, Sasaki Y, Takino R, Pütz B, Yoshioka T, Kawato M (2000)

Human cerebellar activity reflecting an acquired internal model of a new tool. Nature

403:192-195.

Jami L (1992) Golgi tendon organs in mammalian skeletal muscle: functional properties and

central actions. Physiol Rev 72:623-666.

Kitazawa S, Kohno T, Uka T (1995) Effects of delayed visual information on the rate and amount

of prism adaptation in the human. J Neurosci 15:7644-7652.

Lee DD, Seung HS (1999) Learning the parts of objects by non-negative matrix factorization.

Nature 401:788-791.

Levin MF, Weiss PL, Keshner EA (2015) Emergence of virtual reality as a tool for upper limb

rehabilitation: incorporation of motor control and motor learning principles. Phys Ther

95:415-425.

Loeb GE (1999) Asymmetry of hindlimb muscle activity and cutaneous reflexes after tendon

transfers in kittens. J Neurophysiol 82:3392-3405.

Luauté J, Schwartz S, Rossetti Y, Spiridon M, Rode G, Boisson D, Vuilleumier P (2009)

Dynamic changes in brain activity during prism adaptation. J Neurosci 29:169-178.

Madarshahian S, Latash ML (2022) Reciprocal and coactivation commands at the level of

individual motor units in an extrinsic finger flexor-extensor muscle pair. Exp Brain Res

240:321-340.

Martin TA, Keating JG, Goodkin HP, Bastian AJ, Thach WT (1996) Throwing while looking

through prisms. II. Specificity and storage of multiple gaze-throw calibrations. Brain 119

(Pt 4):1199-1211.

Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M (2018)

DeepLabCut: markerless pose estimation of user-defined body parts with deep learning.

Nature Neuroscience 21:1281-1289.

Murata Y, Higo N, Oishi T, Yamashita A, Matsuda K, Hayashi M, Yamane S (2008) Effects of

motor training on the recovery of manual dexterity after primary motor cortex lesion in

macaque monkeys. J Neurophysiol 99:773-786.

Mussa-Ivaldi FA, Bizzi E (1995) The Modular Organization of Motor Control: What Frogs Can

Teach Us About Adaptive Learning. IFAC Proceedings Volumes 28:413-418.

Nath T, Mathis A, Chen AC, Patel A, Bethge M, Mathis MW (2019) Using DeepLabCut for 3D
1029   markerless pose estimation across species and behaviors. Nature Protocols 14:2152-
1030   2176.

1031   Nordez A, Gallot T, Catheline S, Guével A, Cornu C, Hug F (2009) Electromechanical delay
1032   revisited using very high frame rate ultrasound. J Appl Physiol (1985) 106:1970-1975.

1033   Nudo RJ, Wise BM, SiFuentes F, Milliken GW (1996) Neural substrates for the effects of
1034   rehabilitative training on motor recovery after ischemic infarct. Science 272:1791-1794.

1035   Overduin SA, d'Avella A, Roh J, Bizzi E (2008) Modulation of Muscle Synergy Recruitment in
1036   Primate Grasping. The Journal of Neuroscience 28:880-892.

1037   Petitet P, O'Reilly JX, O'Shea J (2018) Towards a neuro-computational account of prism
1038   adaptation. Neuropsychologia 115:188-203.

1039   Power JD, Schlaggar BL (2017) Neural plasticity across the lifespan. Wiley Interdiscip Rev Dev
1040   Biol 6.

1041   Ranganathan R, Adewuyi A, Mussa-Ivaldi FA (2013) Learning to be lazy: exploiting redundancy
1042   in a novel task to minimize movement-related effort. J Neurosci 33:2754-2760.

1043   Sattin D, Parma C, Lunetta C, Zulueta A, Lanzone J, Giani L, Vassallo M, Picozzi M, Parati EA
1044   (2023) An Overview of the Body Schema and Body Image: Theoretical Models,
1045   Methodological Settings and Pitfalls for Rehabilitation of Persons with Neurological
1046   Disorders. Brain Sci 13.

1047   Schärli A, Hecht H, Mast FW, Hossner EJ (2024) How spotting technique affects dizziness and
1048   postural stability after full-body rotations in dancers. Hum Mov Sci 95:103211.

1049   Schoultz TW, Swett JE (1972) The fine structure of the Golgi tendon organ. J Neurocytol 1:1-26.

1050   Slawinska U, Kasicki S (2002) Altered Electromyographic Activity Pattern of Rat Soleus Muscle
1051   Transposed into the Bed of Antagonist Muscle. The Journal of Neuroscience 22:5808-
1052   5812.

1053   Smith MA, Ghazizadeh A, Shadmehr R (2006) Interacting adaptive processes with different
1054   timescales underlie short-term motor learning. PLoS Biol 4:e179.

1055   Sperry RW (1940) The functional results of muscle transposition in the hind limb of the rat.
1056   Journal of Comparative Neurology 73:379-404.

1057   Sperry RW (1942) Transplantation of motor nerves and muscles in the forelimb of the rat.
1058   Journal of Comparative Neurology 76.

1059   Sugita Y (1996) Global plasticity in adult visual cortex following reversal of visual input. Nature
1060   380:523-526.

1061   Takei T, Seki K (2010) Spinal interneurons facilitate coactivation of hand muscles during a
1062   precision grip task in monkeys. J Neurosci 30:17041-17050.

Takei T, Confais J, Tomatsu S, Oya T, Seki K (2017) Neural basis for hand muscle synergies in

the primate spinal cord. Proceedings of the National Academy of Sciences 114:8643-

8648.

Takeuchi N, Izumi S (2012) Maladaptive plasticity for motor recovery after stroke: mechanisms

and approaches. Neural Plast 2012:359728.

Thompson AK, Wolpaw JR (2014) Operant conditioning of spinal reflexes: from basic science to

clinical therapy. Front Integr Neurosci 8:25.

Vallbo AB, Hagbarth KE, Torebjörk HE, Wallin BG (1979) Somatosensory, proprioceptive, and

sympathetic activity in human peripheral nerves. Physiol Rev 59:919-957.

van Vliet P, Carey LM, Turton A, Kwakkel G, Palazzi K, Oldmeadow C, Searles A, Lavis H,

Middleton S, Galloway M, Dimech-Betancourt B, O'Keefe S, Tavener M (2023) Task-

specific training versus usual care to improve upper limb function after stroke: the

"Task-AT Home" randomised controlled trial protocol. Frontiers in Neurology 14.

Vanhoof MJM, van Leeuwen T, Galletta L, Vereecke EE (2021) The forearm and hand

musculature of semi-terrestrial rhesus macaques (Macaca mulatta) and arboreal gibbons

(fam.Hylobatidae). Part II. Quantitative analysis. Journal of Anatomy 238:321-337.

Walker JG, Jackson HJ, Littlejohn GO (2004) Models of adjustment to chronic illness: using the

example of rheumatoid arthritis. Clin Psychol Rev 24:461-488.

Wester K, Hove LM, Barndon R, Craven AR, Hugdahl K (2018) Cortical Plasticity After Surgical

Tendon Transfer in Tetraplegics. Front Hum Neurosci 12:234.

Yan Y, Sobinov AR, Bensmaia SJ (2022) Prehension kinematics in humans and macaques. J

Neurophysiol 127:1669-1678.

Yumiya H, Larsen KD, Asanuma H (1979) Motor readjustment and input-output relationship of

motor cortex following cross-connection of forearm muscles in cats. Brain Research

177:566-570.

Zajac FE (1992) How musculotendon architecture and joint geometry affect the capacity of

muscles to move and exert force on objects: a review with application to arm and forearm

tendon transfer design. J Hand Surg Am 17:799-804.

Zeiler SR, Krakauer JW (2013) The interaction between training and plasticity in the poststroke

brain. Curr Opin Neurol 26:609-616.

Zemková E (2022) Physiological Mechanisms of Exercise and Its Effects on Postural Sway: Does

Sport Make a Difference? Front Physiol 13:792875.

1095

1096

1097

Fig.1: Muscle anatomy of the forearm of the macaque monkey and surgical procedure.

All muscles implanted with electromyography electrodes are labeled. Muscle abbreviations (Monkey A): brachioradialis (BRD), extensor carpi radialis (ECR), extensor carpi ulnaris (ECU), extensor digitorum-2,3 (ED23), extensor digitorum communis (EDC), flexor carpi radialis (FCR), flexor carpi ulnaris (FCU), flexor digitorum profundus (FDP), flexor digitorum superficialis (FDS), and palmaris longus (PL). Additionally implanted muscles in Monkey B: deltoid (DEL) and extensor digitorum-4,5 (ED45). Tendons of EDC and FDS muscles were cross-transferred.

Fig. 2. *Long term confirmation of tendon surgery effectiveness to alter mechanical properties.*

(A) Set-up for the ultrasound measurement and video recordings of the stimulation induced movements of the EDC and FDS tendons. (B) Sonogram of the FDS muscle and its intramuscular tendons. Left side (B-mode, i.e., brightness mode) shows the still image of the monkeys forearm at a given point in time. Right side (M-mode, i.e., motion mode) shows the stagged images of the FDS tendon displacement induced by muscle stimulation (50mA). White arrows demarcate the FDS tendon which was used for the measurement. Grayscale gradations correspond to tissue densities: hyperechoic regions (white) denote denser structures like surface of bones and tendons, while hypoechoic areas (black) signify less dense tissues such as adipose tissue and musculature. Inset demonstrates the area measurement. The area of the displacement waves was measured in the M-mode, representing the strength of muscle contraction. We measured the duration (a, sec) and amplitude (b, cm) of three waves and calculated the average. Area = a*b/2(cm/sec) for days 0, 7 and 105 after tendon transfer. (C) Areas under the wave measured in the M-mode for 3 experimental days (0, 7 and 105 days post-TT) and regression lines in red and blue for FDS

and EDC, respectively. R2>0.5 for FDS. The data suggested that muscle contractions induced by direct electrical stimulation were nearly constant. (D) Markers placed on the index, middle and ring finger nails (A) were used to measure finger displacement in xyz-dimensions. We calculated the sum of the Euclidean distances of each marker from the origin of the 3D coordinate system as a scalar quantity. Observing the movement along the Z-axis, it became reversed post-surgery indicating a reversal from finger flexion to extension due to tendon transfer (D, blue = pre-TT at surgery day; dark brown = post-TT at surgery day; light brown = 1wk post-TT; red = 3wks post-TT). The scalar quantity of the fingers during muscle stimulation did not change much at day 0, 7 and 105 days (E), suggesting that there was no tendon rupture or slackening of the tendons postoperatively (EDC stimulation, left; FDS stimulation, right).

1145

Fig. 3. *Experimental set-up, task sequence and typical EMG (monkey A).*

(A) Schematic of the behavioral sequence (hook → grasp → release). (B) Schematic of the task object using a rod requiring monkey A to perform a controlled grasp. (C) Typical EMG traces of a control session (high-pass filtered) for all recorded muscles. Gray boxes represent the task sequence. Obj 1 ON: start of the hold period of object 1. Obj 1 OFF: end of the hold period of object 1, i.e., object release. Obj 2 ON: start of the hold period of object 2. Obj 2 OFF: end of the hold period of object 2, i.e., object release. Muscle abbreviations: Brachioradialis (BRD), Extensor Carpi Radialis (ECR), Extensor Carpi Ulnaris

(ECU), Extensor Digitorum-2,3 (ED23), Extensor Digitorum Communis (*EDC), Flexor Carpi Radialis (FCR), Flexor Carpi Ulnaris (FCU), Flexor Digitorum Profundus (FDP), Flexor Digitorum Superficialis (*FDS), Palmaris Longus (PL). Tendons of the muscles marked with * were cross-transferred. (D) Rectified and smoothed EMG for all recorded muscles (average for one recording session; amplitude [μV] over task sequence [%]). Horizontal bars illustrate the corresponding behavioral periods, red vertical lines indicate peak amplitude for each muscle). (E) The time the monkey spent on the left side of the yellow dotted line while moving from object 1 to object 2 was measured and used to quantify the mal-adaptive behavior.

1167

Post surgery days

Fig. 4. *Behavioral analysis and mal-adaptation (monkey A)*

Monkey A: (A) Contact times [ms] with the object for the controlled grasp. (B)
Quantification of mal-coordinated behavior. Plotted is the duration [ms] the
monkey spent on the left side of the yellow dotted line while moving from object
1 to object 2 (see Fig. 3 E). Monkey B: (C) Contact times with food [ms] during
the natural grasp. Each data point represents the average of 20 observations
(±SD envelope) plotted in milliseconds over days with respect to TT (day 0). Open
squares: n.s., filled squares: $p<0.05$, two-sample t-test. (D) Contact times [ms]
with the object plate used to quantify the mal-coordinated reaching behavior.
Each data point represents the average of 10 observations (±SD envelope)
plotted over days with respect to TT (day 0). Open circles: n.s., filled circles:
$p<0.05$, two-sample t-test.

1180

1181

Fig. 5: Task sequence, typical electromyography, *EMG activity profiles of 'Landmark days' and Cross-correlation coefficients of transferred muscles for all recording sessions (monkey A).*

(A) Schematic of the behavioral sequence (hook → grasp → release). (B) Typical electromyography (EMG) traces of a control session (high-pass filtered) for all recorded muscles. Gray boxes represent the task sequence. Obj 1 ON: start of hold period of object 1; Obj 1 OFF: end of hold period of object 1 (i.e., object release); Obj 2 ON: start of hold period of object 2; Obj 2 OFF: end of hold period

of object 2 (i.e., object release). Tendons of muscles marked * were cross-transferred. (C, D) Electromyography (EMG) activity profiles (µV) for selected 'landmark days' pre- and post-tendon transfer (TT) of extensor digitorum communis (EDC) (C and D, respectively) in Monkey A. Profiles are aligned on hold off-set (object 1) at 0% task range (vertical dashed lines). Horizontal gray bars indicate three behavioral periods: hook, grasp, and release. (E, F) As above but for flexor digitorum superficialis (FDS). Filled triangles in (C) and (F) indicate maximum EMG amplitudes for EDC before TT and FDS (during extension) immediately after TT. Open triangles indicate maximum EMG amplitudes of FDS before and EDC immediately after TT (E and D, respectively). (G–J) Cross-correlation coefficients plotted over experimental days with respect to tendon surgery. G: EDC pre/post; H: FDS pre/post; I: FDS pre/EDC post; and J: EDC pre/FDS post. Dashed lines indicate recovery periods with no recordings. Small triangles (G–J) indicate landmark days.

1208

Fig. 6: Synergy weights & Temporal activation coefficients of monkey A.

(A-D) Synergy weights for each extracted muscle synergy. Plotted are the synergy weights for each muscle for one control day before (left column) and 5 landmark days after TT surgery (right column). (E) 'Variance Accounted For' (VAF) for the original and shuffled data. Four muscle synergies that accounted for more than 80% of the variance were extracted from the data. Black horizontal lines demarcate the 80% threshold. (F-I) Plotted are the time varying activation coefficients of the muscle synergies over task range [%] before (left column) and after TT (right column; shown are the activation coefficients for 5 landmark days). Horizontal gray bars mark the three behavioral periods (hook, grasp and release, respectively). Filled triangles in (F) and (G) indicate the maximum amplitude for synergy A (finger flexor synergy) just after TT (finger extension) and synergy B (finger extensor synergy) before TT (finger extension), while open triangles indicate the maximum amplitude for synergy A before (finger flexion) and synergy B just after TT (during flexion). Small triangles indicate the landmark days. Profiles are aligned on "object 1 hold off" at zero percent task range [%].

1225
1226

Fig. 7: Cross-correlation of muscle synergies.
Cross-correlation coefficients were plotted over experimental days with respect to tendon surgery (Monkey A). The // represent the recovery period in which no recordings were taken. Synergies were cross-correlated with either synergy A (finger flexor synergy: C, D, G, H in red) or synergy B (finger extensor synergy, A, B, E, F in blue).

1235

Fig. 8: Aggregated and averaged electromyography.
Aggregated and averaged electromyography (EMG) activities of the main muscles contributing to each of the four synergies for each recording day (Monkey A). EMG activity between -15% and +15% task range was summed for the contributing muscles and then averaged for each recording session. Left column: (A) synergy A, contributing muscles: flexor digitorum superficialis (FDS) and flexor carpi ulnaris (FCU); (B) synergy B, contributing muscles: extensor

digitorum communis (EDC), extensor digitorum-2,3 (ED23), and extensor carpi ulnaris (ECU); (C) synergy C, contributing muscles: palmaris longus (PL) and flexor carpi radialis (FCR); and (D) synergy D, contributing muscles: flexor digitorum profundus (FDP), brachioradialis (BRD), and extensor carpi radialis (ECR). Vertical bars represent landmark days (red numbers at the top show actual recording days with respect to tendon transfer surgery). Right column: mean and standard error of the mean for landmark days. *$p < 0.01$, **$p < 0.001$, ***$p < 0.0001$ after Bonferroni correction ($\alpha = 0.01$).

1253

Fig. 9. *Changes in Monkey B's Joint Angles Following Tendon Transfer*
(TT).

Shown are the changes in the angle of two key joints in monkey B before and
after TT surgery (mean of 20 trials ±SD for each landmark day, taken 83ms before
food touch). The angle at the metacarpophalangeal (MCP) joint (A), responsible
for finger extension, steadily increased following surgery (indicating increased
finger extension). Concurrently, the wrist joint angle (B) decreased, indicating
wrist flexion. Statistical analysis confirmed these changes were significant
(ANOVA, α = 0.01, p < 0.0001). This coordinated movement pattern suggests a
potential compensatory strategy employed by the monkey to achieve finger
extension during the grasping tasks. // represents the monkeys' recovery period
in which no recordings were taken.

1266

1267

Fig. 10: Sensorimotor adaptation model following tendon transfer.

1268
Schematic depicting the hypothesized interaction between rapid and gradual
adaptation processes in the central nervous system following a significant
sensory prediction error (SPE) induced by tendon transfer (TT). Our model
incorporates multiple TT adaptation processes with varying timescales,
encompassing both fast and slow dynamics. Tendon transfer surgery (black
arrow) necessitates adjustments in finger muscle coordination, particularly a shift
in the timing of activity between flexor and extensor synergies (left blue arrow).
These changes exceed a critical threshold (blue dotted line), triggering slow
dynamic adaptations (A and B, blue and red arrow 1). Adaptation A is faster and
aims to maintain the results of the initial fast adaptation (potentially maladaptive).
Meanwhile, a slower process B (background) gradually builds a new behavioral
component for "good enough" task performance. This new component only
becomes apparent after a second fast adaptation (second blue arrow) is triggered
by process A reaching another threshold (2). At this point, the cumulative costs
of the maladaptive behavior become unacceptable. The gray line represents the
observed changes of cross-correlation coefficients (electromyography and
temporal activation coefficients of muscle synergies).

1286