1 **Multi-timescale neural adaptation underlying long-term**
2 **musculoskeletal reorganization.**
3 **(**Multi-timescale neural adaptation**)**
4

5 *Roland Philipp[1,2], Yuki Hara[3], Naohito Ohta[1,2], Naoki Uchida[1,2], Tomomichi
6 Oya[1,4], Tetsuro Funato[2], and *Kazuhiko Seki[1]
7
[1]National Center of Neurology and Psychiatry, Department of Neurophysiology,
4-1-1 Ogawa-Higashi, Kodaira, Tokyo 1878502, Japan
[2]University of ElectroCommunications, Graduate School of Informatics and
Engineering, Dept. of Mechanical and Intelligent Systems Engineering. W2-217,
1-5-1 Chofugaoka, Chofu, Tokyo, Japan
[3]National Center of Neurology and Psychiatry, Department of Orthopaedic
Surgery, 4-1-1 Ogawa-Higashi, Kodaira, Tokyo 1878502, Japan
[4]Western Institute for Neuroscience, University of Western Ontario, London,
N6A 3K7, Canada
17
18 *Corresponding authors:
19 Kazuhiko Seki and Roland Philipp
20 Email: seki@ncnp.go.jp and roland@ncnp.go.jp

21
22 Author Contributions: R.P. and K.S. designed all experiments and wrote the
23 paper. R.P., Y.H., N.U., and N.O. performed the experiments and analyzed the
24 data. R.P., Y.H., T.O., and K.S. performed the surgeries. T.F. provided
25 conceptual advice and wrote the paper. All authors reviewed the results and
26 approved the final version of the manuscript.
27
28 Conflict of Interest Statement: The authors declare no competing interests.
29

Number of pages: 83 (92, including supplementary information)
Number of figures: 14 (24, including supplementary figures)
Number of multimedia: 7
Number of words for abstract: 173
Number of words in introduction: 740
Number of words in discussion: 2309

ABSTRACT

The central nervous system (CNS) can effectively control body movements

despite environmental changes. While much is known about adaptation to

external environmental changes, less is known about responses to internal bodily

changes. This study investigates how the CNS adapts to long-term alterations in

the musculoskeletal system using a tendon transfer model in non-human

primates. We surgically relocated finger flexor and extensor muscles to examine

how the CNS adapts its strategy for finger movement control by measuring

muscle activities during grasping tasks. Two months post-surgery, the monkeys

demonstrated significant recovery of grasping function despite the initial

disruption. Our findings suggest a two-phase CNS adaptation process: an initial

phase enabling function with the transferred muscles, followed by a later phase

abandoning this enabled function and restoring a control strategy that, while

potentially less conflicted than the maladaptive state, resembled the original

pattern, possibly representing a 'good enough' solution. These results highlight a

multi-phase CNS adaptation process with distinct time constants in response to

sudden bodily changes, offering potential insights into understanding and treating

movement disorders.

SIGNIFICANCE STATEMENT

After major changes to the body's mechanics, the nervous system adapts using strategies on multiple timescales. Our primate tendon transfer study shows that core muscle synergy groupings remain stable, reflecting a default to modular control. However, the activation of these synergies changes dramatically; an initial, rapid 'swap' of their timing proves to be **maladaptive**, impairing motor function. This conflict is only resolved through the gradual development of **slower, compensatory strategies** over several weeks. This process highlights the fundamental tension the CNS faces when its reliance on stable motor modules conflicts with the need for flexible control, offering insights into neural plasticity and staged rehabilitation.

Introduction

The central nervous system (CNS) continuously adapts bodily functions in response to both external and internal challenges. Experimental models based on external perturbations, such as altered gravitational fields or distorted sensory feedback, have illuminated mechanisms of sensorimotor adaptation (Sugita, 1996; Davidson and Wolpert, 2003; Luauté et al., 2009; Fleury et al., 2019). Because the changes in the external environment can be controlled by the experimental design, these models provide an opportunity to assess how the CNS adapts to them. However, the transient and predictable nature of these changes may not fully capture the demands posed by internal, long-lasting, and unpredictable alterations to the body's internal environment

In contrast, the internal changes such as developmental growth (Power and Schlaggar, 2017), fatigue (Green, 1997), postural sway (Zemková, 2022), and sensory disorientation (Schärli et al., 2024) impose distinct challenges. Because identifying the source, extent, and time-constant of changes in the internal environment is usually difficult, assessing the corresponding CNS adaptation is also challenging. Particularly, structural alterations to the musculoskeletal system, whether due to injury, disease, or surgery, fundamentally change the body's

biomechanics and sensorimotor associations, but the quantification of these

changes is usually difficult. Accordingly, the way CNS remaps its motor control

strategies corresponding to the changes is not yet well understood (Walker et al.,

2004), although understanding such adaptations is crucial for elucidating the

pathophysiology of motor impairments observed in chronic musculoskeletal

conditions such as osteoarthritis, a degenerative joint disease, (Hunter and

Eckstein, 2009) and muscular dystrophy, which involves progressive muscle

weakness, (Mercuri and Muntoni, 2013).

To address this question, in this study, we employed a tendon transfer (TT)

surgery model (Sperry, 1940), which introduces a controlled, sustained change

to the musculoskeletal structure. TT surgery is a clinically well-established

procedure (Gardenier et al., 2020) that surgically re-attaches the tendon of a

specific muscle to that of a surrounding one. This procedure relocates a specific

muscle, so that its contraction generates a new mechanical action and,

consequently, novel somatosensory feedback. Because the internal change is

controlled by the experimenter, the TT model provides a powerful platform to

investigate how the CNS adapts to a new internal state. Another unique feature

of TT model is that it places permanent changes of the internal environment while

leaving the CNS anatomically intact. Unlike CNS lesion models (Hoffmann et al.,

2009), where the injury itself disrupts neural circuits and thereby complicates the

assessment of adaptive capacity of the CNS, TT offers a distinct advantage: it

allows investigation of CNS-driven adaptation without the confounding effects of

direct neural damage. Tendon transfer has been used in various species, with

studies suggesting different adaptive capacities, ranging from limited adaptation

in adult rodents (Sperry, 1940; Sperry, 1942; Slawinska and Kasicki, 2002;

Bowlus et al., 2003) to more substantial adjustments in cats (Yumiya et al., 1979;

Loeb, 1999) and primates, including humans (Lee and Seung, 1999; Wester et

al., 2018; Gaetz et al., 2023). Therefore, adaptability to TT appears enhanced in

primates and humans. Altogether, this approach provides a controlled platform

to examine how the CNS adapts to musculoskeletal changes in primates.

According to earlier reports, there are two motor control strategies that could be

involved when the CNS needs to face the sensorimotor remapping posed by

structural alterations to the musculoskeletal system. First, the CNS may employ

modular building blocks such as muscle synergies, coordinated activations of

muscle groups that reduce the dimensionality of motor control (d'Avella et al.,

2003; Bizzi and Cheung, 2013). Second, skilled behaviors like fine finger

movements often require fractionation, the capacity to activate muscles

independently. When confronted with structural changes to the musculoskeletal

system, does the CNS adapt by modulating existing synergies, or by shifting

toward more fractionated control strategies? Addressing this question requires

examining the evolution of neural control strategies over extended periods of

recovery to capture both immediate adjustments and long-term learning.

This study aimed to identify long-term adaptive mechanisms of the primate CNS

following structural changes to the musculoskeletal system. We investigated

whether the CNS adapts by modulating existing muscle synergies or by altering

the fractionated control of muscles specifically affected by the surgical alteration.

We surgically altered the limb structure by performing crossed tendon transfers

of finger extensor and flexor muscle tendons. Using a trained finger-grasping task

as our behavioral readout, we examined how the CNS recalibrates muscle activity

to regain skilled motor function. Our findings provide new insights into the

dynamic reorganization of motor control following structural changes of the body.

136
137

RESULTS

In this study, we developed a novel crossed tendon transfer animal model. This

involved surgically swapping the tendons of two antagonistic finger muscles in a

macaque monkeys' forearm (Fig. 1; A complete list of all muscle abbreviations,

their full names, and their assigned synergies for each monkey is provided in

Supplementary Table 1). The effectiveness and consistency of the surgery was

confirmed by measuring: (i) the distance traveled by muscle fibers and their

intramuscular tendons in the transferred muscle (Fig. 2B, C); and (ii) the amount

of fingertip displacement (Fig. 2D, E). Both measurements were induced by

percutaneous electrical stimulation over the transferred muscles (Fig. 2A). For a

detailed description of the results of these measurements see the Methods

section. The two monkeys were assigned slightly different tasks, allowing us to

examine controlled grasping in Monkey A and a more natural grasp in Monkey B

(refer to the Methods section, and Figs. 3 and Fig. 4 for further details). The two

monkeys ultimately performed related but distinct grasping tasks, a

methodological divergence that provided a valuable opportunity to test the

generality of the core adaptive mechanisms. Monkey A performed a controlled

grasping task requiring a fine precision grip, designed to study adaptation of fine

motor control (Fig. 3). While the same task was initially planned for both animals,

Monkey B performed this task inconsistently, frequently varying its grip strategy,

and we could not reinforce this monkey to perform in single strategy consistently.

Therefore, to ensure reliable task engagement and in accordance with the ethical

principle of Reduction and Refinement (the '3Rs'; (Tannenbaum and Bennett,

2015)), the task for Monkey B was modified to a more naturalistic food retrieval

grasp that the animal performed consistently (Fig. 4).

Once the monkeys had mastered their respective tasks, we recorded control

sessions and then performed the crossed TT. Post-surgery, and once the

monkeys had fully recovered and were able to perform the task independently

(~3–4 weeks), EMG recordings and behavior were resumed.

Functional Recovery Follows a Biphasic Trajectory

Despite the now reversed roles of the transferred muscles, the monkeys were

able to recover their grasping performance within two months, assessed by both

the return of key behavioral metrics to pre-surgical levels (Fig. 5A, D) and their

consistent ability to perform the task independently (Supplementary Videos S3,

S4, S7). However, this recovery was not immediate. The monkeys spent a period

of 1-2 weeks in their home cages for recovery, followed by several weeks of

assisted task practice. Formal recordings began once the monkeys could perform

the task consistently without assistance. The initial post-surgical period was

defined by a phase of significant motor impairment, primarily characterized by off-

target reaching trajectories (Fig. 5B, E) and prolonged grip formation times (Fig.

5A, D). These off-target reaching movements included inefficient, 'explorative'

trajectories in Monkey A and target overshoots in Monkey B (Supplementary

Videos S2, S3, S6). This recovery process was quantified across several

behavioral and kinematic metrics (Fig. 5). A comparison of pre-TT behavioral

variability on the grip formation task (Fig. 5D) revealed that Monkey B was

substantially more variable (CV = 81.93%) than Monkey A (CV = 46.62%). This

difference in variance was highly statistically significant (Ansari-Bradley test, p <

0.0001), confirming a difference in baseline stability between the subjects likely

due to the less constrained nature of Monkey B's task. Post-surgery, variability in

grip formation time initially increased dramatically for Monkey A (Early Phase CV

= 132.98%), while remaining high for Monkey B (Early Phase CV = 76.46%). In

the late phase, Monkey A's variability stabilized slightly above baseline (Late

Phase CV = 54.79%), whereas Monkey B's variability increased further,

exceeding its pre-TT level (Late Phase CV = 96.59%), suggesting persistent

inconsistency in grasp timing for this animal. Variability patterns differed for the

other metrics: Monkey A's pull time variability decreased below baseline in the

mid and late phases (Pre CV = 64.97%, Mid CV = 41.30%, Late CV = 43.01%),

indicating a refinement of this action, while Monkey B's grasp aperture variability

remained consistently low throughout recovery (Pre CV = 26.37%, Early CV =

23.80%, Mid CV = 19.78%, Late CV = 19.35%). The initial post-surgical period

was defined by motor performance that we classify as maladaptive due to the

persistence of inefficient, off-target reaching and prolonged movement durations.

First, task-related grip formation times were significantly longer immediately post-

TT (Monkey A: 197.7 ±92.2 vs. 660.6 ±221 ms, p = 0.014; Monkey B: 169.7

±14 vs. 316.3 ±31.9 ms, p = 0.016), taking approximately 40 days for Monkey

A and 48 days for Monkey B to return to and stabilize at pre-surgical levels (Fig.

5A, D). Second, the duration of off-target reaching was substantially elevated,

stabilizing approximately after 70 days in Monkey A and 55 days in Monkey B

(Fig. 5B, E). Following this phase, motor performance showed a gradual recovery

in efficiency (Fig. 5C) and kinematics (Fig. 5F). Ultimately, overcoming these

initial maladaptive behaviors through the gradual refinement of motor

performance led to the stabilization of grasping by the end of the experimental

period.

A two-phase adaptation is observed in the EMG activity of individual

muscles

Initially, we investigated whether EMG activity exhibited any changes post-TT.

Given that we interchanged the tendons of an antagonistic muscle pair,

functionally rendering the FDS an extensor and the EDC a flexor (as confirmed

mechanically, Fig. 2), a biomechanically sound adaptation would be to use the

former finger extensor (EDC) for finger flexion, and the former finger flexor (FDS)

for finger extension during a grasping task. We therefore analyzed EMG activity

from both transferred and non-transferred muscles to determine if these expected

functional changes occurred. Figure 6 shows a comparison of EMG activity

profiles for the two transferred muscles, EDC and FDS, in Monkey A (Fig. 6A–D).

Prior to surgery, the control showed distinct and contrasting EMG profiles in both

muscles. For instance, peak activity in EDC (▼) occurred long after completion

of the grasping action (indicated by vertical dashed lines at 0% task range) at

15% task time (Fig. 6A), coinciding with the animal pre-shaping its hand to

prepare for the subsequent grasp. Whereas peak activity in FDS ($\triangledown$) was

observed immediately after completion of the grasping action (0.49%; Fig. 6C).

The question we posed was whether the post-surgery activity of EDC resembled

its original profile (without adaptation to surgery) or the activity of FDS (the

anticipated profile following surgical adaptation), and vice versa. Our findings

suggested that EMG activity patterns largely shifted in a manner consistent with

the new mechanical function imposed by the transfer. Early after TT, peak activity

of EDC occurred at -0.58% task time (Fig. 6B; days 29 [red line] and 64 [orange]),

which aligns with peak activity of FDS prior to TT (Fig. 6C; black line). Also, for

FDS, peak activity occurred at 11% task time (Fig. 6D; ▼), which aligned with

peak activity of EDC (Fig. 6A).

This was corroborated by the results in Monkey B (Fig. 6, panels K-M). Although

the EMG patterns of both muscles varied significantly as the two monkeys

performed different types of grasping actions, we found that the EMG of EDC

peaked at -6.12% before (Fig. 6K; ▼) and 0.35% and 1.05% after TT (Fig. 6L; $\triangledown$,

days 22 and 36, respectively), closely matching the timing for FDS control data

(Fig. 6M). The corresponding analysis for FDS could not be performed in Monkey

B, since the EMG signal of FDS was lost early after TT (see Methods for details).

We were able to extend our recordings to 122- and 64-days post-TT for monkeys

A and B, respectively. This allowed us to examine whether the adaptations

observed in the early period were consistent throughout the experimental period.

These additional days (Fig. 6B, D; 69 [cyan], 79 [blue], and 99 [black] post-TT;

see Fig. S1 for recordings from all days) showed that the EMG activity profiles for

both muscles unexpectedly returned to their pre-TT state. For example, as

already described, the EMG signal of EDC at days 29 and 64 shared the

characteristics of FDS, but by the following day (only 5 days later, day 69 [light

blue] post-TT) already exhibited pre-TT characteristics of EDC, continuing as

such until the end of the recording period (day 79 [blue] and 99 [black]). This was

confirmed for the EDC muscle of Monkey B (Fig. 6K, L; Fig. S1U-AO for

recordings from all days).

To relate these neural changes to functional recovery, we overlaid the behavioral

error metric (off-target reaching time; gray dashed lines) onto the cross-

correlation plots (Fig. 6G–J). Figure 6H (red line) illustrates cross-correlation

coefficients between the EDC EMG profile post-TT (EDC-post) and the original

EDC control data (EDC-pre) in Monkey A. Figure 6I shows coefficients for EDC-

post and FDS-pre correlations in the same monkey. The coefficient

corresponding to the original profile decreased to between -0.3 and 0.2 over

approximately 65 days (Fig. 6H). In contrast, the coefficient corresponding to the

expected EMG profile exhibited an inverse pattern, with low coefficients pre-TT

and a high coefficient (of 0.9) post-TT (Fig. 6J). Cross-correlation analysis for

FDS (Fig. 6G, I) revealed a similar pattern. Cross-correlation coefficients for FDS-

pre- vs. FDS-post analysis declined from 1 to negative values, only to rebound to

their original values within a month (day 79; Fig. 6G). This outcome aligns with

our previous observations (Fig. 6B, D), and were corroborated by the results from

Monkey B (Fig. 6K-Q). Here, cross-correlation coefficients dropped from +0.95 to

+0.1 for EDC-pre vs. EDC-post comparison, and increased from -0.5 to 0.9 for

the FDS-pre vs. EDC-post comparison (Fig. 6P, Q). After 42 days post-TT, cross-

correlation coefficients started to gradually increase or decrease to +0.9 and -0.7,

respectively. This indicates that the anticipated initial changes in EMG activity

profiles are not permanent. Instead, they revert to their original patterns two

months post-TT. A supplementary analysis of the time lag at peak correlation

(Supplementary Figure S1) confirmed that the optimal lag for individual muscles

also fluctuated significantly during the early adaptive phase before stabilizing,

reflecting the process of temporal re-coordination of individual muscle commands.

Interestingly, changes in EMG pattern were not isolated to the transferred

muscles; non-transferred muscles exhibited a variety of complex adaptations

(Figs. 6, S2-S4). A comprehensive overview of the average EMG profiles for all

recorded muscles across all recording sessions and landmark days is provided

in Supplementary Figures S2 (Monkey A) and S3 (Monkey B), respectively. As

shown in these figures, many agonists adapted in concert with their transferred

synergist, following the same two-phase "swap-and-revert" pattern seen in Figure

6. In Monkey A, these included extensors ED23 and ECU, while in Monkey B,

they included ED23, ED45, and ECU (see Fig. S2 for details). In contrast, other

muscles showed patterns that were incompatible with a simple swap. For

example, the non-transferred flexor carpi radialis (FCR) in Monkey A showed a

distinct adaptive profile characterized by a drastic initial decrease in one activity

peak and a consistent increase in a later peak (Fig. 6E, F). A similar incompatible

pattern was seen in palmaris longus (PL) (Fig. S1H, I, R, S). Finally, some

muscles, like the extensor carpi radialis (ECR) in Monkey B, remained relatively

stable post-surgery (Fig. 6N, O). These diverse patterns in non-transferred

muscles do not necessarily contradict a modular control strategy (d'Avella et al.,

2003; Bizzi and Cheung, 2013); rather, they likely reflect the fact that different

muscles are primary members of different synergies (i.e., they possess the

highest weightings within specific modules; e.g., FCR and PL in Synergy C), each

undergoing its own task-specific adaptation.

Adaptation occurs through modulating the activation of stable muscle
304

synergies

To determine whether the CNS adapted by fractionating muscle control or

modulating existing modules, we extracted muscle synergies using non-negative

matrix factorization (NMF) (Lee and Seung, 1999). Four synergies accounted for

>80% (Cheung et al., 2012) of the variance in both monkeys (see Supplementary

Figure S5 for VAF plots; >85% for both monkeys; the results of the original and

shuffled data sets are shown. See also the synergy weights [W] and activation

profiles [C] for both monkeys and all synergies in figs. S6 and S7, respectively).

We first focused on the two primary synergies responsible for the main task axis:

the finger flexor synergy (Synergy A) and the finger extensor synergy (Synergy

B) (Fig. 7).

We first analyzed whether the spatial structure of these primary synergies

changed. To distinguish genuine adaptation from natural variability, we

established a baseline by calculating the cosine similarity of spatial synergy

weights (W) across all pre-surgery days (Fig. 7I, J, blue traces). This revealed

remarkably high stability (>0.99). Post-surgery, the core structure remained

conserved. The cosine distance between pre- and post-surgery weights for the

same synergy pairs was low (Monkey A: 0.03 ± 0.03; Monkey B: 0.09 ± 0.11),

whereas for different pairs it was high (>0.60). This confirms that the CNS

maintained the stable, neurally constrained building blocks of the primary flexor

and extensor modules despite the altered biomechanics.

In contrast to the stable spatial structure, the temporal activation profiles of these

primary synergies underwent drastic changes. In Monkey A, the flexor Synergy

A (originally peaking at +0.97% task time) and extensor Synergy B (originally

peaking at 12.62%) swapped their timing profiles shortly after surgery (Fig. 7C,

D). Specifically, the extensor synergy shifted to peak during the flexion phase,

and vice versa. This "swap" persisted during the early impairment phase before

reverting toward the original timing in the late phase. A similar "swap-and-revert"

pattern was observed in Monkey B (Fig. 7G, H), where the extensor Synergy B

shifted from a pre-surgery peak at -8.75% to a post-surgery peak of +3.5%

(matching the flexor phase) before recovering.

To formally quantify the quality of this reversion, we compared the pre-surgery

activation profiles with those from the final day of recording. We found that while

the temporal shape was highly preserved (Cosine Similarity >0.90), the profiles

remained statistically distinct (Permutation Test, $p < 0.0001$; Fig. S8). This

confirms that the recovered motor program represents a "good enough"

functional approximation rather than a perfect mathematical restoration of the

baseline.

We next examined the secondary synergies involved in wrist control and

stabilization (Fig. 8). In contrast to the coherent "swap" seen in the primary

synergies, these modules exhibited distinct, task-specific adaptations.

In Monkey A, Synergy C (a wrist flexor synergy involving PL and FCR) showed a

temporal shift that was incompatible with a simple swap (Fig. 8C). While its spatial

weights remained stable (Fig. 8A, I), its activation peak shifted from the post-

grasp phase (+1.46%) to the pre-grasp phase, increasing in magnitude over time.

This pattern corresponds to the recruitment of wrist flexors to enable the

compensatory tenodesis grasp. Synergy D (wrist extensor) underwent a single

notable change on day 29 but quickly reverted (Fig. 8D). In Monkey B, Synergy

C showed no discernible trend (Fig. 8G), while Synergy D exhibited a complex

pattern of immediate change followed by a gradual shift in the late adaptation

period (Fig. 8H). Notably, the activation amplitude of Synergy D remained

significantly suppressed compared to baseline (Fig. 8H; Fig. S6H). This

suppression of the wrist extensor likely reflects the specific biomechanical

requirements of the tenodesis strategy employed by Monkey B: relaxing the wrist

extensors facilitates the necessary wrist flexion for hand opening. These

independent changes in secondary synergies likely served to reinforce the overall

functional recovery driven by the primary modules.

Again, the characteristics of these changes in activation profiles of muscle

synergies were also quantitatively confirmed by cross-correlation analysis of all

four muscle synergies in both monkeys. The coefficients were plotted over the

course of experimental days in relation to tendon surgery together with the

overlaid behavioral error metric (off-target reaching time; gray dashed lines) (Fig.

9). In this analysis, the activation profile of each synergy was cross-correlated

with either the one from synergy A (finger flexor synergy; Fig. 9C, D, G, H) or

synergy B (extensor synergy; Fig. 9A, B, E, F) before TT. As previously

demonstrated for the FDS and EDC muscles, the temporal activation profiles of

muscle synergies A and B (flexor and extensor) displayed a distinct pattern. After

cross-correlation with their own control data, both the flexor and extensor

synergies showed coefficients of 1 prior to surgery. These became negative for

66 days post-surgery before reverting close to their original values (Fig. 9B, C).

However, after cross-correlation with control data of the antagonistic muscle

synergy (i.e., extensor for flexor synergy and vice versa, Fig. 9A, D), the pattern

was reversed. Coefficients started with negative values prior to TT and shifted to

positive coefficients near 1 shortly after surgery. In Monkey A, coefficients

returned to negative values around day 66 post-TT. In Monkey B, this reversal

occurred earlier, around day 48. In short, these results mirror our findings

between the transferred muscles (Fig. 6G, J). Furthermore, to determine if low

correlations were driven by simple timing mismatches, we analyzed the optimal

time lag yielding the maximum cross-correlation (Supplementary Figure S9). A

positive lag indicates the post-surgery activity is delayed relative to baseline,

while a negative lag indicates it is advanced. This analysis revealed significant

fluctuations in timing (switching between delays and advances) during the early

and mid-adaptation phases, particularly around the 'switch-back' period, before

stabilizing closer to zero lag in the late phase. For synergies C (main contributing

muscles: ECR, BRD, DEL) and D (main contributing muscles: ECU), the changes

in cross-correlation coefficients over time were markedly different. For synergy C, coefficients begin to steadily increase when cross-correlated with control data of the extensor synergy (synergy B, Fig. 10E), and decrease with the flexor synergy (synergy A, Fig. 10G). This suggests an increasing contribution to finger extension. In contrast, synergy D does not exhibit any specific pattern following TT.

In Monkey B, this cross-correlation analysis revealed a more varied, or differential, pattern of adaptation across the four synergies (Figs. 9 and 10). The primary extensor, Synergy B, mirrored the results in Monkey A, showing a clear swap-and-revert pattern (Fig. 9F, H). In contrast, Synergy A did not show a clear reversal; its correlation coefficients gradually converged towards zero, likely due to the absence of the FDS EMG signal in the analysis (Fig. 9E, G). The secondary synergies, Synergy C and D, showed no discernible trend (Fig. 10E-H).

In summary, both monkeys exhibited a distinct two-phase adaptation following TT. We define the 'early phase' as the period from initial post-surgical recovery up to the reversal of the swapped activation patterns (approximately days 20/29 to ~65 post-TT, see Fig. 9), characterized by the transferred muscles/synergies adopting antagonistic temporal profiles. We also identify a 'mid-adaptation' phase

covering the transitional period around the 'switch-back' event, characterized by

high variability. The 'late phase' encompasses the period following this switch-

back (after ~day 66 post-TT), where original activation timings were largely

restored. While this 'reversion' towards original activation timings in the late phase

marked the abandonment of the initial maladaptive strategy, behavioral recovery

involved more than just restoring original patterns. Concurrent with this neural

shift in the primary synergies, a distinct compensatory strategy began to emerge,

utilizing secondary synergies and biomechanical coupling to achieve functional

hand opening, as detailed below.

The Early Adaptation Phase is Characterized by a Maladaptive Neural and

Behavioral Profile

The early adaptation phase was defined by a distinct neural strategy that

correlated with significant behavioral deficits. First, the period dominated by off-

target reaching movements and prolonged grip times (Fig. 5B, E) was precisely

when the swapped activation of Synergy A and B was most prominent (Fig. 9;

see behavioral overlay). This temporal link provides strong evidence that this

initial 'swap' strategy was, in fact, maladaptive, as the flawed neural control

directly underpinned the impairments in movement efficiency and precision.

Second, we found that the net activity of muscles representing certain muscle

synergies (aggregated average EMG; aaEMG) showed distinct, synergy-specific

changes over time (Fig. 11). A key feature was the evolution of the activity of

finger extensor synergy (Synergy B), which appeared to undergo a

neuromuscular 'arms race', a rapid escalation of antagonistic co-activation, in

both animals, albeit on different timescales (Fig. 11B, J). In Monkey A, which

struggled for a longer period to regain behavioral proficiency according to our

subjective evaluation, our recordings now captured this process unfolding: we

observed a steady and significant increase in Synergy B's aaEMG post-TT until

day 64 ($p < 0.0001$) before the strategy changed (Fig. 11B, F). In contrast,

Monkey B, which adapted its behavior more rapidly according to our subjective

evaluation, showed a different profile; its Synergy B activity peaked with a

significant surge early in the recording period (day 22: $p < 0.0001$) and

subsequently declined toward and below baseline (day 64: $p < 0.001$) (Fig. 11J,

N). This suggests the initial, rapid escalation of the 'arms race' in this animal had

already occurred and peaked prior to our first post-operative recording session.

Therefore, the apparent discrepancy in the evolution of Synergy B is likely not a

fundamental difference in strategy, but rather a reflection of the different

observational windows and overall adaptation rates of the two animals.

For the finger flexor synergy (Synergy A), Monkey A showed a general decrease

in activity ($p = 0.0004$ at day 79; Fig. 11A, E), while Monkey B showed a

consistent and significant increase throughout the experiment (day 22: $p =$

0.0004; day 64: $p = 0.0008$). This was expected, as Monkey B's Synergy A relied

on FDP and PL to compensate for the loss of FDS function (Fig. 11I, M). Synergy

C exhibited a general increase in activity during the post-TT phase in both

monkeys (Fig. 11G, O), while Synergy D showed no significant differences in

Monkey B and only minor changes in Monkey A (Fig. 11H, P). All statistical

comparisons were made against the pre-TT control period using a two-sample t-

test with Bonferroni correction ($\alpha = 0.01$).

Taken together, these distinct patterns of aggregated EMG activity, especially the

escalating co-activation within the conflicted Synergy B, a known marker of

inefficient muscle recruitment (Thoroughman and Shadmehr, 1999; Osu et al.,

2002), further illustrate that this early adaptive phase, which coincided with poor

behavioral performance (Fig. 5), was characterized by an unstable and inefficient

neural control strategy.

Distinct Neural Implementations of a Compensatory Tenodesis Strategy

The final phase of adaptation was characterized by the development of a

compensatory tenodesis strategy. This effect describes the passive coupling of

finger and wrist movements due to the routing of tendons across multiple joints

(Zajac, 1992): because the finger extensor tendons pass over the back of the

wrist, actively flexing the wrist passively tightens these tendons, which in turn

pulls the fingers into extension (Valero-Cuevas and Hentz, 2002; Cash and Jones,

2011). For Monkey B, we found direct kinematic evidence for the acquisition of

this strategy. Post-surgery, the monkey learned to significantly increase finger

extension at the metacarpophalangeal (MCP) joint (Fig. 12A) by concurrently

flexing the wrist (Fig. 12B; $p < 0.0001$, ANOVA; $\alpha = 0.01$). This coordinated

movement pattern is characteristic of an active tenodesis effect. Furthermore,

kinematic analysis revealed that this was a gradually learned skill rather than an

immediate mechanical consequence. The trial-by-trial coupling between wrist

angle and MCP angle (Fig. 13) was initially absent ($R^2 = 0.00$) but strengthened

over weeks, peaking in the mid-adaptation phase (R^2 = 0.58). This period

coincided with the stabilization of grasp aperture (Fig. 5F) and the resolution of

the maladaptive neural patterns (Fig. 9). This kinematic strategy was supported

by a precise neural implementation. Unlike the scaling strategy observed in

Monkey A (see below), Monkey B did not rely on a massive increase in total

muscle activity; the aggregated activation of its primary flexor synergy (Synergy

A) showed consistent but moderate increases (Fig. 11I, M). Instead, the strategy

was one of temporal refinement. The activation profile of Synergy A shifted to

increase specifically during the pre-contact phase (Fig. 7G), providing the

necessary wrist flexion for the tenodesis grasp. This was achieved by a precise

sequential activation of muscles within the synergy, with the wrist flexor

component (PL) peaking just before contact and the finger flexor component

(FDP) peaking just after (Fig. S2). For Monkey A, which successfully restored its

primary extensor synergy, the tenodesis grasp likely served as a similar

compensatory driver. While we could not perform a kinematic analysis for this

animal due to low-resolution video images, strong evidence for this strategy is

provided by the neural data. We observed a clear temporal shift in the activation

of its dedicated wrist flexor synergy (Synergy C). The peak of this synergy's

activation moved from occurring just after object contact to just before it (Fig. 8C),

a re-timing well-suited to enable a tenodesis grasp. This increasing contribution

is further supported by cross-correlation analysis, which shows its activation

pattern became progressively more similar to that of the pre-TT extensor synergy

over time (Fig. 10A; $p < 0.05$, two-sample t-test).

Discussion

The central nervous system's (CNS) response to profound musculoskeletal change is a fundamental problem in motor control. This study sought to determine whether the primate CNS adapts to such change by flexibly modulating stable muscle synergies or by developing more fractionated, independent muscle control. We found that the CNS initially defaulted to a modular strategy, repurposing entire synergies by swapping their activation timings. This simple solution, however, proved to be maladaptive, creating a mechanical conflict that impaired motor function under tendon transfer surgery used in this study. This early maladaptive phase was ultimately resolved through the gradual development of compensatory movements, leading to a "good enough" functional recovery. This multi-stage process, operating on different timescales, highlights the intricate balance between modularity and flexibility in neural adaptation.

Neural Learning, Not Physical Healing, as the Rate-Limiting Factor

A crucial consideration in interpreting our long-term results is the potential confound of the tendon's physical healing process. However, the surgical

technique employed, a multi-weave Pulvertaft transfer, is designed to provide

immediate mechanical strength comparable to that of a native tendon (Graham

et al., 2023). While fibrous integration matures over approximately six weeks, the

repair was biomechanically sound from the beginning. This allows us to conclude

that the prolonged, multi-month recovery period, characterized by a complex two-

phase neural reorganization, was not limited by the tendon's force-bearing

capacity but rather reflects the significant challenge of a purely neural learning

process.

The CNS Defaults to a Modular Strategy, Leading to a Maladaptive Conflict

Our primary finding provides a direct answer to the question posed in the

introduction: the CNS adapted to the tendon transfer not by developing fine-

grained, fractionated control, but by implementing a modular strategy by

repurposing entire co-activation modules (Overduin et al., 2008; Berger et al.,

2013; Bizzi and Cheung, 2013; Takei et al., 2017). The initial and most immediate

neural change was a wholesale swap of the temporal activation patterns of the

primary flexor and extensor synergies (Synergy A and B). The fact that the spatial

structure of all four synergies remained remarkably stable throughout the months-

long experiment (Fig. 7I, S6), in the face of drastically altered biomechanics,

strongly supports the hypothesis that these synergies represent stable, neurally

constrained building blocks (d'Avella et al., 2003; Bizzi and Cheung, 2013; Takei

et al., 2017). While our baseline analysis (Fig. 7I) confirmed that natural variability

is negligible (>0.99 similarity), the specific, transient structural deviations

observed post-surgery (e.g., in the compensatory wrist flexor Synergy C) suggest

that the CNS retains a limited capacity to fine-tune the internal structure of these

modules when driven by strong functional demands. Consequently, the distinct

adaptive patterns observed in individual non-transferred muscles (e.g., FCR, PL

in Monkey A) likely reflect this task-specific tuning of secondary synergies, rather

than a breakdown of modular control.

Our result indicated that this modular approach is likely the default strategy. One

potential reason is that modulating pre-existing modules may be computationally

simpler and metabolically less costly than developing entirely new, fractionated

control patterns (Flash and Sejnowski, 2001; Bizzi and Cheung, 2013). The latter

would require extensive synaptic plasticity, potentially involving cortical

remapping to selectively uncouple previously co-activated muscles (Kitago and

Krakauer, 2013). Alternatively, or perhaps complementarily, the preservation of

synergies may reflect inherent constraints on neural plasticity, suggesting that the

underlying neural circuits encoding these modules are relatively immutable, even

when faced with significant changes in peripheral mechanics (Makin and

Krakauer, 2023). Regardless of the underlying reason, be it computational

efficiency, constraints on plasticity, or both, the CNS appears to prioritize

modulating the activation of existing modules when dealing with acute alterations

to the musculoskeletal system.

However, this adherence to established synergy structures created a

fundamental conflict. Activating the original extensor synergy (Synergy B) after

the transfer now inevitably co-activated the surgically transferred EDC, which

functioned mechanically as a flexor, alongside non-transferred muscles like ED23

and ECU, which remained anatomical extensors. This internal mechanical

antagonism appears to be the root cause of the early maladaptive phase. This

interpretation is supported by two key lines of evidence from our results. First, the

period of severe behavioral impairment, characterized by off-target reaching and

inefficient grasping (Fig. 5B, E), occurred precisely when this flawed "swap"

strategy was active. Second, the aggregated EMG activity revealed a sustained

and significant increase in the total activation of the conflicted Synergy B in

Monkey A (Fig. 11B, F), which we interpret as an energetically costly effort to

overcome the internal mechanical antagonism. This scenario can be viewed

through the lens of optimal control and cost-benefit analysis, where a cost

function is minimized (Wolpert, 1997). The initial "swap" strategy, while

computationally cheap to select, incurred an unacceptably high operational cost

in terms of both poor task performance (high error) and what was likely excessive

energy expenditure. We hypothesize that this profoundly unfavorable cost-benefit

ratio likely served as the critical error signal that drove the CNS to abandon this

initial strategy.

Resolution Through Slower Compensatory Adaptations

The CNS did not persist in the failed swap-based strategy. The high metabolic

and computational cost of activating a mechanically conflicted synergy likely

triggered the second rapid adaptation: the "switch-back" of synergy activation

timings toward their original patterns (Fig. 7, 9). This rapid reversion, occurring

over just a few days, is characteristic of an error-based learning mechanism, a

form of adaptation that is profoundly impaired by cerebellar damage across a

range of tasks, including adaptation to force fields (Smith and Shadmehr, 2005),

prismatic shifts (Martin et al., 1996), and split-belt treadmills (Morton and Bastian,

2006). It is plausible that the CNS operates with an implicit threshold for an

acceptable cost/performance ratio; once the persistent task failure and high

muscular co-contraction of the swap strategy exceeded this threshold, a swift

recalibration was initiated. This reversion, despite the remaining mechanical

antagonism from the transferred tendons, represents a "good enough" solution

where functional success is prioritized over perfect efficiency (Mussa-Ivaldi and

Bizzi, 1995; Ranganathan et al., 2013; Gijsberts et al., 2014). Our quantitative

analysis supports this interpretation: while the CNS successfully restored the

temporal structure of the motor commands (high cosine similarity), it did not

perfectly replicate the pre-surgery state. As shown in Figure S8, significant

differences in activation amplitude persisted in specific phases of the movement,

and permutation tests confirmed the profiles remained statistically distinct. This

suggests the system settled into a stable, functional attractor that was sufficiently

close to the original manifold to execute the task, without expending the

computational or metabolic cost required for a perfect restoration. The rapid

timescale of this change is highly consistent with the cerebellum's proposed role

as a forward model, predicting the sensory consequences of motor commands

and driving rapid learning in response to sensory prediction errors (Popa et al.,

2016). Our lag analysis (Fig. S2) confirmed that the timing of motor commands

fluctuated significantly before stabilizing.

Crucially, this switch-back to a less conflicted state, presumably representing the

involvement of the error-based learning process, was likely enabled only by the

concurrent development of slower, compensatory strategies that provided an

alternative means to achieve the task goal. The primary compensation was a

learned use of the tenodesis effect. Over weeks, the monkeys gradually

increased the activation of the wrist flexor synergy (Synergy C) during hand

opening. This wrist flexion biomechanically generates passive finger extension

(Zajac, 1992; Cash and Jones, 2011), providing a viable new method for hand

pre-shaping. This learned, compensatory behavior, confirmed by kinematic

analysis (Fig. 12), ultimately allowed the CNS to abandon the maladaptive

synergy swap. This gradual, exploratory process is distinct from the following

rapid adaptation and represents a form of motor skill acquisition, a process which

may be associated with plasticity in cortical structures like the motor cortex and

basal ganglia. (Kitago and Krakauer, 2013).

A Multi-Timescale Model Reconciles Smooth Behavioral Recovery and

Abrupt Neural Reorganization

The apparent paradox between the smooth, gradual recovery of motor function

and the abrupt, switch-like reorganization of the underlying neural control strategy

suggests that adaptation is not a single process. Instead, we propose it results

from the interaction of at least two distinct adaptive systems operating in parallel

on different timescales, a concept aligned with established two-state models of

motor learning (Smith et al., 2006a; Trewartha et al., 2014).

We map the gradual acquisition of the compensatory tenodesis strategy to the

slow process of this model. This iterative component is responsible for building

the robust, stable motor skills necessary for long-term recovery. In contrast, we

propose that the fast process drives the initial 'swap' strategy and its subsequent

abandonment. This system appears to be sensitive to immediate performance

errors and metabolic costs, triggering the rapid 'switch-back' to the original

synergy timings (recalibration) once the initial strategy proves inefficient.

The critical insight from our data lies in the interaction between these timescales.

We propose that the slow system effectively 'gates' the fast system: the abrupt

neural reorganization (the switch-back) is likely not an independent event, but a

transition enabled only when the slow learning (tenodesis) reaches a "good

enough" threshold to support function. This hierarchical interaction resolves the

tension between the smooth behavioral curve and the sharp neural transition,

ensuring a stable long-term motor plan (Fig. 14).

Broader Implications and Limitations

Mechanisms of Neural Adaptation. Our findings of a multi-timescale adaptation

process involving stable muscle synergies resonate with, yet also extend,

previous work on motor learning. Early studies in primates demonstrated a

capacity for eventual positive functional readaptation after nerve crossing,

moving beyond the initial maladaptive reversals also seen in our monkeys (Sperry,

1947). However, EMG studies in human patients suggest this adaptation may be

incomplete, as original 'old' muscle activation patterns often persist alongside

newly learned ones (Illert et al., 1986). This aligns with our observation of synergy

timing modulation rather than structural reorganization, a finding strongly

supported by 'virtual surgery' studies indicating the CNS preferentially adapts by

recombining existing motor modules (Berger et al., 2013).

Biomechanical Constraints and Sensory Feedback. The maladaptive conflict

('arms race') observed during the early phase may be exacerbated by peripheral

biomechanical factors. Rodent models show that epimuscular myofascial force

transmission can cause transferred muscles to generate unexpected antagonistic

forces depending on joint angle (Maas and Huijing, 2012). Such biomechanical

conflicts likely contributed to the cost and ultimate abandonment of the initial

swap strategy. Furthermore, the initial, rapid synergy swap may be linked to a

'sensory surprise' from altered proprioceptive feedback, driving a fast but flawed

response (Kitazawa et al., 1995; Martin et al., 1996; Smith et al., 2006b; Edwards

et al., 2012; Petitet et al., 2018). This multi-level network reorganization presents

a more complex challenge than the gain modulation within single or specific CNS

pathways seen in studies of spinal reflex conditioning (Thompson and Wolpaw,

2014). A comprehensive model must ultimately connect the global synergy

reorganization with these local, interacting feedback loops, considering factors

such as the roles of spinal reflex pathways (Smith et al., 2006c) and fusimotor

drive (Vallbo et al., 1979; Hospod et al., 2007).

Task-Dependent Differences. The observation that both monkeys, engaged in both a highly controlled learned task and a more naturalistic grasping task, showed similar adaptive patterns strengthens the generalizability of our findings (Berniker et al., 2013). However, the difference in adaptation rates likely reflects the distinct constraints of these tasks. Monkey A's precision task required exact force vector production, potentially prolonging the neuromuscular conflict. In contrast, Monkey B's naturalistic task allowed for more flexible kinematic solutions, such as the tenodesis grasp, to be discovered and employed more rapidly. This suggests that while the core neural mechanism (swap-and-revert) is conserved, the timescale of its resolution is sensitive to task constraints.

Motor Memory and Unlearning. Our paradigm raises critical questions about the distinction between unlearning a maladaptive strategy and relearning a functional one (Kitago et al., 2013). The disappearance of the initial "swap" strategy could be due to passive decay, active inhibition (extinction), or interference from a new, stronger memory (Villalta et al., 2013). The literature increasingly suggests that unlearning is an active process, and that the new, successful compensatory strategy likely supplants the maladaptive one through interference. Our results

lend weight to this view, suggesting functional recovery is a process of

competitive memory formation.

Future Directions. A valuable future direction would be to combine the empirical

findings of this study with theoretical biomechanical modeling. A subject-specific

musculoskeletal model could provide a more precise mapping of how the tendon

transfer altered the biomechanical plant. Integrating these empirical and

theoretical approaches represents a critical next step in understanding the

complex interplay between biomechanics and neural adaptation (Nakajima et al.,

2023).

Implications for Sensorimotor Rehabilitation

These findings may hold implications for sensorimotor rehabilitation. The

observation of a distinct maladaptive phase followed by compensation and

recovery suggests a staged approach could be beneficial (Hatem et al., 2016;

Kakavas et al., 2025). An early phase of therapy might focus on minimizing

sensory prediction errors caused by the altered proprioceptive feedback, and

reducing maladaptive responses through techniques aimed at integrating the

altered body schema (e.g., using visual feedback or task-oriented practice) (Nudo

et al., 1996; Takeuchi and Izumi, 2012; Zeiler and Krakauer, 2013; Foell et al.,

2014; Levin et al., 2015; De Nunzio et al., 2018; Sattin et al., 2023; van Vliet et

al., 2023). This stage would guide the fast, error-driven learning system. A later

phase could then employ task-specific training to promote effective

compensatory strategies, facilitating the abandonment of maladaptive patterns

and the consolidation of a stable motor strategy (Murata et al., 2008; Maier et al.,

2019). This second stage engages the slow, skill-acquisition system by creating

a rich, problem-solving environment. Finally, a third stage would focus on

intensive, high-repetition practice to automate the new functional motor plan,

making it robust for real-world use through dosage and generalization

(Winterbottom and Nilsen, 2024). This neurobiologically-informed staging,

guiding fast adaptation, then facilitating slow skill learning, then driving

consolidation, may offer a more logical and effective path to functional recovery.

METHODS

Animals

Data were collected using two male macaque monkeys (*Macaca fuscata*; Monkey A: 7.8 kg and Monkey B: 9.9 kg) purpose-bred at the National Bioresource Project (NBRP). They were kept in custom-made primate cages, allowing for potential pair housing. Both monkeys were trained to perform a simple grasping task. During experimental recordings, the monkeys were seated in a primate chair without further restraint. Their head movement remained unrestricted at all times.

Procedures

All procedures were designed to minimize discomfort and pain and approved by the institutional Animal Care and Use Committees at the National Center of Neurology and Psychiatry (NCNP), Tokyo, Japan. Details of the surgical operations, experimental setup, and procedures for electromyography (EMG) recordings have been previously described (Takei and Seki, 2010).

EMG implant surgery

Both animals were familiarized with the experimental setup and trained to perform

the behavioral task prior to surgery. After an initial training period, EMG

electrodes were chronically implanted subcutaneously into several muscles of the

left forearm including the FDS, FDP, and EDC (Figs. 1, 3C, and 4C). Individual

muscles were localized and confirmed using electrical microstimulation before

implanting EMG electrodes chronically by two methods. For muscles involved in

the crossed tendon transfer procedure, muscle fascia was cut and EMG wires

with looped ends were placed on top of the muscle belly. These were then

secured by reclosing the fascia with absorbable suture threaded through the loop

(AS-361, stainless-steel Cooner wire, Conner Wire Co., Chatsworth, CA, USA).

All other muscles were directly implanted into the muscle belly using a 22-gauge

hypodermic needle. Each wire was threaded into the needle tip, folded back along

the shaft, and inserted into the muscle before carefully retracting the needle.

Electrode separation for bipolar recordings was approximately 5–10 mm. Note

that some EMG recording sites were lost over time. Surgical procedures were

carried out under deep general anesthesia (sevoflurane 1.5 to 2.5% in 2:1

O_2/N_2O) and with full aseptic precautions. Heart rate, blood pressure, body

temperature, and blood oxygen saturation were monitored throughout surgery.

Analgesics and antibiotics were administered intramuscularly for at least one

week postoperatively.

Tendon-Transfer Surgery

After a minimum four-week recovery period from EMG implant surgery, TT

surgery was performed. The FDS and EDC tendons were cut as distally as

possible (immediately below and above the wrist for FDS and EDC, respectively)

to avoid damaging the Golgi tendon organ located near the junction between the

muscle fibers and tendon (Schoultz and Swett, 1972; Jami, 1992). Tendons were

then guided either through the gap between the radius and ulnar bone (Monkey

A), or around the wrist and then re-attached to the tendon of the antagonist

muscle using a tendon graft harvested from the plantaris tendon of the lower limb

(Monkey B; EDC → FDS: direct connection; and FDS → EDC: tendon grafting).

The tendon coaptation was performed using a Pulvertaft weave technique

(weaves > 2) with high-strength non-absorbable suture material to ensure

immediate and robust mechanical stability post-surgery (Graham et al., 2023).

This was only necessary in Monkey B since the tendons used for the cross-

transfer were too short. This surgical procedure aimed to reverse the primary

mechanical actions of the manipulated muscles, making the FDS tendon an

effective finger extensor and the EDC tendon an effective finger flexor. The

success and nature of this mechanical rearrangement were verified post-

operatively via direct muscle stimulation (see Methods subsection 'Tendon

transfer confirmation' and Fig. 2). The general organization and function of the

major forearm muscles involved in finger flexion and extension in *Macaca fuscata*

are broadly comparable to those in humans (Vanhoof et al., 2021; Yan et al.,

2022).

Surgical procedures were carried out under deep general anesthesia

(sevoflurane 1.5 to 2.5% in 2:1 O_2/N_2O) and with full aseptic precautions. Heart

rate, blood pressure, body temperature, and blood oxygen saturation were

monitored throughout surgery. Analgesics and antibiotics were administered

intramuscularly for at least one week postoperatively. Both monkeys wore a

plaster cast post-surgery, but its effectiveness was limited (lasting approximately

one week).

Tendon transfer confirmation

To verify the success and long-term stability of TT surgery, two procedures (Fig.

2) were implemented. During the first procedure, the monkey was sedated and

its arm was secured to a metal frame (Fig.2A). Electrical stimulation was then

applied to either the FDS or EDC muscle (50 mA, DS8R, Digitimer, Welwyn

Garden City, UK), while ultrasound scans of tendon movements were

concurrently captured (SONIMAGE MX1, Konica Minolta, Inc., Tokyo, Japan

(Nordez et al., 2009; Dieterich et al., 2017).

Figure 2B shows a sonogram of the FDS muscle and its intramuscular tendons.

The left side displays a static image of the FDS muscle at a specific moment, with

white arrows indicating the FDS tendon used for measurement. The right side

shows staggered images (top white box) of the FDS tendon displacement

triggered by muscle stimulation. The lower right inset illustrates the measurement

area. The displacement wave area represents the intensity of muscle contraction,

and was calculated by measuring the average duration (a, in seconds) and

amplitude (b, in cm) of three successive waves. The area (a*b/2) and regression

line for FDS (red) and EDC (blue) ($R^2 > 0.5$ for FDS) for days 0, 7, and 105 post-

TT are shown (Fig. 2C). The results suggest that the muscle contractions induced

by direct electrical stimulation remained nearly constant.

For the second procedure, high-speed videos (37U Series Color Industrial

Camera, The Imaging Source, Charlotte, NC, USA) of the monkey's finger

movements were recorded while electrically stimulating the FDS or EDC. Markers

were placed on the nails of the index, middle, and ring fingers (Fig. 2A) to

measure finger displacement in xyz-dimensions using DeepLabCut software

(Mathis et al., 2018; Nath et al., 2019). The sum of Euclidean distances of each

marker from the origin of the three-dimensional (3D) coordinate system was

computed as a scalar quantity. Post-surgery, movement along the z-axis

reversed, indicating a shift from finger flexion to extension due to TT (Fig. 2D;

blue: pre-TT on surgery day, dark brown: post-TT on surgery day, light brown: 1

wk post-TT, and red: 3 wk post-TT). The scalar quantities of each finger did not

significantly change during muscle stimulation on days 0-, 7-, and 105-days post-

TT (Fig. 2E), suggesting there was no postoperative tendon rupture or slackening.

Data recordings

EMG signals were recorded using the AlphaLab SnR system (Alpha Omega

Engineering, Hamerkava St. 6, Ziporit Industrial Zone, P.O.Box 810, Nof HaGalil

(Nazareth Illit) 1789062, Israel), displayed online and then stored on a hard drive

for later off-line analysis using MATLAB (MathWorks, Natick, MA, USA). Data

were recorded at 11 kHz and processed for analysis as follows. 1. Down-sampled

to 5 kHz; 2. 50 Hz high-pass filtered (6th order Butterworth filter); 3. rectified; 4.

20 Hz low-pass filtered (6th order Butterworth filter); and 5. down-sampled to 100

Hz. Behavioral task events were recorded as transistor–transistor logic (TTL)

signals using the AlphaLab SnR system. Monkey behavior was further recorded

by two cameras (Sanyo VPC-WH1, 60 fps [Sanyo, Osaka, Japan]; and Casio EX-

100F, 240 fps [Casio, Tokyo, Japan] for monkeys A and B, respectively) from two

different angles (top and side view). The images were later used to detect

additional behavioral events (contact time with the object and pull onset for

Monkey A; and food touch and food lift onset for Monkey B), and detection of

possible changes in the animal's general movement pattern. Kinovea software

was used for this video analysis (free and open-source software;

https://www.kinovea.org/).

 Behavioral task

Before and after EMG surgery, the monkeys were trained on a simple grasping

task that involved a small object attached to a rod (Fig. 3A for Monkey A; and Fig.

4A for Monkey B). For Monkey A, object 1 was a small rod placed between two

side walls encouraging the monkey to grasp the rod with a precision grip using

the tips of the index finger and thumb (controlled grasp). The force used to

compress the spring while pulling was low and the distance moved short. Object

2 had to be grasped in the same way. However, the force to compress the spring

was higher and the distance moved longer (object 1: 300 cN, 4 mm; and object

2: 800 cN, 30 mm). The monkey was asked to grasp and hold the objects for

approximately 300–500 ms (Fig. 3A–D). After completing the trial, the monkey

was rewarded with a piece of fruit that had to be taken from the experimenter's

palm. The task sequence for Monkey A ranged from object 1 hold onset to object

2 hold offset (Fig. 3C, D; and 'Data Analysis' below). For Monkey B, object 1 was

a small rod that was grasped and pulled (300 cN, 4 mm) using a lateral

prehension grip (like using a door key), and resembling a power grip rather than

a natural grasp. However, instead of grasping object 2, Monkey B picked up a

piece of fruit from an allocated location. In doing so, the monkey had to cross the

path of a photo cell which was detected by the recording system (Fig. 4A–D). The

task sequence to be analyzed for Monkey B ranged from object 1 hold onset to

food touch, which was indicated by a LED (Fig. 4C, D; and 'Data Analysis' below).

For Monkey A, the task sequence was as follows (Fig. 3B; and Supplementary

Video S1). From a starting position in front of the monkey, it had to lift its arm and

move towards object 1, grasp, pull, and then hold the object for 500 ms.

Immediately after releasing object 1, Monkey A proceeded to move its hand

towards object 2, which was located to the right of object 1. Again, the monkey

had to grasp, pull, and then hold the object for another 500 ms. Each hold period

was accompanied by an audio signal which stopped once the hold duration was

sufficient. After completing the task, which was signaled by another acoustic

signal, the monkey was rewarded with a piece of fruit presented by the

experimenter and taken by the monkey.

For Monkey B, the first part of the sequence was identical. However, after

releasing object 1, Monkey B was required to pick up a piece of fruit from an

allocated location (Fig. 4A, B; and Supplementary Video S5). In doing so, the

monkey passed a photocell in front of the food well. This event was detected by

the recording system and a video camera (a red LED triggered by activation of

the photocell). For the analysis, only object 1 and the food grasp were considered

in monkeys A and B, respectively.

Data analysis

<u>*EMG analysis*</u>

EMG data were normalized to the average time for each monkey to complete a

trial (object 1 hold onset ➔ object 2 hold offset/LED offset = 100%). Data were

cropped and aligned according to the time stamps (Obj1 hold offset ± 15%; and

LED onset [food touch] ± 15% for monkeys A and B, respectively). Recorded

EMGs from the pre- and post-tendon surgery period were averaged for each

recording and compared over experimental sessions. For Monkey A, FDS and

EDC were analyzed both pre- and post-TT. For Monkey B, FDS and EDC were

analyzed pre-TT, but only EDC was analyzed post-TT due to signal

deterioration.

Both monkeys required time to recover from surgery. Post-TT recordings were

resumed once the monkeys were able to perform the task independently (i.e.,

without assistance from the experimenter) and met specific trial count criteria

(starting 29 and 20 days post-surgery for monkeys A and B, respectively). Post-

operative recording sessions were conducted frequently throughout the recovery

period, though not at fixed daily intervals, reflecting the practical constraints of

long-term behavioral experiments (e.g., animal cooperation, experimental

scheduling) and the aim to capture data during key phases of adaptation. Within

any given session included in the analysis, behavioral (video) and EMG data were

collected concurrently. To be included in the behavioral analyses (e.g., grip

formation times, off-target reaching, kinematics), a session required a minimum

of 20 successful trials. For the more demanding muscle synergy analyses, a

higher threshold of at least 100 successful trials was required to ensure robust

factorization. This difference in criteria may have resulted in slightly different sets

of recording days being represented across behavioral versus synergy-related

figures. After tendon surgery, EMG signals for FDS, FCU, and FCR deteriorated

in Monkey B. Therefore, there is no experimental data for these muscles in this

monkey and they were excluded from the muscle synergy analysis.

 Synergy analysis

The EMG envelopes obtained from pre-processing raw EMG data were divided

by the mean value to normalize activity. Muscle synergies were then extracted

for each session using NMF (d'Avella et al., 2003). NMF decomposes the EMG

data matrix M, as a product of two matrices C and W:

$M = CW^T + re = \Sigma c_i w_i{}^T + re$

where the vector wi and column of matrix W are muscle synergies; the vector ci,

and column of matrix C are their temporal coefficients; and re represents the

residuals. Multiplicative update rules were used for decomposition (Lee and

Seung, 1999). Updating matrices for decomposition from 20 different random

initial matrices was started, and factorization results with least residuals among

the 20 results were used. To improve robustness of the muscle synergies, one

set of muscle synergies was extracted from EMG data of multiple trials of each

day. A k-fold cross validation was used, with k = 4 i.e., EMG data set of multiple

trials during the day was split into four data sets, in which three data sets were

for training and one data set was for testing. Synergy matrix W was calculated

from three training data sets using NMF, and then a coefficient matrix C was

calculated by decomposing the EMG matrix M of the test data set with a fixed W

matrix of the training data set. This operation was repeated by changing the test

data set for each four data sets. Finally, the extracted four C matrices and four W

matrices were averaged to obtain daily synergies.

The number of synergies was determined as the number where VAF exceeded

a threshold. This threshold was set as 0.8. To clarify the effect of TT, the number

of synergies for each monkey was determined, assuming that trial-by-trial

variation in synergies within the same monkey was small. MATLAB 'nnmf'

function was used for NMF.

To determine whether the weights of muscle synergies changed after TT, the

following procedure was used. First, the average of each pre-TT spatial synergy

was calculated and used as control data to calculate the cosine distance for all

post-TT spatial synergies. This generated four distance relationships for each

pre-TT synergy. If the change in spatial synergy before and after TT was small,

then only one of the four cosine distances (e.g., synergy A before TT and synergy

A after TT) should be significantly smaller. Two-way ANOVA was conducted

using the type of synergy and session as factors for the cosine distance. The

significance of synergy pairs was identified using the Bonferroni post-hoc test.

Quantitative Comparison of Synergy Profiles:

To quantify the degree of 'reversion' in the late adaptation phase, we compared

the synergy activation profiles from the pre-surgery period with those from the

final recording day using three distinct metrics. First, Cosine Similarity was

calculated to assess the similarity in the shape of the temporal profiles

independent of amplitude. Second, a Permutation Test (n=10,000 iterations) was

performed to test whether the specific trajectory of the post-surgery profile was

statistically distinguishable from the pre-surgery distribution. Third, to identify

specific phases of the task where activation differences persisted, we conducted

a point-by-point Wilcoxon rank-sum test at each time point of the normalized task

cycle, applying a Bonferroni correction for multiple comparisons.

Cross-correlation analysis

Cross-correlation analysis was performed to examine the similarity between pre-

and post-surgery EMG signals as well as temporal activation profiles of extracted

muscle synergies. For the EMG signal analysis presented in Figure 6, the cross-

correlation coefficient at zero-time lag was calculated. For the synergy analysis

presented in the main figures (Figs. 9, 10), the cross-correlation coefficient at

zero-time lag was calculated using MATLAB (MathWorks, corrcoef function) to

quantify similarity without temporal shifts. These zero-lag coefficients were

plotted over experimental sessions. In a supplementary analysis, we also

computed the cross-correlation allowing for variable time lags using MATLAB's

xcorr function (normalized using the 'coeff' option). From this, we extracted the

specific time lag that yielded the maximum cross-correlation coefficient (optimal

lag) for each session for both the muscle synergies (Supplementary Figure S9)

and the individual EMG signals (Supplementary Figure S1). In this analysis, a

positive lag indicates that the post-surgery activation profile is delayed (occurs

later) relative to the pre-surgery baseline, while a negative lag indicates it is

advanced (occurs earlier).

Based on cross-correlation analysis, five 'landmark days' were identified for

further analysis (Fig. 6P). These days represent distinct stages in the recovery

process and excluded pre-TT control data. The first landmark day was chosen

from one of the initial recording sessions after TT surgery (days 29 and 22 in

monkeys A and B, respectively) when the cross-correlation coefficients had

changed significantly compared with control data. The second day was then

chosen from a time period just before the switch-back, when the cross-correlation

coefficients had started to return to their original values (days 64 and 36). Another

day was then picked from the period when the coefficients were still changing

significantly (days 69 and 44) and before starting to saturate. At this point the next

day was defined (days 79 and 48). Finally, the final landmark day was chosen

from one of the last recording sessions when the behavior had fully recovered

(days 99 and day 64 for monkeys A and B, respectively).

1003

Behavioral analysis

To examine behavioral recovery, the duration, onset, and offset of object and

food grasps were analyzed. Event times extracted from the video analysis were

used for alignment of EMG data and subsequent cross-correlation and synergy

analyses, along with recorded event TTL signals. For each experimental session,

the first 20 trials of each video recording were analyzed. In Monkey A, the

following events were detected and in-between times stored in ms: touch onset,

pull onset, and hold onset (resulting in grasp duration and pull time) for object 1

and object 2. In Monkey B, the following events were detected and in-between

times stored: touch onset, pull onset, and hold onset (resulting in grasp duration

and pull time) for object 1; and for the food grasp component of the task, LED

onset, food-touch onset, food-lift onset, LED offset, and movement end (contact

of food with mouth).

Contact times (touch onset ➔ pull onset) for Monkey A were plotted in ms (mean

± SD) over number of days from tendon surgery for object 1. For Monkey B,

contact times with food (food-touch ➔ food-lift onset) were plotted in ms (mean

± SD).

ANOVA was performed to compare control data recorded before and after tendon

surgery. In total, 5 and 3 control sessions vs. 35 and 22 experimental sessions

were used for analyses in monkeys A and B, respectively.

To quantify and compare the baseline behavioral variability of these metrics, we

analyzed all pre-TT trials for each animal's respective tasks. We calculated the

Coefficient of Variation (CV), defined as the standard deviation divided by the

mean, for each metric (grip formation time, pull time, and grasp aperture). To

formally test the difference in variance for the grip formation task (Fig. 5A, D), we

used the non-parametric Ansari-Bradley test. To quantify the observed off-target

reaching behavior, video footage was analyzed and the means of 10 consecutive

trials calculated for each session as follows. For Monkey A, time spent within a

specific spatial window covering the area behind and between the two objects

was measured; the monkey passed through this space almost exclusively while

exhibiting the impairment (see Fig. 3E, crossing of the yellow dotted line). In

Monkey B, the time spent in contact with the rear plate of the object was

measured (see Fig. 4E).

In addition to the primary behavioral metrics, we performed a more detailed video

analysis of movement kinematics and performance. For Monkey A, we quantified

the 'pull time' for each trial, defined as the duration from the moment the monkey

started pulling the object to the moment hold onset. For Monkey B, we quantified

the 'grasp aperture'. This was defined as the distance between the tips of the

index finger and thumb, measured just before the monkey made contact with the

food pellet.

Joint kinematics analysis

To determine whether compensatory movements occurred, steady state changes

in joint angles were examined before and after recovery. Using DeepLabCut

(Mathis et al., 2018; Nath et al., 2019), key points on the fingers and wrists were

tracked from experimental videos. From the obtained key points, the vector v_2

connecting the wrist and fingers and vector v_2 connecting the arm and wrist were

estimated (Fig. S10). The joint angle θ was calculated as follows:

1051

1052
$$cos(\Theta) = \frac{v_2 \cdot v_3}{|v_2| |v_3|} \rightarrow \Theta = cos^{-1}\left(\frac{v_2 \cdot v_3}{|v_2| |v_3|}\right)$$

To quantify the learning and refinement of the compensatory tenodesis strategy

over time, we analyzed the trial-by-trial relationship between wrist and MCP joint

angles for Monkey B. For each recorded day, we performed a linear regression

between the wrist angle (predictor) and the MCP angle (outcome) across all

successful trials (mean of 20 trials ±SD for each landmark day, taken 83ms before

food touch). The strength of the coupling was quantified using the coefficient of

determination (R^2). These daily scatter plots and their corresponding R^2 values

were visualized to track the evolution of the kinematic coupling from the pre-TT

baseline through the post-surgery adaptation period.

1062

1063 **DATA, MATERIALS, AND SOFTWARE AVAILABILITY**

1064

1065 Matlab scripts and raw data will be made available online. All study data are

1066 included in the article and/or SI Appendix.

ACKNOWLEDGEMENTS:

We would like to express our sincere gratitude to Drs. Francisco Valero-Cuevas and Andrea d'Avella for their invaluable comments and constructive suggestions on the earlier version of the manuscript, and Drs Nicholas Schweighofer and Gerald E. Loeb for stimulating discussion. We would also like to thank Dr. Naomichi Ogihara and Prof. Toshiyasu Nakamura for their valuable insights and technical advice on tendon surgery in the hand and forearm of primates. We are grateful to Drs. Kumiko Oida and Chika Sasaki for their medical care of the experimental animals, vital help during surgery, and training of the animals. Lastly, we would like to thank Masahi Koizumi for designing parts of the experimental setup and Drs. Joachim Confais, Shinji Kubota, Saeka Tomatsu, Tatsuya Umeda, and Natsumi Uchida for their support and assistance during the proceeding pilot study. We thank Rachel James, PhD, from Edanz (https://jp.edanz.com/ac) for editing a draft of this manuscript.

This work was supported by Grants-in-Aid for Scientific Research from the Ministry of Education, Culture, Sports, Science, and Technology of Japan (26120003, 26250013, 15K21754, 19H05724, 19H01092, 23H05488, and 24K21313), the Japan Agency for Medical Research and Development

1095

REFERENCES

Berger DJ, Gentner R, Edmunds T, Pai DK, d'Avella A (2013) Differences in Adaptation Rates
after Virtual Surgeries Provide Direct Evidence for Modularity. The Journal of
Neuroscience 33:12384-12394.

Berniker M, O'Brien MK, Kording KP, Ahmed AA (2013) An Examination of the
Generalizability of Motor Costs. PLOS ONE 8:e53759.

Bizzi E, Cheung VC (2013) The neural origin of muscle synergies. Frontiers in Computational
Neuroscience Volume 7 - 2013.

Bowlus TH, Lane RD, Stojic AS, Johnston M, Pluto CP, Chan M, Chiaia NL, Rhoades RW
(2003) Comparison of reorganization of the somatosensory system in rats that sustained
forelimb removal as neonates and as adults. J Comp Neurol 465:335-348.

Cash DJW, Jones JWM (2011) The role of tenodesis in surgery of the upper limb. The Journal of
Bone & Joint Surgery British Volume 93-B:285-292.

Cheung VC, Turolla A, Agostini M, Silvoni S, Bennis C, Kasi P, Paganoni S, Bonato P, Bizzi E
(2012) Muscle synergy patterns as physiological markers of motor cortical damage. Proc
Natl Acad Sci U S A 109:14652-14656.

d'Avella A, Saltiel P, Bizzi E (2003) Combinations of muscle synergies in the construction of a
natural motor behavior. Nature Neuroscience 6:300-308.

Davidson PR, Wolpert DM (2003) Motor learning and prediction in a variable environment.
Curr Opin Neurobiol 13:232-237.

De Nunzio AM, Schweisfurth MA, Ge N, Falla D, Hahne J, Gödecke K, Petzke F, Siebertz M,
Dechent P, Weiss T, Flor H, Graimann B, Aszmann OC, Farina D (2018) Relieving
phantom limb pain with multimodal sensory-motor training. J Neural Eng 15:066022.

Dieterich AV, Botter A, Vieira TM, Peolsson A, Petzke F, Davey P, Falla D (2017) Spatial
variation and inconsistency between estimates of onset of muscle activation from EMG
and ultrasound. Sci Rep 7:42011.

Edwards MJ, Adams RA, Brown H, Pareés I, Friston KJ (2012) A Bayesian account of 'hysteria'.
Brain 135:3495-3512.

Flash T, Sejnowski TJ (2001) Computational approaches to motor control. Curr Opin Neurobiol
11:655-662.

Fleury L, Prablanc C, Priot AE (2019) Do prism and other adaptation paradigms really measure
the same processes? Cortex 119:480-496.

Foell J, Bekrater-Bodmann R, Diers M, Flor H (2014) Mirror therapy for phantom limb pain:
brain changes and the role of body representation. Eur J Pain 18:729-739.

Gaetz W, Dockstader C, Furlong PL, Amaral S, Vossough A, Schwartz ES, Roberts TPL, Scott

1131   Levin L (2023) Somatosensory and motor representations following bilateral transplants
1132   of the hands: A 6-year longitudinal case report on the first pediatric bilateral hand
1133   transplant patient. Brain Res 1804:148262.

1134 Gardenier J, Garg R, Mudgal C (2020) Upper Extremity Tendon Transfers: A Brief Review of
1135   History, Common Applications, and Technical Tips. Indian J Plast Surg 53:177-190.

1136 Gijsberts A, Bohra R, Sierra González D, Werner A, Nowak M, Caputo B, Roa MA, Castellini C
1137   (2014) Stable myoelectric control of a hand prosthesis using non-linear incremental
1138   learning. Front Neurorobot 8:8.

1139 Graham EM, Oliver JD, Hendrycks R, Maglic D, Mendenhall SD (2023) Alternative Tendon
1140   Coaptations to the Pulvertaft Weave Technique: A Systematic Review and Meta-Analysis
1141   of Biomechanical Studies. HAND 18:446-455.

1142 Green HJ (1997) Mechanisms of muscle fatigue in intense exercise. J Sports Sci 15:247-256.

1143 Hatem SM, Saussez G, Della Faille M, Prist V, Zhang X, Dispa D, Bleyenheuft Y (2016)
1144   Rehabilitation of Motor Function after Stroke: A Multiple Systematic Review Focused on
1145   Techniques to Stimulate Upper Extremity Recovery. Front Hum Neurosci 10:442.

1146 Hoffmann G, Kamper DG, Kahn JH, Rymer WZ, Schmit BD (2009) Modulation of stretch
1147   reflexes of the finger flexors by sensory feedback from the proximal upper limb
1148   poststroke. J Neurophysiol 102:1420-1429.

1149 Hospod V, Aimonetti J-M, Roll J-P, Ribot-Ciscar E (2007) Changes in Human Muscle Spindle
1150   Sensitivity during a Proprioceptive Attention Task. The Journal of Neuroscience
1151   27:5172-5178.

1152 Hunter DJ, Eckstein F (2009) Exercise and osteoarthritis. J Anat 214:197-207.

1153 Illert M, Trauner M, Weller E, Wiedemann E (1986) Forearm muscles of man can reverse their
1154   function after tendon transfers: An electromyographic study. Neuroscience Letters
1155   67:129-134.

1156 Jami L (1992) Golgi tendon organs in mammalian skeletal muscle: functional properties and
1157   central actions. Physiol Rev 72:623-666.

1158 Kakavas G, Brilakis E, Papatzikou M, Malliaropoulos N, Mazeas J, Forelli F (2025) Reverse
1159   Linear Neuro Periodization Model for Rehabilitation After Arthroscopic Rotator Cuff
1160   Repair: A Narrative Review. Clinics and Practice 15:105.

1161 Kitago T, Krakauer JW (2013) Motor learning principles for neurorehabilitation. Handb Clin
1162   Neurol 110:93-103.

1163 Kitago T, Ryan SL, Mazzoni P, Krakauer JW, Haith AM (2013) Unlearning versus savings in
1164   visuomotor adaptation: comparing effects of washout, passage of time, and removal of
1165   errors on motor memory. Front Hum Neurosci 7:307.

Kitazawa S, Kohno T, Uka T (1995) Effects of delayed visual information on the rate and amount

of prism adaptation in the human. J Neurosci 15:7644-7652.

Lee DD, Seung HS (1999) Learning the parts of objects by non-negative matrix factorization.

Nature 401:788-791.

Levin MF, Weiss PL, Keshner EA (2015) Emergence of virtual reality as a tool for upper limb

rehabilitation: incorporation of motor control and motor learning principles. Phys Ther

95:415-425.

Loeb GE (1999) Asymmetry of hindlimb muscle activity and cutaneous reflexes after tendon

transfers in kittens. J Neurophysiol 82:3392-3405.

Luauté J, Schwartz S, Rossetti Y, Spiridon M, Rode G, Boisson D, Vuilleumier P (2009)

Dynamic changes in brain activity during prism adaptation. J Neurosci 29:169-178.

Maas H, Huijing PA (2012) Mechanical effect of rat flexor carpi ulnaris muscle after tendon

transfer: does it generate a wrist extension moment? Journal of Applied Physiology

112:607-614.

Maier M, Ballester BR, Verschure PFMJ (2019) Principles of Neurorehabilitation After Stroke

Based on Motor Learning and Brain Plasticity Mechanisms. Frontiers in Systems

Neuroscience Volume 13 - 2019.

Makin TR, Krakauer JW (2023) Against cortical reorganisation. eLife 12:e84716.

Martin TA, Keating JG, Goodkin HP, Bastian AJ, Thach WT (1996) Throwing while looking

through prisms. II. Specificity and storage of multiple gaze-throw calibrations. Brain 119

(Pt 4):1199-1211.

Mathis A, Mamidanna P, Cury KM, Abe T, Murthy VN, Mathis MW, Bethge M (2018)

DeepLabCut: markerless pose estimation of user-defined body parts with deep learning.

Nature Neuroscience 21:1281-1289.

Mercuri E, Muntoni F (2013) Muscular dystrophies. Lancet 381:845-860.

Morton SM, Bastian AJ (2006) Cerebellar contributions to locomotor adaptations during

splitbelt treadmill walking. J Neurosci 26:9107-9116.

Murata Y, Higo N, Oishi T, Yamashita A, Matsuda K, Hayashi M, Yamane S (2008) Effects of

motor training on the recovery of manual dexterity after primary motor cortex lesion in

macaque monkeys. J Neurophysiol 99:773-786.

Mussa-Ivaldi FA, Bizzi E (1995) The Modular Organization of Motor Control: What Frogs Can

Teach Us About Adaptive Learning. IFAC Proceedings Volumes 28:413-418.

Nakajima N, Wang S, Ogihara N, Oya T, Seki K, Funato T (2023) Upper Limb Musculoskeletal

Model of Macaque Monkey for Approaching Adaptation Mechanism to Tendon Transfer.

Neuroscience 2023 Abstracts, Washington, DC: Society for Neuroscience.

Nath T, Mathis A, Chen AC, Patel A, Bethge M, Mathis MW (2019) Using DeepLabCut for 3D

markerless pose estimation across species and behaviors. Nature Protocols 14:2152-

2176.

Nordez A, Gallot T, Catheline S, Guével A, Cornu C, Hug F (2009) Electromechanical delay
revisited using very high frame rate ultrasound. J Appl Physiol (1985) 106:1970-1975.

Nudo RJ, Wise BM, SiFuentes F, Milliken GW (1996) Neural substrates for the effects of
rehabilitative training on motor recovery after ischemic infarct. Science 272:1791-1794.

Osu R, Franklin DW, Kato H, Gomi H, Domen K, Yoshioka T, Kawato M (2002) Short- and
long-term changes in joint co-contraction associated with motor learning as revealed
from surface EMG. J Neurophysiol 88:991-1004.

Overduin SA, d'Avella A, Roh J, Bizzi E (2008) Modulation of Muscle Synergy Recruitment in
Primate Grasping. The Journal of Neuroscience 28:880-892.

Petitet P, O'Reilly JX, O'Shea J (2018) Towards a neuro-computational account of prism
adaptation. Neuropsychologia 115:188-203.

Popa LS, Streng ML, Hewitt AL, Ebner TJ (2016) The Errors of Our Ways: Understanding Error
Representations in Cerebellar-Dependent Motor Learning. Cerebellum 15:93-103.

Power JD, Schlaggar BL (2017) Neural plasticity across the lifespan. Wiley Interdiscip Rev Dev
Biol 6.

Ranganathan R, Adewuyi A, Mussa-Ivaldi FA (2013) Learning to be lazy: exploiting redundancy
in a novel task to minimize movement-related effort. J Neurosci 33:2754-2760.

Sattin D, Parma C, Lunetta C, Zulueta A, Lanzone J, Giani L, Vassallo M, Picozzi M, Parati EA
(2023) An Overview of the Body Schema and Body Image: Theoretical Models,
Methodological Settings and Pitfalls for Rehabilitation of Persons with Neurological
Disorders. Brain Sci 13.

Schärli A, Hecht H, Mast FW, Hossner EJ (2024) How spotting technique affects dizziness and
postural stability after full-body rotations in dancers. Hum Mov Sci 95:103211.

Schoultz TW, Swett JE (1972) The fine structure of the Golgi tendon organ. J Neurocytol 1:1-26.

Slawinska U, Kasicki S (2002) Altered Electromyographic Activity Pattern of Rat Soleus Muscle
Transposed into the Bed of Antagonist Muscle. The Journal of Neuroscience 22:5808-
5812.

Smith MA, Shadmehr R (2005) Intact ability to learn internal models of arm dynamics in
Huntington's disease but not cerebellar degeneration. J Neurophysiol 93:2809-2821.

Smith MA, Ghazizadeh A, Shadmehr R (2006a) Interacting Adaptive Processes with Different
Timescales Underlie Short-Term Motor Learning. PLOS Biology 4:e179.

Smith MA, Ghazizadeh A, Shadmehr R (2006b) Interacting adaptive processes with different
timescales underlie short-term motor learning. PLoS Biol 4:e179.

Smith SA, Mitchell JH, Garry MG (2006c) The mammalian exercise pressor reflex in health and
disease. Exp Physiol 91:89-102.

Sperry RW (1940) The functional results of muscle transposition in the hind limb of the rat.
1240   Journal of Comparative Neurology 73:379-404.

1241   Sperry RW (1942) Transplantation of motor nerves and muscles in the forelimb of the rat.
1242   Journal of Comparative Neurology 76.

1243   Sperry RW (1947) Effect of crossing nerves to antagonistic limb muscles in the monkey. Arch
1244   Neurol Psychiatry 58:452-473.

1245   Sugita Y (1996) Global plasticity in adult visual cortex following reversal of visual input. Nature
1246   380:523-526.

1247   Takei T, Seki K (2010) Spinal interneurons facilitate coactivation of hand muscles during a
1248   precision grip task in monkeys. J Neurosci 30:17041-17050.

1249   Takei T, Confais J, Tomatsu S, Oya T, Seki K (2017) Neural basis for hand muscle synergies in
1250   the primate spinal cord. Proceedings of the National Academy of Sciences 114:8643-
1251   8648.

1252   Takeuchi N, Izumi S (2012) Maladaptive plasticity for motor recovery after stroke: mechanisms
1253   and approaches. Neural Plast 2012:359728.

1254   Tannenbaum J, Bennett BT (2015) Russell and Burch's 3Rs then and now: the need for clarity in
1255   definition and purpose. J Am Assoc Lab Anim Sci 54:120-132.

1256   Thompson AK, Wolpaw JR (2014) Operant conditioning of spinal reflexes: from basic science to
1257   clinical therapy. Front Integr Neurosci 8:25.

1258   Thoroughman KA, Shadmehr R (1999) Electromyographic correlates of learning an internal
1259   model of reaching movements. J Neurosci 19:8573-8588.

1260   Trewartha KM, Garcia A, Wolpert DM, Flanagan JR (2014) Fast but fleeting: adaptive motor
1261   learning processes associated with aging and cognitive decline. J Neurosci 34:13411-
1262   13421.

1263   Valero-Cuevas FJ, Hentz VR (2002) Releasing the A3 pulley and leaving flexor superficialis intact
1264   increases pinch force following the Zancolli lasso procedures to prevent claw deformity
1265   in the intrinsic palsied finger. J Orthop Res 20:902-909.

1266   Vallbo AB, Hagbarth KE, Torebjörk HE, Wallin BG (1979) Somatosensory, proprioceptive, and
1267   sympathetic activity in human peripheral nerves. Physiol Rev 59:919-957.

1268   van Vliet P, Carey LM, Turton A, Kwakkel G, Palazzi K, Oldmeadow C, Searles A, Lavis H,
1269   Middleton S, Galloway M, Dimech-Betancourt B, O'Keefe S, Tavener M (2023) Task-
1270   specific training versus usual care to improve upper limb function after stroke: the
1271   "Task-AT Home" randomised controlled trial protocol. Frontiers in Neurology 14.

1272   Vanhoof MJM, van Leeuwen T, Galletta L, Vereecke EE (2021) The forearm and hand
1273   musculature of semi-terrestrial rhesus macaques (Macaca mulatta) and arboreal gibbons
1274   (fam.Hylobatidae). Part II. Quantitative analysis. Journal of Anatomy 238:321-337.

1275   Villalta JI, Landi SM, Fló A, Della-Maggiore V (2013) Extinction Interferes with the Retrieval of
1276   Visuomotor Memories Through a Mechanism Involving the Sensorimotor Cortex.
1277   Cerebral Cortex 25:1535-1543.
1278   Walker JG, Jackson HJ, Littlejohn GO (2004) Models of adjustment to chronic illness: using the
1279   example of rheumatoid arthritis. Clin Psychol Rev 24:461-488.
1280   Wester K, Hove LM, Barndon R, Craven AR, Hugdahl K (2018) Cortical Plasticity After Surgical
1281   Tendon Transfer in Tetraplegics. Front Hum Neurosci 12:234.
1282   Winterbottom L, Nilsen DM (2024) Motor Learning Following Stroke: Mechanisms of Learning
1283   and Techniques to Augment Neuroplasticity. Phys Med Rehabil Clin N Am 35:277-291.
1284   Wolpert DM (1997) Computational approaches to motor control. Trends Cogn Sci 1:209-216.
1285   Yan Y, Sobinov AR, Bensmaia SJ (2022) Prehension kinematics in humans and macaques. J
1286   Neurophysiol 127:1669-1678.
1287   Yumiya H, Larsen KD, Asanuma H (1979) Motor readjustment and input-output relationship of
1288   motor cortex following cross-connection of forearm muscles in cats. Brain Research
1289   177:566-570.
1290   Zajac FE (1992) How musculotendon architecture and joint geometry affect the capacity of
1291   muscles to move and exert force on objects: a review with application to arm and forearm
1292   tendon transfer design. J Hand Surg Am 17:799-804.
1293   Zeiler SR, Krakauer JW (2013) The interaction between training and plasticity in the poststroke
1294   brain. Curr Opin Neurol 26:609-616.
1295   Zemková E (2022) Physiological Mechanisms of Exercise and Its Effects on Postural Sway: Does
1296   Sport Make a Difference? Front Physiol 13:792875.

1297

1298

1299

Figure 1: Muscle anatomy of the macaque forearm and the tendon transfer procedure.

A schematic of the primary forearm muscles involved in the study, showing both the dorsal and volar views. The diagram illustrates the surgical crossed tendon transfer of the extensor digitorum communis (EDC) and flexor digitorum superficialis (FDS) tendons. All labeled muscles were implanted with EMG electrodes. Muscle Abbreviations: BRD: brachioradialis, ECR: extensor carpi radialis, ECU: extensor carpi ulnaris, ED2,3: extensor digitorum-2,3, ED4,5: extensor digitorum-4,5, EPL: extensor pollicis longus (not implanted), FCR: flexor carpi radialis, FCU: flexor carpi ulnaris, FDP: flexor digitorum profundus, PL: palmaris longus, PT: pronator teres (not implanted), *(The deltoid (DEL) muscle was also implanted in Monkey B but is not shown as it is a shoulder muscle). See also Supplementary Table 1 for a complete list of all muscle abbreviations, their full names, and their assigned synergies.*

1317

Figure 2: *Long term confirmation of tendon surgery effectiveness to alter mechanical properties.*

(A) Set-up for the ultrasound measurement and video recordings of the stimulation induced movements of the EDC and FDS tendons. (B) Sonogram of the FDS muscle and its intramuscular tendons. Left side (B-mode, i.e., brightness mode) shows the still image of the monkeys forearm at a given point in time. Right side (M-mode, i.e., motion mode) shows the stagged images of the FDS tendon displacement induced by muscle stimulation (50mA). White arrows demarcate the FDS tendon which was used for the measurement. Grayscale gradations correspond to tissue densities: hyperechoic regions (white) denote denser structures like surface of bones and tendons, while hypoechoic areas (black) signify less dense tissues such as adipose tissue and musculature. Inset demonstrates the area measurement. The area of the displacement waves was measured in the M-mode, representing the strength of muscle contraction. We measured the duration (a, sec) and amplitude (b, cm) of three waves and calculated the average. Area = a*b/2(cm/sec) for days 0, 7 and 105 after tendon transfer. (C) Areas under the wave measured in the M-mode for 3 experimental days (0, 7 and 105 days post-TT) and regression lines in red and blue for FDS

and EDC, respectively. $R^2>0.5$ for FDS. The data suggested that muscle contractions induced by direct electrical stimulation were nearly constant. (D) Markers placed on the index, middle and ring finger nails (A) were used to measure finger displacement in xyz-dimensions. We calculated the sum of the Euclidean distances of each marker from the origin of the 3D coordinate system as a scalar quantity. Observing the movement along the Z-axis, it became reversed post-surgery indicating a reversal from finger flexion to extension due to tendon transfer (D, blue = pre-TT at surgery day; dark brown = post-TT at surgery day; light brown = 1wk post-TT; red = 3wks post-TT). The scalar quantity of the fingers during muscle stimulation did not change much at day 0, 7 and 105 days (E), suggesting that there was no tendon rupture or slackening of the tendons postoperatively (EDC stimulation, left; FDS stimulation, right). Data were collected in monkey A.

Figure 3: *Experimental set-up, task sequence and typical EMG (monkey A).*

(A) Schematic of the task object using a rod requiring monkey A to perform a controlled grasp. (B) Schematic of the behavioral sequence (hook → grasp → release). (C) Typical EMG traces of a control session (high-pass filtered) for all recorded muscles. Gray boxes represent the task sequence. Obj 1 ON: start of the hold period of object 1. Obj 1 OFF: end of the hold period of object 1, i.e., object release. Obj 2 ON: start of the hold period of object 2. Obj 2 OFF: end of

the hold period of object 2, i.e., object release. Tendons of the muscles marked with * were cross-transferred. (D) Rectified and smoothed EMG for all recorded muscles (average for one recording session; amplitude [µV] over task sequence [%]). Horizontal bars illustrate the corresponding behavioral periods; red vertical lines indicate peak amplitude for each muscle). (E) The time the monkey spent on the left side of the yellow dotted line while moving from object 1 to object 2 was measured and used to quantify the maladaptive behavior.

1371

Figure 4: Experimental set-up, task sequence and typical EMG (monkey B).

(A) Schematic of the task requiring monkey B to pick up food from a groove allowing for a more natural grasp. (B) Schematic of the task sequence (picking up food). (C) Typical EMG traces of a control session (high-pass filtered) for all recorded muscles. Gray boxes represent the task. Obj 1 ON: start of the hold period of object 1. Obj 1 OFF: end of object 1's hold period, i.e. object release. LED ON: approximate start of the food touch. LED OFF: approximate time of food retrieval. Tendons of the muscles marked with * were cross-transferred. (D)

rectified and smoothed EMG for all recorded muscles (average for one recording session; amplitude [µV] over task sequence [%]). Horizontal bars illustrate the corresponding behavioral periods, red vertical lines indicate peak amplitude for each muscle). (E) Example for maladaptive behavior in monkey B. The time the monkey spent in contact with or behind the object plate was measured and used to quantify the maladaptive behavior.

1387

Figure 5: *Behavioral and kinematic metrics of motor recovery.*

(A, D) Grip formation times (mean ± SD; n = 20 trials) for Monkey A (A) and Monkey B (D). (B, E) Duration of off-target reaching movements (mean ± SD; n = 10 trials) for Monkey A (B) and Monkey B (E). (C) Pull time duration for Monkey A. (F) Grasp aperture size for Monkey B. Filled squares indicate significant difference from pre-TT baseline (p < 0.05, two-sample t-test). All data are plotted over days relative to tendon transfer (TT).

1397

1398

1399 **Figure 6: Temporal EMG profiles and cross-correlation analysis.**

1400 Temporal EMG profiles for Monkey A (left, A-J) and Monkey B (right, K-Q). (A-F,

1401 K-O) Average EMG activity profiles aligned to task events (0%; object release for

1402 A, food touch for B). Shaded envelopes represent standard deviations. Triangles

1403 indicate peak activity during extension (▼) or flexion (▽). Colored traces denote

post-surgery landmark days. (A-D, K-M) Comparison of transferred muscles. Note the temporal shift in the post-surgery profile (B, L) relative to the pre-TT baseline (dashed lines). (E-F, N-O) Profiles of non-transferred muscles. (G-J, P-Q) Zero-lag cross-correlation coefficients between post-surgery EMG profiles and the pre-TT baseline profiles plotted over time. (H, I, P) Correlation coefficients calculated against the muscle's *own* pre-TT baseline. (G, J, Q) Correlation coefficients calculated against the *antagonist's* pre-TT baseline (e.g., Post-EDC vs. Pre-FDS). Black dashed lines on the right y-axis indicate behavioral error metrics (Off-target reaching time for Monkey A; Contact duration for Monkey B; gray shading represents SD). The // represents the recovery period.

1414

1415

Figure 7: Spatial structure and temporal activation of Primary Synergies.

Analysis of the two primary synergies: Synergy A (Flexor) and Synergy B (Extensor). (A, B, E, F) Spatial synergy weights (*W*) showing the contribution of each muscle to the synergy. Bar plots represent the average weights across all recording days. (C, D, G, H) Temporal activation profiles (*C*) aligned to task events (0%). Dashed lines with shaded tubes indicate the average pre-TT EMG profiles of the key contributing muscles (FDS, EDC, FDP) for visual comparison with the synergy profile. Symbols and alignment are as described in Figure 6. (I,

1424 J) Quantification of spatial stability. Cosine similarity of spatial synergy weights
1425 (W) calculated between individual recording days and the pre-TT average. Blue
1426 markers indicate pre-TT control days; Red markers indicate post-TT days. The
1427 horizontal gray shaded region (0.95-1.0) denotes the range of high baseline
1428 stability.

Figure 8: Analysis of Secondary Muscle Synergies.

(A-H) Analysis of Secondary Synergies C (Wrist Flexor) and D (Wrist Extensor).
(A, B, E, F) Spatial synergy weights (*W*) showing the contribution of each muscle.
(C, D, G, H) Temporal activation profiles (*C*) aligned to task events (0%). Colored
traces denote post-surgery landmark days. Layout and symbols are as described
in Figure 6. (I, J) Quantification of spatial stability for Synergies C and D. Cosine
similarity of spatial weights calculated between individual recording days and the
pre-TT average. Blue markers indicate pre-TT control days; Red markers indicate

1446   post-TT days. The horizontal gray shaded region (0.95-1.0) denotes the range of
1447   high baseline stability.
1448

Monkey A

A ── Synergy A ──
B ── Synergy B ──
C
D

Monkey B

E ── Synergy A ──
F ── Synergy B ──
G
H

vs Synergy B (pre-surgery)

vs Synergy A (pre-surgery)

Post surgery days

1449

Figure 9: Cross-correlation analysis of Primary Synergy activation.

Zero-lag cross-correlation coefficients plotted over post-surgery days for Monkey A (A-D) and Monkey B (E-H). Activation patterns of the Primary Flexor (Synergy A) and Primary Extensor (Synergy B) were cross-correlated with pre-TT baseline profiles. (Top Row: A, B, E, F) Correlations calculated against the pre-TT *Extensor* Synergy (Synergy B, blue traces). (Bottom Row: C, D, G, H) Correlations calculated against the pre-TT *Flexor* Synergy (Synergy A, red traces). (C, G) Correlation of Synergy A with its own pre-TT baseline. (B, F) Correlation of Synergy B with its own pre-TT baseline. (A, E, D, H) Cross-correlations between antagonistic synergies (e.g., A is Synergy A vs. Pre-Synergy B). Black dashed lines on the right y-axis indicate behavioral error metrics (gray shading represents SD). The // represents the recovery period. Triangles indicate landmark days.

1461

Figure 10: Cross-correlation analysis of Secondary Synergy activation.

Zero-lag cross-correlation coefficients for the secondary synergies (C and D) plotted over post-surgery days for Monkey A (A-D) and Monkey B (E-H). Activation patterns were cross-correlated with the pre-TT profiles of the Primary Synergies to assess changing affiliations. (Top Row: A, B, E, F) Correlations calculated against the pre-TT *Extensor* Synergy (Synergy B, blue traces). (Bottom Row: C, D, G, H) Correlations calculated against the pre-TT *Flexor* Synergy (Synergy A, red traces). (Left Columns: A, C, E, G) Synergy C correlations. (Right Columns: B, D, F, H) Synergy D correlations. Black dashed lines on the right y-axis indicate behavioral error metrics.

1476

1477

Figure 11: Aggregated and averaged EMG (aaEMG).

Aggregated and averaged electromyography (EMG) activities for the main
contributing muscles of each synergy for Monkey A (A-H) and Monkey B (I-P).
(A-D, I-L) Time course of aaEMG activity (summed within ±15% task range)
plotted over post-surgery days. Black dashed lines on the right y-axis indicate
behavioral error metrics. (E-H, M-P) Bar plots showing the mean (± SEM) aaEMG
for the pre-TT period ("pre") and the five selected landmark days. Vertical colored
bars on the time-series plots indicate the corresponding landmark days. Asterisks
indicate significant difference from the pre-TT control period (*p < 0.01, **p <
0.001, ***p < 0.0001; two-sample t-test with Bonferroni correction).

1488

1489

1490

Figure 12:

Kinematic analysis of joint angles (Monkey B).

Changes in joint angles (mean of 20 trials ± SD) for each landmark day. (A)
Metacarpophalangeal (MCP) joint angle. (B) Wrist joint angle. Asterisks indicate
significant difference from pre-TT baseline (**$p < 0.001$, ***$p < 0.0001$; ANOVA).
(C) Schematic indicating the timing of the kinematic snapshot relative to the task
timeline (dotted line; 83 ms before food touch), capturing the hand configuration
during the pre-shaping phase.

1499

1500

1501

Figure 13: Kinematic analysis reveals gradual refinement of the compensatory tenodesis strategy over time in Monkey B.

(A) Each subplot shows the trial-by-trial relationship between wrist angle (X-axis) and MCP angle (Y-axis) for a single recording day (n=20 trials per day). Points are color-coded based on the day relative to surgery (colorbar). Pre-TT (Day -4), no correlation exists (R^2=0.00). Post-surgery, a negative correlation emerges and strengthens over time, peaking around Day 56 (R^2=0.58), indicating the learned exploitation of the tenodesis effect where wrist flexion predicts finger extension. The tightening of the scatter plots and increase in R^2 over weeks provide direct evidence for a gradual motor skill learning process. (B) All data points combined.

1513

Figure 14: A Proposed Model of Multi-Timescale Adaptation Following Tendon Transfer.

This schematic illustrates the hypothesized interaction between fast and slow adaptive processes driving recovery. The initial Tendon Transfer triggers a rapid but maladaptive 'swap' of motor commands (Fast Adaptation 1), leading to a maladaptive state. During this phase, two slower processes are hypothesized to occur in parallel: a costly 'arms race' within the conflicted synergy (Slow Process A, red curve) and the gradual development of a functional compensatory strategy (Slow Process B, green curve). When the 'arms race' reaches a threshold of unsustainable cost (dashed blue line), a second Fast Adaptation ('Switch-Back', 2) is triggered. This allows for the abandonment of the flawed strategy and the adoption of a stable, 'good enough' solution, which is now supported by the newly learned compensatory strategy. The gray line represents the observed neural data (e.g., cross-correlation coefficients of electromyography and temporal activation profiles of muscle synergies), which reflects this two-phase process.

1529

1530