Crossover in Aromatic Amino Acid Interaction Strength: Tyrosine vs. Phenylalanine in Biomolecular Condensates

David De Sancho[1*] **and Xabier López**[1*]

*For correspondence:
david.desancho@ehu.eus (DS);
xabier.lopez@ehu.eus (XL)

[1]Polimero eta Material Aurreratuak: Fisika, Kimika eta Teknologia, Kimika Fakultatea, UPV/EHU & Donostia International Physics Center (DIPC), PK 1072, 20018 Donostia-San Sebastian, Euskadi, Spain

Abstract   Biomolecular condensates often form through the self-assembly of disordered proteins with low-complexity sequences. In these polypeptides, the aromatic amino acids phenylalanine and tyrosine act as key "sticker" residues, driving the cohesion of dense phases. Recent studies on condensates suggest a hierarchy in sticker strength, with tyrosine being more adhesive than phenylalanine. This hierarchy aligns with experimental data on amino acid solubilities and potentials of mean force derived from atomistic simulations. However, it contradicts conventional chemical intuition based on hydrophobicity scales and pairwise contact statistics from folded protein structures, which suggest phenylalanine should be the stronger sticker. In this work, we use molecular dynamics simulations and quantum chemistry calculations to resolve this apparent discrepancy. Using simple model peptides and side-chain analogues, we demonstrate that the experimentally observed hierarchy arises both from the possibility of tyrosine forming hydrogen bonds and the lower free energy of transfer of tyrosine into the condensate. The high level of hydration of polypeptides in dense phases drives this effect. Notably, as the dielectric constant of the surrounding environment approaches that of an apolar solvent, the trend reverses, with phenylalanine becoming the stronger sticker. These findings highlight the role of hydration in modulating aromatic interactions and provide a clear explanation for the crossover in sticker strength between tyrosine and phenylalanine in different media.

Introduction

For decades, cellular organization was primarily understood through the lens of membrane-bound compartmentalization. However, it has become increasingly clear that membrane-less organelles –such as stress granules, nuclear speckles, and Cajal bodies– are also widespread within cells (*Banani et al., 2017*). At least in some cases, these organelles form by phase separation of their components (*Shin and Brangwynne, 2017*), primarily disordered proteins and nucleic acids, which partition into dense and dilute phases. The condensates thus formed retain properties of liquids, like fusion, dripping and wetting (*Brangwynne et al., 2009*). This phenomenon has generated significant interest in the physical mechanisms governing the phase behavior of biomolecular mixtures (*Choi et al., 2020*).

Many proteins undergoing phase separation share common characteristics, including devia-

tions from typical compositions of folding proteins and simple sequences with multiple amino acid repeats (*Martin and Mittag, 2018*). Specifically, in the sequences of proteins that experience upper critical solution temperature (UCST) transitions, like the low-complexity regions of FUS, hnRNPA1, Ddx4, LAF-1 or atGRP7, we observe long stretches rich in polar residues interspersed with aromatics or positively charged residues. Borrowing language from the theory of associative polymers (*Semenov and Rubinstein, 1998*), these units of sequence have been termed "stickers", in the case of aromatics and positively charged residues, and "spacers", for the polar residue repeats (*Martin et al., 2020*; *Bremer et al., 2022*; *Mittag and Pappu, 2022*). Spacers act as linkers that lend flexibility to the polypeptide mesh in the protein-dense phase. In contrast, stickers play a crucial role in determining both the single-chain properties of the polymer and the phase behaviour of the condensate through interactions involving their aromatic or charged groups (*Wang et al., 2018*). Recent experiments have quantified the influence of different types of stickers in the polymer properties and phase behaviour (*Bremer et al., 2022*). This raises a crucial question: what is the fundamental origin of the relative strengths of different stickers?

This question has recently been addressed for the cationic amino acids lysine (Lys) and arginine (Arg) (*Das et al., 2020*; *Schuster et al., 2020*; *Fisher and Elbaum-Garfinkle, 2020*; *Greig et al., 2020*; *Paloni et al., 2021*). Despite having the same net charge, these residues turn out not to be interchangeable. Mutagenesis experiments on LAF-1 have shown that substituting arginine with lysine completely suppresses phase separation (*Schuster et al., 2020*). This distinction is functionally relevant, as Arg to Lys substitutions affect speckle formation (*Greig et al., 2020*). Additionally, experiments on the intrinsically disordered region (IDR) of Ddx4 indicate that LLPS is favoured by Arg relative to Lys (*Brady et al., 2017*; *Das et al., 2020*; *Schuster et al., 2020*). Coarse-grained simulation models have failed to account for the greater propensity of arginine to promote phase separation in Ddx4 variants with Arg to Lys mutations (*Das et al., 2020*). These models are often parametrized using hydrophobicity scales that capture the "stickiness" of different amino acid residues (*Dignon et al., 2018b,a*; *Regy et al., 2021*; *Tesei et al., 2021*). However, by replacing the hydrophobicity scales in the coarse-grained model with interaction energies from amino acid contact matrices –derived from statistical analysis of the PDB– the correct trends were captured (*Das et al., 2020*). A key to the greater propensity for LLPS in the case of Arg may derive from the pseudo-aromaticity of this residue (*Gobbi and Frenking, 1993*; *Wang et al., 2018*; *Hong et al., 2022*).

Here we focus on the distinct roles of the main aromatic residues acting as stickers –tyrosine (Tyr) and phenylalanine (Phe)– excluding tryptophan due to its much lower abundance (*Maraldo et al., 2024*). Given that they only differ in a hydroxyl group, one could expect Tyr and Phe to be equally relevant to phase separation, especially considering the finding from a double-mutant cycle that Tyr-Tyr and Phe-Phe pairs make nearly identical contributions to protein stability (*Serrano et al., 1991*). However, experimental evidence on various proteins suggests that Tyr is a stronger driver of condensation (*Lin et al., 2017*; *Wang et al., 2018*; *Schuster et al., 2020*; *Bremer et al., 2022*). This is demonstrated by the reduced propensity to phase separate of Tyr-to-Phe mutants of FUS and LAF-1 (*Lin et al., 2017*; *Wang et al., 2018*; *Schuster et al., 2020*) and the opposite effect in Phe-to-Tyr mutants in hnRNPA1 (*Wang et al., 2018*; *Bremer et al., 2022*). Understanding the origin of the different sticker strengths of Phe and Tyr is important due to its functional relevance. In a large sample of hnRNPA1 variants, the fractions of Phe and Tyr have been found to co-vary, suggesting that evolutionary control of composition fine-tunes the properties of condensates (*Bremer et al., 2022*).

The properties of Phe and Tyr may give important insights on the distinct behaviour as stickers in biomolecular condensates. Hydrophobicity scales typically rank Phe as the most hydrophobic residue (*Kyte and Doolittle, 1982*; *Tesei et al., 2021*), consistent with the greater hydration free energy of Tyr relative to Phe (*Wolfenden et al., 1981*; *Chang et al., 2007*) (see Figure 1A-B). As in the case of Arg and Lys, using hydrophobicity as a proxy for interaction energy in bead simulation models proved insufficient to explain the relative strengths of Phe and Tyr as stickers (*Das et al., 2020*). However, in this case statistical contact matrices also could not capture the correct order

Figure 1. Bibliographic values for different properties of phenylalanine and tyrosine. (A) Solvation free energy (ΔG_{solv}) (**_Chang et al., 2007_**). (B) Probability distributions of min-maxed normalized hydropathy values λ from bibliographic hydrophobicity scales (**_Tesei et al., 2021_**). (C) Self-interaction energy (ε_{ii}) from the Miyazawa-Jernigan contact matrix (**_Miyazawa and Jernigan, 1996_**). (D) Solubility in water at 25°C (**_Nozaki and Tanford, 1971_**).

---

of stickiness. In the Miyazawa and Jernigan statistical potential, Tyr-Tyr contacts are weaker than Phe-Phe (with energies of -4.17 and -7.26 in RT units, respectively; see Figure 1C) (**_Miyazawa and Jernigan, 1996_**). We note that a different hydrophobicity scale based on peptides undergoing inverse temperature transitions (the Urry scale (**_Urry et al., 1992_**), where Tyr is more hydrophobic) can account for the correct rank order in saturation concentration (**_Regy et al., 2021_**). On the other hand, the potentials of mean force between amino acids calculated with atomistic force fields have a deeper free energy well for the Tyr-Tyr pair than for Phe-Phe (**_Chelli et al., 2002_**; **_Joseph et al., 2021_**). A final data point of interest is the extremely low solubility of Tyr, over an order of magnitude smaller than that of Phe (0.045 and 2.79 g/100 g of water, respectively; see Figure 1D) (**_Nozaki and Tanford, 1971_**). This has led to the suggestion that tyrosine-tyrosine hydrogen bonds are stronger than those formed by tyrosine and water (**_Pace et al., 2001_**).

In summary, experimental results on condensates containing Phe/Tyr variants do not seem to align with either solvation free energies, most hydrophobicity scales or statistical contact potentials, but are consistent with atomistic force fields in solution and solubilities in water. One possible explanation for these conflicting findings is that, due to their level of hydration, molecular condensates may differ significantly from the tightly-packed cores of folded protein structures (**_Lin et al., 2017_**; **_Das et al., 2020_**). Here we address this paradox using a combination of classical molecular dynamics (MD) simulations and quantum chemical calculations. First, we estimate transfer free energies of peptides including these aromatic residues into model peptide condensates and find that they are more favourable for Tyr than for Phe, an effect that is reversed when we perform the same transformation in apolar media. DFT calculations confirm that the interaction energies in Tyr-Tyr pairs are stronger than those between Phe residues. However, the transfer free energy contribution dominates at sufficiently low dielectric constants, making Phe-Phe pairs more favourable. These findings recapitulate the right rank order of interaction strengths of aromatic stickers in biomolecular condensates but also their crossover in low dielectric media like the

116 hydrophobic cores of folded proteins.

Materials and Methods

Classical molecular dynamics simulations

Molecular models

In this work, we report simulations of terminally-capped GGXGG peptides with X=F/Y in different
media, including water, organic solvents and molecular condensates. This peptide family has
been characterized extensively in the past, both from experiment (*Plaxco et al., 1997*) and sim-
ulation(*Workman and Pettitt, 2021*). We have built structures for the peptides using Ambertools
(*Salomon-Ferrer et al., 2013*) and used the Amber 99sb★-ILDN force field, including corrections
for backbone torsions (*Lindorff-Larsen et al., 2010*), and TIP3P (*Jorgensen et al., 1983*) water. Our
condensates are formed by an equimolar mixture of Gly, Ser and either Tyr or Phe "dipeptides" (in
fact, terminally capped amino acids), as before (*De Sancho, 2022*). For the simulations with differ-
ent organic solvents, we used the GAFF parameters derived by Mobley and co-workers (*Bannan*
et al., 2016).

Thermodynamic cycle

To estimate differences in the transfer free energy ($\Delta\Delta G_{\text{Transfer}}$) of our short peptides into a differ-
ent medium –either a condensate or a different solvent– we use an alchemical (i.e. non-physical)
transformation (*Mey et al., 2020*). Specifically, the thermodynamic cycle we construct involves four
different states for the GGXGG peptide, with X=F ($\lambda = 0$) or X=Y ($\lambda = 1$), which is inserted either in
water or in the alternative medium. We illustrate this thermodynamic cycle in the case of transfer
into a condensate in Figure 2A. The vertical branches of the thermodynamic cycle are associated
to the free energy of transfer of the model peptides from water into the condensate ($\Delta G_{\text{Transfer}}^{\text{F}}$ and
$\Delta G_{\text{Transfer}}^{\text{Y}}$), while the horizontal branches involve an alchemical transformation ($\Delta G_{\text{F}\rightarrow\text{Y}}(\text{H}_2\text{O})$ and
$\Delta G_{\text{F}\rightarrow\text{Y}}(\text{Cond})$). Using simulations, we can easily calculate the free energies corresponding to the
horizontal branches. This enables us to estimate the desired quantity as

$$\Delta\Delta G_{\text{Transfer}} = \Delta G_{\text{Transfer}}^{\text{Y}} - \Delta G_{\text{Transfer}}^{\text{F}} = \Delta G_{\text{F}\rightarrow\text{Y}}(\text{Cond}) - \Delta G_{\text{F}\rightarrow\text{Y}}(\text{H}_2\text{O}) \tag{1}$$

Negative $\Delta\Delta G_{\text{Transfer}}$ values indicate that transferring the Tyr peptide into the condensate is more
favourable than transferring the Phe peptide, whereas positive values indicate the opposite.

Simulation setup

For the simulation runs in pure solvents, we simply inserted the peptide in an octahedral box leav-
ing at least 1 nm of distance to each wall. Then the boxes were filled with the corresponding solvent
molecules. For the simulations in the model condensate, we first inserted the peptide at the central
position of a rectangular box with dimensions 8.5 nm×4.5 nm×4.5 nm, containing an equimolar
mixture of Gly, Ser and either Tyr or Phe dipeptides, which we term GSY and GSF, respectively.
Then, the box was expanded up to 20 nm and filled with water molecules. This results in a simula-
tion box where the peptide of interest is immersed in a protein-dense slab in contact with a dilute
phase.

The equilibration protocol is identical in all cases. First, we ran an energy minimization using a
steepest descent algorithm, followed by short simulations in the NVT and NPT ensembles, using
the Berendsen (*Berendsen et al., 1984*) and velocity rescaling (*Bussi et al., 2007*) thermostats, re-
spectively, to set the temperature at 298 K, and the Berendsen barostat (*Berendsen et al., 1984*) to
set the pressure at 1 bar on the latter. Finally, we performed equilibrium simulations of in the NPT
ensemble using a leap-frog stochastic dynamics integrator with a time-step of 2 fs at 298 K and
setting the pressure to 1 bar with the Parrinello-Rahman barostat (*Parrinello and Rahman, 1980*).
The duration of the equilibrium runs was 1-2 μs for the condensate simulations and 500 ns for
all other solvents. Note that for each system-solvent combination, two different sets of simulations

Figure 2. (a) Thermodynamic cycle used in this study for the estimation of free energy differences upon mutation for the insertion of a peptide in a molecular condensate. (B) Schematic of the process of contact formation between two molecules i and j used in the quantum chemical calculations. We consider that contact formation involves the transfer from water (blue) to a different medium (orange) and the interaction in this medium between the entities involved.

had to be performed, one for each of the alchemical (λ) states corresponding to the central peptide residue being Phe ($\lambda = 0$) or Tyr ($\lambda = 1$). For the alchemical transitions, we use a non-equilibrium switching method (*Seeliger and de Groot, 2010*; *Gapsys et al., 2015*; *Aldeghi et al., 2019*) often used for the calculation of differences in binding affinities (*Aldeghi et al., 2018*; *Gapsys et al., 2020*) or free energy changes upon mutation (*Gapsys et al., 2016*; *Martinez-Martin et al., 2023*). We run short (50 ps) simulation trajectories where the λ-state was switched forward ($\lambda = 0 \rightarrow 1$) or backward ($\lambda = 1 \rightarrow 0$). We have run additional simulations of the solvents in the absence of the solute using the same protocols. To study interactions in the biomolecular condensates, we run simulations of equimolar GSY and GSF mixtures in a slab configuration, which we later sliced to keep only the dense phase. This was then briefly re-equilibrated as described above. Production simulations of the condensates were run for 1 μs.

For all simulations, we have used the Gromacs software package (*Abraham et al., 2015*) (v.2024). To generate hybrid topologies for the different λ-states, we have used the PMX software (*Seeliger and de Groot, 2010*; *Gapsys et al., 2015*; *Aldeghi et al., 2019*) (v. 3.0).

Analysis

We have analyzed the non-equilibrium switching simulations using the PMX package tools (*Seeliger and de Groot, 2010*; *Gapsys et al., 2015*; *Aldeghi et al., 2019*). Specifically, we calculate the work values from the forward and backward transitions (i.e. $W = \int_{\lambda=0}^{\lambda=1} \frac{\partial H}{\partial \lambda} d\lambda$) and then estimate free energy differences between the initial and final states using Bennet's Acceptance Ratio (*Bennett, 1976*) (for more details, we refer to the PMX papers (*Seeliger and de Groot, 2010*; *Gapsys et al., 2015*; *Aldeghi et al., 2019*)). For our free energy estimates, we report bootstrap errors.

The simulations have been analyzed using a combination of analysis programs in the Gromacs suite (*Abraham et al., 2015*) and in-house Python scripts that make extensive use of the MDTraj library (*McGibbon et al., 2015*). Specifically, we used `gmx dipoles` to calculate the dielectric constant ε for condensates and different solvents from the fluctuations in the dipole moment of the simulation box, $\mathbf{M}$, which is defined as (*Neumann, 1983*)

$$\varepsilon = 1 + \frac{\langle \mathbf{M}^2 \rangle - \langle \mathbf{M} \rangle^2}{3\varepsilon_0 k_B T \langle V \rangle}. \tag{2}$$

In this expression, ε_0 is the vacuum permittivity, k_B is Boltzmann's constant and V is the volume of the simulation box. Error analysis has been performed by blocking.

Quantum chemical calculations

We performed quantum mechanical calculations using the Gaussian 16 program to study the interaction energy of the different complexes. All calculations were made at the Density Functional Theory (DFT) level, using the ωB97XD functional (*Chai and Head-Gordon, 2008*) and Pople's 6-311++G(d,p) basis set (*Hehre et al., 1972*) for geometry optimization. Energetics were refined with the 6-311++G(3df,2p) basis set. Geometry optimizations were done in different solvents, using the Polarizable Continuum Model approach (*Mennucci, 2012*). The interaction energy in a given solvent was calculated according to

$$\Delta E_{\text{Int}}^{ij} = E^{ij} - \left(E^i + E^j \right) \tag{3}$$

where i and j specify the interacting monomers. We consider p-cresol to represent the sidechain of tyrosine (Y, hereafter), toluene to represent the sidechain of phenylalanine (F hereafter) and N-methyl-acetamide to represent an amide bond group (A, hereafter). Dimers ij include different combinations of Y, F, and A in various orientations.

To calculate the free energy transfer of a monomer from water to a given solvent, we used the SMD solvation model (*Marenich et al., 2009*) to calculate accurate solvation-free energies of the monomers in different solvents, using B3LYP/6-31+G(d,p) level of theory (*Becke, 1993*; *Chengteh et al., 1988*). Thus, the transfer free energy from water to a given solvent s is calculated according to

$$\Delta G_{\text{Transfer}}^i = \Delta G^i(s) - \Delta G^i(\text{H}_2\text{O}) \tag{4}$$

As solvents, we have considered three different groups:

1. Alcohols: methanol, ethanol, 1-propanol, 1-butanol, 1-poentanol, 1-hexanol, 1-nonanol and 1-decanol.
2. Aliphatic solvents: 2-nitropropane, 2-bromopropane, 1-bromopropane, 1-bromopentane, 1-fluorooctane, n-pentane, n-decane nitrobenzene, o-dichlorobenzene, nromobenzene, iodobenzene, ethylbenzene, benzene, toluene and cyclohexane.
3. Water with varying dielectric values.

These solvents span a broad range of values of the dielectric constant and can be viewed to represent different chemical environments where contact formation can take place.

Results and Discussion

Transfer free energies into a condensate slab

Our first goal is to find whether differences in transfer free energy for Phe and Tyr variants capture the hierarchy in sticker strength in biomolecular condensates. Ideally one would calculate the transfer free energies for the protein of interest inside a realistic protein condensate. However, this would involve simulations of multiple copies of an IDR with tens or hundreds of amino acids, resulting in a system with very slow conformational dynamics (*Paloni et al., 2020*; *Zheng et al., 2020*; *Galvanetto et al., 2023*). As a computationally efficient alternative, we use a simple model peptide with sequence GGXGG, with X being either F or Y, which is transferred from water into the dense mixtures of short peptides we have recently characterized (*De Sancho, 2022*). Peptide condensates are useful model systems as they retain many properties of protein condensates but allow for much greater computational efficiency (*Paloni et al., 2020*; *Tang et al., 2021*; *De Sancho, 2022*; *Brown and Potoyan, 2024*).

Free energy estimates for the horizontal branches in the thermodynamic cycle in Figure 2 were obtained from alchemical transformations, which we performed using a state-of-the-art non-equilibrium switching method (*Seeliger and de Groot, 2010*; *Gapsys et al., 2015*; *Aldeghi et al., 2019*) (see

Figure 3. (A) Time series for the accessible surface area of the GGXGG in the GSY condensate for different values of λ. (B) Time series for the density profiles of all peptides (blue) and the GGXGG peptide (red) in the coexistence simulations. (C) Representative simulation box with a fully solvated condensate in slab geometry including a GGXGG peptide (spheres) and the capped amino acid mixture (G: white, S: yellow and Y: green).

Methodology). The first step in this procedure involves running long equilibrium simulations at the initial λ-states (i.e. $\lambda = 0$ and $\lambda = 1$). While these calculations are straightforward for a peptide in solution, within the condensate we must first ensure that the condensate slab has converged to a stable density. Also the peptide that experiences the transformation must remain buried within the condensate for all the snapshots that we use as initial frames, so that we do not average the work in the dilute and dense phases. In Figure 3A we show the solvent accessible surface area ($SASA$) of all protein residues in a GSY condensate with the model peptide submerged. After an initial increase in $SASA$, at around 100 ns the values stabilize and remain flat for the remainder of the simulation, suggesting that equilibrium between the dense and dilute phases has been reached. We also show the density profiles for all the amino acid residues in the condensate and for the GGXGG peptide alone (Fig. 3B). For both λ values, we find that during the whole duration of the simulation, the GGXGG peptide remains buried within the condensate (see a representative snapshot in Fig. 3C). The protein density in the GSY condensate is consistent with the values reported before (i.e. ~ 1000 mg/mL protein and $\sim$200 mg/mL water) (***De Sancho, 2022***), and slightly larger than in full-length IDP condensates (***Zheng et al., 2020***).

After preparing our system, we selected initial configurations for non-equilibrium switching from evenly spaced snapshots of the $\lambda = 0$ and $\lambda = 1$ trajectories starting at 100 ns. We ran 200 independent simulations forward and backwards, collecting the values of $\langle \partial H / \partial \lambda \rangle$. Alchemical transformations were also performed in the same way for the peptide in TIP3P water (see Methods). Then we estimate transfer free energy differences by combining data from the transformations

inside the condensate and in water. For the GSY condensate, we find that $\Delta\Delta G_{\text{Transfer}}$ is small and negative (-1.91 ± 0.53 kJ/mol), indicating that transfer into a peptide condensate is slightly more favourable for the Tyr-containing peptide than for the Phe variant. This result is consistent with mounting experimental evidence that points to a greater propensity to form condensates for Tyr rather than Phe in variants of proteins like FUS, LAF1 and hnRNPA.

Similar protein-protein interactions in Tyr and Phe condensates

One possible explanation for this result is the formation of preferential sticker-sticker interactions between the GGYGG peptide and the GSY condensate. While the hydroxyl group in tyrosine in the pentapeptide may form hydrogen bonds with other tyrosines in GSY, these should be depleted for the GGFGG variant. Conversely, if the condensate were composed of Phe instead of Tyr (i.e. the GSF condensate), there would be no reason for a preference for GGYGG. Additionally, from hydrophobicity arguments alone, transfer to the GSF condensate should be favoured for Phe. To test this hypothesis, we have run the same procedure for GSF instead of GSY. We show the time series data for $SASA$ and protein density for the GSF simulations in Appendix 2. We find that $\Delta\Delta G_{\text{Transfer}} = -2.75 \pm 0.58$ kJ/mol, hence also favourable to Tyr and of magnitude comparable to the result for transfer to GSY.

To gain further insight into the dominant interactions within condensates formed by different stickers, we have run additional simulations of isolated dense phases for GSY and GSF condensates (see Figure 4A). First we look at the relative number of different interaction pairs inside the condensates. We find that the interaction patterns are very similar in both cases, with a greater number of contacts formed by the aromatic stickers and smaller contributions from Gly and Ser (see Fig. 4B). This result recapitulates observations in atomistic simulations of full-length IDRs (*Zheng et al., 2020*) and is consistent with the scaffold-client relationship between aromatic and polar residues (*Dignon et al., 2020*). To focus on $\pi - \pi$ interactions between pairs of aromatic side chains, we calculated the mean inter-side chain distances and the angles between vectors normal to the aromatic rings. Figure 4C shows their distribution, which is similar in both condensates. Even though aromatic $\pi - \pi$ stacking is feasible, it does not play a dominant role (see the regions with low inter-residue distance and angles close to 0 or π). We also examine $sp^2 - \pi$ contacts formed between aromatic and peptide bond groups (*Vernon et al., 2018*), which are much more common. This is consistent with previous simulation work (*Zheng et al., 2020*; *Murthy et al., 2021*; *Rekhi et al., 2024*) and also with a statistical analysis of experimental PDB structures which showed that these interactions are particularly prevalent in disordered regions of folded proteins (*Vernon et al., 2018*). Interestingly, also for these $sp^2 - \pi$ interactions, no significant differences were found between GSY and GSF condensates.

These findings suggest that the preference for Tyr over Phe to form condensates, as captured by $\Delta\Delta G_{\text{Transfer}}$, is not primarily driven by differences in protein-protein interaction patterns. Although interactions may be slightly more favourable for Tyr than for Phe, as suggested by atomistic potentials of mean force (*Chelli et al., 2002*; *Joseph et al., 2021*), these residues have similar contact statistics both for sticker-sticker and for sticker-spacer contacts, and both exhibit a preference for $sp^2 - \pi$ rather than $\pi - \pi$ interactions within the condensate.

Condensates as hydrated media

An alternative explanation could involve the solvent effects on the Phe/Tyr peptides. To investigate this, we begin by analysing the distribution of water molecules around the distinct side chains in the GSF and GSY dense phase simulations, revealing notable differences. Figure 5A presents the radial distribution function ($g(r)$) for water oxygen around the C^ζ in Phe and Tyr side chains. In the case of Tyr, a prominent peak at short distances suggests the presence of water molecules involved in hydrogen bonds, which are enthalpically favourable and entropically unfavourable due to the water structuring around the Tyr hydroxyl group (*Lin et al., 2017*). These thermodynamic contributions are intertwined in the calculated $\Delta\Delta G_{\text{Transfer}}$ and will vary depending on whether transfer of

Figure 4. (A) Representative snapshots from the simulations of dense phases. (B) Interaction matrix for the normalized number of contacts between different pairs of amino acid residues in the condensates (top) and for each type of amino acid (bottom). (C) Density plots for side chain interactions between aromatic side chains, as characterised by the mean inter-residue distance and the angle θ between the vectors normal to the rings (*Calinsky and Levy, 2024*). (D) Density plots for sp^2-π interactions between amide bonds and aromatic side chains, as characterised by the mean inter-group distance and the angle θ between the vectors normal to the peptide bond and ring planes. In all panels, results for GSY and GSF condensates are shown on the top and bottom, respectively. Representative snapshots of relevant interactions for each type of pair are shown.

the peptide from water occurs into a condensate, into the core of a folded protein or an altogether
different solvent, capable or not of hydrogen bonding.

To investigate this dependence, we performed the same type of alchemical calculations using
solvents such as octanol or cyclohexane –used in the past as models for the interior of folded pro-
teins (*Kauzmann, 1959*; *Nozaki and Tanford, 1971*; *Radzicka and Wolfenden, 1988*; *Wimley et al.,*
1996; *Pace et al., 2011*)– as well as shorter alcohols or acetone. The results of these calculations
are shown in Figure 5B. In the case of a purely non-polar solvent like cyclohexane, the difference
in free energy from Phe to Tyr is large and positive ($\Delta\Delta G_{\mathrm{Transfer}} = 13.29 \pm 0.20$ kJ/mol), indicating that
Phe is favoured. This is also the case for aromatic solvents like benzene and toluene. This result
is consistent with the stronger interaction strength of Phe-Phe interactions in the Miyazawa and
Jernigan potential (*Miyazawa and Jernigan, 1996*) (see Fig. 1C), derived from contacts in the apolar
cores of proteins where hydrophobicity dominates. The difference in free energy is smaller for
alcohols and decreases gradually as we reduce the length of the aliphatic chain. For acetone, Tyr
is slightly more favourable ($\Delta\Delta G_{\mathrm{Transfer}} = -0.91 \pm 0.25$ kJ/mol). Clearly, the $\Delta\Delta G_{\mathrm{Transfer}}$ values have a
dependence on the dielectric constant (ε), which we have also calculated for the GSY and GSF con-
densates (see Figure 5C). Our ε values for the condensates (39 ± 5 for GSY and 47 ± 3 for GSF) are
surprisingly close to that derived from experiments on Ddx condensates (45 ± 13) (*Nott et al., 2015*),
and appear at the rightmost part of the diagram. In summary, for non-polar solvents with low ε
Phe is preferred to Tyr, while for polar solvents, including the condensates, the trend is reversed.
At intermediate values of ε, there is a crossover between a regime dominated by hydrophobicity
and one favoured by hydrogen bonding by Tyr.

Quantum calculations confirm a crossover in interaction strength

322   To characterize these subtle effects in greater detail we resort to quantum chemical calculations,
323   which can more accurately determine interaction strengths for specific configurations involving
324   aromatic residues (*Carter-Fenk et al., 2023*; *Calinsky and Levy, 2024*). We consider different modes

Figure 5. (A) Radial distribution function for water oxygen around the C^ζ in Phe/Tyr for GSF and GSY condensates. We show a representative overlay of simulation snapshots where water molecules are hydrogen-bonded to the Tyr side chain. (B) Transfer free energy differences from water to a different medium between Tyr and Phe. We consider condensates (green), polar solvents (orange) and apolar solvents (blue). (C) Same as a function of calculated dielectric constant, ε.

of interaction between the sidechains of Phe and Tyr, represented by toluene (F) and p-cresol (Y), respectively. We considered various orientations in the optimization process and selected the two most stable structures for each case. For F-F, the most stable structures correspond to *cross* and *antiparallel* configurations, whereas in the case of Y-Y, *H-bonded* and *antiparallel* are most favoured (see Fig. 6A and C). Our results are consistent with previous work on the toluene dimer, where the energy minima also corresponded to the cross and antiparallel structures, with only minor differences in their energies (*Tsuzuki et al., 2005*; *Rogers et al., 2006*). Other structures, like parallel or T-shaped, are less stable for toluene, although in the benzene dimer, T-shaped conformers can be favored (*Chipot et al., 1996*). In our optimizations, T-shaped F-F structures tended to collapse to less stable parallel structures, and hence we do not include them here. The interaction energy values we obtained are qualitatively similar to those in the literature (*Calinsky and Levy, 2024*), though slightly more negative, and they follow the same rank order (i.e. Y-Y H-bonded < antiparallel Y-Y < cross F-F). We also calculate the energies for Phe and Tyr sidechains interacting with an amide group (A) representing a peptide bond in the prevalent sp^2-π interactions (*Vernon et al., 2018*) (see Fig. 6C). In the case of Tyr, we have also analyzed the configuration where it forms a hydrogen bond with the amide through the alcohol group of p-cresol.

To explore the environmental dependence in contact formation for amino acid pairs, we have calculated values of the interaction energies ($\Delta E_{\mathrm{Int}}^{ij}$) considering a set of protic and non-protic solvents (see Fig. 6A). Additionally, we have run the same calculations using a modified water with the dielectric constant as a free parameter. The overall trends in interaction energies are very similar across interaction types. In all cases, the interaction energy tends to become stronger as the value of the dielectric decreases. Also, the relative strength of interactions is consistent across solvents. The strongest interaction corresponds to the Y-A hydrogen-bonded interaction. Next comes a series of partial π-stacking configurations with contributions of hydrogen bonds and sp^2-

Figure 6. (A) ΔE^{ij}_{Int} and (B) $\Delta E^{ij}_{contact}$ for different interaction pairs ij and different solvents as a function of the dielectric constant, ε. (C) Optimized geometries from quantum chemical calculations for Phe and Tyr interaction pairs. (D) $\Delta\Delta E^{Solvent}_{Contact}$ with respect to the cross-F-F interaction for different solvents as a function of the dielectric. Different symbols correspond to the different solvent types. Filled: alcohols; empty: alkanes/benzenes; lines: water with different dielectric.

π interactions, namely, Y-Y H-bond, Y-A sp^2-π and F-A sp^2-π structures, which we have found to be particularly abundant in our MD runs. Finally, the π-stacked F-F cross is the weakest interaction among those included in this analysis.

Therefore, if we consider these interaction energies as the primary driving force for condensate formation, Tyr should be more adhesive than Phe independently of the environment. However, as realized by Miyazawa and Jernigan (***Miyazawa and Jernigan, 1996***; ***Thomas and Dill, 1996***), contact formation, either within the core of folded proteins or in the case of molecular condensates, involves two processes: first, the transfer of these groups from water to an environment with a lower dielectric; and second, the interaction between side chain groups (or with the backbone), which is captured by ΔE^{ij}_{Int} (see Figure 2B). Hence, for a given solvent, the total energy can be calculated from the sum of these two contributions as

$$\Delta E^{ij}_{Contact} = \Delta E^{ij}_{Int} + \Delta G^{i}_{Transfer} + \Delta G^{j}_{Transfer} \tag{5}$$

The last two terms on the right-hand side of this expression capture the propensity of a monomer to be transferred from water to the solvent, which can be modelled using polarizable continuum models such as SMD (***Marenich et al., 2009***). Notice that when the solvent is a low dielectric non-protic solvent such as cyclohexane, $\Delta G^{i}_{Transfer}$ recapitulates the results from a standard hydrophobicity scale (***Kyte and Doolittle, 1982***).

When both contributions are considered, the picture changes dramatically (see Fig. 6B). In non-protic aliphatic solvents, such as alkanes and benzene derivatives, interaction for F-F pairs becomes more favourable than in Y-Y pairs, irrespective of the dielectric constant. On the other hand, in alcohols, $\Delta E^{ij}_{Contact}$ tends to favour Y-A interactions at high ε and F-F pairs at low ε. In other words, at intermediate dielectrics, we find a crossover between F and Y in the propensity to form pairwise contacts in solvents. In our calculations, we have also considered a water-like solvent where the dielectric constant can be tuned as a free parameter (see Fig. 6C). Also, we find a similar tendency

in this case, but now the crossover is obtained at slightly lower values of ϵ. In Figure 6D, we show

$\Delta\Delta E_{\text{Contact}}^{ij}$ values, where we are using as reference the energy of the F-F cross configuration, for

three selected interactions: Y-Y H-bond, Y-A H-bond, and F-A, for the three types of environments.

Negative values of $\Delta\Delta E_{\text{Contact}}^{ij}$ indicate a preference of the corresponding interaction relative to F-F

cross, while positive values indicate that cross-F-F is stronger. Clearly, there is a crossover between

a regime where toluene forms stronger interactions and one where p-cresol is preferred, which is

highly influenced by the type of environment. In a non-protic solvent, where hydrogen bonding

is not possible, or in a protic solvent with low dielectric constant, we find that F-F interactions are

favoured. In contrast, in high dielectric environments, Y-Y and Y-A interactions dominate.

Conclusions

In this work, we have aimed to reconcile the apparent contradictions between the lower solvation-

free energy (*Wolfenden et al., 1981*; *Chang et al., 2007*) and stronger hydrophobicity of Phe (*Kyte*

and Doolittle, 1982; *Tesei et al., 2021*) and the greater contact energies assigned to Phe-Phe interac-

tions in proteins (*Miyazawa and Jernigan, 1996*) with the higher sticker strength of Tyr in biomolec-

ular condensates (*Lin et al., 2017*; *Wang et al., 2018*; *Schuster et al., 2020*; *Bremer et al., 2022*). In

the past, the different propensities of positively charged stickers to form condensates had been

explained from the fundamental properties of Lys and Arg (*Wang et al., 2018*; *Das et al., 2020*;

Gobbi and Frenking, 1993; *Hong et al., 2022*). However, an explanation for the different strengths

of Phe and Tyr remained elusive (*Das et al., 2020*).

Through a combination of classical MD simulations and quantum mechanical calculations, we

have found that, in addition to the distinct types of interactions formed by Phe and Tyr, a crucial fac-

tor in their overall propensity for contact formation across different environments is the transfer

free energy. At one extreme, we find environments such as the well-packed core of a protein, where

the dielectric constant can be as low as 2-4 (*Pitera et al., 2001*). This type of environment can be

modelled using aliphatic solvents, an approach followed extensively in the past (*Kauzmann, 1959*;

Nozaki and Tanford, 1971; *Radzicka and Wolfenden, 1988*; *Wimley et al., 1996*; *Pace et al., 2011*). In

this low-dielectric regime, Phe-Phe interactions are the most favourable, as evidenced by both our

classical transfer free energies (Fig. 5C) and the estimated quantum $\Delta\Delta E_{\text{Contact}}^{ij}$ values (Fig. 6). This

explains why Phe forms the strongest interactions in contact matrices like that by Miyazawa and

Jernigan, which were derived from statistics from 3D structures of proteins (*Miyazawa and Jerni-*

gan, 1996) and are known to be dominated by hydrophobicity effects (*Pokarowski et al., 2005*). At

the other extreme, in environments with high dielectric constants such as aqueous or polar solvent

conditions, Tyr-Tyr hydrogen bonds become more favourable. Biomolecular condensates, due to

their high water content (~ 200 mg/mL in our simulations, but up to ~600 mg/mL in experiments

(*Zheng et al., 2020*)), are best modelled as a solvent with an intermediate dielectric. Hence, they op-

erate in the second regime, although the influence of the dielectric is subtle, due to the possibility

of crossover as a function of the dielectric.

The conclusions from our work are necessarily limited due to our reductionist approach. First,

in our calculations, we are considering very short peptides including a single sticker, and hence

we cannot account for the influence of multivalency in protein IDRs (*Martin et al., 2020*). Also,

we have used a minimal alphabet of possible interaction pairs limited to two types of amino acid

residues. We do not consider effects relative to cation-π interactions that have been characterized

by others (*Brady et al., 2017*; *Das et al., 2020*; *Aldeghi et al., 2018*; *Hazra and Levy, 2023*; *Calinsky*

and Levy, 2024). Also, results may vary somewhat when considering different classical force fields.

For example, our condensates are more protein-dense than found in experiments, possibly due to

the propensity of the TIP3P water model to induce compact states (*Piana et al., 2015*). However the

methodology proposed in the present work can be easily extended to other types of interactions.

The overall qualitative conclusions on the crossover in interaction strengths as a function of the

environment are likely to be maintained if different quantum approaches are used for contact

calculations or different force fields are used for the classical simulations.

Acknowledgments

The authors gratefully acknowledge conversations with Priya R. Banerjee, Rohit Pappu and Tanja Mittag, which inspired this research. The work has been financed by the Spanish Ministry of Science and Innovation through project PID2021-127907NB-I00, and the Basque Government (Project IT1584-22). The authors also thank the IZO-SGI SGIker (UPV/EHU/ERDF,EU) and DIPC for technical and human support and for the allocation of computational resources.

References

Abraham MJ, Murtola T, Schulz R, Páll S, Smith JC, Hess B, Lindahl E. GROMACS: High performance molecular simulations through multi-level parallelism from laptops to supercomputers. SoftwareX. 2015; 1-2:19 – 25. doi: 10.1016/j.softx.2015.06.001.

Aldeghi M, Gapsys V, de Groot BL. Accurate estimation of ligand binding affinity changes upon protein mutation. ACS Cent Sci. 2018; 4(12):1708–1718.

Aldeghi M, de Groot BL, Gapsys V. In: Sikosek T, editor. Accurate Calculation of Free Energy Changes upon Amino Acid Mutation New York, NY: Springer New York; 2019. p. 19–47. doi: 10.1007/978-1-4939-8736-8_2.

Banani SF, Lee HO, Hyman AA, Rosen MK. Biomolecular condensates: organizers of cellular biochemistry. Nat Rev Mol Cell Biol. 2017; 18(5):285–298. doi: 10.1038/nrm.2017.7.

Bannan CC, Calabró G, Kyu DY, Mobley DL. Calculating Partition Coefficients of Small Molecules in Octanol/Water and Cyclohexane/Water. J Chem Theory Comput. 2016; 12(8):4015–4024. doi: 10.1021/acs.jctc.6b00449.

Becke AD. A new mixing of Hartree-Fock and local density-functional theories. J Chem Phys. 1993; 98(2):1372–1377. doi: 10.1063/1.464304.

Bennett CH. Efficient estimation of free energy differences from Monte Carlo data. J Comput Phys. 1976; 22(2):245–268.

Berendsen HJ, Postma Jv, Van Gunsteren WF, DiNola A, Haak JR. Molecular dynamics with coupling to an external bath. J Chem Phys. 1984; 81(8):3684–3690.

Brady JP, Farber PJ, Sekhar A, Lin YH, Huang R, Bah A, Nott TJ, Chan HS, Baldwin AJ, Forman-Kay JD, et al. Structural and hydrodynamic properties of an intrinsically disordered region of a germ cell-specific protein on phase separation. Proc Natl Acad Sci USA. 2017; 114(39):E8194–E8203.

Brangwynne CP, Eckmann CR, Courson DS, Rybarska A, Hoege C, Gharakhani J, Jülicher F, Hyman AA. Germline P granules are liquid droplets that localize by controlled dissolution/condensation. Science. 2009; 324(5935):1729–1732.

Bremer A, Farag M, Borcherds WM, Peran I, Martin EW, Pappu RV, Mittag T. Deciphering how naturally occurring sequence features impact the phase behaviours of disordered prion-like domains. Nat Chem. 2022; (14):196–207. doi: 10.1038/s41557-021-00840-w.

Brown WH, Potoyan DA. Phase separation of multicomponent peptide mixtures into dehydrated clusters with hydrophilic cores. Biophys J. 2024; 123(3):349–360.

Bussi G, Donadio D, Parrinello M. Canonical sampling through velocity rescaling. J Chem Phys. 2007; 126(1):014101.

Calinsky R, Levy Y. Aromatic Residues in Proteins: Re-Evaluating the Geometry and Energetics of π–π, Cation-π, and CH-π Interactions. J Phys Chem B. 2024; .

Carter-Fenk K, Liu M, Pujal L, Loipersberger M, Tsanai M, Vernon RM, Forman-Kay JD, Head-Gordon M, Heidar-Zadeh F, Head-Gordon T. The Energetic Origins of Pi–Pi Contacts in Proteins. J Am Chem Soc. 2023; 145(45):24836–24851.

Chai JD, Head-Gordon M. Long-range corrected hybrid density functionals with damped atom–atom dispersion corrections. Phys Chem Chem Phys. 2008; 10(44):6615–6620.

Chang J, Lenhoff AM, Sandler SI. Solvation free energy of amino acids and side-chain analogues. J Phys Chem B. 2007; 111(8):2098–2106.

Chelli R, Gervasio FL, Procacci P, Schettino V. Stacking and T-shape competition in aromatic- aromatic amino
acid interactions. J Am Chem Soc. 2002; 124(21):6133–6143.

Chengteh L, Yang W, Parr RG. Development of the Colic-Salvetti correlation-energy formula into a functional
of the electron density. Phys Rev B. 1988; 37(2):785–789.

Chipot C, Jaffe R, Maigret B, Pearlman DA, Kollman PA. Benzene dimer: A good model for π-π interactions in
proteins? A comparison between the benzene and the toluene dimers in the gas phase and in an aqueous
solution. J Am Chem Soc. 1996; 118(45):11217–11224. doi: 10.1021/ja961379l.

Choi JM, Holehouse AS, Pappu RV. Physical Principles Underlying the Complex Biology of Intracellular Phase
Transitions. Annu Rev Biophys. 2020; 49(1):107–133. doi: 10.1146/annurev-biophys-121219-081629.

Das S, Lin YH, Vernon RM, Forman-Kay JD, Chan HS. Comparative roles of charge, π, and hydrophobic interac-
tions in sequence-dependent phase separation of intrinsically disordered proteins. Proc Natl Acad Sci USA.
2020; 117(46):28795–28805.

De Sancho D. Phase separation in amino acid mixtures is governed by composition. Biophys J. 2022;
121(21):4119–4127.

Dignon GL, Best RB, Mittal J. Biomolecular Phase Separation: From Molecular Driving Forces to Macroscopic
Properties. Annu Rev Phys Chem. 2020; 71(1):53–75. doi: 10.1146/annurev-physchem-071819-113553.

Dignon GL, Zheng W, Best RB, Kim YC, Mittal J. Relation between single-molecule properties and phase
behavior of intrinsically disordered proteins. Proc Natl Acad Sci USA. 2018; 115(40):9929–9934. doi:
10.1073/pnas.1804177115.

Dignon GL, Zheng W, Kim YC, Best RB, Mittal J. Sequence determinants of protein phase behavior from a
coarse-grained model. PLoS Comput Biol. 2018 01; 14(1):1–23. doi: 10.1371/journal.pcbi.1005941.

Fisher RS, Elbaum-Garfinkle S. Tunable multiphase dynamics of arginine and lysine liquid condensates. Nat
Commun. 2020; 11(1):4628.

Galvanetto N, Ivanović MT, Chowdhury A, Sottini A, Nüesch MF, Nettels D, Best RB, Schuler B. Extreme dynam-
ics in a biomolecular condensate. Nature. 2023; 619(7971):876–883.

Gapsys V, Michielssens S, Seeliger D, de Groot BL. pmx: Automated protein structure and topology generation
for alchemical perturbations. J Comput Chem. 2015; 36(5):348–354. doi: https://doi.org/10.1002/jcc.23804.

Gapsys V, Michielssens S, Seeliger D, de Groot BL. Accurate and rigorous prediction of the changes in protein
free energies in a large-scale mutation scan. Angew Chem Int Ed. 2016; 55(26):7364–7368.

Gapsys V, Pérez-Benito L, Aldeghi M, Seeliger D, Van Vlijmen H, Tresadern G, De Groot BL. Large scale relative
protein ligand binding affinities using non-equilibrium alchemy. Chem Sci. 2020; 11(4):1140–1152.

Gobbi A, Frenking G. Y-conjugated compounds: the equilibrium geometries and electronic structures of guani-
dine, guanidinium cation, urea, and 1, 1-diaminoethylene. J Am Chem Soc. 1993; 115(6):2362–2372.

Greig JA, Nguyen TA, Lee M, Holehouse AS, Posey AE, Pappu RV, Jedd G. Arginine-enriched mixed-charge do-
mains provide cohesion for nuclear speckle condensation. Mol Cell. 2020; 77(6):1237–1250.

Hazra MK, Levy Y. Cross-talk of cation- π interactions with electrostatic and aromatic interactions: A salt-
dependent trade-off in biomolecular condensates. J Phys Chem Lett. 2023; 14(38):8460–8469.

Hehre WJ, Ditchfield K, Pople JA. Self-consistent molecular orbital methods. XII. Further extensions of gaussian-
type basis sets for use in molecular orbital studies of organic molecules. J Chem Phys. 1972; 56(5):2257–2261.
doi: 10.1063/1.1677527.

Hong Y, Najafi S, Casey T, Shea JE, Han SI, Hwang DS. Hydrophobicity of arginine leads to reentrant liquid-liquid
phase separation behaviors of arginine-rich proteins. Nat Commun. 2022; 13(1):7326.

Jorgensen WL, Chandrasekhar J, Madura JD, Impey RW, Klein ML. Comparison of Simple Potential Functions
for Simulating Liquid Water. J Chem Phys. 1983; 79(2):926–935.

Joseph JA, Reinhardt A, Aguirre A, Chew PY, Russell KO, Espinosa JR, Garaizar A, Collepardo-Guevara R. Physics-
driven coarse-grained model for biomolecular phase separation with near-quantitative accuracy. Nat Com-
put Sci. 2021; 1(11):732–743. doi: 10.1038/s43588-021-00155-3.

Kauzmann W. Some factors in the interpretation of protein denaturation. In: *Adv. Protein Chem.*, vol. 14 Elsevier;
1959.p. 1–63.

Kyte J, Doolittle RF. A simple method for displaying the hydropathic character of a protein. J Mol Biol. 1982;
157(1):105–132.

Lin Y, Currie SL, Rosen MK. Intrinsically disordered sequences enable modulation of protein phase separation
through distributed tyrosine motifs. J Biol Chem. 2017; 292(46):19110–19120.

Lindorff-Larsen K, Piana S, Palmo K, Maragakis P, Klepeis JL, Dror RO, Shaw DE. Improved side-chain torsion po-
tentials for the Amber ff99SB protein force field. Proteins. 2010; 78(8):1950–1958. doi: 10.1002/prot.22711.

Maraldo A, Rnjak-Kovacina J, Marquis C. Tyrosine–a structural glue for hierarchical protein assembly. Trends
Biochem Sci. 2024; .

Marenich AV, Cramer CJ, Truhlar DG. Universal solvation model based on solute electron density and on a
continuum model of the solvent defined by the bulk dielectric constant and atomic surface tensions. J Phys
Chem B. 2009; 113(18):6378–6396.

Martin EW, Holehouse AS, Peran I, Farag M, Incicco JJ, Bremer A, Grace CR, Soranno A, Pappu RV, Mittag T.
Valence and patterning of aromatic residues determine the phase behavior of prion-like domains. Science.
2020; 367(6478):694–699. doi: 10.1126/science.aaw8653.

Martin EW, Mittag T. Relationship of Sequence and Phase Separation in Protein Low-Complexity Regions.
Biochemistry. 2018; 57(17):2478–2487. doi: 10.1021/acs.biochem.8b00008.

Martinez-Martin I, Crousilles A, Herrero-Galan E, Velazquez-Carreras D, Mortensen SA, Ochoa JP, Garcia-Pavia
P, de Sancho D, Wilmanns M, Alegre-Cebollada J. Structural basis of domain destabilization by dilated-
cardiomyopathy-causing missense mutations in titin. Biophys J. 2023; 122(3):172a.

McGibbon RT, Beauchamp KA, Harrigan MP, Klein C, Swails JM, Hernández CX, Schwantes CR, Wang LP, Lane
TJ, Pande VS. MDTraj: A Modern Open Library for the Analysis of Molecular Dynamics Trajectories. Biophys
J. 2015; 109(8):1528 – 1532. doi: 10.1016/j.bpj.2015.08.015.

Mennucci B. Polarizable continuum model. Wiley Interdiscip Rev Comput Mol Sci. 2012; 2(3):386–404.

Mey ASJS, Allen BK, Bruce McDonald HE, Chodera JD, Hahn DF, Kuhn M, Michel J, Mobley DL, Naden LN, Prasad
S, Rizzi A, Scheen J, Shirts MR, Tresadern G, Xu H. Best Practices for Alchemical Free Energy Calculations
[Article v1.0]. Living J Comp Mol Sci. 2020; 2(1):18378. doi: 10.33011/livecoms.2.1.18378.

Mittag T, Pappu RV. A conceptual framework for understanding phase separation and addressing open ques-
tions and challenges. Mol Cell. 2022; 82(12):2201–2214. doi: https://doi.org/10.1016/j.molcel.2022.05.018.

Miyazawa S, Jernigan RL. Residue–residue potentials with a favorable contact pair term and an unfavorable
high packing density term, for simulation and threading. J Mol Biol. 1996; 256(3):623–644.

Murthy AC, Tang WS, Jovic N, Janke AM, Seo DH, Perdikari TM, Mittal J, Fawzi NL. Molecular interactions con-
tributing to FUS SYGQ LC-RGG phase separation and co-partitioning with RNA polymerase II heptads. Nat
Struct Mol Biol. 2021; 28(11):923–935.

Neumann M. Dipole moment fluctuation formulas in computer simulations of polar systems. Mol Phys. 1983;
50(4):841–858.

Nott TJ, Petsalaki E, Farber P, Jervis D, Fussner E, Plochowietz A, Craggs TD, Bazett-Jones DP, Pawson T, Forman-
Kay JD, Baldwin AJ. Phase transition of a disordered nuage protein generates environmentally responsive
membraneless organelles. Mol Cell. 2015; 57(5):936–947.

Nozaki Y, Tanford C. The Solubility of Amino Acids and Two Glycine Peptides in Aqueous Ethanol and Dioxane
Solutions: establishment of a hydrophobicity scale. J Biol Chem. 1971; 246(7):2211–2217. doi: 10.1016/S0021-
9258(19)77210-X.

Pace CN, Fu H, Fryar KL, Landua J, Trevino SR, Shirley BA, Hendricks MM, Iimura S, Gajiwala K, Scholtz JM,
Grimsley GR. Contribution of Hydrophobic Interactions to Protein Stability. JMol Biol. 2011; 408(3):514–528.
doi: https://doi.org/10.1016/j.jmb.2011.02.053.

Pace CN, Horn G, Hebert EJ, Bechert J, Shaw K, Urbanikova L, Scholtz JM, Sevcik J. Tyrosine hydrogen bonds
make a large contribution to protein stability. J Mol Biol. 2001; 312(2):393–404.

Paloni M, Bailly R, Ciandrini L, Barducci A. Unraveling Molecular Interactions in Liquid–Liquid Phase Sepa-
ration of Disordered Proteins by Atomistic Simulations. J Phys Chem B. 2020; 124(41):9009–9016. doi:
10.1021/acs.jpcb.0c06288.

Paloni M, Bussi G, Barducci A. Arginine multivalency stabilizes protein/RNA condensates. Protein Sci. 2021;
30(7):1418–1426.

Parrinello M, Rahman A. Crystal structure and pair potentials: A molecular-dynamics study. Phys Rev Lett.
1980; 45(14):1196.

Piana S, Donchev AG, Robustelli P, Shaw DE. Water Dispersion Interactions Strongly Influence Simulated
Structural Properties of Disordered Protein States. J Phys Chem B. 2015; 119(16):5113–5123. doi:
10.1021/jp508971m.

Pitera JW, Falta M, van Gunsteren WF. Dielectric properties of proteins from simulation: the effects of solvent,
ligands, pH, and temperature. Biophys J. 2001; 80(6):2546–2555.

Plaxco KW, Morton CJ, Grimshaw SB, Jones JA, Pitkeathly M, Campbell ID, Dobson CM. The effects of guanidine
hydrochloride on the'random coil 'conformations and NMR chemical shifts of the peptide series GGXGG. J
Biomol NMR. 1997; 10:221–230.

Pokarowski P, Kloczkowski A, Jernigan RL, Kothari NS, Pokarowska M, Kolinski A. Inferring ideal amino acid
interaction forms from statistical protein contact potentials. Proteins. 2005; 59(1):49–57.

Radzicka A, Wolfenden R. Comparing the polarities of the amino acids: side-chain distribution coefficients
between the vapor phase, cyclohexane, 1-octanol, and neutral aqueous solution. Biochemistry. 1988;
27(5):1664–1670.

Regy RM, Thompson J, Kim YC, Mittal J. Improved coarse-grained model for studying sequence dependent
phase separation of disordered proteins. Protein Sci. 2021; 30(7):1371–1379. doi: 10.1002/pro.4094.

Rekhi S, Garcia CG, Barai M, Rizuan A, Schuster BS, Kiick KL, Mittal J. Expanding the molecular language of
protein liquid–liquid phase separation. Nat Chem. 2024; p. 1–12.

Rogers DM, Hirst JD, Lee EPF, Wright TG. Ab initio study of the toluene dimer. Chem Phys Lett. 2006; 427(4-
6):410–413. doi: 10.1016/j.cplett.2006.07.022.

Salomon-Ferrer R, Case DA, Walker RC. An overview of the Amber biomolecular simulation package. Wiley
Interdiscip Rev Comput Mol Sci. 2013; 3(2):198–210.

Schuster BS, Dignon GL, Tang WS, Kelley FM, Ranganath AK, Jahnke CN, Simpkins AG, Regy RM, Hammer
DA, Good MC, Mittal J. Identifying sequence perturbations to an intrinsically disordered protein that
determine its phase-separation behavior. Proc Natl Acad Sci USA. 2020; 117(21):11421–11431. doi:
10.1073/pnas.2000223117.

Seeliger D, de Groot BL. Protein Thermostability Calculations Using Alchemical Free Energy Simulations. Bio-
phys J. 2010; 98(10):2309–2316. doi: 10.1016/j.bpj.2010.01.051.

Semenov AN, Rubinstein M. Thermoreversible gelation in solutions of associative polymers. 1. Statics. Macro-
molecules. 1998; 31(4):1373–1385.

Serrano L, Bycroft M, Fersht AR. Aromatic-aromatic interactions and protein stability: investigation by double-
mutant cycles. J Mol Biol. 1991; 218(2):465–475.

Shin Y, Brangwynne CP. Liquid phase condensation in cell physiology and disease. Science. 2017;
357(6357):eaaf4382. doi: 10.1126/science.aaf4382.

Tang Y, Bera S, Yao Y, Zeng J, Lao Z, Dong X, Gazit E, Wei G. Prediction and characterization of liquid-liquid phase
separation of minimalistic peptides. Cell Rep Phys Sci. 2021; 2(9):100579. doi: 10.1016/j.xcrp.2021.100579.

Tesei G, Schulze TK, Crehuet R, Lindorff-Larsen K. Accurate model of liquid-liquid phase behavior of intrin-
sically disordered proteins from optimization of single-chain properties. Proc Natl Acad Sci U S A. 2021;
118(44):e2111696118. doi: 10.1073/pnas.2111696118.

Thomas PD, Dill KA. An iterative method for extracting energy-like quantities from protein structures. Proc
Natl Acad Sci USA. 1996; 93(21):11628–11633.

Tsuzuki S, Honda K, Uchimaru T, Mikami M. Ab initio calculations of structures and interaction energies of
toluene dimers including CCSD(T) level electron correlation correction. J Chem Phys. 2005; 122(14). doi:
10.1063/1.1876092.

Urry DW, Gowda DC, Parker TM, Luan CH, Reid MC, Harris CM, Pattanaik A, Harris RD. Hydrophobicity scale
for proteins based on inverse temperature transitions. Biopolymers. 1992; 32(9):1243–1250.

Vernon RM, Chong PA, Tsang B, Kim TH, Bah A, Farber P, Lin H, Forman-Kay JD. Pi-Pi contacts are an overlooked
protein feature relevant to phase separation. eLife. 2018; 7:e31486.

Wang J, Choi JM, Holehouse AS, Lee HO, Zhang X, Jahnel M, Maharana S, Lemaitre R, Pozniakovsky A, Drechsel
D, Poser I, Pappu RV, Alberti S, Hyman AA. A Molecular Grammar Governing the Driving Forces for Phase
Separation of Prion-like RNA Binding Proteins. Cell. 2018; 174(3):688–699.e16.

Wimley WC, Creamer TP, White SH. Solvation energies of amino acid side chains and backbone in a family of
host- guest pentapeptides. Biochemistry. 1996; 35(16):5109–5124.

Wolfenden R, Andersson L, Cullis P, Southgate C. Affinities of amino acid side chains for solvent water. Bio-
chemistry. 1981; 20(4):849–855.

Workman RJ, Pettitt BM. Thermodynamic Compensation in Peptides Following Liquid–Liquid Phase Separation.
J Phys Chem B. 2021; 125(24):6431–6439.

Zheng W, Dignon GL, Jovic N, Xu X, Regy RM, Fawzi NL, Kim YC, Best RB, Mittal J. Molecular Details of Protein
Condensates Probed by Microsecond Long Atomistic Simulations. J Phys Chem B. 2020; 124(51):11671–
11679. doi: 10.1021/acs.jpcb.0c10489.

Appendix 1

Supporting Tables

System	Number of molecules
Cyclohexane	500
Benzene	500
Toluene	500
Methanol	1000
Ethanol	1000
Hexanol	500
Octanol	500
Acetone	1000

Appendix 1—table 1. Number of molecules used in pure solvent simulations for the determination of dielectric constants.

Supporting Figures

Appendix 2—figure 1. (A) Time series for the accessible surface area of the GGXGG in the GSF condensate for different values of λ. (B) Time series for the density profiles of all peptides (blue) and the GGXGG peptide (red) in the coexistence simulations for both values of the switching parameter λ. In the case of $\lambda = 0$, the peptide experiencing the transformation transiently scapes from the core of the condensate. These frames were excluded as initial snapshots of the alchemical transformations.

Appendix 2—figure 2. Work values for the forward and reverse alchemical transitions for different initial snapshots (left) and their distributions (right). The BAR estimator is shown in the legend. Each panel corresponds to a different solvent. (A) TIP3P water. (B) GSY condensates. (C) GSF condensates.

Appendix 2—figure 3. Work values for the forward and reverse alchemical transitions for different initial snapshots (left) and their distributions (right). The BAR estimator is shown in the legend. Each panel corresponds to a different solvent: (A) Acetone. (B) Benzene. (C) Cyclohexane. (D) Ethanol. (E) Hexanol. (F) Methanol. (G) Octanol. (H) Toluene.