Integrating genetic and gene expression data in network-based stratification analysis of cancers

Kenny Liou[1*], Ji-Ping Wang[1**]

1 Department of Statistics and Data Science, Northwestern University, Evanston, Illinois, United States of America

* kliou@ini.usc.edu ** jzwang@northwestern.edu

Abstract

Cancers are complex diseases that have heterogeneous genetic drivers and varying clinical outcomes. A critical area of cancer research is organizing patient cohorts into subtypes and associating subtypes with clinical and biological outcomes for more effective prognosis and treatment. Large-scale studies have collected a plethora of omics data across multiple tumor types. These studies provide an extensive dataset for stratifying patient cohorts. Network-based stratification (NBS) approaches have been presented to classify cancer tumors using somatic mutation data. A challenge in cancer stratification is integrating omics data to yield clinically meaningful subtypes. In this study, we integrate somatic mutation data with RNA sequencing data within the NBS framework and investigate the effectiveness of integrated NBS on three cancers: ovarian, bladder, and uterine cancer. We show that integrated NBS subtypes are more significantly associated with overall survival or histology. Integrated NBS networks also reveal highly influential genes that drive cancer initiation and progression. This comprehensive approach underscores the significance of integrating genomic data types in cancer subtyping, offering profound implications for personalized prognosis and treatment strategies.

Introduction

Cancer, a multifaceted disease, exhibits remarkable genetic diversity among patients, rendering the development of effective treatments a formidable challenge. To gain valuable insights into this complexity, large-scale initiatives like the International Cancer Genome Consortium (ICGC) [1] and The Cancer Genome Atlas (TCGA) [2] meticulously profile genomic, transcriptomic, and epigenomic data. This includes DNA methylation, microRNA expression, and protein expression data, unveiling a treasure trove of information. Consequently, the massive amount of data from these projects has spurred a demand for informatics solutions to unearth molecular pathways dictating tumor progression, understanding disease populations, and designing precision-targeted treatment strategies [3].

In cancer informatics, a pivotal objective is to categorize or stratify heterogeneous cancer tumors into clinically and biologically meaningful subtypes using molecular profiling data. Historically, this endeavor primarily leveraged mRNA expression data, yielding valuable insights into cancers like ovarian cancer [4] and breast cancer [5]. However, it has fallen short in other cancer types, such as colorectal cancer, where the association between molecular subtypes and clinical phenotypes remains elusive [6]. This limitation may be attributed to issues like sample quality and overfitting, inherent to gene expression analysis [7].

As the landscape of genome-scale data diversifies and expands, there emerges an urgent need to develop methodologies that effectively integrate multi-omics data for more robust tumor subtyping [8]. Multi-omic research in cancer genetics offers promising results, such as in breast cancer [9, 10]. Typically, multi-omic integration methods synthesize data after processing -omics data types separately or using different -omics data types for different parts of the methodology pipeline [8]. The advantage of multi-omics is that the potential interactions between various molecular layers are considered, providing a holistic perspective that can lead to different results compared to single-data-type methods [11]. In this paper, we integrate molecular profiles before processing these data types to explore if pre-process integration can yield meaningful results.

Recent years have witnessed the widespread adoption of gene interaction networks,

with cancer being increasingly recognized as a network-driven disease [12]. Somatic   ³²
mutation profiles harbor cancer-driver genes capable of instigating genetic mutations in   ³³
other genes. Network-based Stratification (NBS) [13] is an approach that melds gene   ³⁴
interaction networks with somatic mutation profiles. This fusion involves mapping   ³⁵
somatic mutation profiles onto a cancer network and propagating these mutations   ³⁶
throughout the network to create smoothed network profiles. NBS uses these profiles to   ³⁷
stratify tumors by clustering patients with similar smoothed network profiles.   ³⁸

In line with these trends in cancer genomics, we propose a multi-omics methodology   ³⁹
grounded in NBS for tumor stratification, fusing somatic mutation profiles with RNA   ⁴⁰
gene expression data. We fuse somatic mutation and gene expression profiles before   ⁴¹
applying network propagation to the integrated profiles to explore if integrating profiles   ⁴²
before network smoothing can generate informative cancer subtypes. By integrating   ⁴³
these two data types before network smoothing, we have successfully generated robust,   ⁴⁴
biologically significant, and clinically informative tumor subtype clusters. Our   ⁴⁵
methodology was tested using the TCGA ovarian, uterine, and bladder carcinoma   ⁴⁶
cohorts, yielding compelling results that can shed new light on understanding cancer   ⁴⁷
population structures through an integrated approach.   ⁴⁸

Materials and methods   ⁴⁹

Integrating genetic profiles   ⁵⁰

Ovarian serous adenocarcinoma, uterine endometrial carcinoma, and bladder urothelial   ⁵¹
carcinoma somatic mutation and gene expression data were downloaded from the   ⁵²
TCGA data portal in June 2023. Only individuals with somatic mutation and gene   ⁵³
expression data were retained for the experiment. 279 ovarian cancer, 318 uterine   ⁵⁴
endometrial carcinoma, and 399 bladder urothelial carcinoma patients from the TCGA   ⁵⁵
cohorts were used in subsequent analysis. Somatic mutation profiles are binary vectors   ⁵⁶
where a '0' or '1' for whether a gene for that individual has no mutation or a mutation,   ⁵⁷
respectively. Gene expression profiles are continuous TPM normalized values where the   ⁵⁸
value represents the level of gene expression. The gene expression profiles were min-max   ⁵⁹
normalized gene by gene to match the 0 to 1 range of somatic profiles. Somatic mutation   ⁶⁰

and gene expression profile integration can be represented through this formula:

$$S_i = \beta \times p_i + (1 - \beta) \times q_i \tag{1}$$

where $0 < \beta < 1$ is a tuned hyperparameter chosen by the user to linearly combine the somatic mutation profile p_i and the normalized gene expression profile q_i to result in the integrated profile S_i for individual i. In the integration process of NBS for ovarian cancer, bladder cancer, and uterine cancer, we utilized tuned β values of 0.8, 0.3, and 0.1, respectively. The value of β for ovarian and bladder cancers were chosen to give the most significant p-value in the log-rank test from Kaplan Meier survival analysis or the log-likelihood ratio test from Cox regression survival analysis. Survival analysis for uterine cancer was impeded by the notably low mortality rate within the cohort, alongside the absence of significant associations with survival observed among both single-data type and integrated subtypes. To assess the association of uterine cancer subtypes, we examined the association of TCGA subtypes to our generated Multi-NBS subtypes (χ^2 association test statistic) to derive the value of β. Additional details about TCGA are provided at www.cancer.gov/ccg/research/genome-sequencing/tcga.

Gene interaction network

The constructed network is derived from PCNet [16], a network with 19,781 genes and 2,724,724 interactions that is then filtered for cancer-specific genes and interactions in at least one of these four sources: [17–20]. The resulting cancer subnetwork contained 2291 nodes. This network was used for all three methods of NBS. Further details on constructing this network can be found in a previous NBS publication [15].

Network propagation

The integrated profiles are mapped onto the gene interaction network and then network propagation [44] is applied to diffuse the signals across the network. Let m be the number of genes, n be the number of patients, F_0 be the initial $n \times m$ (patient $\times$ gene matrix), and A be the symmetric adjacency matrix ($m \times m$) representing the gene-gene interaction network obtained above. Network propagation follows the following iterative

procedure:

$$F_{t+1} = \alpha F_t A + (1 - \alpha) F_0 \qquad (2)$$

where $\alpha = 0.7$, which is derived from benchmarking results reported in earlier NBS

publications [13]. The network is propagated till F_t converges ($|F_{t+1} - F_t < 0.001|$).

After convergence, the resulting matrix F_t was quantile normalized by row (patient) to

ensure each patient followed the same distribution. F represents the final normalized

and smoothed integrated matrix.

Network-regularized NMF

Non-negative matrix factorization (NMF) decomposes a matrix into two non-negative

matrices whose product results in the original matrix. Network-regularized NMF is an

extension of NMF that constrains NMF to respect the network structure [45–47]. The

following objective function is minimized:

$$min_{W,H>0}\big\{\|F - WH\|^2 + trace(W^t JW)\big\} \qquad (3)$$

F is approximately decomposed into the product of non-negative matrices W (m by K

matrix) and H (K by n matrix). W is a collection of basis vectors ("meta-genes") and

H represents the basis loadings. K controls the dimension reduction and we used values

of $K = 2, 3, 4, 5, 6, 7, 8$ in the following context. The $trace(W^t JW)$ function is

responsible for constraining the basis vectors in W to respect the local neighboring

network structure. J represents the graph Laplacian of the k-nearest neighbor network

and we used $k = 11$ as described in previous work [13].

Consensus clustering

We used consensus clustering [48] to ensure robust clustering to achieve the final patient

cluster assignments. We performed network-regulated NMF using a random sampling

without replacement of 80% of patients. This was repeated 100 times as described

previously [13]. The collection of 100 clustering results was used to construct a

similarity matrix that recorded the frequency with which patient pairs had the same

cluster assignment from all iterations where both patients in the pair were sampled.

Implementation of Integrated Network-Based Stratification

The implementation of integrated network-based stratification is a Python 3.10 version based on an existing Python 2.7 implementation of NBS [15].

Cluster analysis

We used Silhouette scores [21] as internal cluster measures. The Silhouette score calculates how well a patient is affiliated with its cluster compared to neighboring clusters. We also used Adjusted Mutual Information (AMI) [22] to assess cluster similarity between clusters formed through different data-type generated clusters.

Survival analysis

Survival analysis was performed through the lifelines [49]and scikit-survival package [50]. Kaplan-Meier survival curves [51] were fitted to the subtypes generated and log-rank tests were performed on single-data type and integrated NBS subtypes to assess the association between the subtypes and survival. We also fitted a semi-parametric Cox proportional hazard model [52]. The Cox hazard model can be represented with a hazard function $h(t|X_i)$ at time t for an individual i, given p covariates, denoted by X_i:

$$h(t|X_i) = h_0(t) \exp \sum_{j=1}^{p} \beta_j X_{ij} \qquad (4)$$

We included covariates such as age, race, and gender in addition to the cluster assignments to assess cluster assignment influence on survival. The log-likelihood ratio test compares the full model with subtype assignments and clinical covariates against a null model. The log-likelihood ratio test and the associated p-value provide an estimate of the predictive power of the full model (clinical covariates and subtype assignments) compared to the null model. The concordance index assesses the discriminatory power of the model by evaluating the ability of the model to correctly rank the survival time of pairs of patients. Maximizing the concordance index indicates the model is discriminative of early events (which are associated with higher-risk patients) and later events.

Association with TCGA subtypes

Association with TCGA subtypes is calculated through Pearson's χ^2 test of
independence between computed integrated subtypes or single-data type subtypes and
documented TCGA subtypes. The documented TCGA subtypes are obtained through
the R "TCGAbiolinks" library [23–25], which are determined through genomic,
transcriptomic, and proteomic characterization of tumors using array and
sequence-based technologies by The Cancer Genome Atlas Research Network [26–28].

Identifying high-scoring genes

The integrated profiles were first propagated through the network propagation process
described above. The propagated integrated profiles are then grouped by the subtype
assignments generated using Multi-NBS. After being grouped by subtype the propagated
integrated profiles are averaged by number of patients in each subtype to find genes
with the highest average hybrid score in the network for each integrated profile subtype.
We refer to the score as a "hybrid score" because the network values for integrated
profile networks are a combination of gene expression and somatic mutation data.

For somatic mutation high-scoring genes, the exact same process is applied except
we only used somatic mutation profiles and the subtype assignments generated from
using somatic mutation profile NBS. We first applied propagation to the somatic
mutation profiles. The propagated somatic mutation profiles are grouped by the
subtype assignments given by NBS that used only somatic mutation profiles. After
being grouped, the propagated somatic mutation profiles are averaged by number of
patients in each subtype to find genes with the highest average mutation score in the
network for each somatic mutation profile subtype. Note that we refer to the somatic
mutation scores as mutation scores because the network values are derived from solely
somatic mutation data.

Results

For the three cancer types under consideration, ovarian, uterine, and bladder carcinoma,
only individuals extracted from the TCGA database with both somatic mutation and
gene expression profiles were retained, giving 279 ovarian carcinoma, 318 uterine

endometrial carcinoma, and 399 bladder urothelial carcinoma patients. We did not remove individuals with fewer than 10 somatic mutations. Multi-omic Network-Based Stratification (Multi-NBS) merges somatic mutation profiles with RNA sequencing (RNA-seq) gene expression data through a linear combination to creating an integrated profile (Fig 1). The resulting integrated matrix is then processed through NBS to generate patient subtype assignments. The underlying network used in this context is a sub-network comprised of cancer-related genes, which was constructed from PCNet [16], but filtered using four databases containing cancer-specific genes and interactions [17–20]. To assess the efficacy of the subtypes generated by Multi-NBS, we conduct a rigorous performance evaluation, comparing them against subtypes derived from single-data type NBS. Additionally, we employ benchmarking procedures to identify the optimal β values for integration, ensuring the robustness and informativeness of our results (See Methods).

Fig 1. Integrated NBS workflow. A workflow of an integrated approach to network-based stratification.

In the following we assess the efficacy of Multi-NBS clusters in terms of clustering

metrics, clinical significance in predicting survival time, and association with $\qquad$ 180

histological characteristics by benchmarking with clusters obtained from single data $\qquad$ 181

type. In addition we will investigate whether the identified Multi-NBS clusters can $\qquad$ 182

reveal new high-scoring genes within the networks to that have cancer relevance. $\qquad$ 183

Cluster evaluation $\qquad$ 184

We first evaluated whether the clusters generated by Multi-NBS are comparable to $\qquad$ 185

single-data-type clusters using Silhouette scores [21]. Higher Silhouette scores (values $\qquad$ 186

closer to 1) indicate well-formed clusters while lower scores (values closer to -1) indicate $\qquad$ 187

poorly-formed clusters. Values close to 0 in Silhouette scores are still generally $\qquad$ 188

considered good classification. We plotted the Silhouette scores as a function of the $\qquad$ 189

number of clusters generated under each method (Fig 2). Somatic mutation clusters $\qquad$ 190

appeared to have the poorest formed clusters while RNA-seq and Multi-NBS clusters $\qquad$ 191

appeared to generate slightly more well-formed clusters (Fig 2A and Fig 2B), while $\qquad$ 192

RNA-seq seemed to generate the most well-formed clusters (Fig 2C). The Multi-NBS $\qquad$ 193

clusters did not show a uniformly better pattern than other two. Instead, in ovarian and $\qquad$ 194

bladder cancers, we see that the Multi-NBS Silhouette score curves were similar to $\qquad$ 195

RNA-seq clusters, while in uterine cancer, the Multi-NBS Silhouette score was observed $\qquad$ 196

to be between the two single-data-type Silhouette scores. $\qquad$ 197

We next compared the similarity of cluster contents using Adjusted Mutual $\qquad$ 198

Information (AMI) [22]. AMI is a measure of cluster independence, where a score of 0 $\qquad$ 199

means the clusters are completely independent and a score of 1 means the clusters are $\qquad$ 200

completely identical. We observed that across all cancer types, somatic mutation and $\qquad$ 201

RNA-seq clusters were mostly independent of each other (Fig 3). The Multi-NBS and $\qquad$ 202

RNA-seq clusters showed relatively higher cluster similarity and the AMI values are $\qquad$ 203

0.28, 0.82 and 0.8 respectively for ovarian, bladder and uterine cases, reflecting the $\qquad$ 204

impact of RNA-seq data in the calculation of blended gene profile S_i (see Eq 1) (i.e., $\qquad$ 205

$1 - \beta$ are 0.2, 0.8 and 0.9 respectively). However, we see the AMI scores between $\qquad$ 206

Multi-NBS and somatic mutation clusters are unproportionally lower in all three cases, $\qquad$ 207

possibly because the somatic mutation profile is binary while RNA-seq profile is $\qquad$ 208

continuous such that the integrated profile more resembles the RNA-seq profile. In the $\qquad$ 209

Fig 2. Silhouette score analysis Silhouette scores of single-data-type generated clusters and Multi-NBS generated clusters. Refer to the legend in subfigure (C) for all subfigures. (A) Silhouette scores of ovarian cancer clusters. (B) Silhouette scores of bladder cancer clusters. (C) Silhouette scores of uterine cancer clusters.

following we shall investigate how the blended gene profile may benefit characterization of the cancer subtypes clinically.

Survival analysis

To further assess the generated subtypes, we first examined the Kaplan Meier estimate of the survival function of the generated cancer subtypes, and then a Cox's proportional hazard regression model to investigate how age, gender, and race as covariates may affect to survival in addition to the cluster assignments. We found that uterine cancer subtypes, whether identified through single-data type NBS or integrated NBS, did not show significant associations with survival due to the low mortality rate within the cohort at significance level $\alpha = 0.05$ (to be used throughout the following context). For instance, uterine cancer subtypes derived solely from somatic mutation data for $K = 3$ did not exhibit a significant association with survival (S3 Fig, log-rank test,

Fig 3. Adjusted mutual information scores Adjusted Mutual Information scores of single-data-type generated clusters and Multi-NBS generated clusters. (A) AMI scores for ovarian cancer clusters. (B) AMI scores for bladder cancer clusters. (C) AMI scores for uterine cancer clusters.

$p = 3.837 \times 10^{-1}$). These findings are consistent with previous work [13]. Consequently, further survival analysis for uterine cancer was not pursued.

In ovarian cancer, we first consider $K = 4$ Multi-NBS subtypes, as it is classified into 4 subtypes based on cancer histology. We note that Multi-NBS subtypes were significantly associated with Kaplan Meier survival functions (Fig 4A, log-rank test, $p = 6 \times 10^{-3}$ for $K = 4$). In contrast, subtypes derived from somatic mutation or RNA-seq profiles for $K = 4$ were not found to be significantly associated with survival (S1 Fig, log-rank test, $p = 1.79 \times 10^{-1}$ and $p = 1.423 \times 10^{-1}$, respectively). However, we did observe ovarian cancer subtypes generated from somatic mutation data alone for $K = 2$ are significantly associated with survival, consistent with prior work (S3 Fig, log-rank test, $p = 9.8 \times 10^{-3}$) [13, 14]. The difference in the p-value is mostly likely due to the different samples used in each of these studies.

In the Cox-regression model, we found the log hazard ratio for cluster subtypes and

Fig 4. Ovarian cancer survival analysis Ovarian cancer survival analysis indicate that integrated subtypes are significantly informative of survival. Refer to the legend in subfigure (D) for subfigures (C) and (D). (A) Survival curves indicating the probability of survival for subtypes generated by Multi-NBS (log-rank test, $p = 6 \times 10^{-3}$). (B) The Cox regression log hazard ratios for $K = 4$ integrated subtypes. (C) Association of NBS ovarian cancer subtypes to survival time through the log-rank test. The black line represents $p = 0.05$. (D) The predictive power of generated OV subtypes on survival through the log-likelihood ratio test. The black line represents $p = 0.05$.

Caucasian covariates were all significantly different from zero in Fig 4B. The observed   235
negative log hazard ratio of the three cluster assignments indicated that subtypes 2   236
through 4 had a significantly lower hazard compared to subtype 1. This result   237
confirmed the Kaplan Meier estimates of the survival function, where group 1 showed   238
the shortest survival time. In contrast for the single-data type clusters, we found the log   239
hazard ratio for the subtypes did not significantly differ from zero (S1 Fig).   240

 When we varied K for the subtypes definition, Multi-NBS subtypes exhibited more   241
informative survival patterns across various values of K, demonstrating comparable or   242
superior performance in associating subtypes with survival outcomes compared to   243
subtypes derived from single-data types (Fig 4C). Notably, subtypes derived solely from   244
somatic mutation profiles also show significant associations with Kaplan Meier survival   245

functions, consistent with prior work on somatic mutation NBS subtypes [13]. When we accounted for clinical covariates such as age and race, we found Multi-NBS continued to consistently yield robust subtypes that remained more predictive of survival than somatic mutation or RNA-seq profiles for $K > 2$ (Fig 4D).

For bladder cancer, we found that Multi-NBS subtypes proved highly predictive of patient survival time (Fig 5A, log-rank test, $p = 2 \times 10^{-3}$ for $K = 4$). Notably, the most aggressive bladder cancer Multi-NBS subtype exhibited an average survival time of 1100 days (36 months), while the least aggressive bladder cancer subtype showed an average survival time exceeding 2500 days (82 months). We found subtypes generated using only somatic mutation profiles or RNA-seq profiles were not significantly associated with survival (S2 Fig log-rank test, $p = 1.989 \times 10^{-1}$ and $p = 7.51 \times 10^{-2}$ for $K = 4$, respectively). Log hazard ratios for $K = 4$ show that all cluster assignments coefficients except for the subtype 2 coefficient significantly differ from zero (Fig 5B). Similar to ovarian cancer, we see all single-data type cluster assignment coefficients do not significantly differ from zero, indicating that the subtypes generated from mutation or RNA-seq data alone are less informative of survival when covariates are taken into account (S2 Fig). We also see Multi-NBS BLCA subtypes had comparable predictive power to RNA-seq generated subtypes for survival time even after adjusting for clinical covariates such as age, gender, and race in Fig 5C. This highlights the potential utility and clinical relevance of Multi-NBS in delineating patient subtypes with distinct survival outcomes across cancer types, yielding subtypes that may be more informative of survival compared to single-data type subtypes.

Association with TCGA database subtypes

To further investigate the biological significance, we applied χ^2 association tests with subtypes derived from other data types in the TCGA, including copy-number variation (CNV), methylation, mRNA expression, microRNA expression and protein profiles [23–25]. These TCGA subtypes provided by the TCGA database are determined through analyzing various molecular and histological factors [26–28]. For more information, refer to the Methods section.

For ovarian cancer, we see none of the generated clusters showed a significant

Fig 5. Bladder cancer survival analysis Bladder cancer survival analysis indicate that integrated subtypes are significantly informative of survival. (A) Survival curves indicating the probability of survival for subtypes generated by Multi-NBS (log-rank test, $p = 2 \times 10^{-3}$). (B) Cox regression log hazard ratios for $K = 4$ integrated subtypes. (C) Concordance index of different clusters for bladder cancer subtypes.

association with TCGA subtypes in Fig 6A. This indicates that the single-data type and Multi-NBS subtypes identified are independent of TCGA subtypes. Cluster similarity analysis between Multi-NBS subtypes and TCGA subtypes (AMI $= -0.0013$) also supports Multi-NBS subtype assignments being independent of TCGA subtypes. For bladder cancer, however, we found both TCGA subtypes and Multi-NBS subtypes are significantly associated with survival, though Multi-NBS is significantly associated with TCGA subtypes more consistently than single-data types with log-rank test, $p = 1.16 \times 10^{-2}$ and $p = 4 \times 10^{-4}$, respectively (Fig 6B, S4 Fig). For uterine cancer, we see Multi-NBS subtypes exhibited stronger and more consistent association with TCGA subtypes than single-data type clusters (Fig 6C). We also note that NBS subtypes produced from only somatic mutation profiles are also significantly associated with TCGA subtypes, which is in line with previous findings [13]. Overall, our analysis has

revealed that integrated profile-generated subtypes through NBS are potentially more 288

capable than single-data type NBS subtypes of forming stronger association with TCGA 289

subtypes for bladder cancer and uterine cancer. 290

Fig 6. Association with TCGA subtypes. Association tests with TCGA provided subtypes across various cancers. The dotted line represents $p = 0.05$. Refer to the legend in subfigure (C) for all subfigures. (A) Generated ovarian cancer subtypes association with TCGA ovarian cancer subtypes. (B) Generated bladder cancer subtypes association with TCGA bladder cancer subtypes. (C) Generated uterine cancer subtypes association with TCGA uterine cancer subtypes.

High-scoring genes 291

Next, we sought to identify genes with high scores across subtypes generated from 292

Multi-NBS using network smoothed integrated profiles. Since the network values for 293

integrated profile networks are a combination of gene expression and somatic mutation 294

data, we refer to these scores as hybrid scores. We identified the genes with the highest 295

average hybrid score per individual for each subtype. We then compared these findings 296

to high-scoring genes across NBS subtypes generated from only somatic mutation 297

profiles using network smoothed somatic mutation profiles. Again, we identified the 298

genes with the highest average mutation score per individual for each subtype. We limited our investigation to the top 10 highest scoring genes (see Methods for further information on high-scoring genes identification). Across all three cancers, we found that integrated profiles identified more high-scoring genes shared by all subtypes than somatic mutation profiles. We see there were also genes with high scores unique to a subtype or a subset of subtypes.

Here we investigate the genes shared across all subtypes in integrated profiles that were not identified in somatic mutation only profiles to assess their impact on tumour initiation and progression. In ovarian cancer, we found integrated profiles identified BBC3 and UBC as two genes that had high scores across all Multi-NBS subtypes. We see somatic mutation only subtypes only identified NFKBIZ as a high-scoring gene shared by 3 of the 4 somatic mutation NBS subtypes. We found the BBC3 gene had the highest score across all 4 Multi-NBS subtypes. BBC3, or p53 upregulated modulator of apoptosis, whose expression is modulated by tumor suppressor p53. BBC3 has been identified as a strong marker for indicators of cancer [29]. UBC, which has a role in maintaining ubiquitin homeostasis under stress, also has a high score across all subtypes. Current research indicates that UBC is upregulated in cancers and plays a role in the increased proliferation rate of cancer cells and their ability to overcome cellular stresses introduced by cancer treatments [30]. We list the high-scoring genes for integrated NBS subtypes and somatic mutation NBS subtypes for ovarian cancer in S1 Table and S2 Table, respectively.

In bladder cancer, we observe integrated profiles identified ABCC1, LEPR, LTB4R, and IL1RAP as high-scoring genes across all 4 Multi-NBS subtypes while somatic mutation profiles identified ABCC1 and IL1RAP as high-scoring genes present across all 4 NBS subtypes generated from somatic mutation profiles only (S3 Table and S4 Table). LEPR is understood to be involved in tumour proliferation and migration through regulating ERK1/ERK2 and JAK2/STAT3 expression by interacting with ANXA7 [31]. Recently, there has been literature surrounding the possibility of a variant of LEPR being involved in bladder cancer due to the role of leptin in obese individuals [32]. LTB4R, or leukotriene B4 receptors, are involved in cell proliferation, survival, and metastasis. Previous literature has identified LTB4R as a potential biomarker for tumour sensitivity [33].

In uterine cancer, we found integrated profiles identified IDO1, NCK2, EDAR, and DUSP8 as high-scoring genes across all 3 Multi-NBS subtypes. We observe somatic mutation profiles only identified IDO1 as a high-scoring gene shared by all 3 NBS subtypes generated from somatic mutation profiles. NCK2 is highly involved in pathways mediating cytoskeleton organization and cell proliferation with previous literature indicating NCK2 expression is linked to metastatic human melanoma tumours [34]. EDAR is part for the Tumour Necrosis Factor Receptor (TNFR) superfamily and is a death receptor. Research has identified EDAR as an oncogene for breast cancer, and could be a potential target for investigation in other cancers [35]. DUSP8 is involved in the mitogen-activated protein kinase (MAPK) signaling pathway and has been shown to be tumour progression and drug resistance in various cancers such as colorectal cancer, lung cancer, and breast cancer [36–38]. These previous findings indicate that DUSP8 could be a potential candidate for its involvement in uterine cancer. We list the high-scoring genes for integrated NBS subtypes and somatic mutation NBS subtypes for uterine cancer in S5 Table and S6 Table, respectively.

In summary, we found integrated NBS subtypes exhibit a potentially enhanced capability in identifying highly significant genes shared across all subtypes compared to somatic mutation NBS subtypes. The observed shared high-scoring genes are unique for each type of cancer, as evidenced by our analysis which revealed no overlap in these genes among the three cancer types studied. Moreover, the genes within the integrated networks have a documented role in promoting tumor initiation, progression, and drug resistance, as supported by our investigation of existing literature. This underscores the potential of integrated NBS to pinpoint cancer-specific genes warranting further investigation.

Discussion

In this study, we proposed a method that integrates RNA-seq and somatic mutation profiles before undergoing the NBS process to stratify tumors in an unsupervised manner. Through subsequent analysis, we identified the integrated clusters form comparably separated clusters compared to single-data type clusters. We found this integrated approach is also more effective compared to single-data type NBS in

producing subtypes associated with survival, even after accounting for clinical 361

covariates. We see this integrated approach also yielded subtypes more significantly 362

associated with TCGA provided subtypes. We also found this integrated approach 363

identified key genes with high mutation scores that were specific to a subtype but also 364

other genes that were highly prevalent across all subtypes. 365

The reason Multi-NBS performs well could be largely due to the more holistic nature 366

of integrating somatic mutation and RNA-seq profiles. Somatic mutation profiles are a 367

relative comparison of healthy and tumor tissue, while RNA-seq profiles are an absolute 368

measure of the cell state. Somatic mutations are useful because they contain casual 369

genetic signals driving tumor progression and are differential measures between cancer 370

tumors and normal tissue, which is more capable of capturing tumor-specific alterations. 371

However, the tumor phenotype is often variable even when causal genes are identified, 372

and RNA-seq profiles potentially offer to bridge this gap between phenotype and causal 373

genes by better capturing highly expressed genes. Notably, we see integrated profiles 374

were able to identify more genes that had high network scores across all subtypes within 375

a cancer type. Thus, integrated profiles provide a more comprehensive view of the 376

molecular processes underlying tumor initiation and progression, resulting in the 377

improved stratification we observed. 378

It is crucial to acknowledge that the choice of β values may not remain consistent 379

across all cancers, revealing a potential vulnerability of the method. This variability 380

underscores the importance of tailoring the integration process to the unique 381

characteristics of each cancer type for subtype analysis. In our study, integrating 382

somatic mutation and gene expression profiles involved tuning the hyperparameter β to 383

balance these two data types which led to Multi-NBS subtypes exhibiting improved 384

biological and clinical relevance. However, determining optimal β values proved 385

challenging and varied across cancers. For ovarian, bladder, and uterine cancers, we 386

utilized tuned β values of 0.8, 0.3, and 0.1, respectively, guided by survival analysis 387

metrics or TCGA subtype association metrics. The fluctuation of β across cancers 388

emphasizes the need for careful consideration and adaptation when implementing 389

Multi-NBS for cancer subtyping to suit the specific biological nuances of each cancer 390

type. 391

There are still many methods to integrate different layers of genetic data within the 392

NBS framework. Multiple additional layers of genetic data such as copy-number ⁠393
variation and protein expression data could be integrated similarly to identify clinically ⁠394
and biologically relevant subtypes. Another avenue of exploration is the integration of ⁠395
different genetic data that is beyond a linear combination of genetic profiles. This could ⁠396
include using protein-protein interaction, metabolic, or signaling networks in ⁠397
combination with genetic profiles. Exploration of NBS-generated subtypes through ⁠398
different genetic data integration methods could lead to more clinically and biologically ⁠399
informative subtypes. Investigating these integrated NBS approaches across a wide ⁠400
variety of cancers can lead to a stronger understanding of tumor initiation and ⁠401
progression. ⁠402

Acknowledgments ⁠403

The research reported in this paper was supported by the NSF Quantitative Biology ⁠404
REU Site at Northwestern (DMS-2150134); Northwestern Research Training Grant in ⁠405
Quantitative Biological Modeling, (NSF DMS-1547394); NSF-Simons Center for ⁠406
Quantitative Biology (Simons Foundation/SFARI 597491-RWC and NSF DMS-1764421). ⁠407
The authors gratefully acknowledge the support provided by these grants. ⁠408

References

1. The International Cancer Genome Consortium. International network of cancer genome projects. *Nature.* 2010;464(7291):993-8. doi: `10.1038/nature08987`.

2. The Cancer Genome Atlas Research Network. Integrated genomic analyses of ovarian carcinoma. *Nature.* 2011;474(7353):609-15. doi: `10.1038/nature10166`.

3. Whiteside TL. The tumor microenvironment and its role in promoting tumor growth. *Oncogene.* 2008;27(45):5904-12. doi: `10.1038/onc.2008.271`.

4. Konstantinopoulos PA, Spentzos D, Cannistra SA. Gene expression profiling in epithelial ovarian cancer. *Nat Rev Clin Oncol.* 2008;5(10):577-87. doi: `10.1038/ncponc1178`.

5. Reis-Filho JS, Pusztai L. Gene expression profiling in breast cancer: classification, prognostication, and prediction. *Lancet.* 2011;378(9805):1812-23. doi: `10.1016/S0140-6736(11)61539-0`.

6. The Cancer Genome Atlas Network. Comprehensive molecular characterization of human colon and rectal cancer. *Nature.* 2012;487(7407):330-7. doi: `10.1038/nature11252`.

7. Raspe E. Gene expression profiling to dissect the complexity of cancer biology: pitfalls and promise. *Semin Cancer Biol.* 2012;22(3):250-60. doi: `10.1016/j.semcancer.2012.02.011`.

8. Hamid JS, Hu P, Roslin NM, Ling V, Greenwood CMT, Beyene J. Data Integration in Genetics and Genomics: Methods and Challenges. *Hum Genomics Proteomics.* 2009;2009:869093. doi: `10.4061/2009/869093`.

9. Yang G, Xuequn S, Zhanhuai L. Identification of cancer subtypes by integrating multiple types of transcriptomics data with deep learning in breast cancer. *Neurocomputing.* 2019;324:20-30. doi: `10.1016/j.neucom.2018.03.072`.

10. Qianxing M Xuequn S, Zhanhuai L. Pattern discovery and cancer gene identification in integrated cancer genomic data. *PNAS.* 2013;110(11):4245-50. doi: `10.1073/pnas.1208949110`.

11. Subramanian I, Verma S, Kumar S, Jere A, Anamika K. Multi-omics Data Integration, Interpretation, and Its Application. *Bioinform Biol Insights.* 2020;14:1177932219899051. doi: `10.1177/1177932219899051`.

12. Kreeger PK, Lauffenburger DA. Cancer systems biology: a network modeling perspective. *Carcinogenesis.* 2010;31(1):2-8. DOI: `10.1093/carcin/bgp261`.

13. Hofree M, Shen JP, Carter H, Gross A, Ideker T. Network-based stratification of tumor mutations. *Nat Methods.* 2013;10(11):1108-15. doi: `10.1038/nmeth.2651`.

14. Zhong X, Yang H, Zhao S, Shyr Y. Li B. Network-based stratification analysis of 13 major cancer types using mutations in panels of cancer genes *BMC Genomics.* 2015;16(Suppl 7). doi: `10.1186/1471-2164-16-S7-S7`.

15. Huang JK, Jia T, Carlin DE, Ideker T. pyNBS: a Python implementation for network-based stratification of tumor mutations. *Bioinformatics.* 2018;34(16):2859-61. doi: `10.1093/bioinformatics/bty186`.

16. Huang JK, Carlin DE, Yu MK, Zhang W, Kresiburg JF, Tamayo P et al. Systematic Evaluation of Molecular Networks for Discovery of Disease Genes. *Cell Systems.* 2018;6(4):484-95.e5. doi: `10.1016/j.cels.2018.03.001`.

17. Forbes SA, Beare D, Gunasekaran P, Leung K, Bindal N, Boutselakis H et al. COSMIC: exploring the world's knowledge of somatic mutations in human cancer. *Nucleic Acids Res.* 2015;43(D1). doi: `10.1093/nar/gku1075`.

18. Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell.* 2011;144(5):646-74. doi: `10.1016/j.cell.2011.02.013`.

19. Iorio F, Knijnenburg TA, Vis DJ, Bignell G, Menden MP, Schubert M et al. A landscape of pharmacogenomic interactions in cancer. *Cell.* 2016;166(3):740-54. doi: `10.1016/j.cell.2016.06.017`.

20. Vogelstein B, Papadopoulos N, Velculescu VE, Zhou S, Diaz Jr LA, Kinzler KW. Cancer genome landscapes. *Science.* 2013;339(6127):1546-58. doi: `10.1126/science.1235122`.

21. Rousseeuw PJ. Silhouettes: A graphical aid to the interpretation and validation of cluster analysis. *J Appl Comput Math.* 1987;20:53-65. doi: `10.1016/0377-0427(87)90125-7`.

22. Abraham A, Pedregosa F, Eickenberg M, Gervais P, Mueller A, Kossaifi J et al. Machine learning for neuroimaging with scikit-learn. *Front Neuroinform.* 2014;8:14. doi: `10.3389/fninf.2014.00014`.

23. Colaprico A, Silva TC, Olsen C, Garofano L, Cava C, Garolini D et al. TCGAbiolinks: an R/Bioconductor package for integrative analysis of TCGA data. *Nucleic Acids Res.* 2016;44(8). doi: `10.1093/nar/gkv1507`.

24. Silva TC, Colaprico A, Olsen C, D'Angelo F, Bontempi G, Ceccarelli M et al. TCGA Workflow: Analyze cancer genomics and epigenomics data using

Bioconductor packages. *F1000Research*. 2016;5:1542. doi: `10.12688/f1000research.8923.2`.

25. Mounir M, Lucchetta M, Silva TC, Olsen C, Bontempi G, Chen X et al. New functionalities in the TCGAbiolinks package for the study and integration of cancer data from GDC and GTEx. *PLoS Comput Biol*. 2019;15(3). doi: `10.1371/journal.pcbi.1006701`.

26. The Cancer Genome Atlas Research Network. Integrated genomic analyses of ovarian carcinoma. *Nature*. 2011;474(7351):609-15. doi: `10.1038/nature10166`.

27. The Cancer Genome Atlas Research Network. Comprehensive molecular characterization of urothelial bladder carcinoma. *Nature*. 2014;507(7493):315-22. doi: `10.1038/nature12965`.

28. Levine DA, The Cancer Genome Atlas Research Network. Integrated genomic characterization of endometrial carcinoma. *Nature*. 2013;497(7447):67-73. DOI: `10.1038/nature12113`.

29. Han J, Flemington C, Houghton AB, Gu Z, Zambetti GP, Lutz RJ et al. Expression of bbc3, a pro-apoptotic BH3-only gene, is regulated by diverse cell death and survival signals. *PNAS*. 2001;98(20):11318-23. DOI: `10.1073/pnas.201208798`.

30. Dingding S, Grossman SR. Ubiquitin becomes ubiquitous in cancer. *Cancer Biol Ther*. 2010;10(8):737-47. doi: `10.4161/cbt.10.8.13417`.

31. Huang H, Zhang J, Ling F, Huang Y, Yang M, Zhang Y et al. Leptin Receptor (LEPR) promotes proliferation, migration, and invasion and inhibits apoptosis in hepatocellular carcinoma by regulating ANXA7. *Cancer Cell Int*. 2021;21(4):4. doi: `10.1186/s12935-020-01641-w`.

32. Alfaqih MA, Elsalem L, Nusier M, Mhedat K, Khader Y, Ababneh E. Serum Leptin Receptor and the rs1137101 Variant of the LEPR Gene Are Associated with Bladder Cancer. *Biomolecules*. 2023;13(10):1498. doi: `10.3390/biom13101498`.

33. Kalinkin AI, Nemtsova MV, Zaletaev DV, Sigin VO, Ignatova E, Kuznetsova EB et al. Leukotriene B4 receptors abnormal gene expression is associated with either shorter or longer survival in breast cancer patients depending on the intrinsic tumour molecular subtype. *Ann Oncol.* 2019;30(Suppl 7) doi: `10.1093/annonc/mdz413.061`.

34. Labelle-Cote M, Dusseault J, Ismaïl Salma, Picard-Cloutier A, Siegel PM, Larose L. Nck2 promotes human melanoma cell proliferation, migration and invasion in vitro and primary melanoma-derived tumor growth in vivo. *BMC Cancer.* 2011;11:443. doi: `10.1186/1471-2407-11-443`.

35. Williams R, Jobling S, Sims AH, Mou C, Wilkinson L, Collu GM et al. Elevated EDAR signalling promotes mammary gland tumourigenesis with squamous metaplasia. *Oncogene.* 2022;41(10):1040-9. doi: `10.1038/s41388-021-01902-6`.

36. Zhang H, Wang M, Chen D, Luo C. Dual-specificity phosphatase 8 (DUSP8) induces drug resistance in breast cancer by regulating MAPK pathways. *J Investig Med.* 2022;70(5):1293-300. doi: `10.1136/jim-2021-002282`.

37. Turkowski K, Herzberg F, Günther S, Weigert A, Haselbauer T, Fink L et al. miR-147b mediated suppression of DUSP8 promotes lung cancer progression. *Oncogene.* 2024;43(2):1178-89. doi: `10.1038/s41388-024-02969-7`.

38. Ding T, Zhou Y, Long R, Chen C, Zhao J, Cui P et al. DUSP8 phosphatase: structure, functions, expression regulation and the role in human diseases. *Cell Biosci.* 2019;9:70. doi: `10.1186/s13578-019-0329-4`.

39. Wang X, Yong L. Tumor necrosis factor and cancer, buddies or foes? *Acta Pharmacol Sin.* 2008;29(10):1275-88. doi: `10.1111/j.1745-7254.2008.00889.x`.

40. Katopodis P, Kerslake R, Zikopoulos A, Beri N, Anikin V. p38B - MAPK11 and its role in female cancers. *J Ovarian Res.* 2021;14:84. doi: `10.1186/s13048-021-00834-9`.

41. Huang L, Guo Z, Wang F, Fu L. KRAS mutation: from undruggable to druggable in cancer. *Signal Transduct Target Ther.* 2021;6:386. doi: `10.1038/s41392-021-00780-4`.

42. Wu XY, Liu WT, Wu ZF, Chen C, Liu JY, Wu GN et al. Identification of HRAS as cancer-promoting gene in gastric carcinoma cell aggressiveness. *Am J Cancer Res.* 2016;6(9):1935-48. PMID: 27725900; PMCID: PMC5043104.

43. Pazik M, Michalska K, Żebrowska-Nawrocka M, Zawadzka I, Łochowski M, Balcerczak E. Clinical significance of HRAS and KRAS genes expression in patients with non–small-cell lung cancer - preliminary findings. *BMC Cancer.* 2021;21:130. doi: `10.1186/s12885-021-07858-w`.

44. Vanunu O, Magger O, Ruppin E, Shlomi T, Sharan R. Associating Genes and Protein Complexes with Disease via Network Propagation. *PLoS Comput Biol.* 2010;6(1) doi: `10.1371/journal.pcbi.1000641`.

45. Lee DD, Seung HS. Learning the parts of objects by non-negative matrix factorization. *Nature.* 1999;401(6755):788–91. doi: `10.1038/44565`.

46. Brunet JP, Tamayo P, Golub TR, Mesirov JP. Metagenes and molecular pattern discovery using matrix factorization. *Proc Natl Acad.* 2004;101(12):4164–9. doi: `10.1073/pnas.0308531101`.

47. Cai et al. Non-negative matrix factorization on manifold. *8th IEEE Int. Conf. Data Mining.* 2008; Pisa, Italy. p. 63–72. doi: `10.1109/ICDM.2008.57`.

48. Monti S et al. Consensus clustering: a resampling-based method for class discovery and visualization of gene expression microarray data. *Mach Learn.* 2003;52:91–118. doi: `10.1023/A:1023949509487`.

49. Davidson-Pilon C. lifelines: survival analysis in Python. *J Open Source Softw.* 2019;4(40):1317. doi: `10.21105/joss.01317`.

50. Pölsterl S, Gupta P, Wang L, Conjeti S, Katouzian A, Navab N. Heterogeneous Ensembles for Predicting Survival of Metastatic, Castrate-Resistant Prostate Cancer Patients. *F1000Research.* 2016;5:2676. doi: `10.12688/f1000research.8231.1`.

51. Kaplan EL, Meier P. Nonparametric Estimation from Incomplete Observations. *J Am Stat Assoc.* 1958;53(282):457–81. doi: `10.1080/01621459.1958.10501452`.

52. Andersen PK, Gill RD. Cox's regression model for counting processes: a large sample study. *Ann Stat.* 1982;10(4):1100–20. doi: `10.1214/aos/1176345976`.

Supporting information

S1 Fig.

Ovarian cancer single-data type survival analysis. (A, C) Ovarian cancer single-data type survival curves for $K = 4$ with the log-rank test p-value above each survival curve. Single-data type subtypes are not significantly associated with survival while integrated profile generated subtypes are significantly associated with survival as shown in the paper. (B, D) Ovarian cancer log hazard ratios for single-data type generated clusters. The cluster coefficients are not significantly nonzero indicating the cluster assignment is not informative of higher or lower survival than subtype 1.

Bladder cancer single-data type survival analysis. (A, C) Bladder cancer single-data type survival curves for $K = 3$ with the log-rank test p-value above each survival curve. Single-data type subtypes are not significantly associated with survival while integrated profile generated subtypes are significantly associated with survival as shown in the paper. (B, D) Bladder cancer log hazard ratios for single-data type generated clusters. The cluster coefficients are not significantly nonzero indicating the cluster assignment is not informative of higher or lower survival than subtype 1.

S3 Fig.

Somatic mutation generated subtypes. NBS subtypes using only somatic mutation profiles for ovarian cancer ($K = 2$) and uterine cancer ($K = 3$) with the associated survival log-rank test p-value.

S4 Fig.

TCGA subtypes and Multi-NBS subtype survival curves. Survival curves from ovarian cancer TCGA subtypes and Multi-NBS generated subtypes ($K = 4$). Both are significantly associated with survival through log-rank test.

S1 Table.

Rank	Subtype 1		Subtype 2		Subtype 3		Subtype 4	
	Gene	Score	Gene	Score	Gene	Score	Gene	Score
1	BBC3	0.159	BBC3	0.146	BBC3	0.166	BBC3	0.157
2	UBC	0.127	MAPK11	0.072	UBC	0.110	UBC	0.121
3	CCNB1	0.099	UBC	0.071	JAK3	0.097	RB1	0.097
4	YWHAB	0.099	ORC5	0.068	HRAS	0.096	KRAS	0.091
5	DHX9	0.098	NOTCH3	0.068	GSK3A	0.093	HRAS	0.091
6	HSP90AB1	0.097	EIF3E	0.066	IL6	0.091	LIF	0.091
7	TNF	0.096	PPP3R1	0.066	RB1	0.091	JAK3	0.091
8	RB1	0.095	NAP1L1	0.065	SMC3	0.090	MYL7	0.089
9	TERT	0.095	IFNG	0.064	STAT3	0.090	A2M	0.088
10	HRAS	0.092	PPP2R5B	0.064	KRAS	0.090	FN1	0.088

Integrated ovarian subtypes hybrid scores. The genes with the highest mean hybrid score across the networks of the patients in ovarian cancer subtypes using integrated profiles. BBC3 and UBC are relevant high-scoring genes across all 4 subtypes.

S2 Table.

Rank	Subtype 1		Subtype 2		Subtype 3		Subtype 4	
	Gene	Score	Gene	Score	Gene	Score	Gene	Score
1	NFKBIZ	0.167	NFKBIZ	0.224	NFKBIZ	0.224	MYCL	0.256
2	MCM8	0.035	GTF2H4	0.028	XRCC6	0.021	ACADM	0.254
3	B2M	0.027	REL	0.015	SIN3A	0.021	ERCC3	0.250
4	WNT5B	0.024	KDM6B	0.015	MED12	0.017	CCL17	0.245
5	RANBP2	0.024	MAFB	0.015	WNT5B	0.015	G6PC	0.242
6	CR2	0.022	MAPK11	0.014	ADH4	0.014	PTPRM	0.241
7	PIK3CA	0.021	CXCR4	0.014	AKT2	0.013	CBLC	0.241
8	CACNA2D4	0.021	AOX1	0.014	ITGA5	0.012	PABPC1	0.239
9	GADD45B	0.021	MSH3	0.014	PDCD1LG2	0.012	FGF9	0.238
10	PLK1	0.021	IL17RB	0.014	CR2	0.011	ELF1	0.235

Somatic mutation ovarian subtypes mutation scores. The genes with the highest mean mutation score across the networks of the patients in ovarian cancer subtypes using only somatic mutation profiles profiles. NFKBIZ is a relevant high-scoring genes

across 3 of the 4 subtypes.

S3 Table.

Rank	Subtype 1		Subtype 2		Subtype 3		Subtype 4	
	Gene	Score	Gene	Score	Gene	Score	Gene	Score
1	ABCC1	0.155	ABCC1	0.157	LEPR	0.090	LTB4R	0.087
2	CCNH	0.105	IL1RAP	0.095	ABCC1	0.077	ABCC1	0.079
3	IL1RAP	0.069	LEPR	0.076	LTB4R	0.067	IL1RAP	0.060
4	AFAP1L2	0.055	CCNH	0.075	IL1RAP	0.060	AFAP1L2	0.049
5	IBSP	0.054	LTB4R	0.066	KLK3	0.057	KLK3	0.044
6	PICALM	0.048	AFAP1L2	0.059	KIR2DS5	0.054	LEPR	0.043
7	THBS3	0.045	PICALM	0.050	ACAD8	0.050	NFE2L2	0.041
8	TICAM2	0.045	HLA-E	0.047	HLA-E	0.046	RBM15	0.038
9	LTB4R	0.045	KIR2DS5	0.046	SF3B1	0.045	PICALM	0.038
10	LEPR	0.042	CXCL1	0.046	NTHL1	0.042	ACAD8	0.036

Integrated bladder subtypes hybrid scores. The genes with the highest mean hybrid score across the networks of the patients in bladder cancer subtypes using integrated profiles. ABCC1, LEPR, LTB4R, and IL1RAP are relevant high-scoring genes across all 4 subtypes.

S4 Table.

Rank	Subtype 1		Subtype 2		Subtype 3		Subtype 4	
	Gene	Score	Gene	Score	Gene	Score	Gene	Score
1	TPO	0.112	LEPR	0.112	ITGA2	0.112	ABCC1	0.143
2	NTHL1	0.087	ABCC1	0.110	ABCC1	0.108	LTB4R	0.112
3	FGF18	0.049	IL1RAP	0.068	CCNH	0.091	IL1RAP	0.092
4	LEPR	0.048	ACAD8	0.063	KLK3	0.076	CCNH	0.067
5	SF3B1	0.047	KIR2DS5	0.058	LTB4R	0.074	AFAP1L2	0.053
6	ABCC1	0.046	AFAP1L2	0.057	MAPK12	0.056	HLA-E	0.048
7	MMP1	0.040	KLK3	0.046	NTN4	0.055	CXCL1	0.048
8	PICALM	0.037	PICALM-E	0.043	RFC5	0.047	PICALM	0.043
9	TYROBP	0.033	THBS3	0.042	IL1RAP	0.047	SF3B1	0.042
10	IL1RAP	0.032	HLA-E	0.040	NTHL1	0.046	LEPR	0.041

Somatic mutation bladder subtypes mutation scores. The genes with the highest mean mutation score across the networks of the patients in bladder cancer subtypes using only somatic mutation profiles profiles.. ABCC1 and IL1RAP are relevant high-scoring genes across all 4 subtypes. LEPR is a high scoring gene across 3 of the 4 subtypes.

S5 Table.

Rank	Subtype 1		Subtype 2		Subtype 3	
	Gene	Score	Gene	Score	Gene	Score
1	IDO1	0.148	NCK2	0.203	NCK2	0.165
2	NCK2	0.146	IDO1	0.159	IDO1	0.159
3	ANAPC7	0.125	EDAR	0.129	EDAR	0.136
4	EDAR	0.123	DUSP8	0.117	CDC42	0.111
5	DUSP8	0.083	VWF	0.108	DUSP8	0.107
6	VWF	0.081	MYL12B	0.099	ANAPC7	0.107
7	EGF	0.076	CXCL1	0.084	MYL12B	0.096
8	CD48	0.075	CCNH	0.079	UBC	0.095
9	UBC	0.075	CDC42	0.075	CXCL1	0.095
10	ARID1B	0.074	EGF	0.072	MSH2	0.093

Integrated uterine subtypes hybrid scores. The genes with the highest mean hybrid score across the networks of the patients in uterine cancer subtypes using integrated profiles. IDO1, NCK2, EDAR, and DUSP8 are relevant high-scoring genes across all 3 subtypes.

S6 Table.

Rank	Subtype 1		Subtype 2		Subtype 3	
	Gene	Score	Gene	Score	Gene	Score
1	ANAPC7	0.247	NCK2	0.187	NCK2	0.219
2	IDO1	0.112	IDO1	0.140	EDAR	0.191
3	TNFRSF13C	0.083	MYL12B	0.119	IDO1	0.183
4	ARID1B	0.058	EDAR	0.110	CDC42	0.179
5	CCL8	0.037	DUSP8	0.088	EGF	0.164
6	DUSP8	0.036	VWF	0.066	UBC	0.162
7	IL17RA	0.031	PAX8	0.034	MSH2	0.161
8	CCR9	0.030	CCNH	0.032	CXCL1	0.161
9	CNBP	0.022	CCL8	0.026	YWHAQ	0.147
10	LRP6	0.016	CXCL16	0.025	CHUK	0.146

Somatic mutation uterine subtypes mutation scores. The genes with the highest mean mutation score across the networks of the patients in uterine cancer subtypes using only somatic mutation profiles. IDO1 is a relevant high-scoring gene across all 3 subtypes. NCK2, EDAR, and DUSP8 are high-scoring genes in 2 out of the 3 subtypes.