# Elevated Ubiquitin Phosphorylation by PINK1 Contributes to Proteasomal Impairment and Promotes Neurodegeneration

Cong Chen[1,#], Tong-Yao Gao[1,#], Hua-Wei Yi[4], Yi Zhang[3], Tong Wang[1], Zhi-Lin Lou[1], Tao-Feng Wei[1], Yun-Bi Lu[1], Ting-Ting Li[3], Chun Tang[2,*], Wei-Ping Zhang[1,5,*]

[1] Department of Pharmacology, Zhejiang University School of Medicine, Hangzhou 310058, China

[2] Beijing National Laboratory for Molecular Sciences, College of Chemistry and Molecular Engineering, Center for Quantitate Biology, Center for Life Science, Academy for Advanced Interdisciplinary Studies, Peking University, Beijing 100871, China

[3] Department of Biomedical Informatics, School of Basic Medical Sciences, Peking University Health Science Center, Beijing 100191, China

[4] The First People's Hospital of Jingzhou, First Affiliated Hospital of Yangtze University, Jingzhou 121001, China

[5] Zhejiang Key Laboratory of Precision psychiatry, Hangzhou, 10003, China

[#] Equally contributed

*Corresponding authors

Chun Tang, Tang_Chun@pku.edu.cn, 292 Chengfu Road, Beijing 100871, China, 011-86-010-62751710

Wei-Ping Zhang, weiping601@zju.edu.cn, 866 Yuhangtang Rd, Hangzhou 310058, China, 011-86-0571-88208223

25
**Abstract**

Ubiquitin (Ub), a central regulator of protein turnover, can be phosphorylated by PINK1

(PTEN-induced putative kinase 1) to generate S65-phosphorylated ubiquitin (pUb). Elevated

pUb levels have been observed in aged human brains and in Parkinson's disease, but the

mechanistic link between pUb elevation and neurodegeneration remains unclear. Here, we

demonstrate that pUb elevation is a common feature under neurodegenerative conditions,

including Alzheimer's disease, aging, and ischemic injury. We show that impaired

proteasomal activity leads to the accumulation of sPINK1, the cytosolic form of PINK1 that

is normally proteasome-degraded rapidly. This accumulation increases ubiquitin

phosphorylation, which then inhibits ubiquitin-dependent proteasomal activity by interfering

with both ubiquitin chain elongation and proteasome-substrate interactions. Specific

expression of sPINK1 in mouse hippocampal neurons induced progressive pUb

accumulation, accompanied by protein aggregation, proteostasis disruption, neuronal injury,

neuroinflammation, and cognitive decline. Conversely, *pink1* knockout mitigated protein

aggregation in both mouse brains and HEK293 cells. Furthermore, the detrimental effects of

sPINK1 could be counteracted by co-expressing Ub/S65A phospho-null mutant but

exacerbated by over-expressing Ub/S65E phospho-mimic mutant. Together, these findings

reveal that pUb elevation, triggered by reduced proteasomal activity, inhibits proteasomal

activity and forms a feedforward loop that drives progressive neurodegeneration.

## Introduction

Neurodegeneration, a hallmark of aging and neurodegenerative diseases, is characterized by irreversible loss of neurons, ultimately leading to cognitive and motor impairments (**Wilson et al., 2023**). A common feature of neurodegeneration is the accumulation of ubiquitinated proteins, often resulting from the impairment of protein degradation machineries, particularly the ubiquitin-proteasome system (UPS) (**Davidson and Pickering, 2023; Kinger et al., 2024; McDade et al., 2024**). The UPS is essential for maintaining cellular proteostasis, as it not only eliminates misfolded and toxic proteins but also dynamically regulates neuronal structure and function through controlled protein degradation (**Costa et al., 2019; Krishna et al., 2022; Soykan et al., 2021**). Substrate proteins targeted for proteasomal degradation are typically conjugated with K48-linked polyubiquitin chains, which interact non-covalently with ubiquitin receptors in the proteasome (**Komander and Rape, 2012; Pohl and Dikic, 2019; Swatek and Komander, 2016**). However, UPS activity can be compromised in neurodegeneration in multiple ways, including proteasome modifications, reduced ATP level, direct inhibition by amyloid fibrils, and oxidative stress (**Kinger et al., 2024; McDade et al., 2024; Tseng et al., 2008; Wilson et al., 2023; Wojcik and Di Napoli, 2004**). While these factors have been extensively studied in order to preserve proteasomal function, the dynamic changes of ubiquitin have been largely overlooked in the context of neurodegeneration.

Ubiquitin itself undergoes post-translational modifications that can alter its structure and function (**Herhaus and Dikic, 2015**). One such modification is phosphorylation at residue S65, installed by the kinase PINK1 (PTEN-induced putative kinase 1) (**Koyano et al., 2014**). PINK1 can exist in two forms: the full-length PINK1 localized to the mitochondrial membrane (**Okatsu et al., 2015; Okatsu et al., 2013**), and sPINK1, a cytosolic fragment

cleaved from full-length PINK1 (*Takatori et al.*, **2008**; *Yamano and Youle*, **2013**). Full-length PINK1 is activated in response to mitochondrial damages, subsequently phosphorylating ubiquitin and Parkin to initiate mitophagy—a neuroprotective process for the selective removal of damaged mitochondria (*Koyano et al.*, **2014**; *Lazarou et al.*, **2015**; *Okatsu et al.*, **2015**; *Wilson et al.*, **2023**). However, severe and persistent mitochondrial stress can also lead to mitophagy failure and sustained elevations of S65-phosphorylated ubiquitin (pUb) levels (*Chin et al.*, **2023**; *Pollock et al.*, **2024**). Given the prevalence of mitochondrial dysfunction under neurodegenerative conditions (*Wilson et al.*, **2023**), pUb accumulation may result from the activation of full-length PINK1 but without effective mitophagy.

Alternatively, elevated pUb levels may arise from inhibited proteasomal degradation. The cytosolic sPINK1, processed from the full-length PINK1 by mitochondrial proteases, is normally rapidly degraded by the proteasome via the N-end rule pathway (*Takatori et al.*, **2008**; *Yamano and Youle*, **2013**). As the UPS function is often compromised in neurodegeneration (*Wilson et al.*, **2023**), sPINK1 may accumulate and phosphorylate ubiquitin, contributing to the increased pUb levels.

Elevated levels of pUb have been observed in aged human brain and in Parkinson's disease (*Fiesel et al.*, **2015**; *Hou et al.*, **2018**; *Shiba-Fukushima et al.*, **2017**). While pUb elevation has been proposed as a potential biomarker of neurodegenerative diseases (*Fiesel et al.*, **2015**; *Hou et al.*, **2018**), its functional role remains poorly understood. *In vitro* studies have suggested that phosphorylation alters ubiquitin's structural dynamics and may affect its function (*Dong et al.*, **2017**; *Tang and Zhang*, **2020**; *Wauer et al.*, **2015a**). Moreover, PINK1 phosphorylation has been shown to inhibit ubiquitin chain elongation (*Wauer et al.*, **2015b**), and the phosphomimic mutant Ub/S65E has been show to inhibit protein turnover

and reduce cell viability under stress conditions in yeast (**Swaney et al.**, **2015**). As such,

elevated pUb levels may actively contribute to the progression of neurodegenerative diseases.

In the current study, we demonstrate that pUb levels are elevated in various

neurodegenerative conditions, including Alzheimer's disease (AD), aging, and ischemic

injury. Furthermore, we reveal that elevated pUb levels inhibit proteasomal activity, leading

to protein aggregation and neuronal damages. Upon investigating the underlying mechanism

at molecular, cellular and animal levels, we have uncovered a general pathogenic pathway

involving pUb that operates across a wide spectrum of neurodegenerative disorders.

**Results**

**Elevated pUb levels are a pervasive feature of neurodegeneration**

Elevated pUb levels have been observed in the brains of individuals with Parkinson's

disease (PD) (**Fiesel et al.**, **2015; Hou et al.**, **2018; Shiba-Fukushima et al.**, **2017**). In the

current study, we extend this observation to Alzheimer's disease (AD), the most prevalent

neurodegenerative dementia. We found a marked elevation of both PINK1 and pUb in brain

samples from AD patients in the cingulate gyrus brain regions with Aβ plaques, when

compared to age- and sex-matched controls (**Figure 1A, B; Figure 1—table supplementary**

**1**). This finding was corroborated in the APP/PS1 mouse model of AD, where increased

PINK1 and pUb levels could be detected in neocortex of mouse brains with Aβ compared to

wildtype mice (**Figure 1C, D**). On the other hand, PINK1 and pUb levels in neocortex of

wildtype and *pink1*-knockout mice (*pink1-/-*) differ little (**Figure 1C, D**), which can be

attributed to the low expression of PINK1 in normal mouse brains under physiological

conditions.

Beyond the specific neurodegenerative diseases, pUb elevation also appears to be

associated with aging process (*Fiesel et al.*, **2015**; *Hou et al.*, **2018**), a special pathological

condition that is often accompanied by gradual neurodegeneration. The difference of PINK1

expression between young and aged mouse cortex was not distinguishable due to the low

expression level and probably also due to specificity of PINK1 antibody used. However, we

observed a significant increase in neuronal pUb levels in aged wildtype mouse compared to

young mouse in neocortex of the brains (*Figure 1E, F*). As a control, pUb levels in *pink1*-/-

mice remained unchanged with age and were notably lower than those in aged wild-type

mice (*Figure 1E, F*). As such, although the pUb antibody may not be highly specific, the

observed increase in immunofluorescence and protein band intensities, when comparing

across different *pink1* genetic background, can only be attributed to elevated pUb levels

during aging.

We further showed that acute neurodegenerative conditions, such as cerebral ischemia,

are also associated with elevated pUb levels. Using mouse middle cerebral artery occlusion

(MCAO) model, we observed a marked increase in both PINK1 and pUb levels in the

penumbra of mouse brains compared to the contralateral cortex (*Figure 1G, figure 1—figure*

*supplementary 1*). To further explore the relationship between pUb and ischemic stress, we

subjected HEK293 cells to oxygen-glucose deprivation (OGD), a cellular model mimicking

ischemic conditions. OGD followed by reperfusion caused a time-dependent increase in

PINK1, sPINK1, and pUb levels (*Figure 1H*). Additionally, we observed an increase of

protein aggregation, as evidenced by the accumulation of ubiquitin in the insoluble protein

fraction (*Figure 1I*).

Taken together, these findings demonstrate that elevated pUb levels are a common

feature across a wide spectrum of neurodegenerative conditions, both chronic and acute.

While our observation support the notion of pUb as a biomarker, the diverse functions of ubiquitin suggest that elevated pUb levels may actively contribute to the pathogenesis of neurodegeneration.

**Figure 1. Elevated pUb levels are widespread across neurodegenerative conditions.**

**(A, B)** Double immunofluorescence staining showing the distribution of PINK1 and Aβ (A), and pUb and Aβ (B), within the cingulate gyrus brain region of AD patients compared to similar brain regions from age-matched controls. Detailed donor information is provided in ***Supplementary Table 1***.

**(C, D)** Double immunofluorescence staining of PINK1 and Aβ (C), and pUb and Aβ (D) in the brains of wildtype, *pink1-/-*, and APP/PS1 transgenic mice. The images were taken for the neocortex of APP/PS1, wildtype, and *pink1-/-* mice.

**(E)** Double immunofluorescence staining for pUb and the neuronal marker NeuN in the neocortex of young and aged brains of both wildtype and *pink1-/-* mice. The images were taken for the layer III-IV of neocortex.

**(F)** Western blot analysis of pUb levels in the cortex of young (Y) and aged (A) wildtype and *pink1-/-* mouse brains, quantitatively comparing protein levels across ages and genotypes. *$P<0.05$, **$P<0.01$ for comparisons with young wildtype mice; [###]$P<0.001$ for comparisons with aged wildtype mice, one-way ANOVA.

**(G)** Immunofluorescence staining for PINK1 and pUb in the contralateral and penumbra of mouse brains subjected to MCAO for 2 hours, followed by 24 hours of reperfusion. Locations of the analyzed brain regions are shown in ***Figure 1-supplemary figure 1***.

**(H)** Western blot analysis of PINK1 and pUb levels in HEK293 cells subjected to OGD for 2 hours, followed by reperfusion at 0, 6, and 12 hours.

**(I)** Western blot analysis of ubiquitin in the insoluble fraction of HEK293 cells post 2-hour OGD and subsequent 0, 6, 12 hours of reperfusion.

## Ubiquitin phosphorylation by sPINK1 impairs proteasomal function in HEK293 cells

Elevated pUb levels can result from either elevated level of full-length PINK1 or sPINK1. PINK1 increase can occur upon mitochondrial injuries with the treatment of CCCP and O/A, and sPINK1 increase can occur upon proteasomal inhibition with the treatment of MG132 (***Figure 2A***). The treatment of MG132 caused a concentration- and time-dependent increase in sPINK1 and pUb levels in wildtype HEK293 cells, with sPINK1 levels plateaued at 6 hours (***Figures 2B-E***). In contrast, PINK1 and pUb levels remained low in *pink1-*knockout HEK293 cells upon the treatment of MG132 (***Figures 2B-E***).

While PINK1 activation typically involves its dimerization at mitochondrial outer membrane (***Gan et al.*, 2022; *Okatsu et al.*, 2013**), we investigated whether cytoplasmic sPINK1 retains activity of phosphorylating ubiquitin. We transiently transfected HEK293

cells with several PINK1 constructs: full-length PINK1, a cytoplasmic variant of sPINK1 that

is more resistant to UPS degradation (sPINK1*, PINK1/F101M/L102-L581, differing from

the native sequence by the N-terminal Met instead of Phe), a kinase-dead version of this

stable variant (sPINK1*-KD, with K219A/D362A/D384A mutations introduced) (**Beilina et**

**al.**, **2005**), and Ub$^{GG}$-sPINK1, a short-lived version of native-sequence sPINK1 with a

ubiquitin appended at the N-terminus (**Yamano and Youle**, **2013**). Western blot analysis

confirmed the expression of PINK1, sPINK1* and sPINK1*-KD following transfection,

while Ub$^{GG}$-sPINK1 was undetectable due to its rapid degradation (**Figure 2F**). The analysis

also showed elevated pUb levels upon the transfection of full-length PINK1 and sPINK1*,

with the latter more pronounced. In contrast, transfection with sPINK1*-KD or Ub$^{GG}$-

sPINK1 had little effect on pUb levels (**Figure 2F**).

Furthermore, immunofluorescence staining revealed a diffuse cytoplasmic ubiquitin

distribution in non-transfected or eGFP-transfected control cells. In contrast, ubiquitin-

positive puncta were observed following MG132 treatment or upon the transfection of

sPINK1* (**Figure 2G**). Weak ubiquitin-positive puncta were also visible in Ub$^{GG}$-sPINK1-

transfected cells (**Figure 3G**, 5$^{th}$ row, indicated by arrows), likely due to a slight elevation in

sPINK1 level. Furthermore, Western blot analysis showed that MG132 treatment and

sPINK1* transfection (**Figure 2H**), but not sPINK1*-KD or Ub$^{GG}$-sPINK1 transfection,

increased ubiquitin levels in the insoluble protein fraction. These findings suggest a positive

correlation between PINK1 and pUb levels and the aggregation of ubiquitinated proteins,

which likely results from an impairment of proteasomal activity, akin to the effect of MG132.

To confirm the inhibitory effect of pUb on proteasomal activity, we transfected cells

with Ub-R-GFP, a model substrate for the proteasome (**Dantuma et al.**, **2000**; **Samant et al.**,

**2018**). GFP was rapidly degraded by functional proteasomes but accumulated upon the

administration of MG132. Transfection with sPINK1* also caused GFP accumulation, but the

transfection of sPINK1*-KD did not (**Figure 2I**). Thus, sPINK1* transfection leads to

increased pUb levels and impairs proteasomal degradation.

Elevated pUb levels may also lead to protein aggregation by inhibiting autophagic

degradation. To assess the involvement of this pathway, we treated HEK293 cells with

puromycin to increase ubiquitinated protein level, which can be degraded through both

autophagic and UPS pathways (**Figure 2I**). The ubiquitin signal was significantly lower in

the sPINK1*-transfected cells compared to neighboring non-transfected cells (**Figure 2J**),

indicating that sPINK1* promotes the degradation of ubiquitinated proteins. However, when

we blocked autophagic degradation using BALA (**Mauvezin and Neufeld, 2015**), ubiquitin

signal was higher in the sPINK1*-transfected cells compared to non-transfected cells (**Figure**

**2J**), indicating sPINK1* inhibits the degradation of ubiquitinated protein. Taken together, the

findings demonstrate that while sPINK1* can enhance autophagic degradation of

ubiquitinated proteins, potentially through p62 phosphorylation as previously reported (**Gao**

**et al., 2016**), elevated PINK1 and pUb levels cause an inhibition of proteasomal degradation.

**Figure 2. Ubiquitin phosphorylation by sPINK1 affects proteasomal activity in HEK293 cells.**

**(A)** Representative Western blot showing the levels of PINK1 following the administration of CCCP, O/A, or MG132. The CCCP was treated for 12 hours, O/A was treated for 2 hours, and the MG132 was treated for 8 hours.

**(B)** Western blot analysis showing the concentration-dependent effect of MG132 (0-5 µM) over an 8-hour period on PINK1 level. $N$=3; *$P < 0.05$, ***$P < 0.001$, compared to 0 µM MG132, one-way ANOVA.

**(C)** Western blot analysis showing the time-dependent effect of 5 µM MG132 on PINK1 levels in 0-24 hours. $N$=4; *$P < 0.05$, ***$P < 0.001$, compared to 0 hours, one-way ANOVA.

**(D)** Western blot analysis of pUb levels under a concentration gradient of MG132 (0-5 µM) for 8 hours. $N$=3; *$P < 0.05$, compared to 0 µM MG132, one-way ANOVA.

**(E)** Western blot analysis of pUb levels over a time course of 0-24 hours with 5 µM MG132 treatment. $N$=3; *$P < 0.05$, compared to 0 hours, one-way ANOVA.

**(F)** Representative Western blot image showing the levels of pUb and PINK1 at 24 hours post-transfection with different PINK1 constructs: sPINK1* (PINK1/F101M/L102-L581), sPINK1*-KD

(kinase-dead sPINK1* with additional K219A/D362A/D384A mutations), and Ub<sup>GG</sup>-sPINK1 (a
short-lived version of native sPINK1 with an appended N-terminal Ub).

**(G)** Representative immunofluorescence images of ubiquitin staining in cells treated with 5μM
MG132 for 8 hours or transfected with different sPINK1 constructs, highlighting differences in
ubiquitin localization and aggregation. The white arrows indicate positively transfected cells.

**(H)** Representative Western blot image of ubiquitin in the insoluble protein fraction of cells. The cells
were collected at 8 hours after treatment with 5 μM MG132, or collected at 24 hours after transfection
with different sPINK1 constructs.

**(I)** Western blot analysis showing GFP degradation in HEK293 cells transfected with Ub-R-GFP.
Cells were harvested at 8 hours after 5 μM MG132 treatment, or 24 hours following sPINK1* or
sPINK1*-KD transfection. $N$=3; *$P < 0.05$, ***$P < 0.001$, compared to the control, one-way ANOVA.

**(J)** Immunofluorescence staining of ubiquitin illustrating how sPINK1 over-expression impacts on
proteasomal and autophagic degradation. Puromycin blocks protein synthesis at the translation
elongation stage, leading to the production of truncated proteins; BALA (bafilomycin A1, a v-ATPase
inhibitor) blocks the degradation of autophagic cargo by inhibiting autophagosome-lysosome fusion,
as illustrated in the left panel. Puromycin (5 μg/ml) was applied for 2 hours before harvesting with or
without the treatment of 0.1 μM BALA. The white arrows indicate positively transfected cells.

**pUb impairs proteasomal activity through both covalent and noncovalent mechanisms**

To elucidate how elevated pUb levels inhibit UPS activity, we first assessed ubiquitin

chain formation *in vitro*. While the unmodified ubiquitin efficiently formed di-, tri-, and

higher-order chains catalyzed by E1 and E2 enzymes, pUb predominantly formed di-Ub

chains (***Figure 3A***), consistent with the previous report  (***Wauer et al.***, **2015b**). The

introduction of S65A (phospho-null) and S65E (phospho-mimetic) ubiquitin mutants further

confirmed the impact of phosphorylation—Ub/S65A mutation did not affect chain elongation,

producing a polyUb ladder similar to unmodified ubiquitin, while Ub/S65E mutation only

formed di-Ub (***Figure 3A***).

Furthermore, we examined ubiquitin chain formation in HEK293 cells by expressing

FLAG-Ub/48K, a ubiquitin mutant that can only form K48-linked ubiquitin chain. Upon

proteasomal inhibition with MG132, K48-linked chains were observed in wildtype and

*pink1-/-* cells, but were absent in cells over-expressing full-length PINK1 or sPINK1*

(*Figure 3B*), This further confirms the inhibitory effect of S65 phosphorylation on ubiquitin

chain elongation.

Since PINK1 can phosphorylate both ubiquitin monomers and polyUb chains (*Shiba-*

*Fukushima et al.*, **2014**), we next investigated how polyUb phosphorylation impacts

proteasomal degradation. Using an *in vitro* degradation assay using purified proteasomes, we

found that the degradation of K48-linked pUb-chain-modified GFP (pK48-polyUb-GFP) was

slower compared to its unmodified counterpart (K48-polyUb-GFP) (*Figure 3C, figure*

*3—figure supplementary 1, 2*). Additionally, we employed total internal reflection

fluorescence (TIRF) microscopy, following an established protocol (*Lu et al.*, **2015**), to

assess interactions between ubiquitinated substrate proteins and immobilized proteasomes.

The assay revealed fewer GFP puncta for pK48-polyUb-GFP compared to K48-polyUb-GFP

(*Figure 3D, E, figure 3—figure supplementary 3*). Furthermore, the GFP puncta for pK48-

polyUb-GFP dissociated more rapidly from the proteasomes, reflecting a shorter dwell time

of the proteasome-substrate complex (*Figure 3F-I, figure 3—figure supplementary 4*). Thus,

phosphorylation impairs the noncovalent interactions between polyUb chains conjugated to

substrate proteins and the ubiquitin receptors residing in the proteasome.

Together, these findings demonstrate that elevated pUb levels disrupt proteasomal

function through two distinct mechanisms: impairing covalent ubiquitin chain elongation and

substrate ubiquitination, as well as weakening non-covalent interactions between

ubiquitinated substrates and the proteasome. Both mechanisms collectively contribute to the

inhibition of ubiquitin-dependent proteasomal activity.

**Figure 3. Ubiquitin phosphorylation inhibits both ubiquitin chain elongation and ubiquitin proteasome interactions.**

**(A)** Coomassie blue staining showing ubiquitin chain formation/elongation using different ubiquitin variants as building blocks: wildtype ubiquitin, phosphorylated ubiquitin (pUb), phospho-null Ub/S65A, and phospho-mimic Ub/S65E.

**(B)** Western blot showing the formation of K48-linked ubiquitin chain in wildtype (WT) and *pink1-/-* HEK293 cells without or with PINK1 and sPINK1* transfection. Cells were transfected with FLAG-tagged Ub/48K to form K48-linked ubiquitin chain only, and 10 µM MG132 was applied at 12 hours post-transfection to prevent the degradation of ubiquitinated substrates.

**(C)** Western blot analysis of *in vitro* proteasomal degradation of GFP modified with K48-linked ubiquitin chains (K48-polyUb-GFP) and GFP modified with phosphorylated K48-linked ubiquitin chains (pK48-polyUb-GFP). $N$=3; **$P$<0.01, ***$P$<0.001 compared to respective controls without added proteasome; [#]$P$<0.05, [##]$P$<0.01, [###]$P$<0.001 compared to K48-polyUb-GFP with proteasome, two-way ANOVA.

**(D)** Representative TIRF microscopy images showing single-molecule association of K48-polyUb-GFP (left) and pK48-polyUb-GFP (right) to the surface-immobilized proteasomes, visualized as bright puncta. Details of TIRF single-molecule visualization is shown in ***Figure 3—figure supplementary 3***.

**(E)** Quantitative analysis of puncta density from TIRF images comparing the number of puncta of K48-polyUb-GFP and pK48-polyUb-GFP associated with surface-immobilized proteasomes. $N$=4; ***$P$<0.001 compared to K48-polyUb-GFP, using a paired *t*-test.

**(F, H)** Representative fluorescence time traces of a single punctum for proteasomal bind K48-polyUb-GFP (F) and pK48-polyUb-GFP (H). More representative figures of single-molecule fluorescence are shown in in ***Figure 3—figure supplementary 4***.

**(G, I)** Analysis of GFP fluorescence dwell time for K48-polyUb-GFP (G) and pK48-polyUb-GFP (I) associated with the proteasome. Data were binned and modeled with a single exponential decay curve (red line).

**312** *pink1* **knockout mitigates protein aggregation under neurodegenerative conditions**

**313**     Having established the causative link between elevated pUb levels and proteasomal

**314** inhibition, we further explored the relationship with protein aggregation and whether *pink1*

**315** knockout mitigates this process. We treated wildtype and *pink1-/-* HEK293 cells with MG132,

**316** which increased insoluble ubiquitin levels in both cell types in a concentration- and time-

**317** dependent manner (*Figure 4A, B*). However, insoluble ubiquitin levels accumulated

**318** significantly slower in *pink1-/-* cells, though eventually reaching levels comparable to those

**319** in wildtype cells (*Figure 4B*). This not only confirms that PINK1/pUb operates within the

**320** same pathway as MG132—the proteasome—but also demonstrates that PINK1 deficiency

**321** delays the accumulation of protein aggregates. MG132 treatment also elevated the levels

**322** LC3-II, a marker of autophagy, in wildtype cells, indicating either autophagy activation or

**323** reduced LC3 degradation in response to proteasomal inhibition (*Figure 4C, D*). However,

**324** LC3-II levels did not increase in *pink1-/-* cells following MG132 treatment (*Figure 4C, D*).

**325** This confirms the critical role of PINK1 in the activation of for autophagy (*Figure 2J*).

**326**     Furthermore, *pink1* knockout mitigated protein aggregation in animal models.  In young

**327** wildtype mouse brains, ubiquitin was evenly distributed in the neurons in neocortex, but

**328** formed ubiquitin positive puncta in aged wildtype mouse brains (*Figure 4E*). Western blot

**329** analysis revealed elevated levels of insoluble ubiquitin in the cortex of aged mice compared

**330** to young mice (*Figure 4F*), further supporting age-related protein aggregation. In contrast,

**331** aged *pink1-/-* mice showed no such age-related changes in neuronal ubiquitin distribution or

**332** insoluble ubiquitin levels (*Figure 4E, F*). Following MCAO, insoluble ubiquitin levels were

**333** significantly elevated in the ipsilateral hemisphere of wildtype mice compared to sham and

**334** contralateral hemisphere, but remained unchanged in *pink1-/-* mice (*Figure 4G*).

Protein aggregation is a hallmark of aging and neurodegenerative diseases such as

AD (***Hegde et al.***, **2023**); it arises from perturbed proteostasis and can impair proteasomal

activity in return. Acute proteasomal inhibition can also result from ischemia or

pharmacological treatments. Since sPINK1 is rapidly degraded by functional proteasomes,

impaired proteasomes can lead to the accumulation of sPINK1 and pUb, leading to further

proteasomal inhibition. Together, our results reveal that *pink1* knockout disrupts this cycle,

thereby alleviating proteasomal inhibition and mitigating protein aggregation under these

conditions, further demonstrating the critical role of pUb in this process.

**Figure 4. *pink1* knockout mitigates protein aggregation upon proteasomal inhibition.**

**(A)** Western blot analysis of ubiquitin levels in the insoluble protein fraction of HEK293 cells following the treatment with 0-5 μM MG132 for 8 hours. $N$=3; ***$P$<0.001 compared to 0 μM MG132; ##$P$<0.01 compared with wildtype cells, one-way ANOVA.

**(B)** Western blot analysis of ubiquitin levels in the insoluble protein fraction of HEK293 cells following the treatment with 5 μM MG132 over 24-hour period. $N$=4; *$P$<0.05, **$P$<0.01, ***$P$<0.001 compared to 0 μM MG132; ##$P$<0.01 compared with wildtype cells, one-way ANOVA.

**(C)** Western blot analysis of LC3 levels in HEK293 cells following the treatment with 0-5 μM MG132 for 8 hours. $N$=4; *$P$<0.05 compared to 0 μM MG132, one-way ANOVA.

**(D)** Western blot analysis of LC3 levels in HEK293 cells following the treatment with 5 μM MG132 treatment over 24 hours. $N$=3; *$P$<0.05 compared to 0 μM MG132, one-way ANOVA.

**(E)** Immunofluorescence staining of ubiquitin taken from layer III-IV in the neocortex of brains from young and aged wildtype and *pink1-/-* mice.

**(F)** Western blot analysis quantifying ubiquitin in the insoluble protein fraction of the cortex from young and aged mouse brains. $N$=4; **$P$<0.01, one-way ANOVA.

**(G)** Western blot analysis of ubiquitin in the insoluble protein fraction of wildtype and *pink1-/-* mouse brains subjected to MCAO for 2 hours followed by 24 hours of reperfusion. Comparison includes the sham-operated group (same procedure without occlusion). Locations of the analyzed regions (contralateral and ipsilateral) are indicated in ***Figure 1—figure supplementary 1***.

## sPINK1* over-expression causes cumulative proteome-wide changes

Given the critical role of ubiquitin phosphorylation in proteasomal inhibition and protein aggregation, we sought to confirm these effects using phosphomimetic Ub/S65E mutant in SH-SY5Y cell, a neuroblast-link cell line. Over-expression of sPINK1* or Ub/S65E induced the formation of ubiquitin-positive puncta in cells, similar to the effects of MG132 treatment (***Figure 5—figure supplementary 1***) and the observations in HEK293 cells (***Figure 2G***).

To extend our findings, we over-expressed sPINK1*, sPINK1* with Ub/S65A (sPINK1*+Ub/S65A), Ub/S65E, or GFP control in mouse hippocampal neurons using AAV2/9. Specific expression was confirmed through GFP co-localization with NeuN fluorescence in the CA1 region (***Figure 5—figure supplementary 2***). While no visible changes were observed at 10 days post-transfection, significant neuronal loss, based on NeuN

immunofluorescence staining, was evident at 30 and 70 days in mice expressing Ub/S65E,

but not in other three groups (*Figure 5—figure supplementary 2*). Additionally, Ub/S65E

induced glial responses at 30- and 70-days post-transfection, based on GFAP/Iba1

immunofluorescence staining and GFAP/CD11b Western blot bands (*Figure 5—figure*

*supplementary 3 and 4*, **respectively**). In contrast, the sPINK1* and sPINK1*+Ub/S65A

groups showed an increase in GFAP levels at 70-days post-transfection with no changes in

Iba1or CD11b levels, suggesting potential neuronal injury (*Figure 5—figure supplementary*

*3, 4*).

To explore the underlying mechanism for neuronal injury, we conducted proteomic

analysis of mouse hippocampus at 30- and 70-days post-transfection (*Figure 5—dataset*

*supplementary 1*). Although the total number of proteins identified was similar between the

GFP and sPINK1* groups (*Figure 5—figure supplementary 5A-D*), notable differences were

observed. For example, the neuronal injury and neuroinflammation marker HMGB1 was

elevated at both time points following sPINK1* over-expression, while the mitochondrial

marker Tom20 showed a delayed decrease at 70 days, indicative of mitochondrial loss.

We also conducted gene set enrichment analysis (GSEA) and found a time-dependent

effect of sPINK1* over-expression on mitochondrial function. At 30-days, changes in

proteins associated with mitophagy, mitochondrial organization, and respiratory electron

transport were observed, suggesting an initial injury and compensatory response to

mitochondrial stress (*Figure 5—figure supplementary 5E*). By 70-days, these changes

expanded to include broader mitochondrial dysfunction and apoptotic processes, ultimately

leading to mitochondrial loss and neuronal injury (*Figure 5—figure supplementary 5E*).

Additional gene ontology (GO) analysis revealed extensive proteomic dysregulation

upon sPINK1* over-expression (***Figure 5—figure supplementary 5F-H***). At 70-days post-

transfection, a large number of proteins were either upregulated or downregulated, indicative

of disrupted proteostasis. This can be attributed to proteasomal inhibition or stress response

to increased protein aggregates. For example, we observed a significant up-regulation of

RNA-binding and RNA-processing proteins. Significantly, many of the up-regulated proteins

were associated with impaired synaptic functions and neuronal projections, while down-

regulated proteins were linked to myelination, synaptic maturation, and lipid metabolism

(***Figure 5—figure supplementary 5F-H***). As these changes were insignificant at 30-days

post-transfection (***Figure 5—figure supplementary 5F-H***), the proteomics data indicate a

systemic and time-dependent impact upon sPINK1* over-expression on neuronal structure

and function, leading to cumulative neuronal injury that mirrors the progressive pathogenesis

of neurodegeneration.

**Elevated pUb levels promote protein aggregation in mouse hippocampal neurons**

To investigate the impact of sPINK1* over-expression on protein homeostasis, we

measured the levels of PINK1 and pUb in the mouse hippocampus and evaluated protein

aggregation at 30- and 70-days post-transfection. At 30-days following sPINK1* transfection,

a distinct band for sPINK1 was observed, accompanied by a significant increase in pUb

levels (***Figure 5A***). At 70-days following sPINK1* transfection, three distinct bands could be

observed that likely correspond to sPINK1, full-length PINK1, and ubiquitinated PINK1

(***Figure 5B***). The presence of full-length PINK1 indicates mitochondrial injury, and

ubiquitinated PINK1 indicates compromised proteasomal activity. The pUb levels were

increased by 43.8 % and 59.9 % compared to the control at 30- and 70-days following

sPINK1* transfection, respectively. In comparison, the pUb level remained at the control

level with the Ub/S65A mutant co-expressed with sPINK1* (*Figure 5C*). Thus, Ub/S65A

counteract the proteasomal inhibition induced by sPINK1*, thereby promoting the

degradation of sPINK1* and prevent pUb accumulation.

Immunofluorescence staining revealed the formation of ubiquitin-positive puncta in

hippocampal neurons over-expressing sPINK1*, sPINK1+Ub/S65A, or Ub/S65E (*Figure*

*5D*). The increase of ubiquitin levels in the soluble protein fraction of the hippocampus of

brains from the sPINK1*+Ub/S65A group but not in the other groups likely reflects the

antibody detection of over-expressed Ub/S65A (*Figure 5E*). On the other hand, the levels of

ubiquitin in the insoluble protein fraction were significantly elevated in the sPINK1*,

sPINK1*+Ub/S65A, and Ub/S65E groups (*Figure 5F*). Notably, proteins smaller than 70

kDa accumulated in the sPINK1* and Ub/S65E groups, indicating an inhibition of ubiquitin

chain elongation, consistent with our findings obtained *in vitro* (*Figure 3B*), but not in the

sPINK1*+Ub/S65A group. Furthermore, over-expression of Ub/S65E resulted in significant

neuronal death and glial proliferation (*Figure 5E and Figure 5-supplementary 2, 3, 4*),

despite that the overall pUb levels appear unchanged at 70 days post-transfection (*Figure*

*5D*).

It should note that over-expression of sPINK1*, sPINK1*+Ub/S65A, or Ub/S65E had

minimal effects on autophagic flux at the animal level, as evidenced by unchanged LC3-II

and p62 levels in both soluble and insoluble protein fractions (*Figure 5—figure*

*supplementary 6*). This contrasts with the findings in cultured cells (*Figure 4F, 4G*), and

suggests that, mechanistically, protein aggregation driven by elevated pUb levels primarily

results from the inhibition of proteasomal degradation in the brains of animal or human with

neurodegenerative conditions.

**Figure 5. Elevated pUb levels induce protein aggregation in mouse hippocampal neurons.**

**(A)** Western blot analysis showing the levels of PINK1 and pUb in mouse hippocampus at 30-days post-transfection. *N*=4; **P<0.01 compared with control, one-way ANOVA.

**(B)** Western blot analysis of PINK1 in the mouse hippocampus at 70-days post-transfection. *N*=4; **P<0.001 compared with control; ###P<0.001 compared with sPINK1, one-way ANOVA.

**(C)** Western blot analysis of pUb in the mouse hippocampus at 70-days post-transfection. *N*=5; *P<0.05 compared with control, one-way ANOVA.

**(D)** Representative immunofluorescence images depicting ubiquitin staining in the CA1 neurons of the mouse hippocampus at 70-days post-transfection.

**(E)** Western blot analysis of ubiquitin in the soluble protein fraction of hippocampal lysates at 70-days post-transfection. *N*=7; *P<0.05 compared with control, one-way ANOVA.

**(F)** Western blot analysis of ubiquitin in the insoluble protein fraction of hippocampal lysates at 70-days post-transfection. Quantitative analysis was conducted for total protein at all molecular weight and for proteins with molecular weight <70 kDa. *N*=7; *P<0.05, **P<0.001 compared with control, one-way ANOVA.

**Elevated pUb levels impair neuronal integrity and cognitive function**

At 70 days post-transfection with sPINK1*, we observed protein aggregation, mitochondrial injury, neuroinflammation and proteome-wide changes (*Figure 5 and Figure 5-supplemenatary 2, 3, 4, 5*). To assess the impact of elevated pUb levels on neuronal integrity and cognitive function, we performed behavioral tests on mice at 70-days post-transfection. Over-expression of sPINK1* or Ub/S65E significantly reduced the number of sniffs toward the novel object in novel object recognition test compared to the control mice or the old object (*Figure 6A, B*). In the fear conditioning test, over-expression of sPINK1* or Ub/S65E also significantly reduced freezing time compared to control mice (*Figure 6C, D*). These findings indicate that both sPINK1* and Ub/S65E over-expression impaired cognitive functions of mice. Moreover, co-expression of Ub/S65A with sPINK1* ameliorated sPINK1*-induced cognitive deficits, suggesting a protective effect against pUb-caused impairments (*Figure 6A-D*).

Consistent with the proteomics analysis (*Figure 5—figure supplementary 5D*), Western blot analysis showed decreased levels of Tom20, a mitochondrial marker, in brains over-expressing sPINK1* (*Figure 6E*). Notably, the mitochondrial deficit induced by sPINK1* over-expression could be reversed by co-expressing Ub/S65A (*Figure 6E*). Furthermore, we observed reduced levels of MAP2, a dendritic marker, and PSD95, a postsynaptic density marker, in the hippocampus of mice over-expressing sPINK1* and Ub/S65E. The reductions could be restored by co-expressing Ub/S65A with sPINK1* (*Figure 6F, G*). Together, these findings indicate that sPINK1* over-expression induced neuronal injury originating from the nerve terminals, recapitulating neurodegenerative changes. The protective effects conferred upon Ub/S65A co-expression further support the mechanism that the neuronal injury is primarily mediated by elevated pUb levels.

Despite the absence of obvious changes in NeuN immunofluorescence at 70 days post-

transfection of sPINK1* (*Figure 5-fingure supplementary 2*), we investigated morphological

alternations in nerve terminals using Golgi staining. The staining revealed a reduction in

dendritic spine density following sPINK1* over-expression, which was reversed by co-

expressing Ub/S65A (*Figure 6H*). In contrast, Ub/S65E over-expression resulted in severe

dendritic and spine loss, characterized by marked slimmer dendrites and a significant

reduction in spine density (*Figure 6H*). These findings are consistent with the Western blot

data showing reduced levels of MAP2 and PSD95 (*Figure 6F, G*).

**Figure 6. Elevated pUb levels induce neuronal injury in mouse brains at 70 days post-transfection.**

**(A)** Novel object recognition test showing the number of sniffs toward two distinct objects in the training phase. $N=10$.

**(B)** Novel object recognition test showing the number of sniffs toward the old object (A1) and the novel object (B) in the testing phase. $N=10$; ***$P<0.001$ compared with the sniff number toward A1 object, paired $t$-test. #$P<0.05$, one-way ANOVA.

**(C, D)** The percentage of freezing time in fear conditioning tests for the evaluation of contextual (C) and cued (D) memory. For contextual memory, the mice were put in the box and received foot shock. For cued memory, the mice were put in a new box, and the same tone accompanying the foot shock was applied. $N=10$; ***$P<0.001$ compared with Box A (contextual) or absence of tone (cued), paired $t$-test. #$P<0.05$, ##$P<0.01$, ###$P<0.001$, one-way ANOVA.

**(E-G)** Western blot analyses to assess levels of mitochondrial, dendritic, and synaptic markers:
Tom20 (E), MAP2 (F), and PSD95 (G) in the mouse hippocampus. $N$=7; *$P$<0.05, **$P$<0.01,
***$P$<0.001 compared with control, one-way ANOVA.

**(H)** Golgi staining to assess the number of neuronal spines in hippocampal neurons. The left panel
shows representative images, and the right panel provides a statistical analysis of spine density on
hippocampal neuron dendrites. $N$=11-20 dendrites from 3 mice for each group. ***$P$<0.001 compared
with control, one-way ANOVA.

**sPINK1 over-expression inhibits CamKII-CREB pathway in the hippocampus**

The inhibition of proteasomal activity not only impairs protein degradation but also

disrupts protein activation by preventing the controlled removal of inhibitory proteins.

Notably, our proteomics analysis revealed a >54-fold increase in CamK2n1

(Calcium/calmodulin-dependent protein kinase II inhibitor 1, Uniprot Q6QWF9) upon

sPINK1* over-expression. CamK2n1 is an endogenous inhibitor of CamKII (***Gouet et al.,***

***2012***), and CamKII is essential for synapse formation, neuronal plasticity, and long-term

potentiation (***Hell, 2014; O'Day, 2022***). As illustrated in ***Figure 7A***, the activation of

CamKII requires the targeted degradation of CamK2n1 by the proteasome (***Buard et al.,***

***2010***). Then, the increase of phosphorylated CamKII subsequently initiate the CREB signal

pathway, representing a key mechanism in synaptic plasticity (***Davidson and Pickering, 2023;***

***Ortega-Martínez, 2015***). The over-expression of sPINK1* increases pUb level, which can

inhibit CamK2n1 degradation and suppress the activation of CamKII-CREB signal pathway.

Immunofluorescence staining confirmed the elevated CamK2n1 levels in hippocampus

CA1 neurons at 70 days post-transfection of sPINK1* compared to the controls (***Figure 7B***).

This increase was accompanied by reduced levels of phosphorylated CamKII (pCamKII)

(***Figure 7C***) and phosphorylated CREB (pCREB) (***Figure 7D***), the latter of which was

confirmed by Western blotting (***Figure 7E***). Furthermore, sPINK1* over-expression led to

reduced levels of BDNF and ERK, all downstream targets of CREB signaling (***Figure 7F, G***). These findings suggest that impaired degradation of CamK2n1 disrupts CamKII-CREB signaling, contributing to the structural deficits in dendritic spines, postsynaptic density, and neuronal projection.

**Figure 7. Elevated pUb levels inhibit CamKII-CREB signaling in mouse hippocampal neurons.**
(A) An illustration of the inhibitory effect on CamK2n1 in the CamKII-CREB1 signaling pathway.
(B-D) Immunofluorescence staining of CamK2n1 (B), pCamKII (C), and pCREB1 (D) in the CA1 region of hippocampus of mice at 70 days post-transfection. (E) Western blot analysis of pCREB1 in mouse hippocampi. $N$=8, independent $t$-test. (F and G) The immunofluorescence staining of BDNF (F) and ERK (G) in the CA1 region of mouse hippocampus.

**Discussion**

In this study, we demonstrate that elevated pUb levels are a common feature in both chronic and acute neurodegenerative conditions. While mitochondrial injury has been established as a mechanism that causes pUb elevation (***Chin et al.*, 2023; *Lambourne et al.*, 2023**), we have identified an additional pathway—the inhibition of proteasomal degradation leads to the stabilization of sPINK1 in the cytoplasm, where it remains active and phosphorylates ubiquitin.

Proteasomal inhibition can arise from many different pathological factors associated with neurodegeneration (***Wilson et al.*, 2023**). For example, cerebral ischemia can cause ATP deficiency, and hypoxia and other secondary effect from oxidative stress can further modify the proteasome and impair its function (***Chen et al.*, 2020; *Wojcik and Di Napoli*, 2004**). In AD, amyloid fibrils can impair proteasomal activities by obstructing the proteasome's substrate entry channel (***McDade et al.*, 2024; *Tseng et al.*, 2008**). Declined proteasomal activity slows the degradation of sPINK1, leading to the elevation of pUb levels in the cytoplasm. Indeed, the elevated pUb levels have been proposed as a biomarker of neurodegeneration (***Fiesel et al.*, 2015; *Hou et al.*, 2018**).

Using both *in vitro* and *in vivo* models, we demonstrate that elevated pUb levels, driven by sPINK1* over-expression, inhibit proteasomal activity and exacerbate protein aggregation. Conversely, *pink1* knockout reduced protein aggregation in aged mouse brains, brains subjected to MCAO, and HEK293 cells treated with a proteasome inhibitor. Mechanistically, we show that ubiquitin phosphorylation disrupts two critical processes: (1) the covalent elongation of ubiquitin chains, as previously reported (***Wauer et al.*, 2015b**), and (2) the non-covalent interaction between K48-linked polyUb chains and the proteasome, a novel finding in this study. This dual inhibitory mechanism interferes with the ubiquitination of substrate

proteins, the recruitment of ubiquitinated proteins to the proteasome, and ultimately, the

proteasomal degradation of substrate proteins. Impaired proteasomal activity further impedes

sPINK1 degradation, creating a self-amplifying cycle that drives pUb elevation and

progressive proteasomal dysfunction. Although cells have built-in stress responses to

perturbation of the ubiquitin-proteasome system (**Hegde et al.**, **1993**; **Rai et al.**, **2025**), the

disruption caused by elevated pUb levels targets ubiquitin—the central player in the

system—and is self-amplifying as proteasomal function deteriorates.

Extensive literature has reported that pUb plays a neuroprotective role, wherein PINK1

activation in response to mitochondrial damage results in the phosphorylation of ubiquitin

and Parkin, initiating mitophagy to remove damaged mitochondria (**Agarwal and Muqit**,

**2022**; **Li et al.**, **2023**; **Shiba-Fukushima et al.**, **2014**). Accordingly, PINK1 deficiency has

been linked to neurodegeneration, prompting the development of small-molecule activators

(**Chin et al.**, **2023**; **Hertz et al.**, **2024**; **Li et al.**, **2023**). However, in the current study, over-

expression of sPINK1 resulted in chronic neuronal injury, driven by the reciprocal

relationship between proteasomal inhibition and pUb elevation. The progressive decline of

proteasomal activity disrupts cellular proteostasis, leading to the accumulation of protein

aggregates. In addition to the mechanism of causing mitochondrial damage and inflammation

by the protein aggregates, impaired proteasomal activity hinders the degradation of inhibitory

proteins and consequently block the activation of key signaling pathways essential for

neurogenesis and memory formation. Collectively, our findings reveal that PINK1 can

function beyond the established role in mitophagy, and elevated pUb levels are not simply a

biomarker of neurodegeneration but a driving force behind progressive neuronal injury and

neuroinflammation.

The neurotoxic effect upon sPINK1 over-expression is mediated through the ubiquitin

phosphorylation. Co-expression of the phospho-null mutant Ub/S65A restored proteasomal

activity and even promoted the degradation of the rather stable sPINK1 variant used in the

current study. Additionally, Ub/S65A co-expression permitted the formation of longer

polyUb chains, further supporting its role in antagonizing ubiquitin phosphorylation. In

contrast, over-expression of the phospho-mimetic Ub/S65E mutant caused more severe

neuronal damage than sPINK1* over-expression. The difference can be attributed to the 100%

increase in pUb mimicry with Ub/S65E, whereas sPINK1*-induced pUb levels are relatively

low and subjected to dephosphorylation (*Wall et al.*, **2019**; *Wang et al.*, **2018**). Similar toxic

effects of Ub/S65E over-expression has been reported in yeast (*Swaney et al.*, **2015**).

Moreover, sPINK1* can phosphorylate both free and substrate-conjugated Ub, whereas

Ub/S65E only mimics monomeric pUb, further contributing to the heightened detrimental

effects of the Ub/S65E mutant.

In conclusion, our findings have revealed a novel pathogenetic mechanism of

neurodegeneration, driven by a self-amplifying cycle of elevated pUb levels and proteasomal

inhibition. This cycle leads to the progressive decline of proteasomal activity and the

accumulation of toxic protein aggregates, which may underlie both the sequelae of acute

insults and the chronic progression of neurodegenerative diseases. Our results demonstrate

that pUb is not merely a consequence of this process but also a critical driver of disease

progression. Therefore, targeting the pUb-mediated self-amplifying cycle represents a

promising therapeutic strategy to slow or halt the progression of neurodegenerative diseases.

## Materials and Methods

**Human brain immunofluorescence staining**

The human brain cingulate gyrus paraffin section samples include two samples from people with Alzheimer's Disease and two age-matched samples as the control. The samples with AD were obtained from the Netherlands Brain Bank (NBB), Netherlands Institute for Neuroscience, Amsterdam (Project number 1613), and the age matched control samples were obtained from the National Health and Disease Human Brain Tissue Resource Center, Hangzhou, China (Project number CN20240454). All material has been collected from donors for or from whom a written informed consent for a brain autopsy and the use of the material and clinical information for research purposes had been obtained by the NBB and CBB. The detailed information of the donors is provided in Supplementary Table 1. The experiments were conducted with approval from the Ethics Committee of the School of Medicine, Zhejiang University, with the approval number ZJU2023-011.

For immunofluorescence staining, the samples was incubated at 62°C overnight to dewax. The samples were then hydrated—Xylene for 3 times, 100% ethanol for 2 times, 95% ethanol for 2 times, 75% ethanol for 2 times, and ddH$_2$O for 2 times. The samples were placed in EDTA antigen retrieval solution, heated to 95-100°C for 30 minutes, and cooled down at room temperature for 30 minutes. After washing with PBS buffer for 3 times, the samples were placed in 3% H$_2$O$_2$ solution for 10 minutes. After washing with PBS buffer for 3 times, the samples were incubated in 0.3% Triton-X PBS for 30 minutes. After washing with PBS buffer for 3 more times, the samples were permeabilized with 0.3% Tween in PBS for 30 minutes, and rinsed 3 times with PBS to remove the detergent. The samples were incubated in a solution with 5% Sudan black B (Aladdin Biochemical Technology Co., Ltd, S109070-25g, Shanghai, China) and 70% ethanol for 1 min at room temperature.

Following PBS rinse, the samples were covered with the blocking solution (5% normal goat serum, 2% bovine serum albumin, 0.1% Tween prepared in TBS-T) for 1 hr. The samples were incubated with rabbit anti-PINK1 antibody (1:200, Novus, Colorado, USA, BC100-494), rabbit anti-pUb antibody (1:200, Millipore, Massachusetts, USA, ABS1513), or mouse anti-Aβ antibody (1:500, Biolegend, California, USA, SIG-39320) at 4 ºC overnight. After washing with PBS for 3 times, the samples were incubated with Cy3-conjugated anti-Rabbit IgG (1:200, Jackson, Pennsylvania, USA, 711-165-152) and fluorescein (FITC)-conjugated anti-mouse IgG (1:200, Jackson, Pennsylvania, USA, 715-096-150) for 2 hours at room temperature. After washing with PBS for 3 times, the samples were mounted on the slides using a ProLong™ Gold Antifade Mountant with DAPI (Invitrogen Corp., Carlsbad, CA, USA). The immunofluorescence images were taken using an Olympus FV100 confocal microscope.

**Animals**

The C57BL/6J and APP/PS1 mice were purchased from Zhejiang Academy of Medical Science. All mice were kept in the Laboratory Animal Center, Zhejiang University School of Medicine. The mice had free access to water and food in air-conditioned rooms (~26 °C, relative humidity ~50%) on a 12-h light/dark cycle. Mice were handled following the Guide for the Care and Use of the Laboratory Animals of the National Institutes of Health. The experimental protocols were approved by the Ethics Committee of Laboratory Animal Care and Welfare, Zhejiang University School of Medicine, with the approval number ZJU20190138.

Using CRISPR-Cas9-mediated genome editing technology (**Cong et al., 2013**), the *pink1* gene knockout mice (C57BL/6J) were customized purchased in Transgenic Mouse Laboratory of Laboratory Animal Center of Zhejiang University. Briefly, two sgRNA

sequences that target exon 6 of the *pink1* gene were cloned into pX330-U6-Chimeric_BB-

CBh-hSpCas9 plasmid (a gift from Feng Zhang (Addgene plasmid # 42230;

http://n2t.net/addgene: 42230; RRID: Addgene 42230) using the *Bbsl* restriction site. The

sequence of sgRNA1 is 5'-GACAGCCATCTGCAGAGAGG-3', and the sequence of

sgRNA2 is 5'-GCAGGCAGGACTCACCTCAG-3'. The knockout of *pink1* gene was

confirmed by using DAN genotype, quantitative PCR, and Western blotting.

**Mouse hippocampus CA1 injection of Adeno-associated virus (AAV)**

The rAAV-EF1a-WPRE-hGH pA (AAV2/9) was used to specifically express proteins

within neurons, a bistronic vector expressing GFP and sPINK1* (PINK1/F101M/L102-L581)

as two separate proteins. Moreover, the rAAV-EF1a-Ub/S65A-P2A-sPINK1-P2A-EGFP-

WPRE-hGH pA can express GFP, sPINK1*, and Ub/S65A as three separate proteins, and

rAAV-EF1a-Ub/S65E-P2A-EGFP-WPRE-hGH pA can express GFP and Ub/S65E as two

separate proteins.

Mice of 9-months old were anesthetized with pentobarbital sodium (80 mg/kg) by

intraperitoneal injection (I.p.). Then the mice were fixed on a mouse stereotaxic holder

(RWD Life Science Co. LTP, China) for the injection of AAV. The distinct AAV was

stereotactically injected bilaterally into the CA1 region using a microinjector (Hamilton

syringe) with a 31-gauge needle and micropump (LEGATO 130 syringe pump, KD

Scientific). After anesthesia using pentobarbital sodium (100 mg/kg, i.p.), the injections were

made at stereotaxic coordinates of Bregma: anterioposterior (AP) = −1.7 mm, mediolateral

(ML) =1.5 mm, dorsoventral (DV) = 1.7 mm. A total 0.3 μl AAV ($2.25 \times 10^{12}$ viral

particles/ml) was injected into each side of the hippocampus.

**Behavioral tests**

Novel object recognition (NOR) was used to evaluate the spatial memory of mice

according to the previous report (***Shen et al.*, 2023**). After handling for 5 days, the test was

performed on the 68[th] day post injection. An open-field test system (ViewPoint Behavior

Technology, France) was used for NOR detection. During the training, two identical objects

(named object A1 and A2, 5 cm' 5 cm' 5 cm blue cone) was placed diagonally. The mouse

was placed at the center of the box, and the movement of the mouse was recorded for 10 min.

After returning to the home cage for 24 hours, the mouse was put in the same box with one of

the objects replaced by a novel object in distinct color and shape (object B, 5 cm' 5 cm' 10 cm

yellow cuboid). The sniff number (number of visits) toward the objects was analyzed.

The fear conditioning test was performed in the Fear Conditioning System (Coulbourn

Instruments, USA) according to the previous report (***Shen et al.*, 2023**). The freezing

percentage was recorded during each phase to analyze their memory. During the training

phase, the mouse was placed in the box (identified as box A) without any stimulation for 1.5

minutes. Then, a tone (3000 Hz, 85 db) was applied for 30 seconds, and an electric shock (1

mA) was applied at the final 2 seconds. After the shock, the mouse was kept in box A for 30

second, and was put back to the home cage. The testing was performed at 24 hours after

training. The mouse was placed back in the box A without any stimulation for 5 minutes,

while the freezing percentage in box A indicates contextual memory. Two hours after the test

in box A, the mouse was placed in a new box (identified to box B). For the first 2 minutes, no

stimulus was given, and subsequently, the same tone (3000 Hz, 85 db) was applied for the

following 3 minutes. The freezing percentage in box B without and with tone was used to

analyze the cue memory of the mice.

**Focal cerebral ischemia/reperfusion**

After anesthetized with pentobarbital sodium (80 mg/kg) by intraperitoneal injection

(I.p.), the mice were operated with middle cerebral artery occlusion (MCAO) for 2 hours

ischemia followed by 24 hours reperfusion. Briefly, a 6-0 nylon monofilament suture was

inserted into the internal carotid to occlude the origin of MCA (*Shi et al.*, 2012). Two hours

after occlusion, the suture was withdrawn to allow reperfusion for 24 hours. Mouse body

temperature was maintained at 37 °C using a thermostatic pad during and after surgery. In

sham-operated mice, the external carotid artery was surgically isolated but the suture was no

inserted.

**Mouse brain tissue collection**

Mice were anesthetized using pentobarbital sodium (150 mg/kg, i.p.). The mouse brains

were removed after transcardially perfused with 4 °C saline and freshly prepared 4%

paraformaldehyde. The brains were further fixed in 4% paraformaldehyde at 4 °C for one day

and transferred to 30% sucrose solution for dehydration. The brains were sliced into 30 μm

thick slices by cryomicrotomy (CM1900, Leica, Wetzlar, Germany). The slices were stored

at -20°C in the solution (30% glycerol, 30% glycol, 40% PBS) for immunofluorescence and

Nissle staining.

For Western blotting, the mouse was transcardially perfused only with 4 °C saline. The

brains were removed, and the hippocampus was isolated. The hippocampus was quickly

frozen in liquid nitrogen and stored at -80 °C until use.

**Cell lines, cell culture, and cell transfection**

HEK293 cells and SH-SYT5Y cells were purchased from the Institute of Cell Biology

of the Chinese Academy of Sciences (Cell Biology of the Chinese Academy of Sciences,

Shanghai, China). The *pink1* gene knockout HEK293 cell was homely made using CRISPR-

Cas9-mediated genome editing technology(***Ran et al., 2013***). PINK1 sgRNA

(CCTCATCGAGGAAAAACAGG) was cloned to U6-sgPINK1-mCherry plasmid (a gift

from John Doench & David Root (Addgene plasmid # 78038; http://n2t.net/addgene:78038 ;

RRID:Addgene_78038). The U6-sgPINK1-mCherry plasmid and PX458-Cas9-EGFP

plasmid (a gift from Prof. Feng Zhang (Broad Institute, Cambridge, MA), Addgene plasmid

#48138) were co-transfected into HEK293 cells. The mCherry and GFP double-positive cells

were single-cell sorted 24 hours post-transfection using an MoFloAstrios EQ cell sorter

(Beckman Coulter, US) and grown in separate cultures that were subsequently screened for

the presence of frameshift mutations leading to nonsense-mediated decay on both alleles. The

*pink1* knockout was confirmed using western blotting.

Cells were grown in Dulbecco's modified essential medium (DMEM, Gibco by Thermo

Fisher Scientific, C11995500BT) supplemented with 10% heat-inactivated fetal bovine

serum (FBS, Zhejiang Tianhang Biotechnology, China, 11011-8611) in the atmosphere of 5%

$CO_2$ at 37 °C.

Lipo3000 (Invitrogen, California, USA, L3000001) was used to transfect plasmids,

following to the manufacturer' protocol. Briefly, an appropriate number of cells were seeded

one day before transfection with 70-80% confluent at the time of transfection. Before

transfection, plasmid DNA was carefully mixed with 250 μl Opti-MEM (Gibco by Life

Technologies, Carlsbad, USA). Subsequently, 250 μl Lipo3000 was carefully mixed with

Opti-MEM. The mixture was incubated at room temperature for 25 min. The freshly formed

DNA/Lipo3000 precipitates were carefully pipetted to the cells. The medium containing

transfection reagents was removed after 6 hours and fresh medium was added.

At the end of treatments, the cells on cover slides were washed with 37 °C PBS and then

fixed using freshly prepared 4% paraformaldehyde. Subsequently, the cells were stored in

PBS under 4 °C for immunofluorescence staining.

**Oxygen glucose deprivation and reperfusion (OGD/R)**

758    Cells were washed twice by using glucose-free DMEM (Procell, PM150270) and

759    incubated at 37 °C with the ventilation of 95% $N_2$ and 5% $CO_2$ for 2 hours. The control cells

760    were washed twice by using high glucose DMEM, and incubated in 95% air and 5% $CO_2$

761    under 37 °C for 2 hours. Then the cells were cultured in the high glucose DMEM with 10%

762    FBS in 95% air and 5% $CO_2$ under 37 °C. This was identified as reperfusion.

**The construction of Ub-R-GFP plasmid**

The Ub-R-GFP is used for the *in vivo* measurement of proteasomal degradation activity, and

constructed according to a previous report(***Dantuma et al.*, 2000**). Briefly, the ubiquitin open

reading frame was amplified by PCR with the following primers:

Sense primer: 5'-GCG GAATTCACCATGCAGATCTTCGTGAAGACT-3'

Antisense primer:

5'-GCG GGATCCTGTCGACCAAGCTTCCCGCGCCCACCTCTGAGACGGAGTAC-3'

The PCR product was cloned into the EcoRI and BamHI sites of the EGFP-N2 vector. Beside

the R residue, a 12 amino acid peptide was inserted between ubiquitin and GFP to increase

the proteasomal degradation. Thus, the final product is Ub-R-GKLGRQDPPVAT-GFP. The

accumulation of GFP protein was determined using Western blot, reflecting proteasomal

degradation activity.

**Nissl staining**

The brain slices were incubated in a mixture of acetone and chloroform (1:1) for 15 min,

and then sequentially incubated in 100%, 95%, 70% alcohol for 5 min. The slices were

stained for 10 min in Nissl staining containing 0.2% purple crystal (Yuanhang Reagent

Factory, Shanghai, China, YHSJ-01- 92) and 0.3% acetic acid. Subsequently, the slices were

dehydrated by incubation in 70%, 95%, 100% alcohol. Finally, the slices were put into

xylene for 5 min and then mounted with a mixture of neutral resin and xylene (1:1). The

images were taken under a microscope (Olympus BX51, Japan).

**Immunofluorescence staining**

The floating brain slices or cells on the cover glasses were incubated in 0.1% Triton-X

PBS for 30 min, and followed by the incubation in 5% donkey serum for 1 hours. The

samples were then incubated with mouse anti-ubiquitin antibody (1:200, Santa Cruz

Biotechnology, Texas, USA; SC-8017), rabbit anti-pUb antibody (1:200, Millipore,

Massachusetts, USA; ABS1513), rabbit anti-PINK1 antibody (1:200, Novus, Colorado, USA;

BC100-494), mouse anti-GFAP antibody (1:1000, CST, Pennsylvania, USA; #3670), rabbit

anti-Iba1 antibody (1:1000, FUJIFILM Wako Pure Chemical Corporation, Tokyo, Japan;

019-19741), rabbit anti-NeuN antibody (1:1000, CST, Pennsylvania, USA; #36662), mouse

anti-A$\beta$ (1:500, Biolegend, California, USA, SIG-39320), rabbit anti-pCREB antibody

(1:1000, ABclonal Technology, Wuhan, China; AP0019), rabbit anti-ERK1/2 antibody

(1:200, ABclonal Technology, Wuhan, China; E-AB-31374), rabbit anti-pCamKII antibody

(1:200, ABclonal Technology, Wuhan, China; AP0255), rabbit anti-CamK2n1 (1:200,

Thermo fisher, Massachusetts, USA; PA5-23740), and rabbit anti-BDNF antibody (1:200,

ABclonal Technology, Wuhan, China; A11028) at 4 °C overnight. After washed with PBS

(10 min' 3 times), the samples were incubated with Cy3 AffiniPure Donkey Anti-Mouse IgG

(H+L) (1:200, Jackson ImmunoResearch, PA, USA;715-605-150) or Cy3 AffiniPure Goat

Anti-Rabbit IgG (H+L) (1:200, Jackson ImmunoResearch, PA, USA;715-605-150; 111-165-

003) or Alexa Fluor488 AffiniPure Goat Anti-Rabbit IgG (H+L) (1:200,

Jackson ImmunoResearch, PA, USA; 111-545-003) or Alexa Fluor488 AffiniPure Goat Anti-

Rabbit IgG (H+L) (1:200, Jackson ImmunoResearch, PA, USA; 715-545-150) for 2 h. After

washing with PBS (10 min each for 3 times), the slices were mounted on slides using a

ProLong Gold Antifade Mountant with DAPI. The images were taken under an Olympus

FV100 confocal microscope (Olympus, Japan) and analyzed using MetaMorph (version

7.8.0.0, Molecular Devices, LLC. San Jose, CA 95134 USA).

**The preparation of soluble and insoluble protein fractions for Western blot analysis**

We prepared the soluble and insoluble protein fractions as described in a previous report

(*Wirths*, **2017**). The cells and brain tissues for the Western blot were lysed using RIPA lysis

buffer (Beyotime Biotechnology Research Institute, Jiangsu, China; P0013B) with protease

inhibitor (Beyotime Biotechnology Research Institute, Jiangsu, China; P1005) and

phosphatase inhibitor (Beyotime Biotechnology Research Institute, Jiangsu, China; P1081).

The cells were lysed on ice for 30 min, and every 10 min, the cells were gently vortexed. The

brain samples were thoroughly homogenized with a precooled TissuePrep instrument (TP-24,

Gering Instrument Company, Tianjin, China) for 1 min at 4 °C. Subsequently, the lysate was

centrifuged at 12,000 rpm for 30 min at 4 °C. The supernatant was collected as the soluble

protein fraction for Western blot analysis.

The precipitate was resuspended in 20 μl SDS buffer (2% SDS, 50 mM Tris-HCl, pH7.5)

for ultrasonic pyrolysis at 4 °C. The ultrasonic pyrolysis cycle included 10-sec ultrasonic

pyrolysis (Diagenode, Seraing, Belgium) and 30-sec interval, for a total of 8 cycles. The

samples were then centrifuged at 12,000 rpm for 30 min at 4 °C. The supernatant was

collected as the insoluble protein fraction for Western blot analysis.

All protein concentration was determined by using BCA Protein Assay Kit according to

the instruction (Beyotime Biotechnology, Shanghai, China; P0009).

**Western blot analysis**

Appropriate protein samples (50-100 μg) were used and separated using SDS-page gel.

The separated proteins were transferred to nitrocellulose membrane. The following antibodies

were used: mouse anti-ubiquitin antibody (1:800, Santa Cruz Biotechnology, Texas, USA;

SC-8017), rabbit anti-pUb antibody (1:1000, Millipore, Massachusetts, USA; ABS1513),

rabbit anti-PINK1 antibody (1:1000, Novus, Colorado, USA; BC100-494), mouse anti-GFAP

antibody (1:1000, CST, Pennsylvania, USA; #3670), rabbit anti-CD11b antibody (1:1000,

Abcam, Cambridge, UK; ab133357), mouse anti-GAPDH antibody (1:5000, Proteintech,

Wuhan, China; 60004-1-Ig), mouse anti-MAP2 antibody (1:2000, Millipore, Billerica, MA,

USA; AB5622), rabbit anti-PSD95 antibody (1:1000, CST, Pennsylvania, USA; 3450S),

rabbit anti-Tom20 antibody (1:1000, Pennsylvania, USA; 42406S), rabbit anti-LC3 antibody

(1:1000, Sigma, Massachusetts, USA; L7543), rabbit anti-p62 antibody (1:1000, Abcam,

California, USA; ab109012), rabbit anti-pCREB antibody (1:1000, ABclonal Technology,

Wuhan, China; AP0019), and mouse anti-FLAG antibody (1:10000, TransGen biotech,

Beijing, China; HT201-01). The secondary antibody was HRP-conjugated goat anti-mouse

IgG (1:3000, Cell Signaling Technology, MA, USA; 7076S) or HRP-conjugated goat anti-

rabbit IgG (1:10000, Jackson ImmunoResearch, PA, USA, 111-035-003).

The immunoblots were then detected using ECL reagents (Potent ECL kit,

MultiSciences Biotech, Hangzhou, China; P1425) and measured using GBOX (LI-COR,

Odyssey-SA-GBOX, NE, USA). The results were normalized to GAPDH or Ponceau

staining (Beyotime Biotechnology Research Institute, Jiangsu, China; P0022), or to the

eGFP-expression control on the same immunoblot membrane.

**Golgi staining**

On the 70[th]-day post-transfection, mice were anesthetized by intraperitoneal injection of

pentobarbital sodium (150 mg/kg) before sacrifice. The mouse brain was quickly removed,

and the dorsal hippocampus was separated. The hippocampus was immediately fixed in the

fixative solution (Servicebio, Wuhan, China, G1101) for over 48 hours.

The dorsal hippocampus was sliced with a thickness of 2-3 mm thickness around the

injection site. The tissue was gently rinsed with PBS at least 3 times. The hippocampus tissue

was then placed in Golgi-cox staining solution (Servicebio, Wuhan, China; G1069), and

incubated in a dark for 14 days. The Golgi-cox staining solution was changed 48 hours after

the first soak, and every 3 days afterwards. On the14[th]-day after staining, the tissues were

immersed in distilled water for 3 times and then incubated in 80% acetic acid overnight until

the tissue became soft. After rinsing with distilled water, the tissue was placed into 30%

sucrose.

The tissue was cut into 100 μm slices with an oscillating microtome, and the slices were

placed on a gelatin slide and dried overnight in the dark. The slices were then treated with

ammonia water for 15 min. After washing with distilled water, the slices were incubated in

the fixing solution for 15 min. After washing, the slices were sealed using glycerin gelatin.

Images were taken using VS120 Virtual Slide Microscope (Olympus, Japan).

**The *in vitro* ubiquitin-dependent proteasome degradation**

Ubiquitin was prepared as previously described (***Liu et al.*, 2015; *Liu et al.*, 2012**).

K48-linked di-ubiquitin and tetra-ubiquitin were prepared following an established protocol

(***Pickart and Raasi*, 2005**), with the conjugation reaction catalyzed by 2.5 μM ubiquitin-

activating enzyme E1 (Uniprot P22314) and 20 μM ubiquitin-conjugating enzyme E2-25K

(Uniprot P61086), in 20 mM pH 8.0 Tris-HCl buffer. Ub/K48R mutant was incorporated as

the distant Ub, and Ub/77D mutant was incorporated as the proximal ubiquitin to prepare

ubiquitin chain of the desired length. The residue D77 at the C-terminus was removed with

hydrolase YUH1. The fusion protein His-TEV-Ub-GFP was prepared recombinantly based

on the design previously described (***Dantuma et al.*, 2000; *Inobe et al.*, 2011**). The *Pediculus*

*humanus corporis* PINK1 kinase (phPINK1, uniport E0W1I1) was prepared and purified as

previously described (***Dong et al.*, 2017; *Wauer et al.*, 2015a**). ubiquitin phosphorylation was

confirmed with electrospray mass spectrometry (Agilent G6530 Q-TOF).

The 26S proteasome from HEK293 cells was prepared following an established

protocol, with the purification tag appended at the C-terminus of Rpn11 (***Guerrero et al.*,**

**2006; *Huang et al.*, 2016**). The proteasome activity was assessed with a fluorogenic peptide,

N-succinyl-Leu-Leu-Val-Tyr-7-amido-4-methyl coumarin (Sigma, Massachusetts, USA;

S6510). A final concentration of 100 nM proteasome and 100 μM proteasome peptide

substrate was prepared in the reaction buffer, containing 50 mM Tris-HCl pH 7.5, 4 mM ATP

(Sigma- Aldrich, Cat# A6559) and 5 mM $MgCl_2$. The fluorescence intensity of the product

was measured by a fluorometer (Horiba Scientific, FluoroMax-4) at an excitation wavelength

of 360 nm.

For Western blot analysis of GFP, 200 nM proteasomal substrate and 10 nM human

26S proteasome were prepared in 20 mM Tris pH 8.0 buffer, with 50 mM NaCl, 5 mM ATP,

and 5 mM $MgCl_2$. The mixture was incubated at 37 ° C for 0, 10, 30, 60, 90, and 120 min. At

each time point, 20 μL samples were taken for Western blot analysis. A mouse anti-GFP

antibody (1:2000, San-Ying Proteintech Group, 66002-1) and an HRP-conjugated goat anti-

mouse IgG (1:3000, CST, Pennsylvania, USA; 7076S) were used. ImageJ was used to

analyze the band intensities, which are normalized to the initial intensity.

The recombinant Ub, pUb (phosphorylated by phPINK1, uniport E0W1I1), Ub/S65A,

and Ub/S65E were reacted with 2.5 μM human ubiquitin-activating enzyme E1 and 20 μM

ubiquitin-conjugating enzyme 25K in a reaction buffer containing 20 mM Tris-HCl, pH 8.0.

The reaction was carried out under standard conditions to promote K48-linked polyubiquitin

chain formation. The resulting polyubiquitin chains were analyzed by SDS-PAGE, followed

by Coomassie blue staining to visualize the chain formation.

**TIRF analysis of ubiquitin-proteasome interaction**

The coverslips were prepared following an established protocol (**_Roy et al._**, **2008**). Each

coverslip was divided into multiple lanes with double-sided tape for parallel detection of the

transient association of K48-polyUb-GFP or pK48-polyUb-GFP in the TIRF experiments,

thus to ensure repeatability with different proteins added to the same slide. Streptavidin

(VWR Life Science, Cat# 97062-808) in imaging buffer at a concentration of 50 μg/ml was

loaded; excess unbound streptavidin was washed away. The imaging buffer contains 25 mM

Tris-HCl (pH 7.5), 50 mM NaCl, 10 mM MgCl2, 40 mM imidazole, 5 m/mL BSA (Sigma-

Aldrich), 2.5 mM ADP (Sigma-Aldrich), and 0.5 mM ATP-γ-S (Sigma-Aldrich), as

previously described (**_Lu et al._**, **2015**).

The purified human 26S proteasome at 20 nM in the imaging buffer was added to the

streptavidin-immobilized coverslip, with excess unbound proteins washed away with the

imaging buffer. The substrate protein, either K48-polyUb-GFP or pK48-polyUb-GFP, was

then added to the coverslip to a final concentration of 400 pM. The imaging was performed

on a Nikon A1 TIRF microscope equipped with a 488 nm laser and an EMCCD camera

(Andor DU-897). The time series were acquired at 300 ms per frame for a total duration of 2

min for each time series. The view area was 512 pixels by 512 pixels (16 by 16 μm), and the

experiments were repeated four times on four different proteasome-immobilized cover slides.

Only when K48-polyUb-GFP or pK48-polyUb-GFP is associated with the immobilized

proteasome, the green puncta can be observed. The puncta density was counted in iSMS

software (***Preus et al.*, 2015**) with default settings.  The time series of puncta with brightness

above a certain threshold ($\geq$ 3000 a.u.) was captured and the dwell time of more than three

consecutive frames are selected for counting the overall number of puncta. The dwell times

were tabulated in 300 ms bins and fitted with single exponential decay.

**Proteomics analysis**

Hippocampus samples were rinsed using PBS and lysed in RIPA lysis buffer. The

samples were then denatured at 95 °C for 5 min, followed by sonication at 4 °C (3 sec on

followed with 3 sec off, with 30% amplitude). The lysate was centrifuged at 16,000 g for 10

min at 4 °C, and the supernatant was collected as whole tissue extract. Protein concentration

was determined by Bradford protein assay. Extracts from the protein sample (2 g) were

trypsin digested.

The enriched peptides sample were precipitated in solution A (0.1% formic acid), and

was centrifuged at 16,000 g for 10 min. The peptides were separated on a reverse-phase

nano-HPLC C18 column (Precolumn, 3 μm, 120 Å, 2x100 i.d.; analysis column, 1.9 μm 120

Å, 30 cm x150 μm, i.d.) at a flow rate of 600 nL/min with a 150 min gradient of 7-95%

solution B (0.1% formic acid in acetonitrile). For peptide ionization, 2100 V was applied, and

a 320 ºC capillary temperature was used. For detection with Q Exactive HF, peptides were

analyzed with one full scan (350–1400 m/z, R = 120,000 at 200 m/z) with an automatic gain

control target of $3x10^6$ ions, with max injection time of 80 ms, followed by up to 30 data-

dependent MS/MS scans with high-energy collision dissociation (target $5 \ x10^4$ ions, max

injection time 19 ms, isolation window 1.6 m/z, normalized collision energy of 27%),

detected in the Orbitrap (R = 15,000 at 200 m/z). Dynamic exclusion time was set as 30 sec.

Raw MS data were searched against the mouse National Center for Biotechnology

Information (NCBI) Refseq protein database (updated on 2013/07/01, 29764 entries) by the

software Thermo Proteome Discoverer 2.1 using Mascot 2.3. The mass tolerances were 20

ppm for precursor and 50 mmu for product ions for Q Exactive HF. The search engine set

Acetyl(Protein N-term), oxidation(M), Phospho(ST), Phospho(Y) as variable modifications.

Trypsin digestion of up to two missed cleavages was allowed. The peptide identifications

were accepted at a false discovery rate (FDR) of 1%. Using a label-free approach, a unique

peptide was used to represent the absolute abundance of a particular peptide across the

sample.

For the proteomic data analysis, quantile normalization was applied to eliminate batch

effect, and missing values were imputed using the minimum value observed in the dataset.

The differentially expressed proteins were defined with the following these criteria: an

increase by more than two-fold or decrease by more than 50%. For Gene Set Enrichment

Analysis (GSEA) analysis, Gene Ontology (GO) analysis was conducted by GSEApy

(version 0.10.5) Python package to evaluate the enriched GO terms. The association of

proteins with GO terms was derived from the DAVID database. For Gene Set Enrichment

Analysis (GSEA), the ranking metric used was the fold change, and significantly enriched

GO terms were filtered with a cutoff $P$-value of 0.05.

**Statistical analysis**

Data are presented as mean±SD. The number in statistical analysis indicates biological

replicates. GraphPad Prism Software (version 6.0, GraphPad Software, San Diego, CA, USA)

was used for statistical analysis and plot graphs. The ROUT test was used to identify outliers.

The Brown-Forsythe test was used to evaluate the equal variances of the data. D'Agonstino-

Pearson omnibus normality test was performed to test the normality of the data. If the data

pass the normality test and equal variance test, *t*-test, parametric one-way ANOVA (Tukey

multiple comparisons test), or two-way ANOVA (Newman-Keuls multiple comparisons test)

was used. If not, nonparametric test (Dunn's multiple comparisons test) was used. Chi-square

test was also used to assess the differences in categorical data distributions. $P<0.05$ was

considered statistically significant.

**Author contributions**

W.P.Z. and C.T. designed the study and wrote the manuscript. C.C., T.Y.G., H.W.Y., Y.Z., T.Y., T.W., and T.F.W. conducted the experiments. Y.B.L. and W.P.Z. analyzed the data at mouse and cellular level. T.T.L. analyzed the proteomic data. C.T. analyzed the data at protein level.

**Acknowledgements**

This study was supported by the National Key R&D Program of China (2023YFF1204400) to C.T. and W.P.Z., and (2021YFF1200900) to T.T.L., the National Natural Science Foundation of China (92353304) to C.T. and (32070666) to T.T.L..

**Data availability**

All data needed to evaluate the conclusions in the paper are present in the paper and/or the Supplementary Materials.

**References**

Agarwal S, and Muqit MMK. 2022. PTEN-induced kinase 1 (PINK1) and Parkin: Unlocking a mitochondrial quality control pathway linked to Parkinson's disease. *Curr Opin Neurobiol* **72**: 111-119. https://doi: 10.1016/j.conb.2021.09.005

Beilina A, Van Der Brug M, Ahmad R, Kesavapany S, Miller DW, Petsko GA, and Cookson MR. 2005. Mutations in PTEN-induced putative kinase 1 associated with recessive parkinsonism have differential effects on protein stability. *Proc Natl Acad Sci U S A* **102**: 5703-5708. https://doi: 10.1073/pnas.0500617102

Buard I, Coultrap SJ, Freund RK, Lee YS, Dell'Acqua ML, Silva AJ, and Bayer KU. 2010. CaMKII "autonomy" is required for initiating but not for maintaining neuronal long-term information storage. *J Neurosci* **30**: 8214-8220. https://doi: 10.1523/jneurosci.1469-10.2010

Chen C, Qin H, Tan J, Hu Z, and Zeng L. 2020. The Role of Ubiquitin-Proteasome Pathway and Autophagy-Lysosome Pathway in Cerebral Ischemia. *Oxid Med Cell Longev* **2020**: 5457049. https://doi: 10.1155/2020/5457049

Chin RM, Rakhit R, Ditsworth D, Wang C, Bartholomeus J, Liu S, Mody A, Laishu A, Eastes A, Tai C, Kim RY, Li J, Hansberry S, Khasnavis S, Rafalski V, Herendeen D, Garda V, Phung J, de Roulet D, Ordureau A, Harper JW, Johnstone S, Stohr J, and Hertz NT. 2023. Pharmacological PINK1 activation ameliorates Pathology in Parkinson's Disease models. *bioRxiv*. https://doi: 10.1101/2023.02.14.528378

Cong L, Ran FA, Cox D, Lin S, Barretto R, Habib N, Hsu PD, Wu X, Jiang W, Marraffini LA, and Zhang F. 2013. Multiplex genome engineering using CRISPR/Cas systems. *Science* **339**: 819-823. https://doi: 10.1126/science.1231143

Costa RO, Martins H, Martins LF, Cwetsch AW, Mele M, Pedro JR, Tome D, Jeon NL, Cancedda L, Jaffrey SR, and Almeida RD. 2019. Synaptogenesis Stimulates a Proteasome-Mediated Ribosome Reduction in Axons. *Cell reports* **28**: 864-876 e866. https://doi: 10.1016/j.celrep.2019.06.080

Dantuma NP, Lindsten K, Glas R, Jellne M, and Masucci MG. 2000. Short-lived green fluorescent proteins for quantifying ubiquitin/proteasome-dependent proteolysis in living cells. *Nat Biotechnol* **18**: 538-543. https://doi: 10.1038/75406

Davidson K, and Pickering AM. 2023. The proteasome: A key modulator of nervous system function, brain aging, and neurodegenerative disease. *Front Cell Dev Biol* **11**: 1124907. https://doi: 10.3389/fcell.2023.1124907

Dong X, Gong Z, Lu YB, Liu K, Qin LY, Ran ML, Zhang CL, Liu Z, Zhang WP, and Tang C. 2017. Ubiquitin S65 phosphorylation engenders a pH-sensitive conformational switch. *Proc Natl Acad Sci U S A* **114**: 6770-6775. https://doi: 10.1073/pnas.1705718114

Fiesel FC, Ando M, Hudec R, Hill AR, Castanedes-Casey M, Caulfield TR, Moussaud-Lamodiere EL, Stankowski JN, Bauer PO, Lorenzo-Betancor O, Ferrer I, Arbelo JM, Siuda J, Chen L, Dawson VL, Dawson TM, Wszolek ZK, Ross OA, Dickson DW, and Springer W. 2015. (Patho-)physiological relevance of PINK1-dependent ubiquitin phosphorylation. *EMBO Rep* **16**: 1114-1130. https://doi: 10.15252/embr.201540514

Gan ZY, Callegari S, Cobbold SA, Cotton TR, Mlodzianoski MJ, Schubert AF, Geoghegan ND, Rogers KL, Leis A, Dewson G, Glukhova A, and Komander D. 2022. Activation mechanism of PINK1. *Nature* **602**: 328-335. https://doi: 10.1038/s41586-021-04340-2

Gao J, Li M, Qin S, Zhang T, Jiang S, Hu Y, Deng Y, Zhang C, You D, Li H, Mu D, Zhang Z, and Jiang C. 2016. Cytosolic PINK1 promotes the targeting of ubiquitinated proteins to the aggresome-autophagy pathway during proteasomal stress. *Autophagy* **12**: 632-647. https://doi: 10.1080/15548627.2016.1147667

Gouet C, Aburto B, Vergara C, and Sanhueza M. 2012. On the mechanism of synaptic depression induced by CaMKIIN, an endogenous inhibitor of CaMKII. *PLoS One* **7**: e49293. https://doi: 10.1371/journal.pone.0049293

Guerrero C, Tagwerker C, Kaiser P, and Huang L. 2006. An integrated mass spectrometry-based proteomic approach: quantitative analysis of tandem affinity-purified in vivo cross-linked protein complexes (QTAX) to decipher the 26 S proteasome-interacting network. *Mol Cell Proteomics* **5**: 366-378. https://doi: 10.1074/mcp.M500303-MCP200

Hegde AN, Duke LM, Timm LE, and Nobles H. 2023. The Proteasome and Ageing. *Subcell Biochem* **102**: 99-112. https://doi: 10.1007/978-3-031-21410-3_5

Hegde AN, Goldberg AL, and Schwartz JH. 1993. Regulatory subunits of cAMP-dependent protein kinases are degraded after conjugation to ubiquitin: a molecular mechanism underlying long-term synaptic plasticity. *Proc Natl Acad Sci U S A* **90**: 7436-7440. https://doi: 10.1073/pnas.90.16.7436

Hell JW. 2014. CaMKII: claiming center stage in postsynaptic function and organization. *Neuron* **81**: 249-265. https://doi: 10.1016/j.neuron.2013.12.024

Herhaus L, and Dikic I. 2015. Expanding the ubiquitin code through post-translational modification. *EMBO reports* **16**: 1071-1083. https://doi: 10.15252/embr.201540891

Hertz N, Chin R, Rakhit R, Ditsworth D, Wang C, Bartholomeus J, Liu S, Mody A, Laihsu A, Eastes A, Tai C, Kim R, Li J, Khasnavis S, Rafalski V, Heerendeen D, Garda V, Phung J, de Roulet D, Ordureau A, Harper JW, Johnstone S, and Stohr J. 2024. Pharmacological PINK1 activation ameliorates Pathology in Parkinson's Disease models. *Res Sq*. https://doi: 10.21203/rs.3.rs-4356493/v1

Hou X, Fiesel FC, Truban D, Castanedes Casey M, Lin WL, Soto AI, Tacik P, Rousseau LG, Diehl NN, Heckman MG, Lorenzo-Betancor O, Ferrer I, Arbelo JM, Steele JC, Farrer MJ, Cornejo-Olivas M, Torres L, Mata IF, Graff-Radford NR, Wszolek ZK, Ross OA, Murray ME, Dickson DW, and Springer W. 2018. Age- and disease-dependent increase of the mitophagy marker phospho-ubiquitin in normal aging and Lewy body disease. *Autophagy* **14**: 1404-1418. https://doi: 10.1080/15548627.2018.1461294

Huang X, Luan B, Wu J, and Shi Y. 2016. An atomic structure of the human 26S proteasome. *Nat Struct Mol Biol* **23**: 778-785. https://doi: 10.1038/nsmb.3273

Inobe T, Fishbain S, Prakash S, and Matouschek A. 2011. Defining the geometry of the two-component proteasome degron. *Nat Chem Biol* **7**: 161-167. https://doi: 10.1038/nchembio.521

Kinger S, Jagtap YA, Kumar P, Choudhary A, Prasad A, Prajapati VK, Kumar A, Mehta G, and Mishra A. 2024. Proteostasis in neurodegenerative diseases. *Adv Clin Chem* **121**: 270-333. https://doi: 10.1016/bs.acc.2024.04.002

Komander D, and Rape M. 2012. The ubiquitin code. *Annu Rev Biochem* **81**: 203-229. https://doi: 10.1146/annurev-biochem-060310-170328

Koyano F, Okatsu K, Kosako H, Tamura Y, Go E, Kimura M, Kimura Y, Tsuchiya H, Yoshihara H, Hirokawa T, Endo T, Fon EA, Trempe JF, Saeki Y, Tanaka K, and Matsuda N. 2014. Ubiquitin is phosphorylated by PINK1 to activate parkin. *Nature* **510**: 162-166. https://doi: 10.1038/nature13392

Krishna KK, Behnisch T, and Sajikumar S. 2022. Modulation of the Ubiquitin-Proteasome System Restores Plasticity in Hippocampal Pyramidal Neurons of the APP/PS1 Alzheimer's Disease-Like Mice. *J Alzheimers Dis* **86**: 1611-1616. https://doi: 10.3233/JAD-215718

Lambourne OA, Bell S, Wilhelm LP, Yarbrough EB, Holly GG, Russell OM, Alghamdi AM, Fdel AM, Varricchio C, Lane EL, Ganley IG, Jones AT, Goldberg MS, and Mehellou Y. 2023. PINK1-Dependent Mitophagy Inhibits Elevated Ubiquitin Phosphorylation Caused by

Mitochondrial Damage. *J Med Chem* **66**: 7645-7656. https://doi:
10.1021/acs.jmedchem.3c00555

Lazarou M, Sliter DA, Kane LA, Sarraf SA, Wang C, Burman JL, Sideris DP, Fogel AI, and
Youle RJ. 2015. The ubiquitin kinase PINK1 recruits autophagy receptors to induce
mitophagy. *Nature* **524**: 309-314. https://doi: 10.1038/nature14893

Li J, Yang D, Li Z, Zhao M, Wang D, Sun Z, Wen P, Dai Y, Gou F, Ji Y, Zhao D, and Yang
L. 2023. PINK1/Parkin-mediated mitophagy in neurodegenerative diseases. *Ageing Res Rev*
**84**: 101817. https://doi: 10.1016/j.arr.2022.101817

Liu Z, Gong Z, Jiang WX, Yang J, Zhu WK, Guo DC, Zhang WP, Liu ML, and Tang C.
2015. Lys63-linked ubiquitin chain adopts multiple conformational states for specific target
recognition. *eLife* **4**: e05767. https://doi: 10.7554/eLife.05767

Liu Z, Zhang WP, Xing Q, Ren X, Liu M, and Tang C. 2012. Noncovalent dimerization of
ubiquitin. *Angew Chem Int Ed Engl* **51**: 469-472. https://doi: 10.1002/anie.201106190

Lu Y, Lee BH, King RW, Finley D, and Kirschner MW. 2015. Substrate degradation by the
proteasome: a single-molecule kinetic analysis. *Science* **348**: 1250834. https://doi:
10.1126/science.1250834

Mauvezin C, and Neufeld TP. 2015. Bafilomycin A1 disrupts autophagic flux by inhibiting
both V-ATPase-dependent acidification and Ca-P60A/SERCA-dependent autophagosome-
lysosome fusion. *Autophagy* **11**: 1437-1438. https://doi: 10.1080/15548627.2015.1066957

McDade E, Liu H, Bui Q, Hassenstab J, Gordon B, Benzinger T, Shen Y, Timsina J, Wang L,
Sung YJ, Karch C, Renton A, Daniels A, Morris J, Xiong C, Ibanez L, Perrin R, Llibre-
Guerra JJ, Day G, Supnet-Bell C, Xu X, Berman S, Chhatwal J, Ikeuchi T, Kasuga K, Niimi
Y, Huey E, Schofield P, Brooks W, Ryan N, Jucker M, Laske C, Levin J, Voglein J, Roh JH,
Lopera F, Bateman R, and Cruchaga C. 2024. Ubiquitin-Proteasome System in the Different
Stages of Dominantly Inherited Alzheimer's Disease. *Res Sq*. https://doi: 10.21203/rs.3.rs-
4202125/v1

O'Day DH. 2022. Calmodulin Binding Domains in Critical Risk Proteins Involved in
Neurodegeneration. *Curr Issues Mol Biol* **44**: 5802-5814. https://doi: 10.3390/cimb44110394

Okatsu K, Kimura M, Oka T, Tanaka K, and Matsuda N. 2015. Unconventional PINK1
localization to the outer membrane of depolarized mitochondria drives Parkin recruitment. *J*
*Cell Sci* **128**: 964-978. https://doi: 10.1242/jcs.161000

Okatsu K, Uno M, Koyano F, Go E, Kimura M, Oka T, Tanaka K, and Matsuda N. 2013. A
dimeric PINK1-containing complex on depolarized mitochondria stimulates Parkin
recruitment. *J Biol Chem* **288**: 36372-36384. https://doi: 10.1074/jbc.M113.509653

Ortega-Martínez S. 2015. A new perspective on the role of the CREB family of transcription
factors in memory consolidation via adult hippocampal neurogenesis. *Front Mol Neurosci* **8**:
46. https://doi: 10.3389/fnmol.2015.00046

Pickart CM, and Raasi S. 2005. Controlled synthesis of polyubiquitin chains. *Methods*
*Enzymol* **399**: 21-36. https://doi: 10.1016/S0076-6879(05)99002-2

Pohl C, and Dikic I. 2019. Cellular quality control by the ubiquitin-proteasome system and
autophagy. *Science* **366**: 818-822. https://doi: 10.1126/science.aax3769

Pollock L, Georgiou LC, Rusilowicz-Jones EV, Clague MJ, and Urbé S. 2024. A long-lived
pool of PINK1 imparts a molecular memory of depolarisation-induced activity. *bioRxiv*.
https://doi: 10.1101/2024.07.03.601901

Preus S, Noer SL, Hildebrandt LL, Gudnason D, and Birkedal V. 2015. iSMS: single-
molecule FRET microscopy software. *Nat Methods* **12**: 593-594. https://doi:
10.1038/nmeth.3435

Rai M, Hunt LC, and Demontis F. 2025. Stress responses induced by perturbation of the ubiquitin-proteasome system. *Trends Biochem Sci* **50**: 175-178. https://doi: 10.1016/j.tibs.2024.12.011

Ran FA, Hsu PD, Wright J, Agarwala V, Scott DA, and Zhang F. 2013. Genome engineering using the CRISPR-Cas9 system. *Nat Protoc* **8**: 2281-2308. https://doi: 10.1038/nprot.2013.143

Roy R, Hohng S, and Ha T. 2008. A practical guide to single-molecule FRET. *Nat Methods* **5**: 507-516. https://doi: 10.1038/nmeth.1208

Samant RS, Livingston CM, Sontag EM, and Frydman J. 2018. Distinct proteostasis circuits cooperate in nuclear and cytoplasmic protein quality control. *Nature* **563**: 407-411. https://doi: 10.1038/s41586-018-0678-x

Shen C, Chen C, Wang T, Gao TY, Zeng M, Lu YB, and Zhang WP. 2023. The Depletion of NAMPT Disturbs Mitochondrial Homeostasis and Causes Neuronal Degeneration in Mouse Hippocampus. *Mol Neurobiol* **60**: 1267-1280. https://doi: 10.1007/s12035-022-03142-5

Shi WZ, Qi LL, Fang SH, Lu YB, Zhang WP, and Wei EQ. 2012. Aggravated chronic brain injury after focal cerebral ischemia in aquaporin-4-deficient mice. *Neurosci Lett* **520**: 121-125. https://doi: 10.1016/j.neulet.2012.05.052

Shiba-Fukushima K, Arano T, Matsumoto G, Inoshita T, Yoshida S, Ishihama Y, Ryu KY, Nukina N, Hattori N, and Imai Y. 2014. Phosphorylation of mitochondrial polyubiquitin by PINK1 promotes Parkin mitochondrial tethering. *PLoS Genet* **10**: e1004861. https://doi: 10.1371/journal.pgen.1004861

Shiba-Fukushima K, Ishikawa KI, Inoshita T, Izawa N, Takanashi M, Sato S, Onodera O, Akamatsu W, Okano H, Imai Y, and Hattori N. 2017. Evidence that phosphorylated ubiquitin signaling is involved in the etiology of Parkinson's disease. *Hum Mol Genet* **26**: 3172-3185. https://doi: 10.1093/hmg/ddx201

Soykan T, Haucke V, and Kuijpers M. 2021. Mechanism of synaptic protein turnover and its regulation by neuronal activity. *Curr Opin Neurobiol* **69**: 76-83. https://doi: 10.1016/j.conb.2021.02.006

Swaney DL, Rodriguez-Mias RA, and Villen J. 2015. Phosphorylation of ubiquitin at Ser65 affects its polymerization, targets, and proteome-wide turnover. *EMBO Rep* **16**: 1131-1144. https://doi: 10.15252/embr.201540298

Swatek KN, and Komander D. 2016. Ubiquitin modifications. *Cell Res* **26**: 399-422. https://doi: 10.1038/cr.2016.39

Takatori S, Ito G, and Iwatsubo T. 2008. Cytoplasmic localization and proteasomal degradation of N-terminally cleaved form of PINK1. *Neurosci Lett* **430**: 13-17. https://doi: 10.1016/j.neulet.2007.10.019

Tang C, and Zhang WP. 2020. How Phosphorylation by PINK1 Remodels the Ubiquitin System: A Perspective from Structure and Dynamics. *Biochemistry* **59**: 26-33. https://doi: 10.1021/acs.biochem.9b00715

Tseng BP, Green KN, Chan JL, Blurton-Jones M, and LaFerla FM. 2008. Abeta inhibits the proteasome and enhances amyloid and tau accumulation. *Neurobiol Aging* **29**: 1607-1618. https://doi: 10.1016/j.neurobiolaging.2007.04.014

Wall CE, Rose CM, Adrian M, Zeng YJ, Kirkpatrick DS, and Bingol B. 2019. PPEF2 Opposes PINK1-Mediated Mitochondrial Quality Control by Dephosphorylating Ubiquitin. *Cell Rep* **29**: 3280-3292 e3287. https://doi: 10.1016/j.celrep.2019.10.130

Wang L, Cho YL, Tang Y, Wang J, Park JE, Wu Y, Wang C, Tong Y, Chawla R, Zhang J, Shi Y, Deng S, Lu G, Wu Y, Tan HW, Pawijit P, Lim GG, Chan HY, Zhang J, Fang L, Yu H,

Liou YC, Karthik M, Bay BH, Lim KL, Sze SK, Yap CT, and Shen HM. 2018. PTEN-L is a
novel protein phosphatase for ubiquitin dephosphorylation to inhibit PINK1-Parkin-mediated
mitophagy. *Cell Res* **28**: 787-802. https://doi: 10.1038/s41422-018-0056-0

Wauer T, Simicek M, Schubert A, and Komander D. 2015a. Mechanism of phospho-
ubiquitin-induced PARKIN activation. *Nature* **524**: 370-374. https://doi:
10.1038/nature14879

Wauer T, Swatek KN, Wagstaff JL, Gladkova C, Pruneda JN, Michel MA, Gersch M,
Johnson CM, Freund SM, and Komander D. 2015b. Ubiquitin Ser65 phosphorylation affects
ubiquitin structure, chain assembly and hydrolysis. *EMBO J* **34**: 307-325. https://doi:
10.15252/embj.201489847

Wilson DM, 3rd, Cookson MR, Van Den Bosch L, Zetterberg H, Holtzman DM, and
Dewachter I. 2023. Hallmarks of neurodegenerative diseases. *Cell* **186**: 693-714. https://doi:
10.1016/j.cell.2022.12.032

Wirths O. 2017. Extraction of Soluble and Insoluble Protein Fractions from Mouse Brains
and Spinal Cords. *Bio Protoc* **7**: e2422. https://doi: 10.21769/BioProtoc.2422

Wojcik C, and Di Napoli M. 2004. Ubiquitin-proteasome system and proteasome inhibition:
new strategies in stroke therapy. *Stroke* **35**: 1506-1518. https://doi:
10.1161/01.STR.0000126891.93919.4e

Yamano K, and Youle RJ. 2013. PINK1 is degraded through the N-end rule pathway.
*Autophagy* **9**: 1758-1769. https://doi: 10.4161/auto.24633

1197