Hippocampal place cell sequences are impaired in a rat model of Fragile X Syndrome

Margaret M. Donahue[1,2], Emma Robson[1,3], Laura Lee Colgin[1,2,3]

[1]Center for Learning and Memory, The University of Texas at Austin, Austin, TX 78712
[2]Institute for Neuroscience, The University of Texas at Austin, Austin, TX 78712
[3]Department of Neuroscience, The University of Texas at Austin, Austin, TX 78712

Abbreviated title: Spatial coding in a Fragile X Syndrome rat model

Corresponding author: Laura Lee Colgin, colgin@mail.clm.utexas.edu

Number of pages: 45; **Number of figures:** 8; **Number of tables:** 4

Number of words: Abstract: 176; Significance statement: 107; Introduction: 647; Discussion: 1495

Conflict of Interest: The authors declare no competing financial interests.

Acknowledgements:
This research was supported by the Department of Defense CDMRP award W81XWH1810314 (to L.L.C.) and National Institutes of Health awards R56MH125655 (to L.L.C), R01MH131317 (to L.L.C.), and F31MH127933 (to M.M.D.). The authors thank Isabella Lee, Misty Hill, Ayomide Akinsooto, and Sirisha Dhavala for technical assistance and Chenguang Zheng, Ernie Hwaun, and John Trimper for providing MATLAB code for some of the analyses. The authors acknowledge the Texas Advanced Computing Center (TACC) at The University of Texas at Austin for providing data storage resources that have contributed to the research described within this article. URL: http://www.tacc.utexas.edu

Author Contributions
M.M.D. and L.L.C. designed research; M.M.D. and E.R. performed research; M.M.D. analyzed data; M.M.D. and L.L.C. wrote the paper.

Abstract

Fragile X Syndrome (FXS) is a neurodevelopmental disorder that can cause impairments in spatial cognition and memory. The hippocampus is thought to support spatial cognition through the activity of place cells, neurons with spatial receptive fields. Coordinated firing of place cell populations is organized by different oscillatory patterns in the hippocampus during specific behavioral states. Theta rhythms organize place cell populations during awake exploration. Sharp wave-ripples organize place cell population reactivation during waking rest. Here, we examined the coordination of CA1 place cell populations during active behavior and subsequent rest in a rat model of FXS (*Fmr1* knockout rats). While the organization of individual place cells by the theta rhythm was normal, the coordinated activation of sequences of place cells during individual theta cycles was impaired in *Fmr1* knockout rats. Further, the subsequent replay of place cell sequences was impaired during waking rest following active exploration. Together, these results expand our understanding of how genetic modifications that model those observed in FXS affect hippocampal physiology and suggest a potential mechanism underlying impaired spatial cognition in FXS.

Significance Statement

Fragile X Syndrome (FXS) is a neurodevelopmental disorder that can cause impaired memory and atypical spatial behaviors such as "elopement" (i.e., wandering off and becoming lost). Activity in the CA1 subregion of the hippocampus supports spatial memory and spatial cognition, making it an important candidate to study in the context of FXS; however, how neuronal population activity in CA1 is affected by FXS is poorly understood. In this study, we found that the coordination of populations of CA1 neurons during active behavior and waking

rest was impaired in a rat model of FXS. These results reveal hippocampal physiological deficits

that may contribute to cognitive impairments in FXS.

Introduction

Fragile X Syndrome (FXS) is a widespread neurodevelopmental disorder that is caused by

epigenetic suppression of the X-chromosome linked *Fmr1* gene and subsequent loss of Fragile

X Messenger Ribonucleoprotein (FMRP) (De Boulle et al., 1993; Kremer et al., 1991; Mikiko et

al., 1995; Pieretti et al., 1991). Patients with FXS are impaired in memory tasks (Cornish et al.,

1998; Gallagher & Hallahan, 2012; Jäkälä et al., 1997) and often display elopement behaviors

(Machalicek et al., 2014; Muller et al., 2019). Mouse and rat models of FXS ("FXS mice" or "FXS

rats", respectively) have been produced by knocking out the *Fmr1* gene (Bakker et al., 1994;

Tian et al., 2017). Spatial memory deficits have been reported for both FXS mice and FXS rats

(Asiminas et al., 2019; Bakker et al., 1994; Mineur et al., 2002; Tian et al., 2017; Till et al., 2015;

Van Dam et al., 2000). FMRP is an RNA-binding protein (Siomi et al., 1993) that interacts with

many neuronal proteins (Darnell et al., 2011; for a review, see Darnell & Klann, 2013) and is

highly expressed in the hippocampus (Ludwig et al., 2014).

The hippocampus is thought to support memory and spatial cognition through the activity of

place cells, neurons with spatial receptive fields known as "place fields" (O'Keefe, 1976;

O'Keefe & Dostrovsky, 1971). Populations of place cells form sequences that represent

trajectories through an environment. These sequential firing patterns are coordinated by

hippocampal rhythms that are differentially associated with behavioral states. During active

exploration of an environment, place cells are coordinated by the theta rhythm, a ~6-10 Hz

oscillation occurring prominently in local field potential (LFP) recordings from the hippocampus

(Buzsáki, 2002). As an animal moves through a cell's place field, place cells fire at progressively

earlier phases of the theta rhythm in a phenomenon known as "theta phase precession"

(O'Keefe & Recce, 1993). Coordinated theta phase precession across multiple place cells with

adjacent place fields would be expected to result in an organized sequence of spikes within an

individual theta cycle that represent a rat's previous, current, and future locations. Such

sequential activation of place cells within a single theta cycle has been observed in rats and

termed a "theta sequence" (Foster & Wilson, 2007). Previous work has suggested that

hippocampal networks are hypersynchronized in FXS (Arbab, Battaglia, et al., 2018; Talbot et

al., 2018), which could impair organization of cells in theta sequences. However, the activity of

theta-coordinated sequences of place cells is yet to be investigated in a rodent model of FXS.

Sequences of place cells that were activated during active exploration of an environment are

reactivated or "replayed" in a time-compressed manner during subsequent waking rest and non-

REM sleep (Kudrimoti et al., 1999; Lee & Wilson, 2002). Place cell replay co-occurs with

characteristic events in the hippocampal LFP called sharp wave-ripples (Kudrimoti et al., 1999;

Lee & Wilson, 2002; Nádasdy et al., 1999). Sharp wave-ripples during sleep are abnormal in

FXS mice (Boone et al., 2018), suggesting that coordination of place cell populations during

sleep and rest may be disrupted in FXS. However, replay of place cell sequences has not been

examined in FXS models.

Here, we examined the activity of coordinated sequences of CA1 place cells in FXS rats

traversing a familiar circular track and subsequently resting. During active behaviors, we found

that the coordination of individual place cells by the theta oscillation was normal in FXS rats.

However, theta-coordinated sequences of place cells in FXS rats were less temporally

compressed and represented shorter paths than those in wild-type (WT) control rats. Further,

replay of place cell sequences during rest was abnormally slow in FXS rats, with replay events

exhibiting longer durations and representing less temporally compressed spatial trajectories

than in WT rats. These findings raise the possibility that the abnormal coordination of sequences

of hippocampal place cells may contribute to impairments in spatial memory and cognition in

FXS.

Materials and Methods

Additional data collected from one of the rats (rat 418) used in this study was presented in a

previous study (Robson et al., 2024). Surgery, data acquisition, histology, and spike sorting

methods in this paper are identical to methods described in that study and re-stated below.

Subjects

Twelve male Sprague-Dawley rats (Inotiv) were used for this study. Six rats were *Fmr1-*

knockout rats (*SD-Fmr1-nulltm1Sage*), and six were littermate WT control rats. As FXS is an X-

chromosome linked disorder, FXS has a higher prevalence and increased symptom severity in

males. Thus, we chose to use male rats for this study. Rats were between the ages of 3-11

months old at the time of surgery. Prior to surgery, rats were double- or triple-housed in

genotype-matched groups. After surgery, rats were singly housed in custom-built acrylic cages

(40 cm x 40 cm x 40 cm) containing enrichment material (wooden blocks, paper towel rolls, etc.)

and maintained on a reverse light cycle (light: 8 p.m. – 8 a.m.). Rats were housed next to their

former cage mates after recovering from surgery and throughout behavioral testing. Rats

recovered from surgery for at least one week before behavioral training resumed. All behavioral

experiments were performed during the dark cycle. When necessary to encourage spatial

exploration, one rat (rat 316) was placed on a food-deprivation regimen. While on the regimen,

this rat maintained ~98% of his free-feeding body weight. Following the completion of all

recording experiments, a small piece of ear tissue was collected from each rat to verify

genotype. All experiments were conducted according to the guidelines of the United States

National Institutes of Health Guide for the Care and Use of Laboratory Animals and under a

protocol approved by the University of Texas at Austin Institutional Animal Care and Use

Committee.

Surgery and tetrode positioning

"Hyperdrives" with 14 independently movable tetrodes were implanted in eight of the rats.

Hyperdrives with 21 independently movable tetrodes were implanted in four of the rats. Implants

were positioned above the right dorsal hippocampus (anterior-posterior -3.8 mm from bregma,

medial-lateral -3.0 mm from bregma). To implant and stabilize the hyperdrives, eleven bone

screws were affixed to the skull, and the base of the implant and the screws were covered in

dental acrylic. Two of the screws were connected to the recording drive ground. Prior to surgery,

tetrodes were built from 17 µm polyimide-coated platinum-iridium (90/10%) wire (California Fine

Wire, Grover Beach, California). The tips of tetrodes designated for single-unit recording were

plated with platinum to reduce single-channel impedances to ~150 to 300 kOhms. All tetrodes

were lowered ~1 mm on the day of surgery. Thereafter, tetrodes were slowly lowered to the

hippocampal pyramidal cell body layer over the course of several weeks except for one tetrode

that was designated for use as a reference for differential recording. This reference tetrode was

placed in an electrically quiet area approximately 1 mm above the hippocampus and adjusted

as needed to ensure quiescence. All four wires of this tetrode were connected to a single

channel on the electrode interface board. The reference signal was duplicated using an MDR-50

breakout board (Neuralynx, Bozeman, Montana) and recorded continuously to ensure that unit

activity or volume conducted signals of interest were not detected. Another tetrode was placed

in the apical dendritic layer of CA1 to monitor LFPs and guide placement of the other tetrodes

using estimated depth and electrophysiological hallmarks of the hippocampus (e.g., sharp

wave-ripples).

Data acquisition

Data were acquired using a Digital Lynx system and Cheetah recording software (Neuralynx,

Bozeman, Montana). The recording setup has been described in detail previously (Hsiao et al.,

2016; Zheng et al., 2016). Briefly, LFP signals from one randomly chosen channel per tetrode

were continuously recorded at a 2000 Hz sampling rate and filtered in the 0.1–500 Hz band.

Input amplitude ranges were adjusted before each recording session to maximize resolution

without signal saturation. Input ranges for LFPs generally fell within ±2,000 to ±3,000 μV. To

detect unit activity, all four channels within each tetrode were bandpass filtered from 600 to

6,000 Hz. Spikes were detected when the filtered continuous signal on one or more of the

channels exceeded a threshold of 55 μV. Detected events were acquired with a 32,000 Hz

sampling rate for 1 ms. For both LFPs and unit activity, signals were recorded differentially

against a dedicated reference channel (see "*Surgery and tetrode positioning*" section above).

Videos of rats' behavior were recorded through the Neuralynx system with a resolution of 720 ×

480 pixels and a frame rate of 29.97 frames/s. Rat position and head direction were tracked via

an array of red and green or red and blue light-emitting diodes (LEDs) on a HS-54 or HS-27

headstage (Neuralynx, Bozeman, Montana), respectively.

Behavior

Rats were trained to run unidirectionally on a 1-meter diameter circular track. The track was 0.5

meters in height. Rats ran four 10-minute sessions on the track per day. Inter-session rest

periods were 10 minutes, and the rat rested in a towel-lined flowerpot in the recording room

during rest sessions. Rats additionally rested in the pot for 10 minutes prior to the start of the

first track running session and after the final session for a total of five rest sessions per day.

Small pieces of sweetened cereal or cookies were placed at one or two locations on the circular

track to encourage running. The reward locations were kept consistent within each day but

changed daily in order to prevent accumulation of place fields at the reward site (Hollup et al.,

2001). To ensure that rats were familiarized with the environment prior to recording, rats

completed a minimum of 2 days of the full recording session (i.e., all five rest sessions and all

four run sessions) before data collection began. To compare behavior between genotypes, we

calculated run speed on the circular track and number of laps completed per session. To ensure

exclusion of pauses in running behavior and periods of immobility, only times when the rat was

traveling at speeds greater than 5 cm/s were included when calculating run speed.

Histology and tetrode localization

Following recording, rats were perfused with 4% paraformaldehyde solution in phosphate-

buffered saline. Brains were cut coronally in 30 µm sections using a cryostat. Brain slices were

immunostained for the CA2 marker Purkinje Cell Protein 4 (PCP4), allowing differentiation of all

subregions of the hippocampus (CA1, CA2, and CA3). Sections were initially washed and

blocked in 10% normal goat serum in TBS. Sections were incubated overnight with rabbit anti-

PCP4 (1:200, Sigma-Aldrich Cat# HPA005792) diluted in TBS containing 0.05% Tween. The

next day, sections were washed and incubated overnight with a secondary fluorescent antibody

(Alexa Flour™-555 anti-rabbit, Thermo Fisher Scientific). All washes and incubations were

performed at room temperature. Slides were mounted using DAPI Fluoromount-G (Fisher

Scientific). Tetrode recording sites were identified by comparing locations across adjacent

sections.

Spike sorting

Spike sorting was performed manually using graphical cluster-cutting software (MClust, A.D.

Redish, University of Minnesota, Minneapolis, Minnesota) run in MATLAB (Mathworks). Spikes

were sorted using two-dimensional representations of waveform properties (i.e., energy, peak,

and peak-to-valley difference) from four channels. A single unit was accepted for further analysis

if the associated cluster was well isolated from, and did not share spikes with, other clusters on

the same tetrode. Units were also required to have a minimum 1 ms refractory period. Units with

mean firing rates above 5 Hz were considered putative interneurons and excluded from further

analyses. To be included in the replay event firing analyses, a unit was required to have valid

clusters in both the active exploration and rest sessions. Place cell yields for each rat for the

active behavior and replay event analyses are reported in Tables 1 and 2, respectively.

Place cell analyses

Only cells from tetrodes in CA1 were included in analyses (see "*Histology and tetrode*

localization" section above). Firing rate maps were created for each single unit using methods

based on those used in our prior studies (Hwaun & Colgin, 2019; Zheng et al., 2021) (see

Figure 1A-B for example rate maps). The circular track was divided into 4-degree bins. The

number of spikes fired by a unit was divided by the time spent in each bin. Spikes that occurred

at times when a rat was moving at speeds less than 5 cm/s were excluded. This rate map was

then smoothed with a Gaussian kernel (standard deviation = 8 degrees). Rate maps were

calculated individually for each run session and for all run sessions in each day. To be included

for further analysis, the day-averaged rate map of a cell had to reach a minimum peak firing rate

of 1 Hz.

To examine stability of place cell rate maps, a Pearson correlation coefficient R ("spatial

correlation") was calculated for each unit between pairs of rate maps from different sessions

(Figure 2A). We additionally calculated rate overlap between pairs of rate maps from different

sessions for each cell to determine whether a cell's firing rate changed significantly across

sessions (Figure 2B). Rate overlap was calculated by taking the ratio of mean firing rates

between two sessions, with the larger firing rate as the denominator (Colgin et al., 2010).

Spatial information was calculated using the following formula (Skaggs et al., 1996):

$$Spatial\ information = \sum P_i \frac{\lambda_i}{\lambda} log2 \frac{\lambda_i}{\lambda}$$

where i is an index of spatial bins, P_i is the probability of a rat being in the ith bin, λ_i is the mean

firing rate in the ith bin, and λ is the overall mean firing rate of the cell (Figure 2C).

To identify place fields (Figure 2D-E), firing rates across locations in the rate map were first z-

scored. Potential place fields were identified as locations where the z-scored firing rate was

greater than or equal to 2. Identified place fields were bounded by locations where z-scored

firing rates fell below 0.5. To be included for further analysis, the peak firing rate in a place field

had to be at least 1 Hz, and the minimum length of a place field had to be at least 18 degrees

(~15 cm).

Phase precession analysis

Phase precession analysis was performed similarly to the analysis described in our previously

published work (Bieri et al., 2014). The LFP signal from every tetrode that recorded CA1 cells

identified as place cells was bandpass filtered in the theta range (i.e., between 6-10 Hz). The

phase of the theta oscillation was then estimated using a Hilbert transform. Locations within

each place field were normalized between 0 and 1. For each spike that a cell fired within its

place field on all sessions within a day, the theta phase at the spike time was estimated using

the transformed signal from the tetrode on which it was recorded. The normalized distance

through the place field at the spike time was also determined. Spikes that occurred while the rat

was traveling at a speed of less than 5 cm/s were excluded. A place cell had to fire a minimum

of 50 spikes within its place field in order to be included in theta phase precession analysis.

Circular-linear regression was then performed with theta phase as the circular variable and

normalized distance through the place field as the linear variable (Figure 3A-B). The correlation

coefficient for the relationship between theta phase and normalized distance (Figure 3C) was

calculated using the *circ_corrcl* function from the Circular Statistics toolbox in MATLAB (Berens,

2009) (https://www.mathworks.com/matlabcentral/fileexchange/10676-circular-statistics-toolbox-

directional-statistics).

Bayesian decoding analysis

We used a Bayesian decoding algorithm (K. Zhang et al., 1998) to estimate posterior probability

distributions of angular positions represented by the spiking activity of CA1 place cell

populations during track running. The probability of a rat being at position *x* given the number of

spikes *n* that occurred within a given time window was defined using Bayes' rule:

243
$$P(x|n) = \frac{P(n|x) * P(x)}{P(n)}$$

where *P(n | x)* was estimated using the averaged position tuning of each unit across all four

sessions. The probability of a rat being at any given position on the circular track *P(x)* was set to

1 to avoid biasing the decoder towards any particular position on the circular track. The

normalizing constant *P(n)* was set such that the posterior probability distribution *P(x | n)*

summed to 1.

Decoding accuracy analysis

To verify that a given day's place cell yields were sufficient to accurately decode a rat's positions

on the circular track, we used a decoding accuracy analysis similar to the methods described in

our previously published work (Hwaun & Colgin, 2019; Zheng et al., 2021). Positions were

decoded for all times when a rat was traveling at speeds over 5 cm/s in 500 ms windows with

100 ms steps. To create confusion matrices (Figure 4A-B), the average decoded probability for

all times when a rat was at a given position was calculated. To determine the decoding error, the

decoded position was defined as the position with maximal decoded probability for each time

bin. The decoding error was then defined as the difference between a rat's actual position and

the corresponding decoded position. The cumulative error distributions for each day were then

determined (Figure 4A-B). For a day of recordings to be included in theta sequence event and

replay event analysis, its cumulative error distribution had to reach 50% at error values less than

20 degrees (Davidson et al., 2009; Hwaun & Colgin, 2019). Days that reached this criterion are

shown in Figure 4A-B. The overall error distributions for rats with days that met this criterion are

shown in Figure 4C.

Detection and analysis of theta sequences

Theta sequences were detected only on days that reached the decoding criterion (see

"*Decoding accuracy analysis"* section above). Theta sequences were detected and

characterized using methods similar to our previously published procedures (Zheng et al.,

2016). LFP signals were band-pass filtered in the theta band (6-10 Hz), and individual theta

cycles were cut at the theta phase with the lowest number of spikes. Bayesian decoding was

then performed on theta cycles in which at least 3 place cells fired and when the rat was

traveling at a speed of 5 cm/s or greater (see "*Bayesian decoding analysis"* section above).

Bayesian decoding was performed across partially overlapping 40 ms windows advanced by a

10 ms step (Figure 5A-B). If spikes did not occur throughout the entirely of the theta cycle,

contiguity was considered to be broken when there were two consecutive time bins (i.e., 50 ms)

without spikes. Sequence property and significance analyses were then performed on the

longest set of contiguous time bins (Figure 5C-F). The t-span of a sequence was defined as the

temporal duration of these time bins. To determine the slope and x-span of the sequence event,

we calculated the center of mass of the posterior probability distribution $P(x \mid n)$ from the

Bayesian decoding for each time bin. A circular-linear regression line was fit to these positions in

order to determine the slope. The x-span was defined as the distance between the first and last

positions of the regression line. To determine if a theta sequence was significant, we compared

the r^2 value from this circular-linear regression to a shuffled distribution. This distribution was

obtained by circularly shuffling each time bin of the weighted probability distribution (i.e., *P(x | n)*

from the Bayesian decoding analysis) a random distance 1000 times and then obtaining an r^2

value for each shuffled distribution. The r^2 value from the theta sequence had to exceed 95% of

the shuffled values in order to be considered significant. Additionally, to ensure the regression

provided an accurate representation of the true decoded probability distribution for the event, at

least 60% of the total posterior probability needed to be within 20 degrees of the fitted circular-

linear regression line (Zheng et al., 2021). In addition, the minimum distance between the fitted

trajectory and the actual position of the rat had to be less than 20 degrees (Zheng et al., 2021).

The number of theta sequences from each rat is shown in Table 3.

Detection and characterization of replay events

Replay events were detected only on days that reached the decoding criterion (see "*Decoding*

accuracy analysis" section above). Replay events were detected using methods similar to

previously published procedures (Hwaun & Colgin, 2019; Pfeiffer & Foster, 2013). Replay

events were detected while the rat rested off of the circular track in a towel-lined flowerpot (see

"*Behavior*" section above). To detect candidate events, a histogram of population firing rates

was constructed using all cells that were classified as active during the track running sessions

(see "*Place cell analyses*" section above). This histogram was then smoothed with a Gaussian

kernel (standard deviation = 10 ms). Candidate events were detected when the population firing

rate exceeded 3 standard deviations above the mean population firing rate and were bounded

by crossing of the mean. Events within 40 ms of each other were combined. Start and end times

were then adjusted inward so that the first and last time bins of a candidate event each

contained at least one spike. To be included for further analysis, an event had to be between 50

and 2000 ms in duration, and at least 5 cells had to fire during an event.

For each candidate event, Bayesian decoding was performed (see "*Bayesian decoding*

analysis" section above) across partially overlapping 20 ms windows advanced by a 10 ms step

(Figure 6A-B). To assess how well a posterior probability distribution represented an actual

trajectory on the circular track, circular-linear regression was performed with the decoded

position as the circular variable and time within the event as the linear variable. The decoded

position for each time bin was defined as the center of mass of each time bin, which was

determined by taking the circular mean of positions weighted by their associated posterior

probability values. The r^2 value of the regression line was used as an assessment of replay

fidelity (Figure 6C), as in previous studies (Davidson et al., 2009; Hwaun & Colgin, 2019;

Karlsson & Frank, 2009). To be classified as a significant replay event, the r^2 value of an event's

regression line had to be at least 0.5 (as in Hwaun & Colgin, 2019), and the decoded positions

between adjacent time bins (i.e., the "jump" distance between adjacent time bins) could not

exceed 25% of the length of the circular track (as in Berners-Lee et al., 2022). Only replay

events that occurred after the first track running session of the day (i.e., in rest sessions 2-5)

were included for further analysis. To quantify the temporal compression of a replay event, we

estimated the slope of the circular-linear regression line (Figure 6E). We calculated the absolute

value of the slope to include both forward and reverse replay events. The path distance

represented within a replay event was calculated as the difference between the decoded

positions of the first and last time bins of a replay event (Figure 6F).

Replay event place cell analyses

To investigate the firing of place cells during replay events, we binned the firing rates of CA1

place cells in 1 ms bins around times of replay event onset for each unit (as in Hwaun & Colgin,

2019). We only included replay events in which a unit participated (as in Boone et al., 2018).

This allowed us to examine differences in place cell firing patterns during replay events while

controlling for differences in the number of replay events in which a unit participates. We

smoothed each firing rate vector with a Gaussian kernel (standard deviation = 10 ms). We

averaged across all replay events to obtain a binned mean firing rate vector for each unit

(Figure 7A). Additionally, we determined the average firing rate for each unit during all replay

events in which a unit participated (Figure 7B) and the average number of spikes per event that

each unit fired (Figure 7C) (as in Boone et al., 2018).

To determine whether the spike timing of place cells during replay events was impaired in FXS

rats, we calculated the interval between spikes in a replay event in two ways (Figure 7D). For

the first method, the "population inter-spike interval", we calculated the average interval between

consecutive spikes from all cells in a replay event for each replay event (Figure 7E). In the

second method, the "first spike inter-spike interval", we only considered the first spike that each

cell fired in a replay event and then computed the average interval between consecutive times

of first spikes for each replay event (Figure 7F).

Replay event LFP analyses

The LFP from the tetrode with the most place cells on a given day was used for each day's

analysis of power associated with replay events. The time-varying power across frequencies

around the time of replay event onset was computed using a wavelet transform (Tallon-Baudry

et al., 1997) as previously described (Mably et al., 2017). Time-varying power around replay

event onset was calculated in 1 Hz steps from 2-250 Hz and averaged across all replay events

within a day. Power was z-scored within each frequency band. A time-varying power vector for

each frequency was created for each rat by averaging across recording days. The plots

presented in Figure 8A-B represent the average time-frequency representations of power

associated with replay events across all rats for each genotype. The peak ripple frequency for a

replay event (Figure 8C) was defined as the frequency with the highest power in the ripple band,

150-250 Hz. To calculate an overall estimate of slow gamma power associated with a replay

event (Figure 8D), we averaged power estimates across the slow gamma band of frequencies

(25-55 Hz) and across time for each replay event.

To assess the relationship between the duration of replay events and number of sharp wave-

ripples (Figure 8E), we used methods adapted from a previous study (Davidson et al., 2009).

The LFP from the tetrode with the most place cells from a given day was bandpass filtered from

150-250 Hz, and a Hilbert transform was then applied to the filtered signal. The absolute value

of the Hilbert transformed signal was then smoothed with a Gaussian kernel (standard deviation

= 8 ms) and z-scored. Ripples were detected at times when the z-score was greater than or

equal to 3. The number of ripples is displayed with a small amount of random jitter on the y-axis

for visualization (Figure 8E) (as in Davidson et al., 2009).

Data visualization

Whenever possible, data is shown for each animal individually. In Figures 2, 3, 6, 7, and 8, each

data point represents one measure (i.e., for an individual place cell or replay event). Boxes

show 95% confidence intervals of the mean values within each rat. Confidence intervals were

calculated with 1000 bootstrapped samples. WT rats are shown with blue data points, and FXS

rats are shown with red data points. In each plot for each genotype, data from individual rats

were presented in an order corresponding to increasing mean values for ease of comparison

across genotypes. In Figures 4 and 5, distributions are shown due to the large number of data

points for each genotype. In Figure 5, shaded areas represent 95% confidence intervals

calculated with 1000 bootstrapped samples.

Experimental design and statistical analyses

Whenever possible, experimenters were blinded to genotype during data acquisition and

analyses. However, due to temporarily limited availability of FXS rats from the supplier,

experiments on two cohorts of rats were performed unblinded (rats 416, 418, 442, and 445).

MATLAB (Mathworks) scripts were custom written for the analyses in this study, based on

algorithms that have been used in prior studies, as described above. All statistical tests were

performed using SPSS Statistics (Version 29.0, IBM) unless otherwise stated. To compare

behavior, place cell properties, decoding errors, and replay properties across genotypes, we

used a generalized linear mixed model design (as in Robson et al., 2024). Individual rats were

subjects, and genotype was included as a fixed factor. When comparing behavior metrics,

sessions were nested within days as repeated measures. When applicable, place cells were

nested within rats (analyses in Figures 2, 3, and 7A-C). When comparing different session pairs

to assess place cell firing rate map stability (Figure 2A-B), session pair was included as a

repeated measure. To compare decoding errors between genotypes (Figure 4C), the error

calculated for each time bin in which a rat was moving faster than 5 cm/s (see "*Decoding*

accuracy analysis" section above) was nested within sessions, and sessions were nested within

days as repeated measures. For analyses of replay event properties (Figures 6, 7E-F, and 8C-

D), individual replay events were nested within the rest session in which they occurred, and rest

sessions were nested within a day as repeated measures (as in Boone et al., 2018).

To compare theta sequence properties between genotypes (Figure 5C-F), we used the non-

parametric Wilcoxon rank-sum test (*ranksum* function in MATLAB).

To account for potential differences in decoding accuracy across rats when analyzing theta

sequence slope and replay event slope, we used a generalized linear model (*fitglm* function in

MATLAB) with a log-link function. The model included genotype and decoding errors for each

session as predictors of slope values to determine if decoding accuracy significantly affected

the results (similar to analysis of effects of decoding accuracy in Hwaun & Colgin, 2019).

To assess whether the correlation between replay duration and number of sharp wave-ripples

differed across genotypes (Figure 8E), we used multiple linear regression. Genotype, replay

event duration, and the interaction between genotype and replay event duration were included

as predictors.

Code and data accessibility

Analysis code is available on GitHub

(https://github.com/mmdonahue/FMR1CircTrack_PublicShare). Data will be made available

upon reasonable request.

Results

Hippocampal place cells showed similar activity in FXS and WT rats

To determine how place cell coding of spatial locations is affected in a rat model of FXS, we

recorded from the CA1 pyramidal cell layer of the hippocampus in adult FXS and WT rats. We

recorded place cell activity while rats ran unidirectionally on a circular track in a familiar room for

four ten-minute sessions per day (Figure 1). WT and FXS rats ran at similar speeds on the

circular track (generalized linear mixed model, WT: estimated mean = 34 cm/s, estimated 95%

CI = 28-39 cm/s; FXS: estimated mean = 27 cm/s, estimated 95% CI = 22-33 cm/s; no

significant main effect of genotype ($F_{(1,98)}$ = 2.5, p = 0.116)), However, FXS rats completed

fewer laps per session on average (generalized linear mixed model, WT: estimated mean = 17

laps, estimated 95% CI = 13-22 laps; FXS: estimated mean = 10 laps, estimated 95% CI = 5-15

laps; significant main effect of genotype ($F_{(1,98)}$ = 4.4, p = 0.039), suggesting that FXS rats

paused on the track more than WT rats.

Previous work in a different FXS rat model (i.e., Long Evans *Fmr1* knockout rats) found no

difference in spatial stability of CA1 place cells when rats explored an initially novel environment

over two days (Asiminas et al., 2022). However, work from FXS mice reported impaired short-

term stability in CA1 place cells (Arbab, Pennartz, et al., 2018). To examine place cell stability in

the present FXS rat model, we computed spatial correlation coefficients between pairs of track

running sessions for each place cell. Spatial correlations were lower in FXS rats than in WT

rats, suggesting impaired place field stability in FXS rats (Figure 2A; generalized linear mixed

model, no significant interaction between genotype and session pair ($F_{(3,2036)}$ = 0.399, p =

0.754), significant main effect of genotype ($F_{(1,2036)}$ = 4.596, p = 0.032)). Further, place cell

firing rates changed more between sessions in FXS rats than in WT rats (Figure 2B; generalized

linear mixed model, no significant interaction between genotype and session pair ($F_{(3,2036)}$ =

0.615, p = 0.605), significant main effect of genotype ($F_{(1,2036)}$ = 69.542, p < 0.001)). The

seeming discrepancy between our results and the relatively stable place fields observed in

another rat model of FXS (Asiminas et al., 2022) may be due to differences in testing

environments or rat strains.

The previous study of CA1 place cell activity in a different FXS rat model than the one used in

the present study showed no impairments in place cells' spatial information, mean firing rates, or

place field size (Asiminas et al., 2022). Here, we also found no differences between FXS and

WT rats in CA1 place cells' spatial information (Figure 2C; generalized linear mixed model, no

significant main effect of genotype ($F_{(1,436)}$ = 3.674, p = 0.056)), peak firing rates (Figure 2D;

generalized linear mixed model, no significant main effect of genotype ($F_{(1,458)}$ = 0.316, p =

0.574)), and place field size (Figure 2E; generalized linear mixed model, no significant main

effect of genotype (F(1,458) < 0.001, p = 0.995)). This indicates that CA1 place cells in FXS rats

can represent different locations within a session in a familiar environment as well as CA1 place

cells in WT control rats.

Theta modulation of place cells was normal in FXS rats

During active exploration, the activity of place cells is coordinated by the ~6-10 Hz theta rhythm

in the hippocampus. Theta phase precession refers to a place cell firing pattern in which

successive place cell spikes occur at progressively earlier phases of the theta cycle as a rat

traverses the cell's place field (O'Keefe & Recce, 1993). Manipulations that reduce theta phase

precession (Robbe & Buzsáki, 2009) and disrupt the precise spike timing of neurons during

theta sequences (Petersen & Buzsáki, 2020) reduce spatial memory performance, raising the

possibility that abnormal theta phase precession contributes to spatial memory deficits that have

been reported in FXS models (Asiminas et al., 2019; Bakker et al., 1994; Mineur et al., 2002;

Tian et al., 2017; Till et al., 2015; Van Dam et al., 2000). We examined whether CA1 place cells

exhibit abnormal phase precession in FXS rats (Figure 3). We found no difference between FXS

and WT rats in the relationship between theta phase and relative position in the place field

(Figure 3C; generalized linear mixed model, no significant main effect of genotype (F(1,423) =

0.795, p = 0.373)), suggesting that theta coordination of spiking is normal in FXS rats at the

level of individual cells.

Theta sequences spanned shorter distances in FXS rats

Other work has led to the hypothesis that individual place cells are largely normal in rodent

models of FXS while coordinated activity of large populations of place cells is impaired (Talbot et

al., 2018). Studies in FXS mice have suggested that hippocampal networks are

hypersynchronized in FXS (Arbab, Battaglia, et al., 2018; Talbot et al., 2018). FXS mice also

showed abnormally low coordination between place cells with overlapping place fields (Talbot et

al., 2018). Together, these prior results suggest that coordinated theta sequences of place cells

may be impaired in FXS rats. To assess theta sequences in FXS rats, we used a Bayesian

decoding approach (K. Zhang et al., 1998). Specifically, we applied Bayesian decoding to

spiking activity from place cell populations during active running sessions to reconstruct

positions on the circular track represented by place cell populations. We first determined how

accurately we were able to reconstruct the rat's positions during running using the place cell

spiking activity. Most rats' recordings (WT rats 326, 392, 416, and 418 and FXS rats 316, 330,

395, and 442) surpassed our criteria for sufficient decoding accuracy (Figure 4A-B; see also

"*Decoding accuracy analysis*" section in Materials and Methods). Also, decoding errors were not

significantly different between genotypes (Figure 4C; generalized linear mixed model, no

significant main effect of genotype (F(1,573941) = 0.17, p = 0.678)).

For the animals that surpassed decoding accuracy criteria, we used Bayesian decoding to

identify trajectories represented by place cell populations within individual theta cycles as WT

and FXS rats ran laps unidirectionally on a circular track (Figure 5A-B; for the number of theta

sequences per rat, see Table 3). We then performed circular-linear regression to fit a regression

line to the spatial path represented within these theta cycles. This regression line allowed us to

quantify the temporal compression (slope, in cm/s), distance ("x-span", in cm), and duration ("t-

span", in s) of each trajectory. We found that the slopes of theta sequences were lower in FXS

rats (Figure 5C; Wilcoxon rank-sum test, significant effect of genotype (Z = 3.7, p < 0.001)). To

account for potential differences in decoding accuracy across different rats and days, we used a

generalized linear model with a log-link function to predict slope values from genotype and the

decoding errors for each session. Including genotype in the model increased prediction

accuracy while including decoding errors had no significant effect on the model (genotype, β =

0.2, p = 0.007; decoding accuracy, β = 0.08, p = 0.32), suggesting that the difference in the

temporal compression of theta sequences was not due to differences in decoding accuracy

between genotypes. The difference in slope values reflected both a decrease in the distances of

paths represented during theta sequences (Figure 5D; Wilcoxon rank-sum test, significant effect

of genotype (Z = 2.2, p = 0.029)) and an increase in the duration of events (Figure 5E; Wilcoxon

rank-sum test, significant effect of genotype (Z = 3.6, p < 0.001)). We also calculated the r^2

value of the regression line fit to decoded sequence representations in order to assess the

extent to which sequences represented coherent trajectories. We found no significant difference

in the r^2 values of the circular-linear regression lines for sequences from WT and FXS rats

(Figure 5F; Wilcoxon rank-sum test, no significant effect of genotype (Z = 1.9, p = 0.062)).

These results suggest that theta sequences represented spatial trajectories in FXS rats.

However, theta sequences represented shorter trajectories, and representations were less

temporally compressed, in FXS rats compared to WT rats.

Replay events were less temporally compressed in FXS rats

Replay events co-occur with sharp wave-ripples in the LFP of the hippocampus during NREM

sleep or waking rest (Davidson et al., 2009; Kudrimoti et al., 1999; Lee & Wilson, 2002;

Nádasdy et al., 1999). Previous work has shown that sharp wave-ripples during NREM sleep

are abnormally long in duration in FXS mice (Boone et al., 2018). To assess replay events in

FXS rats, we examined the firing of place cell sequences while rats rested quietly in a location

off the circular track after running. We used Bayesian decoding to reconstruct positions on the

circular track represented by CA1 place cell populations during replay events in WT and FXS

rats (Figure 6A-B; for the number of replay events recorded from each rat, see Table 4). We

quantified replay fidelity, duration, and temporal compression of each replay event using

protocols similar to previously published procedures (Davidson et al., 2009; Hwaun & Colgin,

2019; Karlsson & Frank, 2009). Specifically, we first applied circular-linear regression analysis to

the posterior probability distributions resulting from Bayesian decoding of place cell spikes

during replay events. We then assessed associated r^2 values as a measure of replay fidelity,

and the slopes of the fitted lines were used to estimate the temporal compression of replay

events. There was no difference in replay fidelity between WT and FXS rats (Figure 6C;

generalized linear mixed model, no significant effect of genotype ($F(1,751) = 2.185$, $p = 0.140$)).

However, the duration of replay events was longer in FXS rats than in WT rats (Figure 6D;

generalized linear mixed model, significant effect of genotype ($F(1,751) = 33.332$, $p < 0.001$)).

Further, the representations during replay events were less temporally compressed, indicating

slower transitions across representations of successive locations, in FXS rats than in WT rats

(Figure 6E; generalized linear mixed model, significant effect of genotype ($F(1,751) = 19.340$, p

< 0.001)). To account for potential differences in decoding accuracy across different rats and

days, we used a generalized linear model with a log-link function to predict slope values from

genotype and the decoding errors for each session. Including genotype in the model increased

prediction accuracy while including decoding errors had no significant effect on the model

(genotype, $\beta = 0.25$, $p < 0.001$; decoding accuracy, $\beta = 0.06$, $p = 0.82$), suggesting that the

difference in temporal compression of replay events was not due to different decoding accuracy

between genotypes. The low temporal compression in FXS rats did not indicate that replay

events represented unusually long paths in FXS rats, however, because path lengths of replay

sequences were similar in FXS and WT rats (Figure 6F; generalized linear mixed model, no

significant effect of genotype ($F(1,751) = 1.3$, $p = 0.253$). Instead, this pattern of results

suggests that replay events in FXS rats represented trajectories of similar lengths and

comparable fidelity as in WT rats but that replay of representations of trajectories occurred more

slowly in FXS rats than in WT rats.

Place cells fired more slowly during replay events in FXS rats

In addition to differences in duration, Boone and colleagues (Boone et al., 2018) observed

abnormal place cell firing during sharp wave-ripples in a mouse model of FXS. They found that

place cells had lower in-event firing rates during sharp wave-ripples recorded during sleep,

although individual cells fired the same number of spikes per event (Boone et al., 2018).

Abnormally slow firing of place cells during replay events in FXS rats may underlie the reduced

temporal compression observed in our data. Indeed, we found that CA1 place cells had lower

peak firing rates after replay event onset in FXS rats than in WT rats (Figure 7A; generalized

linear mixed model, significant effect of genotype on peak firing rate after event onset (F(1,444)

= 8.877, p = 0.003)). In-event firing rates were also lower in FXS rats than in WT rats (Figure

7B; generalized linear mixed model, significant effect of genotype (F(1,444) = 23.140, p <

0.001)), while the number of spikes each cell fired during a replay event did not differ between

FXS and WT rats (Figure 7C; generalized linear mixed model, no significant effect of genotype

(F(1,444) = 1.020, p = 0.313)) likely because of replay events' relatively long durations in FXS

rats (Figure 6D). The low replay-associated firing rates in individual place cells in FXS rats

suggest that dynamics of CA1 place cell sequence firing may occur more slowly during replay

events in FXS rats than in WT rats. To test this hypothesis, we calculated the distribution of

intervals between successive spikes in a replay event in two ways (Figure 7D). First, we

considered all spikes that occurred across all cells during a replay event and found that these

population inter-spike intervals were longer in FXS rats than in WT rats (Figure 7E; generalized

linear mixed model, significant effect of genotype (F(1,751) = 58.337, p < 0.001)). Next, we only

considered the first spike that each cell in a sequence fired in a replay event (Figure 7F). These

first spike inter-spike intervals were also longer in FXS rats than in WT rats (generalized linear

mixed model, significant effect of genotype (F(1,751) = 56.843, p < 0.001)).

Properties of ripples and slow gamma rhythms during replay events in FXS rats

We further examined properties of oscillatory patterns in the LFP, specifically sharp wave-ripples

and slow gamma oscillations, that normally co-occur with replay events in WT rats (Davidson et

al., 2009; Kudrimoti et al., 1999; Lee & Wilson, 2002; Nádasdy et al., 1999; Carr et al., 2012;

Bieri et al., 2014). In both genotypes, we saw increases in power in the ripple and slow gamma

bands at event onset (Figure 8A-B), suggesting that ripples and slow gamma rhythms co-occur

with replay events in FXS rats as in WT rats. However, the peak ripple frequency during replay

events was lower in FXS rats than in WT rats (Figure 8C; generalized linear mixed model,

significant effect of genotype ($F_{(1,751)}$ = 21.168, $p < 0.001$)), consistent with results from FXS

mice showing lower peak ripple frequency during sharp wave-ripples during sleep (Boone et al.,

2018). Although the frequency of ripples was lower in FXS rats, the number of sharp wave-

ripples that co-occurred with a replay event of a given duration was similar in FXS and WT rats

(Figure 8E; significant multiple linear regression ($F_{(3,749)}$ = 145.149, $p < 0.001$); no significant

interaction between genotype and replay event duration ($t = 0.369$, $p = 0.712$)), likely due to the

longer duration of replay events in FXS rats than in WT rats (Figure 6D).

Previous work has shown increased slow gamma power in CA1 during sleep-associated sharp

wave-ripples in a FXS mouse model (Boone et al., 2018). Here, we found no difference in slow

gamma power during rest-associated replay events in WT and FXS rats (Figure 8D; generalized

linear mixed model, no significant effect of genotype ($F_{(1,751)}$ = 0.095, $p = 0.758$)). Replay

events during sleep and waking rest may have different functions and characteristics (Roumis &

Frank, 2015; Tang et al., 2017) and may involve input from different upstream structures to CA1

(Yamamoto & Tonegawa, 2017). Thus, differences in rest and sleep states may underlie the

seemingly discrepant results for replay-associated slow gamma in FXS rat and mouse models.

Discussion

To our knowledge, this is the first study that examines spatial representations coded by large populations of hippocampal place cells in a rodent model of FXS. Here, we present data showing that coding of spatial trajectories by coordinated place cell populations is impaired in FXS rats. Specifically, we found that theta sequences coded less temporally compressed representations and represented shorter path distances in FXS rats, while theta phase precession in individual place cells was normal. Further, we found that place cell sequences fired abnormally slowly during hippocampal replay events in FXS rats. This slow place cell firing was associated with reduced temporal compression of representations of track trajectories during replay in FXS rats. Together, these results suggest that coordinated place cell sequences code spatial trajectories abnormally during both active running and during subsequent rest in a rat model of FXS.

Previous studies examining the activity of place cells during active exploration in FXS rodents have yielded mixed results. Studies from FXS mice and rats have shown no difference in spatial information in CA1 place cells (Arbab, Pennartz, et al., 2018; Asiminas et al., 2022; Talbot et al., 2018), although one study has shown abnormally large place field sizes and excessive out-of-field firing in FXS mice (Arbab, Pennartz, et al., 2018). Discrepancies between these results and the present findings may result from different experimental paradigms, including different testing arenas, experimental time courses, and the degree of environmental novelty. Regarding the latter point, work from FXS rats has shown that experience-dependent reductions in firing rates and sharpening of spatial tuning of place cells are lacking in FXS rats after introduction to a novel environment (Asiminas et al., 2022). These results suggest that the initial formation of a spatial memory may be stunted in FXS rats. We did not observe such differences in our study, likely due to our use of a testing environment with which rats had already been familiarized.

Work from FXS mice has shown normal theta modulation in CA1 place cells but reduced

correlated firing of pairs of place cells (Talbot et al., 2018). This suggested that coordinated

population coding would be impaired in FXS rodents. Consistent with this result, we show that

theta-coordinated sequences of place cells during active exploration represented shorter

trajectories in FXS rats, while theta phase precession in individual place cells remained normal.

While theta phase precession and theta sequences appear to be related phenomena, it is

possible to observe the absence of theta sequences in populations of place cells that

individually show intact phase precession (Feng et al., 2015; Middleton & McHugh, 2016).

Computational models have suggested that alterations in the coupling between place cells and

local CA1 interneurons may disrupt the temporal compression of theta sequences while leaving

phase precession intact (Chadwick et al., 2016). Interneurons are less modulated by theta in

FXS mice (Talbot et al., 2018), and CA1 interneurons and pyramidal cells are less correlated in

FXS mice (Arbab, Battaglia, et al., 2018). This interneuron-pyramidal cell dysfunction may result

in hyper-synchronization between CA1 pyramidal neurons and thereby cause place cells that

represent multiple locations along a trajectory to fire more synchronously in FXS rats. This could

flatten slopes of theta sequences and shorten distances of represented trajectories.

Disrupting the input from CA3 to CA1 may also affect the path length of theta sequences

(Chadwick et al., 2016; Middleton & McHugh 2016). *In vitro* work examining the Schaffer

collateral pathway from CA3 to CA1 in FXS has yielded mixed results. Work from FXS mice has

shown an abnormally low induction threshold for long-term potentiation (LTP) in FXS mice when

pre- and post-synaptic neurons are simultaneously activated (Routh et al., 2013). However,

other work from FXS mice (Lauterborn et al., 2007) and rats (Tian et al., 2017) has reported

reduced Schaffer collateral LTP. FXS mouse models have also shown increased metabotropic

glutamate receptor (mGluR) long-term depression (LTD) (Huber et al., 2002; Iliff et al., 2013; J.

Zhang et al., 2009), suggesting that hippocampal synapses may be relatively weak in rodent

models of FXS. Such a reduction in synaptic input from CA3 to CA1 may affect the temporal

compression of theta sequences in FXS rats, resulting in representation of relatively short paths.

Our work shows that CA1 place cells fired more slowly in FXS rats than in WT rats during replay

events that occurred during waking rest. This finding is consistent with abnormally slow firing of

pyramidal cells during sharp wave-ripples in non-REM sleep reported in FXS mice (Boone et al.,

2018). *In vitro* work from FXS rats has also shown lower frequency multi-unit activity during

sharp wave-ripples specifically in dorsal hippocampus (Leontiadis et al., 2023). Parvalbumin-

expressing inhibitory interneurons in the hippocampus are important for pacing pyramidal cell

spiking during sharp wave-ripples (Schlingloff et al., 2014; Stark et al., 2014), and hippocampal

network models have suggested that inhibition within CA1 is important for controlling cell

participation within replay events (Ramirez-Villegas et al., 2018). Reported weaknesses in

pyramidal cell-interneuron coupling in CA1 have only been examined during active behavior in

FXS models (Arbab, Battaglia, et al., 2018). However, considering that ripple oscillations are

locally generated in CA1 (Csicsvari et al., 1999), the lower peak ripple frequencies that we

found during replay events may suggest that firing of CA1 inhibitory interneurons is also slowed

during ripples in FXS rats. Due to the limited number of interneurons in our dataset, we were not

able to test this hypothesis directly in the present study.

Both CA3 and CA2 contribute to generation of sharp wave-ripples and replay events in CA1

(Csicsvari et al., 1999, 2000; Oliva et al., 2016). Thus, deficits in CA1 place cell firing in FXS

rats during replay events may be inherited from these upstream regions or due to local deficits

in CA1. Future studies involving simultaneous recordings from CA3, CA2, and CA1 are

necessary to shed light on circuit mechanisms underlying slowed firing of place cells during

replay events in FXS rats.

In addition to inputs to CA1 from CA3 and CA2, input from the medial entorhinal cortex (MEC) to

CA1 can be important for replay events, particularly replay events that span multiple sharp

wave-ripples (Yamamoto & Tonegawa, 2017). Previous work has suggested that synaptic inputs

from MEC to CA1 pyramidal cells are reduced in FXS models (Asiminas et al., 2022; Ordemann

et al., 2021). Although we did not observe a difference in the number of ripples co-occurring with

replay events of a given duration between FXS and WT rats (Figure 8E), a reduction in MEC

input during replay events may affect temporal compression of replayed sequences for

extended replay events in CA1.

Downstream targets of the hippocampus may be affected by impaired temporal compression of

awake replay events. During replay events, activity between hippocampus and prefrontal cortex

is coordinated (Berners-Lee et al., 2021; Harvey et al., 2023; Jadhav et al., 2016; Peyrache et

al., 2009; Shin et al., 2019; Tang et al., 2017). This coordinated activity has been hypothesized

to support memory retrieval processes that can be used to guide future decision making

(Jadhav et al., 2016; Zielinski et al., 2020). The strength of excitatory drive from the

hippocampus to the prefrontal cortex during sharp wave-ripples can affect the response of the

prefrontal cortex (Wierzynski et al., 2009), suggesting that abnormally slowed spike timing

during replay events in CA1 of FXS rats may alter subsequent prefrontal responses.

The work presented shows novel evidence for specific physiological impairments in the

hippocampus in a rat model of Fragile X Syndrome. However, these impairments were

characterized using a simple behavioral protocol with no memory component. Rats were

familiarized to the environment before recording, and food rewards were randomly given without

any motivational salience for specific spatial trajectories. Previous work suggests that the

temporal compression of place cell sequences during both active behavior and awake rest can

contribute to spatial learning and memory. The slopes and strength of theta sequences have

been reported to increase during learning of new environments or trajectories to a new goal

location (Feng et al., 2015; Igata et al., 2021; Pfeiffer, 2022), and theta sequences exhibited

higher slopes during correct trials than error trials of a spatial memory task (Zheng et al., 2021).

Replay duration and temporal compression have been linked to learning and memory in studies

of healthy WT rats (Berners-Lee et al., 2022; Fernández-Ruiz et al., 2019). Replay duration

increased, and temporal compression of replay events decreased, across the first several paths

that rats took during learning of a new environment (Berners-Lee et al., 2022). However, on a

longer time scale, the duration of replay events decreased, while the lengths of trajectory

representations increased, across sessions in rats learning a spatial memory task (Shin et al.,

2019). These results raise the possibility that less temporally compressed theta sequences and

slow replay could contribute to impaired spatial learning and memory in FXS rats. Future studies

of place cell population activity in FXS rats engaged in learning and memory tasks will be

important to shed light on this question.

Tables

Table 1. Total number of place cells recorded from each rat during active behavior.

Genotype	Rat	Number of place cells
WT	326 ("Zachariah")	94
WT	334 ("Chickpea")	4
WT	335 ("Couscous")	5
WT	392 ("Danish")	29
WT	416 ("Desmond")	60
WT	418 ("Hugo")	71
FXS	316 ("Yuki")	21

FXS	330 ("Aries")	99
FXS	394 ("Mr. Eko")	3
FXS	395 ("Elijah")	36
FXS	445 ("Paddy")	20
FXS	442 ("Pippin")	69

742   Table 2. Total number of place cells recorded from each rat during replay events.

Genotype	Rat	Number of place cells
WT	326 ("Zachariah")	93
WT	392 ("Danish")	16
WT	416 ("Desmond")	60
WT	418 ("Hugo")	71
FXS	316 ("Yuki")	21
FXS	330 ("Aries")	98
FXS	395 ("Elijah")	18
FXS	442 ("Pippin")	69

744   Table 3. Total number of theta sequences recorded from each rat.

Genotype	Rat	Number of sequences
WT	326 ("Zachariah")	3151
WT	392 ("Danish")	110
WT	416 ("Desmond")	1617
WT	418 ("Hugo")	2505
FXS	316 ("Yuki")	600
FXS	330 ("Aries")	5032

| FXS | 395 ("Elijah") | 461 |
| FXS | 442 ("Pippin") | 2392 |

746   Table 4. Total number of replay events from each rat.

Genotype	Rat	Number of replay events
WT	326 ("Zachariah")	114
WT	392 ("Danish")	28
WT	416 ("Desmond")	116
WT	418 ("Hugo")	166
FXS	316 ("Yuki")	23
FXS	330 ("Aries")	110
FXS	395 ("Elijah")	24
FXS	442 ("Pippin")	172

References

Arbab, T., Battaglia, F. P., Pennartz, C. M. A., & Bosman, C. A. (2018). Abnormal hippocampal

theta and gamma hypersynchrony produces network and spike timing disturbances in the

Fmr1-KO mouse model of Fragile X syndrome. *Neurobiology of Disease*, *114*(1), 65–73.

https://doi.org/10.1016/j.nbd.2018.02.011

Arbab, T., Pennartz, C. M. A., & Battaglia, F. P. (2018). Impaired hippocampal representation of

place in the Fmr1-knockout mouse model of fragile X syndrome. *Scientific Reports*, *8*(1),

1–9. https://doi.org/10.1038/s41598-018-26853-z

Asiminas, A., Booker, S. A., Dando, O. R., Kozic, Z., Arkell, D., Inkpen, F. H., Sumera, A.,

Akyel, I., Kind, P. C., & Wood, E. R. (2022). Experience-dependent changes in

hippocampal spatial activity and hippocampal circuit function are disrupted in a rat model of

Fragile X Syndrome. *Molecular Autism*, *13*(1), 1–29. https://doi.org/10.1186/s13229-022-00528-z

Asiminas, A., Jackson, A. D., Louros, S. R., Till, S. M., Spano, T., Dando, O., Bear, M. F., Chattarji, S., Hardingham, G. E., Osterweil, E. K., Wyllie, D. J. A., Wood, E. R., & Kind, P. C. (2019). Sustained correction of associative learning deficits after brief, early treatment in a rat model of Fragile X Syndrome. *Science Translational Medicine*, *11*(494), 1–10. https://doi.org/10.1126/scitranslmed.aao0498

Bakker, C. E., Verheij, C., Willemsen, R., van der Helm, R., Oerlemans, F., Vermey, M., Bygrave, A., Hoogeveen, A. T., Oostra, B. A., Reyniers, E., De Boule, K., D'Hooge, R., Cras, P., van Velzen, D., Nagels, G., Martin, J. J., De Deyn, P. P., Darby, J. K., & Willems, P. J. (1994). Fmr1 knockout mice: A model to study fragile X mental retardation. *Cell*, *78*(1), 23–33. https://doi.org/10.1016/0092-8674(94)90569-X

Berens, P. (2009). CircStat: A MATLAB Toolbox for Circular Statistics. *Journal of Statistical Software*, *31*(10), 1–21. https://doi.org/10.18637/jss.v031.i10

Berners-Lee, A., Feng, T., Silva, D., Wu, X., Ambrose, E. R., Pfeiffer, B. E., & Foster, D. J. (2022). Hippocampal replays appear after a single experience and incorporate greater detail with more experience. *Neuron*, *110*(11), 1829-1842.e5. https://doi.org/10.1016/j.neuron.2022.03.010

Berners-Lee, A., Wu, X., & Foster, D. J. (2021). Prefrontal cortical neurons are selective for non-local hippocampal representations during replay and behavior. *Journal of Neuroscience*, *41*(27), 5894–5908. https://doi.org/10.1523/JNEUROSCI.1158-20.2021

Bieri, K. W., Bobbitt, K. N., & Colgin, L. L. (2014). Slow and fast gamma rhythms coordinate different spatial coding modes in hippocampal place cells. *Neuron*, *82*(3), 670–681. https://doi.org/10.1016/j.neuron.2014.03.013

Boone, C. E., Davoudi, H., Harrold, J. B., & Foster, D. J. (2018). Abnormal sleep architecture and hippocampal circuit dysfunction in a mouse model of Fragile X Syndrome.

Neuroscience, *384*, 275–289. https://doi.org/10.1016/j.neuroscience.2018.05.012

Buzsáki, G. (2002). Theta oscillations in the hippocampus. *Neuron*, *33*(3), 325–340.

https://doi.org/10.1016/S0896-6273(02)00586-X

Chadwick, A., Van Rossum, M. C. W., & Nolan, M. F. (2016). Flexible theta sequence

compression mediated via phase precessing interneurons. *ELife*, *5*(DECEMBER2016).

https://doi.org/10.7554/ELIFE.20349

Colgin, L. L., Leutgeb, S., Jezek, K., Leutgeb, J. K., Moser, E. I., McNaughton, B. L., & Moser,

M. B. (2010). Attractor-map versus autoassociation based attractor dynamics in the

hippocampal network. *Journal of Neurophysiology*, *104*(1), 35–50.

https://doi.org/10.1152/jn.00202.2010

Cornish, K. M., Munir, F., & Cross, G. (1998). The nature of the spatial deficit in young females

with Fragile-X syndrome: A neuropsychological and molecular perspective.

Neuropsychologia, *36*(11), 1239–1246. https://doi.org/10.1016/S0028-3932(97)00162-0

Csicsvari, J., Hirase, H., Czurkó, A., Mamiya, A., & Buzsáki, G. (1999). Fast network oscillations

in the hippocampal CA1 region of the behaving rat. *The Journal of Neuroscience : The*

Official Journal of the Society for Neuroscience, *19*(16), RC20.

https://doi.org/10.1523/jneurosci.19-16-j0001.1999

Csicsvari, J., Hirase, H., Mamiya, A., & Buzsáki, G. (2000). Ensemble patterns of hippocampal

CA3-CA1 neurons during sharp wave-associated population events. *Neuron*, *28*(2), 585–

594. https://doi.org/10.1016/S0896-6273(00)00135-5

Darnell, J. C., & Klann, E. (2013). The translation of translational control by FMRP: therapeutic

targets for FXS. *Nature Neuroscience 2013 16:11*, *16*(11), 1530–1536.

https://doi.org/10.1038/nn.3379

Darnell, J. C., Van Driesche, S. J., Zhang, C., Hung, K. Y. S., Mele, A., Fraser, C. E., Stone, E.

F., Chen, C., Fak, J. J., Chi, S. W., Licatalosi, D. D., Richter, J. D., & Darnell, R. B. (2011).

FMRP stalls ribosomal translocation on mRNAs linked to synaptic function and autism.

Cell, *146*(2), 247–261. https://doi.org/10.1016/j.cell.2011.06.013

Davidson, T. J., Kloosterman, F., & Wilson, M. A. (2009). Hippocampal replay of extended experience. *Neuron*, *63*(4), 497–507. https://doi.org/10.1016/j.neuron.2009.07.027

De Boulle, K., Verkerk, A. J. M. H., Reyniers, E., Vits, L., Hendrickx, J., Van Roy, B., Van Den Bos, F., de Graaff, E., Oostra, B. A., & Willems, P. J. (1993). A point mutation in the FMR-1 gene associated with fragile X mental retardation. *Nature Genetics*, *3*(1), 31–35. https://doi.org/10.1038/ng0193-31

Feng, T., Silva, D., & Foster, D. J. (2015). Dissociation between the experience-dependent development of hippocampal theta sequences and single-trial phase precession. *Journal of Neuroscience*, *35*(12), 4890–4902. https://doi.org/10.1523/JNEUROSCI.2614-14.2015

Fernández-Ruiz, A., Oliva, A., de Oliveira, E. F., Rocha-Almeida, F., Tingley, D., & Buzsáki, G. (2019). Long-duration hippocampal sharp wave ripples improve memory. *Science*, *364*(6445), 1082–1086. https://doi.org/10.1126/science.aax0758

Foster, D. J., & Wilson, M. A. (2007). Hippocampal theta sequences. *Hippocampus*, *17*(11), 1093–1099. https://doi.org/10.1002/hipo.20345

Gallagher, A., & Hallahan, B. (2012). Fragile X-associated disorders: A clinical overview. In *Journal of Neurology* (Vol. 259, Issue 3, pp. 401–413). Springer. https://doi.org/10.1007/s00415-011-6161-3

Harvey, R. E., Robinson, H. L., Liu, C., Oliva, A., & Fernandez-Ruiz, A. (2023). Hippocampo-cortical circuits for selective memory encoding, routing, and replay. *Neuron*, *111*(13), 2076-2090.e9. https://doi.org/10.1016/j.neuron.2023.04.015

Hollup, S. A., Molden, S., Donnett, J. G., Moser, M. B., & Moser, E. I. (2001). Accumulation of hippocampal place fields at the goal location in an annular watermaze task. *Journal of Neuroscience*, *21*(5), 1635–1644. https://doi.org/10.1523/jneurosci.21-05-01635.2001

Huber, K. M., Gallagher, S. M., Warren, S. T., & Bear, M. F. (2002). Altered synaptic plasticity in a mouse model of fragile X mental retardation. *Proceedings of the National Academy of*

Sciences of the United States of America, *99*(11), 7746–7750.

https://doi.org/10.1073/PNAS.122205699/ASSET/4D74A266-B9D1-4EEC-A089-

AB9F614C8070/ASSETS/GRAPHIC/PQ1222056004.JPEG

Hwaun, E., & Colgin, L. L. (2019). CA3 place cells that represent a novel waking experience are

preferentially reactivated during sharp wave-ripples in subsequent sleep. *Hippocampus*,

29(10), 921–938. https://doi.org/10.1002/hipo.23090

Igata, H., Ikegaya, Y., & Sasaki, T. (2021). Prioritized experience replays on a hippocampal

predictive map for learning. *Proceedings of the National Academy of Sciences of the*

United States of America, *118*(1), e20112.

https://doi.org/10.1073/PNAS.2011266118/SUPPL_FILE/PNAS.2011266118.SM02.MP4

Iliff, A. J., Renoux, A. J., Krans, A., Usdin, K. U., Sutton, M. A., & Todd, P. K. (2013). Impaired

activity-dependent FMRP translation and enhanced mGluR-dependent LTD in Fragile X

premutation mice. *Human Molecular Genetics*, *22*(6), 1180–1192.

https://doi.org/10.1093/HMG/DDS525

Jadhav, S. P. P., Rothschild, G., Roumis, D. K. K., & Frank, L. M. M. (2016). Coordinated

excitation and inhibition of prefrontal ensembles during awake hippocampal sharp-wave

ripple events. *Neuron*, *90*(1), 113–127. https://doi.org/10.1016/j.neuron.2016.02.010

Jäkälä, P., Hänninen, T., Ryynänen, M., Laakso, M., Partanen, K., Mannermaa, A., & Soininen,

H. (1997). Fragile-X: Neuropsychological test performance, CGG triplet repeat lengths, and

hippocampal volumes. *Journal of Clinical Investigation*, *100*(2), 331–338.

https://doi.org/10.1172/JCI119538

Karlsson, M. P., & Frank, L. M. (2009). Awake replay of remote experiences in the

hippocampus. *Nature Neuroscience*, *12*(7), 913–918. https://doi.org/10.1038/nn.2344

Kremer, E. J., Pritchard, M., Lynch, M., Yu, S., Holman, K., Baker, E., Warren, S. T.,

Schlessinger, D., Sutherland, G. R., & Richards, R. I. (1991). Mapping of DNA instability at

the fragile X to a trinucleotide repeat sequence p(CCG)n. *Science*, *252*(5013), 1711–1714.

https://doi.org/10.1126/science.1675488

Kudrimoti, H. S., Barnes, C. A., & McNaughton, B. L. (1999). Reactivation of hippocampal cell assemblies: Effects of behavioral state, experience, and EEG dynamics. *Journal of Neuroscience*, *19*(10), 4090–4101. https://doi.org/10.1523/jneurosci.19-10-04090.1999

Lauterborn, J. C., Rex, C. S., Kramár, E., Chen, L. Y., Pandyarajan, V., Lynch, G., & Gall, C. M. (2007). Brain-derived neurotrophic factor rescues synaptic plasticity in a mouse model of Fragile X Syndrome. *Journal of Neuroscience*, *27*(40), 10685–10694. https://doi.org/10.1523/JNEUROSCI.2624-07.2007

Lee, A. K., & Wilson, M. A. (2002). Memory of sequential experience in the hippocampus during slow wave sleep. *Neuron*, *36*(6), 1183–1194. https://doi.org/10.1016/S0896-6273(02)01096-6

Leontiadis, L. J., Trompoukis, G., Tsotsokou, G., Miliou, A., Felemegkas, P., & Papatheodoropoulos, C. (2023). Rescue of sharp wave-ripples and prevention of network hyperexcitability in the ventral but not the dorsal hippocampus of a rat model of fragile X syndrome. *Frontiers in Cellular Neuroscience*, *17*, 1296235. https://doi.org/10.3389/FNCEL.2023.1296235/BIBTEX

Ludwig, A. L., Espinal, G. M., Pretto, D. I., Jamal, A. L., Arque, G., Tassone, F., Berman, R. F., & Hagerman, P. J. (2014). CNS expression of murine fragile X protein (FMRP) as a function of CGG-repeat size. *Human Molecular Genetics*, *23*(12), 3228–3238. https://doi.org/10.1093/HMG/DDU032

Mably, A. J., Gereke, B. J., Jones, D. T., & Colgin, L. L. (2017). Impairments in spatial representations and rhythmic coordination of place cells in the 3xTg mouse model of Alzheimer's disease. *Hippocampus*, *27*(4), 378–392. https://doi.org/10.1002/hipo.22697

Machalicek, W., McDuffie, A., Oakes, A., Ma, M., Thurman, A. J., Rispoli, M. J., & Abbeduto, L. (2014). Examining the operant function of challenging behavior in young males with fragile X syndrome: A summary of 12 cases. *Research in Developmental Disabilities*, *35*(7),

1694–1704. https://doi.org/10.1016/j.ridd.2014.03.014

Middleton, S. J., & McHugh, T. J. (2016). Silencing CA3 disrupts temporal coding in the CA1

ensemble. *Nature Neuroscience*, *19*(7), 945–951. https://doi.org/10.1038/nn.4311

Mikiko, C. S., Siomi, H., Sauer, W. H., Srinivasan, S., Nussbaum, R. L., & Dreyfuss, G. (1995).

FXR1, an autosomal homolog of the fragile X mental retardation gene. *EMBO Journal*,

14(11), 2401–2408. https://doi.org/10.1002/j.1460-2075.1995.tb07237.x

Mineur, Y. S., Sluyter, F., De Wit, S., Oostra, B. A., & Crusio, W. E. (2002). Behavioral and

neuroanatomical characterization of the Fmr1 knockout mouse. *Hippocampus*, *12*(1), 39–

46. https://doi.org/10.1002/hipo.10005

Muller, K., Brady, N. C., Warren, S. F., & Fleming, K. K. (2019). Mothers' perspectives on

challenging behaviours in their children with fragile X syndrome. *Journal of Intellectual and*

Developmental Disability, *44*(4), 481–491. https://doi.org/10.3109/13668250.2018.1496379

Nádasdy, Z., Hirase, H., Czurkó, A., Csicsvari, J., & Buzsáki, G. (1999). Replay and time

compression of recurring spike sequences in the hippocampus. *Journal of Neuroscience*,

19(21), 9497–9507. https://doi.org/10.1523/jneurosci.19-21-09497.1999

O'Keefe, J. (1976). Place units in the hippocampus of the freely moving rat. *Experimental*

Neurology, *51*(1), 78–109. https://doi.org/10.1016/0014-4886(76)90055-8

O'Keefe, J., & Dostrovsky, J. (1971). The hippocampus as a spatial map. Preliminary evidence

from unit activity in the freely-moving rat. *Brain Research*, *34*(1), 171–175.

https://doi.org/10.1016/0006-8993(71)90358-1

O'Keefe, J., & Recce, M. L. (1993). Phase relationship between hippocampal place units and

the EEG theta rhythm. *Hippocampus*, *3*(3), 317–330.

https://doi.org/10.1002/hipo.450030307

Oliva, A., Fernández-Ruiz, A., Buzsáki, G., & Berényi, A. (2016). Role of hippocampal CA2

region in triggering sharp-wave ripples. *Neuron*, *91*(6), 1342–1355.

https://doi.org/10.1016/j.neuron.2016.08.008

Ordemann, G. J., Apgar, C. J., Chitwood, R. A., & Brager, D. H. (2021). Altered A-type

potassium channel function impairs dendritic spike initiation and temporoammonic long-

term potentiation in Fragile X syndrome. *Journal of Neuroscience*, *41*(27).

https://doi.org/10.1523/JNEUROSCI.0082-21.2021

Peyrache, A., Khamassi, M., Benchenane, K., Wiener, S. I., & Battaglia, F. P. (2009). Replay of

rule-learning related neural patterns in the prefrontal cortex during sleep. *Nature*

Neuroscience, *12*(7), 919–926. https://doi.org/10.1038/nn.2337

Pfeiffer, B. E. (2022). Spatial learning drives rapid goal representation in hippocampal ripples

without place field accumulation or goal-oriented theta sequences. *Journal of*

Neuroscience, *42*(19), 3975–3988. https://doi.org/10.1523/JNEUROSCI.2479-21.2022

Pfeiffer, B. E., & Foster, D. J. (2013). Hippocampal place-cell sequences depict future paths to

remembered goals. *Nature*, *497*(7447), 74–79. https://doi.org/10.1038/nature12112

Pieretti, M., Zhang, F., Fu, Y. H., Warren, S. T., Oostra, B. A., Caskey, C. T., & Nelson, D. L.

(1991). Absence of expression of the FMR-1 gene in fragile X syndrome. *Cell*, *66*(4), 817–

822. https://doi.org/10.1016/0092-8674(91)90125-I

Ramirez-Villegas, J. F., Willeke, K. F., Logothetis, N. K., & Besserve, M. (2018). Dissecting the

synapse- and frequency-dependent network mechanisms of in vivo hippocampal sharp

wave-ripples. *Neuron*, *100*(5), 1224-1240.e13.

https://doi.org/10.1016/j.neuron.2018.09.041

Robson, E., Donahue, M. M., Mably, A. J., Demetrovich, P. G., Hewitt, L. T., & Colgin, L. L.

(2024). Social odors drive hippocampal CA2 place cell responses to social stimuli.

Progress in Neurobiology, 102708. https://doi.org/10.1016/j.pneurobio.2024.102708

Roumis, D. K., & Frank, L. M. (2015). Hippocampal sharp-wave ripples in waking and sleeping

states. *Current Opinion in Neurobiology*, *35*, 6–12.

https://doi.org/10.1016/j.conb.2015.05.001

Routh, B. N., Johnston, D., & Brager, D. H. (2013). Loss of functional A-type potassium

channels in the dendrites of CA1 pyramidal neurons from a mouse model of Fragile X

Syndrome. *Journal of Neuroscience*, *33*(50), 19442–19450.

https://doi.org/10.1523/JNEUROSCI.3256-13.2013

Schlingloff, D., Káli, S., Freund, T. F., Hájos, N., & Gulyás, A. I. (2014). Mechanisms of sharp

wave initiation and ripple generation. *Journal of Neuroscience*, *34*(34), 11385–11398.

https://doi.org/10.1523/JNEUROSCI.0867-14.2014

Shin, J. D., Tang, W., & Jadhav, S. P. (2019). Dynamics of awake hippocampal-prefrontal

replay for spatial learning and memory-guided decision making. *Neuron*, *104*(6), 1110-

1125.e7. https://doi.org/10.1016/j.neuron.2019.09.012

Siomi, H., Siomi, M. C., Nussbaum, R. L., & Dreyfuss, G. (1993). The protein product of the

fragile X gene, FMR1, has characteristics of an RNA-binding protein. *Cell*, *74*(2), 291–298.

https://doi.org/10.1016/0092-8674(93)90420-U

Skaggs, W. E., McNaughton, B. L., Wilson, M. A., & Barnes, C. A. (1996). Theta phase

precession in hippocampal neuronal populations and the compression of temporal

sequences. *Hippocampus*, *6*(2), 149–172. https://doi.org/10.1002/(SICI)1098-

1063(1996)6:2<149::AID-HIPO6>3.0.CO;2-K

Stark, E., Roux, L., Eichler, R., Senzai, Y., Royer, S., & Buzsáki, G. (2014). Pyramidal cell-

interneuron interactions underlie hippocampal ripple oscillations. *Neuron*, *83*(2), 467–480.

https://doi.org/10.1016/j.neuron.2014.06.023

Talbot, Z. N., Sparks, F. T., Dvorak, D., Curran, B. M., Alarcon, J. M., & Fenton, A. A. (2018).

Normal CA1 place fields but discoordinated network discharge in a Fmr1-null mouse model

of Fragile X Syndrome. *Neuron*, *97*(3), 684-697.e4.

https://doi.org/10.1016/j.neuron.2017.12.043

Tallon-Baudry, C., Bertrand, O., Delpuech, C., & Pernier, J. (1997). Oscillatory γ-band (30–70

Hz) activity induced by a visual search task in humans. *Journal of Neuroscience*, *17*(2),

722–734. https://doi.org/10.1523/JNEUROSCI.17-02-00722.1997

Tang, W., Shin, J. D., Frank, L. M., & Jadhav, S. P. (2017). Hippocampal-prefrontal reactivation

during learning is stronger in awake compared with sleep states. *Journal of Neuroscience*,

37(49), 11789–11805. https://doi.org/10.1523/JNEUROSCI.2291-17.2017

Tian, Y., Yang, C., Shang, S., Cai, Y., Deng, X., Zhang, J., Shao, F., Zhu, D., Liu, Y., Chen, G.,

Liang, J., Sun, Q., Qiu, Z., & Zhang, C. (2017). Loss of FMRP impaired hippocampal long-

term plasticity and spatial learning in rats. *Frontiers in Molecular Neuroscience*, *10*, 269.

https://doi.org/10.3389/fnmol.2017.00269

Till, S. M., Asiminas, A., Jackson, A. D., Katsanevaki, D., Barnes, S. A., Osterweil, E. K., Bear,

M. F., Chattarji, S., Wood, E. R., Wyllie, D. J. A., & Kind, P. C. (2015). Conserved

hippocampal cellular pathophysiology but distinct behavioural deficits in a new rat model of

FXS. *Human Molecular Genetics*, *24*(21), 5977–5984.

https://doi.org/10.1093/HMG/DDV299

Van Dam, D., D'Hooge, R., Hauben, E., Reyniers, E., Gantois, I., Bakker, C. E., Oostra, B. A.,

Kooy, R. F., & De Deyn, P. P. (2000). Spatial learning, contextual fear conditioning and

conditioned emotional response in Fmr1 knockout mice. *Behavioural Brain Research*,

117(1–2), 127–136. https://doi.org/10.1016/S0166-4328(00)00296-5

Wierzynski, C. M., Lubenov, E. V., Gu, M., & Siapas, A. G. (2009). State-dependent spike-

timing relationships between hippocampal and prefrontal circuits during sleep. *Neuron*,

61(4), 587–596. https://doi.org/10.1016/j.neuron.2009.01.011

Yamamoto, J., & Tonegawa, S. (2017). Direct medial entorhinal cortex input to hippocampal

CA1 is crucial for extended quiet awake replay. *Neuron*, *96*(1), 217-227.e4.

https://doi.org/10.1016/J.NEURON.2017.09.017

Zhang, J., Hou, L., Klann, E., & Nelson, D. L. (2009). Altered hippocampal synaptic plasticity in

the Fmr1 gene family knockout mouse models. *Journal of Neurophysiology*, *101*(5), 2572–

2580. https://doi.org/10.1152/jn.90558.2008

Zhang, K., Ginzburg, I., McNaughton, B. L., & Sejnowski, T. J. (1998). Interpreting neuronal

population activity by reconstruction: Unified framework with application to hippocampal

place cells. *Journal of Neurophysiology*, *79*(2), 1017–1044.

https://doi.org/10.1152/jn.1998.79.2.1017

Zheng, C., Bieri, K. W., Hsiao, Y. T., & Colgin, L. L. (2016). Spatial sequence coding differs

during slow and fast gamma rhythms in the hippocampus. *Neuron, 89*(2), 398–408.

https://doi.org/10.1016/j.neuron.2015.12.005

Zheng, C., Hwaun, E., Loza, C. A., & Colgin, L. L. (2021). Hippocampal place cell sequences

differ during correct and error trials in a spatial memory task. *Nature Communications*,

12(1), 3373. https://doi.org/10.1038/s41467-021-23765-x

Zielinski, M. C., Tang, W., & Jadhav, S. P. (2020). The role of replay and theta sequences in

mediating hippocampal-prefrontal interactions for memory and cognition. *Hippocampus*,

30(1), 60–72. https://doi.org/10.1002/hipo.22821

1005

1006   **Figure Legends**

1007

1008   Figure 1. Example CA1 place cells across four recording sessions from WT and FXS rats. (A-B)

1009   Firing rates across positions on the circular track are shown for all place cells (rows) recorded

1010   from one example tetrode across all four sessions (columns) in an example day from a WT (A)

1011   and a FXS (B) rat. (C-D) Example spike raster plots for successive laps within a day for a WT

1012   (C) and a FXS (D) place cell. Dashed lines separate different sessions.

1013

Figure 2. Place cell firing rate maps in FXS rats were unstable across sessions but otherwise

exhibit normal place field properties. (A) Spatial correlation coefficients are shown across

sessions pairs for each rat. Spatial correlation values were lower in FXS rats than WT rats,

indicating impaired stability of place cell responses in FXS rats. (B) Rate overlap values are

shown across session pairs for each rat. Rate overlap values were lower in FXS rats than in WT

rats, indicating highly variable firing rates of place cells across sessions in FXS rats. (C) There

was no difference in spatial information of place cells between WT and FXS rats. (D) Peak firing

rates of place cells did not differ between WT and FXS rats. (E) Place field size did not differ

between WT and FXS rats. For all plots, each dot represents a measure from one place cell.

Boxes represent 95% confidence intervals of the mean for each rat.

1024

Figure 3. Theta phase precession was preserved in FXS rats. (A-B) Place cell phase precession

plots are shown for two example place cells from (A) WT and (B) FXS rats. Each dot represents

the theta phase associated with each spike and corresponding normalized position in the cell's

place field. The solid line represents the correlation between the theta phase and normalized

position in place field. The magnitude of the correlation (r) is shown (top right) for each cell. (C)

The correlation between theta phase and normalized position in a place field did not differ for

place cells from WT and FXS rats. Each dot represents the correlation measure for one cell.

Boxes represent 95% confidence intervals of the mean for each rat.

1033

Figure 4. Decoding accuracy in WT and FXS rats. (A-B) Cumulative decoding error and

confusion matrices are shown for decoded place cell populations from WT (A) and FXS (B) rats.

Individual lines on cumulative error plots represent individual days. Insets show confusion

matrices from each day for each rat. Only days that met the decoding criteria are shown. (C)

There was no difference in decoding error values between WT and FXS rats. Each line

represents the distribution of decoding error values from one rat.

1040

Figure 5. Theta sequence events coded paths that were less temporally compressed and

shorter in FXS rats than in WT rats. (A-B) Example theta sequence events are shown for WT (A)

and FXS (B) rats. Position (on the y-axis) is shown relative to the mean of the rat's actual

position during the theta sequence event (indicated by white dashed line). The associated r^2 and

slope values are shown for each event. (C) Theta sequences' slopes were lower in FXS rats

than WT rats, indicating that theta sequences were less temporally compressed in FXS rats. (D)

The x-span values of theta sequences were lower in FXS rats than in WT rats, indicating that

theta sequences represented relatively short spatial paths in FXS rats. (E) The duration (t-span)

of theta sequences was higher in FXS rats than in WT rats. (F) There was no difference in r^2

values between FXS and WT rats, indicating that theta sequences represented coherent paths

through an environment in FXS rats despite the reduced temporal compression of

representations of these paths. (C-F) The distributions for theta sequence properties are shown

with shaded areas representing 95% confidence intervals of the distributions for each genotype.

1054

Figure 6. Replay events were less temporally compressed in FXS rats than in WT rats. (A-B)

Example replay events are shown for WT (A) and FXS (B) rats. The r^2 value and slope of each

replay event is shown above each plot. (C) Replay events' r^2 values did not differ between WT

and FXS rats. (D) Replay event durations were longer in FXS rats than in WT rats. (E) The

slopes of the regression lines fit to posterior probability distributions of replay events were lower

in FXS rats than in WT rats. (F) Path distances of replayed trajectories did not differ between

WT and FXS rats. (C-F) Each dot represents a measure for one replay event. Boxes represent

95% confidence intervals for the mean values from each rat.

1063

Figure 7. Place cells fired more slowly during replay events in FXS rats than in WT rats. (A)

Place cells from WT rats reached a higher peak firing rate after replay event onset than place

cells from FXS rats. Shaded areas represent 95% confidence intervals of the binned firing rate

distributions. (B) Place cell firing rates during replay events were higher in WT rats than in FXS

rats. (C) The number of spikes a place cell fired during replay events did not differ between WT

and FXS rats. (D) Schematic illustrating how population inter-spike interval (ISI) and first spike

ISI were calculated. An LFP recording during an example replay event from a WT rat is shown

(top). A raster plot shows spiking activity of 8 place cells that participated in the replay event,

with spikes from different cells represented by different colored tick marks (middle). The bottom

two rows show spikes included when calculating the average ISI for all spikes (population ISI)

and only the first spike from each cell (first spike ISI). (E) Population ISIs were higher in FXS

rats than in WT rats. (F) First spike ISIs were higher in FXS rats than in WT rats. (B-C) Each dot

represents a measure for an individual place cell. Boxes represent 95% confidence intervals for

the mean values for each rat. (E-F) Each dot represents one replay event. Boxes represent 95%

confidence intervals for the mean values for each rat.

1079

Figure 8. Peak ripple frequency was lower in FXS rats than in WT rats. (A-B) Time-frequency

representations of power during replay events in WT (A) and FXS (B) rats. (C) Peak ripple

frequency was lower in FXS rats than in WT rats. (D) Slow gamma power during replay events

(25-55 Hz) did not differ between WT and FXS rats. (C-D) Each dot represents a measure for

one replay event. Boxes represent 95% confidence intervals for the mean values for each rat.

(E) The number of ripples that co-occurred with replay events of a given duration did not differ

between WT (left) and FXS (right) rats. Each dot represents a measure for one replay event.

The dashed line represents the regression line fit to the distribution of points for each genotype.

1088

A

Sessions 1-2   Sessions 2-3   Sessions 3-4   Sessions 1-4

Spatial correlation

● WT
● FXS   *

Rat #   Rat #   Rat #   Rat #

B

Sessions 1-2   Sessions 2-3   Sessions 3-4   Sessions 1-4

Rate overlap

● WT
● FXS   ***

Rat #   Rat #   Rat #   Rat #

C   ● WT ● FXS

n.s.

Spatial information (bits/spike)

Rat #

D   ● WT ● FXS

n.s.

Peak firing rate (Hz)

Rat #

E   ● WT ● FXS

n.s.

Place field size (cm)

Rat #

A WT

rat 326

rat 416

rat 392

rat 418

Probability

B FXS

rat 316

rat 395

rat 330

rat 442

Probability

C

n.s.

WT   FXS
— rat 326   — rat 316
— — rat 392 — — rat 330
— rat 416   — rat 395
— — rat 418 — — rat 442