Host genetics and the Skin Microbiome Independently Predict Parasite Resistance

Rachael D. Kramp[1*], Mary J. Janecka[1,5], Nadine Tardent[2,3], Jukka Jokela[2,3], Kevin D. Kohl[1], Jessica F. Stephenson[1,2,3,4]

1. Department of Biological Sciences, University of Pittsburgh, Pittsburgh, PA 15260, USA

2. Department of Aquatic Ecology, EAWAG, Swiss Federal Institute of Aquatic Sci & Tech, Dübendorf 8600, Switzerland.

3. Institute for Integrative Biology, ETH Zürich, Zürich 8092, Switzerland.

4. Center for Adaptation to a Changing Environment, ETH Zürich, Zürich 8092, Switzerland.

5. Department of Biological Sciences, University of Texas at El Paso, El Paso, TX 79968, USA

*Author for correspondence: rdkramp@gmail.com

Abstract

Host responses to parasite infection involve several interacting systems. Host genetics determine

much of the response, but it is increasingly clear that the host-associated microbiome also plays a

role. Genetically determined systems and the microbiome can also interact; for example, the

microbiome can modulate the immune response, and vice versa. However, it remains unclear

how such interactions between the host immune system and the microbiome may influence the

host's overall response to parasites. To investigate how host genetics and the microbiome interact

to shape responses to parasites, we imposed truncation selection on Trinidadian guppies

(*Poecilia reticulata*) for low and high resistance to the specialist ectoparasite *Gyrodactylus*

turnbulli. After 3-6 generations of breeding without parasites, we sampled the skin-associated

microbiome and infected fish from each line. We applied Dirichlet Multinomial Modeling

(DMM) machine-learning to identify bacterial community types across lines and evaluated how

selection line and community type explained variations in infection severity. Our findings

showed that among females, the resistant line had significantly lower infection severity, while

the susceptible line had higher infection severity. Among males, only the susceptible line

experienced higher infection severity compared to the other lines. Line did not explain skin

microbial diversity, structure or composition. Our DMM analysis revealed three distinct bacterial

community types, independent of artificial selection lines, which explained just as much

variation in infection load as selection line. Overall, we found that the microbiome and host

genetics independently predict infection severity, highlighting the microbiome's active role in

host-parasite interactions.

1. Introduction

Hosts use multiple strategies to defend themselves against parasites and pathogens (parasites hereafter). Host genetic background is commonly credited with much of the defense against parasites through the immune system, behavior, or other genetic/protein alterations, e.g., thalassemia (Labro, 2022). Recent research has highlighted that the host-associated microbial communities, or 'microbiome', may also influence parasite defense indirectly through the host immune system or directly via niche occupation or secreting anti-parasitic compounds (Bernardo-Cravo et al., 2020), and may also exhibit heritable components. Indeed, the host-associated microbiome can dramatically impact host survival and defense against parasites. However, the question of whether the microbiome plays an active role in host-parasite infection or merely reflects the host's health/immune status is commonly posed (Trevelline et al., 2020). While there is abundant evidence that the microbial community can have a direct or indirect role in host-parasite interactions (Becker et al., 2009; Nelson & May, 2020), there is also evidence that the microbiome is merely a bystander in some diseases and a reflection of the host's genetics, diet, and general health (Ni et al., 2017).

Host genetics and the microbiome interact throughout host development, with complex feedbacks complicating the separation of their roles in traits like parasite defense. On one hand, hosts exert control over their symbionts, shaping their associated microbiomes, via immune responses, barrier function, and physiological regulation (Wilde et al., 2024). On the other hand, the host immune system relies on early microbial colonization for proper development and ultimately shapes immune cell composition (Amenyogbe et al., 2017). The host's genetic makeup therefore influences microbiome composition, and may lead to the selection of specific microbial

communities that can be inherited across generations (Bordenstein & Theis, 2015). However,

distinguishing host genetic effects on the microbiome from factors such as priority effects, drift,

and other stochastic elements, is an ongoing challenge (Debray et al., 2022), and the influence of

external factors such as environment and temporal changes in shaping host-associated

microbiomes must also be carefully considered. Direct manipulation of the host genetics in traits

like parasite resistance, where the microbiome likely plays a role, thus provides a useful tool in

disentangling host genetic effects from microbial ones.

Both the host-associated microbiome and the immune system commonly differ between host

sexes (Markle & Fish, 2014). These differences are likely due to other correlated differences, for

example, high testosterone levels in the blood both suppress the immune system and significantly

alter gut microbial communities (Santos-Marcos et al., 2023), and both the immune system and

microbiome change during pregnancy and menstruation (Chen et al., 2023; Gorczyca et al.,

2022). Given that host sexes across taxa also differ in their response to parasites (Morrow, 2015),

it is essential to test for sex-specific interactions between genetic resistance factors, the

microbiome, and parasites.

Trinidadian guppies (*Poecilia reticulata*) and their specialist ectoparasite *Gyrodactylus turnbulli*

provide a rich context to study host-parasite dynamics and interactions with the microbiome.

Gyrodactylus spp. feed and reproduce (both sexually and asexually, generation time 24hrs) on

host skin. Guppies show a large variation in parasite resistance across populations, sexes, and

ecological contexts (Dargent et al., 2016; Phillips et al., 2021; Stephenson, van Oosterhout, &

Cable, 2015; Van Oosterhout et al., 2003). Guppy resistance is at least partly genetically

determined and heritable (Dargent et al., 2016; Phillips et al., 2018, 2021; Van Oosterhout et al.,

2003; Weiler et al., 2025). Innate and acquired immunity are important for *G. turnbulli*

resistance, however, the precise mechanisms involved in guppy parasite resistance remain

unclear (Cable & Van Oosterhout, 2007; Mohammed et al., 2016). Research into genetically

based resistance to *G. turnbulli* has largely focused on the major histocompatibility complex

(MHC) (Fraser et al., 2010; Fraser & Neff, 2009; Phillips et al., 2018). Notably, skin bacterial

alpha diversity correlates with MHC heterozygosity in frogs (Cortazar-Chinarro et al., 2024), and

probiotic supplementation results in significant upregulation of MHC genes in fish (Tanpichai et

al., 2025). Similar associations and covariation between host genetics and the skin microbiome

of fish would sets up a clear avenue for the skin microbiome and host genetics to interactively

shape parasite resistance phenotypes in our system.

Here, we tested how host genetics and host-associated microbiomes affect the interaction

between guppies and *G. turnbulli*. We imposed truncation selection for host resistance and

susceptibility to the parasite, quantified as the mean number of parasites over infection. While

we did not specifically select for tissue-specific tolerance (tolerance hereafter), or the ability to

minimize the fitness losses caused by a given parasite load (Stephenson & Adelman, 2022), we

did examine the levels of tolerance within our lines. Previous literature has demonstrated that

tolerance is an essential factor to consider in guppy response to *Gyrodactylus* parasites (Jog et

al., 2022; Stephenson, van Oosterhout, & Cable, 2015; Tadiri et al., 2021). However, whether

tolerance of infection is heritable has rarely been explored.

We found that host genetics and the microbiome significantly and independently predicted the

severity of parasite infection. Parasite resistance and tolerance showed sex-specific responses to

selection. These results suggest that the microbiome independently influences host-parasite

interactions rather than merely reflecting the host's genetics, highlighting the complex interplay

among host genetics, the microbiome, and sex-specific responses to infections.

2. Materials and methods

2. Materials and methods

2.1 Data and sample collection

2.1.1 Truncation selection to establish guppy lines that differ in genetically-based resistance to

Gyrodactylus turnbulli

We used laboratory-bred descendants of wild guppies from a high-predation population on the

Caura River, Trinidad (UTM: 20 P 679527.7 m E, 1180376.4 m N; elevation 112 m). Wild

guppies (n ~600) were transported to Cardiff University in June 2012 (Cefas APB authorization

number CW054-D-187A) and treated prophylactically with Binox® (Nitrofurazone). The site

was previously confirmed *Gyrodactylus* spp.-free by previous surveys, and no *Gyrodactylus* spp.

were found in a subset of 80 fish examined. We collected fin clips from 66 parental-generation

fish for microsatellite genotyping. Fish were housed in 70 L mixed-sex tanks under a 12h

light:12h dark cycle at 24±1°C and fed daily with Aquarian® tropical fish flakes, supplemented

with Artemia and bloodworm. Fry were transferred to juvenile tanks soon after birth, checked for

ectoparasitic infection, and reared until sex determination (6-8 weeks). They were then kept in

sex-specific tanks for at least another month before use, ensuring all fish were sexually mature

and unmated. *Gyrodactylus turnbulli* were from an established isogenic culture in the lab (*Gt3*),

maintained on commercially sourced guppies ('culture fish').

We experimentally infected 100 male and 100 female F1 guppies to evaluate their resistance and

select founders for our 'resistant' and 'susceptible' lines. To initiate infections, a heavily infected

culture fish was euthanized with MS222 (0.02% tricaine methanesulfonate) and placed next to an

anesthetized experimental fish until two *G. turnbulli* parasites transmitted, as observed under a

dissecting scope. Experimental fish were housed individually in 1 L tanks, and parasites were

counted every other day on anesthetized fish. After nine days, all fish were treated with

Levamisole and screened under anesthetic several times to confirm parasite clearance. Fin clips

were collected for genotyping. The 30 males and 30 females with the highest (lowest) mean

number of worms founded the 'susceptible' ('resistant') line, and randomly selected uninfected

F1 fish (30 of each sex) founded the control line (*Figure S1*). Each line was allowed to breed

freely in separate 120 L tanks. Tanks were monitored for fry, which were moved to F2 rearing

tanks. The F2 generation was split into two subpopulations (named 'A' and 'B' within each of the

susceptible, resistant, and control lines), maintaining 60 fish per tank. This population size was

kept consistent across the six subpopulations, with new tanks set up for each subsequent

generation as above. During preliminary analysis, we confirmed that these subpopulations did

not differ in either infection severity or parasite tolerance (see Supporting Information) and

therefore combined them for the subsequent analyses described below.

After roughly 3 generations post-selection, the fish moved with one of the authors to a new

institution, at which the husbandry conditions were different. Here, guppies were housed in

mixed-sex groups at densities of 1-2 fish per L in 4.5 L tanks in a recirculating system at 24±1°C,

on a 12 h light: 12 h dark lighting schedule (overhead fluorescent lighting) and fed daily on

commercial flake food supplemented with *Artemia*. Each subpopulation was therefore housed

across at least 10 separate tanks, again with multiple new tanks set up for each subsequent

generation. The fish were held and bred under these conditions from August 2015 until their

experimental infections in August-November 2017.

2.1.2 A note on replication in our study design

As described above, we imposed one round of truncation selection for resistance and

susceptibility, on one group of F1 fish. Typically, selection experiments using guppies to

characterize the genes underlying phenotypic variation (e.g., (van Wijk et al., 2013)), or

evolutionary change in correlated traits (e.g., (Bartuseviciute et al., 2022; Kotrschal et al., 2013))

replicate this process three times because the scale of inference is at the level of the selection

event: to draw conclusions about how underlying genes or correlated traits respond to selection,

the selection event should be replicated (Kawecki et al., 2012).

By contrast, we do not identify genes, or focus on correlated evolutionary trait changes: instead,

we use truncation selection to maximize the variation in response to infection in our focal

population, and to compare the role of host genetics and associated microbiomes in infection

phenotypes. That resistance to *Gyrodactylus* spp. is heritable is not our focus, and has already

been established several times (Dargent et al., 2013; Madhavi & Anderson, 1985; Phillips et al.,

2018, 2021; Weiler et al., 2025). To ensure that the difference between our lines was due to

selection, and thus underlying genetic differences, we kept consistent population sizes across

generations, split each line into two subpopulations, housed fish across several replicate tanks

per line, bred the fish for at least 3 generations in the absence of parasites, and used genotyping

to eliminate the role of inbreeding differences between them. With our design, therefore, we can

rule out potential confounding effects (from tanks, stochastic differences post selection,

inbreeding differences, and maternal effects).

Our selection process thus produced individual guppies that differed in genetically-based

resistance to our focal parasite; here we show that the microbiomes associated with these

individuals contributed independently to their infection phenotype. Our main conclusion, that the

host-associated microbiome predicts infection phenotype independently of and just as strongly as

host genetics, is therefore robustly supported by our study design.

2.1.3 Experimental infection and microbiome sampling protocol

Before infection began, we collected samples from 51 fish split between the infection and sham-

infection treatment groups: control (infected n=10, sham n=5), Resistant (infected n=12, sham

n=6), and Susceptible (infected n=9, sham n=8). We used the methods previously published to

collect and inventory the microbial communities on the fish skin.

We infected 88 guppies 3-7 generations post truncation selection with *G. turnbulli* in five

batches, following the same procedure as for the F1 generation, described above. 14 guppies

from each line were 'sham' infected and were used in microbiome analysis. *G. turnbulli for this*

infection were collected from a local pet shop in June 2017. An isogenic strain was initiated by

infecting a parasite-naïve ornamental guppy with a single parasite individual. This strain was

subsequently maintained on parasite-naïve culture fish, and identified to species level using

molecular methods.

***2.2 What is the magnitude of host genetic contribution to guppy resistance to* Gyrodactylus turnbulli?**

2.2.1 Statistical analysis

We used R v.4.3.2 (R Core Team, 2020) for these analyses and provide our code, output, and validation in the Supporting Information (SI). We used ggplot2 (Wickham, 2016) to plot all the figures. For all models, we assessed collinearity among predictor variables using the pairs function (Zuur et al., 2009), validated model fits using the DHARMa package (Hartig, 2022), and used Type II Wald chi-square tests to test for significance using the *Anova()* function from the 'car' package (Fox & Weisberg, 2019).

We quantified infection resistance/susceptibility as the area under the curve of parasite number over time, or "infection severity." To test response to selection, we used infection severity as the response variable in a generalized linear mixed model (GLMM) in the glmmTMB package, v. 1.1.2.3 (Brooks et al., 2017) (Gamma error family and logit link function). As fixed effects, we included selection line (resistant n=30, control n=28, susceptible n=29), fish sex, length (pre-infection, using residuals from a regression on sex to control for sex differences), body condition (scaled mass index; (Peig & Green, 2009)), generation, number of parasites establishing the infection (dose), and trial date (Julian date). Length and body condition were calculated with entire population of fish including sham infected fish (resistant n=14, control n=14, susceptible n=14), to obtain a more accurate distribution. We also included the interaction between sex and line.

We quantified tolerance as the host's relative per-parasite percentage mass change during

infection, calculated by dividing the percentage mass change from day 0 to day 20 by the

maximum number of worms on the fish, relative to the highest observed percentage change. To

assess the impact of selection on tolerance, we used tolerance as a response variable in a GLMM,

including the fixed effects of line, sex, length, dose, and Julian date. We excluded scaled mass

index due to its overlap with mass change. Furthermore, we did not include infection severity

due to high collinearity with other fixed effects within the model (line & dose). Additionally,

infection severity within the original model caused model fit issues due to the non-linear

relationship between infection severity and ppmass.

In a different model, we tested for a tradeoff between resistance and tolerance with a quadratic

polynomial regression model to account for non-linear effects. We used tolerance as the response

variable and log-transformed infection severity (our resistance metric), sex, their interaction,

length, and generation as fixed effects.

2.2.2 Calculating heritability

We calculated narrow-sense heritability (assuming that the response to selection is primarily due

to additive genetic variance) of resistance (using mean worm – the resistance metric we used

during truncation selection) and tolerance following the breeder's equation (Lush, 1937). For

mean worm, we used the average number of parasites between 1-9 days of infection, centered to

a mean of zero to account for the differences in worm growth between the infections of the F1

and subsequent generations. Tolerance was centered identically.

P_b: Average starting phenotype (mean phenotype of the population before selection)

P_s: Average selected phenotype (mean phenotype of the selected parents)

P_r: Average response phenotype (mean phenotype of the offspring generation)

h^2: Narrow-sense heritability

$h^2 = P_r - P_b P_s - P_b$

2.2.3 Do differences in inbreeding contribute to line differences?

We evaluated the genetic diversity and variance in inbreeding within the wild-caught

populations, F1, and F3-F6 generations, to determine if the observed difference between the

resistant and susceptible lines could be attributed to differences in inbreeding. We extracted DNA

from fin clips with the HotSHOT method: using 50 µl of alkaline lysis reagent and neutralizing

solution followed by incubation at 95 °C for 30 minutes (Truett et al., 2000). We used two

multiplex assays to amplify eight previously designed microsatellite markers (Becher et al.,

2002; Nater et al., 2008; Paterson et al., 2005; van Oosterhout et al., 2006; Watanabe et al., 2005)

at the Genetic Diversity Centre (GDC), ETH Zurich. Samples were genotyped on an ABI

Genetic Analyzer 3730 (Applied Biosystems, USA), scored using GeneMarker (Version 2.6.4;

Softgenetics), and binned using the package MSatAllele (Alberto, 2013) in R (R Core Team,

2020). See the supporting information for more details.

We tested allelic richness (A_R), gene diversity (H_S), $F_{IS,}$ and identity disequilibrium (g2) among

our artificial selection lines to evaluate whether the differences in phenotype were attributable to

variation in inbreeding (Smallbone et al., 2016). Allelic richness is the first population genetic

metric to reflect bottleneck effects, which decrease the effective population size and facilitate

reductions in heterozygosity due to inbreeding (Nei et al., 1975). Significant differences in allelic

richness among populations are indicative of variation in increases in the probability that

inbreeding may occur. Gene diversity measures the mean heterozygosity per individual per

population, under inbreeding conditions homozygosity increases but gene diversity is not

immediately lost (Charlesworth, 2003). Positive values in F_{IS} can be generated by inbreeding but

may also be artificially elevated by technical artifacts such as large allele drop out or null alleles,

which can artificially inflate a deficiency of heterozygotes (De Meeûs, 2018; Nei et al., 1975).

We therefore also estimated identity disequilibrium (g_2), the correlation of heterozygosity across

loci resulting from inbreeding, which is robust in the presence of null alleles (David et al.,

2007). Concurrence among population genetic metrics consistent with patterns inbreeding are

needed to determine if phenotypic differences among populations are attributable to inbreeding.

Allelic richness (A_R) (rarefied to the lowest population sample size), gene diversity (H_S) and F_{IS},

was calculated in FSTAT 2.39 (Goudet, 01 2001). Significance levels for F_{IS} were assessed in

GENEPOP (web version 4.2;(Raymond & Rousset, 1995) with 5000 iterations followed by a

Bonferroni correction for multiple comparisons. Tests for statistically significant differences

allelic richness (A_R), gene diversity (H_S) and F_{IS} among groups were calculated in FSTAT with

2000 permutations. Identity disequilibrium (g^2) is the difference between joint identity by

descent and the product of the separate probabilities of identity by decent for two loci. g^2 is

estimated with microsatellite markers as the excess of double heterozygous genotypes at all pairs

of loci relative to a random distribution of heterozygous genotypes among individuals at each

locus (Cockerham & Weir, 1973; David et al., 2007). g_2 values significantly greater than 0 are

indicative of identity disequilibrium (ID). ID was calculated in inbreedR (Stoffel et al., 2016).

2.3 Does the host-associated microbiome contribute to guppy resistance to Gyrodactylus

infection?

2.3.1 Microbial alpha and beta diversity

Raw sequence data were processed using QIIME2 version 2019.7 (Bolyen et al., 2019). We

rarefied the resulting ASV table to 3,654 sequences per sample which resulted in excluding one

sample and kit controls, and used the DADA2 pipeline to calculate the unweighted and weighted

UniFrac distances (Lozupone & Knight, 2005) pairwise between all samples to create distance

matrices. We used a PERMANOVA from the adonis package in R (Anderson, 2001) to test for

significant differences in microbial community across fish line, sex, length, and body condition.

We used QIIME2 to calculate the number observed number of ASVs, Shannon's diversity index,

and Faith's phylogenetic diversity, and compare the mean values of these metrics across line and

sex using Pairwise Kruskal-Wallis tests.

2.3.2 Bacterial community typing

We used Dirichlet multinomial mixture models (DMM) (Holmes et al., 2012) in the Dirichlet

Multinomial v1.16.0 package (Morgan, 2023) to assign all samples (n=50) to three different

bacterial community types based on the relative frequency of taxa. We compared the Laplace

approximation of the negative log models to identify the number of Dirichlet components that

resulted in the best model fit (see SI).

2.3.3 Does bacterial community type predict the severity of Gyrodactylus *infection?*

To test if the bacterial community type significantly predicted infection severity, we used

bacterial community type as a fixed effect in the GLMM model used for the selection line

resistance analysis described in section 2.1.2, with the data subset to just swabbed fish.

2.4 What is the relative importance of host genetics and the microbiome to guppy resistance to

Gyrodactylus infection?

To compare how well our GLMM models with a) bacterial community type only, b) artificial

selection line only, and c) both factors explained variation in infection severity, we used the

anova() function. All three models shared the same fixed effects, differing only in whether line or

bacterial community type or both were included as a fixed effect. Models involving skin

microbiome and infection were run on the reduced dataset using only the infected and swabbed

fish (n=31).

3. Results

3.1 Host genetics contribute to guppy resistance to, and tolerance of, Gyrodactylus turnbulli

infection

3.1.1 Resistance responded to selection

Selection line was a significant predictor of infection severity (Figure 1; $\chi^2(2) =37.74$, p=6.4e-

09) Additionally, an interaction between selection line and sex was observed ($\chi^2(2) =13.51$,

p=0.0012). Females had significantly higher infections than males ($\chi^2(1) =24.29$, p=8.3e-07) and

high starting worm 'dose' resulted in higher infection severity ($\chi^2(1) =5.11$, p=0.024). Post hoc

tests to evaluate whether selection line significantly explained infection severity among each sex

found that line was still significant in both sexes (female; $\chi^2(2) =24.33$, p=5.2e-06 & male; $\chi^2(2)$

=17.57, p=0.00015). Post hoc Tukey tests also revealed that the female guppies from the

susceptible line experienced significantly higher (Cohen's d=3.49, p<0.001), and from the

resistant line experienced significantly lower infection severity (Cohen's d=2.74, p=0.016) than

the control. Male guppies from the susceptible line experienced significantly higher infection

severity than control line males (Cohen's d=4.19, p<0.001), but resistant line males did not differ

significantly from control line males (Cohen's d=1.39, p=0.15). The heritability of the average

number of parasites throughout infection was h^2=0.79 in the susceptible line, and h^2=0.41 in the

resistant line.

3.1.2 Tolerance responded to selection and trades off with resistance

The selection lines significantly differed in relative tolerance (Figure 2; χ^2(2)=14.56, p=0.00069)

and significantly interacted with sex (χ^2(2)=10.28, p=0.00586). Additionally, larger fish were

more tolerant (χ^2(1)=5.94, p=0.014). A post hoc test found that selection line was only significant

in female guppies (female; χ^2(2)=23.81, p=6.7e-06), and smaller females were less tolerant (χ^2(2)

=6.96, p=0.0083). Post hoc Tukey tests revealed that resistant line females were significantly less

tolerant than control and susceptible line females (control; Cohen's d=7.04, p=0.0003;

susceptible; Cohen's d=8.54, p<0.00016). However, no fixed effect explained variation in

tolerance among males and we did not find that the bacterial community types significantly

predicted tolerance. The heritability of tolerance was h^2=0.52 in the susceptible line and h^2=0.24

in the resistant line.

The polynomial regression model revealed a tradeoff between tolerance and resistance at the

individual level: higher tolerance was correlated with lower resistance (Figure S1; F= 55.49,

p<7e-15). No other fixed effects included explained significant variation in tolerance.

3.1.3 Selection lines did not differ in the extent of inbreeding

We saw a significant reduction in allelic richness among all experimental infection groups

relative to the wild-caught founders, likely reflecting a post-capture reduction in population size

(Table S1). However, we found no significant difference in AR (p= 0.589), Hs (p=0.73), FIS

(p=0.42), or g2 (Table S2) among the three lines.

3.2 The host-associated microbiome is associated with guppy resistance to Gyrodactylus

turnbulli infection, independently and to a similar degree as host genetics

DMM modeling found that the lowest Laplace approximation model fit was three bacterial

clusters (*Figure S2*). All samples had a posterior probability of 98% or higher. Each bacterial

community type contained 10.0–66.8% of all samples (n=5-33), and no samples were undefined.

The three bacterial community types (A, B, and C) in our fish skin samples were distinct

(PERMANOVA: F = 7.37, p = 0.0003, Figure 3a & Figures S3-4). Males and females were

balanced among the bacterial community types A and B, and bacterial community types clustered

evenly across the artificial selection lines (Figure 3a; Fisher's exact test, p=0.8). Additionally, we

found no interaction between line and bacterial community types in our model ($\chi 2(4)=3.07$,

p=0.546). All skin swab samples were assigned to a community type with the greatest posterior

probability. For top contributing bacterial genera, see *Figure S3*.

The selection lines did not significantly differ in any skin microbial diversity metrics tested. The microbiomes associated with females had significantly higher alpha diversity than those associated with males (Kruskal–Wallis pairwise test, Shannon's index: H=4.06, q=0.04; observed ASVs: H=4.88, q=0.03). Consistent with Kramp et al. (2022), the microbiomes associated with the two sexes also had significantly different beta diversity (unweighted unifrac PERMANOVA; F=1.56, p=0.006).

We found that the bacterial community type A of the skin microbiome on Trinidadian guppies had significantly reduced infection severity compared to bacterial community type B & C ($\chi^2(2)$ = 19.83, p=4.9e-5, Figure 3b). A post-hoc test showed that a model including both selection line and bacterial community type is significantly better at explaining variation in infection severity than models with either variable alone (Figure 4; $\chi^2(14)$=13.40, p=2e-16).

4. Discussion

Our selection lines demonstrated striking heritability of parasite infection resistance and tolerance. Despite no significant differences in microbiome between the three lines, we found that bacterial community type, found by DMM machine-learning, predicted infection severity. Host genetics and the host-associated microbiome thus appear to independently predict infection severity, with similar effect sizes. We also observed notable differences between males and females in skin microbiome, and in response to selection. We discuss each of these results in turn.

We found that both guppy resistance and susceptibility to *G. turnbulli* responded readily to a

single generation of truncation selection. We found no notable differences between the three

selection lines in our genetic metrics, indicating that the differences we found were not due to

differences in inbreeding among them. Our narrow-sense heritability estimates for resistance and

tolerance were high, particularly for our susceptible lines. However, our values are comparable to

heritability estimates in previous studies (Weiler et al., 2025) and to the heritability of male

guppy ornaments (Morris et al., 2020), which are correlated with *G. turnbulli* resistance

(Stephenson et al., 2020).

Parasite resistance and susceptibility likely responded to selection through different mechanisms.

The response to selection for susceptibility could have involved either an increase in tolerance, a

loss of resistance, or some combination. Indeed, our experimental design may have favored

tolerance in the susceptible line, as only individuals surviving severe infections would have

contributed to future generations. Tolerance and resistance likely have a different genetic bases,

and hosts appear to invest in one strategy at the expense of the other (Råberg et al., 2007). This

observation is supported by the fact that our resistant line fish had lower tolerance than control

and susceptible line fish – at least among females – and that the two defense strategies were

negatively correlated at the level of individual fish in our experiment. However, these tradeoffs

do not always occur, and both tolerance and resistance traits can coexist within a single

individual host and a population (Hardy et al., 2024; Pagán & García-Arenal, 2018).

Our observed high heritability of susceptibility (and potentially tolerance) has implications for

our understanding of how guppy defenses might evolve in natural conditions (Walsman et al.,

2022, 2024). High heritability suggests natural selection can act rapidly on host defense traits in

populations under high parasite pressure, and indeed host defense appears to have the capacity to

evolve rapidly in this system (Dargent et al., 2016; Phillips et al., 2018). In some populations,

infection prevalence is very high (>90%; (Clark et al., 2023; Stephenson, van Oosterhout,

Mohammed, et al., 2015)), potentially selecting for increased infection tolerance and decreased

resistance, a phenomenon known as the 'resistance is futile' effect (Walsman et al., 2023).

However, we (and all previous studies of host defense traits in this system) deliberately excluded

any potential role of parental effects of infection, when recent findings suggest that

transgenerational immune priming may be important in this system, which would reduce the

strength of selection on these traits, particularly when prevalence is high (Weiler et al., 2025).

We found that bacterial community type predicted infection severity (but interestingly not

tolerance) independently, and to a similar degree, as host genetics. Despite the fact that host

characteristics such as the immune system, metabolism, and growth rate have been shown to

interact with the microbiome (Speer, 2022; Zheng et al., 2020), we found no difference between

our lines in skin microbiome composition, alpha diversity, or bacterial community types.

Potentially this is because fish skin microbiomes are highly variable and may not reflect the

host's genetics but instead drift stochastically (Jones et al., 2022; Wang et al., 2023). However,

recently it has come to light that different organ may provide unique insights of host health, i.e.,

eye microbiome predicted disease outcome but gut microbiome did not in Island foxes

(DeCandia et al., 2025). Therefore, it is possible that microbiome from other tissue could have

been modified due to our selection, though, we did not test for this. By contrast, artificial

selection for body size in gilthead sea bream (*Sparus aurata*) significantly altered the host-

associated microbiome, and the microbiome in the fast-growing line had greater parasite

colonization resistance (Piazzon et al., 2020), and size selection experiments indirectly affect

guppy resistance to *G. turnbulli* (Bartuseviciute et al., 2022) potentially mediated through the

microbiome. These results suggest that selection on host growth does influence the associated

microbiome composition, potentially through effects on host metabolism, with downstream

effects on parasite resistance (Bartuseviciute et al., 2022; Piazzon et al., 2020), whereas selection

on parasite resistance appears not to have direct effects on skin-associated microbial

communities (this study).

Our findings thus contribute to the growing literature linking host-associated microbiomes and

infection severity, pointing to important interactions between the skin microbiome and parasites

(Scheifler et al., 2022; Speer, 2022; Trevelline et al., 2020). More broadly, the microbiome's role

in determining resistance has been seen in several taxa, where pre-infection microbiome

composition, alpha diversity, or microbiome function is a powerful tool to predict individuals

that are most susceptible to parasites within a population (Bates et al., 2022; Lee et al., 2019).

Intriguingly, we found five of the eight ASVs reported to be indicator taxa of epizootic dynamics

in a recent study of skin microbial metabolism function in amphibian chytridiomycosis (Bates et

al., 2022): *Burkholderiales*, *Pseudomonadales*, *Sphingomonadales*, *Rhizobiales*,

Flavobacteriales (see Figure S2 for all contributing species).

Sex differences are pervasive throughout our results. Sexual dimorphism in immunocompetence

has been reported in many taxa, including Trinidadian guppies. Our results reveal additional

evidence of sex-specificity in defenses against the specialist parasite *G. turnbulli*: resistance

responded more strongly to selection among female than male guppies, as has been found

previously (Dargent et al., 2016). Potentially, this result reflects the fact that males in nature and

in our experiment were subject to sexual selection in addition to direct selection for parasite

resistance. Females in many populations prefer males with larger orange ornaments (Endler et

al., 2001; Houde, 1988, 1997), with implications for disease dynamics (Rovenolt et al., 2024).

Larger orange ornaments indicate male resistance to this parasite (Stephenson et al., 2020).

Males tend to develop lower infection loads in general in the lab (Johnson et al., 2011), are less

likely, and less heavily infected in surveys of natural populations (Clark et al., 2023; Gotanda et

al., 2013; Stephenson, van Oosterhout, Mohammed, et al., 2015). Thus, if female preference has

already eroded genetic variation in parasite resistance among males, and females across our lines

selected the most resistant males as mates, this may explain the overall high and invariable

resistance we observe in males. By contrast, and again in line with previous results from field

surveys (Stephenson, van Oosterhout, & Cable, 2015), females – particularly large ones – appear

to be more tolerant of infection than males. Notably, we found evidence for a tradeoff between

resistance and tolerance at the line level among females, but not among males, perhaps indicating

different trade-offs for each sex, though there was no sex difference in the tradeoff at the level of

individual fish.

The strong sex-specificity in host defense traits we observe here highlights the need to

investigate aspects of guppy response to *Gyrodactylus turnbulli* beyond the MHC. Specifically, if

the primary mechanism of resistance were driven by MHC parasite resistance, we would not see

such pronounced sex differences in response to selection: MHC resistance is a mixture of both

mother and father MHC genes. While our data indicate that the microbiome is not driving this

sex difference in response to selection, the microbiome clearly contributes to parasite resistance

in this system and appears to differ between the sexes: they differed in both alpha and beta

diversity metrics of bacterial communities, similar to the skin microbiome in other species

(Rodríguez-Barreto et al., 2023). The observed sex-specific differences highlight potential co-

evolutionary interactions between host sex and microbiome in relation to parasite resistance.

These possibilities will hopefully inspire exciting and novel exploration of the evolutionary

dynamics of resistance and tolerance between the sexes.

References

Alberto, F. (2013). MsatAllele: Visualizes the scoring and binning of microsatellite fragment

sizes. *R Package Version*.

Amenyogbe, N., Kollmann, T. R., & Ben-Othman, R. (2017). Early-Life Host-Microbiome

Interphase: The Key Frontier for Immune Development. *Frontiers in Pediatrics*, *5*, 111.

Anderson, M. J. (2001). A new method for non-parametric multivariate analysis of variance.

Austral Ecology, *26*(1), 32–46.

Bartuseviciute, V., Diaz Pauli, B., Salvanes, A. G. V., & Heino, M. (2022). Size-selective

harvesting affects the immunocompetence of guppies exposed to the parasite

Gyrodactylus. *Proceedings. Biological Sciences / The Royal Society*, *289*(1981),

20220534.

Bates, K. A., Sommer, U., Hopkins, K. P., Shelton, J. M. G., Wierzbicki, C., Sergeant, C., Tapley,

B., Michaels, C. J., Schmeller, D. S., Loyau, A., Bosch, J., Viant, M. R., Harrison, X. A.,

Garner, T. W. J., & Fisher, M. C. (2022). Microbiome function predicts amphibian

chytridiomycosis disease dynamics. *Microbiome*, *10*(1), 44.

Becher, S. A., Russell, S. T., & Magurran, A. E. (2002). Isolation and characterization of

polymorphic microsatellites in the Trinidadian guppy (Poecilia reticulata). *Molecular*

Ecology Notes, *2*(4), 456–458.

Becker, M. H., Brucker, R. M., Schwantes, C. R., Harris, R. N., & Minbiole, K. P. C. (2009). The

bacterially produced metabolite violacein is associated with survival of amphibians

infected with a lethal fungus. *Applied and Environmental Microbiology*, *75*(21), 6635–

6638.

Bernardo-Cravo, A. P., Schmeller, D. S., Chatzinotas, A., Vredenburg, V. T., & Loyau, A. (2020).

Environmental Factors and Host Microbiomes Shape Host-Pathogen Dynamics. *Trends in*

Parasitology, *36*(7), 616–633.

Bolyen, E., Rideout, J. R., Dillon, M. R., Bokulich, N. A., Abnet, C. C., Al-Ghalith, G. A.,

Alexander, H., Alm, E. J., Arumugam, M., Asnicar, F., Bai, Y., Bisanz, J. E., Bittinger, K.,

Brejnrod, A., Brislawn, C. J., Brown, C. T., Callahan, B. J., Caraballo-Rodríguez, A. M.,

Chase, J., … Caporaso, J. G. (2019). Reproducible, interactive, scalable and extensible

microbiome data science using QIIME 2. *Nature Biotechnology*, *37*(8), 852–857.

Bordenstein, S. R., & Theis, K. R. (2015). Host Biology in Light of the Microbiome: Ten

Principles of Holobionts and Hologenomes. *PLoS Biology*, *13*(8), e1002226.

Brooks, M. E., Kristensen, K., van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A.,

Skaug, H. J., Maechler, M., & Bolker, B. M. (2017). glmmTMB Balances Speed and

Flexibility Among Packages for Zero-inflated Generalized Linear Mixed Modeling. In

The R Journal (Vol. 9, Issue 2, pp. 378–400). https://doi.org/10.32614/RJ-2017-066

Cable, J., & Van Oosterhout, C. (2007). The role of innate and acquired resistance in two natural

populations of guppies (Poecilia reticulata) infected with the ectoparasite Gyrodactylus

turnbulli. *Biological Journal of the Linnean Society. Linnean Society of London*, *90*(4),

647–655.

Charlesworth, D. (2003). Effects of inbreeding on the genetic diversity of populations.

Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences,

358(1434), 1051–1070.

Chen, X., Wu, R., Li, L., Zeng, Y., Chen, J., Wei, M., Feng, Y., Chen, G., Wang, Y., Lin, L., Luo,

H., Chen, A., Zeng, Z., He, F., Bai, Y., Zhang, S., Han, Y., Wang, Z., Zhao, X., … Gong,

S. (2023). Pregnancy-induced changes to the gut microbiota drive macrophage pyroptosis

and exacerbate septic inflammation. *Immunity*, *56*(2), 336-352.e9.

Clark, D. R., Young, K. A., Kitzes, J., Moore, P. J., Evans, A. J., & Stephenson, J. F. (2023).

Aggregation of symbionts on hosts depends on interaction type and host traits.

Ecography. https://doi.org/10.1111/ecog.06858

Cockerham, C. C., & Weir, B. S. (1973). Descent measures for two loci with some applications.

Theoretical Population Biology, *4*(3), 300–330.

Cortazar-Chinarro, M., Richter-Boix, A., Rödin-Mörch, P., Halvarsson, P., Logue, J. B., Laurila,

A., & Höglund, J. (2024). Association between the skin microbiome and MHC class II

diversity in an amphibian. *Molecular Ecology*, *33*(1), e17198.

Dargent, F., Rolshausen, G., Hendry, A. P., Scott, M. E., & Fussmann, G. F. (2016). Parting

ways: parasite release in nature leads to sex-specific evolution of defence. *Journal of*

Evolutionary Biology, *29*(1), 23–34.

Dargent, F., Scott, M. E., Hendry, A. P., & Fussmann, G. F. (2013). Experimental elimination of

parasites in nature leads to the evolution of increased resistance in hosts. *Proceedings.*

Biological Sciences / The Royal Society, *280*(1773), 20132371.

David, P., Pujol, B., Viard, F., Castella, V., & Goudet, J. (2007). Reliable selfing rate estimates

from imperfect population genetic data. *Molecular Ecology*, *16*(12), 2474–2487.

De Meeûs, T. (2018). Revisiting FIS, FST, Wahlund Effects, and Null Alleles. *The Journal of*

Heredity, *109*(4), 446–456.

Debray, R., Herbert, R. A., Jaffe, A. L., Crits-Christoph, A., Power, M. E., & Koskella, B. (2022).

Priority effects in microbiome assembly. *Nature Reviews. Microbiology*, *20*(2), 109–121.

DeCandia, A. L., Lu, J., Hamblen, E. E., Brenner, L. J., King, J. L., Gagorik, C. N., Schamel, J.

T., Baker, S. S., Ferrara, F. J., Booker, M., Bridges, A., Carrasco, C., vonHoldt, B. M.,

Koepfli, K.-P., & Maldonado, J. E. (2025). Phylosymbiosis and elevated cancer risk in

genetically depauperate channel island foxes. *Molecular Ecology*, *34*(2), e17610.

Endler, J. A., Basolo, A., Glowacki, S., & Zerr, J. (2001). Variation in response to artificial

selection for light sensitivity in guppies (Poecilia reticulata). *The American Naturalist*,

158(1), 36–48.

Fox, J., & Weisberg, S. (2019). Companion to Applied Regression [R package car version 3.1-2].

Sage, *3rd*. https://cran.r-project.org/web/packages/car/index.html

Fraser, B. A., & Neff, B. D. (2009). MHC class IIB additive and non-additive effects on fitness

measures in the guppy *Poecilia reticulata*. *Journal of Fish Biology*, *75*(9), 2299–2312.

Fraser, B. A., Ramnarine, I. W., & Neff, B. D. (2010). TEMPORAL VARIATION AT THE MHC

CLASS IIB IN WILD POPULATIONS OF THE GUPPY (*POECILIA RETICULATA*).

Evolution; International Journal of Organic Evolution, *64*(7), 2086–2096.

Gorczyca, K., Obuchowska, A., Kimber-Trojnar, Ż., Wierzchowska-Opoka, M., & Leszczyńska-
Gorzelak, B. (2022). Changes in the Gut Microbiome and Pathologies in Pregnancy.
International Journal of Environmental Research and Public Health, *19*(16).
https://doi.org/10.3390/ijerph19169961

Gotanda, K. M., Delaire, L. C., Raeymaekers, J. A. M., Pérez-Jvostov, F., Dargent, F., Bentzen,
P., Scott, M. E., Fussmann, G. F., & Hendry, A. P. (2013). Adding parasites to the guppy-
predation story: insights from field surveys. *Oecologia*, *172*(1), 155–166.

Goudet, J. (01 2001). FSTAT, Version. *1. 2: A Computer Program to Calculate F-Statistics.*
Heredity, *86*, 485–486.

Hardy, B. M., Muths, E., Funk, W. C., & Bailey, L. L. (2024). Quantifying intraspecific variation
in host resistance and tolerance to a lethal pathogen. *The Journal of Animal Ecology*.
https://doi.org/10.1111/1365-2656.14106

Hartig, F. (2022). *DHARMa: residual diagnostics for hierarchical (multi-level/mixed) regression*
models. R package version 0.4. 6.

Holmes, I., Harris, K., & Quince, C. (2012). Dirichlet multinomial mixtures: generative models
for microbial metagenomics. *PloS One*, *7*(2), e30126.

Houde, A. E. (1988). Genetic difference in female choice between two guppy populations.
Animal Behaviour, *36*(2), 510–516.

Houde, A. E. (1997). Effect of artificial selection on male colour patterns on mating preference
of female guppies. *Proceedings of the Royal Society of London. Series B: Biological*
Sciences, *256*(1346), 125–130.

Jog, M. G., Sackett, M. E., Kisty, S. D., Hansen, J. A., & Stephenson, J. F. (2022). The behaviour

of infected guppies depends on social context, parasite tolerance and host sex. *Animal*

Behaviour, *187*, 97–104.

Johnson, M. B., Lafferty, K. D., van Oosterhout, C., & Cable, J. (2011). Parasite transmission in

social interacting hosts: monogenean epidemics in guppies. *PloS One*, *6*(8), e22634.

Jones, E. W., Carlson, J. M., Sivak, D. A., & Ludington, W. B. (2022). Stochastic microbiome

assembly depends on context. *Proceedings of the National Academy of Sciences of the*

United States of America, *119*(7), e2115877119.

Kawecki, T. J., Lenski, R. E., Ebert, D., Hollis, B., Olivieri, I., & Whitlock, M. C. (2012).

Experimental evolution. *Trends in Ecology & Evolution*, *27*(10), 547–560.

Kotrschal, A., Rogell, B., Bundsen, A., Svensson, B., Zajitschek, S., Brännström, I., Immler, S.,

Maklakov, A. A., & Kolm, N. (2013). Artificial selection on relative brain size in the

guppy reveals costs and benefits of evolving a larger brain. *Current Biology: CB*, *23*(2),

168–171.

Labro, M. T. (2022). *Host defense and infection*. CRC Press.

Lee, K. H., Gordon, A., Shedden, K., Kuan, G., Ng, S., Balmaseda, A., & Foxman, B. (2019).

The respiratory microbiome and susceptibility to influenza virus infection. *PloS One*,

14(1), e0207898.

Lozupone, C., & Knight, R. (2005). UniFrac: a new phylogenetic method for comparing

microbial communities. *Applied and Environmental Microbiology*, *71*(12), 8228–8235.

Lush, J. L. (1937). *Animal Breeding Plans - Scholar's Choice Edition*. Bibliolife DBA of Bibilio

Bazaar II LLC.

Madhavi, R., & Anderson, R. M. (1985). Variability in the susceptibility of the fish host, *Poecilia*

reticulata, to infection with *Gyrodactylus bullatarudis* (Monogenea). *Parasitology*, *91*(3),

531–544.

Markle, J. G., & Fish, E. N. (2014). SeXX matters in immunity. *Trends in Immunology*, *35*(3),

97–104.

Mohammed, R. S., Reynolds, M., James, J., Williams, C., Mohammed, A., Ramsubhag, A., van

Oosterhout, C., & Cable, J. (2016). Getting into hot water: sick guppies frequent warmer

thermal conditions. *Oecologia*, *181*(3), 911–917.

Morgan, M. (2023). *DirichletMultinomial: Dirichlet-Multinomial Mixture Model Machine*

Learning for Microbiome Data. https://doi.org/10.18129/B9.bioc.DirichletMultinomial

Morris, J., Darolti, I., van der Bijl, W., & Mank, J. E. (2020). High-resolution characterization of

male ornamentation and re-evaluation of sex linkage in guppies. *Proceedings. Biological*

Sciences / The Royal Society, *287*(1937), 20201677.

Morrow, E. H. (2015). The evolution of sex differences in disease. *Biology of Sex Differences*,

6(1), 5.

Nater, A., Krützen, M., & Lindholm, A. K. (2008). Development of polymorphic microsatellite

markers for the livebearing fish Poecilia parae. *Molecular Ecology Resources*, *8*(4), 857–

860.

Nei, M., Maruyama, T., & Chakraborty, R. (1975). The Bottleneck Effect and Genetic Variability

in Populations. *Evolution; International Journal of Organic Evolution*, *29*(1), 1–10.

Nelson, P., & May, G. (2020). Defensive Symbiosis and the Evolution of Virulence. *The*

American Naturalist, *196*(3), 333–343.

Ni, J., Wu, G. D., Albenberg, L., & Tomov, V. T. (2017). Gut microbiota and IBD: causation or

correlation? *Nature Reviews. Gastroenterology & Hepatology*, *14*(10), 573–584.

Pagán, I., & García-Arenal, F. (2018). Tolerance to plant pathogens: Theory and experimental

evidence. *International Journal of Molecular Sciences*, *19*(3).

https://doi.org/10.3390/ijms19030810

Paterson, I. G., Crispo, E., Kinnison, M. T., Hendry, A. P., & Bentzen, P. (2005). Characterization

of tetranucleotide microsatellite markers in guppy (Poecilia reticulata). *Molecular

Ecology Notes*, *5*(2), 269–271.

Peig, J., & Green, A. J. (2009). New perspectives for estimating body condition from

mass/length data: the scaled mass index as an alternative method. *Oikos* , *118*(12), 1883–

1891.

Phillips, K. P., Cable, J., Mohammed, R. S., Chmielewski, S., Przesmycka, K. J., van Oosterhout,

C., & Radwan, J. (2021). Functional immunogenetic variation, rather than local

adaptation, predicts ectoparasite infection intensity in a model fish species. *Molecular

Ecology*, *30*(21), 5588–5604.

Phillips, K. P., Cable, J., Mohammed, R. S., Herdegen-Radwan, M., Raubic, J., Przesmycka, K.

J., van Oosterhout, C., & Radwan, J. (2018). Immunogenetic novelty confers a selective

advantage in host-pathogen coevolution. *Proceedings of the National Academy of

Sciences of the United States of America*, *115*(7), 1552–1557.

Piazzon, M. C., Naya-Català, F., Perera, E., Palenzuela, O., Sitjà-Bobadilla, A., & Pérez-

Sánchez, J. (2020). Genetic selection for growth drives differences in intestinal

microbiota composition and parasite disease resistance in gilthead sea bream.

Microbiome, *8*(1), 168.

R Core Team. (2020). *R: A language and environment for statistical computing* (Version v.4.1.0)

[R Foundation for Statistical Computing, Vienna, Austria.].

Råberg, L., Sim, D., & Read, A. F. (2007). Disentangling genetic variation for resistance and

tolerance to infectious diseases in animals. *Science*, *318*(5851), 812–814.

Raymond, M., & Rousset, F. (1995). GENEPOP (Version 1.2): Population Genetics Software for

Exact Tests and Ecumenicism. *The Journal of Heredity*, *86*(3), 248–249.

Rodríguez-Barreto, D., Sanz-González, J. C., Martín, M. V., Arrieta, J. M., & Almansa, E.

(2023). Sex-specific bacterial microbiome variation in octopus vulgaris skin. *Frontiers in*

Microbiology, *14*, 1233661.

Rovenolt, F., Walsman, J., Martin, C., Gotanda, K., & Stephenson, J. (2024). Parasite prevalence

depends on female preference: Integrating parasite-mediated sexual selection and

infectious   disease   dynamics.   In   *Authorea   Inc.*

https://doi.org/10.22541/au.172447480.06236699/v1

Santos-Marcos, J. A., Mora-Ortiz, M., Tena-Sempere, M., Lopez-Miranda, J., & Camargo, A.

(2023). Interaction between gut microbiota and sex hormones and their relation to sexual

dimorphism in metabolic diseases. *Biology of Sex Differences*, *14*(1), 4.

Scheifler, M., Sanchez-Brosseau, S., Magnanou, E., & Desdevises, Y. (2022). Diversity and

structure of sparids external microbiota (Teleostei) and its link with monogenean

ectoparasites. *Animal Microbiome*, *4*(1), 27.

Smallbone, W., Van Oosterhout, C., & Cable, J. (2016). The effects of inbreeding on disease

susceptibility: *Gyrodactylus turnbulli* infection of guppies, *Poecilia reticulata*.

Experimental Parasitology, *167*, 32–37.

Speer, K. A. (2022). [Review of *Microbiomes mediate host-parasite interactions*]. *Molecular Ecology*, *31*(7), 1925–1927.

Stephenson, J. F., & Adelman, J. S. (2022). The behavior of infected hosts: Behavioral tolerance, behavioral resilience, and their implications for behavioral competence. *Animal Behavior and Parasitism.* https://books.google.ca/books?hl=en&lr=&id=xSZ_EAAAQBAJ&oi=fnd&pg=PA287&ots=xW36pEEkt3&sig=asg6-91ifwE5LAwCypVYASxqwO4

Stephenson, J. F., Stevens, M., Troscianko, J., & Jokela, J. (2020). The size, symmetry, and color saturation of a male guppy's ornaments forecast his resistance to parasites. *The American Naturalist*, *196*(5), 597–608.

Stephenson, J. F., van Oosterhout, C., & Cable, J. (2015). Pace of life, predators and parasites: predator-induced life-history evolution in Trinidadian guppies predicts decrease in parasite tolerance. *Biology Letters*, *11*(11), 20150806.

Stephenson, J. F., van Oosterhout, C., Mohammed, R. S., & Cable, J. (2015). Parasites of Trinidadian guppies: evidence for sex- and age-specific trait-mediated indirect effects of predators. *Ecology*, *96*(2), 489–498.

Stoffel, M. A., Esser, M., Kardos, M., Humble, E., Nichols, H., David, P., & Hoffman, J. I. (2016). inbreedR: an R package for the analysis of inbreeding based on genetic markers. *Methods in Ecology and Evolution*, *7*(11), 1331–1339.

Tadiri, C. P., Fussmann, G. F., & Scott, M. E. (2021). Parasite spread in experimental metapopulations: resistance, tolerance and host competence. *Oikos (Copenhagen, Denmark)*, *130*(9), 1562–1571.

Tanpichai, P., Kusonmano, K., Jantrakajorn, S., Piamsomboon, P., & Wongtavatchai, J. (2025). Supplementation of Bacillus subtilis probiotics supports health and gut microbiome in barramundi (Lates calcarifer, Bloch 1790). *Aquaculture (Amsterdam, Netherlands)*, *596*(741815), 741815.

Trevelline, B. K., Stephenson, J. F., & Kohl, K. D. (2020). [Review of *Two's company, three's a crowd: Exploring how host-parasite-microbiota interactions may influence disease susceptibility and conservation of wildlife*]. *Molecular Ecology*, *29*(8), 1402–1405.

Van Oosterhout, C., Harris, P. D., & Cable, J. (2003). Marked variation in parasite resistance between two wild populations of the Trinidadian guppy, *Poecilia reticulata* (Pisces: Poeciliidae). *Biological Journal of the Linnean Society. Linnean Society of London*, *79*(4), 645–651.

van Oosterhout, C., Joyce, D. A., Cummings, S. M., Blais, J., Barson, N. J., Ramnarine, I. W., Mohammed, R. S., Persad, N., & Cable, J. (2006). Balancing selection, random genetic drift, and genetic variation at the major histocompatibility complex in two wild populations of guppies (*Poecilia reticulata*). *Evolution; International Journal of Organic Evolution*, *60*(12), 2562–2574.

van Wijk, S. J., Taylor, M. I., Creer, S., Dreyer, C., Rodrigues, F. M., Ramnarine, I. W., van Oosterhout, C., & Carvalho, G. R. (2013). Experimental harvesting of fish populations drives genetically based shifts in body size and maturation. *Frontiers in Ecology and the Environment*, *11*(4), 181–187.

Walsman, J. C., Duffy, M. A., Cáceres, C. E., & Hall, S. R. (2023). "Resistance Is Futile": Weaker Selection for Resistance by Abundant Parasites Increases Prevalence and Depresses Host Density. *The American Naturalist*, *201*(6), 864–879.

Walsman, J. C., Janecka, M. J., Clark, D. R., Kramp, R. D., Rovenolt, F., Patrick, R.,

Mohammed, R. S., Konczal, M., Cressler, C. E., & Stephenson, J. F. (2022). Shoaling

guppies evade predation but have deadlier parasites. *Nature Ecology & Evolution*, *6*(7),

945–954.

Walsman, J. C., Lambe, M., & Stephenson, J. F. (2024). Associating with kin selects for disease

resistance and against tolerance. *Proceedings. Biological Sciences*, *291*(2023), 20240356.

Wang, L.-C., Chen, L.-H., Chiu, Y.-C., Liou, C.-Y., Chen, H.-C., Lu, C.-Y., & Chen, J.-L. (2023).

Teleost skin microbiome: An intimate interplay between the environment and the host

immunity. *Fish & Shellfish Immunology*, *139*, 108869.

Watanabe, T., Yoshida, M., Nakajima, M., & Taniguchi, N. (2005). Linkage mapping of AFLP

and microsatellite DNA markers with the body color- and sex-determining loci in the

guppy (Poecilia reticulata). *Zoological Science*, *22*(8), 883–889.

Weiler, I. L. G., Kramp, R. D., Rovenolt, F., & Stephenson, J. F. (2025). Sex-dependent effects of

infection on guppy reproductive fitness and offspring parasite resistance. *The Journal of*

Animal Ecology. https://doi.org/10.1111/1365-2656.70006

Wickham, H. (2016). *ggplot2: Elegant Graphics for Data Analysis*. Springer.

Wilde, J., Slack, E., & Foster, K. R. (2024). Host control of the microbiome: Mechanisms,

evolution, and disease. *Science (New York, N.Y.)*, *385*(6706), eadi3338.

Zheng, D., Liwinski, T., & Elinav, E. (2020). Interaction between microbiota and immunity in

health and disease. *Cell Research*, *30*(6), 492–506.

Zuur, A. F., Ieno, E. N., Walker, N., Saveliev, A. A., & Smith, G. M. (2009). *Mixed Effects*

Models and Extensions in Ecology with R. Springer New York.

Conflict of Interest: No conflicts of interest to report for any authors

Author Contributions: Rachael Kramp and Jessica Stephenson conceived the ideas and

designed methodology; Jessica Stephenson and Nadine Tardent collected the data; Rachael

Kramp, Jessica Stephenson and Mary J. Janecka analysed the data; Rachael Kramp, Jessica

Stephenson, Kevin Kohl and Jukka Jokela led the writing and editing of the manuscript. All

authors contributed critically to the drafts and gave final approval for publication.

Animal use ethics statement: The permit to collect and export guppies was granted by the

Director of Fisheries in the Ministry of Agriculture, Land and Fisheries Division, Aquaculture

Unit of the Republic of Trinidad and Tobago. Import to Cardiff University was under Cefas APB

authorisation number CW054-D—187A. Fish husbandry and experimental infections of the F1

generation at Cardiff were under UK Home Office licence PPL 302876, with approval by the

Cardiff University Animal Ethics Committee. Guppies were shipped to Zürich, and their

husbandry and experimental infections of the F3-7 generations were approved by the

Veterinäramt of Kanton Zürich (ZH177/15).

Data availability statement: The raw experimental data, description of the data and analysis

code are available from the Dryad Digital Repository:

http://datadryad.org/share/UUQY9h2hXgGiWvvPM_SC6D2ii3ssYFAq71k3w4dFKlI

Funding: This work was supported by the Fisheries Society of the British Isles (studentship to

JFS), the Center for Adaptation to a Changing Environment at ETH Zürich (fellowship to JFS),

the University of Pittsburgh, and Howard Hughes Medical Institute (Gilliam Fellowship for

Advanced Study to RDK).

Figures

Figure 1. Artificial selection lines significantly predicted G. turnbulli infection severity in

Trinidadian guppies. Infection severity (worm days) reflects the area under the infection load

curve. Female guppies (n=44) in susceptible and resistant lines differed from the control line.

Male guppies (n=43) only showed differences in the susceptible line. Box plots illustrate median,

interquartile range, and values within 1.5× the interquartile range of infection severity.

807 ***Figure 2. Artificial selection lines also predicted relative tissue-specific tolerance in***

808 ***Trinidadian Guppies***. Tolerance measured as per-parasite mass change across infection

809 minus max tolerance. Female guppies (n=44) in resistant lines significantly differed from control

810 and susceptible lines. Male guppies (n=43) did not show significant differences. Box plots

811 illustrate the same metrics as in Figure 1.

Figure 3. The distribution of skin bacterial community types across guppy selection lines
showed no significant association. A. mosaic plot presents bacterial community types (A, B,
C) across control (c), resistant (r), and susceptible (s) lines, with no significant distribution
differences (Fisher's exact test, p=0.8). B. PCoA analysis (n=52) indicated bacterial community
type was significant in variation (PERMANOVA: F=7.85, P=0.001), but no significant clustering
was detected for selection lines (PERMANOVA: F=0.81, P=0.95).

Figure 4. Selection lines and bacterial community types predicted G. turnbulli infection

severity. A. Lines significantly predicted severity in swabbed fish (n=31; ANOVA: χ²=8.73,

P=0.013). B. Community type A had lower infection severity than types B and C (A-B, z=3.23,

P=0.0034; A-C, z=4.92, P<0.001). No interactions between community type and selection line

were detected (ANOVA: χ²=7.4, P=0.116). A combined model of line and bacterial types

explained infection severity better than either factor alone (χ²=13.40, P<0.001).

834